Growth of Turaev-Viro invariants and cabling

Renaud Detcherry

Abstract

The Chen-Yang volume conjecture [\[4\]](#page-5-0) states that the growth rate of the Turaev-Viro invariants of a compact oriented 3-manifold determines its simplicial volume. In this paper we prove that the Chen-Yang conjecture is stable under $(2n + 1, 2)$ cabling.

1 Introduction

For M a 3-manifold that is either closed or with boundary, the Turaev-Viro invariants $TV_r(M)$ are real valued topological invariants that can be computed using state sums over a triangulation of M . They depend of the choice of an $2r$ -th root of unity q whose square is a primitive r-th root; in this paper we will always choose $q = e^{\frac{2i\pi}{r}}$ and assume r odd. Moreover, when M is a manifold with empty or toroidal boundary, the invariants $TV_r(M)$ are always non-negative.

The geometric meaning of the Turaev-Viro invariants is hard to understand from their state sum defini-tion. However, a conjecture of Chen and Yang [\[4\]](#page-5-0) states that the asymptotics of the TV_r invariants at root $q = e^{\frac{2i\pi}{r}}$ is related to hyperbolic volume:

Conjecture 1.1. *[\[4\]](#page-5-0) For any hyperbolic manifold* M *(closed or with boundary), we have*

$$
\lim_{r \to \infty, r \text{ odd}} \frac{2\pi}{r} \log |TV_r(M, q = e^{\frac{2i\pi}{r}})| = \text{Vol}(M)
$$

where Vol(M) *is the hyperbolic volume of* M.

Conjecture [1.1](#page-0-0) is reminiscent of the Volume Conjecture of Kashaev and Murakami-Murakami [\[7\]](#page-5-1)[\[9\]](#page-5-2) where the TV_r invariants are replaced with evaluations $J_n(K, e^{\frac{2i\pi}{n}})$ of the normalized colored Jones polynomials of an hyperbolic knot.

In [\[6\]](#page-5-3), Yang, Kalfagianni and the author gave a formula relating the Turaev-Viro invariants of a link complement to colored Jones polynomials, establishing a connection between the two conjectures. In the same paper, Conjecture [1.1](#page-0-0) was proved for the complements of the figure-eight knot, the borromean link and knots of Gromov norm zero.

Let us define the growth rate of Turaev-Viro invariants by:

Definition 1.2. *Let* M *be a* 3*-manifold, closed or with boundary. Then the Turaev-Viro growth rate is*

$$
LTV(M) = \limsup_{r \to \infty, r \text{ odd}} \frac{2\pi}{r} \log |TV_r(M, q = e^{\frac{2i\pi}{r}})|.
$$

A way to restate Conjecture [1.1](#page-0-0) for general 3-manifold (not necessarily hyperbolic), is the following: Conjecture 1.3. *For any compact oriented* 3*-manifold* M, *we have*

$$
LTV(M) = Vol(M),
$$

where Vol(M) *is the simplicial volume of* M.

We recall that the simplicial volume of M can be thought either as the sum of the hyperbolic volumes of the hyperbolic pieces in the JSJ decomposition of M, or as $v_3||M||$ where $||M||$ is the Gromov norm of M . Note that the Turaev-Viro invariants sometimes vanish on lens spaces, thus replacing the limit by an upper limit is necessary.

In [\[5\]](#page-5-4), Kalfagianni and the author investigated the growth rate of Turaev-Viro invariants. They showed that the growth rate of Turaev-Viro invariants satisfies properties similar to that of the simplicial volume:

Theorem 1.4. *[\[5\]](#page-5-4) Let* M *be a compact oriented* 3*-manifold, with empty or toroidal boundary.*

- *(1)* If M is a Seifert manifold, then there exists constants $B > 0$ and N such that for any odd $r \ge 3$, *we have* $TV_r(M) \leqslant Br^N$ *and* $LTV(M) \leqslant 0$.
- (2) If M is a Dehn-filling of M', then $TV_r(M) \leq TV_r(M')$ and $LTV(M) \leq LTV(M')$.
- (3) If $M = M_1 \cup_{T} M_2$ is obtained by gluing two 3-manifolds M_1 and M_2 along a torus boundary *component, then* $TV_r(M) \leq TV_r(M_1)TV_r(M_2)$ *and* $LTV(M) \leq LTV(M_1) + LTV(M_2)$.

These properties are parallel to the properties of the simplicial volume: the simplicial volume of Seifert manifolds is 0, the simplicial volume decreases under Dehn-filling and is subadditive under gluing along tori [\[12\]](#page-5-5).

Let p, q be coprime integers, the (p, q) -cabling space is the complement of a (p, q) -torus knot standardly embedded in a solid torus. A (p, q) -cabling of a manifold M with toroidal boundary is a manifold M' obtained by gluing a (p, q) -cabling space to a boundary component of M. In this paper, we will investigate the compatibility of Conjecture [1.1](#page-0-0) with $(p, 2)$ -cabling. We will show the following:

Theorem 1.5. *Let* M *be a manifold with toroidal boundary and* M′ *be obtained by gluing a* (p, 2) *cabling space* $C_{p,2}$ *to a boundary component of M. Then there are constants* $B > 0$ *and* $N > 0$ *such that*

$$
\frac{1}{Br^N}TV_r(M') \leq TV_r(M) \leq Br^NTV_r(M').
$$

In particular, this means that if Conjecture [1.1](#page-0-0) is true for M then it is true for M' .

2 Preliminaries

2.1 Reshetikhin-Turaev SO_3 -TQFTs and TQFT basis

We briefly sketch the properties of Reshetikhin-Turaev SO_3 -TQFTs, defined by Reshetikhin and Turaev in [\[10\]](#page-5-6). We will introduce them in the skein-theoretic framework of Blanchet, Habegger, Masbaum and Vogel [\[3\]](#page-5-7). We refer to [\[2\]](#page-5-8)[\[3\]](#page-5-7) for the details of these constructions.

For any odd integer $r \geqslant 3$, and primitive 2r-th root of unity A, there is an associated TQFT functor RT_r , with the following properties:

- For Σ a closed compact oriented surface, $RT_r(\Sigma)$ is a finite dimensional C-vector space, with a natural Hermitian form. Moreover for a disjoint union $\Sigma \coprod \Sigma'$ one has $RT_r(\Sigma \coprod \Sigma') = RT_r(\Sigma) \otimes$ $RT_r(\Sigma').$
- For M a compact oriented closed 3-manifold, $RT_r(M)$ is the SO₃ Reshetikhin-Turaev invariant, a complex valued topological invariant of 3-manifolds, and for M with boundary, $RT_r(M)$ is a vector in $RT_r(\partial M)$.

- If (M, Σ_1, Σ_2) is a cobordism, $RT_r(M)$: $RT_r(\Sigma_1) \rightarrow RT_r(\Sigma_2)$ is a linear map. Moreover, the composition of cobordisms is sent to the composition of linear maps, up to a power of A.

Moreover, some basis of the TQFT spaces $RT_r(\Sigma)$ of surfaces has been explicitely described in [\[3\]](#page-5-7). We recall that for Σ a surface, $RT_r(\Sigma)$ is a quotient of the Kauffman module of a handlebody of the same genus.

In the case of a torus we get the following picture: the torus T^2 is the boundary of a solid torus $D^2 \times S^1$. One gets a family of elements of the Kauffman module of $D^2 \times S^1$ by taking the core $\{0\} \times S^1$ and coloring it by the $i - 1$ -th Jones-Wenzl idempotents, thus obtaining an element $e_i \in RT_r(T^2)$. For a definition of Jones-Wenzl idempotents we refer to [\[3\]](#page-5-7). For $r = 2m + 1$, and A a 2r-th root of unity, only finitely many Jones-Wenzl idemptotents can be defined, thus only the elements e_1, \ldots, e_{2m-1} are well defined (see [\[3\]](#page-5-7)[Lemma 3.2]).

As elements of the Kauffman module of the solid torus, the e_i 's can also be considered as elements of $RT_r(T^2)$. One gets a basis of $RT_r(T^2)$ consisting of elements e_i :

Theorem 2.1. *[\[3\]](#page-5-7)[Theorem 4.10] If* $r = 2m + 1 \ge 3$ *, then the family* $e_1, e_2, \ldots e_m$ *is an orthonormal basis of* $RT_r(T^2)$. *Moreover one has* $e_{m-i} = e_{m+1+i}$ *for* $0 \leq i \leq m-1$.

Note that the last part implies that the family $e_1, e_3, \ldots e_{2m-1}$ is the same basis of $RT_r(T^2)$ as the family $e_1, e_2, \ldots e_m$, in a different order.

As a consequence of TQFTs axioms, the RT_r vectors associated to link complements can be tied to values of colored Jones polynomials. Indeed, if $M = S^3 \setminus L$ is a link complement where L has n components, $RT_r(M)$ will be a vector in $RT_r(T^2)^{\otimes n}$ whose coefficient along $e_{i_1} \otimes e_{i_2} \dots \otimes e_{i_n}$ is obtained by gluing solid tori with cores colored by Jones-Wenzl idempotents to the boundary components of M and taking the Reshetikhin-Turaev invariant of that. Thus the coefficient we get is $\eta_r J_i(L, A^4)$, where $J_i(L, t)$ is the *i*-th unnormalized colored Jones polynomial of L, *i* is a multi-index of colors, and

$$
\eta_r = RT_r(S^3) = \frac{A^2 - A^{-2}}{\sqrt{-r}}.
$$

2.2 Relationship with the Turaev-Viro invariants

While the Turaev-Viro invariants of compact oriented 3-manifolds M are defined as state sums over triangulations of M (see [\[13\]](#page-5-9)), we will only use a well-known identity relating the Turaev-Viro invariants and Reshetikhin-Turaev invariants. This property was first proved by Roberts [\[11\]](#page-5-10) in the case of closed 3-manifolds, and extended to manifolds with boundary by Benedetti and Petronio [\[1\]](#page-5-11). For simplicity we state it only in the case of manifolds with toroidal boundary:

Theorem 2.2. Let M be a compact oriented manifold with toroidal boundary, let $r \geq 3$ be an odd *integer and let* A *be a primitive* 2r*-th root of unity. Then have*

$$
TV_r(M, A^2) = ||RT_r(M, A)||^2
$$

where $|| \cdot ||$ *is the natural Hermitian norm on* $RT_r(\partial M)$.

For T^2 a torus, the natural Hermitian form on $RT_r(T^2)$ is definite positive for any A, thus $TV_r(M)$ is non-negative.

2.3 The cabling formula

For $r = 2m + 1$, it is convenient to extend the definition of vectors $e_i \in RT_r(T^2)$ to all possible values of $i \in \mathbb{Z}$ as follows: we formally set $e_{-l} = -e_l$ for any $l \ge 0$ (and particular $e_0 = 0$) and

$$
e_{l+kr} = (-1)^k e_l
$$

for any $k \in \mathbb{Z}$. This is compatible with the above mentioned symmetry of the vectors e_i 's.

Recall that for p and q coprime integers with $q > 0$, the (p, q) -cabling space $C_{p,q}$ is the complement in a solid torus of a standardly embedded (p, q) -torus knot. The cobordism $C_{p,q}$ gives rise in TQFT to a linear map

$$
RT_r(C_{p,q}): RT_r(T^2) \to RT_r(T^2).
$$

We now describe the action of this map on the basis $\{e_i\}_{1\leq i\leq m}$ of $RT_r(T^2)$. By TQFT axioms, the map $RT_r(C_{p,q})$ sends the element e_i to the element of $RT_r(T^2)$ corresponding to a (p,q) -torus knot embedded in the solid torus and colored by the $i - 1$ -th Jones-Wenzl idempotent.

Morton computed these elements using skein calculus, yielding the following formula:

Theorem 2.3. *[\[8\]](#page-5-12) (Cabling formula) For any odd* $r = 2m + 1 \ge 3$, *for any* $1 \le i \le m$, *for any coprime integers* (p, q) *, one has:*

$$
RT_r(C_{p,2})(e_i) = A^{pq(i^2-1)/2} \sum_{k \in S_i} A^{-2pk(qk+1)} e_{2qk+1}.
$$

where S_i is the set

$$
\mathcal{S}_i = \{-\frac{i-1}{2}, -\frac{i-3}{2}, \dots, \frac{i-3}{2}, \frac{i-1}{2}\}.
$$

3 Stability under $(p, 2)$ -cabling

In this section, we will let $r = 2m + 1 \geq 3$ be an odd integer and let $A = e^{\frac{i\pi}{r}}$, which is a primitive 2r-th root of unity.

Recall that a (p, q) -cabling M' of a manifold M consist of gluing the exterior torus boundary component of $C_{p,q}$ to a boundary component of M. The JSJ decomposition of M' will consist of the pieces in the JSJ decomposition of M, plus an extra piece that is the cabling space $C_{p,q}$. As $C_{p,q}$ is a Seifert manifold, M and M' have the same simplicial volume.

Theorem 3.1. *Let* M *be a manifold with toroidal boundary, let* p *be an odd integer and let* M′ *be a* $(p, 2)$ -cabling of M. Then there exists constants $B > 0$ and N such that

$$
\frac{r^{-N}}{B}TV_r(M) \leq TV_r(M') \leqslant Br^NTV_r(M)
$$

In particular, we have $LTV(M) = LTV(M')$.

As M and M' in the theorem have the same volume, the conclusion implies that if Conjecture [1.1](#page-0-0) is true for M it is true for M' .

Proof. First, M' is obtained by gluing $C_{p,2}$, which is Seifert and thus has volume 0, to M along a torus. We know by Theorem [1.4](#page-1-0) that

$$
TV_r(M') \leq TV_r(M)TV_r(C_{p,2}).
$$

But as $C_{p,2}$ has volume 0, one has $TV_r(C_{p,2}) \leqslant Br^N$ for some constants $B > 0$ and N. To prove the other inequality, we study the map induced by $C_{p,2}$ in the RT_r -TQFT. If T is the boundary component coming from the $(p, 2)$ -torus knot and T' is the exterior boundary component, then $C_{p,2}$ induces a map

$$
RT_r(C_{p,2}): RT_r(T) \to RT_r(T').
$$

If M has only one boundary component, as RT_r is a TQFT, we have that $RT_r(M') = RT_r(C_{p,2})(RT_r(M))$. If M has other boundary components than the one used to glue $C_{p,2}$, then for any coloring i of the other boundary components of M , we have

$$
RT_r(M',i) = RT_r(C_{p,2})(RT_r(M,i)).
$$

In all cases, if $RT_r(C_{p,2})$ is invertible, we can write

$$
||RT_r(M)|| \le |||RT_r(C_{p,2})^{-1}||| \cdot ||RT_r(M')||,
$$

where $|| \cdot ||$ is the norm induced by the Hermitian form on $RT_r(\partial M)$ or $RT_r(\partial M')$ and $|||\cdot|||$ is the corresponding operator norm.

To conclude the proof of the theorem, we thus only need to prove that $RT_r(C_{p,2})$ is invertible, and that $|||RT_r(C_{p,2})^{-1}|||$ grows at most polynomially.

We can compute the matrix of $RT_r(C_{n,2})$ in the basis $e_1, \ldots e_m$ of $RT_r(T)$ by the cabling formula recalled above as Theorem [2.3.](#page-3-0)

For $q = 2$, the formula states:

$$
RT_r(C_{p,2})(e_i) = A^{p(i^2-1)} \sum_{k \in S_i} A^{-2pk(2k+1)} e_{4k+1}
$$

The cabling formula implies that the image lies in the vector space spanned by $e_1, e_3, \ldots, e_{2m-1}$, but by the Symmetry Principle recalled in [2.1,](#page-2-0) $e_{m-i} = e_{m+1+i}$ for all $0 \leq i \leq m-1$. Thus $\{e_1, e_3, \ldots, e_{2m-1}\}$ is actually the basis $\{e_1, \ldots e_m\}$ in a different order.

From the cabling formula we get that $RT_r(C_{p,2})(e_i)$ lies in $Span(e_1, e_3, \ldots e_{2i-1})$ and that the coefficient in e_{2j-1} in $RT_r(C_{p,2})(e_i)$ is $A^{p(i^2-1)} \cdot A^{p(j-j^2)}$ if j has same parity as i, and $-A^{p(i^2-1)} \cdot A^{p(j-j^2)}$ else. We can write $RT_r(C_{p,2})$ in the basis $\{e_1, \ldots e_m\}$ and $\{e_1, e_3, \ldots, e_{2m-1}\}$ as a product of two diagonal matrices and a triangular matrix:

$$
RT_r(C_{p,2}) = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & A^{p(2-2^2)} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & A^{p(m-m^2)} \end{pmatrix} \begin{pmatrix} 1 & -1 & 1 & \cdots \\ 0 & 1 & -1 & & \\ \vdots & & 1 & \ddots & \vdots \\ 0 & & & \ddots & -1 \\ 0 & & & & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & A^{p(2^2-1)} & \ddots & \vdots \\ \vdots & & \ddots & \ddots & 0 \\ 0 & & & 0 & A^{p(m^2-1)} \end{pmatrix}
$$

And thus we have:

$$
RT_r(C_{p,2})^{-1} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & A^{p(1-2^2)} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & A^{p(1-m^2)} \end{pmatrix} \begin{pmatrix} 1 & 1 & 0 & \dots & 0 \\ 0 & 1 & 1 & \vdots \\ \vdots & \ddots & 1 & \ddots \\ 0 & \dots & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & A^{p(2^2-2)} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & A^{p(m^2-m)} \end{pmatrix}
$$

The two diagonal matrices are isometries and the middle matrix clearly has norm bounded by a polynomial in r , which concludes the proof of Theorem [3.1.](#page-3-1) \Box

References

- [1] R. Benedetti and C. Petronio, *On Roberts' proof of the Turaev-Walker theorem*, J. Knot Theory Ramifications 5 (1996), no. 4, 427–439.
- [2] C. Blanchet, N. Habegger, G. Masbaum, and P. Vogel, *Remarks on the three-manifold invariants* θ_p , Operator algebras, mathematical physics, and low-dimensional topology (Istanbul, 1991), Res. Notes Math., vol. 5, A K Peters, Wellesley, MA, 1993, pp. 39–59.
- [3] , *Topological quantum field theories derived from the Kauffman bracket*, Topology 34 (1995), no. 4, 883–927.
- [4] Q. Chen and T. Yang, *Volume conjectures for the Reshetikhin-Turaev and the Turaev-Viro invariants*, arXiv:1503.02547.
- [5] R. Detcherry and E. Kalfagianni, *Gromov norm and Turaev-Viro invariants of 3-manifolds*, Arxiv 1705.09964.
- [6] R. Detcherry, E. Kalfagianni, and T. Yang, *Turaev Viro knot invariants and the colored Jones polynomials*, Arxiv 1701.07818.
- [7] R. M. Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. 39 (1997), no. 3, 269–275.
- [8] Morton, H. R., *The coloured Jones function and Alexander polynomial for torus knots*, Math. Proc. Cambridge Philos. Soc. 117 (1995), no. 1, 129–135.
- [9] H. Murakami and J. Murakami, *The colored Jones polynomials and the simplicial volume of a knot*, Acta Math. 186 (2001), no. 1, 85–104.
- [10] N. Reshetikhin and V. G. Turaev, *Invariants of* 3*-manifolds via link polynomials and quantum groups*, Invent. Math. 103 (1991), no. 3, 547–597.
- [11] J. Roberts, *Skein theory and Turaev-Viro invariants*, Topology 34 (1995), no. 4, 771–787.
- [12] W. P. Thurston, *The geometry and topology of three-manifolds*, Princeton Univ. Math. Dept. Notes, 1979.
- [13] V. G. Turaev and O. Y. Viro, *State sum invariants of* 3*-manifolds and quantum* 6j*-symbols*, Topology 4 (1992), 865—902.

Renaud Detcherry Department of Mathematics, Michigan State University East Lansing, MI 48824 (detcherry@math.msu.edu)