FOUR GENERA OF LINKS AND HEEGAARD FLOER HOMOLOGY

BEIBEI LIU

ABSTRACT. For links with vanishing pairwise linking numbers, the link components bound pairwise disjoint surfaces in B^4 . In this paper, we describe the set of genera of such surfaces in terms of the *h*-function, which is a link invariant from Heegaard Floer homology. In particular, we use the *h*-function to give lower bounds for the 4-genus of the link. For *L*-space links, the *h*-function is explicitly determined by Alexander polynomials of the link and sublinks. We show some *L*-space links where the lower bounds are sharp, and also describe all possible genera of disjoint surfaces bounded by such links.

1. INTRODUCTION

Let $\mathcal{L} = L_1 \cup L_2 \cdots \cup L_n$ be an oriented *n*-component link in S^3 with all linking numbers 0. Recall that a link bounds disjointly embedded surfaces in B^4 if and only if it has vanishing pairwise linking numbers. The 4-genus of \mathcal{L} is defined as:

$$g_4(\mathcal{L}) = \min\{\sum_{i=1}^n g_i \mid g_i = g(\Sigma_i), \Sigma_1 \sqcup \cdots \sqcup \Sigma_n \hookrightarrow B^4, \partial \Sigma_i = L_i\}.$$

If \mathcal{L} is a knot, the 4-genus is also known as the slice genus. Powell, Murasugi and Livingston showed lower bounds for the 4-genera of links in terms of the Levine-Tristram signatures, see [9, 10, 18]. Rasmussen defined the *h*-function (as an analogue of the Frøyshov invariant in Seiberg-Witten theory) for knots, and used it to obtain nontrivial lower bounds for the slice genus of a knot [19, 20]. We generalize Rasmussen's result and obtain lower bounds for the 4-genera of links with vanishing pairwise linking numbers. The *h*-function for links was introduced by Gorsky and Némethi [3]. It is closely related to *d*-invariants of large surgeries on links. For details, see Section 2.

We obtain lower bounds for the 4-genera of links in terms of the h-function. When the link has one component, we recover the lower bound for the slice genus given by Rasmussen. Here is our main result:

Theorem 1.1. Let $\mathcal{L} = L_1 \cup \cdots \cup L_n \subseteq S^3$ be an oriented link with vanishing pairwise linking numbers. Assume that the link components L_i bound pairwise disjoint, smoothly embedded surfaces $\Sigma_i \subseteq B^4$ of genera g_i . Then for any $\mathbf{v} = (v_1, \cdots, v_n) \in \mathbb{Z}^n$,

$$h(\boldsymbol{v}) \leq \sum_{i=1}^{n} f_{g_i}(v_i).$$

where $h(\mathbf{v})$ is the h-function of \mathcal{L} , and $f_{q_i} : \mathbb{Z} \to \mathbb{Z}$ is defined as follows:

$$f_{g_i}(v_i) = \begin{cases} \left\lceil \frac{g_i - |v_i|}{2} \right\rceil & |v_i| \le g_i \\ 0 & |v_i| > g_i \end{cases}$$

Corollary 1.2. For the link \mathcal{L} in Theorem 1.1, if $\mathbf{v} \succeq \mathbf{g}$, then $h(\mathbf{v}) = 0$ where $\mathbf{g} = (g_1, \dots, g_n)$.

The proof of Theorem 1.1 is inspired by Rasmussen's argument for knots [20]. We construct a non-positive definite Spin^c-cobordism from large surgeries on the link to the connected sum of circle bundles over closed, oriented surfaces of genera g_i . Ozsváth and Szabó proved the *d*-invariant inequality for a negative definite Spin^c-cobordism between rational homology spheres [11]. The *d*-invariant was generalized to *standard* 3-manifolds, and Rasmussen proved the *d*-invariant inequality for a non-positive definite Spin^c-cobordism between standard 3-manifolds [6, 20] (see Subsection 2.2). We apply the result, and obtain the inequality between the *d*-invariants of large surgeries on the link and *d*-invariants of circle bundles. By using the *h*-function of the link to compute *d*-invariants of large surgeries, we prove the inequality.

Theorem 1.3. If $\mathcal{L} = L_1 \cup \cdots \cup L_n \subset S^3$ is a (smoothly) slice L-space link, then \mathcal{L} is an unlink.

The idea of the proof goes as follows: the 4-genus for the slice link \mathcal{L} is 0. By Theorem 1.1, the *h*-function is identically 0. We compute the dual Thurston polytope of \mathcal{L} by using the properties of *L*-space links and prove that \mathcal{L} is an unlink. For details, see Subsection 3.2.

As an application of the inequality in Theorem 1.1, we can compare the following two sets. Let

$$\mathfrak{G}(\mathcal{L}) = \{ \boldsymbol{g} = (g_1, g_2, \cdots, g_n) \mid g_i = g(\Sigma_i), \Sigma_1 \sqcup \cdots \sqcup \Sigma_n \hookrightarrow B^4, \partial \Sigma_i = L_i \}.$$

and

$$\mathfrak{G}_{HF}(\mathcal{L}) = \{ \boldsymbol{v} = (v_1, \cdots, v_n) \mid h(\boldsymbol{v}) = 0 \text{ and } \boldsymbol{v} \succeq \boldsymbol{0} \}.$$

The 4-genus of the link \mathcal{L} equals $\min_{\boldsymbol{g} \in \mathfrak{G}(\mathcal{L})} (g_1 + \cdots + g_n)$. By Theorem 1.1, $\mathfrak{G}(\mathcal{L}) \subseteq \mathfrak{G}_{HF}(\mathcal{L})$.

If \mathcal{L} is an *L*-space link (see Definition 2.16), the *h*-function is explicitly determined by Alexander polynomials of the link and sublinks [1, Section 3.3]. We can describe the set $\mathfrak{G}_{HF}(\mathcal{L})$ in terms of Alexander polynomials explicitly (see Lemma 2.19). Moreover, let p_i, q_i be coprime positive integers where $1 \leq i \leq n$, and let $L_{(p_i,q_i)}$ denote the (p_i, q_i) -cable of L_i . Then the link $\mathcal{L}_{cab} = L_{(p_1,q_1)} \cup \cdots \cup L_{(p_n,q_n)}$ is also an *L*-space link if q_i/p_i is sufficiently large for each $1 \leq i \leq n$, [1, Proposition 2.8]. For example, let \mathcal{L} denote the 2-bridge link $b(4k^2 + 4k, -2k - 1)$ which is an *L*-space link [7]. Then for sufficiently large surgery coefficients, \mathcal{L}_{cab} is also an *L*-space link, and $\mathfrak{G}(\mathcal{L}_{cab}) = \mathfrak{G}_{HF}(\mathcal{L}_{cab})$ is shown as in Figure 1. For details, see Section 4.



FIGURE 1. The set $\mathfrak{G}(\mathcal{L}_{cab})$ for the cable link

Proposition 1.4. If $\mathcal{L} \subset S^3$ is an L-space link such that $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L})$, then for sufficiently large cables, \mathcal{L}_{cab} also satisfies that $\mathfrak{G}(\mathcal{L}_{cab}) = \mathfrak{G}_{HF}(\mathcal{L}_{cab})$.

Organization of the paper. In Subsection 2.1 and Subsection 2.2, we review the definitions of the *h*-function for links in S^3 and the *d*-invariants for standard 3-manifolds. In Subsection 2.3, we review the definition of *L*-space links, and the explicit formula to compute the *h*-function in terms of the Alexander polynomials of the link and sublinks. In Subsection 2.4, we review the Heegaard Floer link homologies of *L*-space links. In Section 3, we prove Theorem 1.1, Theorem 1.3, and give some lower bounds for the 4-genera of links. In Section 4, we show some examples of *L*-space links including the 2-bridge links $b(4k^2 + 4k, -2k - 1)$ where k is some positive integer, and prove that $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L})$ in these examples. Then the 4-genus is determined by the Alexander polynomials. We also show the proof of Proposition 1.4.

Notations and Conventions. In this paper, all the links are assumed to be oriented. We use \mathcal{L} to denote a link in S^3 , and L_1, \dots, L_n to denote the link components. Then \mathcal{L}_1 and \mathcal{L}_2 denote different links in S^3 , and L_1 and L_2 denote different components in the same link. We denote vectors in the *n*-dimension lattice \mathbb{Z}^n by bold letters. For two vectors $\boldsymbol{u} = (u_1, u_2, \dots, u_n)$ and $\boldsymbol{v} = (v_1, \dots, v_n)$ in \mathbb{Z}^n , we write $\boldsymbol{u} \leq \boldsymbol{v}$ if $u_i \leq v_i$ for each $1 \leq i \leq n$, and $\boldsymbol{u} \prec \boldsymbol{v}$ if $\boldsymbol{u} \leq \boldsymbol{v}$ and $\boldsymbol{u} \neq \boldsymbol{v}$. Let \boldsymbol{e}_i denote a vector in \mathbb{Z}^n where the *i*th-entry is 1 and other entries are 0. For a subset $B \subset \{1, \dots, n\}$, let $\boldsymbol{e}_B = \sum_{i \in B} \boldsymbol{e}_i$. Similarly, we use $L_B \subset \mathcal{L}$ to denote the sublink $\cup_{i \in B} L_i$. Assume $\{1, \dots, n\} - B = \{i_1, \dots, i_k\}$. For $\boldsymbol{y} = (y_1, \dots, y_n) \in \mathbb{Z}^n$, let $\boldsymbol{y} \setminus y_B = (y_{i_1}, \dots, y_{i_k})$. Let $\Delta_{\mathcal{L}}(t_1, \dots, t_n)$ denote the symmetrized Alexander polynomial of \mathcal{L} . Throughout this paper, we work over the field $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$.

Acknowledgements. I deeply appreciate Eugene Gorsky for introducing this interesting question to me and his patient guidance and helpful discussions during the project. I am also grateful to Allison Moore, Robert Lipschtiz, Jacob Rasmussen, Zhongtao Wu for useful discussions. The project is partially supported by NSF grant DMS-1700814.

2. Background

2.1. The *h*-function. Ozsváth and Szabó associated chain complexes $CF^{-}(M), CF(M), CF^{\infty}(M)$ and $CF^{+}(M)$ to an admissible Heegaard diagram for a closed oriented connected 3-manifold M [13]. The homologies of these chain complexes are called Heegaard Floer homologies $HF^{-}(M), \widehat{HF}(M), HF^{\infty}(M)$ and $HF^{+}(M)$, which are 3-manifold invariants. A link $\mathcal{L} = L_1 \cup \cdots \cup L_n$ in M defines a filtration on the link Floer complex $CF^{-}(M)$ [8, 16]. For links in S^3 , the filtration is indexed by an *n*-dimensional lattice \mathbb{H} which is defined as follows:

Definition 2.1. For an oriented link $\mathcal{L} = L_1 \cup \cdots \cup L_n \subset S^3$, define $\mathbb{H}(\mathcal{L})$ to be an affine lattice over \mathbb{Z}^n :

$$\mathbb{H}(\mathcal{L}) = \oplus_{i=1}^{n} \mathbb{H}_{i}(\mathcal{L}), \quad \mathbb{H}_{i}(\mathcal{L}) = \mathbb{Z} + \frac{lk(L_{i}, \mathcal{L} \setminus L_{i})}{2}$$

where $lk(L_i, \mathcal{L} \setminus L_i)$ denotes the linking number of L_i and $\mathcal{L} \setminus L_i$.

Given $\mathbf{s} = (s_1, \dots, s_n) \in \mathbb{H}(\mathcal{L})$, the generalized Heegaard Floer complex $A^-(\mathcal{L}, s)$ is defined to be a subcomplex of $CF^-(S^3)$ corresponding to the filtration indexed by \mathbf{s} [8]. For $\mathbf{v} \leq \mathbf{s}$, $A^-(\mathcal{L}, \mathbf{v}) \subseteq A^-(\mathcal{L}, \mathbf{s})$. The link homology HFL^- is defined as the homology of the associated graded complex:

(2.1)
$$HFL^{-}(\mathcal{L}, \boldsymbol{s}) = H_{*}\left(A^{-}(\mathcal{L}, \boldsymbol{s}) / \sum_{\boldsymbol{v} \prec \boldsymbol{s}} A^{-}(\mathcal{L}, \boldsymbol{v})\right).$$

The complex $A^{-}(\mathcal{L}, \mathbf{s})$ is a finitely generated module over the polynomial ring $\mathbb{F}[U_1, \dots, U_n]$ where the action of U_i drops the homological grading by 2 and drops the *i*-th filtration A_i by 1 [16]. Hence, $U_i A^{-}(\mathcal{L}, \mathbf{s}) \subseteq A^{-}(\mathcal{L}, \mathbf{s} - \mathbf{e}_i)$. All the actions U_i are homotopic to each other on each $A^{-}(\mathcal{L}, \mathbf{s})$, and the homology of $A^{-}(\mathcal{L}, \mathbf{s})$ can be regarded as a $\mathbb{F}[U]$ -module where U acts as U_1 [3, 16].

By the large surgery theorem [8], the homology of $A^{-}(\mathcal{L}, \mathbf{s})$ is isomorphic to the Heegaard Floer homology of a large surgery on the link \mathcal{L} equipped with some Spin^c-structure as a $\mathbb{F}[U]$ -module [8]. Then the homology of $A^{-}(\mathcal{L}, \mathbf{s})$ consists of the free part which is a direct sum of one copy of $\mathbb{F}[U]$ and some U-torsion.

Definition 2.2. [1, Definition 3.9] For an oriented link $\mathcal{L} \subseteq S^3$, we define the *H*-function $H_{\mathcal{L}}(s)$ by saying that $-2H_{\mathcal{L}}(s)$ is the maximal homological degree of the free part of $H_*(A^-(\mathcal{L}, s))$ where $s \in \mathbb{H}$.

Lemma 2.3. [1, Proposition 3.10] For an oriented link $\mathcal{L} \subseteq S^3$, the *H*-function $H_{\mathcal{L}}(s)$ takes nonnegative values, and $H_{\mathcal{L}}(s - e_i) = H_{\mathcal{L}}(s)$ or $H_{\mathcal{L}}(s - e_i) = H_{\mathcal{L}}(s) + 1$ where $s \in \mathbb{H}$.

For an *n*-component link \mathcal{L} with vanishing pairwise linking numbers, $\mathbb{H}(\mathcal{L}) = \mathbb{Z}^n$. The *h*-function $h_{\mathcal{L}}(s)$ is defined as

$$h_{\mathcal{L}}(\boldsymbol{s}) = H_{\mathcal{L}}(\boldsymbol{s}) - H_O(\boldsymbol{s})$$

where O denotes the unlink with n components, and $\mathbf{s} \in \mathbb{Z}^n$. Recall that for split links \mathcal{L} , the H-function $H(\mathcal{L}, \mathbf{s}) = H_{L_1}(s_1) + \cdots + H_{L_n}(s_n)$ where $H_{L_i}(s_i)$ is the H-function of the link component L_i , [1, Proposition 3.11]. Then $H_O(\mathbf{s}) = H(s_1) + \cdots + H(s_n)$ where $H(s_i)$ denotes the H-function of the unknot. More precisely, $H_O(\mathbf{s}) = \sum_{i=1}^n (|s_i| - s_i)/2$. Then $H_{\mathcal{L}}(\mathbf{s}) = h_{\mathcal{L}}(\mathbf{s})$ for all $\mathbf{s} \succeq \mathbf{0}$.

For the rest of this subsection, we use $\mathcal{L} = L_1 \cup \cdots \cup L_n \subset S^3$ to denote links with vanishing pairwise linking numbers. Consider the set

$$\mathfrak{G}_{HF}(\mathcal{L}) = \{ \boldsymbol{s} = (s_1, \cdots, s_n) \in \mathbb{Z}^n \mid h(\boldsymbol{s}) = 0, \boldsymbol{s} \succeq \boldsymbol{0} \}.$$

We obtain the following properties of the set $\mathfrak{G}_{HF}(\mathcal{L})$.

Lemma 2.4. If $x \in \mathfrak{G}_{HF}(\mathcal{L})$ and $y \succeq x$, then $y \in \mathfrak{G}_{HF}(\mathcal{L})$. Equivalently, if $x \notin \mathfrak{G}_{HF}(\mathcal{L})$ and $y \preceq x$, then $y \notin \mathfrak{G}_{HF}(\mathcal{L})$.

Proof. This is straightforward from Lemma 2.3.

Lemma 2.5. If $s = (s_1, \dots, s_n) \in \mathfrak{G}_{HF}(\mathcal{L})$, then $s \setminus s_i \in \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$ for all $1 \leq i \leq n$. Moreover, if $s \setminus s_i \in \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$, then for s_i sufficiently large, $s = (s_1, \dots, s_n) \in \mathfrak{G}_{HF}(\mathcal{L})$.

Proof. For an oriented link \mathcal{L} , there exists a natural forgetful map $\pi_i : \mathbb{H}(\mathcal{L}) \to \mathbb{H}(\mathcal{L} \setminus L_i)$ [8]. If \mathcal{L} has vanishing pairwise linking numbers, $\pi_i(\mathbf{s}) = \mathbf{s} \setminus s_i$ where $\mathbf{s} \in \mathbb{Z}^n$. Suppose that $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$. Then $h_{\mathcal{L}}(\mathbf{s}) = H_{\mathcal{L}}(\mathbf{s}) = 0$. By Lemma 2.3, $H_{\mathcal{L}}(\mathbf{s} + t\mathbf{e}_i) = 0$ for all i and t > 0. Recall that $H_{\mathcal{L}}(\mathbf{s} + t\mathbf{e}_i) = H_{\mathcal{L}\setminus L_i}(\mathbf{s} \setminus s_i)$ for sufficiently large t, [1, Proposition 3.12]. Then $H_{\mathcal{L}\setminus L_i}(\mathbf{s}\setminus s_i) = 0$. Thus, $\mathbf{s} \setminus s_i \in \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$.

Conversely, if $\mathbf{s} \setminus s_i \in \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$, and s_i is sufficiently large, then $H_{\mathcal{L}}(\mathbf{s}) = H_{\mathcal{L} \setminus L_i}(\mathbf{s} \setminus s_i) = 0$, which implies that $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$.

Definition 2.6. A lattice point $s \in \mathbb{Z}^n$ is maximal if $s \notin \mathfrak{G}_{HF}(\mathcal{L})$, but $s + e_i \in \mathfrak{G}_{HF}(\mathcal{L})$ for all $1 \leq i \leq n$.

Lemma 2.7. The set $\mathfrak{G}_{HF}(\mathcal{L})$ is determined by the set of maximal lattice points and $\mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$ for all $1 \leq i \leq n$.

Proof. We claim that $\boldsymbol{x} \notin \mathfrak{G}_{HF}(L)$ if and only if either $\boldsymbol{x} \leq \boldsymbol{z}$ for some maximal lattice point $\boldsymbol{z} \in \mathbb{Z}^n$ or $\boldsymbol{x} \setminus x_i \notin \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$ for some *i* where $\boldsymbol{x} = (x_1, \cdots, x_n)$. For the "if" part, assume that $\boldsymbol{x} \in \mathfrak{G}_{HF}(\mathcal{L})$. Then $\boldsymbol{z} \in \mathfrak{G}_{HF}(\mathcal{L})$ if $\boldsymbol{z} \succeq \boldsymbol{x}$ and $\boldsymbol{x} \setminus x_i \in \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$ for all *i* by Lemma 2.4 and Lemma 2.5, which contradicts to the assumption.

For the "only if" part, assume that $\boldsymbol{x} \notin \mathfrak{G}_{HF}(\mathcal{L})$ and $\boldsymbol{x} \setminus x_i \in \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$ for all *i*. It suffices to find a maximal lattice point \boldsymbol{z} such that $\boldsymbol{x} \preceq \boldsymbol{z}$. If $H_{\mathcal{L}}(\boldsymbol{x} + \boldsymbol{e}_i) = 0$ for all *i*, we let $\boldsymbol{z} = \boldsymbol{x}$. Otherwise, suppose $H_{\mathcal{L}}(\boldsymbol{x} + \boldsymbol{e}_i) \neq 0$ for some $1 \leq i \leq n$. There exists some constant t_i such that $H_{\mathcal{L}}(\boldsymbol{x} + t_i \boldsymbol{e}_i) \neq 0$, and $H_{\mathcal{L}}(\boldsymbol{x} + (t_i + 1)\boldsymbol{e}_i) = 0$ since $\boldsymbol{x} \setminus x_i \in \mathfrak{G}_{HF}(\mathcal{L} \setminus L_i)$. If for all $j \neq i$, $H_{\mathcal{L}}(\boldsymbol{x} + t_i \boldsymbol{e}_i + \boldsymbol{e}_j) = 0$, we let $\boldsymbol{z} = \boldsymbol{x} + t_i \boldsymbol{e}_i$. Otherwise, we repeat this process. The process stops after finite steps. Thus, there exists a maximal lattice point \boldsymbol{z} such that $\boldsymbol{x} \preceq \boldsymbol{z}$.

2.2. The *d*-invariant. For a rational homology sphere M with a Spin^c-structure \mathfrak{s} , the Heegaard Floer homology $HF^{\infty}(M,\mathfrak{s}) \cong \mathbb{F}[U, U^{-1}]$ and $HF^+(M,\mathfrak{s})$ is absolutely graded where the free part is isomorphic to $\mathbb{F}[U^{-1}]$. Define the *d*-invariant of (M,\mathfrak{s}) to be the absolute grading of $1 \in \mathbb{F}[U^{-1}]$ [11]. Equivalently, the *d*-invariant of (M,\mathfrak{s}) is the maximum grading of $x \in HF^-(M,\mathfrak{s})$, which has a nontrivial image in $HF^{\infty}(M,\mathfrak{s})$.

We define *standard* 3-manifolds following [11, Section 9]:

Definition 2.8. A closed, oriented 3-manifold M is *standard* if for each torsion Spin^c-structure \mathfrak{s} ,

$$HF^{\infty}(M,\mathfrak{s}) \cong (\wedge^{b}H^{1}(M,\mathbb{F})) \otimes_{\mathbb{F}} \mathbb{F}[U,U^{-1}],$$

where $b = b_1(M)$.

Remark 2.9. If M is standard, then $\operatorname{rk} HF^{\infty}(M, \mathfrak{s}) = 2^{b}$ as a $\mathbb{F}[U, U^{-1}]$ -module.

Let M_1 and M_2 be a pair of oriented closed 3-manifolds equipped with Spin^c-structures \mathfrak{s}_1 and \mathfrak{s}_2 respectively. There is a connected sum formula for the Heegaard Floer homology [13, Theorem 6.2]:

$$HF^{\infty}(M_1 \# M_2, \mathfrak{s}_1 \# \mathfrak{s}_2) \cong H_*(CF^{\infty}(M_1, \mathfrak{s}_1) \otimes_{\mathbb{F}[U, U^{-1}]} CF^{\infty}(M_2, \mathfrak{s}_2)).$$

By the algebraic Künneth theorem, if M_1 and M_2 are standard, then $M_1 \# M_2$ is also standard.

If a 3-manifold M has a positive first Betti number (i.e $b_1(M) > 0$), the exterior algebra $\Lambda^*(H_1(M;\mathbb{F}))$ acts on the homology groups $HF^{\infty}(M,\mathfrak{s}), HF^+(M,\mathfrak{s}), HF^-(M,\mathfrak{s})$ and $\widehat{HF}(M,\mathfrak{s}), [13, \text{Section 4.2.5}]$. Define the subgroup $\mathfrak{A}_{\mathfrak{s}} \subset HF^{\infty}(M,\mathfrak{s})$ as follows:

$$\mathfrak{A}_{\mathfrak{s}} = \{ x \in HF^{\infty}(M, \mathfrak{s}) \mid \gamma \cdot x = 0 \quad \forall \gamma \in H_1(M, \mathbb{F}) \}.$$

If M is standard, $\mathfrak{A}_{\mathfrak{s}} \cong \mathbb{F}[U, U^{-1}]$, and its image under the map $\pi : HF^{\infty}(M, \mathfrak{s}) \to HF^+(M, \mathfrak{s})$ is isomorphic to $\mathbb{F}[U^{-1}]$.

Definition 2.10. For a standard 3-manifold M equipped with a torsion Spin^c-structure \mathfrak{s} , the *d*-invariant $d(M, \mathfrak{s})$ is defined as the absolute grading of $1 \in \pi(\mathfrak{A}_{\mathfrak{s}}) \cong \mathbb{F}[U^{-1}]$.

Given rational homology spheres Y_1, Y_2 and a negative definite Spin^c-cobordism (W, \mathfrak{t}) from (Y_1, \mathfrak{t}_1) to (Y_2, \mathfrak{t}_2) , the *d*-invariant of Y_2 is no smaller than the *d*-invariant of Y_1 up to the degree shift of the cobordism [11]. Rasmussen generalized this inequality for nonpositive definite Spin^c-cobordisms from rational homology spheres to standard 3-manifolds [20].

Proposition 2.11. [20] Suppose that W is a connected cobordism from closed, oriented and connected 3-manifolds Y_1 to Y_2 such that $b_1(Y_1) = b_1(W) = b_2^+(W) = 0$, and Y_2 is standard. Let \mathfrak{s} be a Spin^c-structure on W whose restriction \mathfrak{s}_i to Y_i is torsion. Then

$$F_{W,\mathfrak{s}}^{\infty}: HF^{\infty}(Y_1,\mathfrak{s}_1) \to HF^{\infty}(Y_2,\mathfrak{s}_2)$$

maps $HF^{\infty}(Y_1,\mathfrak{s}_1)$ isomorphically onto $\mathfrak{A}_{\mathfrak{s}_2} \subset HF^{\infty}(Y_2,\mathfrak{s}_2)$. Moreover, we have

$$d(Y_1,\mathfrak{s}_1) + \deg F_{W\mathfrak{s}}^+ \le d(Y_2,\mathfrak{s}_2)$$

where the degree of $F_{W,\mathfrak{s}}^+$: $HF^+(Y_1,\mathfrak{s}_1) \to HF^+(Y_2,\mathfrak{s}_2)$ is $\frac{1}{4}(c_1^2(\mathfrak{s}) - 3\sigma(W) - 2\chi(W)).$

The *d*-invariants of large surgeries on a link $\mathcal{L} = L_1 \cup \cdots \cup L_n \subset S^3$ can be computed in terms of the *H*-function of the link by the large surgery theorem [8]. Choose a framing vector $\boldsymbol{q} = (q_1, \cdots, q_n)$ where q_1, \cdots, q_n are sufficiently large. Let Λ denote the linking matrix where Λ_{ij} is the linking number of L_i and L_j when $i \neq j$, and $\Lambda_{ii} = q_i$.

Attach *n* 2-handles to the 4-ball B^4 along $L_1, L_2 \cdots, L_n$ with framings q_1, \cdots, q_n . We obtain a 2-handlebody W with boundary $\partial W = S^3_q(\mathcal{L})$ which is the 3-manifold obtained by doing surgery along L_1, L_2, \cdots, L_n with surgery coefficients q_1, \cdots, q_n respectively. Assume that $\det(\Lambda) \neq 0$, then $S^3_q(\mathcal{L})$ is a rational homology sphere with $|H_1(S^3_q(\mathcal{L}))| = |\det(\Lambda)|$. Note that if \mathcal{L} has vanishing pairwise linking numbers, then Λ is a diagonal matrix with $\Lambda_{ii} = q_i$, and $\det(\Lambda) = q_1 \cdots q_n \neq 0$. The Spin^{*c*}-structures on $S^3_q(\mathcal{L})$ are enumerated as follows:

Lemma 2.12. [8, Section 9.3] There are natural identifications:

$$H^2(S^3_{\boldsymbol{a}}(\mathcal{L})) \cong H_1(S^3_{\boldsymbol{a}}(\mathcal{L})) \cong \mathbb{Z}^n / \mathbb{Z}^n \Lambda$$

such that $c_1(\mathfrak{s}) = [2\mathfrak{s}]$ for any $\mathfrak{s} \in \operatorname{Spin}^c(S^3_{\mathfrak{q}}(\mathcal{L})) \cong \mathbb{H}(\mathcal{L})/\mathbb{Z}^n\Lambda$.

Fix $\zeta = (\zeta_1, \dots, \zeta_n)$, a small real vector whose entries are linearly independent over \mathbb{Q} . Let $P(\Lambda)$ be the hyper-parallelepiped with vertices

$$\zeta + \frac{1}{2}(\pm\Lambda_1,\pm\Lambda_2,\cdots,\pm\Lambda_n),$$

where all combinations of the signs are used and $\Lambda_1, \dots, \Lambda_n$ are column vectors of the matrix Λ . Denote

$$P_{\mathbb{H}}(\Lambda) = P(\Lambda) \cap \mathbb{H}(\mathcal{L}),$$

where $\mathbb{H}(\mathcal{L})$ is the lattice for \mathcal{L} .

Proposition 2.13. [8, Section 10.1] For any $v \in P_{\mathbb{H}}(\Lambda)$ there exists a unique Spin^c-structure \mathfrak{s}_{v} on $S_{q}(\mathcal{L})$ which extends to a Spin^c-structure \mathfrak{t}_{v} on W with $c_{1}(\mathfrak{t}_{v}) = 2v - (\Lambda_{1} + \cdots + \Lambda_{n})$.

Remove a ball B^4 from the 2-handlebody W. We obtain a Spin^c-cobordism \mathcal{U} from (S^3, \mathfrak{s}_0) to $(S^3_{\boldsymbol{q}}(\mathcal{L}), \mathfrak{s}_{\boldsymbol{v}})$. By reversing the orientation of \mathcal{U} , we obtain a Spin^c-cobordism \mathcal{U}' equipped with the Spin^c-structure $\mathfrak{t}_{\boldsymbol{v}}$ from $(S^3_{\boldsymbol{q}}(\mathcal{L}), \mathfrak{s}_{\boldsymbol{v}})$ to (S^3, \mathfrak{s}_0) .

Theorem 2.14. [1, 8] For $v \in P_{\mathbb{H}}(\Lambda)$, the d-invariant of a large surgery with surgery coefficients q on \mathcal{L} is given by

$$d(S^{3}_{\boldsymbol{q}}(\mathcal{L}), \boldsymbol{\mathfrak{s}}_{\boldsymbol{v}}) = -\mathrm{deg}F_{(\mathcal{U}', \boldsymbol{\mathfrak{t}}_{\boldsymbol{v}})} - 2H(\boldsymbol{v}),$$

where $\deg F_{\mathcal{U}',\mathfrak{t}_{v}}$ is the grading shift of the cobordism \mathcal{U}' with $Spin^{c}$ -structure \mathfrak{t}_{v} . It does not depend on the link, but depends on the linking matrix Λ .

7

2.3. The *h*-function of *L*-space links. In [14], Ozsváth and Szabó introduced the concept of *L*-spaces.

Definition 2.15. A 3-manifold M is an L-space if it is a rational homology sphere and its Heegaard Floer homology has minimal possible rank: for any Spin^c -structure \mathfrak{s} , $\widehat{HF}(M, \mathfrak{s}) = \mathbb{F}$, and $HF^-(Y, \mathfrak{s})$ is a free $\mathbb{F}[U]$ -module of rank 1.

In terms of the large surgery, Gorsky and Némethi defined L-space links in [3].

Definition 2.16. An oriented *n*-component link $\mathcal{L} \subset S^3$ is an *L*-space link if there exists $\mathbf{0} \prec \mathbf{p} \in \mathbb{Z}^n$ such that the surgery manifold $S^3_{\mathbf{q}}$ is an *L*-space for any $\mathbf{q} \succeq \mathbf{p}$.

For L-space links \mathcal{L} , $H_*(A^-(\mathcal{L}, \mathbf{s})) = \mathbb{F}[[U]]$ [7]. By Equation (2.1) and the inclusionexclusion formula, one can write [1]:

(2.2)
$$\chi(HFL^{-}(\mathcal{L}, \mathbf{s})) = \sum_{B \subset \{1, \cdots, n\}} (-1)^{|B|-1} H_{\mathcal{L}}(\mathbf{s} - \mathbf{e}_{B}).$$

The Euler characteristic $\chi(HFL^{-}(\mathcal{L}, \mathbf{s}))$ was computed in [16]:

(2.3)
$$\tilde{\Delta}(t_1,\cdots,t_n) := \sum_{\boldsymbol{s}\in\mathbb{H}(\mathcal{L})} \chi(HFL^{-}(\mathcal{L},\boldsymbol{s}))t_1^{s_1}\cdots t_n^{s_n}$$

where $\boldsymbol{s} = (s_1, \cdots, s_n)$, and

$$\tilde{\Delta}_{\mathcal{L}}(t_1,\cdots,t_n) := \begin{cases} (t_1\cdots t_n)^{1/2} \Delta_{\mathcal{L}}(t_1,\cdots,t_n) & \text{if } n > 1, \\ \Delta_{\mathcal{L}}(t)/(1-t^{-1}) & \text{if } n = 1. \end{cases}$$

Theorem 2.17. [3] The H-function of an L-space link is determined by the Alexander polynomials of its sublinks as follows:

(2.4)
$$H_{\mathcal{L}}(\boldsymbol{s}) = \sum_{\mathcal{L}' \subset \mathcal{L}} (-1)^{\# \mathcal{L}' - 1} \sum_{\boldsymbol{u}' \succeq \pi_{\mathcal{L}'}(\boldsymbol{s} + 1)} \chi(HFL^{-}(\mathcal{L}', \boldsymbol{u}')),$$

where $1 = (1, \dots, 1)$.

Remark 2.18. For *L*-space links with two components, the explicit formula for the *H*-function can also be found in [7].

Consider *L*-space links \mathcal{L} with vanishing pairwise linking numbers. The set $\mathfrak{G}_{HF}(\mathcal{L})$ can also be described in terms of Alexander polynomials of the link and sublinks.

Lemma 2.19. For an n-component L-space link $\mathcal{L} \subseteq S^3$ with vanishing pairwise linking numbers, $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$ if and only if for all $\mathbf{y} = (y_1, \dots, y_n) \succ \mathbf{s}$, the coefficients of $t_1^{y_1} \cdots t_n^{y_n}$ in $\tilde{\Delta}_{\mathcal{L}}(t_1, \dots, t_n)$ are 0, and the coefficients corresponding to $\mathbf{y} \setminus y_B$ in $\tilde{\Delta}_{\mathcal{L} \setminus L_B}(t_{i_1}, \dots, t_{i_k})$ are also 0 for all $B \subset \{1, \dots, n\}$.

Proof. For the "if" part, observe that $\chi(HFL^{-}(\mathcal{L}, \boldsymbol{y})) = 0$ and $\chi(HFL^{-}(\mathcal{L}\backslash L_{B}), \boldsymbol{y}\backslash y_{B}) = 0$ for all $\boldsymbol{y} \succ \boldsymbol{s}$ and $B \subset \{1, \dots, n\}$ by Equation (2.3). Then $H_{\mathcal{L}}(\boldsymbol{s}) = 0$ by Theorem 2.17, and $\boldsymbol{s} \in \mathfrak{G}_{HF}(\mathcal{L})$. For the "only if" part, suppose that $\boldsymbol{s} \notin \mathfrak{G}_{HF}(\mathcal{L})$. By Lemma 2.7, either there exists a maximal vector $\boldsymbol{z} \notin \mathfrak{G}_{HF}(\mathcal{L})$ such that $\boldsymbol{s} \preceq \boldsymbol{z}$ or there exists some $1 \leq j \leq n$, such that $\boldsymbol{s} \backslash s_{j} \notin \mathfrak{G}_{HF}(\mathcal{L} \backslash L_{j})$. We claim that for all maximal lattice points \boldsymbol{z} , $\chi(HFL^{-}(\mathcal{L}, \boldsymbol{z}+\boldsymbol{1})) \neq 0$. Since \boldsymbol{z} is maximal, $h_{\mathcal{L}}(\boldsymbol{z}) = 1$, and for any subset $B \subset \{1, \dots, n\}$, $h_{\mathcal{L}}(\boldsymbol{z} + \boldsymbol{e}_{B}) = 0$. By Equation (2.2), $\chi(HFL^{-}(\mathcal{L}, \boldsymbol{z} + \boldsymbol{1})) = (-1)^{n} \neq 0$. If $\boldsymbol{s} \preceq \boldsymbol{z}$, the coefficient of $\boldsymbol{z} + \boldsymbol{1} \succ \boldsymbol{s}$ in $\tilde{\Delta}_{\mathcal{L}}(t_{1}, \dots, t_{n})$ equals 0 which contradicts to our assumption. If $\boldsymbol{s} \backslash s_{i} \notin \mathfrak{G}_{HF}(\mathcal{L} \backslash L_{i})$, we use the induction to get a contradiction.

2.4. The Heegaard Floer link homology. Ozsváth and Szabó associated the multigraded link invariants $HFL^{-}(\mathcal{L})$ and $\widehat{HFL}(\mathcal{L})$ to links $\mathcal{L} \subset S^{3}$ where $HFL^{-}(L)$ was defined in Equation (2.1), and $\widehat{HFL}(L)$ is defined as follows [2, 16]:

$$\widehat{HFL}(\mathcal{L}, \boldsymbol{s}) = H_*\left(A^-(\mathcal{L}, \boldsymbol{s}) / \left[\sum_{i=1}^n A^-(\boldsymbol{s} - \boldsymbol{e}_i) \oplus \sum_{i=1}^n U_i A^-(\boldsymbol{s} + \boldsymbol{e}_i)\right]\right)$$

If \mathcal{L} is an *L*-space link, there exist spectral sequences converging to $HFL^{-}(\mathcal{L})$ and $\widehat{HFL}(\mathcal{L})$ respectively [2, 3].

Proposition 2.20. [3, Theorem 1.5.1] For an oriented L-space link $\mathcal{L} \subset S^3$ with n components and $s \in \mathbb{H}(\mathcal{L})$, there exists a spectral sequence with $E_{\infty} = HFL^{-}(\mathcal{L}, s)$ and

$$E_1 = \bigoplus_{B \subset \{1, \cdots, n\}} H_*(A^-(\mathcal{L}, \boldsymbol{s} - \boldsymbol{e}_B)),$$

where the differential in E_1 is induced by inclusions.

Remark 2.21. Precisely, the differential ∂_1 in the E_1 -page is

$$\partial_1(z(\boldsymbol{s}-\boldsymbol{e}_B)) = \sum_{i\in B} U^{H(\boldsymbol{s}-\boldsymbol{e}_B)-H(\boldsymbol{s}-\boldsymbol{e}_B+\boldsymbol{e}_i)} z(\boldsymbol{s}-\boldsymbol{e}_B+\boldsymbol{e}_i),$$

where $z(s - e_B)$ denotes the unique generator in $H_*(A^-(\mathcal{L}, s - e_B))$ with the homological grading $-2H(s - e_B)$.

Proposition 2.22. [2, Proposition 3.8] For an L-space link $\mathcal{L} \subset S^3$ with n components and $s \in \mathbb{H}(\mathcal{L})$, there exists a spectral sequence converging to $\hat{E}_{\infty} = \widehat{HFL}(\mathcal{L}, s)$ with E_1 page

$$\hat{E}_1 = \bigoplus_{B \subset \{1, \cdots, n\}} HFL^-(\mathcal{L}, s + e_B).$$

There is a nice symmetric property of $\widehat{HFL}(\mathcal{L})$ proved by Oszváth and Szabó [15]:

(2.5)
$$\widehat{HFL}_*(\mathcal{L}, \boldsymbol{s}) \cong \widehat{HFL}_*(\mathcal{L}, -\boldsymbol{s}).$$

3. The proof of the Main Theorem

3.1. The Spin^c-cobordism. In this section, we use $\mathcal{L} = L_1 \cup \cdots \cup L_n \subset S^3$ to denote an oriented link with vanishing pairwise linking numbers. Suppose that link components L_i bound pairwise disjoint smoothly embedded surfaces Σ_i of genera g_i in B^4 for all $1 \leq i \leq n$. Attach n 2-handles to the 4-ball B^4 along $L_1, L_2 \cdots, L_n$ with framings $-p_1, -p_2, \cdots, -p_n$. We obtain a 2-handlebody W with boundary $\partial W = S^3_{-p_1, \cdots, -p_n}(\mathcal{L})$ which is the 3-manifold obtained by doing surgery along L_1, L_2, \cdots, L_n with surgery coefficients $-p_1, -p_2, \cdots, -p_n$ respectively. The linking matrix Λ is a diagonal matrix with $\lambda_{ii} = -p_i$. Observe that $\det(\Lambda) \neq 0$, then $S^3_{-p_1, \cdots, -p_n}(\mathcal{L})$ is a rational homology sphere.

Let Σ'_i be the closed surface in W which is the union of Σ_i and the core of the 2handle attached along L_i . Then Σ'_i are also pairwise disjoint. Observe that W is homotopy equivalent to the wedge of n copies of S^2 . Thus, $H_2(W) = \mathbb{Z}^n$ and $[\Sigma'_i]$ are generators of $H_2(W)$. The self intersection number of each Σ'_i in W is $-p_i$.

Take small tubular neighborhoods $nd(\Sigma'_i)$ of Σ'_i such that they are also pairwise disjoint. Then $nd(\Sigma'_i)$ is a disk bundle over Σ'_i and its boundary $\partial(nd(\Sigma'_i))$ is a circle bundle B_{-p_i} with Euler number $-p_i$. The boundary connected sum \mathfrak{D} of the disk bundles over Σ'_i in W is obtained by identifying smoothly embedded balls $B_i^3 \subset B_{-p_i}$ and $B_{i+1}^3 \subset B_{-p_{i+1}}$ for $1 \leq i \leq n-1$, and \mathfrak{D} is also a smooth oriented manifold [5, Section 6.3]. Observe that \mathfrak{D} has the homotopy type of $D_{-p_1} \vee \cdots \vee D_{-p_n}$ where D_{-p_i} denotes the disk bundle over Σ'_i . Since D_{-p_i} is homotopy equivalent to Σ'_i , then

$$\widetilde{H}_i(\mathfrak{D}) \cong \bigoplus_{i=1}^n \widetilde{H}_i(\Sigma'_i).$$

Let X denote the complement of \mathfrak{D} in W. It is a cobordism from $B_{-p_1} \# \cdots \# B_{-p_n}$ to $S^3_{-p_1,\cdots,-p_n}(\mathcal{L})$. Let \bar{X} be the cobordism from $S^3_{-p_1,\cdots,-p_n}(\mathcal{L})$ to $B_{-p_1} \# \cdots \# B_{-p_n}$ obtained by reversing the orientation of X.

Proposition 3.1. For a circle bundle B_{-m} over a closed oriented surface of genus g and Euler number -m < 0, its cohomology is the following:

$$H^1(B_{-m}) \cong \mathbb{Z}^{2g}, \quad H^2(B_{-m}) \cong \mathbb{Z}^{2g} \oplus \mathbb{Z}_m, \quad H^3(B_{-m}) \cong \mathbb{Z}.$$

Proof. For the circle bundle B_{-m} , we have the following long exact sequence by using the Leray sequence:

$$0 \to H^1(\Sigma_g) \to H^1(B_{-m}) \to H^0(\Sigma_g) \xrightarrow{\wedge e} H^2(\Sigma_g) \to H^2(B_{-m}) \to H^1(\Sigma_g) \to 0$$

where e is the Euler class. Then we compute that

$$0 \to \mathbb{Z}^{2g} \to H^1(B_{-m}) \to \mathbb{Z} \xrightarrow{\cdot m} \mathbb{Z} \to H^2(B_{-m}) \to \mathbb{Z}^{2g} \to 0.$$

Thus, $H^1(B_{-m}) \cong \mathbb{Z}^{2g}$ and we have the following short exact sequence:

$$0 \to \mathbb{Z}_m \to H^2(B_{-m}) \to \mathbb{Z}^{2g} \to 0.$$

Since \mathbb{Z}^{2g} is free, the exact sequence splits and $H^2(B_{-m}) \cong \mathbb{Z}^{2g} \oplus \mathbb{Z}_m$. The circle bundle B_{-m} is oriented and closed, so $H^3(B_{-m}) \cong \mathbb{Z}$.

Lemma 3.2. Suppose that M_1 and M_2 are closed, connected and oriented smooth ndimensional manifolds. Then

$$H^i(M_1 \# M_2) \cong H^i(M_1) \oplus H^i(M_2)$$
, for $i \neq 0$ and n_i

and $H^0(M_1 \# M_2) \cong H^n(M_1 \# M_2) \cong \mathbb{Z}$.

Corollary 3.3. The cohomology of $\#_{i=1}^n B_{-p_i}$ is as follows:

$$H^{1}(\#_{i=1}^{n}B_{-p_{i}}) \cong \mathbb{Z}^{2g_{1}+\dots+2g_{n}}, \quad H^{2}(\#_{i=1}^{n}B_{-p_{i}}) \cong \mathbb{Z}^{2g_{1}+\dots+2g_{n}} \oplus \mathbb{Z}_{p_{1}} \dots \oplus \mathbb{Z}_{p_{n}},$$

and $H^{0}(\#_{i=1}^{n}B_{-p_{i}}) \cong H^{3}(\#_{i=1}^{n}B_{-p_{i}}) \cong \mathbb{Z}.$

Proposition 3.4. For the cobordism \overline{X} , its second cohomology has the following form:

$$H^{2}(X) \cong H^{2}(\#_{i=1}^{n}B_{-p_{i}}) \cong \mathbb{Z}^{2g_{1}+\cdots+2g_{n}} \oplus \mathbb{Z}_{p_{1}} \oplus \cdots \oplus \mathbb{Z}_{p_{n}}.$$

Proof. We use the Mayer-Vietoris sequence to compute the cohomology of \bar{X} . Observe that W is the union of \bar{X} and boundary connected sum \mathfrak{D} , and the intersection of \bar{X} and \mathfrak{D} is $\#_{i=1}^{n} B_{-p_{i}}$. Then we have the following long exact sequence:

$$0 \to H^1(W) \to H^1(\bar{X}) \oplus H^1(\Sigma'_1) \dots \oplus H^1(\Sigma'_n) \xrightarrow{i^*} \oplus_{i=1}^n H^1(B_{-p_i}) \to H^2(W)$$

$$\rightarrow \bigoplus_{i=1}^{n} H^2(\Sigma'_i) \oplus H^2(X) \rightarrow H^2(\#_{i=1}^{n} B_{-p_i}) \rightarrow H^3(W) \rightarrow H^3(X) \rightarrow H^3(\#_{i=1}^{n} B_{-p_2}) \rightarrow 0.$$
Becall that W is homotopy equivalent to $S^2 \lor \cdots \lor S^2$. Then $H^1(W) \simeq H^3(W) \in S^2$.

Recall that W is homotopy equivalent to $S^2 \vee \cdots \vee S^2$. Then $H^1(W) \cong H^3(W) \cong H^4(W) \cong 0$. Thus, we have

$$H^{3}(\bar{X}) \cong H^{3}(\#_{i=1}^{n}B_{-p_{i}}) \cong \mathbb{Z}, \quad H^{4}(\bar{X}) = 0,$$

and

$$0 \to H^1(\bar{X}) \oplus \mathbb{Z}^{2g_1} \dots \oplus \mathbb{Z}^{2g_n} \xrightarrow{i^*} \oplus_{i=1}^n \mathbb{Z}^{2g_i} \to \mathbb{Z}^n \to H^2(\bar{X}) \oplus \mathbb{Z}^n \to \oplus_{i=1}^n (\mathbb{Z}^{2g_i} \oplus \mathbb{Z}_{p_i}) \to 0.$$

We claim that $H^1(\bar{X}) = 0$. Let $j^* : H^1(\Sigma'_i) \to H^1(B_{-p_i})$. Observe that $H^1(\Sigma'_i) \cong H_1(\Sigma'_i)$ and $H^1(B_{-p_i}) \cong H_2(B_{-p_i})$ by the Poincare duality. Each generator in $H_1(\Sigma'_i)$ is represented by a simple closed curve in Σ'_i . The curve along with its circle fiber is a generator in $H_2(B_{-p_i})$. Thus $H_1(\Sigma'_i) \cong H_2(B_{-p_i})$. Therefore, j^* is an isomorphism and $H^1(\bar{X}) = 0$. We have the following short exact sequence:

(3.1)
$$0 \to \mathbb{Z}^n \to H^2(\bar{X}) \oplus \mathbb{Z}^n \to \bigoplus_{i=1}^p (\mathbb{Z}^{2g_i} \oplus \mathbb{Z}_{p_i}) \to 0.$$

We claim that $H^2(W) \cong \bigoplus_{i=1}^n H^2(\Sigma'_i)$. Each \mathbb{Z} -summand in $H_2(W, \partial W) \cong H^2(W)$ is represented by the surface Σ_i and it corresponds to the generator of $H^2(\Sigma'_i) \cong H_0(\Sigma'_i)$. Thus $H^2(\bar{X}) \cong H^2(\#_{i=1}^n B_{-p_i}) \cong \bigoplus_{i=1}^p (\mathbb{Z}^{2g_i} \oplus \mathbb{Z}_{p_i})$.

Remark 3.5. From the computation in the proof, $\chi(X) = 2g_1 + \cdots + 2g_n$.

Proposition 3.6. The intersection form \mathcal{Q} : $H^2(\bar{X})/Tor \times H^2(\bar{X})/Tor \to \mathbb{Q}$ vanishes.

Proof. For two elements $s, t \in H^2(\bar{X})/T$ or $\cong H_2(\bar{X}), \mathcal{Q}(s,t) = \langle \bar{s}, \mathrm{PD}(t) \rangle$ where \bar{s} is the image of s under the map $p_* : H_2(\bar{X}) \to H_2(\bar{X}, \partial \bar{X})$ induced by the projection and $\mathrm{PD}(t) \in H^2(\bar{X}, \partial \bar{X})$. By Proposition 3.4, the map $i_* : H_2(\partial \bar{X}) \to H_2(\bar{X})$ induced by the inclusion is surjective. Then $p_* = 0$ by the long exact sequence of homology. Hence, $\bar{s} = 0$ and $\mathcal{Q}(s,t) = 0$. Therefore the intersection form \mathcal{Q} vanishes in \bar{X} .

Corollary 3.7. The signature $\sigma(\bar{X}) = 0$.

By Proposition 3.4, $H^2(\bar{X}) \cong H^2(\#_{i=1}^n B_{-p_i}) \cong \mathbb{Z}^{2g_1 \cdots + 2g_n} \oplus \mathbb{Z}_{p_1} \cdots \oplus \mathbb{Z}_{p_n}$, and the restriction map to $H^2(S^3_{-p_1,\cdots,-p_n}(\mathcal{L}))$ is the projection onto the torsion factors. For $s = (s_1, \cdots, s_n) \in \mathbb{Z}^n/\mathbb{Z}^n \Lambda$, it corresponds to a Spin^c-structure \mathfrak{s} on $S^3_{-p_1,\cdots,-p_n}(\mathcal{L})$ which can be extended to W by Proposition 2.13. We denote its restrictions to \bar{X} and $\#_{i=1}^n B_{-p_i}$ both by \mathfrak{s}' . Moreover, we let s'_i denote the restriction of the Spin^c-structure on \bar{X} to B_{-p_i} . By an argument similar to the one in [20, Lemma 3.1], we have $c_1(s'_i) = 2s_i$.

Rasmussen proved that each circle bundle B_{-p_i} is standard since

$$HF^{\infty}(B_{-p_i}, s'_i) \cong HF^{\infty}(\#^{2g}S^1 \times S^2, \mathfrak{s}_0)$$

where s'_i is a torsion Spin^c-structure on B_{-p_i} and \mathfrak{s}_0 is the unique torsion Spin^c-structure on $\#^{2g}(S^1 \times S^2)$ [20]. Thus $HF^{\infty}(\#^n_{i=1}B_{-p_i},\mathfrak{s}')$ is also standard, and by the additivity property of the *d*-invariants [6, Proposition 4.3],

$$d(\#_{i=1}^{n}B_{-p_{i}},\mathfrak{s}') = d(B_{-p_{1}},s_{1}') + \dots + d(B_{-p_{n}},s_{n}')$$

Next, we can use Proposition 2.11 to prove the following d-invariant inequality:

Proposition 3.8.

$$d(S^{3}_{-p_{1},\cdots,-p_{n}}(\mathcal{L}),\mathfrak{s}) \leq \sum_{i=1}^{n} d(B_{-p_{i}},s'_{i}) + g_{1} + \dots + g_{n}.$$

Proof. The argument is similar to the one in [20, Lemma 3.3]. Consider the map $F_{\bar{X},\mathfrak{s}'}^+$: $HF^+(S^3_{-p_1,\cdots,-p_n}(\mathcal{L}),\mathfrak{s}) \to HF^+(\#^n_{i=1}B_{-p_i},\mathfrak{s}')$. This map is U-equivariant and agrees with $F^{\infty}_{\bar{X},\mathfrak{s}'}$ in high degrees, which implies that it takes $\pi(HF^{\infty}(S^3_{-p_1,\cdots,-p_n}(\mathcal{L}),\mathfrak{s}))$ onto $\pi(\mathfrak{A}_{\mathfrak{s}'})$ by Proposition 2.11. Thus if $1 \in \pi(HF^{\infty}(S^3_{-p_1,\cdots,-p_n}(\mathcal{L}),\mathfrak{s}) \otimes \mathbb{F} \cong \mathbb{F}[U^{-1}]$ is the element with the lowest absolute grading, we must have

$$gr(F_{\bar{X},\mathfrak{s}'}^+(1)) \le d(\#_{i=1}^n B_{-p_i},\mathfrak{s}').$$

Observe that $gr(1) = d(S^3_{-p_1,\dots,-p_n}(\mathcal{L}),\mathfrak{s})$ and $F^+_{\bar{X},\mathfrak{s}'}$ shifts the absolute grading by

$$degF_{\bar{X},\mathfrak{s}'}^+ = \frac{c_1^2(\mathfrak{s}') - 2\chi(X) - 3\sigma(X)}{4} = \frac{0 - 2(2g_1 + \dots + 2g_n) - 3\cdot 0}{4} = -g_1 - \dots - g_n$$

where $c_1^2(\mathfrak{s}') = \mathcal{Q}(c_1(\mathfrak{s}'), c_1(\mathfrak{s}')) = 0$. Thus,

$$d(S^{3}_{-p_{1},\cdots,-p_{n}}(\mathcal{L}),\mathfrak{s}) \leq \sum_{i=1}^{n} d(B_{-p_{i}},s'_{i}) + g_{1} + \dots + g_{n}.$$

Let \mathcal{L}^* denote the mirror of \mathcal{L} . Observe that $S^3_{-p_1,\cdots,-p_n}(\mathcal{L})$ is obtained from $S^3_{p_1,\cdots,p_n}(\mathcal{L}^*)$ by reversing the orientation. For any $s \in \mathbb{Z}^n$, choose sufficiently large $p_i \gg 0$ so that $s \in P_{\mathbb{H}}(\Lambda)$. Let \mathfrak{s} denote the Spin^c-structure on $S^3_{-p_1,\cdots,-p_n}(\mathcal{L})$ corresponding to s. By Theorem 2.14

$$d(S^3_{-p_1,\cdots,-p_n}(\mathcal{L}),\mathfrak{s}) = -d(S^3_{p_1,\cdots,p_n}(\mathcal{L}^*),\mathfrak{s}) = \deg F_{\mathcal{U}',\mathfrak{s}} + 2H_{L^*}(\boldsymbol{s})$$

where $H_{\mathcal{L}^*}$ is the *H*-function of \mathcal{L}^* . Let *O* denote the unlink with *n* components. Similarly, we have

$$d(S^3_{-p_1,\cdots,-p_n}(O),\mathfrak{s}) = -d(S^3_{p_1,\cdots,p_n}(O),\mathfrak{s}) = \deg F_{\mathcal{U}',\mathfrak{s}} + 2H_O(\mathbf{s})$$

Thus,

$$d(S^{3}_{-p_{1},\cdots,-p_{n}}(\mathcal{L}),\mathfrak{s}) - d(S^{3}_{-p_{1},\cdots,-p_{n}}(O),\mathfrak{s}) = 2H_{L^{*}}(s) - 2H_{O}(s) = 2h_{\mathcal{L}^{*}}(s).$$

Recall that for a circle bundle B_{-m} with Euler number -m over a closed, oriented genus g surface, $H^2(B_{-m}) \cong \mathbb{Z}^{2g} \oplus \mathbb{Z}_m$. There is a natural way to label the torsion Spin^c-structures on B_{-m} . Let \mathfrak{t}_k denote the torsion Spin^c-structure on B_{-m} such that $c_1(\mathfrak{t}_k) = 2k \in \mathbb{Z}_m$ where $-m/2 \leq k \leq m/2$.

Proposition 3.9. [20, Proposition 3.4] Let B_{-m} denote a circle bundle equipped with a torsion Spin^c-structure \mathfrak{t}_k over a closed oriented surface Σ_q . For $m \gg 0$,

$$d(B_{-m}, \mathfrak{t}_k) = \begin{cases} E(m, k) - g + 2 \left\lceil \frac{g - |k|}{2} \right\rceil & |k| \le g \\ E(m, k) - g & |k| > g \end{cases}$$

where $\{E(m,k) \mid k \in \mathbb{Z}_m\}$ is the set of d-invariants of the lens space L(m,1).

Proof. The circle bundle B_{-m} can be obtained by doing -m-surgery on "the Borromean knot" $B \subset \#^{2g}(S^1 \times S^2)$. For the large surgery, $HF(B_{-m}, \mathfrak{t}_k)$ can be obtained from the knot Floer homology $HFK^{\infty}(B)$. Ozsváth and Szabó proved that [15]

$$\widehat{H}F\widetilde{K}(B,i) \cong \wedge^{g+i}(H^1(\Sigma_g)) \quad \text{ if } |i| \le g.$$

Otherwise, $\widehat{HFK}(B,i) = 0$. The rest of the proof can be found in [20, Proposition 3.4].

3.2. **Proofs of main theorems.** We prove Theorem 1.1 and Theorem 1.3 in this subsection.

Proof of Theorem 1.1: By Proposition 3.8 and Proposition 3.9,

$$d(S^{3}_{-p_{1},\cdots,-p_{n}}(\mathcal{L}),\mathfrak{s}) \leq \sum_{i=1}^{n} (E(p_{i},s_{i}) - g_{i} + 2f_{g_{i}}(s_{i})) + g_{1} + \dots + g_{n}.$$

Recall that $d(S^3_{-p_1,\dots,-p_n}(\mathcal{L}),\mathfrak{s}) = 2h_{\mathcal{L}^*}(\mathfrak{s}) + d(S^3_{-p_1,\dots,-p_n}(O),\mathfrak{s})$. For lens spaces, our orientation convention is the one used in [20], namely, that -p surgery on the unknot produces the oriented space L(p,1). Then $S^3_{-p_1,\dots,-p_n}(O) = L(p_1,1)\#\cdots\#L(p_n,1)$, and

$$d(S^{3}_{-p_{1},\cdots,-p_{n}}(O),\mathfrak{s}) = \sum_{i=1}^{n} d(L(p_{i},1),s_{i}) = \sum_{i=1}^{n} E(p_{i},s_{i}).$$

Hence,

$$h_{\mathcal{L}^*} \le \sum_{i=1}^n f_{g_i}(s_i).$$

The surfaces Σ_i bounded by L_i are pairwise disjoint. Then the corresponding link components of the mirror link \mathcal{L}^* bound the mirrors of Σ_i which have the same genera as Σ_i . Thus we have $h_L(\mathbf{s}) \leq \sum_{i=1}^n f_{g_i}(s_i)$.

Corollary 3.10. For an oriented n-component link $\mathcal{L} \subset S^3$, $\mathfrak{G}(\mathcal{L}) \subset \mathfrak{G}_{HF}(\mathcal{L})$.

Proof. Suppose that the link components of \mathcal{L} bound pairwise disjoint surfaces in B^4 of genera g_i . By Theorem 1.1, $h_{\mathcal{L}}(s) = 0$ if $s \succeq g$ where $g = (g_1, \dots, g_n)$.

Definition 3.11. An oriented *n*-component link $\mathcal{L} \subset S^3$ is (smoothly) *slice* if there exist *n* disjoint, smoothly embedded disks in B^4 with boundary \mathcal{L} .

Proof of Theorem 1.3: If \mathcal{L} is slice, then $h_{\mathcal{L}} = 0$ by Theorem 1.1. Thus $H_{\mathcal{L}}(\boldsymbol{v}) = H_O(\boldsymbol{v}) = \sum_{i=1}^n H(v_i)$ where $H(v_i)$ is the *H*-function for the unknot and $\boldsymbol{v} = (v_1, \dots, v_n) \in \mathbb{Z}^n$. We claim that $HFL^-(\mathcal{L}, \boldsymbol{v}) = 0$ if there exists a component $v_j > 0$. By Proposition 2.20, there exists a spectral sequence converging to $HFL^-(\mathcal{L})$ with the E_1 -page

$$E_1(\boldsymbol{v}) = \bigoplus_{B \subset \{1, \cdots, n\}} H_*(A^-(\mathcal{L}, \boldsymbol{v} - \boldsymbol{e}_B)).$$

Let $\mathcal{K} = \{1, \cdots, n\} \setminus \{j\}$, and

$$E'(\boldsymbol{v}) = \bigoplus_{B \subset \mathcal{K}} H_*(A^-(\mathcal{L}, \boldsymbol{v} - \boldsymbol{e}_B)), \quad E''(\boldsymbol{v}) = \bigoplus_{B \subset \mathcal{K}} H_*(A^-(\mathcal{L}, \boldsymbol{v} - \boldsymbol{e}_B - \boldsymbol{e}_j)).$$

Then $E_1(\boldsymbol{v}) = E'(\boldsymbol{v}) \oplus E''(\boldsymbol{v})$. Recall that for each $B \subset \{1, \dots, n\}$, $H_*(A^-(\mathcal{L}, \boldsymbol{v} - \boldsymbol{e}_B)) \cong \mathbb{F}[U]$, [7]. Let $\partial_1, \partial', \partial''$ denote the differentials in $E_1(\boldsymbol{v}), E'(\boldsymbol{v})$ and $E''(\boldsymbol{v})$, respectively. Let z denote the generator of $H_*(A^-(\mathcal{L}, \boldsymbol{v} - \boldsymbol{e}_B - \boldsymbol{e}_j)) \in E''(\boldsymbol{v})$ with homological grading $-2H(\boldsymbol{v} - \boldsymbol{e}_B - \boldsymbol{e}_j)$. Observe that $H(\boldsymbol{v} - \boldsymbol{e}_B - \boldsymbol{e}_j) = H(\boldsymbol{v} - \boldsymbol{e}_B)$ since $H(v_j - 1) = H(v_j)$ for $v_j > 0$. Then $\partial_1(z) = \partial''(z) + z'$ where z' is the generator of $H_*(A^-(\mathcal{L}, \boldsymbol{v} - \boldsymbol{e}_B))$ with homological grading $-2H(\boldsymbol{v} - \boldsymbol{e}_B)$. Let \mathcal{D} be an acyclic chain complex with two generators a and b, and the differential $\partial_D(a) = b$. Then the chain complex $(E_1(\boldsymbol{v}), \partial_1)$ is isomorphic to $(E'(\boldsymbol{v}) \otimes \mathcal{D}, \partial_1 \otimes \partial_D)$. Thus $E_2 = 0$, and the spectral sequence collapes at E_2 . Therefore, $HFL^-(\mathcal{L}, \boldsymbol{v}) = 0$ if there exists $v_j > 0$.

We also have $\widehat{HFL}(\mathcal{L}, \boldsymbol{v}) = 0$ if there exists $v_j > 0$ by the spectral sequence in Proposition 2.22. By the symmetric property [15], $\widehat{HFL}(\mathcal{L}, -\boldsymbol{v}) = \widehat{HFL}(\mathcal{L}, \boldsymbol{v}) = 0$. Hence, $\widehat{HFL}(\mathcal{L}, \boldsymbol{v}) = 0$ if $\boldsymbol{v} \neq \boldsymbol{0}$. The dual Thurston polytope of \mathcal{L} is a point at the origin [17, Theorem 1.1]. Thus, the link \mathcal{L} bounds disjoint disks in S^3 , and \mathcal{L} is an unlink. \Box

3.3. Lower bounds for the 4-genera. In this subsection, we use $\mathcal{L} \subset S^3$ to denote an *n*-component link with vanishing pairwise linking numbers. The inequality in Theorem 1.1 produces some lower bounds for the 4-genus of \mathcal{L} .

Corollary 3.12. For the link \mathcal{L} ,

(3.2)
$$g_4(\mathcal{L}) \ge \min\{s_1 + \dots + s_2 \mid h(\boldsymbol{x}) = 0 \text{ if } \boldsymbol{x} \succeq \boldsymbol{s} = (s_1, \dots, s_n)\}.$$

Proof. This is straightforward from Corollary 3.10

Corollary 3.13. For the link \mathcal{L} , $g_4(\mathcal{L}) \geq 2 \max_{s \in \mathbb{Z}^n} h_{\mathcal{L}}(s) - n$. In particular,

(3.3)
$$g_4(\mathcal{L}) \ge 2h_{\mathcal{L}}(\mathbf{0}) - n$$

Proof. By Theorem 1.1, for all $s \in \mathbb{Z}^n$, $h_{\mathcal{L}}(s) \leq \lceil g_1/2 \rceil + \cdots + \lceil g_n/2 \rceil$. Observe that $\lceil g_i/2 \rceil \leq (g_i+1)/2$. Then

$$g_1 + \dots + g_n + n \ge 2 \max_{\boldsymbol{s} \in \mathbb{Z}^n} h_{\mathcal{L}}(\boldsymbol{s}).$$

Hence $g_4(\mathcal{L}) \geq 2 \max_{\boldsymbol{s} \in \mathbb{Z}^n} h_{\mathcal{L}}(\boldsymbol{s}) - n.$

Corollary 3.14. Let $g_4(L_i)$ denote the 4-genus of the link component L_i . Then

(3.4)
$$g_4(\mathcal{L}) \ge 2h_{\mathcal{L}}(s) - n + |s_1| + \dots + |s_2|,$$

where $s = (s_1, \dots, s_n)$, and $|s_i| \le g_4(L_i)$.

Proof. Suppose that \mathcal{L} bound pairwise disjoint surfaces Σ_i in B^4 of genera g_i . Then $g_i \geq g_4(L_i)$ for all *i*. If $|s_i| \leq g_4(L_i)$, then by Theorem 1.1,

$$h_{\mathcal{L}}(\boldsymbol{s}) \leq \sum_{i=1}^{n} \left\lceil \frac{g_i - |s_i|}{2} \right\rceil.$$

Since $\lceil (g_i - |s_i|)/2 \rceil \le (g_i - |s_i| + 1)/2$, we have $g_1 + \dots + g_n \ge 2h_{\mathcal{L}}(s) - n + |s_1| + \dots + |s_2|$. Hence, $g_4(\mathcal{L}) \ge 2h_{\mathcal{L}}(s) - n + |s_1| + \dots + |s_2|$.

For the rest of the subsection, we prove that the analogues of Lemma 2.4, Lemma 2.5 and Lemma 2.7 hold for the set $\mathfrak{G}(\mathcal{L})$. For an oriented link \mathcal{L} with vanishing pairwise linking numbers, we use the *cancellation process* to find pairwise disjoint surfaces in B^4 bounded by \mathcal{L} . Let $\Sigma_i \subset S^3$ denote a Seifert surface bounded by L_i . Then Σ_i and Σ_j intersect transversely at even number of points in B^4 since the linking number equals 0. We remove the tubular neighborhoods of a positive crossing and a negative crossing in Σ_i and obtain a new surface with two punctures. Add a tube along an arc in Σ_j which connects the two intersection points to the punctured surface where the attaching circles are boundaries of these two punctures, as in Figure 2. Then we obtain a new surface Σ'_i with fewer intersection points with Σ_j and higher genus compared with Σ_i . The tube can also be attached to the surface Σ_j along an arc connecting the intersection points in Σ_i . We repeat the process until we get pairwise disjoint surfaces in B^4 bounded by \mathcal{L} . We call the process of adding tubes to eliminate intersection points as *cancellation process*.



FIGURE 2. Cancellation process

Lemma 3.15. If $g \in \mathfrak{G}(\mathcal{L})$ and $y \succeq g$, then $y \in \mathfrak{G}(\mathcal{L})$. Equivalently, if $g \notin \mathfrak{G}(\mathcal{L})$ and $y \preceq g$, then $y \notin \mathfrak{G}(\mathcal{L})$.

Proof. If $\boldsymbol{g} = (g_1, \dots, g_n) \in \mathfrak{G}(\mathcal{L})$, there exist pairwise disjoint surfaces Σ_i embedded in B^4 of genera g_i and $\partial \Sigma_i = L_i$. We can attach tubes to the surfaces Σ_i to increase the genera. Thus $\boldsymbol{y} \in \mathfrak{G}(\mathcal{L})$ if $\boldsymbol{y} \succeq \boldsymbol{g}$.

Lemma 3.16. If $\boldsymbol{g} = (g_1, \dots, g_n) \in \mathfrak{G}(\mathcal{L})$, then $\boldsymbol{g} \setminus g_i \in \mathfrak{G}(\mathcal{L} \setminus L_i)$ for all $1 \leq i \leq n$. Moreover, if $\boldsymbol{g} \setminus g_i \in \mathfrak{G}(\mathcal{L} \setminus L_i)$, then for g_i sufficiently large, $\boldsymbol{g} = (g_1, \dots, g_n) \in \mathfrak{G}(\mathcal{L})$.

Proof. If $\mathbf{g} \in \mathfrak{G}(\mathcal{L})$, it is easy to obtain that $\mathbf{g} \setminus g_i \in \mathfrak{G}(\mathcal{L} \setminus L_i)$. Conversely, if $\mathbf{g} \setminus g_i \in \mathfrak{G}(\mathcal{L} \setminus L_i)$, for sufficiently large $g_i \gg 0$, we claim that $\mathbf{g} \in \mathfrak{G}(\mathcal{L})$. Suppose that $\mathcal{L} \setminus L_i$ bounds pairwise disjoint surfaces Σ_j in B^4 . Let Σ_i in S^3 denote a Seifert surface bounded by L_i . Then Σ_i intersects with Σ_j transversely at even number of points in B^4 since the linking number equals 0. By the cancellation process, we add tubes to Σ_i until the new surface is disjoint from all the surfaces Σ_j . Thus for sufficiently large $g_i, \mathbf{g} \in \mathfrak{G}(\mathcal{L})$.

Lemma 3.17. The set $\mathfrak{G}(\mathcal{L})$ is determined by the set of maximal lattice points and $\mathfrak{G}(\mathcal{L} \setminus L_i)$ for all $1 \leq i \leq n$.

Proof. The proof is similar to the one in Lemma 2.7 by using Lemma 3.15 and Lemma 3.16. $\hfill \Box$

4. Examples

4.1. **Examples.** For *L*-space links, the *H*-function can be computed explicitly by the Alexander polynomials of the link and sublinks. The lower bounds for 4-genus of the link in Section 3 can also be computed explicitly. In this section, we will show examples of *L*-space links where $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L})$.

Example 4.1. Let k be a positive integer. The two bridge link $\mathcal{L}_k = b(4k^2 + 4k, -2k - 1)$ is a 2-component L-space link with linking number 0 [7], and both link components are unknots, see Figure 3. The Alexander polynomial of \mathcal{L}_k is computed in [7, Section 3],

$$\Delta_{\mathcal{L}_k}(t_1, t_2) = (-1)^k \sum_{|i+1/2| + |j+1/2| \le k} (-1)^{i+j} t_1^{i+1/2} t_2^{j+1/2}.$$



FIGURE 3. Two bridge link $b(4k^2 + 4k, -2k - 1)$.

The *H*-function of \mathcal{L}_k is computed in [7, Proposition 6.12]. Then the *h*-function of \mathcal{L}_k is shown in Figure 4, where h(k, 0) = 0, and h(k - 1, 0) = 1. For the shaded area bounded by the "stairs", the *h*-function is nonzero, and h(s) = 0 for all lattice points *s* on the "stairs"

and outside of the shaded area in Figure 4. Thus, $\mathfrak{G}_{HF}(\mathcal{L})$ consists of all the lattice points on the "stairs" and outside the shaded area in the first quadrant. By the inequality (3.3),

$$g_4(\mathcal{L}_k) \ge \min\{s_1 + s_2 \mid h(x) = 0 \text{ if } x \succeq s = (s_1, s_2)\} = k.$$

Observe that the components of \mathcal{L}_k bound disks D_1 and D_2 in S^3 . Push the disks into B^4 . Then they intersect transversely at 2k points in B^4 . By the cancellation process of crossings, we obtain disjoint surfaces Σ'_1 and Σ'_2 in B^4 bounded by the link components. Assume that the genus of Σ'_1 is k and Σ'_2 is still a disk of genus 0. Then $g_4(\mathcal{L}_k) \leq k$. Thus, $g_4(\mathcal{L}_k) = k$. We can add tubes to either D_1 or D_2 in the cancellation process. Thus, for all $\mathbf{g} = (g_1, g_2)$ with $g_1 + g_2 = k$, we find disjoint surfaces in B^4 of genera g_1, g_2 , respectively. Therefore, $\mathfrak{G}(\mathcal{L}_k) = \mathfrak{G}_{HF}(\mathcal{L}_k)$.



FIGURE 4. The *h*-function of \mathcal{L}_k .

Remark 4.2. For k = 1 we get the Whitehead link \mathcal{L}_1 , and the 4-genus $g_4(\mathcal{L}_1) = 1$.

Example 4.3. The Borromean link $\mathcal{L} = L_1 \cup L_2 \cup L_3$ is a 3-component *L*-space link with vanishing pairwise linking numbers [7]. Its Alexander polynomial equals

$$\Delta_{\mathcal{L}}(t_1, t_2, t_3) = (t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2})(t_3^{1/2} - t_3^{-1/2}).$$

By Equation (2.4), if $\boldsymbol{v} \succ \boldsymbol{0}$, $h_{\mathcal{L}}(\boldsymbol{v}) = 0$ and $h_{\mathcal{L}}(\boldsymbol{0}) = 1$. Thus, $\mathfrak{G}_{HF}(\mathcal{L}) = \{\boldsymbol{v} \in \mathbb{Z}^n \mid \boldsymbol{v} \succ \boldsymbol{0}\}.$

Each component L_i is an unknot and bounds a disk D_i in B^4 . The disks D_1 and D_2 intersect transversely at two points in $B^4 \setminus L_3$. By the cancellation process, L_1 and L_2 bound disjoint surfaces D'_1 and D_2 of genera 1 and 0 respectively in $B^4 \setminus L_3$. The disk D_3 intersects D'_1 and D_2 transversely in B^4 . Since D'_1 and D_2 are disjoint, by an isotopy in B^4 , we can make D_3 disjoint from D'_1 and D_2 . Thus, $g_4(\mathcal{L}) = 1$, and $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L})$.

Example 4.4. The mirror of L7a3 is a 2-component L-space link $\mathcal{L} = L_1 \cup L_2$ with linking number 0, where L_1 is the right-handed trefoil and L_2 is the unknot [7]. Its Alexander polynomial equals

$$\Delta_{\mathcal{L}}(t_1, t_2) = -(t_1^{1/2} - t_1^{-1/2})(t_2^{1/2} - t_2^{-1/2})(t_2 + t_2^{-1}).$$

The *h*-function in the first quadrant is shown as in Figure 5 by Equation (2.4) or the formula in [7]. Then the shaded area is $\mathfrak{G}_{HF}(\mathcal{L})$ and $g_4(\mathcal{L}) \geq 2$.

Observe that the right-handed trefoil and the unknot bound Seifert surfaces of genera 1 and 0 respectively in S^3 . They intersect transversely at two points in B^4 . By the cancellation process, we can obtain disjoint surfaces of genera (2,0) or (1,1) bounded by the link. Thus, $g_4(\mathcal{L}) = 2$, and $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L})$.



FIGURE 5. The h-function for the mirror of L7a3

Example 4.5. Let \mathcal{L} denote the disjoint union of two links \mathcal{L}_1 and \mathcal{L}_2 . Then $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}(\mathcal{L}_1) \times \mathfrak{G}(\mathcal{L}_2)$ and $\mathfrak{G}_{HF}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L}_1) \times \mathfrak{G}_{HF}(\mathcal{L}_2)$.

Proof. Suppose that \mathcal{L}_1 has n_1 components and \mathcal{L}_2 has n_2 components. If $\mathbf{g}_1 \in \mathfrak{G}(\mathcal{L}_1)$ and $\mathbf{g}_2 \in \mathfrak{G}(\mathcal{L}_2)$, then $(\mathbf{g}_1, \mathbf{g}_2) \in \mathfrak{G}(\mathcal{L})$ where $\mathcal{L} = \mathcal{L}_1 \sqcup \mathcal{L}_2$. Conversely, if

$$\boldsymbol{g} = (g_1, \cdots, g_{n_1}, \cdots, g_{n_1+n_2}) \in \mathfrak{G}(\mathcal{L}),$$

it is straightforward to obtain that $(g_1, \dots, g_{n_1}) \in \mathfrak{G}(\mathcal{L}_1)$ and $(g_{n_1+1}, \dots, g_{n_1+n_2}) \in \mathfrak{G}(\mathcal{L}_2)$. Thus, $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}(\mathcal{L}_1) \times \mathfrak{G}(\mathcal{L}_2)$.

For the set $\mathfrak{G}_{HF}(\mathcal{L})$, recall that $H_{\mathcal{L}}(\mathbf{s}) = H_{\mathcal{L}_1}(\mathbf{s}_1) + H_{\mathcal{L}_2}(\mathbf{s}_2)$ [1, Proposition 3.11] where $\mathbf{s} = (s_1, \cdots, s_{n_1}, \cdots, s_{n_1+s_{n_2}})$, $\mathbf{s_1} = (s_1, \cdots, s_{n_1})$ and $\mathbf{s_2} = (s_{n_1+1}, \cdots, s_{n_1+n_2})$. Since H-functions take nonnegative values, $H_{\mathcal{L}}(\mathbf{s}) = 0$ if and only if $H_{\mathcal{L}_1}(\mathbf{s}_1) = H_{\mathcal{L}_2}(\mathbf{s}_2) = 0$. Thus, $\mathfrak{G}_{HF}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L}_1) \times \mathfrak{G}_{HF}(\mathcal{L}_2)$.

4.2. Cables of *L*-space links. Let $\mathcal{L} = L_1 \cup \cdots \cup L_n \subset S^3$ be an *L*-space link with vanishing pairwise linking numbers. For each $1 \leq i \leq n$, let p_i and q_i be coprime positive integers. Consider the link $\mathcal{L}_{cab} = L_{(p_1,q_1)} \cup \cdots \cup L_{(p_n,q_n)}$ where $L_{(p_i,q_i)}$ is the (p_i,q_i) -cable on L_i . If for all $i, p_i/q_i$ are sufficiently large, then \mathcal{L}_{cab} is also an *L*-space link [1, Proposition 2.8].

Theorem 4.6. For such cable links \mathcal{L}_{cab} ,

 $\mathfrak{G}_{HF}(\mathcal{L}_{cab}) = \{ \boldsymbol{u} \in \mathbb{Z}^n \mid \boldsymbol{u} \succeq T(\boldsymbol{s}) \text{ for some } \boldsymbol{s} \in \mathfrak{G}_{HF}(\mathcal{L}) \}$

where $T: \mathbb{Z}^n \to \mathbb{Z}^n$ is defined as

$$T(s) = \mathbf{p} \cdot \mathbf{s} + ((p_1 - 1) (q_1 - 1) / 2, \cdots, (p_n - 1) (q_n - 1) / 2),$$

$$\mathbf{p} = (p_1, \cdots, p_n), \mathbf{s} = \{s_1, \cdots, s_n\} \text{ and } \mathbf{p} \cdot \mathbf{s} = p_1 s_1 + \cdots + p_n s_n.$$

Proof. By Lemma 2.19, $\mathfrak{G}_{HF}(\mathcal{L})$ is determined by Alexander polynomials of the link and sublinks. Let $\Delta_{\mathcal{L}}(t_1, \dots, t_n)$ denote the Alexander polynomial of \mathcal{L} . Then the Alexander polynomial of the cable link \mathcal{L}_{cab} is computed by Turaev in [21, Theorem 1.3.1],

(4.1)
$$\Delta_{\mathcal{L}_{cab}}(t_1,\cdots,t_n) = \Delta_{\mathcal{L}}(t_1^{p_1},\cdots,t_n^{p_n}) \prod_{i=1}^n \frac{t_i^{p_i q_i/2} - t_i^{-p_i q_i/2}}{t_i^{q_i/2} - t_i^{-q_i/2}}$$

If \mathcal{L} is a knot, we should replace the Alexander polynomials by $\Delta_{\mathcal{L}}(t)/(t-1)$. Then

(4.2)
$$\tilde{\Delta}_{\mathcal{L}_{cab}}(t_1,\cdots,t_n) = t_1^{1/2-p_1/2}\cdots t_n^{1/2-p_n/2}\tilde{\Delta}_{\mathcal{L}}(t_1^{p_1},\cdots,t_n^{p_n})\prod_{i=1}^n \frac{t_i^{p_iq_i/2}-t_i^{-p_iq_i/2}}{t_i^{q_i/2}-t_i^{-q_i/2}}$$

Observe that $T(\mathbf{s}) = \mathbf{p} \cdot \mathbf{s} + (1/2 - p_1/2, \cdots, 1/2 - p_n/2) + ((p_1q_1 - q_1)/2, \cdots, (p_nq_n - q_n)/2)$ for any $\mathbf{s} \in \mathbb{Z}^n$. We claim that the coefficients of $t_1^{y_1} \cdots t_n^{y_n}$ in $\tilde{\Delta}_{\mathcal{L}}(t_1, \cdots, t_n)$ are 0 for all $\mathbf{y} \succ \mathbf{s}$ if and only if for all $\mathbf{y}' \succ T(\mathbf{s})$, the coefficients of $t_1^{y_1'} \cdots t_n^{y_n'}$ are 0 in $\tilde{\Delta}_{\mathcal{L}_{cab}}(t_1, \cdots, t_n)$. By Equation (4.2), the coefficient of $t_1^{y_1'} \cdots t_n^{y_n'}$ is 0 if \mathbf{y}' is not in the image of T and $\mathbf{y}' \succ T(\mathbf{s})$. If $\mathbf{y}' = T(\mathbf{y})$ for some $\mathbf{y} \succ \mathbf{s}$, then the coefficient of $t_1^{y_1'} \cdots t_n^{y_n'}$ equals the coefficient of $t_1^{y_1'} \cdots t_n^{y_n'}$ in $\tilde{\Delta}_{\mathcal{L}}(t_1, \cdots, t_n)$ which is 0. This proves the "only if " part. For the "if " part, suppose there exists $\mathbf{y} \succ \mathbf{s}$ such that the coefficient of $t_1^{y_1} \cdots t_n^{y_n}$ is nonzero. Then the coefficient corresponding to $T(\mathbf{y})$ in $\tilde{\Delta}_{\mathcal{L}_{cab}}(t_1, \cdots, t_n)$ is 0 which contradicts to our assumption. For all subsets $B \subset \{1, \cdots, n\}$, the similar statement holds for Alexander polynomials $\tilde{\Delta}_{\mathcal{L}\setminus L_B}(t_{i_1}, \cdots, t_{i_k})$ and $\tilde{\Delta}_{\mathcal{L}_{cab}\setminus (\mathcal{L}_{cab})_B}(t_{i_1}, \cdots, t_{i_k})$. By Lemma 2.19, $\mathbf{y}' \in \mathfrak{G}_{HF}(\mathcal{L}_{cab})$ if $\mathbf{y}' \succ T(\mathbf{s})$ for some $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$. Thus, $\mathfrak{G}_{HF}(\mathcal{L}_{cab}) \supset \{\mathbf{u} \in \mathbb{Z}^n \mid \mathbf{u} \succeq T(\mathbf{s})$ for some $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$.

Conversely, suppose $\mathbf{y}' \in \mathfrak{G}_{HF}(\mathcal{L}_{cab})$. If $\mathbf{y}' = T(\mathbf{s})$ for some $\mathbf{s} \in \mathbb{Z}^n$, by Lemma 2.19 and the claim, $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$. If \mathbf{y}' is not in the image of T, then there exists $\mathbf{s} \in \mathbb{Z}^n$ such that $\mathbf{y}' \succ T(\mathbf{s})$ and $\mathbf{y}' \prec T(\mathbf{y})$ for all $\mathbf{y} \succ \mathbf{s}$. We claim that $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$. If there exists $\mathbf{y} \succ \mathbf{s}$ such that the coefficient corresponding to \mathbf{y} in $\tilde{\Delta}_{\mathcal{L}}(t_1, \cdots, t_n)$ is not 0, then the coefficient corresponding to $T(\mathbf{y})$ in $\tilde{\Delta}_{\mathcal{L}_{cab}}(t_1, \cdots, t_n)$ is also not 0 which contradicts to our assumption. Similarly, we prove that for all subsets $B \subset \{1, \cdots, n\}$ and all $\mathbf{y} \succ \mathbf{s}$, the coefficients corresponding to $\mathbf{y} \setminus y_B$ in $\tilde{\Delta}_{\mathcal{L} \setminus L_B}(t_{i_1}, \cdots, t_{i_k})$ are all 0. By Lemma 2.19, $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})$. Thus, $\mathfrak{G}_{HF}(\mathcal{L}_{cab}) = \{\mathbf{u} \in \mathbb{Z}^n \mid \mathbf{u} \succeq T(\mathbf{s})$ for some $\mathbf{s} \in \mathfrak{G}_{HF}(\mathcal{L})\}$.

$$\square$$

Lemma 4.7. For such cable links \mathcal{L}_{cab} , $\mathfrak{G}(\mathcal{L}_{cab}) \supset \{ u \in \mathbb{Z}^n \mid u \succeq T(g) \text{ for some } g \in \mathfrak{G}(\mathcal{L}) \}$.

Proof. Suppose that the link components in \mathcal{L} bound pairwise disjoint surfaces Σ_i in B^4 of genera g_i . The cable knot $L_{(p_i,q_i)}$ bounds a surface of genus $p_ig_i + (p_i - 1)(q_i - 1)/2$. We start with p_i copies of Σ_i and use $(p_i - 1)(q_i - 1)$ half-twisted bands to connect them. Since Σ_i are pairwise disjoint, the new surfaces are also pairwise disjoint.

Proof of Proposition 1.4: Let $\mathfrak{G}' = \{ \boldsymbol{u} \in \mathbb{Z}^n \mid \boldsymbol{u} \succeq T(\boldsymbol{g}) \text{ for some } \boldsymbol{g} \in \mathfrak{G}(\mathcal{L}) \}$. By assumption, $\mathfrak{G}(\mathcal{L}) = \mathfrak{G}_{HF}(\mathcal{L})$. Then $\mathfrak{G}' = \mathfrak{G}_{HF}(\mathcal{L}_{cab})$ by Theorem 4.6. Since $\mathfrak{G}_{HF}(\mathcal{L}_{cab}) \supset \mathfrak{G}(\mathcal{L}_{cab}) \supset \mathfrak{G}'$ by Lemma 4.7, we have $\mathfrak{G}_{HF}(\mathcal{L}_{cab}) = \mathfrak{G}(\mathcal{L}_{cab})$.

Remark 4.8. Proposition 1.4 also holds if we only replace some link components in \mathcal{L} by their cables.

By Proposition 1.4, we can apply cables on all L-space links in Examples of Subsection 4.1.

Example 4.9. Cables on the Whitehead link.

Let $Wh_{p,q}$ denote the link consisting of the (p,q)-cable on one component of the Whitehead link and the unchanged second component. The linking number is 0, and $Wh_{p,q}$ is an *L*-space link if p, q are coprime with $q/p \ge 3$, [1].

By Proposition 1.4, the *h*-function of $Wh_{p,q}$ in the first quadrant is shown as in Figure 6. The shaded area is $\mathfrak{G}_{HF}(Wh_{p,q}) = \mathfrak{G}(Wh_{p,q})$. Thus $g_4(Wh_{p,q}) = g_1 + g_2 = (p-1)(q-1)/2 + 1$. The link $Wh_{p,q}$ bounds disjoint surfaces of genera (p-1)(q-1)/2 and 1 as in Figure 6. The red line denotes the tube attached to the disk bounded by the unknot.



FIGURE 6. The h-function of $Wh_{p,q}$

Example 4.10. For the 2-bridge link $\mathcal{L}_k = b(4k^2 + 4k, -2k - 1) = L_1 \cup L_2$, consider the cable link $\mathcal{L}_{cab} = L_{(p_1,q_1)} \cup L_{(p_2,q_2)}$ where p_i, q_i are coprime positive integers with q_i/p_i sufficiently large. By proposition 1.4, $\mathfrak{G}_{HF}(\mathcal{L}_{cab}) = \mathfrak{G}(\mathcal{L}_{cab})$ is shown as in Figure 7 where all the horizontal segments in the "stair" have length p_1 , vertical segments have length p_2 , and there are k steps.



FIGURE 7. The set $\mathfrak{G}(\mathcal{L}_{cab})$ for the cable link

References

 M.Borodzik, E.Gorsky, Immersed concordances of links and Heegaard Floer homology, preprint 2016, arXiv: 1601.07507v1.

- [2] E. Gorsky, J. Hom, Cable links and L-space surgeries, Quantum Topol. 8 (2017), no. 4, 629-666.
- [3] E. Gorsky, A. Némethi, Lattice and Heegaard Floer homologies of algebraic links, Int. Math. Res. Not. 2015, no. 23, 12737-12780.
- [4] A. Hatcher, "Algebraic topology." Cambridge University Press, 2002.
- [5] A. Kosinski, "Differential manifolds." Academic Press, Inc., 1993.
- [6] A. Levine, D. Ruberman, Generalized Heegaard Floer correction terms, Proceedings of the G'okova Geometry/Topology Conference, G'okova, 2014.
- [7] Y. Liu, L-space surgeries on links, Quantum Topol. 8 (2017), no. 3, 505-570.
- [8] C.Manolescu, P. Ozsváth, Heegaard Floer homology and integer surgeries on links, preprint 2010, arXiv:1011.11317.
- [9] C. Livingston, Knot 4-genus and the rank of classes in $W(\mathbb{Q}(t))$, Pacific J. Math., 252(1): 113-126, 2011.
- [10] K. Murasugi, On a certain numerical invariant of link types, Trans. Amer. Math. Soc., 117:387-422, 1965.
- P. Ozsváth, Z. Szabó, Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary, Adv. Math. 173 (2003) 179-261 MR1988284.
- [12] P. Ozsváth, Z. Szabó, Holomorphic disks and knot invariants, Adv. Math. 186 (2004) 58-116.
- [13] P. Ozsváth, Z. Szabó, Holomorphic disks and three-manifold invariants: Properties and applications, Annal of Mathematics, 159 (2004), 1159-1245.
- [14] P. Ozsváth, Z. Szabó, On knot Floer homology and lens space surgeries, Topology 44 (2005), no.6, 1281-1300.
- [15] P. Ozsváth, Z. Szabó, *Heegaard diagram and Floer homology*, International Congress of Mathematicians. Vol. II, 1083-1099, Eur. Math. Soc., Zürich, 2006.
- [16] P. Ozsváth, Z. Szabó. Holomorphic discs, link invariants and the multi-variable Alexander polynomial, Geom. Topol. 8 (2008), no. 2, 615-692.
- [17] P. Ozsváth, Z. Szabó, Link Floer homology and the Thurston norm, J. Amer. Math. Soc. 21 (2008) 671-709.
- [18] M. Powell, The four-genus of a link, Levine-Tristram signatures and satellites, J. Knot Theory Ramifications 26 (2017), no. 2, 1740008, 28 pp.
- [19] J. Rasmussen, Floer homology and knot complements, Ph.D. thesis, Harvard University, 2003.
- [20] J. Rasmussen, Lens space surgeries and a conjecture of Goda and Teragaito, Geom. Topol. 8 (2004), 1013-1031.
- [21] V. Turaev, Reidemeister torsion in knot theory, Russian Math. Surveys 41(1986), No. 1, 119-182.

B.L.: DEPARTMENT OF MATHEMATICS, UC DAVIS, ONE SHIELDS AVENUE, DAVIS CA 95616, USA *E-mail address*: bxliu@math.ucdavis.edu