

# THE ALON–MILMAN THEOREM FOR NON-SYMMETRIC BODIES

MÁRTON NASZÓDI

ABSTRACT. A classical theorem of Alon and Milman states that any  $d$  dimensional centrally symmetric convex body has a projection of dimension  $m \geq e^{c\sqrt{\ln d}}$  which is either close to the  $m$ -dimensional Euclidean ball or to the  $m$ -dimensional cross-polytope. We extended this result to non-symmetric convex bodies.

## 1. INTRODUCTION

Some fundamental results from the theory of normed spaces have been shown to hold in the more general setting of non-symmetric convex bodies. Dvoretzky's theorem [Dvo61, Mil71] was extended in [LM75] and [Gor88]; Milman's Quotient of Subspace theorem [Mil85] and duality of entropy results were extended in [MP00]. In this note, we extend the Alon–Milman Theorem.

A *convex body* is a compact convex set in  $\mathbb{R}^d$  with non-empty interior. We denote the orthogonal projection onto a linear subspace  $H$  or  $\mathbb{R}^d$  by  $P_H$ . For  $p = 1, 2, \infty$ , the closed unit ball of  $\ell_p^d$  centered at the origin is denoted by  $\mathbf{B}_p^d$ . Let  $K$  and  $L$  be convex bodies in  $\mathbb{R}^d$  with  $L = -L$ . We define their *distance* as

$$d(K, L) = \inf\{\lambda > 0: L \subset T(K - a) \subset \lambda L \text{ for some } a \in \mathbb{R}^d \text{ and } T \in GL(\mathbb{R}^d)\}.$$

By compactness, this infimum is attained, and when  $K = -K$ , it is attained with  $a = 0$ .

Alon and Milman [AM83] proved the following theorem in the case when  $K$  is centrally symmetric.

**Theorem 1.** *For every  $\varepsilon > 0$  there is a constant  $C(\varepsilon) > 0$  with the property that in any dimension  $d \in \mathbb{Z}^+$ , and for any convex body  $K$  in  $\mathbb{R}^d$ , at least one of the following two statements hold:*

- (i) *there is an  $m$ -dimensional linear subspace  $H$  of  $\mathbb{R}^d$  such that  $d(P_H(K), \mathbf{B}_2^m) < 1 + \varepsilon$ , for some  $m$  satisfying  $\ln \ln m \geq \frac{1}{2} \ln \ln d$ , or*
- (ii) *there is an  $m$ -dimensional linear subspace  $H$  such that  $d(P_H(K), \mathbf{B}_1^m) < 1 + \varepsilon$ , for some  $m$  satisfying  $\ln \ln m \geq \frac{1}{2} \ln \ln d - C(\varepsilon)$ .*

The main contribution of the present note is a way to deduce Theorem 1 from the original result of Alon and Milman, that is, the centrally symmetric case. By polarity, one immediately obtains

---

2010 *Mathematics Subject Classification.* (Primary) 52A21, 52A21 (Secondary) 46B20.

*Key words and phrases.* Section, projection of convex body, Banach–Mazur distance, irreducible set, almost Euclidean section.

The author thanks Alexander Litvak and Nicole Tomczak-Jaegermann for the discussions on the topic we had some time ago. The research was partially supported by the National Research, Development and Innovation Office (NKFIH) grant NKFI-K119670.

**Corollary 1.** *For every  $\varepsilon > 0$  there is a constant  $C(\varepsilon) > 0$  with the property that in any dimension  $d \in \mathbb{Z}^+$ , and for any convex body  $K$  in  $\mathbb{R}^d$  containing the origin in its interior, at least one of the following two statements hold:*

- (i) *there is an  $m$ -dimensional linear subspace  $H$  of  $\mathbb{R}^d$  such that  $d(H \cap K, \mathbf{B}_2^m) < 1 + \varepsilon$ , for some  $m$  satisfying  $\ln \ln m \geq \frac{1}{2} \ln \ln d$ , or*
- (ii) *there is an  $m$ -dimensional linear subspace  $H$  such that  $d(H \cap K, \mathbf{B}_\infty^m) < 1 + \varepsilon$ , for some  $m$  satisfying  $\ln \ln m \geq \frac{1}{2} \ln \ln d - C(\varepsilon)$ .*

## 2. PROOF OF THEOREM 1

For a convex body  $K$  in  $\mathbb{R}^d$ , we denote its polar by  $K^* = \{x \in \mathbb{R}^d : \langle x, y \rangle \leq 1 \text{ for all } y \in K\}$ . The *support function* of  $K$  is  $h_K(x) = \sup\{\langle x, y \rangle : y \in K\}$ . For basic properties, see [Sch14, BGVV14].

First in Lemma 2, by a standard argument, we show that if the *difference body*  $L - L$  of a convex body  $L$  is close to the Euclidean ball, then so is some linear dimensional section of  $L$ . For this, we need Milman's theorem whose proof (cf. [Mil71, FLM77, MS86]) does not use the symmetry of  $K$  even if it is stated with that assumption.

**Lemma 1** (Milman's Theorem). *For every  $\varepsilon > 0$  there is a constant  $C(\varepsilon) > 0$  with the property that in any dimension  $d \in \mathbb{Z}^+$ , and for any convex body  $K$  in  $\mathbb{R}^d$  with  $\mathbf{B}_2^d \subseteq K$ , there is an  $m$ -dimensional linear subspace  $H$  of  $\mathbb{R}^d$  such that  $(1 - \varepsilon)r(\mathbf{B}_2^d \cap H) \subseteq K \subseteq (1 + \varepsilon)r(\mathbf{B}_2^d \cap H)$ , for some  $m$  satisfying  $m \geq C(\varepsilon)M^2d$ , where*

$$M = M(K) = \int_{\mathbb{S}_2^{d-1}} \|x\|_K d\sigma(x),$$

and  $r = \frac{1}{M}$ .

**Lemma 2.** *Let  $\alpha, \varepsilon > 0$  be given. Then there is a constant  $c = c(\alpha, \varepsilon)$  with the property that in any dimension  $m \in \mathbb{Z}^+$ , and for any convex body  $L$  in  $\mathbb{R}^m$  with  $d(L - L, \mathbf{B}_2^m) < 1 + \alpha$ , there is a  $k$  dimensional linear subspace  $F$  of  $\mathbb{R}^m$  such that  $d(P_F(L), \mathbf{B}_2^k) < 1 + \varepsilon$  for some  $k \geq cm$ .*

*Proof.* Let  $\delta = d(L - L, \mathbf{B}_2^m)$ . We may assume that  $\frac{1}{\delta}\mathbf{B}_2^m \subseteq L - L \subseteq \mathbf{B}_2^m$ . Thus,  $h_{L-L} \geq \frac{1}{\delta}$ . With the notations of Lemma 1, we have

$$\begin{aligned} (1) \quad M(L^*) &= \int_{\mathbb{S}_2^{d-1}} \|x\|_{L^*} d\sigma(x) = \frac{1}{2} \int_{\mathbb{S}_2^{d-1}} h_L(x) + h_L(-x) d\sigma(x) \\ &= \frac{1}{2} \int_{\mathbb{S}_2^{d-1}} h_{L-L}(x) d\sigma(x) \geq \frac{1}{2\delta} \geq \frac{1}{2(1+\alpha)}. \end{aligned}$$

Note that  $L^* \supset (L - L)^* \supset \mathbf{B}_2^d$ , thus, by Lemma 1 and polarity, we obtain that  $L$  has a  $k$  dimensional projection  $P_F$  with  $d(P_F L, \mathbf{B}_2^d \cap F) \leq 1 + \varepsilon$  and  $k \geq C(\varepsilon) \frac{1}{4(1+\alpha)^2} m$ . Here,  $C(\varepsilon)$  is the same as in Lemma 1.  $\square$

The novel geometric idea of our proof is the following. We call a convex body  $T = \text{conv}(T_1 \cup \{\pm e\})$  in  $\mathbb{R}^m$  a *double cone* if  $T_1 = -T_1$  is convex set,  $\text{span} T_1$  is an  $(m-1)$ -dimensional linear subspace, and  $e \in \mathbb{R}^m \setminus \text{span} T_1$ . Double cones are *irreducible convex bodies*, that is, for any double cone  $T$ , if  $T = L - L$  then  $L = T/2$ , see [NV03, Yos91]. We prove a stability version of this fact.

**Lemma 3** (Stability of irreducibility of double cones). *Let  $L$  be a convex body in  $\mathbb{R}^m$  with  $m \geq 2$ , and  $T$  be a double cone of the form  $T = \text{conv}(T_1 \cup \{\pm e\})$ . Assume that  $T \subseteq L - L \subseteq \delta T$  for some  $1 \leq \delta < \frac{3}{2}$ . Then*

$$\left(\frac{3}{2} - \delta\right) T \subseteq L - a \subseteq \left(\delta - \frac{1}{2}\right) T.$$

for some  $a \in \mathbb{R}^m$ .

*Proof.* By the assumptions,  $e \in T \subseteq L - L$ , thus, by translating  $L$ , we may assume that  $0, e \in L$ . Thus,

$$(2) \quad L \subseteq (L - L) \cap (L - L + e) \subseteq \delta T \cap (\delta T + e).$$

We claim that

$$(3) \quad \delta T \cap (\delta T + e) = \frac{e}{2} + \left(\delta - \frac{1}{2}\right) T.$$

Indeed, let  $H_\lambda$  denote the hyperplane  $H_\lambda = \lambda e + e^\perp$ . To prove (3), we describe the sections of the right hand side and the left hand side by the hyperplanes  $H_\lambda$  for all relevant values of  $\lambda$ . For any  $\lambda \in [-\delta, \delta]$ , we have

$$\delta T \cap H_\lambda = \delta(T \cap H_{\lambda/\delta}) = \lambda e + \delta \left(1 - \frac{|\lambda|}{\delta}\right) T_1.$$

For any  $\lambda \in [-\delta + 1, \delta + 1]$ , we have

$$(\delta T + e) \cap H_\lambda = e + (\delta T \cap H_{\lambda-1}) = \lambda e + \delta \left(1 - \frac{|\lambda-1|}{\delta}\right) T_1.$$

Thus, for any  $\lambda \in [-\delta + 1, \delta]$ , we have

$$\delta T \cap (\delta T + e) \cap H_\lambda = \lambda e + \delta \left(1 - \frac{1}{\delta} \max\{|\lambda|, |\lambda-1|\}\right) T_1.$$

On the other hand, for any  $\lambda \in [-\delta + 1, \delta]$ , we have

$$(e/2 + (\delta - 1/2)T) \cap H_\lambda = \lambda e + (\delta - 1/2) \left(1 - \frac{|\lambda - 1/2|}{\delta - 1/2}\right) T_1.$$

Combining these two equations yields (3).

Thus,

$$T \subseteq L - L = \left(L - \frac{e}{2}\right) - \left(L - \frac{e}{2}\right) \subseteq \left(L - \frac{e}{2}\right) - \left(\delta - \frac{1}{2}\right) T.$$

Using the fact that  $T = -T$ , and  $1 \leq \delta < 3/2$ , we obtain

$$\left(\frac{3}{2} - \delta\right) T \subseteq L - \frac{e}{2},$$

finishing the proof of Lemma 3. □

Now, we are ready to prove Theorem 1. With the notations of the theorem, let  $D = K - K$ , and apply the symmetric version of the theorem for  $D$  in place of  $K$ . We may assume that  $\varepsilon < 1/2$ . In case (i), we use Lemma 2 and loose a linear factor in the dimension of the almost-euclidean projection. In case (ii), we use Lemma 3 with  $T = \mathbf{B}_1^m$  and obtain the same dimension for the almost- $\ell_1^m$  projection.

#### REFERENCES

- [AM83] N. Alon and V. D. Milman. Embedding of  $l_\infty^k$  in finite-dimensional Banach spaces. *Israel J. Math.*, 45(4):265–280, 1983.
- [BGVV14] Silouanos Brazitikos, Apostolos Giannopoulos, Petros Valettas, and Beatrice-Helen Vritsiou. *Geometry of isotropic convex bodies*, volume 196 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2014.
- [Dvo61] Aryeh Dvoretzky. Some results on convex bodies and Banach spaces. In *Proc. Internat. Sympos. Linear Spaces (Jerusalem, 1960)*, pages 123–160. Jerusalem Academic Press, Jerusalem; Pergamon, Oxford, 1961.
- [FLM77] T. Figiel, J. Lindenstrauss, and V. D. Milman. The dimension of almost spherical sections of convex bodies. *Acta Math.*, 139(1-2):53–94, 1977.
- [Gor88] Yehoram Gordon. Gaussian processes and almost spherical sections of convex bodies. *Ann. Probab.*, 16(1):180–188, 1988.
- [LM75] D. G. Larman and P. Mani. Almost ellipsoidal sections and projections of convex bodies. *Math. Proc. Cambridge Philos. Soc.*, 77:529–546, 1975.
- [Mil71] V. D. Milman. A new proof of A. Dvoretzky’s theorem on cross-sections of convex bodies. *Funkcional. Anal. i Priložen.*, 5(4):28–37, 1971.
- [Mil85] V. D. Milman. Almost Euclidean quotient spaces of subspaces of a finite-dimensional normed space. *Proc. Amer. Math. Soc.*, 94(3):445–449, 1985.
- [MP00] V. D. Milman and A. Pajor. Entropy and asymptotic geometry of non-symmetric convex bodies. *Adv. Math.*, 152(2):314–335, 2000.
- [MS86] Vitali D. Milman and Gideon Schechtman. *Asymptotic theory of finite-dimensional normed spaces*, volume 1200 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986. With an appendix by M. Gromov.
- [NV03] Márton Naszódi and Balázs Visy. Sets with a unique extension to a set of constant width. In *Discrete geometry*, volume 253 of *Monogr. Textbooks Pure Appl. Math.*, pages 373–380. Dekker, New York, 2003.
- [Sch14] Rolf Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 151 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, expanded edition, 2014.
- [Yos91] David Yost. Irreducible convex sets. *Mathematika*, 38(1):134–155, 1991.

DEPARTMENT OF GEOMETRY, EÖTVÖS LORÁND UNIVERSITY, BUDAPEST, HUNGARY  
*E-mail address:* marton.naszodi@math.elte.hu