

# Monotone flows with dense periodic orbits

Morris W. Hirsch<sup>\*</sup>  
University of Wisconsin

November 13, 2018

## Abstract

Give  $\mathbb{R}^n$  the (partial) order  $\geq$  determined by a closed convex cone  $K$  having nonempty interior:  $y \geq x \iff y - x \in K$ . Let  $X \subset \mathbb{R}^n$  be a connected open set and  $\varphi$  a flow on  $X$  that is monotone for this order: if  $y \geq x$  and  $t \geq 0$ , then  $\varphi^t y \geq \varphi^t x$ .

*Theorem.* If periodic points of  $\varphi$  are dense in  $X$ , then  $\varphi^t$  is the identity map of  $X$  for some  $t \geq 0$ .

## Contents

<b>1 Introduction</b>	<b>1</b>
Terminology . . . . .	2
<b>2 Resonance</b>	<b>4</b>
<b>3 Proof of Theorem 1</b>	<b>6</b>

## 1 Introduction

Many dynamical systems, especially those used as models in applied fields, are *monotone*: the state space has an order relation that is preserved in positive time. A recurrent theme is that bounded orbits tend toward periodic orbits. For a sampling of the large literature on monotone dynamics, consult the following works and the references therein: [2, 1, 3, 4, 6, 7, 8, 9, 11, 13, 15, 16, 20, 21, 23, 24, 25, 28, 30, 32, 33, 34, 35, 37, 39, 38].

Another common dynamical property is *dense periodicity*: periodic points are dense in the state space. Often considered typical of chaotic dynamics, this condition is closely connected to many other important dynamical topics, such as structural stability, ergodic theory, Hamiltonian mechanics, smoothness of flows and diffeomorphisms.

The main result in this article, Theorem 1, is that a large class of monotone, densely periodic flows  $\varphi := \{\varphi^t\}_{t \in \mathbb{R}}$  on open subsets of  $\mathbb{R}^n$  are (globally) *periodic*: There exists  $t > 0$  such that  $\varphi^t$  is the identity map.

---

<sup>\*</sup>I thank ERIC BACH, BAS LEMMENS and JANUSZ MIERCZYŃSKI for helpful discussions.

## Terminology

Throughout this paper  $X$  and  $Y$  denote (topological) spaces. The closure of a subset of a space is  $\bar{S}$ . The group of homeomorphisms of  $X$  is  $\mathcal{H}(X)$ .

$\mathbb{Z}$  denotes the integers,  $\mathbb{N}$  the nonnegative integers, and  $\mathbb{N}_+$  the positive integers.  $\mathbb{R}$  denotes the reals,  $\mathbb{Q}$  the rationals, and  $\mathbb{Q}_+$  the positive rationals.  $\mathbb{R}^n$  is Euclidean  $n$ -space.

Every manifold  $M$  is assumed metrizable with empty boundary  $\partial M$ , unless otherwise indicated.

Every map is assumed continuous unless otherwise described.  $f: X \approx Y$  means  $f$  maps  $X$  homeomorphically onto  $Y$ . The identity map of  $X$  is  $id_X$ .

If  $f$  and  $g$  denote maps, the composition of  $g$  following  $f$  is the map  $g \circ f: x \mapsto g(f(x))$  (which may be empty). We set  $f^0 := id_X$ , and recursively define the  $k$ 'th iterate  $f^k$  of  $f$  as  $f^k := f \circ f^{k-1}$ , ( $k \in \mathbb{N}_+$ ).

The *orbit* of  $p$  under  $f$  is the set

$$O(p) := \{f^k(p) : k \in \mathbb{N}\},$$

denoted as  $O(p, f)$  to record  $f$ .

When  $f(X) \subset X$ , the orbit of  $P \subset X$  is defined as

$$O(P) = O(P, f) := \bigcup_{p \in P} O(p).$$

A set  $A \subset X$  is *invariant* under  $f$  if  $f(A) = A = f^{-1}(A)$ . The *fixed set* and *periodic set* of  $f$  are the respective invariant sets

$$\mathcal{F}(f) := \{x : f(x) = x\}, \quad \mathcal{P}(f) := \bigcup_{n \in \mathbb{N}_+} \mathcal{F}(f^n).$$

**Flows** A *flow*  $\psi := \{\psi^t\}_{t \in \mathbb{R}}$  on  $Y$ , denoted formally by  $(\psi, Y)$ , is a continuous action of the group  $\mathbb{R}$  on  $Y$ : a family of homeomorphisms  $\psi^t: Y \approx Y$  such that the map

$$\mathbb{R} \rightarrow \mathcal{H}(Y), \quad t \mapsto \psi^t$$

is a homomorphism, and the *evaluation map*

$$\text{ev}_\psi: \mathbb{R} \times Y \rightarrow Y, \quad (t, x) \mapsto \psi^t x \tag{1}$$

is continuous.

A set  $A \subset Y$  is *invariant under  $\psi$*  if it is invariant under every  $\psi^t$ . When this holds, the *restricted flow*  $(\psi|_A)$  on  $A$  has the evaluation map  $\text{ev}_\psi|_{\mathbb{R} \times A}$ .

The *orbit* of  $y \in Y$  under  $\psi$  is the invariant set

$$O(y) := \{\psi^t y : t \in \mathbb{R}\},$$

denoted formally by  $O(y, \psi)$ . Orbits of distinct points either coincide or are disjoint.

The *periodic* and *equilibrium* sets of  $\psi$  are the respective invariant sets

$$\mathcal{P}(\psi) := \bigcup_{t \in \mathbb{R}} \mathcal{F}(\psi^t), \quad \mathcal{E}(\psi) := \bigcap_{t \in \mathbb{R}} \mathcal{F}(\psi^t),$$

denoted by  $\mathcal{P}$  and  $\mathcal{E}$  if  $\psi$  is clear from the context. Points in  $\mathcal{E}$  are called *stationary*. The orbit of a nonstationary periodic point is a *cycle*.

The *period* of  $p \in Y$  is  $r = \text{per}(p) > 0$ , provided

$$p \in \mathcal{P} \setminus \mathcal{E}, \quad r = \min \{t > 0: \psi^t p = p\}$$

The set of *r-periodic* points is

$$\mathcal{P}^r(\Psi) = \mathcal{P}^r := \{p \in \mathcal{P}: \text{per}(p) = r\}$$

When  $p$  is  $r$ -periodic, the restricted flow  $\psi|_{O(p)}$  is topologically conjugate to the rotational flow on the topological circle  $\mathbb{R}/r\mathbb{Z}$ , covered by the translational flow on  $\mathbb{R}$  whose evaluation map is  $(t, x) \mapsto t + x$ .

The flow  $(\psi, Y)$  is:

- *trivial* if  $\mathcal{E} = Y$ ,
- *densely periodic* if  $\mathcal{P}$  is dense in  $Y$ ,
- *pointwise periodic* if  $\mathcal{P} = Y$ ,
- *periodic* if there exists  $l > 0$  such that  $\psi^l = id_Y$ . Equivalently: the homomorphism  $\psi: \mathbb{R} \rightarrow \mathcal{H}(Y)$  factors through a homomorphism of the circle group  $\mathbb{R}/\mathbb{Z}$ .

**Order** Recall that a (partial) *order* on  $Y$  is a binary relation  $\leq$  on  $Y$  that is reflexive, transitive and antisymmetric. In this paper all orders are *closed*: the set  $\{(u, v) \in Y \times Y: u \leq v\}$  is closed. The *trivial order* is:  $x \leq y \iff x = y$ .

The pair  $(Y, \leq)$  is an *ordered space*, denoted by  $Y$  if the order is clear from the context.

We write  $y \geq x$  to mean  $x \leq y$ . If  $x \leq y$  and  $x \neq y$ , we write  $x < y$  and  $y > x$ . For sets  $A, B \subset Y$ , the condition  $A \leq B$  means  $a \leq b$  for all  $a \in A, b \in B$ , and similarly for the relations  $<, \geq, >$ .

The *closed order interval* spanned by  $a, b \in Y$  is the closed set

$$[a, b] := \{y \in Y: a \leq y \leq b\},$$

and its interior is the *open order interval*  $[[a, b]]$ . We write  $a \ll b$  to indicate  $[[a, b]] \neq \emptyset$ .

An *order cone*  $K \subset \mathbb{R}^n$  is a closed convex cone that is pointed (contains no affine line) and solid (has nonempty interior).  $K$  is *polyhedral* if is the intersection of finitely many closed linear halfspaces.

The *K-order*, denoted by  $\leq_K$  for clarity, is the order defined on every subset of  $\mathbb{R}^n$  by

$$x \leq_K y \iff y - x \in K.$$

For the  $K$ -order on  $\mathbb{R}^n$ , every order interval is a convex  $n$ -cell.

A map  $f: X \rightarrow Y$  between ordered spaces is *monotone* when

$$x \leq_X x' \implies f(x) \leq_Y f(x').$$

When  $Y$  is ordered and  $g: Y' \rightarrow Y$  is injective,  $Y'$  has a unique *induced order* making  $g$  monotone.

Every subspace  $S \subset Y$  is given the order induced by the inclusion map  $S \hookrightarrow Y$ . When this order is trivial,  $S$  is *unordered*.

A flow  $\psi$  on an ordered space is *monotone* iff the maps  $\psi^t$  are monotone. It is easy to see that every cycle for a monotone flow is unordered.

This is the chief result:

**Theorem 1.** *Assume:*

**(H1)**  $K \subset \mathbb{R}^n$  is an order cone.

**(H2)**  $X \subset \mathbb{R}^n$  is open and connected.

**(H3)**  $X$  is  $K$ -ordered.

**(H4)** The flow  $(\varphi, X)$  is monotone and densely periodic.

Then  $\varphi$  is periodic.

The proof will be given after results about various types of flows.

## 2 Resonance

In this section:

- $Y$  is an ordered space,
- $(\psi, Y)$  is a monotone flow,
- $O(y)$  is the orbit of  $y$  under  $\psi$ ,
- $O(y, \psi^t)$  is the orbit of  $y$  under  $\psi^t$ .

**Lemma 2.** *Assume:*

**(i)**  $\{p_k\}$  and  $\{q_k\}$  are sequences in  $Y$  converging to  $y \in Y$ ,

**(ii)**  $O(p_k) < O(q_k)$ ,  $(k \geq 1)$ .

Then  $y \in \mathcal{E}$ .

*Proof.* By (i) and continuity of  $\psi$  we have

$$\lim_{k \rightarrow \infty} \psi^t p_k = \lim_{k \rightarrow \infty} \psi^t q_k = \psi^t y, \quad (t \in \mathbb{R}) \quad (2)$$

while (ii) and monotonicity of  $\psi$  imply

$$k \in \mathbb{N}, t \in \mathbb{R} \implies p_k \leq \psi^t q_k. \quad (3)$$

From (i), (2), (3) and continuity of  $\psi$  we infer:

$$y \leq \psi^t y, \quad (t \in \mathbb{R}).$$

Therefore monotonicity shows that  $\psi^t y = y$  for all  $t \in \mathbb{R}$ . ■

**Rationality** Rational numbers play a surprising role in monotone flows:

**Theorem 3.** *Assume  $r, s > 0$ . If*

$$p \in \mathcal{P}^r, \quad q \in \mathcal{P}^s, \quad p < q, \quad \text{and} \quad O(p) \not\prec O(q),$$

*then  $r/s$  is rational.*

*Proof.* We have

$$O(p) \cap O(q) = \emptyset \tag{4}$$

because  $p < q$  and every cycle in a monotone flow is unordered.

Note that the restriction of  $\psi^s$  to the circle  $O(p)$  is conjugate to the rotation  $R_{2\pi r/s}$  of the unit circle  $\mathbb{C} \subset \mathbb{R}^2$  through  $2\pi r/s$  radians.

Arguing by contradiction, we provisionally assume  $r/s$  is irrational. Then the orbit of  $R_{2\pi r/s}$  is dense in  $\mathbb{C}$ ,<sup>1</sup> and the conjugacy described above implies

$$O(p) = \overline{O(p, \psi^r)}. \tag{5}$$

Since  $p < q$ , monotonicity

$$(\psi^r)^k p \leq (\psi^s)^k q = q, \quad (k \in \mathbb{N}_+),$$

whence

$$O(p; \psi^r) \leq \{q\},$$

and (5) implies

$$O(p) < \{q\}. \tag{6}$$

Therefore  $O(p) < O(q)$  by monotonicity (4) and monotonicity. But this contradicts the hypothesis. ■

**Definition 4.** A set  $S$  is *resonant* for the flow  $(\psi, Y)$  if  $S \subset Y$  and

$$u, v \in S \cap (\mathcal{P} \setminus \mathcal{E}) \implies \frac{\text{per}(u)}{\text{per}(v)} \in \mathbb{Q}_+.$$

It is easy to prove:

**Lemma 5.** *If  $S$  is resonant, so is its orbit.* ■

**Proposition 6.**  *$S$  is resonant provided there exists*

$$q \in S \cap (\mathcal{P} \setminus \mathcal{E})$$

*such that*

$$z \in S \cap (\mathcal{P} \setminus \mathcal{E}) \implies \frac{\text{per}(z)}{\text{per}(q)} \in \mathbb{Q}_+. \tag{7}$$

---

<sup>1</sup>This was discovered— in the 14th century!— by NICOLE ORESME. See GRANT [10], KAR [17]. A short proof based on the pigeon-hole principle is due to SPEYER [36]. Stronger density theorems are given in BOHR [5], KRONECKER [19], and WEYL [40, 41].

*Proof.* If  $u, v \in S \cap \mathcal{P}(\psi) \setminus \mathcal{E}$ , then

$$\frac{\text{per}(u)}{\text{per}(v)} = \frac{\text{per}(u)}{\text{per}(q)} \cdot \frac{\text{per}(q)}{\text{per}(v)},$$

which lies in  $\mathbb{Q}_+$  by (7). ■

**Theorem 7.** *Assume*

$$p, q \in \mathcal{P} \setminus \mathcal{E}, \quad p < q, \quad O(p) \neq O(q). \quad (8)$$

*Then  $[p, q]$  is resonant.*

*Proof.* Note that

$$p \leq z < q \implies O(z) \neq O(q). \quad (9)$$

For if this is false, there exists  $z \in [p, q]$  such that

$$O(z) < O(q),$$

whence monotonicity implies

$$(\forall w \in O(p)) (\exists w' \in O(z)): \quad w \leq w' < O(q).$$

It follows that  $O(p) < O(q)$ , contrary to hypothesis. Resonance of  $[p, q]$  now follows from Equation (9), and Theorem 3 with the parameters  $a := p$ ,  $b := q$ . ■

The next result will be used to derive the Main Theorem from the analogous result for homeomorphisms, Theorem 9.

**Theorem 8.** *Assume  $Y$  is resonant,  $s > 0$  and  $\mathcal{P}^s \neq \emptyset$ . Then  $\mathcal{P}(\psi) = \mathcal{P}(\psi^s)$ .*

*Proof.* We fix  $p \in \mathcal{P}^s$  and show that every  $q \in \mathcal{P}(\psi)$  lies  $\mathcal{P}(\psi^s)$ . If then  $q$  is stationary then  $q \in \mathcal{P}(\psi^s)$ . If  $q \in \mathcal{P}^r$  then  $r > 0$  and resonance implies  $sk = rl$  with  $k, l \in \mathbb{N}_+$ .

Since  $\psi^r q = q$ , we have  $T^k q = (\psi^s)^k q = (\psi^r)^l q = q$ , and again  $q \in \mathcal{P}(\psi^s)$ . ■

### 3 Proof of Theorem 1

Recall the statement of the Theorem:

*Assume:*

- (H1)  $K \subset \mathbb{R}^n$  is an order cone,
- (H2)  $X \subset \mathbb{R}^n$  is open and connected,
- (H3)  $X$  is  $K$ -ordered,
- (H4) the flow  $(\varphi, X)$  monotone and densely periodic.

Then  $\varphi$  is periodic.

The proof relies on two sufficient conditions for periodicity of monotone homeomorphisms, Theorems 9 and 10 below.

**Definitions.** A homeomorphism  $T: W \approx W$  is:

- *densely periodic* if  $\mathcal{P}(T)$  is dense in  $W$ ,
- *pointwise periodic* if  $\mathcal{P}(T) = W$ ,
- *periodic* if  $T^k = id_W$  for some  $k \in \mathbb{N}_+$ .

The key to Theorem 1 is this striking result:

**Theorem 9** (B. LEMMENS *et al.*, [22]). *Assume (H1), (H2), (H3). Then a monotone homeomorphism  $T: X \approx X$  is periodic provided it is densely periodic.*<sup>2</sup>

We also use an elegant result from the early days of transformation groups:

**Theorem 10** (D. MONTGOMERY, [26, 27]). *A homeomorphism of a connected topological manifold is periodic provided it is pointwise periodic.*<sup>3</sup>

**Lemma 11.**  *$\varphi$  is pointwise periodic*

*Proof.* Since  $\mathcal{E} \subset \mathcal{P}(\varphi)$ , it suffices to prove that the restriction of  $\varphi$  to each component of  $X \setminus \mathcal{E}$  is pointwise periodic. Therefore we assume  $\mathcal{E} = \emptyset$ .

Let  $x \in X$  be arbitrary. As  $\mathcal{P}(\varphi)$  is dense, Lemma 2 and Theorem 7 show there exist  $p, q \in \mathcal{P}(\varphi)$  such that the open set  $[[p, q]]$  is connected and resonant, and contains  $x$ . Therefore the open set  $Y := \mathcal{O}([[p, q]])$ , which is invariant and connected, is resonant by Lemma 5.

By Theorem 8 there exists  $s > 0$  such that  $\mathcal{P}(\varphi^s) = \mathcal{P}(\varphi) \cap Y$ . Consequently  $\varphi^s$  is densely periodic, hence periodic by Theorem 9. Therefore  $x \in \mathcal{P}(\varphi)$ . ■

Theorem 1 follows from Lemma 11 and Theorem 10.

## References

- [1] R. Anguelov, Y. Dumont, J. Lubuma, *Mathematical modeling of sterile insect technology for control of anopheles mosquito*, Computers & Mathematics with Applications **64** (2012), 374–389,
- [2] D. Angeli, M. Hirsch, E. Sontag, *Attractors in coherent systems of differential equations*, Journal of Differential Equations **246** (2009), 3058–3076.
- [3] M. Benâim & M. Hirsch, *Stochastic approximation algorithms with constant step size whose average is cooperative*, Annals of Applied Probability **9** (1999), 216–241.

<sup>2</sup>Conjectured, and proved for polyhedral cones, in M. HIRSCH [14].

<sup>3</sup>There are analogs of Montgomery’s Theorem for countable transformation groups; see KAUL [18], ROBERTS [31], YANG [42]. Pointwise periodic homeomorphisms on compact metric spaces are investigated in HALL & SCHWEIGERT [12].

- [4] M. Benaïm & M. Hirsch, *Mixed equilibria and dynamical systems arising from fictitious play in repeated games*, Games & Economic Behavior **29** (1999), 36–72.
- [5] H. Bohr, *Another proof of Kronecker’s theorem*, Proceedings of London Mathematical Society **2-21** (1923), 315–316.
- [6] P. De Leenheer, *The puzzle of partial migration*, Journal of Theoretical Biology **412** (2017), 172–185.
- [7] G. Dirr, H. Ito, A. Rantzer & B. Rüffer, *Separable Lyapunov functions for monotone systems: constructions and limitations*, Discrete & Continuous Dynamical Systems Series B **20** (2015), 2497–2526.
- [8] E. Balreira, S. Elaydi & R. Luis, *Global stability of higher dimensional monotone maps*, J. Difference Equations & Applications **23** (2017), 2037–2071.
- [9] G. Enciso & E. Sontag, *Global attractivity, I/O monotone small-gain theorems, and biological delay systems*, Discrete & Continuous Dynamical Systems **14** (2006), 249–578
- [10] E. Grant, “Nicole Oresme and the kinematics of circular motion: Tractatus de commensurabilitate vel incommensurabilitate motuum celi”, University of Wisconsin Press 1971
- [11] S. Grossberg, *Competition, decision and consensus*, Journal of Mathematical Analysis & Applications, **66** (1978), 470–493.
- [12] D. Hall & G. Schweigert, *Properties of invariant sets under pointwise periodic homeomorphisms*, Duke Math. J. **4** (1938), 719–724.
- [13] P. Hess & P. Polacik, *Boundedness of prime periods of stable cycles and convergence to fixed points in discrete monotone dynamical systems*, SIAM J. Math. Anal. **24** (1993), 1312–1330.
- [14] M. Hirsch, *Monotone dynamical systems with polyhedral order cones and dense periodic points*, AIMS Mathematics **2** (2017), 24–27. Also in: <http://arxiv.org/abs/1611.09251> (2016)
- [15] M. W. Hirsch & H. L. Smith, *Monotone Dynamical Systems*, “Handbook of Differential Equations: Ordinary Differential Equations, Vol. 2,” A. Cañada and P. Drábek and A. Fonda editors, 239–258. Elsevier North Holland, Boston, Massachusetts (2005)
- [16] E. Kamke, *Zur Theorie der Systeme gewöhnlicher differential-gleichungen, II*, Acta Math., **58** (1932), 57–85.
- [17] A. Kar, *Weyl’s Equidistribution Theorem*, Resonance **8** (2003), 30–37.
- [18] S. Kaul, *On pointwise periodic transformation groups*, Proceedings of American Mathematical Society, **27** (1971), 391–394.
- [19] L. Kronecker, *Näherungsweise ganzzahlige Auflösung linearer Gleichungen*, Werke 3, Chelsea reprint (1968), 47–109.



- [20] A. Lajmanovich & J. Yorke, *A deterministic model for gonorrhoea in a non-homogeneous population*, *Mathematical Biosciences* **28** (1976), 221–236.
- [21] A. Landsberg & E. Friedman, *Dynamical Effects of Partial Orderings in Physical Systems*, *Physical Review E*, **54** (1996), 3135–3141
- [22] B. Lemmens, O. van Gaans & H. van Imhoff, *Monotone dynamical systems with dense periodic points*, <https://arxiv.org/abs/1712.09223> (2017).
- [23] W. Leonard & R. May, *Nonlinear aspects of competition between species*, *SIAM Journal of Applied Mathematics* **29** (1975), 243–275.
- [24] H. Matano, *Strongly order-preserving local semi-dynamical systems-Theory and Applications*. *Semigroups, Theory and Applications*, Volume 1 (Editors: H. Brezis, M. Crandall, F. Kappel). *Research Notes in Mathematics* **141**, Longman Scientific & Technical, London, 178–185 (1986).
- [25] J. Mierczynski, *P-arcs in strongly monotone discrete-time dynamical systems*, *Differential & Integral Equations*, **7** (1994), 1473–1494.
- [26] D. Montgomery, *Pointwise periodic homeomorphisms*, *American Journal of Mathematics* **59** (1937), 118–120.
- [27] D. Montgomery, “Topological Transformation Groups,” Interscience (1955).
- [28] C. Potzsche, *Order-preserving nonautonomous discrete dynamics: Attractors and entire solutions*, *Positivity* **19** (2015), 547–576.
- [29] R. Redheffer & W. Walter, *Flow-invariant sets and differential inequalities in normed spaces*, *Applicable Analysis* **5** (1975) 149–1611.
- [30] R. Redheffer & Walter, *Remarks on ordinary differential equations in ordered Banach spaces*, *Monatshefte Mathematik*. **102** (1986), 237–249
- [31] J. Roberts, *Pointwise finite families of mappings*, *Canadian Mathematical Bulletin* **18** (1975), 767–768.
- [32] J. Selgrade, *Mathematical analysis of a cellular control process with positive feedback*, *SIAM Journal of Applied Mathematics* **36** (1979), 219–229.
- [33] S. Smale, *On the differential equations of species in competition*, *Journal of Mathematical Biology* **3** (1976), 5–7.
- [34] H. L. Smith, “Monotone Dynamical Systems, an introduction to the theory of competitive and cooperative systems,” *Math. Surveys & Monographs*, No. 41, American Mathematical Society, Providence, Rhode Island (1995).
- [35] H. L. Smith, *Monotone dynamical systems: reflections on new advances & applications*, *Discrete & Continuous Dynamical Systems Series B*, **37** (2017), 485–504.
- [36] D. Speyer, <https://mathoverflow.net/questions/75777> (2017). s

- [37] P. Volkmann, *Gewöhnliche Differentialungleichungen mit quasimonoton wachsenden Funktionen in topologischen Vektorräumen*, *Mathematische Zeitschrift* **17** (1972), 157–164.
- [38] S. Walcher, *On cooperative systems with respect to arbitrary orderings*, *Journal of Mathematical Analysis & Applications* **2** (2001), 543–554.
- [39] L. Feng, Y. Wang & J. Wu, *Semiflows monotone with respect to high-rank cones on a Banach space*, *SIAM Journal of Mathematical Analysis* **49** (2017), 142–161.
- [40] H. Weyl, *Über die Gibbs'sche Erscheinung und verwandte Konvergenz Phänomene*, *Rendiconti del Circolo Matematico di Palermo* **330** (1910), 377–407.
- [41] H. Weyl, *Über die Gleichverteilung von Zahlen mod. Eins*, *Mathematischen Annalen* **77** (1916), 313–352.
- [42] J. Yang, *Pointwise periodic transformation groups*, *Notices of American Mathematical Society* **18** (1971), 830.