

Decidability of the existential fragment of some infinitely generated trace monoids: an application to ordinals

Alexis Bès

Université Paris-Est, LACL (EA 4219), UPEC, Créteil, France

Christian Choffrut

IRIF, CNRS and Université Paris 7 Denis Diderot, France

November 4, 2021

Abstract

Diekert, Matiyasevich and Muscholl proved that the existential first-order theory of a trace monoid over a finite alphabet is decidable. We extend this result to a natural class of trace monoids with infinitely many generators. As an application, we prove that for every ordinal λ less than ε_0 , the existential theory of the set of successor ordinals less than λ equipped with multiplication is decidable.

1 Introduction

Since the publication of the surprising result of Makanin showing that it is decidable whether or not an equation in a free monoid with constants has a solution, research continued in different directions. Notably, Schulz showed that the decidability still holds when each variable is bound to be interpreted in a predefined regular subset of the free monoid, i.e., as subset recognized by some finite automaton, [10]. This allowed to extend Makanin's result to trace monoids which can be viewed as free monoids where some pairs of generators may commute. Indeed, building on top of the results of Makanin and Schulz, Diekert et al. were able to prove that it can be decided whether or not a system of equations in a trace monoid has a solution and more generally whether or not a sentence of the existential fragment the theory of trace monoids provided with the concatenation is valid.

Our contribution consists of going one step further by considering some trace monoids over a countable alphabet, namely those that are the inverse image in a generator-to-generator substitution into a finitely generated trace monoid and to show that the above result still holds in this framework.

The idea can be summarized as follows. Considering infinite generators is no problem as far as equations are concerned because a solution, if it exists, can always be assumed to map into the submonoid of the generators appearing in the constants. In contrast, the existential fragment requires the possibility of expressing the negation of an equality. When the monoid is finitely generated this can be done by a finite disjunction of equations containing new constants. For infinitely generated trace monoids, this leads to an infinite disjunction with infinite constants. The idea is then to observe that the actual values of these constants are irrelevant and can be reduced to a number of values that can be bounded a priori, allowing thus to resort to the case of finitely generated trace monoids.

We apply our result to an issue concerning the multiplicative structure of ordinals. Let us recall that the first-order theory of an ordinal α with the multiplication as unique operation is undecidable if and only if this ordinal is greater than or equal to ω^ω , see [1]. We are not aware of any investigation on which fragment, if any, of such an ordinal is decidable, except a paper of the present authors, [2]. It happens that the successor ordinals less than a multiplicatively closed ordinal λ ($\alpha, \beta < \lambda$ implies $\alpha \times \beta < \lambda$) form a monoid \mathcal{S}_λ which is a free product of an infinitely generated free commutative monoid (more precisely the monoid generated by the ordinary prime integers) and an infinitely generated free monoid. The result on traces implies that the existential fragment of \mathcal{S}_λ is decidable. More precisely we prove that if λ is less than ε_0 then the existential theory of the structure $\langle \mathcal{S}_\lambda : \times, \{\alpha\}_{\alpha \in \mathcal{S}_\lambda} \rangle$ is decidable. Recall that ε_0 is the least fixed point of the function $x \mapsto \omega^x$. The condition $\lambda < \varepsilon_0$ is here to ensure that one can perform effectively operations on constants given by their Cantor normal form.

This does not settle the problem of the decidability of the multiplicative structure of λ but we hope it arises the curiosity of some researchers.

2 Trace monoids

Our purpose in this section is to show how the result of Diekert and al. [4] for finitely generated trace monoid extends to some type of infinitely generated trace monoids. We first give a definition of trace monoids with possibly infinite generators.

2.1 Traces over possibly infinite alphabets

The notion of finitely generated trace monoids can be recovered from the next definition.

Definition 1. Let Σ be a countable set, $n > 0$ an integer, $\Sigma = \bigcup_{i=1}^n \Sigma_i$ a decomposition and $I \subseteq \{1, \dots, n\} \times \{1, \dots, n\}$ a symmetric relation. The

relation

$$\{(ab, ba) \mid a \in \Sigma_i, b \in \Sigma_j, (i, j) \in I\}$$

is the *independence relation* on Σ induced by I . We denote by \equiv_I the congruence on the free monoid Σ^* generated by the independence relation and by $\mathbb{M}(\Sigma, I)$ the quotient monoid Σ^* / \equiv_I , also known as the *trace monoid*. The *canonical morphism* is the mapping which associates with an element of Σ^* its class modulo \equiv_I . When Σ and I are clear from the context we simply write \mathbb{M} .

Observe that in contrast to the finite case, we do not impose the relation I to be irreflexive. If it were, we would necessarily have a free (noncommutative) submonoid.

Example 2. The free product of a free monoid generated by Σ_1 and a free commutative monoid generated by a disjoint subset Σ_2 is defined by the conditions $\Sigma = \Sigma_1 \cup \Sigma_2$, $n = 2$ and $I = \{(2, 2)\}$. The free product of three free commutative monoids is defined by taking three disjoint subsets $\Sigma_1, \Sigma_2, \Sigma_3$ and considering the independence relation $I = \{(i, i) \mid i = 1, 2, 3\}$. The direct product of two free monoids is defined by taking two disjoint subsets Σ_1, Σ_2 and considering the independence relation $I = \{(1, 2), (2, 1)\}$.

We will need no sophisticated result on trace monoids, only the following simple combinatorial result stating under which conditions two traces are different, see [4, page 8]. Further reading on this theory is referred, for example, to [5, 6].

Lemma 3. *Two elements $u, v \in \mathbb{M}(\Sigma, I)$ are different exactly under one of the following conditions (up to exchanging the roles of u and v)*

1. u is a strict prefix of v .
2. there exist $w, w_1, w_2, w_3 \in \mathbb{M}(\Sigma, I)$, $a \in \Sigma$ such that $u = waw_1, v = ww_2aw_3$ and w_2 contains no occurrence of a but contains an occurrence of some $b \neq a$ which does not commute with a , i.e., $(a, b) \notin I$.
3. there exist $w, w_1, w_2 \in \mathbb{M}(\Sigma, I)$, $a \in \Sigma$ such that $u = waw_1, v = ww_2$ and w_2 contains no a .

Proof. Clearly, the conditions 1, 2 and 3 are sufficient. We prove that they are necessary. Let w be the longest common prefix of u and v (such a prefix is guaranteed by Levi's Lemma, see [3, Prop. 1.3]). If $w = u$ this is condition 1. If $u = waw_1$ and $v = ww_2$ where w_2 contains no a this is item 3. Otherwise set $v = ww_2aw_3$ where w_2 contains no occurrence of a . Then w_2 certainly contains some b which does not commute with a . \square

2.2 Trace monoids as logical structures

The logical structures on trace monoid considered in this work contain at least the equality as nonlogical symbol, the product of traces as function and all their elements as constants.

We consider the structure $\mathcal{M} = \langle \mathbb{M}(\Sigma, I); =, \cdot, \{u\}_{u \in \mathbb{M}(\Sigma, I)} \rangle$. We are given a countable set of elements called *variables*. The family of *terms* is defined inductively by the conditions: a variable is a term, a constant is a term and if t and t' are two terms, so is $t \cdot t'$. If needed, we may write a term as $t(x_1, \dots, x_n)$ to signify that the variables occurring in t are among x_1, \dots, x_n . A *basic predicate* is of the form $t = t'$ where t and t' are two terms.

As much as possible, we use the lower case a, b, \dots for elements in Σ which we do not distinguish from generators of $\mathbb{M}(\Sigma, I)$, and lower case letters u, v, \dots for traces.

2.3 The case of finitely generated trace monoids

We recall the result of Diekert et al. [4]. In this case, the generator set Σ is finite and its decomposition consists of the union of all singletons $\{a\}$ for $a \in \Sigma$. The structure has additional predicates whose definition requires the following notion.

Definition 4. A subset K of $\mathbb{M}(\Sigma, I)$ is *regular* if it is the image, in the canonical mapping of Σ^* onto $\mathbb{M}(\Sigma, I)$, of a regular subset $K_0 \subseteq \Sigma^*$ which is *I-closed*, in the sense that the condition ($u \in K_0$ and $u \equiv_I v$) implies $v \in K_0$. The family of regular subsets is denoted by \mathcal{K} .

Observe that each singleton of $\mathbb{M}(\Sigma, I)$ is regular since it is the canonical image of a finite subset of Σ^* .

Theorem 5 ([4]). *Given a finitely generated trace monoid, the existential fragment of the first-order theory of the structure $\langle \mathbb{M}(\Sigma, I); =, \cdot, \{K\}_{K \in \mathcal{K}} \rangle$, where each K is viewed as a unary predicate, is decidable.*

2.4 The case of infinitely generated trace monoids

The objective is to prove the following.

Theorem 6. *Given a trace monoid as in Definition 1, the existential first-order theory of the structure $\langle \mathbb{M}(\Sigma, I); =, \cdot, \{u\}_{u \in \mathbb{M}(\Sigma, I)} \rangle$ is decidable.*

Proof. We fix the decomposition $\Sigma = \bigcup_{i=1}^n \Sigma_i$ and the independence relation I as in Definition 1 and we write \mathbb{M} for $\mathbb{M}(\Sigma, I)$ whenever no confusion may arise. A formula of the existential fragment of the theory of the trace monoid

is equivalent to a formula of the form

$$\begin{aligned}
& \exists x_1, \dots, x_n \bigvee_{k=1}^p (\mathcal{E}_k^+ \wedge \mathcal{E}_k^-) \\
& \text{where each } \mathcal{E}_k^+ \text{ is of the form} \\
& \bigwedge_{i \in I} t_i(x_1, \dots, x_n) = s_i(x_1, \dots, x_n) \\
& \text{and where each } \mathcal{E}_k^- \text{ is of the form} \\
& \bigwedge_{j \in J} t'_j(x_1, \dots, x_n) \neq s'_j(x_1, \dots, x_n)
\end{aligned} \tag{1}$$

Such a formula is *satisfiable* if there exists an interpretation

$$\theta : \{x_1, \dots, x_n\} \rightarrow \mathbb{M}(\Sigma, I),$$

such that $\theta(t_i) = \theta(s_i)$ for $i \in I$ and $\theta(t'_j) \neq \theta(s'_j)$ for $j \in J$.

For simplification purposes we replace the basic predicates by simpler predicates by applying the following rules.

- Constants: a constant $u \neq 1$ in a term is replaced by a new variable x and by adding the new condition $x = u$. All occurrences of the empty trace 1 are deleted except if the left- or right-hand side is reduced to 1.
- Equation $y_1 \cdots y_n = y_{n+1} \cdots y_m$ is replaced by the condition $\bigwedge_{i=1}^n y_i = 1$ if the right-hand side is reduced to the empty trace. Similarly if the left-hand side is reduced to 1. Otherwise the equation is of the form $y_1 \cdots y_n = y_{n+1} \cdots y_m$ with $1 < n < m$. Then we introduce m new variables z_i and replace the equation by a conjunction of simple predicates

$$\begin{aligned}
& z_1 = y_1 \wedge z_2 = z_1 y_2 \wedge \cdots \wedge z_n = z_{n-1} y_n \\
& \wedge z_{n+1} = y_{n+1} \wedge z_{n+2} = z_{n+1} y_{n+2} \wedge \cdots \wedge z_m = z_{m-1} y_m \\
& \wedge z_n = z_m
\end{aligned} \tag{2}$$

- Inequation $y_1 \cdots y_n \neq y_{n+1} \cdots y_m$ is replaced by the condition $\bigvee_{i=1}^n y_i \neq 1$ if the right-hand side is reduced to the empty trace. Similarly if the left-hand side is reduced to 1. Otherwise the inequation is of the form $y_1 \cdots y_n \neq y_{n+1} \cdots y_m$ with $1 < n < m$. As in the case of equality, we introduce m new variables z_i and replace the equation by a conjunction of simple predicates

$$\begin{aligned}
& z_1 = y_1 \wedge z_2 = z_1 y_2 \wedge \cdots \wedge z_n = z_{n-1} y_n \\
& \wedge z_{n+1} = y_{n+1} \wedge z_{n+2} = z_{n+1} y_{n+2} \wedge \cdots \wedge z_m = z_{m-1} y_m \\
& \wedge z_n \neq z_m
\end{aligned} \tag{3}$$

Equalities between two variables can be deleted by keeping one of the two variables only. By applying the previous rules and the routine Boolean rules, we may rewrite expression **1** in such a way that all its basic predicates are of the form $x = u$ or $xy = z$ where x, y, z are variables and u is a constant, possibly equal to 1, formally

$$\mathcal{E}_k^+ \equiv \bigwedge_{i \in J_1^{(k)}} x_i = u_i \wedge \bigwedge_{(i,j,h) \in J_2^{(k)}} x_i y_j = z_h \quad (4)$$

and

$$\mathcal{E}_k^- \equiv \bigwedge_{i \in J_3^{(k)}} x_i \neq 1 \wedge \bigwedge_{(i,j) \in J_4^{(k)}} x_i \neq y_j \quad (5)$$

Consequently, we may assume that we start off with a disjunction of **4** and **5**, i.e., of the form $\Phi \equiv \exists x_1, \dots, x_n (\mathcal{E}^+ \wedge \mathcal{E}^-)$ by omitting the index k .

Let $\Delta \subseteq \Sigma$ be the set of generators appearing as a factor of some constant in Φ , i.e., the smallest subset $\Sigma' \subseteq \Sigma$ satisfying the inclusion

$$\{u_i \mid i \in J_1\} \subseteq \mathbb{M}(\Sigma', I \cap (\Sigma' \times \Sigma')).$$

We claim that there exists a finite collection of finite subsets $\Delta \subseteq \Gamma_i \subseteq \Sigma$, $i = 1 \dots, \ell$, and existential formulas Φ_i , $i = 1 \dots, \ell$ in $\langle \mathbb{M}_i, \cdot, \mathcal{K}_i \rangle$ where $\mathbb{M}_i = \mathbb{M}(\Gamma_i, I \cap (\Gamma_i \times \Gamma_i))$ and $\mathcal{K}_i = \{L \subseteq \mathbb{M}_i \mid L \in \mathcal{K}\}$, such that the following property holds for all assignments $\theta : \{x_1, \dots, x_n\} \rightarrow \mathbb{M}(\Sigma, I)$

$$\begin{aligned} & \langle \mathbb{M}(\Sigma, I); \cdot, \{u\}_{u \in \mathbb{M}(\Sigma, I)} \rangle \models \Phi(\theta(x_1), \dots, \theta(x_n)) \\ & \text{iff there exists some } i \in \{1, \dots, \ell\} \text{ and some interpretation} \\ & \quad \theta_i : \{x_1, \dots, x_n\} \rightarrow \mathbb{M}_i \text{ such that} \\ & \quad \langle \mathbb{M}_i; \cdot, \mathcal{K}_i \rangle \models \Phi_i(\theta_i(x_1), \dots, \theta_i(x_n)) \end{aligned} \quad (6)$$

The idea is to restate the different conditions of inequality for traces. We fix an assignment θ . For each inequality $x \neq 1$ in **5**, the condition $\theta(x) \neq 1$ is satisfied if and only if there exist a generator $a \in \Sigma$ and an element $v \in \mathbb{M}$ such that

$$\theta(x) = av \quad (7)$$

Define

- $\Delta_\theta^{(1)}$ is the subset consisting of all $a \in \Sigma$ appearing in **7** when $x \neq 1$ ranges over all inequalities of this form in **5**
- $U_\theta^{(1)}$ is the subset consisting of all $v \in \mathbb{M}$ appearing in **7** when $x \neq 1$ ranges over all inequalities of this form in **5**

Similarly, for each inequality of the form $x \neq y$ in 5, by Lemma 3 the condition $\theta(x) \neq \theta(y)$ is equivalent to the disjunction of the following three cases.

Case 1: there exist an element $v \in \mathbb{M}$ and a generator $a \in \Sigma$ such that

$$\theta(y) = \theta(x)av \quad (8)$$

Case 2: there exist two integers $(i, j) \notin I$, two generators $a, b \in \Sigma$ and five elements $u, v, w, r, t \in \mathbb{M}$ such that

$$\begin{aligned} \theta(x) = uav, \theta(y) = urbtaw, a \in \Sigma_i, b \in \Sigma_j \text{ and } r, t \\ \text{contain no occurrence of } a \end{aligned} \quad (9)$$

Case 3: there exist $a \in \Sigma$, $u, v, w \in \mathbb{M}$ such that w contains no occurrence of a and

$$\theta(x) = uav \text{ and } \theta(y) = uw \quad (10)$$

Define

- $\Delta_\theta^{(2)}$ is the subset consisting of all a and b in Σ appearing in 8, 9 and 10, when $x \neq y$ ranges over all inequalities of this form in 5
- $U_\theta^{(2)}$ is the subset consisting of all $r, t, u, v, w \in \mathbb{M}$ appearing in 8, 9 and 10 when $x \neq y$ ranges over all inequalities of this form in 5

Set

$$\Delta_\theta = \Delta_\theta^{(1)} \cup \Delta_\theta^{(2)} \subseteq \Sigma, \quad U_\theta = U_\theta^{(1)} \cup U_\theta^{(2)} \subseteq \mathbb{M}$$

The cardinality of Δ_θ is bounded by the integer K which is 4 times the number of inequalities in expression 5 because each inequality of the form $x \neq 1$ introduces one generator and each inequality of the form $x \neq y$ introduces four not necessarily different generators. Observe that θ satisfies Φ if and only if so does $\pi \circ \theta$ where π maps all generators in $\Sigma \setminus (\Delta \cup \Delta_\theta)$ to the empty trace, so that we can, from now on, assume that θ maps each variable to the submonoid generated by $\Delta \cup \Delta_\theta$.

Let R_θ be the predicate that specifies for all pairs of generators (e, f) in $\Delta \cup \Delta_\theta$ whether they are equal or different and which sub-alphabet Σ_i they belong to. A permutation σ of Σ is *respectful* if it fixes each element of Δ and if it respects the membership to a specific Σ_i , i.e., if 1) $\sigma(e) = e$ if $e \in \Delta$ and 2) for all $e \in \Sigma$ and for all $i = 1, \dots, n$ we have $e \in \Sigma_i$ if and only if $\sigma(e) \in \Sigma_i$. Now observe that up to a respectful permutation, the number of possible predicates R_θ is finite: this is due to the fact that R_θ involves a number of generators bounded by a function of the size of the formula Φ . Furthermore, if θ satisfies Φ so does $\sigma \circ \theta$. Consequently, when θ ranges over the possible assignments satisfying Φ , up to a respectful permutation there

exists a finite number of different predicates R_θ , say R_1, \dots, R_ℓ , and each predicate involves a finite number of elements of Σ .

Consequently, if Φ is satisfiable, it is satisfiable by some assignment θ which maps the variables into the submonoid generated by a finite subset Γ_i containing Δ and satisfying a predicate R_i for some $i \in \{1, \dots, \ell\}$. We show that the existence of such an assignment is decidable. Indeed, observe that the actual subset Γ_i is irrelevant as long as it satisfies R_i . This means that we can consider the elements a and b as in expressions 7, 8, 9 and 10 as fixed constants of Σ . It remains to define the existential formulas Φ_i in the structure $\langle \mathbb{M}(\Gamma_i, I \cap (\Gamma_i \times \Gamma_i)); \cdot, \mathcal{K}_i \rangle$ as in the above claim 6. This is achieved as follows. We keep the clauses 4, modify the clauses in 5 as below and prefix the resulting formula by as many existential quantifiers as there are new variables. Concerning the modification of the clauses in 5, each inequality $x \neq 1$ is replaced by a condition

$$x = az_1$$

where z_1 is a new variable. Similarly, each inequality $x \neq y$ in 5 is replaced by a disjunction

$$(y = xaz_1) \vee (x = z_1az_2, \wedge y = z_1z_3bz_4az_5 \wedge z_3, z_4 \in \mathbb{M}(\Gamma_a, I \cap (\Gamma_a \times \Gamma_a))) \\ \vee (x = z_1az_2 \wedge y = z_1z_3 \wedge z_3 \in \mathbb{M}(\Gamma_a, I \cap (\Gamma_a \times \Gamma_a)))$$

where $\Gamma_a = \Gamma_i \setminus \{a\}$ and z_1, \dots, z_4 are new variables. Observe that no condition on the generators such as a and b above is required because these conditions are already covered by the predicate R_i .

We may now safely apply the result [4] because the only new predicates $\mathbb{M}(\Gamma_a, I \cap (\Gamma_a \times \Gamma_a))$ are clearly regular in the finitely generated trace monoid $\mathbb{M}(\Gamma_i, I \cap (\Gamma_i \times \Gamma_i))$.

□

3 An application to ordinals

We denote by Ord the class of ordinals. For a thorough exposition of ordinals we refer to the classical handbooks such as [9] and [8].

3.1 Arithmetic operations on the ordinals

The following definition of the *Cantor normal form*, abbreviated CNF, is actually a property in its own right.

Definition 7. Every nonzero ordinal α has a unique form as a sum of ω -powers with integer coefficients, namely

$$\alpha = \omega^{\lambda_r} a_r + \dots + \omega^{\lambda_1} a_1,$$

where $\lambda_r > \dots > \lambda_1 \geq 0$ are ordinals and $a_r, \dots, a_1 > 0$ are integers. A nonzero ordinal is a *successor* if $\lambda_1 = 0$, otherwise it is a *limit*.

We recall the definition of the multiplication on ordinals by use of their Cantor normal form.

Definition 8. Let

$$\alpha = \omega^{\lambda_r} a_r + \cdots + \omega^{\lambda_1} a_1, \quad \beta = \omega^{\mu_s} b_s + \cdots + \omega^{\mu_1} b_1,$$

be two nonzero ordinals written in CNF. If $\mu_1 > 0$ we have

$$\alpha \times \beta = \omega^{\lambda_r + \mu_s} b_s + \cdots + \omega^{\lambda_r + \mu_1} b_1. \quad (11)$$

If $\mu_1 = 0$ we have

$$\begin{aligned} \alpha \times \beta = \omega^{\lambda_r + \mu_s} b_s + \omega^{\lambda_r + \mu_{s-1}} b_{s-1} + \cdots + \omega^{\lambda_r + \mu_2} b_2 + \\ \omega^{\lambda_r} a_r b_1 + \omega^{\lambda_{r-1}} a_{r-1} + \cdots + \omega^{\lambda_1} a_1. \end{aligned} \quad (12)$$

Remark 9. The multiplication is associative, has a neutral element 1, is noncommutative, is left- (but not right-) *cancellative* ($x \times y = x \times z \Rightarrow y = z$) and left- (but not right-) distributes over the addition. With the definition of the multiplication it can be easily verified that an ordinal λ is closed under multiplication ($\alpha, \beta < \lambda$ implies $\alpha \times \beta < \lambda$) if and only if it is of the form ω^{ω^ξ} for some ordinal $\xi \geq 0$. Also since we are concerned with effectivity, we assume that λ is less than the ordinal ε_0 so that providing a Cantor Normal Form and performing operations such as comparing ordinals and finding divisors make sense.

3.2 Primes

Definition 10. An ordinal x is a *prime* if it has exactly two right divisors, i.e., two ordinals $z_1 \neq z_2$ for which there exist y_1, y_2 with $x = y_1 z_1 = y_2 z_2$.

This definition of prime is equivalent, as can be readily verified, to the standard definition which stipulates that it has exactly two right divisors 1 and x .

Definition 11. There are three kinds of primes, [9, p. 336].

- finite primes: the ordinary prime natural numbers
- non-finite successor primes: of the form $\omega^\lambda + 1$, $\lambda \in \text{Ord}$.
- limit primes: of the form ω^{ω^ξ} , $\xi \in \text{Ord}$

The main result concerning primes is the following

Theorem 12 (the prime factorization [7]). *Every ordinal has a unique factorization of the form*

$$(\omega^{\omega^{\xi_1}})^{n_1} \dots (\omega^{\omega^{\xi_r}})^{n_r} a_k (\omega^{\lambda_{k-1}} + 1) a_{k-1} (\omega^{\lambda_{k-1}} + 1) \dots a_1 (\omega^{\lambda_1} + 1) a_0$$

with $\xi_1 > \xi_2 > \dots > \xi_r$ and $\lambda_1, \lambda_2, \dots, \lambda_k \geq 1$ (Greek letters are arbitrary ordinals and Latin letters are finite ordinals).

Observe that the condition on the exponents of limit primes is necessary: ω and ω^ω are primes and $\omega^\omega = \omega \times \omega^\omega$.

3.3 The monoid of successor ordinals as a trace monoid

For every countable ordinal λ closed under multiplication and less than ε_0 , let \mathcal{S}_λ denote the set of all successor ordinals less than λ . The rule of multiplication on ordinals show that \mathcal{S}_λ forms a multiplicative submonoid.

Theorem 12 can be interpreted as follows. The ordinal ω is the submonoid generated by the finite primes (the ordinary prime integers). Let \mathcal{P}_λ be the submonoid generated by the infinite successor primes less than λ . Then Theorem 12 claims that \mathcal{S}_λ is the free product of the (infinitely generated) free commutative monoid ω and the (infinitely generated) free monoid \mathcal{P}_λ . We have

Theorem 13. *Given an ordinal λ less than ε_0 and closed under multiplication, the existential theory of the structure $\langle \mathcal{S}_\lambda : \times, \{\alpha\}_{\alpha \in \mathcal{S}_\lambda} \rangle$ is decidable.*

Proof. Indeed, this is an immediate consequence of Theorem 6 once we have observe the simple following relationship between the Cantor normal form and the prime factorization

$$\omega^{\lambda_r} a_r + \dots + \omega^{\lambda_1} a_1 + a_0 = a_0 (\omega^{\lambda_1} + 1) a_1 (\omega^{\lambda_2 - \lambda_1} + 1) \dots (\omega^{\lambda_r - \lambda_{r-1}} + 1) a_r$$

□

4 Open questions

Via the unicity of the factorization for ordinals, the monoid of successor ordinals bears a strong resemblance to the free monoid: if one ignores the finite ordinals, one is left with a free infinitely generated monoid. This explains why, in the end, thanks to the improvements in [10] and [4], we could resort to Makanin's result to obtain Theorem 13. What about the monoid of all ordinals? Or less ambitiously, what about solving equations in the monoid of ordinals with constants? More precisely, we are given an equation

$$L(x_1, \dots, x_n, \alpha_1, \dots, \alpha_p) = R(x_1, \dots, x_n, \alpha_1, \dots, \alpha_p)$$

where L and R are products of variables in $\{x_1, \dots, x_n\}$ and constants in $\{\alpha_1, \dots, \alpha_p\}$ which belong to an ordinal λ closed under multiplication. We are looking for solutions where all variables take on nonzero values.

Without Makanin's result, the authors do not know how to answer the question even in the specific case of the monoid of successor ordinals. But this is maybe no indication that solving equations in the structure $\langle \lambda; \times \rangle$ with arbitrary constants is at least as conceptually difficult as proving Makanin's result from scratch. We just make a couple of more or less trivial observations which tend to show that the similarity of solving solutions in two structures (finite free monoids and multiplicative ordinals) is maybe delusive. For example, even if all constants are successor ordinals, it might be the case that the equation has no solution in the monoid of successor ordinals but has a solution in λ , for example $2x = x$. More generally it is not difficult, but boring, to prove that given an equation where the constants are limit ordinals, it is decidable in polynomial time relative to the number of unknowns whether or not it has a solution where the unknowns are themselves limit ordinals. Of course the equation could have only solutions in the successor ordinals even if the constants were limit ordinals, see $\omega x = \omega(\omega + 1)$. More generally, considering specific submonoids Γ of λ for the constants such as the successors, the ω -powers, the limit ordinals etc . . . , and specific submonoids Ξ for the values assumed by the variables, one can investigate whether or not an equation with constants in Γ has a solution in Ξ .

References

- [1] A. Bès. Definability and decidability results related to the elementary theory of ordinal multiplication. In *Fund.Math.*, number 171, pages 197–211, 2002.
- [2] A. Bès and C. Choffrut. Complexity and (un)decidability results of fragments of $\langle \omega^{\omega^\lambda}; \times \rangle$. to appear In *Fund. Inf.*.
- [3] R. Cori and D. Perrin. Automates et Commutations Partielles. *in: ITA*, 19:1, 21–32, 1985.
- [4] V. Diekert and Y. V. Matiyasevich and A. Muscholl. Solving Word Equations modulo Partial Commutations, *Theor. Comput. Sci.*, 224(1-2), 215–235, 1999.
- [5] V. Diekert. Combinatorics on Traces, *Lecture Notes in Computer Science 454*, Springer, 1990.
- [6] V. Diekert and G. Rozenberg (Eds.). *The Book of Traces*, World Scientific, Singapore, 1995.

- [7] E. Jacobsthal. Über der Aufbau der Transfiniten Arithmetik. *Math. Ann.*, 66:145-194, 1909.
- [8] J. G. Rosenstein. *Linear ordering*. Academic Press, New-York, 1982.
- [9] W. Sierpinski. *Cardinal and Ordinal Numbers*. Warsaw: PWN, 1958.
- [10] K. U. Schulz. *Makanin's Algorithm for Word Equations - Two Improvements and a Generalization*, In *Word Equations and Related Topics, First International Workshop, IWVERT, Proceedings*, 85–150,1990.