

Ergodicity for Neutral Type SDEs with Infinite Length of Memory *

Jianhai Bao^{b),c)}, Feng-Yu Wang^{a),c)}, Chenggui Yuan^{c)}

^{a)}Center for Applied Mathematics, Tianjin University, Tianjin 300072, China

^{b)}School of Mathematics and Statistics, Central South University, Changsha 410083, China

^{c)}Department of Mathematics, Swansea University, Singleton Park, SA2 8PP, UK

jianhaibao@csu.edu.cn, wangfy@bnu.edu.cn, C.Yuan@swansea.ac.uk

March 3, 2022

Abstract

In this paper, the weak Harris theorem developed in [18] is illustrated by using a straightforward Wasserstein coupling, which implies the exponential ergodicity of the functional solutions to a range of neutral type SDEs with infinite length of memory. A concrete example is presented to illustrate the main result.

AMS Subject Classification: 34K50, 37A30, 60J05

Keywords: ergodicity, neutral type stochastic differential equation, infinite memory, weak Harris' theorem, Wasserstein coupling

1 Introduction

The ergodicity theory is a rich and active area in the study of Markov processes and related topics. Existing results include both qualitative characterizations (for instance, existence and uniqueness of invariant probability measures, strong Feller property, irreducibility) and quantitative estimates (convergence rate of Markov transition semigroups, gradient and heat kernel estimates, etc.). Among many other references, we would like to mention [9, 11, 15, 16, 23, 29] for the study of non-degenerate stochastic differential equations (SDEs) and stochastic partial differential equations (SPDEs), and [1, 12, 13, 14, 17, 22, 27, 31] for degenerate SDEs/SPDEs. In these references, several different probability

*This work is supported in part by NNSFC (11771326, 11431014, 11726627).

distances (for example, total variational distance, L^2 distance, and Wasserstein distance) have been adopted to measure the convergence rate of Markov transition semigroups. Efficient tools developed in the literature include functional inequalities (for instance, weak Poincaré, Poincaré, and log-Sobolev inequalities), Lyapunov type criteria, Harris' theorem, and coupling method, etc.

For path-dependent SDEs (i.e., the coefficients depend on the history), which are also called functional SDEs or SDEs with memory, the solutions are no longer Markovian. In this case, one investigates the functional solutions (i.e., the segment process, also called window process, of the solutions), which are Markov processes on the path space determined by the length of memory. However, the above tools mentioned are very hard to apply to this type infinite-dimensional Markov processes:

- Due to the lack of characterization on Dirichlet forms, functional inequalities are not yet established;
- The Lyapunov condition on the path space is less explicit since the formulation of infinitesimal generator is not yet available; on account of the same reason, the classical coupling argument via coupling operator is invalid;
- Since the functional solutions are highly degenerate (infinite-dimensional Markov processes with finite-dimensional noises), the classical Harris' theorem does not apply.

In recent years, some new approaches have been developed to investigate the ergodicity and related properties for path-dependent SDEs. When the noise term is path-independent and the drift depends only on a finite segment of path, the ergodicity under the total variational distance was investigated in [8], while gradient estimates and Harnack type inequalities (which in particular imply the strong Feller and irreducibility) have been established in [2, 3, 32, 30], to name a few, by using coupling by change of measures. When the noise part is also path-dependent, but both drift and noise parts depend only on a fixed length of past path, a weak Harris' theorem has been established in [18] to derive the exponential ergodicity under the Wasserstein distance. In case the noise is path-dependent, we would like to emphasize that the ergodicity under the total variational distance and the strong Feller property are not available since the laws of functional solutions with different initial data are mutually singular. The weak Harris' theorem has been applied in, e.g., [7, 10, 20, 26] to establish the ergodicity for highly degenerate stochastic dynamical systems including Markov processes with random switching.

In this paper, we aim to investigate the exponential ergodicity for path-dependent SDEs of neutral type and with infinite length of memory. Such kind of model fits more real world systems whose time evolution depends on the whole history (cf. [21, Chapter 6]). Intuitively, the farther the history, the weaker the influence to the evolution of the system. So, in the following we will take a reference norm on the path space which indicates that the influence of history exponentially decay when the time goes to $-\infty$.

For an integer $d \geq 1$, let $(\mathbb{R}^d, \langle \cdot, \cdot \rangle, |\cdot|)$ be the standard d -dimensional Euclidean space, and $\mathbb{R}^d \otimes \mathbb{R}^d$ the family of all $d \times d$ -matrices equipped with the Hilbert-Schmidt norm

$\|\cdot\|_{\text{HS}}$. $\mathcal{C} = C((-\infty, 0]; \mathbb{R}^d)$ stands for the space of all continuous maps $f : (-\infty, 0] \rightarrow \mathbb{R}^d$. For a map $f(\cdot) : (-\infty, \infty) \rightarrow \mathbb{R}^d$, define its segment map $f : [0, \infty) \rightarrow \mathcal{C}$ by setting

$$f_t(\theta) = f(t + \theta), \quad t \geq 0, \theta \in (-\infty, 0].$$

For a fixed number $r \in (0, \infty)$, let

$$\mathcal{C}_r = \left\{ \phi \in \mathcal{C} : \|\phi\|_r := \sup_{-\infty < \theta \leq 0} (e^{r\theta} |\phi(\theta)|) < \infty \right\}.$$

Then, $(\mathcal{C}_r, \|\cdot\|_r)$ is a Polish space. The norm $\|\cdot\|_r$ fits the intuition of exponential decay with regard to the influence of history; that is, the contribution to the norm from the history at time $\theta < 0$ has a minus exponential discount $e^{\theta r}$. For any $\theta \in (-\infty, 0]$, let $\xi_0(\theta) \equiv \mathbf{0}$, a d -dimensional zero vector.

Consider the following path-dependent SDE on \mathbb{R}^d of neutral type

$$(1.1) \quad d\{X(t) - G(X_t)\} = b(X_t)dt + \sigma(X_t)dW(t), \quad t > 0, \quad X_0 = \xi \in \mathcal{C}_r,$$

where $G, b : \mathcal{C}_r \rightarrow \mathbb{R}^d$ and $\sigma : \mathcal{C}_r \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d$ are measurable with $G(\xi_0) = \mathbf{0}$, $(X_t)_{t \geq 0}$ is the segment process associated with $(X(t))_{t \geq 0}$, $(W(t))_{t \geq 0}$ is the d -dimensional Brownian motion on a complete filtration probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$. For more motivating examples of (1.1), please refer to [21, p.201-2].

A continuous adapted process $(X(t))_{t \geq 0}$ is called a solution to (1.1) with the initial value X_0 , if \mathbb{P} -a.s.

$$X(t) = X(0) + G(X_t) - G(X_0) + \int_0^t b(X_s)ds + \int_0^t \sigma(X_s)dW(s), \quad t \geq 0.$$

We call $(X_t^\xi)_{t \geq 0}$ a functional solution to (1.1) with the initial value $X_0^\xi = \xi \in \mathcal{C}_r$.

To investigate the exponential convergence of $P_t(\xi, \cdot)$ under the metric \mathbb{W}_{ρ_r} , we impose the following assumptions.

(A0) b and σ are continuous and bounded on bounded subsets of \mathcal{C}_r , and there exists $\alpha \in (0, 1)$ such that

$$|G(\xi) - G(\eta)| \leq \alpha \|\xi - \eta\|_r, \quad \xi, \eta \in \mathcal{C}_r.$$

(A1) There exists a constant $L_0 > 0$ such that

$$\langle \xi(0) - \eta(0) + G(\eta) - G(\xi), b(\xi) - b(\eta) \rangle^+ + \|\sigma(\xi) - \sigma(\eta)\|_{\text{HS}}^2 \leq L_0 \|\xi - \eta\|_r^2, \quad \xi, \eta \in \mathcal{C}_r.$$

(A2) For each $\xi \in \mathcal{C}_r$, $\sigma(\xi)$ is invertible, and $\sup_{\xi \in \mathcal{C}_r} \{\|\sigma(\xi)\| + \|\sigma(\xi)^{-1}\|\} < \infty$;

(A3) There exists a continuous function $V : \mathcal{C}_r \rightarrow \mathbb{R}_+$ with $\lim_{\|\xi\|_r \rightarrow \infty} V(\xi) = \infty$ such that

$$P_t V(\xi) \leq K e^{-\gamma t} V(\xi) + K$$

holds for some constants $K, \gamma > 0$.

Theorem 1.1. *Let (A0) and (A1) hold. Then (1.1) has a unique solution. Moreover, there exists a constant $C > 0$ such that*

$$\mathbb{E}\|X_t^\xi\|_r^2 \leq C(1 + \|\xi\|_r^2)e^{Ct}, \quad t \geq 0, \quad \xi \in \mathcal{C}_r.$$

In the case $G \equiv \mathbf{0}$, the existence and uniqueness of solutions has been investigated in, e.g., [21, 24, 33] under the local Lipschitz and coercive conditions. When the length of memory is finite, the local Lipschitz condition was developed in [28] by the local weak monotone condition. Theorem 1.1 extends the result of [28] to path-dependent SDEs of neutral type and with infinite length of memory. Since we are interesting in ergodicity of the functional solutions, we are not going for giving a proof for this Theorem.

Next, we consider the ergodicity of the functional solutions to (1.1). In the situation of Theorem 1.1, for any $\xi \in \mathcal{C}_r$, let $(X^\xi(t))_{t \geq 0}$ be the unique solution to (1.1) with the initial datum $X_0^\xi = \xi$. Then, the functional solution $(X_t^\xi)_{t \geq 0}$ is a Markov process with the semigroup

$$P_t f(\xi) := \mathbb{E}f(X_t^\xi) = \int_{\mathcal{C}_r} f(\eta) P_t(\xi, d\eta), \quad t \geq 0, \quad f \in \mathcal{B}_b(\mathcal{C}_r), \quad \xi \in \mathcal{C}_r,$$

where $P_t(\xi, \cdot)$ is the distribution of X_t^ξ . As explained above, $P_t(\xi, \cdot)$ does not converges in the total variational distance. To investigate the ergodicity, we will take the Wasserstein distance induced by the distance

$$(1.2) \quad \rho_r(\xi, \eta) := 1 \wedge \|\xi - \eta\|_r, \quad \xi, \eta \in \mathcal{C}_r.$$

For any $\mu, \nu \in \mathcal{P}(\mathcal{C}_r)$, the collection of all probability measures on \mathcal{C}_r , the L^1 -Wasserstein distance between μ and ν induced by ρ_r is defined by

$$(1.3) \quad \mathbb{W}_{\rho_r}(\mu, \nu) = \inf_{\pi \in \mathcal{C}(\mu, \nu)} \int_{\mathcal{C}_r \times \mathcal{C}_r} \rho_r(\xi, \eta) \pi(d\xi, d\eta),$$

where $\mathcal{C}(\mu, \nu)$ is the set of all couplings of μ and ν ; that is, $\pi \in \mathcal{C}(\mu, \nu)$ if and only if π is a probability measure on $\mathcal{C}_r \times \mathcal{C}_r$ such that $\pi(\cdot \times \mathcal{C}_r) = \mu$ and $\pi(\mathcal{C}_r \times \cdot) = \nu$.

Condition (A1) with $G(\xi_0) = \mathbf{0}$ implies (H2) and (H3) and according to Theorem 1.1 yields the existence and uniqueness of (1.1). For the Lyapunov function V in (A3), let

$$\rho_{r,V}(\xi, \eta) := \sqrt{\rho_r(\xi, \eta)(1 + V(\xi) + V(\eta))}, \quad \xi, \eta \in \mathcal{C}_r.$$

Theorem 1.2. *Assume (H1) and (A1)-(A3). Then P_t has a unique invariant probability measure π , and there exist constants $c, \lambda > 0$ such that*

$$(1.4) \quad \mathbb{W}_{\rho_{r,V}}(\mu P_t, \nu P_t) \leq c e^{-\lambda t} \mathbb{W}_{\rho_{r,V}}(\mu, \nu), \quad \mu, \nu \in \mathcal{P}(\mathcal{C}_r), \quad t \geq 0.$$

Consequently, there exists a constant $C > 0$ such that

$$(1.5) \quad \mathbb{W}_{\rho_{r,V}}(P_t(\xi, \cdot), \pi) \leq C e^{-\lambda t} \sqrt{1 + V(\xi)}, \quad t \geq 0.$$

The proof of Theorem 1.2 is based on the weak Harris' theorem developed in [18]; see Theorem 2.1 below for more details. To meet the conditions of this theorem, one has to overcome the difficulties caused by the infinite length of memory.

Unlike conditions **(A1)** and **(A2)** which are explicitly imposed on the coefficients, the Lyapunov condition **(A3)** is set by means of the semigroup P_t which is less explicit. In many cases, one may verify **(A3)** by using the Lyapunov condition

$$(1.6) \quad \mathcal{L}V(\xi) \leq -\lambda V(\xi) + c, \quad \xi \in \mathcal{C}_r$$

for some constants $c, \lambda > 0$, where \mathcal{L} is the extended generator corresponding to the semigroup $(P_t)_{t \geq 0}$. However, as already explained before, \mathcal{L} is not yet available for the present model. Indeed, there are a number of examples satisfying **(A3)** but not (1.6); see, e.g., [5] for such an example on path-dependent SDEs without dissipativity. In this spirit, we present below explicit conditions for **(A3)**.

Proposition 1.3. *Let $\mu_0 \in \mathcal{P}((-\infty, 0])$ such that*

$$(1.7) \quad \delta_r(\mu_0) := \int_{-\infty}^0 e^{-2r\theta} \mu_0(d\theta) < \infty,$$

and set

$$(1.8) \quad \beta := \left(1 + \sqrt{\alpha_1 + \alpha_2 \delta_r(\mu_0)}\right)^2.$$

Then **(A3)** holds for $V(\xi) := \|\xi\|_r^2$ provided that the following two conditions hold:

(i) For any $\xi \in \mathcal{C}_r$, there exist constants $\alpha_1, \alpha_2 > 0$ with $\alpha_1 + \alpha_2 \delta_r(\mu_0) < 1$ such that

$$(1.9) \quad |G(\xi)|^2 \leq \alpha_1 |\xi(0)|^2 + \alpha_2 \int_{-\infty}^0 |\xi(\theta)|^2 \mu_0(d\theta).$$

(ii) There exist constants $c_0, \lambda_1, \lambda_2 > 0$ with $\gamma := \lambda_1 - 2r\beta - \lambda_2 \delta_r(\mu_0) > 0$ such that

$$(1.10) \quad 2\langle \xi(0) - G(\xi), b(\xi) \rangle + \|\sigma(\xi)\|_{\text{HS}}^2 \leq c_0 - \lambda_1 |\xi(0)|^2 + \lambda_2 \int_{-\infty}^0 |\xi(\theta)|^2 \mu_0(d\theta),$$

$$(1.11) \quad \|\sigma(\xi)\|_{\text{HS}}^2 \leq c_0 \left(1 + |\xi(0)|^2 + \int_{-\infty}^0 |\xi(\theta)|^2 \mu_0(d\theta)\right).$$

To conclude this section, we present below a concrete example to illustrate Theorem 1.2.

Example 1.4. Let $\mu_0(d\theta) = \frac{1}{r_0}e^{r_0\theta}d\theta \in \mathcal{P}((-\infty, 0])$ for some $r_0 > 2r$ and let

$$G(\xi) = \gamma_1 \int_{-\infty}^0 \xi(\theta)\mu_0(d\theta), \quad \sigma(\xi) = 1 + \gamma_2 \int_{-\infty}^0 (1 \wedge |\xi(\theta)|)\mu_0(d\theta),$$

$$b(\xi) = -\gamma_3\xi(0) - \gamma_4\left(\xi(0) - \gamma_1 \int_{-\infty}^0 \xi(\theta)\mu_0(d\theta)\right)^{\frac{1}{3}} + \gamma_5 \int_{-\infty}^0 \xi(\theta)\mu_0(d\theta)$$

for some constants $\gamma_i > 0, i = 1, \dots, 5$. If

(1.12)

$$\gamma_1^2 < r_0(r_0 - 2r) \quad \text{and} \quad 2\gamma_3 > 2r\left(1 + \frac{\gamma_1}{\sqrt{r_0(r_0 - 2r)}}\right)^2 + \frac{\gamma_2^2}{r_0(r_0 - 2r)} + \frac{2(\gamma_5 + \gamma_1\gamma_3)}{\sqrt{r_0(r_0 - 2r)}},$$

assertions in Theorem 1.2 hold.

The remainder of this paper is organized as follows. In Section 2, Theorem 1.2 is proved by using weak Harris' theorem and asymptotic coupling. In Section 4, we prove Proposition 1.3 and Example 1.4.

2 Proof of Theorem 1.2

For simplicity, we introduce the following notation. Let $f, g \in C(\mathbb{R}; \mathbb{R}^d)$, define

$$(2.1) \quad \Lambda^f(t) = f(t) - G(f_t), \quad \Lambda^{f,g}(t) = \Lambda^f(t) - \Lambda^g(t), \quad t \geq 0.$$

By **(A0)** and using $\varepsilon = \frac{\alpha}{1-\alpha}$, we obtain

$$|f(t) - g(t)|^2 \leq (1 + \varepsilon)|\Lambda^{f,g}(t)|^2 + (1 + \varepsilon^{-1})\alpha^2\|f_t - g_t\|_r^2 = \frac{|\Lambda^{f,g}(t)|^2}{1 - \alpha} + \alpha\|f_t - g_t\|_r^2.$$

When $f_0 = g_0$, this implies

$$e^{2rt}\|f_t - g_t\|_r^2 = \sup_{0 \leq s \leq t} (e^{2rs}|f(s) - g(s)|^2) \leq \frac{1}{1 - \alpha} \sup_{0 \leq s \leq t} (e^{2rs}|\Lambda^{f,g}(s)|^2) + \alpha e^{2rt}\|f_t - g_t\|_r^2,$$

so that

$$(2.2) \quad e^{2rt}\|f_t - g_t\|_r^2 \leq \frac{1}{(1 - \alpha)^2} \sup_{0 \leq s \leq t} (e^{2rs}|\Lambda^{f,g}(s)|^2), \quad f_0 = g_0.$$

Similarly,

$$(2.3) \quad e^{2rt}\|f_t\|_r^2 \leq \frac{1}{1 - \alpha}\|f_0\|_r^2 + \frac{1}{(1 - \alpha)^2} \sup_{0 \leq s \leq t} (e^{2rs}|\Lambda^f(s)|^2).$$

We shall complete the proof of Theorem 1.2 by the aid of weak Harris' theorem introduced in [18]. For readers' convenience, we state it below in details. We first recall some notions.

Definition 2.1. Let \mathbb{X} be a Polish space, and $(P_t)_{t \geq 0}$ a Markov semigroup with transition kernel $P_t(\xi, \cdot)$ on \mathbb{X} .

- (1) A continuous function $V : \mathbb{X} \rightarrow \mathbb{R}_+$ is called a Lyapunov function for $(P_t)_{t \geq 0}$, if there exist constants $\gamma, K > 0$ such that

$$(2.4) \quad P_t V(\xi) := \int_{\mathbb{X}} V(\eta) P_t(\xi, d\eta) \leq K e^{-\gamma t} V(\xi) + K, \quad \xi \in \mathbb{X}, \quad t \geq 0.$$

- (2) A function $\rho : \mathbb{X} \times \mathbb{X} \rightarrow [0, 1]$ is said to be distance-like if it is symmetric, lower semi-continuous, and $\rho(\xi, \eta) = 0$ if and only if $\xi = \eta$.
- (3) A set $A \subset \mathbb{X}$ is said to be ρ -small for P_t , if there exists $\varepsilon \in (0, 1)$ such that

$$\mathbb{W}_\rho(P_t(\xi, \cdot), P_t(\eta, \cdot)) \leq 1 - \varepsilon, \quad \xi, \eta \in A,$$

where \mathbb{W}_ρ is defined as in (1.3) for (\mathbb{X}, ρ) replacing (\mathcal{C}_r, ρ_r) .

- (4) ρ is said to be contractive for P_t , if there exists $\varepsilon \in (0, 1)$ such that

$$\mathbb{W}_\rho(P_t(\xi, \cdot), P_t(\eta, \cdot)) \leq \varepsilon \rho(\xi, \eta), \quad \xi, \eta \in \mathbb{X} \text{ with } \rho(\xi, \eta) < 1.$$

The following result is due to [18, Theorem 4.8].

Theorem 2.1. *Let ρ be a distance-like function on $\mathbb{X} \times \mathbb{X}$, and V a Lyapunov function such that (2.4) holds for some constants $\gamma, K > 0$. If there exists a constant $t^* > 0$ such that $\{V \leq 4K\}$ is ρ -small and ρ is contractive for P_{t^*} , then there exists a constant $t > 0$ such that*

$$\mathbb{W}_{\rho_V}(\mu P_t, \nu P_t) \leq \frac{1}{2} \mathbb{W}_{\rho_V}(\mu, \nu), \quad \forall \mu, \nu \in \mathcal{P}(\mathbb{X}),$$

where $\rho_V(\xi, \eta) := \sqrt{\rho(\xi, \eta)(1 + V(\xi) + V(\eta))}$, $\xi, \eta \in \mathbb{X}$.

To apply this result to the present model, for any $\delta > 0$ and $R > 0$, let

$$\rho_{r,\delta} = 1 \wedge (\delta^{-1} \rho_r), \quad B_R = \{\xi \in \mathcal{C}_r : \|\xi\|_r \leq R\},$$

$$(2.5) \quad t_{R,\delta} := 1 + \frac{1}{2r} \log \left(\frac{3}{2\delta^2} \left(\frac{2e^{2r}}{(1-\alpha)^2} \left(2R + \frac{\delta}{3} \right)^2 + \frac{R^2}{1-\alpha} \right) \right),$$

where ρ_r was given in (1.2). Obviously, the metric $\rho_{r,\delta}$ is equivalent to ρ_r . To check the conditions in Theorem 2.1 for the present setup, we need to prepare the following four lemmas concerned, respectively, with the (local) irreducibility, the continuity with respect to the initial variable, $\rho_{r,\delta}$ -small property, and $\rho_{r,\delta}$ -contractive property for the Markov transition kernel.

Lemma 2.2. *Under the conditions of Theorem 1.2, for any $R, \delta > 0$,*

$$(2.6) \quad \inf_{\xi \in B_R} \mathbb{P}(X_t^\xi \in B_\delta) > 0, \quad t \geq t_{R,\delta}.$$

Proof. The crucial point of the proof is to apply a standard result (e.g., [6, Lemma I.8.3]) that a uniform elliptic diffusion process is irreducible. So, below we will compare the radial process $|X^\xi|(s)$ with an elliptic diffusion process. For any $\xi \in B_R$ and $\delta > 0$, let $h \in C_b^\infty(\mathbb{R}_+; \mathbb{R}^d)$ such that

$$(2.7) \quad h(0) = \xi(0) - G(\xi) - \frac{\delta(1-\alpha)}{3}(1, 0, \dots, 0), \quad |h| \leq |h(0)|, \quad \text{and } h(s) = 0 \text{ for } s \geq 1,$$

where $\alpha \in (0, 1)$ was introduced in **(H1)**. According to (2.3), we have

$$(2.8) \quad (1-\alpha)^2 \sup_{0 \leq u \leq s} (e^{2ru} |X^\xi(u)|^2) \leq (1-\alpha) \|\xi\|_r^2 + \sup_{0 \leq u \leq s} (e^{2ru} |\Lambda^{X^\xi}(u)|^2), \quad s \geq 0,$$

in which Λ^{X^ξ} was defined as in (2.1) with $f = X^\xi$. Consider the following radial process

$$(2.9) \quad D(s) := |\Lambda^{X^\xi}(s) - h(s)|^2 - \frac{\delta^2(1-\alpha)^2}{9}, \quad s \geq 0.$$

By Itô's formula, it follows that

$$(2.10) \quad \begin{aligned} d(e^{2rs} D(s)) &= 2r e^{2rs} D(s) ds + e^{2rs} dD(s) \\ &= e^{2rs} \{ 2r D(s) + 2\langle \Lambda^{X^\xi}(s) - h(s), b(X_s^\xi) - h'(s) \rangle + \|\sigma(X_s^\xi)\|_{\text{HS}}^2 \} ds \\ &\quad + 2e^{2rs} \langle \Lambda^{X^\xi}(s) - h(s), \sigma(X_s^\xi) dW(s) \rangle, \quad s \geq 0. \end{aligned}$$

Define the stopping time

$$(2.11) \quad \tau = \inf \left\{ s \geq 0 : e^{2rs} |D(s)| \geq \frac{\delta^2(1-\alpha)^2}{18} \right\}.$$

Since $D(0) = 0$ and $D(s)$ is continuous with respect to s , we have $\mathbb{P}(\tau > 0) = 1$. In terms (2.9) and (2.11), we therefore have

$$\frac{\delta^2(1-\alpha)^2}{18} \geq e^{2rs} |D(s)| \geq |D(s)| \geq \frac{\delta^2(1-\alpha)^2}{9} - |\Lambda^{X^\xi}(s) - h(s)|^2, \quad s \in [0, \tau].$$

As a consequence, we arrive at

$$(2.12) \quad |\Lambda^{X^\xi}(s) - h(s)|^2 \geq \frac{\delta^2(1-\alpha)^2}{18}, \quad s \in [0, \tau].$$

Combining (2.12) with **(A2)**, we obtain from (2.10) that

$$(2.13) \quad \frac{d}{ds} \langle e^{2rs} D(s) \rangle = 4e^{4rs} |\sigma^*(X_s^\xi) (\Lambda^{X^\xi}(s) - h(s))|^2 \in [c_1, c_2], \quad s \in [0, \tau \wedge t]$$

for some constants $c_2 > c_1 > 0$. Herein $\langle \cdot \rangle$ means the quadratic variation of a continuous semi-martingale and $t \geq t_{R,\delta}$, which is to be fixed in what follows. Next, we are going to claim that (2.13) implies that

$$(2.14) \quad \mathbb{P}\left(\sup_{0 \leq u \leq s} (e^{2ru}|D(u)|) < \frac{\delta^2(1-\alpha)^2}{18}\right) > 0, \quad s \geq 0.$$

To achieve (2.14), we extend $(e^{2rs}D(s))_{s \in [0,\tau]}$ into $(e^{2rs}D(s))_{s \geq 0}$ in the following manner

$$(2.15) \quad Y(s) := e^{2r(s \wedge \tau)}D(s \wedge \tau) + \mathbf{1}_{\{s > \tau\}}(W^1(s) - W^1(\tau)), \quad s \geq 0,$$

where $(W^1(s))_{s \geq 0}$ stands for the first component of $(W(s))_{s \geq 0}$. Consequently, (2.13) gives that

$$\frac{d}{ds}\langle Y(s) \rangle \in [c_1 \wedge 1, c_2 \vee 1].$$

By [6, Lemma I.8.3] and using $Y(0) = 0$, this yields

$$(2.16) \quad \mathbb{P}\left(\sup_{0 \leq u \leq s} |Y(u)| < c\right) > 0, \quad sc > 0.$$

Combining this with (2.11) and (2.15), we obtain

$$\begin{aligned} \mathbb{P}\left(\sup_{0 \leq u \leq s} (e^{2ru}|D(u)|) < \frac{\delta^2(1-\alpha)^2}{18}\right) &= \mathbb{P}\left(\sup_{0 \leq u \leq s \wedge \tau} (e^{2ru}|D(u)|) < \frac{\delta^2(1-\alpha)^2}{18}, s < \tau\right) \\ &= \mathbb{P}\left(\sup_{0 \leq u \leq s} |Y(u)| < \frac{\delta^2(1-\alpha)^2}{18}\right) > 0. \end{aligned}$$

So, (2.14) holds true.

By using the fundamental inequality: $\frac{1}{2}|u|^2 - |v|^2 \leq |u - v|^2$, $u, v \in \mathbb{R}^d$, and recalling that $h(s) = 0$ for $s \geq 1$, we deduce from (2.8) and $\xi \in B_R$ that

$$\begin{aligned} &\mathbb{P}\left(\sup_{0 \leq s \leq t} (e^{2rs}|D(s)|) \leq \frac{\delta^2(1-\alpha)^2}{18}\right) \\ &\leq \mathbb{P}\left(\sup_{0 \leq s \leq t} (e^{2rs}|\Lambda^{X^\xi}(s) - h(s)|^2) \leq \frac{\delta^2(1-\alpha)^2}{6}e^{2rt}\right) \\ (2.17) \quad &\leq \mathbb{P}\left(\sup_{0 \leq s \leq t} \left(\frac{1}{2}e^{2rs}|\Lambda^{X^\xi}(s)|^2 - e^{2rs}|h(s)|^2\right) \leq \frac{\delta^2(1-\alpha)^2}{6}e^{2rt}\right) \\ &\leq \mathbb{P}\left(\sup_{0 \leq s \leq t} (e^{2rs}|\Lambda^{X^\xi}(s)|^2) \leq 2 \sup_{0 \leq s \leq 1} (e^{2rs}|h(s)|^2) + \frac{\delta^2(1-\alpha)^2}{3}e^{2rt}\right) \\ &\leq \mathbb{P}\left(\sup_{0 \leq s \leq t} (e^{2rs}|X^\xi(s)|^2) \leq \frac{1}{1-\alpha}R^2 + \frac{2}{(1-\alpha)^2} \sup_{0 \leq s \leq 1} (e^{2rs}|h(s)|^2) + \frac{\delta^2}{3}e^{2rt}\right). \end{aligned}$$

On the other hand, we observe that

$$\begin{aligned}
\mathbb{P}(X_t^\xi \in B_\delta) &= \mathbb{P}\left(e^{-2rt} \sup_{-\infty < s \leq t} (e^{2rs} |X^\xi(s)|^2) \leq \delta^2\right) \\
&= \mathbb{P}\left(\|\xi\|_r^2 \vee \sup_{0 \leq s \leq t} (e^{2rs} |X^\xi(s)|^2) \leq e^{2rt} \delta^2\right) \\
&\geq \mathbb{P}\left(\sup_{0 \leq s \leq t} (e^{2rs} |X^\xi(s)|^2) \leq e^{2rt} \delta^2\right).
\end{aligned}$$

Combining this with (2.14) and (2.17), it follows that

$$(2.18) \quad \frac{1}{1-\alpha} R^2 + \frac{2}{(1-\alpha)^2} \sup_{0 \leq s \leq 1} (e^{2rs} |h(s)|^2) + \frac{\delta^2}{3} e^{2rt} \leq e^{2rt} \delta^2, \quad t \geq t_{R,\delta}.$$

So it remains to prove (2.18). By (2.7), **(H1)**, $G(\xi_0) = \mathbf{0}$ and $\xi \in B_R$, we infer that

$$|h|^2 \leq |h(0)|^2 \leq \left(2R + \frac{\delta}{3}\right)^2,$$

which incurs

$$\sup_{0 \leq s \leq 1} (e^{2rs} |h(s)|^2) \leq e^{2r} \left(2R + \frac{\delta}{3}\right)^2.$$

Then (2.18) holds definitely provided

$$\frac{2\delta^2}{3} e^{2rt} \geq \frac{2e^{2r}}{(1-\alpha)^2} \left(2R + \frac{\delta}{3}\right)^2 + \frac{R^2}{1-\alpha},$$

which indeed is true for $t \geq t_{R,\delta}$. \square

Lemma 2.3. *Under conditions of Theorem 1.2, there exists a constant $K > 0$ such that*

$$(2.19) \quad \mathbb{E}\|X_t^\xi - X_t^\eta\|_r^2 \leq K e^{Kt} \|\xi - \eta\|_r^2, \quad t \geq 0, \quad \xi, \eta \in \mathcal{C}_r.$$

Proof. Let $\Lambda^{\xi,\eta}(t) = \Lambda^{X^\xi, X^\eta}(t)$ be defined by (2.1) with $f = X^\xi$ and $g = X^\eta$. Then, as in (2.2) we have

$$(2.20) \quad e^{2rt} \|X_t^\xi - X_t^\eta\|_r^2 \leq \frac{1}{1-\alpha} \|\xi - \eta\|_r^2 + \frac{1}{(1-\alpha)^2} \sup_{0 \leq s \leq t} (e^{2rs} |\Lambda^{\xi,\eta}(s)|^2).$$

Thus, to obtain the desired assertion (2.19), it is sufficient to show that

$$(2.21) \quad \Gamma(t) := \mathbb{E}\left(\sup_{0 \leq s \leq t} (e^{2rs} |\Lambda^{\xi,\eta}(s)|^2)\right) \leq J e^{Jt} \|\xi - \eta\|_r^2, \quad t \geq 0, \quad \xi, \eta \in \mathcal{C}_r$$

for some constant $J > 0$. Applying Itô's formula and using **(A1)** and **(A2)**, we obtain

$$\begin{aligned}
e^{2rt} |\Lambda^{\xi,\eta}(t)|^2 &\leq 4\|\xi - \eta\|_r^2 + \int_0^t e^{2rs} \{2r |\Lambda^{\xi,\eta}(s)|^2 + L_0 \|X_s^\xi - X_s^\eta\|_r^2\} ds \\
&\quad + 2 \int_0^t e^{2rs} \langle \Lambda^{\xi,\eta}(s), (\sigma(X_s^\xi) - \sigma(X_s^\eta)) dW(s) \rangle.
\end{aligned}$$

Combining this with (2.20), **(A2)** and BDG's inequality, we find out constants $c_1, c_2, c_3 > 0$ such that

$$\begin{aligned}
\Gamma(t) &\leq 4\|\xi - \eta\|_r^2 + c_1 \int_0^t \{\|\xi - \eta\|_r^2 + \Gamma(s)\} ds \\
&\quad + \mathbb{E} \left(\sup_{0 \leq s \leq t} (e^{2rs} |\Lambda^{\xi, \eta}(s)|^2) \int_0^t e^{2rs} \|\sigma(X_s^\xi) - \sigma(X_s^\eta)\|_{\text{HS}}^2 ds \right)^{1/2} \\
&\leq 4\|\xi - \eta\|_r^2 + c_1 \int_0^t \{\|\xi - \eta\|_r^2 + \Gamma(s)\} ds + \frac{1}{2}\Gamma(t) + c_2 \int_0^t \mathbb{E}(e^{2rs} \|X_s^\xi - X_s^\eta\|_r^2) ds \\
&\leq \frac{1}{2}\Gamma(t) + c_3(1+t)\|\xi - \eta\|_r^2 + c_3 \int_0^t \Gamma(s) ds.
\end{aligned}$$

Consequently,

$$\Gamma(t) \leq 2c_3(1+t)\|\xi - \eta\|_r^2 + 2c_3 \int_0^t \Gamma(s) ds.$$

By Gronwall's inequality, we obtain

$$\Gamma(t) \leq 2c_3(1+t)e^{2c_3 t} \|\xi - \eta\|_r^2.$$

Therefore, (2.21) follows from (2.20) immediately. \square

Lemma 2.4. *Under conditions of Theorem 1.2, for any $R, \delta > 0$,*

$$(2.22) \quad \mathbb{W}_{\rho_r, \delta}(P_t(\xi, \cdot), P_t(\eta, \cdot)) \leq 1 - \frac{\alpha_t^2}{2} < 1, \quad t \geq t_{R, \delta/4}, \quad \xi, \eta \in B_R,$$

holds for $t_{R, \delta/4}$ in (2.5) and $\alpha_t := \inf_{\xi \in B_R} \mathbb{P}(X_t^\xi \in B_{\delta/4})$.

Proof. For any $\xi, \eta \in B_R$, let $(X_t^\xi)_{t \geq 0}$ be the functional solution to (1.1) with the initial value $X_0^\xi = \xi \in \mathcal{C}_r$, and $(\tilde{X}_t^\eta)_{t \geq 0}$ the functional solution to (1.1) with the initial datum $\tilde{X}_0^\eta = \eta$ but for an independent Brownian motion $(\tilde{W}(t))_{t \geq 0}$ replacing $(W(t))_{t \geq 0}$. We call $(X_t^\xi, \tilde{X}_t^\eta)$ an independent coupling of the functional solutions to (1.1). In view of the independence of $(X_t^\xi)_{t \geq 0}$ and $(\tilde{X}_t^\eta)_{t \geq 0}$, we deduce that

$$\begin{aligned}
\mathbb{W}_{\rho_r, \delta}(P_t(\xi, \cdot), P_t(\eta, \cdot)) &\leq \mathbb{E}(1 \wedge (\delta^{-1} \|X_t^\xi - X_t^\eta\|_r)) \\
&\leq \frac{1}{2} \mathbb{P}(X_t^\xi \in B_{\delta/4}, X_t^\eta \in B_{\delta/4}) + \mathbb{P}(\{X_t^\xi \notin B_{\delta/4}\} \cup \{X_t^\eta \notin B_{\delta/4}\}) \\
&= 1 - \frac{1}{2} \mathbb{P}(X_t^\xi \in B_{\delta/4}) \mathbb{P}(X_t^\eta \in B_{\delta/4}) \\
&\leq 1 - \frac{\alpha_t^2}{2}.
\end{aligned}$$

Hence, (2.22) holds true due to Lemma 2.2. \square

Lemma 2.5. *Under the conditions of Theorem 1.2, for any $\beta \in (0, 1)$ there exist constants $\delta_\beta, t_\beta > 0$ such that*

$$(2.23) \quad \mathbb{W}_{\rho_{r,\delta}}(P_t(\xi, \cdot), P_t(\eta, \cdot)) \leq \beta \rho_{r,\delta}(\xi, \eta), \quad t \geq t_\beta, \delta \in (0, \delta_\beta]$$

for any $\xi, \eta \in \mathcal{C}_r$ with $\rho_{r,\delta}(\xi, \eta) < 1$.

Proof. Our proof is based on the Girsanov transform and Warsserstein coupling, which is more straightforward than the “binding construction” argument adopted in [18, p.254-257]. For $\xi, \eta \in \mathcal{C}_r$, let $(X_s^\xi)_{s \geq 0}$ be the functional solution to (1.1), and let $(Y_s^\eta)_{s \geq 0}$ solve the following SDE

$$(2.24) \quad d\{Y_s^\eta - G(Y_s^\eta)\} = \{b(Y_s^\eta) + \lambda \Lambda^{\xi,\eta}(s)\} ds + \sigma(Y_s^\eta) dW(s), \quad s \geq 0, \quad Y_0^\eta = \eta,$$

where $\lambda > 0$ is a constant, $\Lambda^{\xi,\eta}(s) := \Lambda^{X_s^\xi, Y_s^\eta}(s)$ is defined in (2.1). For $\lambda > 0$ sufficiently large, the additional drift $\lambda \Lambda^{\xi,\eta}(s)$ strongly pushes Y_s^η moving toward to X_s^ξ whenever $s \uparrow \infty$. Indeed, when $\lambda > 0$ is sufficiently large, for any $r_0 \in (0, r)$, there exists a constant $c > 0$ such that

$$(2.25) \quad \mathbb{E} \|X_s^\xi - Y_s^\eta\|_r^2 \leq c e^{-r_0 s} \|\xi - \eta\|_r^2, \quad s \geq 0, \xi, \eta \in \mathcal{C}_r,$$

and, for any stopping time τ ,

$$(2.26) \quad \mathbb{E} \|X_{s \wedge \tau}^\xi - Y_{s \wedge \tau}^\eta\|_r^2 \leq c \|\xi - \eta\|_r^2, \quad s \geq 0, \xi, \eta \in \mathcal{C}_r$$

as shown in the proof of [4, (3.11)],

To compare Y_s^η with X_s^η via the Girsanov theorem, let $h(s) = \lambda \sigma^{-1}(Y_s^\eta) \Lambda^{\xi,\eta}(s)$ and set

$$R_t := \exp \left(- \int_0^t \langle h(s), dW(s) \rangle - \frac{1}{2} \int_0^t |h(s)|^2 ds \right).$$

Generally, R_t may not be a well defined probability density, so we shall restrict it by the following stopping time

$$\tau_\varepsilon = \inf \left\{ s \geq 0 : \int_0^s |h(s)|^2 ds \geq \varepsilon^{-1} \|\xi - \eta\|_r^2 \right\}$$

for some constant $\varepsilon \in (0, 1)$ sufficiently small to be determined later. By Girsanov theorem, $d\mathbb{Q}_\varepsilon := R_{t \wedge \tau_\varepsilon} d\mathbb{P}$ is a probability measure on (Ω, \mathcal{F}) under which

$$\widetilde{W}(s) := W(s) + \int_0^{s \wedge \tau_\varepsilon} h(u) du, \quad s \geq 0$$

is a d -dimensional Brownian motion. Let $\widetilde{Y}^\eta(s)$ solve the SDE

$$d\{\widetilde{Y}^\eta(s) - G(\widetilde{Y}^\eta)\} = \{b(\widetilde{Y}^\eta) + \mathbf{1}_{\{\tau_\varepsilon \geq s\}} \lambda \widetilde{\Lambda}^{\xi,\eta}(s)\} ds + \sigma(\widetilde{Y}^\eta) dW(s), \quad s \geq 0, \quad \widetilde{Y}_0^\eta = \eta,$$

where $\tilde{\Lambda}^{\xi, \eta}(s) := \Lambda^{X^\xi, \tilde{Y}^\eta}(s)$. By the weak uniqueness of solutions to (2.24) up to time $t \wedge \tau_\varepsilon$, we have

$$(2.27) \quad \mathbb{P}(X_t^\eta \in \cdot) = \mathbb{Q}_\varepsilon(\tilde{Y}_t^\eta \in \cdot), \quad \tilde{Y}_{t \wedge \tau_\varepsilon}^\eta = Y_{t \wedge \tau_\varepsilon}^\eta, \quad t \geq 0.$$

To estimate $\mathbb{W}_{\rho_{r, \delta}}(P_t(\xi, \cdot), P_t(\eta, \cdot))$, we take the following Wasserstein coupling of \mathbb{P} and \mathbb{Q}_ε :

$$(2.28) \quad \begin{aligned} \Pi(d\omega, d\tilde{\omega}) &= (1 \wedge R_{t \wedge \tau_\varepsilon})(\omega) \mathbb{P}(d\omega) \delta_\omega(d\tilde{\omega}) \\ &+ \frac{(1 - R_{t \wedge \tau_\varepsilon})^+(\omega) (R_{t \wedge \tau_\varepsilon} - 1)^+(\tilde{\omega})}{\mathbb{E}[(1 - R_{t \wedge \tau_\varepsilon})^+]} \mathbb{P}(d\omega) \mathbb{P}(d\tilde{\omega}), \end{aligned}$$

where δ_ω is the Dirac measure at point ω , and the last term vanishes if $\mathbb{E}[(1 - R_{t \wedge \tau_\varepsilon})^+] = 0$ which is only possible when $\xi = \eta$. Combining this coupling with (2.27), and noting that $\rho_{r, \delta} \leq 1$, we obtain that

$$(2.29) \quad \begin{aligned} &\mathbb{W}_{\rho_{r, \delta}}(P_t(\xi, \cdot), P_t(\eta, \cdot)) \\ &\leq \int_{\Omega \times \Omega} \rho_{r, \delta}(X_t^\xi(\omega), \tilde{Y}_t^\eta(\tilde{\omega})) \Pi(d\omega, d\tilde{\omega}) \\ &\leq \mathbb{E}[\rho_{r, \delta}(X_t^\xi, \tilde{Y}_t^\eta)(1 \wedge R_{t \wedge \tau_\varepsilon})] + \mathbb{E}[(R_{t \wedge \tau_\varepsilon} - 1)^+] \\ &\leq \mathbb{E}[\mathbf{1}_{\{t \leq \tau_\varepsilon\}} \rho_{r, \delta}(X_t^\xi, Y_t^\eta)] + \mathbb{E}[\mathbf{1}_{\{t > \tau_\varepsilon\}} \rho_{r, \delta}(X_t^\xi, \tilde{Y}_t^\eta)] + \mathbb{E}[(R_{t \wedge \tau_\varepsilon} - 1)^+] \\ &=: I_1(t) + I_2(t) + I_3(t). \end{aligned}$$

Next we are going to estimate three terms above, one-by-one.

Firstly, by (2.25), there exists a constant $c > 0$ such that

$$(2.30) \quad I_1(t) \leq c e^{-r_0 t/2} \delta^{-1} \|\xi - \eta\|_r = c e^{-r_0 t/2} \rho_{r, \delta}(\xi, \eta)$$

for arbitrary $\xi, \eta \in \mathcal{C}_r$ with $\rho_{r, \delta}(\xi, \eta) < 1$. Next, by Hölder's inequality, the strong Markov property, (2.19), (2.25), (2.26), $\tilde{Y}_{t \wedge \tau_\varepsilon}^\eta = Y_{t \wedge \tau_\varepsilon}^\eta$ due to (2.27), and noting that the SDE for \tilde{Y}_s^η coincides with (1.1) when $s \geq \tau_\varepsilon$, we obtain that

$$(2.31) \quad \begin{aligned} I_2(t) &\leq \delta^{-1} \mathbb{E}[\|X_t^\xi - \tilde{Y}_t^\eta\|_r \mathbf{1}_{\{\tau_\varepsilon < t\}}] \\ &= \delta^{-1} \mathbb{E}[\mathbf{1}_{\{\tau_\varepsilon < t\}} \{\mathbb{E}[\|X_{t-\tau_\varepsilon}^{\xi'} - \tilde{Y}_{t-\tau_\varepsilon}^{\eta'}\|_r] \mid (\xi', \eta') = (X_{t \wedge \tau_\varepsilon}^\xi, Y_{t \wedge \tau_\varepsilon}^\eta)\}] \\ &\leq \delta^{-1} \sqrt{\mathbb{P}(\tau_\varepsilon < t)} K e^{Kt} \mathbb{E}[\|X_{t \wedge \tau_\varepsilon}^\xi - Y_{t \wedge \tau_\varepsilon}^\eta\|_r^2] \\ &\leq \sqrt{c K e^{Kt} \mathbb{P}(\tau_\varepsilon < t)} \rho_{r, \delta}(\xi, \eta) \end{aligned}$$

for any $\xi, \eta \in \mathcal{C}_r$ with $\rho_{r, \delta}(\xi, \eta) < 1$. On the other hand, Chebyshev's inequality, **(H1)**, **(A1)**, **(A2)** and (2.25) imply

$$(2.32) \quad \begin{aligned} \mathbb{P}(\tau_\varepsilon < t) &\leq \mathbb{P}\left(\int_0^t |h(s)|^2 ds \geq \varepsilon^{-1} \|\xi - \eta\|_r^2\right) \\ &\leq c_1 \varepsilon \|\xi - \eta\|_r^{-2} \int_0^t \mathbb{E}[\|X_s^\xi - Y_s^\eta\|_r^2] ds \\ &\leq c_2 \varepsilon \int_0^t e^{-r_0 s} ds \leq \frac{c_2 \varepsilon}{r_0} \end{aligned}$$

for some constants $c_1, c_2 > 0$. Combining (2.32) with (2.31), we may find out a constant $c_3 > 0$ such that

$$(2.33) \quad I_2(t) \leq c_3 \sqrt{\varepsilon} e^{c_3 t} \rho_{r,\delta}(\xi, \eta), \quad \xi, \eta \in \mathcal{C}_r, \quad \rho_{r,\delta}(\xi, \eta) < 1.$$

By Hölder's inequality and the definition of the stopping time τ_ε , we obtain

$$\begin{aligned} I_3^2(t) &\leq \mathbb{E} R_{t \wedge \tau_\varepsilon}^2 - 1 \\ &\leq \mathbb{E} \exp \left(-2 \int_0^{t \wedge \tau_\varepsilon} \langle h(s), dW(s) \rangle - \int_0^{t \wedge \tau_\varepsilon} |h(s)|^2 ds \right) - 1 \\ &\leq \left(\mathbb{E} \exp \left(6 \int_0^{t \wedge \tau_\varepsilon} |h(s)|^2 ds \right) \right)^{1/2} - 1 \\ &\leq e^{3\varepsilon^{-1} \|\xi - \eta\|_r^2} - 1 \leq 3\varepsilon^{-1} \|\xi - \eta\|_r^2 e^{3\varepsilon^{-1} \|\xi - \eta\|_r^2}, \end{aligned}$$

where the last step is due to the inequality: $e^x - 1 \leq x e^x$, $x \geq 0$. Hence, for any $\xi, \eta \in \mathcal{C}_r$ with $\rho_{r,\delta}(\xi, \eta) < 1$, we infer that

$$I_3(t) \leq \sqrt{3\varepsilon}^{-1/2} e^{\frac{3}{2}\varepsilon^{-1}\delta^2} \|\xi - \eta\|_r = \sqrt{3\varepsilon}^{-1/2} \delta e^{\frac{3}{2}\varepsilon^{-1}\delta^2} \rho_{r,\delta}(\xi, \eta).$$

Combining this with (2.29), (2.30) and (2.33), we arrive at

$$\mathbb{W}_{\rho_{r,\delta}}(P_t(\xi, \cdot), P_t(\eta, \cdot)) \leq c_3 \left\{ e^{-r_0 t/2} + e^{Kt/2} \varepsilon + \varepsilon^{-1/2} \delta e^{\frac{3}{2}\varepsilon^{-1}\delta^2} \right\} \rho_{r,\delta}(\xi, \eta)$$

for any $\xi, \eta \in \mathcal{C}_r$ with $\rho_{r,\delta}(\xi, \eta) < 1$. Thus, (2.23) holds by taking $t > 0$ sufficiently large and $\varepsilon = \delta \in (0, 1)$ sufficiently small. \square

Proof of Theorem 1.2. Since $\lim_{\|\xi\|_r \rightarrow \infty} V(\xi) = \infty$, there is a constant $R > 0$ such that $\{V \leq 4K\} \subset B_R$. By Lemmas 2.3 and 2.4, there exists $t_0 \geq t_{R,\delta/4}$ such that $\{V \leq 4K\}$ is $\rho_{r,\delta}$ -small and $\rho_{r,\delta}$ is contractive for P_t for any $t \geq t_0$ and $\delta > 0$. So, in terms of Theorem 2.1, P_t has a unique probability measure π , and there exists a constant $t_1 > 0$ such that

$$(2.34) \quad \mathbb{W}_{\rho_{r,\delta,V}}(\mu P_{t_1}, \nu P_{t_1}) \leq \frac{1}{2} \mathbb{W}_{\rho_{r,\delta,V}}(\mu, \nu), \quad \mu, \nu \in \mathcal{P}(\mathcal{C}_r).$$

Combining this with the semigroup property, to prove (1.4) it suffices to find out a constant $C > 0$ such that

$$(2.35) \quad \mathbb{W}_{\rho_{r,\delta,V}}(\delta_\xi P_t, \delta_\eta P_t) \leq C \rho_{r,\delta,V}(\xi, \eta), \quad t \in [0, t_1], \xi, \eta \in \mathcal{C}_r.$$

By (2.4) and (2.19), there exists from Hölder's inequality a constant $C > 0$ such that

$$\begin{aligned} \mathbb{W}_{\rho_{r,\delta,V}}(\delta_\xi P_t, \delta_\eta P_t) &\leq \mathbb{E} \sqrt{\rho_{r,\delta}(X_t^\xi, X_t^\eta) (1 + V(X_t^\xi) + V(X_t^\eta))} \\ &\leq \sqrt{\mathbb{E} \rho_{r,\delta}(X_t^\xi, X_t^\eta) \mathbb{E} (1 + V(X_t^\xi) + V(X_t^\eta))} \\ &\leq C \sqrt{\rho_r(\xi, \eta) (1 + V(\xi) + V(\eta))} \\ &= C \rho_{r,\delta,V}(\xi, \eta) \end{aligned}$$

for any $t \in [0, t_1]$, $\xi, \eta \in \mathcal{C}_r$. Therefore, (2.35) holds true so that (1.4) is available by in addition taking the equivalence of ρ_r and $\rho_{r,\delta}$.

Next, by (2.4), we have $\pi(V) := \int_{\mathcal{C}_r} V d\pi < \infty$, so that (1.4) implies

$$\mathbb{W}_{\rho_{r,V}}(P_t(\xi, \cdot), \pi) = \mathbb{W}_{\rho_{r,V}}(\delta_\xi P_t, \pi P_t) \leq c e^{-\lambda t} \int_{\mathcal{C}_r} \rho_{r,V}(\xi, \eta) \pi(d\eta) \leq C e^{-\lambda t} \sqrt{1 + V(\xi)}$$

for some constant $C > 0$. □

3 Proofs of Proposition 1.3 and Example 1.4

Proof of Proposition 1.3. For simplicity, we write $X(t) = X^\xi(t)$ and $X_t = X_t^\xi$. By (2.3), it is sufficient to find out a constant $c > 0$ such that

$$(3.1) \quad \mathbb{E} \left(\sup_{0 \leq s \leq t} (e^{2rs} |\Lambda^X(s)|^2) \right) \leq c \left((1+t) \|\xi\|_r^2 + e^{2rt} \right).$$

By Fubini's theorem and integration by substitution, we deduce from (1.7) that

$$(3.2) \quad \begin{aligned} & \int_0^t \int_{-\infty}^0 e^{2rs} |X(s+\theta)|^2 \mu_0(d\theta) ds \\ &= \int_0^t \int_{-\infty}^{-s} e^{-2r\theta} e^{2r(s+\theta)} |X(s+\theta)|^2 \mu_0(d\theta) ds + \int_{-t}^0 e^{-2\theta} \int_0^{t+\theta} e^{2rs} |X(s)|^2 ds \mu_0(d\theta) \\ &\leq \delta_r(\mu_0) \|\xi\|_r^2 t + \delta_r(\mu_0) \int_0^t e^{2rs} |X(s)|^2 ds, \end{aligned}$$

which, together with (1.9), leads to: for any $\varepsilon > 0$, there exists a constant $c_\varepsilon > 0$ such that

$$\begin{aligned} \int_0^t e^{2rs} |\Lambda^X(s)|^2 ds &\leq \left(1 + \varepsilon + (1 + 1/\varepsilon) \alpha_1 \right) \int_0^t e^{2rs} |X(s)|^2 ds \\ &\quad + (1 + 1/\varepsilon) \alpha_2 \int_0^t \int_{-\infty}^0 e^{2rs} |X(s+\theta)|^2 \mu_0(d\theta) ds \\ &\leq c_\varepsilon \|\xi\|_r^2 t + \left(1 + \alpha_1 + \alpha_2 \delta_r(\mu_0) + \varepsilon + (\alpha_1 + \alpha_2 \delta_r(\mu_0)) / \varepsilon \right) \int_0^t e^{2rs} |X(s)|^2 ds. \end{aligned}$$

Taking $\varepsilon = (\alpha_1 + \alpha_2 \delta_r(\mu_0))^{1/2}$, we find out a constant $c_1 > 0$ such that

$$(3.3) \quad \int_0^t e^{2rs} |\Lambda^X(s)|^2 ds \leq c_1 \|\xi\|_r^2 t + \beta_1 \int_0^t e^{2rs} |X(s)|^2 ds$$

where $\beta_1 > 0$ is in (1.8). Now, by Itô's formula, it follows from (1.10) that

$$(3.4) \quad \begin{aligned} e^{2rt} |\Lambda^X(t)|^2 &\leq |\Lambda^\xi(0)|^2 + \int_0^t e^{2rs} \left\{ c_0 + 2r |\Lambda^X(s)|^2 - \lambda_1 |X(s)|^2 \right. \\ &\quad \left. + \lambda_2 \int_{-\infty}^0 |X(s+\theta)|^2 \mu_0(d\theta) \right\} ds + 2 \int_0^t e^{2rs} \langle \Lambda^X(s), \sigma(X_s) dW(s) \rangle. \end{aligned}$$

Plugging (3.2) and (3.3) into (3.4) and utilizing **(H1)** gives

$$e^{2rt}\mathbb{E}|X(t)|^2 \leq c_2\left((1+t)\|\xi\|_r^2 + e^{2rt}\right) - \left(\lambda_1 - 2r\beta_1 - \lambda_2\delta_r(\mu_0)\right) \int_0^t e^{2rs}\mathbb{E}|X(s)|^2 ds$$

for some constant $c_2 > 0$. Since $\lambda_1 - 2r\beta_1 - \lambda_2\delta_r(\mu_0) > 0$, this implies

$$(3.5) \quad \int_0^t e^{2rs}\mathbb{E}|X(s)|^2 ds \leq c_3(1+t)\|\xi\|_r^2 + c_3e^{2rt}$$

for some constant $c_3 > 0$. On the other hand, by BDG's inequality, we deduce from (1.11) and (3.2) that

$$(3.6) \quad \begin{aligned} & 2 \sup_{0 \leq s \leq t} \left| \int_0^s e^{2ru} \langle \Lambda^X(u), \sigma(X_u) dW(u) \rangle \right| \\ & \leq 8\sqrt{2} \mathbb{E} \left(\sup_{0 \leq s \leq t} (e^{2rs} |\Lambda^X(s)|^2) \int_0^t e^{2rs} \|\sigma(X_s)\|_{\text{HS}}^2 ds \right)^{1/2} \\ & \leq \frac{1}{2} \mathbb{E} \left(\sup_{0 \leq s \leq t} (e^{2rs} |\Lambda^X(s)|^2) \right) + c_4 \left(e^{2rt} + t \|\xi\|_r^2 + \int_0^t e^{2rs} \mathbb{E}|X(s)|^2 ds \right) \end{aligned}$$

for some $c_4 > 0$. Thus, by taking (3.3) and (3.4) into account and making use of (3.6) and **(H1)**, there exists a constant $c_5 > 0$ such that

$$\mathbb{E} \left(\sup_{0 \leq s \leq t} (e^{2rs} |\Lambda^X(s)|^2) \right) \leq c_5 \left((1+t)\|\xi\|_r^2 + e^{2rt} + \int_0^t e^{2rs} \mathbb{E}|X(s)|^2 ds \right).$$

Henceforth, (3.1) follows directly from (3.5). \square

Proof of Example 1.4. It suffices to verify **(H1)**, **(A1)**-**(A2)** and conditions in Proposition 1.3.

By Hölder's inequality, one has

$$(3.7) \quad |G(\xi) - G(\eta)|^2 \leq \gamma_1^2 \int_{-\infty}^0 |\xi(\theta) - \eta(\theta)|^2 \mu_0(d\theta) \leq \gamma_1^2 \delta_r(\mu_0) \|\xi - \eta\|_r^2, \quad \xi, \eta \in \mathcal{C}_r.$$

Therefore, **(H1)** holds for $\alpha = \gamma_1 \sqrt{\delta_r(\mu_0)} < 1$ owing to (1.12). By Hölder's inequality and (3.7), we can find some constants $c_1, c_2 > 0$ such that

$$\begin{aligned} & \langle \xi(0) - \eta(0) - (G(\xi) - G(\eta)), b(\xi) - b(\eta) \rangle^+ + |\sigma(\xi) - \sigma(\eta)|^2 \\ & \leq \left((\xi(0) - \eta(0) - (G(\xi) - G(\eta))) \left\{ -\gamma_3(\xi(0) - \eta(0)) + \gamma_5 \int_{-\infty}^0 (\xi(\theta) - \eta(\theta)) \mu_0(d\theta) \right\} \right)^+ \\ & \quad + \gamma_2^2 \left(\int_{-\infty}^0 |\xi(\theta) - \eta(\theta)| \mu_0(d\theta) \right)^2 \\ & \leq c_1 \left\{ |\xi(0) - \eta(0)|^2 + |G(\xi) - G(\eta)|^2 \right\} \\ & \leq c_2 \|\xi - \eta\|_r^2, \quad \xi, \eta \in \mathcal{C}_r, \end{aligned}$$

where the first inequality is due to

$$-\gamma_4 \langle \xi(0) - G(\xi) - (\eta(0) - G(\eta)), (\xi(0) - G(\xi))^{1/3} - (\eta(0) - G(\eta))^{1/3} \rangle \leq 0.$$

Consequently, **(A1)** holds true. According to the formula of $\sigma(\xi)$, **(A1)** holds trivially.

Finally, we verify conditions in Proposition 1.3. Obviously, (1.7) holds with $\delta_r(\mu_0) = \frac{1}{r_0(r_0-2r)} < \infty$ due to $r_0 > 2r$. Thanks to $G(\xi_0) = \mathbf{0}$ and (3.7), (1.9) holds for $\alpha_1 = 0$ and $\alpha_2 = \gamma_1^2$. By Hölder's inequality, it follows that

$$(3.8) \quad |\sigma(\xi)|^2 \leq 1 + \frac{1}{\alpha} + (1 + \alpha)\gamma_2^2 \int_{-\infty}^0 |\xi(\theta)|^2 \mu_0(d\theta), \quad \alpha > 0, \xi \in \mathcal{C}_r.$$

Next, by using (3.8), for any $\varepsilon > 0$ and $\xi \in \mathcal{C}_r$,

$$\begin{aligned} & 2(\xi(0) - G(\xi))b(\xi) + |\sigma(\xi)|^2 \\ & \leq 1 + \frac{1}{\alpha} - 2\gamma_3\xi^2(0) + 2(\gamma_5 + \gamma_1\gamma_3)\xi(0) \int_{-\infty}^0 \xi(\theta)\mu_0(d\theta) + (1 + \alpha)\gamma_2^2 \int_{-\infty}^0 |\xi(\theta)|^2 \mu_0(d\theta) \\ & \leq 1 + \frac{1}{\alpha} - \left(2\gamma_3 - (\gamma_5 + \gamma_1\gamma_3)\varepsilon\right)\xi^2(0) + \left(\frac{1}{\varepsilon}(\gamma_5 + \gamma_1\gamma_3) + (1 + \alpha)\gamma_2^2\right) \int_{-\infty}^0 |\xi(\theta)|^2 \mu_0(d\theta). \end{aligned}$$

Taking $\varepsilon = \sqrt{\delta_r(\mu_0)}$ and $\alpha \in (0, 1)$ sufficiently small and combining (1.12), we conclude that (1.10) holds. □

References

- [1] Banerjee, S., Gordina, M., Mariano, P., Coupling in the Heisenberg group and its applications to gradient estimates, arXiv:1610.06430v3.
- [2] Bao, J., Wang, F.-Y., Yuan, C., Derivative formula and Harnack inequality for degenerate functional SDEs, *Stoch. Dyn.*, **13** (2013), 1250013, 22pp.
- [3] Bao, J., Wang, F.-Y., Yuan, C., *Bismut Formulae and Applications for Functional SPDEs*, *Bull. Math. Sci.*, **137** (2013), 509–522.
- [4] Bao, J., Wang, F.-Y., Yuan, C., Asymptotic Log-Harnack Inequality and Applications for Stochastic Systems of Infinite Memory, arXiv: 1710.01042v1.
- [5] Bao, J., Yin, G., Yuan, C., Stationary distributions for retarded stochastic differential equations without dissipativity, *Stochastics*, **89** (2017), 530–549.
- [6] Bass, R., *Diffusions and elliptic operators*, Springer, New York, 1998.
- [7] Butkovsky, O., Subgeometric rates of convergence of Markov processes in the Wasserstein metric, *Ann. Appl. Probab.*, **24** (2014), 526–552.
- [8] Butkovsky, O., Scheutzow, M., Invariant measures for stochastic functional differential equations, *Electron. J. Probab.*, **22** (2017), 1–23
- [9] Chen, M.-F., Li, S., Coupling methods for multidimensional diffusion processes, *Ann. Probab.*, **17** (1989), 151–177.

- [10] Cloez, B. and Hairer, M.: Exponential ergodicity for Markov processes with random switching, *Bernoulli*, **21** (2015), 505–536.
- [11] Da Prato, G., Zabczyk, J., Ergodicity for Infinite-Dimensional Systems, in: London Mathematical Society, Lecture Note Series, vol. 229, Cambridge University Press, Cambridge, 1996.
- [12] Dolbeault, J., Mouhot, C., Schmeiser, C., Hypocoercivity for linear kinetic equations conserving mass, *Trans. Amer. Math. Soc.*, **367** (2015), 3807–3828.
- [13] Grothaus, M., Stilgenbauer, P., Hypocoercivity for kolmogorov backward evolution equations and applications, *J. Funct. Anal.*, **267** (2014), 3515–3556.
- [14] Grothaus, M., Wang, F. -Y., Weak Poincaré inequalities for convergence rate of degenerate diffusion processes, arXiv:1703.04821.
- [15] Hairer, M., Convergence of Markov Processes, <http://www.hairer.org/notes/Convergence.pdf>.
- [16] Hairer, M., Ergodicity for stochastic PDEs, 2008. www.hairer.org/notes/Imperial.pdf
- [17] Hairer, M., Mattingly, Jonathan C., Ergodicity of the 2D Navier-Stokes equations with degenerate stochastic forcing, *Ann. of Math.*, **164** (2006), 993–1032.
- [18] Hairer, M., Mattingly, J. C., Scheutzow, M., Asymptotic coupling and a general form of Harris theorem with applications to stochastic delay equations, *Probab. Theory Related Fields*, **149** (2011), 223–259.
- [19] Krylov, N. V., On Kolmogorov’s equations for finite-dimensional diffusions, Stochastic PDE’s and Kolmogorov equations in infinite dimensions (Cetraro, 1998), Lecture Notes in Math., vol. 1715, Springer, Berlin, 1999, pp. 1–63.
- [20] Majda, A. J., Tong, X. T., Geometric ergodicity for piecewise contracting processes with applications for tropical stochastic lattice models, *Comm. Pure Appl. Math.*, **LXIX** (2016), 1110–1153.
- [21] Mao, X., *Stochastic Differential Equations and Applications*, Second Ed., Horwood Publishing Limited, Chichester, 2008.
- [22] Mattingly, J. C., Stuart, A. M., Higham, D. J., Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise, *Stochastic Process. Appl.*, **101** (2002), 185–232.
- [23] Meyn, S. P., Tweedie, R. L., *Markov Chains and Stochastic Stability*, Springer-Verlag, Berlin, 1993.
- [24] Mohammed, S.-E. A., *Stochastic Functional Differential Equations*, Pitman, Boston, 1984.
- [25] Prévôt, C., Röckner, M., A concise course on stochastic partial differential equations. Lecture Notes in Mathematics, 1905. Springer, Berlin, 2007.
- [26] Tong, X. T., Majda, A. J., Moment bounds and geometric ergodicity of diffusions with random switching and unbounded transition rates, *Res. Math. Sci.*, **3** (2016), Paper No. 41, 33 pp.
- [27] Villani, C., Hypocoercivity, *Mem. Amer. Math. Soc.*, **202** (2009), iv+141.
- [28] von Renesse, M.-K., Scheutzow, M., Existence and uniqueness of solutions of stochastic functional differential equations, *Random Oper. Stoch. Equ.*, **18** (2010), 267–284.
- [29] Wang, F.-Y., *Functional Inequalities, Markov Semigroups and Spectral Theory*, Science Press, 2005, Beijing.
- [30] Wang, F.-Y., *Harnack inequalities and Applications for Stochastic Partial Differential Equations*, Springer, 2013, Berlin.
- [31] Wang, F.-Y., Hypercontractivity and applications for stochastic Hamiltonian systems, *J. Funct. Anal.*, **272** (2017), 5360–5383.

- [32] Wang, F.-Y., Yuan, C., Harnack inequalities for functional SDEs with multiplicative noise and applications, *Stochastic Process. Appl.*, **121** (2011), 2692–2710.
- [33] Wu, F., Yin, G., Mei, H., Stochastic functional differential equations with infinite delay: existence and uniqueness of solutions, solution maps, Markov properties, and ergodicity, *J. Differential Equations*, **262** (2017), 1226–1252.