

# A PROBLEM FOR HANKEL MEASURES ON HARDY SPACE

GUANLONG BAO AND FANGQIN YE

ABSTRACT. In this note, we investigate a condition related to the characterization of Hankel measures on Hardy space. We address a problem mentioned by J. Xiao in 2000.

## 1. INTRODUCTION

Let  $\mathbb{D}$  be the open unit disk in the complex plan  $\mathbb{C}$  and let  $H(\mathbb{D})$  be the space of analytic functions in  $\mathbb{D}$ . Recall that for  $0 < p < \infty$  the Hardy space  $H^p$  consists of functions  $f \in H(\mathbb{D})$  such that

$$\|f\|_{H^p} = \sup_{0 < r < 1} \left( \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p} < \infty.$$

It is well known that every function  $f \in H^p$  has non-tangential limits  $f(\zeta)$  for almost every  $\zeta$  on the unit circle  $\mathbb{T}$ . See [8] for the theory of Hardy spaces.

Let  $BMOA$  be the space of analytic functions of bounded mean oscillation. It is well known (cf. [4, 9]) that the space  $BMOA$  can be defined as the set of functions  $f \in H(\mathbb{D})$  satisfying

$$\|f\|_{BMOA}^2 = \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^2 (1 - |\sigma_a(z)|^2) dA(z) < \infty,$$

where

$$dA(z) = \frac{1}{\pi} dx dy = \frac{r}{\pi} dr d\theta, \quad z = x + iy = re^{i\theta},$$

and

$$\sigma_a(z) = \frac{a - z}{1 - \bar{a}z}$$

is the Möbius transformation of  $\mathbb{D}$  interchanging  $a$  and 0. The Fefferman-Stein duality theorem tells us that the dual space of  $H^1$  is  $BMOA$ . Also,  $BMOA$  is a proper subset of the Bloch space  $\mathcal{B}$  consisting of functions  $f \in H(\mathbb{D})$  for which

$$\|f\|_{\mathcal{B}} = \sup_{z \in \mathbb{D}} (1 - |z|^2) |f'(z)| < \infty.$$

Given an arc  $I$  of  $\mathbb{T}$  with arclength  $|I|$ , the Carleson box  $S(I)$  is

$$S(I) = \left\{ r\zeta \in \mathbb{D} : 1 - \frac{|I|}{2\pi} < r < 1, \zeta \in I \right\}.$$

---

2010 *Mathematics Subject Classification.* 30H10, 47B35.

*Key words and phrases.* Hankel measure; Hardy space; Hankel matrix.

The work was supported by NNSF of China (No. 11720101003 and No. 11571217), Department of Education of Guangdong Province (No. 2017KQNCX078) and STU Scientific Research Foundation for Talents (No. NTF17020 and No. STF17005).

A complex measure  $\mu$  on  $\mathbb{D}$  is called a *Carleson measure* if

$$\sup_{I \subseteq \mathbb{T}} \frac{|\mu|(S(I))}{|I|} < \infty.$$

It is well known (cf. [8]) that  $\mu$  is a Carleson measure if and only if there exists a positive constant  $C$  such that

$$\int_{\mathbb{D}} |f(z)|^2 d|\mu|(z) \leq C \|f\|_{H^2}^2$$

for all  $f \in H^2$ .

Following J. Xiao [15], a complex measure  $\mu$  on  $\mathbb{D}$  is said to be a *Hankel measure* if there exists a positive constant  $C$  such that

$$(1.1) \quad \left| \int_{\mathbb{D}} f^2(z) d\mu(z) \right| \leq C \|f\|_{H^2}^2$$

for all  $f$  in  $H^2$ . It is clear that if  $\mu$  is a Hankel measure, then  $|\mu(\mathbb{D})| < \infty$ . The name of Hankel measure has its root in the study of Hankel matrices (see also [11, 14]). Clearly, a Carleson measure must be a Hankel measure. Answering a problem posed by J. Peetre, J. Xiao [15] gave a series of complete characterizations of Hankel measures involving Carleson measures, the duality between  $H^1$  and  $BMOA$ , and Hankel operators.

We refer to [1, p. 11] and [16] for the study of complex measure inequalities similar to (1.1) in the setting of the classical Dirichlet space and weighted Bergman spaces, respectively.

From [15], a necessary condition for a complex measure  $\mu$  on  $\mathbb{D}$  to be a Hankel measure is

$$(1.2) \quad \sup_{w \in \mathbb{D}} \left| \int_{\mathbb{D}} \frac{1 - |w|^2}{(1 - \bar{w}z)^2} d\mu(z) \right| < \infty.$$

By this observation, J. Xiao [15, p. 139] mentioned the following problem: Is condition (1.2) sufficient for  $\mu$  to be a Hankel measure too? In this note, we show that in general the answer to this problem is negative. We give a complex measure  $\mu$  on  $\mathbb{D}$  satisfying condition (1.2) but  $\mu$  is not a Hankel measure. A positive Borel measure  $\mu$  on  $[0, 1)$  can be seen as a Borel measure on  $\mathbb{D}$  by identifying it with the measure  $\tilde{\mu}$  defined by

$$\tilde{\mu}(E) = \mu(E \cap [0, 1)),$$

for any Borel subset  $E$  of  $\mathbb{D}$ . If  $\mu$  is a positive Borel measure on  $[0, 1)$ , we obtain that condition (1.2) holds if and only if  $\mu$  is a Hankel measure, which also implies some new information of a Hankel matrix acting on Hardy spaces and Dirichlet type spaces.

## 2. CONDITION (1.2) AND HANKEL MEASURES

The section is devoted to consider whether condition (1.2) is an equivalent description of Hankel measure. For a complex measure  $\mu$  on  $\mathbb{D}$ , the function  $P_\mu$  is given by

$$P_\mu(w) = \int_{\mathbb{D}} \frac{1}{1 - w\bar{z}} d\mu(z), \quad w \in \mathbb{D}.$$

We will show that the problem considered in this note is equivalent to investigate those functions  $P_{\tilde{\mu}} \in \mathcal{B}$  must actually satisfying  $P_{\tilde{\mu}} \in BMOA$ .

There exist several complete descriptions of Hankel measures in [15]. Now we cite only some of them as follows.

**Theorem A.** *Let  $\mu$  be a complex measure on  $\mathbb{D}$ . Then the following conditions are equivalent.*

- (1)  $\mu$  is a Hankel measure.
- (2) There exists a positive constant  $C$  such that

$$\left| \int_{\mathbb{D}} f(z) d\mu(z) \right| \leq C \|f\|_{H^1},$$

for all  $f \in H^1$ .

- (3)  $P_{\bar{\mu}}$  is in  $BMOA$ .
- (4)

$$\sup_{I \subseteq \mathbb{T}} \frac{1}{|I|} \int_{S(I)} \left| \int_{\mathbb{D}} \frac{\bar{z} d\bar{\mu}(z)}{(1-w\bar{z})^2} \right|^2 (1-|w|^2) dA(w) < \infty.$$

The following theorem characterizes lacunary series in the Bloch space  $\mathcal{B}$  and  $BMOA$  (see [9] and [12]).

**Theorem B.** *Let  $f \in H(\mathbb{D})$  with the power series expansion  $f(z) = \sum_{k=1}^{\infty} a_k z^{n_k}$  and suppose there exists  $\lambda > 1$  such that  $n_{k+1} \geq \lambda n_k$  for all  $k$ . Then the following assertions hold.*

- (1)  $f \in \mathcal{B}$  if and only if the sequence  $\{a_k\}$  is bounded.
- (2)  $f \in BMOA$  if and only if

$$\sum_{k=1}^{\infty} |a_k|^2 < \infty.$$

In 1995, R. Aulaskari, J. Xiao and R. Zhao [3] introduced  $\mathcal{Q}_p$  spaces which attracted a lot of attention in recent years. For  $0 < p < \infty$ , the space  $\mathcal{Q}_p$  consists of functions  $f \in H(\mathbb{D})$  with

$$\|f\|_{\mathcal{Q}_p}^2 = \sup_{a \in \mathbb{D}} \int_{\mathbb{D}} |f'(z)|^2 (1-|\sigma_a(z)|^2)^p dA(z) < \infty.$$

Clearly,  $\mathcal{Q}_1 = BMOA$ . By [2], we know that for  $1 < p < \infty$ , all  $\mathcal{Q}_p$  spaces are the same and equal to the Bloch space  $\mathcal{B}$ . See J. Xiao's monographs [17, 18] for more results of  $\mathcal{Q}_p$  spaces.

The following result can be founded in [17, p. 29].

**Theorem C.** *Suppose  $0 < p < \infty$ . Let  $f(z) = \sum_{n=0}^{\infty} a_n z^n$  with  $\{a_n\}$  being a decreasing sequence of nonnegative numbers. Then  $f \in \mathcal{Q}_p$  if and only if  $a_n = O(\frac{1}{n})$ .*

We first show that the problem considered in this note is associated with certain self-improving property of functions  $P_{\bar{\mu}}$ .

**Theorem 2.1.** *Let  $\mu$  be a complex measure on  $\mathbb{D}$ . Then the following two statements are equivalent.*

- (1) If  $\mu$  satisfies condition (1.2), then  $\mu$  is a Hankel measure.
- (2) If  $P_{\bar{\mu}} \in \mathcal{B}$ , then  $P_{\bar{\mu}} \in BMOA$ .

*Proof.* For  $w \in \mathbb{D}$ , one gets

$$\begin{aligned} wP_{\bar{\mu}}(w) &= \int_{\mathbb{D}} \frac{w}{1-w\bar{z}} d\bar{\mu}(z) \\ &= \sum_{n=0}^{\infty} \left( \int_{\mathbb{D}} \bar{z}^n d\bar{\mu}(z) \right) w^{n+1}. \end{aligned}$$

Then

$$\begin{aligned}
& \sup_{w \in \mathbb{D}} (1 - |w|^2) |(wP_{\bar{\mu}}(w))'| \\
&= \sup_{w \in \mathbb{D}} (1 - |w|^2) \left| \sum_{n=0}^{\infty} (n+1) \left( \int_{\mathbb{D}} \bar{z}^n d\bar{\mu}(z) \right) w^n \right| \\
&= \sup_{w \in \mathbb{D}} \left| \int_{\mathbb{D}} \frac{1 - |w|^2}{(1 - w\bar{z})^2} d\bar{\mu}(z) \right| \\
&= \sup_{w \in \mathbb{D}} \left| \int_{\mathbb{D}} \frac{1 - |w|^2}{(1 - \bar{w}z)^2} d\mu(z) \right|.
\end{aligned}$$

Consequently, the function  $w \rightarrow wP_{\bar{\mu}}(w)$  belongs to  $\mathcal{B}$  if and only if condition (1.2) holds for  $\mu$ .

By the growth estimates of functions in  $\mathcal{B}$  (see [17, p. 6] for example), one gets that there exists a positive  $C$  such that

$$|f(z) - f(0)| \leq C \|f\|_{\mathcal{B}} \log \frac{1}{1 - |z|}$$

for all  $f \in \mathcal{B}$ . Bearing in mind this estimate, we see that  $P_{\bar{\mu}} \in \mathcal{B}$  if and only if the function  $w \rightarrow wP_{\bar{\mu}}(w)$  belongs to  $\mathcal{B}$ .

Thus we proved that  $P_{\bar{\mu}} \in \mathcal{B}$  if and only if  $\mu$  satisfies condition (1.2). By Theorem A,  $\mu$  is a Hankel measure if and only if  $P_{\bar{\mu}} \in BMOA$ . These imply the desired result.  $\square$

The following result shows that in general the answer to J. Xiao's problem is negative.

**Proposition 2.2.** *There exists a complex measure  $\mu$  on  $\mathbb{D}$  such that condition (1.2) holds but  $\mu$  is not a Hankel measure.*

*Proof.* Set  $d\bar{\mu}(z) = f(z)dA(z)$  for  $z \in \mathbb{D}$ , where  $f(z) = 1 + \sum_{k=1}^{\infty} (1 + 2^k)z^{2^k}$ . Then  $\bar{\mu}(\mathbb{D}) = 1$ . By the Parseval formula, for any positive integer  $n$ ,

$$\begin{aligned}
\int_{\mathbb{D}} \bar{z}^n d\bar{\mu}(z) &= \frac{1}{\pi} \int_0^1 r^{n+1} dr \int_0^{2\pi} e^{-in\theta} \left[ 1 + \sum_{k=1}^{\infty} (1 + 2^k) r^{2^k} e^{i2^k\theta} \right] d\theta \\
&= \begin{cases} 0, & \text{if } n \notin \{2^k : k = 1, 2, 3, \dots\}, \\ 1, & \text{if } n \in \{2^k : k = 1, 2, 3, \dots\}. \end{cases}
\end{aligned}$$

Hence, for any  $w \in \mathbb{D}$ ,

$$\begin{aligned}
P_{\bar{\mu}}(w) &= \sum_{n=0}^{\infty} \left( \int_{\mathbb{D}} \bar{z}^n d\bar{\mu}(z) \right) w^n \\
&= 1 + \sum_{k=1}^{\infty} w^{2^k}.
\end{aligned}$$

By Theorem B,  $P_{\bar{\mu}} \in \mathcal{B}$  but  $P_{\bar{\mu}} \notin BMOA$ . Via Theorem A and the proof of Theorem 2.1, we see that the measure  $\mu$  satisfies condition (1.2), but  $\mu$  is not a Hankel measure.  $\square$

For a positive Borel measure  $\mu$  on  $[0, 1)$  and a nonnegative integer  $n$ , denote by  $\mu[n]$  the moment of order  $n$  of  $\mu$ , that is,

$$\mu[n] = \int_0^1 t^n d\mu(t).$$

The following result shows that there exist measures  $\mu$  such that if  $P_{\bar{\mu}} \in \mathcal{B}$ , then  $P_{\bar{\mu}} \in BMOA$ . It follows from Theorem 2.1 that in this case  $\mu$  is a Hankel measure if and only if  $\mu$  satisfies condition (1.2).

**Theorem 2.3.** *Let  $\mu$  be a positive Borel measure on  $[0, 1)$ . Then the following conditions are equivalent.*

- (1)  $\mu$  is a Hankel measure.
- (2)  $\mu$  satisfies condition (1.2), that is,

$$\sup_{w \in \mathbb{D}} \left| \int_0^1 \frac{1 - |w|^2}{(1 - \bar{w}t)^2} d\mu(t) \right| < \infty.$$

- (3)  $P_{\mu} \in \mathcal{Q}_p$  for some  $0 < p < \infty$ .
- (4)  $P_{\mu} \in \mathcal{Q}_p$  for all  $0 < p < \infty$ .
- (5)  $\mu[n] = O(\frac{1}{n})$ .

*Proof.* If  $\mu$  is a positive Borel measure on  $[0, 1)$ , then for  $w \in \mathbb{D}$ ,

$$P_{\bar{\mu}}(w) = P_{\mu}(w) = \sum_{n=0}^{\infty} \left( \int_0^1 t^n d\mu(t) \right) w^n = \sum_{n=0}^{\infty} \mu[n] w^n.$$

Clearly,  $\{\mu[n]\}$  is a decreasing sequence with nonnegative numbers. By Theorem C, we see that (3) $\Leftrightarrow$ (4) $\Leftrightarrow$ (5). By the proof of Theorem 2.1,  $P_{\bar{\mu}} \in \mathcal{B}$  if and only if  $\mu$  satisfies condition (1.2). From Theorem A,  $P_{\bar{\mu}} \in BMOA$  if and only if  $\mu$  is Hankel measure. Bear in mind that  $\mathcal{B} = \mathcal{Q}_2$  and  $BMOA = \mathcal{Q}_1$ . Thus both (1) and (2) are equivalent to (5). The proof is complete.  $\square$

Let  $\mu$  be a positive Borel measure on  $[0, 1)$ . Denote by  $\mathcal{H}_{\mu}$  the Hankel matrix  $(\mu[n+k])_{n,k \geq 0}$ . This matrix induces formally an operator, denoted also by  $\mathcal{H}_{\mu}$ , on  $H(\mathbb{D})$  in the following sense. For  $f(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{D})$ , by multiplication of the matrix with the sequence of Taylor coefficients of the function  $f$ ,

$$\{a_n\}_{n \geq 0} \mapsto \left\{ \sum_{k=0}^{\infty} \mu[n+k] a_k \right\}_{n \geq 0},$$

define

$$\mathcal{H}_{\mu}(f)(z) = \sum_{n=0}^{\infty} \left( \sum_{k=0}^{\infty} \mu[n+k] a_k \right) z^n.$$

If  $\mu$  is the Lebesgue measure on  $[0, 1)$ , the matrix  $\mathcal{H}_{\mu}$  reduces to the classical Hilbert matrix  $\mathcal{H} = ((n+k+1)^{-1})_{n,k \geq 0}$ , which induces the classical Hilbert operator  $\mathcal{H}$ . The Hankel matrix  $\mathcal{H}_{\mu}$  acting on  $H^2$  was studied in [13, 14]. For all Hardy spaces  $H^p$ , the operator  $\mathcal{H}_{\mu}$  was investigated in [6, 10]. The Dirichlet type space  $\mathcal{D}_{\alpha}$ ,  $\alpha \in \mathbb{R}$ , consists of functions  $f(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{D})$  for which

$$\|f\|_{\mathcal{D}_{\alpha}}^2 = \sum_{n=0}^{\infty} (n+1)^{1-\alpha} |a_n|^2 < \infty.$$

The Hankel matrix  $\mathcal{H}_\mu$  acting on  $\mathcal{D}_\alpha$  spaces was considered in [5, 7]. It is worth referring to [10] for the recent development of  $\mathcal{H}_\mu$  acting on spaces of analytic functions. The following theorem is from the related literatures mentioned before.

**Theorem D.** *Let  $\mu$  be a positive Borel measure on  $[0, 1)$ . Suppose  $1 < p < \infty$  and  $0 < \alpha < 2$ . Then the following conditions are equivalent.*

- (1)  $\mu$  is a Carleson measure.
- (2)

$$\sup_{w \in \mathbb{D}} \int_0^1 \frac{1 - |w|^2}{|1 - \bar{w}t|^2} d\mu(t) < \infty.$$

- (3)  $\mu[n] = O(\frac{1}{n})$ .
- (4)  $\mathcal{H}_\mu$  is bounded on  $H^p$ .
- (5)  $\mathcal{H}_\mu$  is bounded on  $\mathcal{D}_\alpha$ .

Combining Theorem 2.3 with Theorem D, we obtain immediately new characterizations of a Hankel matrix  $\mathcal{H}_\mu$  acting on Hardy spaces and Dirichlet type spaces in terms of Hankel measures and  $\mathcal{Q}_p$  functions.

#### REFERENCES

- [1] N. Arcozzi, R. Rochberg, E. Sawyer and B. Wick, *Function spaces related to the Dirichlet space*, J. Lond. Math. Soc., 83 (2011), 1-18.
- [2] R. Aulaskari and P. Lappan, *Criteria for an analytic function to be Bloch and a harmonic or meromorphic function to be normal*, Complex analysis and its applications, Pitman Res. Notes in Math., 305, Longman Sci. Tec., Harlow, 1994, 136-146.
- [3] R. Aulaskari, J. Xiao and R. Zhao, *On subspaces and subsets of BMOA and UBC*, Analysis, 15 (1995), 101-121.
- [4] A. Baernstein, *Analytic functions of bounded mean oscillation*, Aspects of Contemporary Complex Analysis, Academic Press, 1980, 3-36.
- [5] G. Bao and H. Wulan, *Hankel matrices acting on Dirichlet spaces*, J. Math. Anal. Appl., 409 (2014), 228-235.
- [6] C. Chatzifountas, D. Girela and J. Peláez, *A generalized Hilbert matrix acting on Hardy spaces*, J. Math. Anal. Appl., 413 (2014), 154-168.
- [7] E. Diamantopoulos, *Operators induced by Hankel matrices on Dirichlet spaces*, Analysis (Munich), 24 (2004), 345-360.
- [8] P. Duren, *Theory of  $H^p$  Spaces*, Academic Press, New York, 1970.
- [9] D. Girela, *Analytic functions of bounded mean oscillation*. In: Complex Function Spaces, Mekrijärvi 1999 Editor: R. Aulaskari. Univ. Joensuu Dept. Math. Rep. Ser. 4, Univ. Joensuu, Joensuu (2001) pp. 61-170.
- [10] D. Girela and N. Merchán, *A Hankel matrix acting on spaces of analytic functions*, Integral Equations Operator Theory, 89 (2017), 581-594.
- [11] S. Janson and J. Peetre, *A new generalization of Hankel operators (the case of higher weights)*, Math. Nachr., 132 (1987), 313-328.
- [12] Ch. Pommerenke, *On Bloch functions*, J. London Math. Soc., 2 (1970), 689-695.
- [13] S. Power, *Vanishing Carleson measures*, Bull. London Math. Soc., 12 (1980), 207-210.
- [14] H. Widom, *Hankel matrices*, Trans. Amer. Math. Soc., 121 (1966), 1-35.
- [15] J. Xiao, *Hankel measures on Hardy space*, Bull. Austral. Math. Soc., 62 (2000), 135-140.
- [16] J. Xiao, *Pseudo-Carleson measures for weighted Bergman spaces*, Michigan Math. J., 47 (2000), 447-452.
- [17] J. Xiao, *Holomorphic  $\mathcal{Q}$  Classes*, Springer, LNM 1767, Berlin, 2001.
- [18] J. Xiao, *Geometric  $\mathcal{Q}_p$  Functions*, Birkhäuser Verlag, Basel-Boston-Berlin, 2006.

GUANLONG BAO, DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA

*E-mail address:* glbao@stu.edu.cn

FANGQIN YE, BUSINESS SCHOOL, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA

*E-mail address:* fqye@stu.edu.cn