Pre-Markov Operators

Hūlya Duru and Serkan Ilter

ABSTRACT. A positive linear operator T between two unital f-algebras, with point separating order duals, A and B is called a Markov operator for which $T(e_1) = e_2$ where e_1, e_2 are the identities of A and B respectively. Let A and B be semiprime f-algebras with point separating order duals such that their second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ are unital *f*-algebras. In this case, we will call a positive linear operator $T: A \to B$ to be a pre-Markov operator, if the second adjoint operator of T is a Markov operator. A positive linear operator T between two semiprime f-algebras, with point separating order duals, A and B is said to be contractive if $Ta \in B \cap [0, I_B]$ whenever $a \in A \cap [0, I_A]$, where I_A and I_B are the identity operators on A and B respectively. In this paper we characterize pre-Markov algebra homomorphisms. In this regard, we show that a pre-Markov operator is an algebra homomorphism if and only if its second adjoint operator is an extreme point in the collection of all Markov operators from $A^{\sim \sim}$ to $B^{\sim \sim}$. Moreover we characterize extreme points of contractive operators from A to B. In addition, we give a condition that makes an order bounded algebra homomorphism is a lattice homomorphism.

1. Introduction

A positive linear operator T between two unital f-algebras, with point separating order duals, A and B is called a Markov operator for which $T(e_1) = e_2$ where e_1 , e_2 are the identities of A and B respectively. Let A and B be semiprime f-algebras with point separating order duals such that their second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ are unital f-algebras. In this case, we will call a positive linear operator $T: A \to B$ to be a pre-Markov operator, if the second adjoint operator of T is a Markov operator. A positive linear operator T between two semiprime f-algebras, with point separating order duals, A and B is said to be contractive if $Ta \in B \cap [0, I_B]$ whenever $a \in A \cap [0, I_A]$, where I_A and I_B are the identity operators on A and B respectively.

The collection of all pre-Markov operators is a convex set. In this paper, first of all, we characterize pre-Markov algebra homomorphisms. In this regard, we show that a pre-Markov operator is an algebra homomorphism if and only if its second adjoint operator is an extreme point in the collection of all Markov operators from $A^{\sim\sim}$ to $B^{\sim\sim}$ (Theorem 1). In addition, we characterize the extreme points of all contractive operators $T: A \to B$ whenever A and B are Archimedean semiprime

²⁰¹⁰ Mathematics Subject Classification. Primary 47B38, 46B42.

Key words and phrases. Markov Operator, f-algebra, algebra homomorphism, lattice homomorphism, contractive operator, Arens Multiplication.

f-algebras provided B is relatively uniformly complete (Proposition 3). For the second aim, let A and B be Archimedean semiprime f-algebras and $T: A \to B$ a linear operator. Huijsman and de Pagter proved in [8] the following:

(1) If T is a positive algebra homomorphism then it s a lattice homomorphism.

(2) In addition, if the domain A is relatively uniformly complete and T is an algebra homomorphism then it s a lattice homomorphism and the assumption that the domain A of T is relatively uniformly complete is not reduntant (Theorem 5.1 and Example 5.2.).

(3) In addition, if the domain A has a unit element and T is an order bounded algebra homomorphism then it s a lattice homomorphism (Theorem 5.3).

We prove that any order bounded algebra homomorphism $T : A \to B$ is a lattice homomorphism, if the region B is relatively uniformly complete (Corollary 3). In this regard, first we give an alternate proof of Lemma 6 in [10] for order bounded operators with the relatively uniformly complete region instead of positive operators with Dedekind complete region (Proposition 4 and 5). In the last part, we give a necessary and sufficient condition for a positive operator to be a lattice homomorphism (Proposition 7).

2. Preliminaries

For unexplained terminology and the basic results on vector lattices and semiprime f-algebras we refer to [1, 11, 13, 15]. The real algebra A is called a Riesz algebra or lattice-ordered algebra if A is a Riesz space such that $ab \in A$ whenever a, b are positive elements in A. The Riesz algebra is called an f-algebra if A satisfies the condition that

$$a \wedge b = 0$$
 implies $ac \wedge b = ca \wedge b = 0$ for all $0 \leq c \in A$.

In an Archimedean *f*-algebra *A*, all nilpotent elements have index 2. Indeed, assume that $a^3 = 0$ for some $0 \le a \in A$. Since the equality $(a^2 - na) \land (a - na^2) = 0$ implies $(a^2 - na) \land a^2 = (a^2 - na) = 0$ we get $a^2 = 0$ as *A* is Archimedean. The same argument is true for all $n \ge 3$. Throughout this paper *A* will show an Archimedean semiprime *f*-algebra with point separating order dual A^{\sim} [15]. By definition, if zero is the unique nilpotent element of *A*, that is, $a^2 = 0$ implies a = 0, *A* is called semiprime *f*-algebra. It is well known that every *f*-algebra with unit element is semiprime.

Let A be a lattice ordered algebra. If A is a lattice ordered space, then the first order dual space A^{\sim} of A is defined as the collection of all order bounded linear functionals on A and A^{\sim} is a Dedekind complete Riesz space. The second order dual space of A is denoted by $A^{\sim\sim}$. Let $a \in A$, $f \in A^{\sim}$ and $F, G \in A^{\sim\sim}$. Define $f \cdot a \in A^{\sim}$, by $(f \cdot a)(b) = f(ab)$

and $F \cdot f \in A^{\sim}$, by

$$(F \cdot f)(a) = F(f \cdot a)$$

and $F \cdot G \in A^{\sim \sim}$, by

$$(F \cdot G)(f) = F(G \cdot f)$$

The last equality is called the Arens multiplication in $A^{\sim \sim}$ [2].

The second order dual space $A^{\sim \sim}$ of a semiprime *f*-algebra *A* is again an *f*-algebra with respect to the Arens multiplication [4]. In the literature, there are

several studies, for example [5, 6, 7, 9], that respond the question "Under what conditions does the *f*-algebra $A^{\sim \sim}$ have a unit element?".

Let A and B be semiprime f-algebras with point separating order duals such that their second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ have unit elements E_1 and E_2 respectively. Let $T: A \to B$ be an order bounded operator. We denote the second adjoint operator of T by T^{**} . Since A and B have point separating order duals, the linear operator $J_1: A \to A^{\sim\sim}$, which assigns to $a \in A$ the linear functional \hat{a} defined on A^{\sim} by $\hat{a}(f) = f(a)$ for all $a \in A$, is an injective algebra homomorphism. Therefore we will identify A with $J_1(A)$, and B with $J_2(B)$ in the similar sense.

DEFINITION 1. Let A and B be semiprime f-algebras with point separating order duals such that their second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ are unital f-algebras. In this case, we call a positive linear operator $T : A \to B$ to be a pre-Markov operator, if the second adjoint operator of T is a Markov operator. That is, the second adjoint operator $T^{**} : A^{\sim\sim} \to B^{\sim\sim}$ of T is a positive linear and $T^{**}(E_1) =$ E_2 , where E_1 and E_2 are the unitals of A and B respectively.

Recall that a positive operator $T : A \to B$ satisfying $0 \le T(a) \le E_2$ whenever $0 \le a \le E_1$ is called a contractive operator.

In this point we remark that , if A and B are semiprime f-algebras with point separating order duals and $T: A \to B$ is a positive linear operator, then T^{**} is positive. Indeed, let $0 \leq F \in A^{\sim\sim}$ and $0 \leq g \in B^{\sim}$. Then $0 \leq g \circ T \in A^{\sim}$ and therefore $F(g \circ T) = T^{**}(F) \geq 0$.

PROPOSITION 1. Let A and B be semiprime f-algebras with point separating order duals such that their second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ have unit elements E_1 and E_2 respectively. $T: A \to B$ is contractive if and only if T^{**} is contractive.

PROOF. Suppose that T is contractive. Then T^{**} is positive. Let $F \in [0, E_1] \cap A^{\sim \sim}$. In order to prove that T^{**} is contractive we shall show that $T^{**}(E_1) \leq E_2$. Due to [9],

$$E_1(f) = \sup f(A \cap [0, E_1]) E_2(g) = \sup g(B \cap [0, E_2])$$

for all $f \in A^{\sim}$ and $g \in B^{\sim}$. Let $a \in A \cap [0, E_1]$ and $0 \leq g \in B^{\sim}$. Since T is contractive, $T(a) \in B \cap [0, E_2]$ so $g(T(a)) \leq E_2(g)$ which implies that $T^{**}E_1(g) = E_1(g \circ T) \leq E_2(g)$. Thus $T^{**}(E_1) \leq E_2$. Conversely, assume that T^{**} is contractive. Let $a \in A \cap [0, E_1]$ and $0 \leq g \in B^{\sim}$. Then $\widehat{Ta}(g) = g(Ta) \leq T^{**}E_1(g) \leq E_2(g)$ Thus $0 \leq Ta = \widehat{Ta} \leq E_2$.

COROLLARY 1. Let A and B be semiprime f-algebras with point separating order duals such that their second order duals A^{\sim} and B^{\sim} have unit elements E_1 and E_2 respectively. If $T: A \to B$ is a pre-Markov operator then T is contractive.

PROOF. Since $T^{**}(E_1) = E_2$ and T^{**} is positive, T^{**} is contractive. By Proposition 1 we have the conclusion.

For $a \in A$, the mapping $\pi_a : A \to OrthA$, defined by $\pi_a(b) = a.b$ is an orthomorphism on A. Since A is a Archimedean semiprime f-algebra, the mapping $\pi : A \to OrthA$, defined by $\pi(a) = \pi_a$ is an injective f-algebra homomorphism. Hence we shall identify A with $\pi(A)$.

3. Main Results

THEOREM 1. Let A and B be semiprime f-algebras with point separating order duals such that their second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ have unit elements E_1 and E_2 respectively. A pre-Markov operator $T: A \to B$ is an algebra homomorphism if and only if its second adjoint operator T^{**} is an algebra homomorphism.

PROOF. Suppose that the pre-Markov operator T is an algebra homomorphism. Since T^{**} is a Markov operator, due to [8], it is enough to show that it is a lattice homomorphism. Let $F, G \in A^{\sim \sim}$ such that $F \wedge G = 0$. Since $A^{\sim \sim}$ and $B^{\sim \sim}$ are semiprime f- algebras, $F \cdot G = 0$. We shall show that $T^{**}(F) \cdot T^{**}(G) = 0$. Let $a, b \in A$ and $f \in B^{\sim}$. Then it follows from the following equations

$$((f \cdot Ta) \circ T)(b) = (f \cdot Ta)(Tb) = f(TaTb) = f(T(ab))$$
$$= (f \circ T)(ab) = ((f \circ T) \cdot a)(b)$$

that

$$(3.1) (f \cdot Ta) \circ T = (f \circ T) \cdot a$$

On the other hand, the following equations

$$((G \circ T^*) \cdot f) \circ T)(a) = ((G \circ T^*) \cdot f)(Ta) = (G \circ T^*)(f \cdot Ta) = G((f \cdot Ta) \circ T)$$

hold. Thus $((G \circ T^*) \cdot f) \circ T)(a) = G((f \cdot Ta) \circ T)$. From here, by setting the equation (3.1), we conclude that

$$\left(\left(G\circ T^*\right)\cdot f\right)\circ T\right)(a)=G\left(\left(f\circ T\right)\cdot a\right)=\left(G\cdot \left(f\circ T\right)\right)(a)$$

which implies

(3.2)
$$((G \circ T^*) \cdot f) \circ T) = (G \cdot (f \circ T))$$

Taking into account the equation (3.2), we get

$$\begin{array}{ll} \left(T^{**}\left(F\right) \cdot T^{**}\left(G\right)\right)\left(f\right) &=& T^{**}\left(F\right)\left(\left(T^{**}\left(G\right) \cdot f\right)\right) = \left(F \circ T^{*}\right)\left(\left(G \circ T^{*}\right) \cdot f\right) \\ &=& F\left(\left(G \circ T^{*}\right) \cdot f\right) \circ T\right) = F\left(G \cdot \left(f \circ T\right)\right) \end{array}$$

thus we have

$$(T^{**}(F) \cdot T^{**}(G))(f) = (F \cdot G)(f \circ T) = 0$$

as desired. Conversely suppose that T^{**} is an algebra homomorphism. Let $a, b \in A$. It follows from

$$T(ab) = \widehat{T(ab)} = T^{**}\left(\widehat{ab}\right) = T^{**}\left(\widehat{a}\cdot\widehat{b}\right) = T^{**}\left(\widehat{a}\right) \cdot T^{**}\left(\widehat{b}\right) = \widehat{Ta}\cdot\widehat{Tb} = Ta.Tb$$

that T is an algebra homomorphism.

COROLLARY 2. Let A, B and their second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ be semiprime f-algebras and $T: A \to B$ a positive algebra homomorphism. Then T^{**} is a lattice homomorphism.

THEOREM 2. Let A and B be semiprime f-algebras with point separating order duals and $T: A \to B$ a positive linear operator. If the second order duals $A^{\sim\sim}$ and $B^{\sim\sim}$ have unit elements and T is an algebra homomorphism, then T is an extreme point of the contractive operators from A to B.

$$\square$$

PRE-MARKOV

PROOF. Suppose that T is a positive algebra homomorphism. Then due to [14, Theorem 4.3], T is a contractive operator. Let $2T = T_1 + T_2$ for some contractive operators T_1, T_2 from A to B. In this case, $2T^{**} = T_1^{**} + T_2^{**}$. By Proposition 1, T^{**} , T_1^{**} and T_2^{**} are contractive and by Corollary 2, T^{**} is a lattice homomorphism. Taking into account [3, Theorem 3.3], we derive that T^{**} is an extreme point in the collection of all contractive operators from $A^{\sim\sim}$ to $B^{\sim\sim}$. Thus $T^{**} = T_1^{**} = T_2^{**}$ and therefore $T = T_1 = T_2$.

At this point, we recall the definition of uniform completion of an Archimedean Riesz space. If A is an Archimedean Riesz space and \widehat{A} is the Dedekind completion of A, then \overline{A} , the closure of A in \widehat{A} with respect to the relatively uniform topology [11], is so called that relatively uniformly completion of A [12]. If A is an semiprime f-algebra then the multiplication in A can be extended in a unique way into a lattice ordered algebra multiplication on \overline{A} such that A becomes a sub-algebra of \overline{A} and \overline{A} is an relatively uniformly complete semiprime f-algebra. In [14, Theorem 3.4] it is shown that a positive operator T from a Riesz space A to a uniformly complete space B, has a unique positive linear extension $\overline{T} : \overline{A} \to B$ to the relatively uniformly completion \overline{A} of A, defined by,

$$\overline{T}(x) = \sup \left\{ T(a) : 0 \le a \le x \right\}$$

for $0 \le x \in \overline{A}$. We also recall that \overline{A} satisfies the Stone condition (that is, $x \wedge nI^* \in \overline{A}$, for all $x \in \overline{A}$, where I denotes the identity on A of OrthA) due to Theorem 2.5 in [7]. For the completeness we give the easy proof of the following proposition.

PROPOSITION 2. Let A and B be Archimedean semiprime f-algebras such that B is relatively uniformly complete. In this case, $T : A \to B$ is contractive if and only if \overline{T} is contractive.

PROOF. Suppose that T is contractive. Let $x \in \overline{A} \cap [0,\overline{I}]$, here \overline{I} is the unique extension to \overline{A} of the identity mapping $I : A \to A$. Since T is contractive, $a \in A \cap [0, x]$ implies that I is an upper bound for the set $\{T(a) : a \leq x, a \in A\}$, so $\overline{T}(x) \leq I$. Therefore \overline{T} is contractive. The converse implication is trivial, since \overline{T} is the extension of T, we get $0 \leq \overline{T}(a) = T(a) \leq I$ whenever $a \in A \cap [0, I]$.

PROPOSITION 3. Let A and B be Archimedean semiprime f-algebras such that B is relatively uniformly complete and let $T : A \to B$ be a contractive operator. Then T is an extreme point in the collection of all contractive operators from A to B if and only if \overline{T} is an extreme point of all contractive operators from \overline{A} to B.

PROOF. Suppose that \overline{T} is an extreme point in the set of all contractive operators from \overline{A} to B. We shall show that for arbitrary $\varepsilon > 0$ and contractive operator S from A to B satisfying $\varepsilon T - S \ge 0$ implies that T = S. Let $0 \le x \in \overline{A}$. Then there exists a positive sequence $(a_n)_n$ in A converging relatively uniformly to x. Since \overline{T} and \overline{S} are relatively uniformly continuos, the sequence $\varepsilon \overline{T}(a_n) - \overline{S}(a_n) = \varepsilon T(a_n) - S(a_n)$ converges to $\varepsilon \overline{T}(x) - \overline{S}(x)$. Therefore, since $(a_n)_n$ is positive sequence and $\varepsilon T - S \ge 0$, we get $\varepsilon \overline{T} - \overline{S} \ge 0$. Since \overline{T} is an extreme point, we have $\overline{T} = \overline{S}$, so that T = S. Conversely assume that T is an extreme point in the set of all contractive operator from A to B. Let $\varepsilon > 0$ be any number and let S be any contractive operator from \overline{A} to B satisfying $\varepsilon \overline{T} - S \ge 0$. Let U be the restriction of S to A. Since S is contractive, by Proposition 2, $S \mid_A = U$

is contractive and by the uniqueess of the extension, we infer that $S = \overline{U}$. Hence $(\varepsilon \overline{T} - S) \mid_A = \varepsilon T - U \ge 0$. Thus $\overline{T} = S$, which shows that \overline{T} is an extreme point.

After proving the following Propositions 4 and 5 for order bounded operators with the relatively uniformly complete region, we remarked that both were proved in [10] for the positive operators with Dedekind complete region. They might be regarded as the alternate proofs.

PROPOSITION 4. Let $T : A \to B$ be an order bounded operator where A and B are Archimedean f-algebras and B is, in addition, relatively uniformly complete. Then T is an algebra homomorphism iff \overline{T} is an algebra homomorphism.

PROOF. Suppose that $T: A \to B$ is an algebra homomorphism and x, y be positive elements in \overline{A} . By [14], since

$$xy = \sup \left\{ R_y(a) : 0 \le a \le x, a \in A \right\}$$

and

$$R_{y}(a) = \sup \left\{ ab : 0 \le b \le y, b \in A \right\}.$$

Now as \overline{T} is relatively uniformly continuous, we get,

$$\overline{T} (R_y (a)) = \sup \left\{ \overline{T} (ab) = T (ab) = T (a) T (b) : 0 \le b \le y, b \in A \right\}$$
$$= T (a) \sup \left\{ T (b) : 0 \le b \le y, b \in A \right\}$$
$$= T (a) \overline{T} (y)$$

and then

$$\overline{T}(xy) = \sup \left\{ \overline{T}(R_y(a)) : 0 \le a \le x, a \in A \right\}$$
$$= \sup \left\{ T(a) \overline{T}(y) : 0 \le a \le x, a \in A \right\}$$
$$= \overline{T}(y) \sup \left\{ T(a) : 0 \le a \le x, a \in A \right\}$$
$$= \overline{T}(x) \overline{T}(y)$$

Hence \overline{T} is an algebra homomorphism. The converse is trivial.

In [8], both were proved that an algebra homomorphism $T: A \to B$ need not be a lattice homomorphism if the domain A is not relatively uniformly complete (Example 5.2) and an order bounded algebra homomorphism $T: A \to B$ is a lattice homomorphism whenever the domain A has a unit element. We remarked that Proposition 4 yields that the second result also holds for an order bounded algebra homomorphism without unitary domain but the region is relatively uniformly complete.

COROLLARY 3. Let A be an Archimedean semiprime f-algebra and B a relatively uniformly complete Archimedean f-algebra. Then any order bounded algebra homomorphism $T: A \to B$ is a lattice homomorphism.

PROOF. By Proposition 4, \overline{T} is an algebra homomorphism and since \overline{A} is relatively uniformly complete, \overline{T} is a lattice homomorphism [8]. Thus T is a lattice homomorphism.

PROPOSITION 5. Let A be an Archimedean f-algebra and let B be a relatively uniformly complete semiprime f-algebra. Then the operator $T: A \to B$ is a lattice homomorphism iff \overline{T} is a lattice homomorphism.

PRE-MARKOV

PROOF. Suppose that T is a lattice homomorphism. Let $x \in \overline{A}$. Let $a \in [0, x^+] \cap A$ and $b \in [0, x^-] \cap A$. Since T is a lattice homomorphism, we have

$$T(a \wedge b) = T(a) \wedge T(b) = 0.$$

On the other hand, it follows from the equality

$$T(a) \wedge \overline{T}(x^{-}) = \sup \left\{ T(a) \wedge T(b) : 0 \le b \le x^{-}, b \in A \right\}$$

that

$$\overline{T}(x^{+})\overline{T}(x^{-}) = \sup\left\{T(a) \land \overline{T}(x^{-}) : 0 \le a \le x^{+}, a \in A\right\} = 0$$

which its turn is equivalent to \overline{T} is a lattice homomorphism, as B is semiprime. Converse is trivial.

In this point, we remark that Lemma 3.1 and Theorem 3.3 in [3] are also true for Archimedean semiprime f-algebras without the Stone condition on the domain A whenever B is relatively uniformly complete.

PROPOSITION 6. Let A and B be Archimedean semiprime f-algebras, B relatively uniformly complete and $T: A \to B$ a contractive operator. Assume that \overline{A} has unit element. For $y \in \overline{A}$, define $H_x(y) = \overline{T}(xy) - \overline{T}(x)\overline{T}(y)$. Then $\overline{T} + H_x$ are contractive mappings for all $x \in \overline{A} \cap [0, I]$.

PROOF. By Proposition 2, \overline{T} is contractive. Since \overline{A} satisfies the Stone condition, due to [3, Lemma 3.1], we have the conclusion.

COROLLARY 4. Let A and B be Archimedean semiprime f-algebras such that B is relatively uniformly complete and let $T : A \to B$ be a contractive operator. If \overline{A} has unit element, then $T + T_a$ are contractive for all $a \in A \cap [0, I]$, here $T_a(b) = T(ab) - T(a)T(b)$.

PROOF. By Proposition 6, $\overline{T} + H_x$ are contractive mappings for all $x \in \overline{A} \cap [0, I]$. Let $a \in A \cap [0, I]$ and $0 \le b \in A$. Then $0 \le (\overline{T} + H_a)$ $(b) = T(b) + T_a(b)$ holds. Thus $T + T_a$ is positive. Let $b \in A \cap [0, I]$. It follows from

$$0 \le \left(\overline{T} + H_a\right)(b) = T(b) + T_a(b) \le I$$

that $T + T_a$ are contractive.

PROPOSITION 7. Let A and B be Archimedean semiprime f-algebras such that B is relatively uniformly complete and let $T : A \to B$ be a positive linear operator. T is contractive and it is an extreme point in the collection of all contractive operators from A to B if and only if T is an algebra homomorphism.

PROOF. Let T be an extreme point in the collection of all contractive operators from A to B. Then by Proposition 3, \overline{T} is an extreme point of all contractive operators from \overline{A} to B. It follows from [3, Theorem 3.3] that \overline{T} is an algebra homomorphism. By Proposition 4, T is an algebra homomorphism. Conversely, if T is an algebra homomorphism, then due to [14, Theorem 4.3], T is a contractive operator. By Proposition 2, \overline{T} is contractive and by Proposition 4, \overline{T} is an algebra homomorphism. Thus \overline{T} is an extreme point in the set of all contractions from \overline{A} to B due to [3, Theorem 3.3]. By using Proposition 3, we have the conclusion. \Box

References

- [1] C. D. Aliprantis, O. Burkinshaw, Positive Operators, Academic Press, Orlando, 1985
- [2] R. Arens, The adjoint of bilinear operation, Proc. Am. Math. Soc. 2 (1951) 839-848
- [3] M. A. Ben Amor, K. Boulabiar, C. El Adeb, Extreme contractive operators on Stone falgebras, Indagationes Mathematicae 25 (2014) 93-103
- [4] S. J. Bernau, C.B. Huijsmans, The order bidual of almost f-algebras and d-algebras, Trans.Amer.Math.Soc. 347 (1995) 4259-4275
- [5] K. Boulabiar, J. Jaber, The order bidual of f-algebras revisited, Positivity 15 (2011) 271-279
- [6] C.B. Huijsmans, The order bidual of lattice-ordered algebras II, J.Operator Theory 22 (1989) 277-290
- [7] C. B. Huijsmans, B. de Pagter, The order bidual of lattice-ordered algebras, J.Funct.Analysis 59 (1984) 41-64
- [8] C. B. Huijsmans, B. de Pagter, Subalgebras and Riesz subspaces of an f-algebra, Proc.Lond.Math.Soc. 48 (1984) 161-174
- [9] J. Jaber, f-algebras with σ -bounded approximate unit, Positivity, 18 (2014) 161-170
- [10] J. Jaber, Contractive operators on semiprime f -algebras, Indagationes Mathematicae, 28 (2017) 1067-1075
- [11] W. A. J. Luxemburg, A. C. Zaanen, Riesz Spaces I, North Holland, Amsterdam, 1971
- [12] J. Quinn, Intermediate Riesz spaces, Pac.J.Math. 56 (1975) 255-263
- [13] H. H. Schaefer, Banach Lattices and Positive Operators, Springer, Berlin, 1974
- [14] A. Triki, Algebra homomorphisms in complex almost f-algebras, Comment Math.Univ.Carol. 43 (2002) 23-31
- [15] A. C. Zaanen, Riesz Spaces II, North- Holland, Amsterdam, 1983

ISTANBUL UNIVERSITY, FACULTY OF SCIENCE, MATHEMATICS DEPARTMENT, VEZNECILER-ISTANBUL, 34134, TURKEY

E-mail address: hduru@istanbul.edu.tr

Current address: Istanbul University, Faculty of Science, Mathematics Department, Vezneciler-Istanbul, 34134, Turkey

E-mail address: ilters@istanbul.edu.tr