

# $M_2$ -Ranks of overpartitions modulo 6 and 10

Helen W.J. Zhang

Center for Applied Mathematics

Tianjin University, Tianjin 300072, P.R. China

wenjingzhang@tju.edu.cn

**Abstract.** In this paper, we obtain inequalities on  $M_2$ -ranks of overpartitions modulo 6. Let  $\overline{N}_2(s, m, n)$  to be the number of overpartitions of  $n$  whose  $M_2$ -rank is congruent to  $s$  modulo  $m$ . For  $M_2$ -ranks modulo 3, Lovejoy and Osburn derived the generating function of  $\overline{N}_2(s, 3, n) - \overline{N}_2(t, 3, n)$ , which implies the inequalities  $\overline{N}_2(0, 3, n) \geq \overline{N}_2(1, 3, n)$ . For  $\ell = 6, 10$ , we consider the generating function  $\overline{R}_{s,t}(d, \ell)$  of the  $M_2$ -rank differences  $\overline{N}_2(s, \ell, \ell n/2 + d) + \overline{N}_2(s+1, \ell, \ell n/2 + d) - \overline{N}_2(t, \ell, \ell n/2 + d) - \overline{N}_2(t+1, \ell, \ell n/2 + d)$ . By the method of Lovejoy and Osburn, we derive a formula for  $\overline{R}_{0,2}(d, 6)$ . This leads to the inequalities for  $n \geq 0$ ,  $\overline{N}_2(0, 6, 3n) \geq \overline{N}_2(2, 6, 3n)$  and  $\overline{N}_2(0, 6, 3n+1) \geq \overline{N}_2(2, 6, 3n+1)$ . Based on the valence formula for modular functions, we compute  $\overline{R}_{0,4}(d, 10)$  and  $\overline{R}_{1,3}(d, 10)$ . In particular, we notice that the generating function  $\overline{R}_{0,2}(2, 6)$  can be expressed in terms of the third order mock theta function  $\rho(q)$ , and the generating functions  $\overline{R}_{0,4}(4, 10)$ ,  $\overline{R}_{1,3}(1, 10)$  and  $\overline{R}_{1,3}(4, 10)$  can also be expressed in terms of the tenth order mock theta functions  $\phi(q)$  and  $\psi(q)$ .

**Keywords:** Overpartition,  $M_2$ -rank, rank difference, generalized  $\eta$ -function, modular function, mock theta function.

**AMS Classifications:** 11P81, 05A17, 33D15.

## 1 Introduction

The rank of a partition was introduced by Dyson [8], which equals to the largest part minus the number of parts. Let  $N(s, \ell, n)$  denote the number of partitions of  $n$  with rank congruent to  $s$  modulo  $\ell$ . Atkin and Swinnerton-Dyer [4] obtained generating functions for rank differences  $N(s, \ell, \ell n + d) - N(t, \ell, \ell n + d)$  with  $\ell = 5$  or  $7$  and  $0 \leq d, s, t < \ell$ , which lead to combinatorial interpretations of Ramanujan's congruences modulo 5 and 7. The generating functions for the rank difference  $N(s, \ell, \ell n + d) - N(t, \ell, \ell n + d)$  with  $\ell = 11$  have also been determined by Atkin and Hussain [3], which implies Ramanujan's congruence modulo 11. Since then, the ranks of partitions with moduli other than 5, 7 and 11 have been extensively studied, see, for example, [14, 15, 21, 25].

As to overpartitions, there are two kinds of ranks defined by Lovejoy:  $D$ -rank [17] and  $M_2$ -rank [18]. An overpartition [7] of a nonnegative integer  $n$  is a partition of  $n$  where the first occurrence of each distinct part may be overlined. To define these ranks, let  $l(\lambda)$  be

the largest part of  $\lambda$ ,  $n(\lambda)$  be the number of parts of  $\lambda$  and  $n(\lambda_o)$  be the number of odd non-overlined parts of  $\lambda$ .

Analogous to the rank of a partition, the  $D$ -rank of an overpartition  $\lambda$  is the largest part  $l(\lambda)$  minus the number of parts  $n(\lambda)$ . Similarly, let  $\overline{N}(s, m, n)$  to be the number of overpartitions of  $n$  with rank congruent to  $s$  modulo  $m$ . Lovejoy and Osburn [19] studied the rank differences  $\overline{N}(s, \ell, \ell n + d) - \overline{N}(t, \ell, \ell n + d)$  with  $\ell = 3$  and  $5$  and  $0 \leq d, s, t < \ell$ .

The  $M_2$ -rank of an overpartition  $\lambda$  is defined by

$$M_2\text{-rank}(\lambda) = \left\lceil \frac{l(\lambda)}{2} \right\rceil - n(\lambda) - n(\lambda_o) - \chi(\lambda),$$

where  $\chi(\lambda) = 1$  if  $l(\lambda)$  is odd and non-overlined and  $\chi(\lambda) = 0$  otherwise. Denote by  $\overline{N}_2(m, n)$  the number of overpartitions of  $n$  with  $M_2$ -rank equal to  $m$ . We adopt the notation  $\overline{R2}(z; q)$  in [13] to denote the generating function of  $\overline{N}_2(m, n)$ , that is,

$$\overline{R2}(z; q) = \sum_{n=0}^{\infty} \sum_{m=-\infty}^{\infty} \overline{N}_2(m, n) z^m q^n. \quad (1.1)$$

Lovejoy [18] obtained the following generating function:

$$\overline{R2}(z; q) = \frac{(-q; q)_{\infty}}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(1-z)(1-z^{-1})(-1)^n q^{n^2+2n}}{(1-zq^{2n})(1-z^{-1}q^{2n})}. \quad (1.2)$$

In this paper, we shall study  $M_2$ -ranks of overpartitions modulo 6 and 10. The main results are presented in Theorems 1.1, 1.2 and 1.3 below. We shall adopt the following common notation

$$\begin{aligned} (x_1, x_2, \dots, x_k; q)_{\infty} &= \prod_{n=0}^{\infty} (1 - x_1 q^n)(1 - x_2 q^n) \cdots (1 - x_k q^n), \\ j(z; q) &= (z; q)_{\infty} (q/z; q)_{\infty} (q; q)_{\infty}, \\ J_{a,m} &= j(q^a, q^m), \quad J_m = (q^m; q^m)_{\infty}, \quad \overline{J}_{a,m} = j(-q^a, q^m). \end{aligned}$$

where we assume that  $|q| < 1$ .

Define

$$\begin{aligned} \overline{R}_{s,t}(d, \ell) &= \sum_{n=0}^{\infty} (\overline{N}_2(s, \ell, \ell n/2 + d) + \overline{N}_2(s+1, \ell, \ell n/2 + d) \\ &\quad - \overline{N}_2(t, \ell, \ell n/2 + d) - \overline{N}_2(t+1, \ell, \ell n/2 + d)) q^n. \end{aligned}$$

**Theorem 1.1.** *We have*

$$\overline{R}_{0,2}(0, 6) = \frac{J_6^9 J_{2,4}^2}{J_{1,6}^6 J_2^3 J_{3,6}}, \quad (1.3)$$

$$\bar{R}_{0,2}(1, 6) = \frac{2J_6^9 J_{2,4}^2}{J_{1,6}^5 J_2^3 J_{3,6}^2}, \quad (1.4)$$

$$\bar{R}_{0,2}(2, 6) = \frac{4J_6^9 J_{2,4}^2}{J_{1,6}^4 J_2^3 J_{3,6}^3} + \frac{2q^2}{J_{3,6}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+6n}}{1+q^{6n+4}}. \quad (1.5)$$

**Theorem 1.2.** *We have*

$$\bar{R}_{0,4}(i, 10) = \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} A_i(q), \quad 0 \leq i \leq 3, \quad (1.6)$$

$$\bar{R}_{0,4}(4, 10) = \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} A_4(q) + \frac{q^4}{J_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2+10n}}{1+q^{10n+8}}, \quad (1.7)$$

where

$$\begin{aligned} A_0(q) = & \frac{J_2^4 J_{3,20}^2 J_{5,20}^8 J_{7,20}^2}{2J_{4,20}^4 J_{6,20}^3 J_{8,20} J_{20}^6} + \frac{2qJ_1^3 J_{10} J_{5,20} J_{20}^5}{J_{1,20}^2 J_{2,20} J_{8,20}^3 J_{9,20}^2} + \frac{8q^2 J_1^2 J_2^4 J_{2,20} J_{3,20} J_{7,20}}{J_{4,20}^2 J_{6,20} J_{8,20} J_{10,20}^2 J_{20}} \\ & - \frac{16q^2 J_1^2 J_{2,20} J_{3,20}^2 J_{6,20} J_{7,20}^2}{J_{4,20} J_{5,20}^2 J_{20}^3}, \end{aligned} \quad (1.8)$$

$$\begin{aligned} A_1(q) = & \frac{J_2^4 J_{3,20}^3 J_{5,20}^6 J_{7,20}^3}{J_{4,20}^4 J_{6,20}^3 J_{8,20} J_{20}^6} + \frac{8qJ_1^2 J_{10}^2 J_{2,20}^2 J_{3,20}^2 J_{7,20}^2 J_{8,20}}{J_{1,20} J_{4,20}^2 J_{6,20} J_{9,20} J_{20}^4} - \frac{8qJ_1^2 J_{2,20}^4 J_{6,20}^2 J_{8,20}^3}{J_{1,20}^2 J_{3,20} J_{7,20} J_{9,20}^2 J_{20}^3} \\ & + \frac{4q^2 J_1 J_2^2 J_{1,20}^2 J_{5,20}^3 J_{9,20}^2}{J_{4,20}^2 J_{8,20} J_{20}^4 J_{10}} - \frac{8q^3 J_1^2 J_{1,20}^2 J_{4,20} J_{6,20} J_{9,20}^2 J_{10,20}}{J_{5,20}^2 J_{8,20}^2 J_{20}^3}, \end{aligned} \quad (1.9)$$

$$\begin{aligned} A_2(q) = & \frac{2J_2^4 J_{3,20}^4 J_{5,20}^4 J_{7,20}^4}{J_{4,20}^4 J_{6,20}^3 J_{8,20} J_{20}^6} + \frac{2qJ_1^2 J_{2,20} J_{4,20}^4 J_{6,20}}{J_{3,20} J_{4,20} J_{7,20} J_{20}^3} - \frac{8qJ_1^2 J_{10}^2 J_{3,20}^3 J_{4,20} J_{6,20} J_{7,20}^3}{J_{1,20} J_{5,20}^2 J_{8,20}^2 J_{9,20} J_{20}^4} \\ & + \frac{4q^2 J_1^2 J_{10}^2 J_{1,20} J_{4,20} J_{6,20} J_{9,20}}{J_{8,20}^2 J_{20}^4} + \frac{8q^2 J_1^2 J_{1,20} J_{2,20} J_{6,20} J_{9,20}}{J_{4,20} J_{20}^3}, \end{aligned} \quad (1.10)$$

$$\begin{aligned} A_3(q) = & -\frac{J_1 J_2^2 J_{5,20}^7}{J_{10} J_{4,20} J_{8,20} J_{20}^4} + \frac{4J_2^4 J_{3,20}^5 J_{5,20}^2 J_{7,20}^5}{J_{4,20}^4 J_{6,20}^3 J_{8,20} J_{20}^6} + \frac{2qJ_1^2 J_{10}^2 J_{2,20}^2 J_{5,20}^2 J_{8,20}}{J_{4,20}^2 J_{6,20} J_{20}^4} \\ & - \frac{12qJ_1^2 J_{2,20} J_{5,20}^2 J_{6,20}}{J_{4,20} J_{20}^3} - \frac{16q^2 J_1^3 J_{20}^6}{J_{4,20}^2 J_{5,20}^3 J_{8,20} J_{10}} + \frac{8q^2 J_1^3 J_{10} J_{2,20} J_{5,20}^2}{J_{4,20}^3 J_{5,20}^3 J_{6,20}^2}, \end{aligned} \quad (1.11)$$

$$\begin{aligned} A_4(q) = & -\frac{2J_1^2 J_{2,20} J_{5,20}^4 J_{6,20}}{J_{1,20} J_{4,20} J_{9,20} J_{20}^3} - \frac{J_1^2 J_{10}^2 J_{2,20}^2 J_{5,20}^4 J_{8,20}}{J_{1,20} J_{4,20}^2 J_{6,20} J_{9,20} J_{20}^4} + \frac{8J_2^4 J_{3,20}^6 J_{7,20}^6}{J_{4,20}^4 J_{6,20}^3 J_{8,20} J_{20}^6} \\ & - \frac{16qJ_1^2 J_{2,20} J_{3,20} J_{6,20} J_{7,20}}{J_{4,20} J_{20}^3} - \frac{8qJ_1^3 J_{10} J_{20}^5 J_{2,20}}{J_{1,20} J_{4,20}^3 J_{5,20} J_{6,20}^2 J_{9,20}} - \frac{16q^3 J_1 J_2^2 J_{1,20}^4 J_{9,20}^4}{J_{10} J_{20}^4 J_{4,20}^2 J_{5,20} J_{8,20}}. \end{aligned} \quad (1.12)$$

**Theorem 1.3.** *We have*

$$\bar{R}_{1,3}(i, 10) = \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_i(q), \quad i = 0, 2, 3, \quad (1.13)$$

$$\bar{R}_{1,3}(1, 10) = \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_1(q) - \frac{q^4}{J_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2+10n}}{1+q^{10n+6}}, \quad (1.14)$$

$$\bar{R}_{1,3}(4, 10) = \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_4(q) - \frac{q^4}{J_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2+10n}}{1+q^{10n+8}}, \quad (1.15)$$

where

$$\begin{aligned} B_0(q) = & -\frac{qJ_1^5 J_{1,20} J_{5,20} J_{9,20}}{J_{10} J_{20} J_{4,20}^2 J_{6,20} J_{8,20}} + \frac{4qJ_1^4 J_{20}^2 J_{3,20} J_{7,20}}{J_{1,20} J_{4,20}^2 J_{8,20} J_{9,20} J_{10,20}} - \frac{16q^2 J_2^{\frac{7}{2}} J_{1,2}^{\frac{1}{2}} J_{1,20} J_{2,20} J_{9,20}}{J_{20} J_{4,20} J_{5,20}^3} \\ & + \frac{16q^2 J_1^2 J_{2,20} J_{3,20}^2 J_{6,20} J_{7,20}^2}{J_{20}^3 J_{4,20} J_{5,20}^2} - \frac{4q^3 J_1^4 J_{20}^2 J_{1,20} J_{9,20}}{J_{3,20} J_{4,20} J_{7,20} J_{8,20}^2 J_{10,20}} - \frac{8q^3 J_1 J_2^2 J_{1,20}^2 J_{5,20}^3 J_{9,20}^2}{J_{10} J_{20}^4 J_{4,20} J_{8,20}^2} \\ & - \frac{16q^4 J_{1,20}^2 J_{2,20} J_{6,20} J_{9,20}^2 J_1^2}{J_{20}^3 J_{5,20}^2 J_{8,20}}, \end{aligned} \quad (1.16)$$

$$\begin{aligned} B_1(q) = & -\frac{2qJ_1^2 J_{10}^2 J_{2,20} J_{5,20}^2}{J_{20}^4 J_{4,20}} + \frac{8qJ_1^3 J_{2,20} J_{3,20}^3 J_{6,20} J_{7,20}^3}{J_{10} J_{4,20} J_{5,20} J_{20}^6} + \frac{2q^2 J_2^4 J_{1,20} J_{2,20} J_{5,20}^2 J_{6,20} J_{9,20}}{J_{20}^6 J_{8,20} J_{10,20}} \\ & + \frac{4q^2 J_2^4 J_{1,20}^2 J_{2,20} J_{6,20} J_{9,20}^2}{J_{20}^6 J_{4,20} J_{10,20}} - \frac{16q^2 J_1^3 J_{10} J_{20}^5}{J_{4,20}^2 J_{5,20}^3 J_{6,20} J_{8,20}} + \frac{12q^3 J_1^2 J_{1,20}^2 J_{2,20} J_{6,20} J_{9,20}^2}{J_{20}^3 J_{4,20} J_{5,20}^2} \\ & + \frac{14q^3 J_1^2 J_{1,20} J_{2,20} J_{6,20} J_{9,20}}{J_{20}^3 J_{8,20}} - \frac{16q^4 J_1^2 J_{10}^2 J_{4,20}^4 J_{2,20} J_{9,20}^4}{J_{20}^4 J_{3,20} J_{4,20} J_{5,20}^4 J_{7,20}}, \end{aligned} \quad (1.17)$$

$$\begin{aligned} B_2(q) = & \frac{qJ_1^3 J_{2,20} J_{5,20}^5 J_{6,20}}{J_{10} J_{20}^6 J_{8,20}} + \frac{4qJ_1^2 J_{10}^2 J_{1,20}^2 J_{4,20}^2 J_{6,20}^3 J_{9,20}^2}{J_{20}^4 J_{2,20}^2 J_{3,20} J_{7,20} J_{8,20}^3} - \frac{8qJ_2^4 J_{1,20} J_{3,20}^5 J_{7,20}^5 J_{9,20}}{J_{20}^6 J_{4,20}^4 J_{6,20}^3 J_{8,20}} \\ & + \frac{12qJ_1 J_2^2 J_{3,20}^4 J_{7,20}^4}{J_{10} J_{20}^4 J_{4,20}^2 J_{5,20} J_{8,20}} + \frac{4q^2 J_1 J_{10} J_{2,20}^6 J_{6,20}^2 J_{8,20}^5}{J_{5,20} J_{20}^{12}} - \frac{12q^2 J_1^2 J_{2,20} J_{5,20}^2 J_{6,20}}{J_{20}^3 J_{8,20}} \\ & - \frac{16q^2 J_1^2 J_{1,20} J_{2,20} J_{6,20} J_{9,20}}{J_{20}^3 J_{4,20}} - \frac{8q^3 J_1 J_2^2 J_{10} J_{2,20}^4 J_{6,20}}{J_{20}^5 J_{4,20} J_{5,20}}, \end{aligned} \quad (1.18)$$

$$\begin{aligned} B_3(q) = & \frac{J_1^3 J_{2,20} J_{5,20}^5 J_{6,20}}{J_{10} J_{20}^6 J_{4,20}} - \frac{8qJ_1^2 J_{10}^2 J_{2,20}^3 J_{3,20}^2 J_{7,20}^2 J_{8,20}^2}{J_{20}^4 J_{1,20} J_{4,20}^3 J_{6,20}^2 J_{9,20}} + \frac{12qJ_1^2 J_{2,20} J_{5,20}^2 J_{6,20}}{J_{20}^3 J_{4,20}} \\ & + \frac{4q^2 J_2^4 J_{1,20}^6 J_{5,20}^2 J_{6,20} J_{9,20}^6}{J_{20}^6 J_{4,20}^4 J_{3,20} J_{7,20} J_{8,20}^5} - \frac{16q^3 J_1^3 J_{20}^6 J_{3,20} J_{7,20}}{J_{10} J_{1,20} J_{5,20}^3 J_{8,20}^3 J_{9,20}} - \frac{16q^4 J_1 J_2^2 J_{1,20}^4 J_{9,20}^4}{J_{10} J_{20}^4 J_{10,20} J_{5,20} J_{8,20}^2}, \end{aligned} \quad (1.19)$$

$$\begin{aligned}
B_4(q) = & \frac{2J_1^2 J_{2,20} J_{5,20}^4 J_{6,20}}{J_{20}^3 J_{1,20} J_{4,20} J_{9,20}} - \frac{2q J_1^2 J_{10,20}^2 J_{2,20}^5 J_{5,20}^2 J_{6,20}^2 J_{8,20}}{J_{20}^4 J_{1,20}^3 J_{3,20} J_{7,20} J_{9,20}^3} + \frac{16q^2 J_1^2 J_{10,20}^2 J_{1,20}^2 J_{2,20} J_{9,20}^2}{J_{20}^4 J_{4,20} J_{5,20}^2} \\
& - \frac{16q^2 J_1^2 J_{2,20} J_{3,20}^2 J_{6,20} J_{7,20}^2}{J_{20}^3 J_{5,20}^2 J_{8,20}} + \frac{8q^3 J_1^2 J_{1,20}^2 J_{2,20} J_{6,20} J_{9,20}^2}{J_{20}^3 J_{3,20} J_{7,20} J_{8,20}} + \frac{16q^3 J_1 J_2^2 J_{1,20}^4 J_{9,20}^4}{J_{10} J_{20}^4 J_{4,20}^2 J_{5,20} J_{8,20}} \\
& + \frac{32q^4 J_1^2 J_{2,20}^3 J_{3,20}^2 J_{7,20}^2}{J_{10}^2 J_{20}^2 J_{4,20} J_{5,20}^2}. \tag{1.20}
\end{aligned}$$

The generating functions of rank differences often lead to rank inequalities, see, for example, Andrews [1], Garvan [9], Mao [21, 22]. In this paper, we obtain the following inequalities between the  $M_2$ -ranks of overpartitions modulo 6 with the aid of the generating functions in Theorem 1.1.

**Theorem 1.4.** *For  $n \geq 1$ , we have*

$$\overline{N}_2(0, 6, 3n) \geq \overline{N}_2(2, 6, 3n), \tag{1.21}$$

$$\overline{N}_2(1, 6, 3n) \geq \overline{N}_2(3, 6, 3n), \tag{1.22}$$

$$\overline{N}_2(0, 6, 3n + 1) \geq \overline{N}_2(2, 6, 3n + 1), \tag{1.23}$$

$$\overline{N}_2(1, 6, 3n + 1) \geq \overline{N}_2(3, 6, 3n + 1), \tag{1.24}$$

$$\overline{N}_2(0, 6, 3n) + \overline{N}_2(3, 6, 3n) \geq \overline{N}_2(1, 6, 3n) + \overline{N}_2(2, 6, 3n), \tag{1.25}$$

$$\overline{N}_2(0, 6, 3n + 1) + \overline{N}_2(3, 6, 3n + 1) \geq \overline{N}_2(1, 6, 3n + 1) + \overline{N}_2(2, 6, 3n + 1), \tag{1.26}$$

$$\overline{N}_2(0, 6, 3n + 2) + \overline{N}_2(3, 6, 3n + 2) \geq \overline{N}_2(1, 6, 3n + 2) + \overline{N}_2(2, 6, 3n + 2). \tag{1.27}$$

The connection between classical mock theta functions and the generating functions of rank differences of partitions have been extensively studied, see [2, 11, 20]. We find a relation between the generating function of  $M_2$ -ranks of overpartitions modulo 6 and the third order mock theta function  $\rho(q)$  defined by

$$\rho(q) = \sum_{n=0}^{\infty} \frac{q^{2n(n+1)}(q; q^2)_{n+1}}{(q^3; q^6)_{n+1}},$$

see [23].

**Theorem 1.5.** *We have*

$$\overline{R}_{0,2}(2, 6) = 2\rho(-q) + M_0(q), \tag{1.28}$$

where  $M_0(q)$  is an (explicit) weakly holomorphic modular form given by

$$M_0(q) = -\frac{2J_6^3 J_2 \overline{J}_{1,6}}{J_{1,6} J_{3,6} \overline{J}_2^2} + \frac{4J_6^9 J_{2,4}^2}{J_{1,6}^4 J_2^3 J_{3,6}^3}.$$

In light of Theorem 1.2 and Theorem 1.3, we deduce the following relations on the generating function of  $M_2$ -ranks of overpartitions modulo 10 and the tenth order mock theta functions  $\phi(q)$  and  $\psi(q)$  are defined as follows,

$$\phi(q) = \sum_{n=0}^{\infty} \frac{q^{\binom{n+1}{2}}}{(q; q^2)_{n+1}}, \quad \psi(q) = \sum_{n=0}^{\infty} \frac{q^{\binom{n+2}{2}}}{(q; q^2)_{n+1}},$$

see [6].

**Theorem 1.6.** *We have*

$$\overline{R}_{0,4}(4, 10) = q^{-1}\psi(-q) + M_1(q), \quad (1.29)$$

$$\overline{R}_{1,3}(1, 10) = \phi(-q) + M_2(q), \quad (1.30)$$

$$\overline{R}_{1,3}(4, 10) = -q^{-1}\psi(-q) + M_3(q), \quad (1.31)$$

where  $M_1(q)$ ,  $M_2(q)$  and  $M_3(q)$  are (explicit) weakly holomorphic modular forms given by

$$\begin{aligned} M_1(q) &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} A_4(q) + \frac{J_{10}^3 J_{1,10} \overline{J}_{4,10}}{J_{4,10} J_{5,10} \overline{J}_{1,10} \overline{J}_{2,10}} - \frac{q J_{10}^3 J_{1,10} \overline{J}_{2,10}}{J_{4,10} J_{5,10} \overline{J}_{2,10} \overline{J}_{3,10}}, \\ M_2(q) &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_1(q) - \frac{J_{10}^3 J_{3,10} \overline{J}_{4,10}}{J_{2,10} J_{5,10} \overline{J}_{1,10} \overline{J}_{4,10}} + \frac{q J_{10}^3 J_{3,10} \overline{J}_{2,10}}{J_{2,10} J_{5,10} \overline{J}_{3,10} \overline{J}_{4,10}}, \\ M_3(q) &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_4(q) - \frac{J_{10}^3 \overline{J}_{4,10} J_{1,10}}{J_{4,10} J_{5,10} \overline{J}_{1,10} \overline{J}_{2,10}} + \frac{q J_{10}^3 J_{1,10} \overline{J}_{2,10}}{J_{4,10} J_{5,10} \overline{J}_{2,10} \overline{J}_{3,10}}, \end{aligned}$$

and  $A_4(q)$  is defined in Theorem 1.2,  $B_1(q)$  and  $B_4(q)$  are defined in Theorem 1.3.

This paper is organized as follows. We prove Theorem 1.1 in Section 2. Theorems 1.2 and 1.3 are proved in Section 3. In Section 4, we derive inequalities on  $M_2$ -ranks of overpartitions mod 6 from the generating function in Theorem 1.1. In Section 5, we obtain relations between the generating functions of  $M_2$ -rank differences of overpartitions and mock theta functions as given in Theorems 1.5 and 1.6.

## 2 Proof of Theorem 1.1

The following lemma plays a central role in the proof of Theorem 1.1.

**Lemma 2.1.** *We have*

$$\begin{aligned} \overline{R}2(\exp(\pi i/3); q) &= \sum_{n=0}^{\infty} (\overline{N}_2(0, 6, n) + \overline{N}_2(1, 6, n) - \overline{N}_2(2, 6, n) - \overline{N}_2(3, 6, n)) q^n \\ &= \frac{2(-q; q)_{\infty}}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{6n}}. \end{aligned} \quad (2.1)$$

*Proof.* Replacing  $z$  by  $\xi_6 = e^{\frac{\pi i}{3}}$ , the right-hand side of (1.1) simplifies to

$$\begin{aligned}
\overline{R2}(\exp(\pi i/3); q) &= \sum_{n=0}^{\infty} \sum_{m=-\infty}^{\infty} \overline{N}_2(m, n) \xi_6^m q^n \\
&= \sum_{n=0}^{\infty} \sum_{t=0}^5 \sum_{m=-\infty}^{\infty} \overline{N}_2(6m+t, n) \xi_6^t q^n \\
&= \sum_{n=0}^{\infty} (\overline{N}_2(0, 6, n) + \xi_6 \overline{N}_2(1, 6, n) + \xi_6^2 \overline{N}_2(2, 6, n) \\
&\quad + \xi_6^3 \overline{N}_2(3, 6, n) + \xi_6^4 \overline{N}_2(4, 6, n) + \xi_6^5 \overline{N}_2(5, 6, n)) q^n. \tag{2.2}
\end{aligned}$$

In view of the symmetry property

$$\overline{N}_2(m, n) = \overline{N}_2(-m, n), \text{ for } m \geq 0$$

due to Lovejoy and Osburn [20], we see that

$$\overline{N}_2(s, m, n) = \overline{N}_2(m-s, m, n). \tag{2.3}$$

In particular, for  $m \geq 0$ ,

$$\overline{N}_2(1, 6, n) = \overline{N}_2(5, 6, n), \tag{2.4}$$

and

$$\overline{N}_2(2, 6, n) = \overline{N}_2(4, 6, n). \tag{2.5}$$

Substituting (2.4) and (2.5) into (2.2), we find that

$$\begin{aligned}
\overline{R2}(\exp(\pi i/3); q) &= \sum_{n=0}^{\infty} (\overline{N}_2(0, 6, n) + (\xi_6 + \xi_6^5) \overline{N}_2(1, 6, n) \\
&\quad + (\xi_6^2 + \xi_6^4) \overline{N}_2(2, 6, n) + \xi_6^3 \overline{N}_2(3, 6, n)) q^n. \tag{2.6}
\end{aligned}$$

Observing that  $1 - \xi_6 + \xi_6^2 = 0$  and  $\xi_6^3 = -1$ , (2.6) yields that

$$\overline{R2}(\exp(\pi i/3); q) = \sum_{n=0}^{\infty} (\overline{N}_2(0, 6, n) + \overline{N}_2(1, 6, n) - \overline{N}_2(2, 6, n) - \overline{N}_2(3, 6, n)) q^n.$$

To evaluate the right-hand side of (1.2) for  $z = \xi_6$ , we get

$$\begin{aligned}
\overline{R2}(\exp(\pi i/3); q) &= \frac{(-q; q)_{\infty}}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(1 - \xi_6)(1 - \xi_6^{-1})(-1)^n q^{n^2+2n}}{(1 - \xi_6 q^{2n})(1 - \xi_6^{-1} q^{2n})} \\
&= \frac{(-q; q)_{\infty}}{(q; q)_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(2 - \xi_6^{-1} - \xi_6)(-1)^n q^{n^2+2n}}{1 - \xi_6^{-1} q^{2n} - \xi_6 q^{2n} + q^{4n}}.
\end{aligned}$$

But  $1 - \xi_6^{-1} - \xi_6 = 0$ , so we obtain that

$$\begin{aligned} \overline{R2}(\exp(\pi i/3); q) &= \frac{(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 - q^{2n} + q^{4n}} \\ &= \frac{(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}(1 + q^{2n})}{1 + q^{6n}} \\ &= \frac{2(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{6n}}, \end{aligned}$$

as desired. This completes the proof of Lemma 2.1.  $\blacksquare$

Now we are in a position to prove Theorem 1.1.

*Proof of Theorem 1.1.* By Lemma 2.1, it suffices to show that

$$\begin{aligned} &\frac{(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{6n}} \\ &= \frac{J_{18}^9 J_{6,12}^2}{2J_{3,18}^6 J_6^3 J_{9,18}} + q \frac{J_{18}^9 J_{6,12}^2}{J_{3,18}^5 J_6^3 J_{9,18}^2} \\ &\quad + q^2 \left\{ \frac{2J_{18}^9 J_{6,12}^2}{J_{3,18}^4 J_6^3 J_{9,18}^3} + \frac{q^6}{J_{9,18}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{9n^2+18n}}{1 + q^{18n+12}} \right\}. \end{aligned} \quad (2.7)$$

The summation on the left hand side of (2.7) can be written as

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{6n}} = \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{9n^2+6n}}{1 + q^{18n}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{9n^2+12n+3}}{1 + q^{18n+6}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{9n^2+18n+8}}{1 + q^{18n+12}}. \quad (2.8)$$

Let  $G_0, G_1$  and  $G_2$  denote the summations on the right hand side of (2.8), respectively. Recall the following identity [20, Lemma 3.1]:

$$\begin{aligned} &\sum_{n=-\infty}^{\infty} (-1)^n q^{n^2+2n} \left[ \frac{\zeta^{-2n}}{1 - z^2 \zeta^{-2} q^{2n}} + \frac{\zeta^{2n+4}}{1 - z^2 \zeta^2 q^{2n}} \right] \\ &= \frac{-2(\zeta^4, q^2 \zeta^{-4}; q^2)_\infty (-q; q)_\infty^2}{(-\zeta^2, -q \zeta^{-2}; q)_\infty (\zeta^{-2}, q^2 \zeta^2; q^2)_\infty} \sum_{n=-\infty}^{\infty} (-1)^n \frac{q^{n^2+2n}}{1 - z^2 q^{2n}} \\ &\quad + \frac{(-z^2, -q z^{-2}; q)_\infty (\zeta^4, q^2 \zeta^{-4}, \zeta^2, q^2 \zeta^{-2}; q^2)_\infty (q^2; q^2)_\infty^2}{(-\zeta^2, -q \zeta^{-2}; q)_\infty (z^2 \zeta^2, q^2 z^{-2} \zeta^{-2}, z^2 \zeta^{-2}, q^2 z^{-2} \zeta^2, z^2, q^2 z^{-2}; q^2)_\infty}. \end{aligned} \quad (2.9)$$

Substituting  $q, z^2$  and  $\zeta^2$  by  $q^9, -q^{12}$  and  $q^{12}$ , respectively, in (2.9), we find that

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{9n^2+18n} \left[ \frac{q^{-12n}}{1 + q^{18n}} + \frac{q^{12n+24}}{1 + q^{18n+24}} \right]$$



$$\begin{aligned}
&= \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{9n^2+6n}}{1+q^{18n}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{9n^2+12n+3}}{1+q^{18n+6}} \\
&= -\frac{2q^9 J_{3,18}}{J_{9,18}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{9n^2+18n}}{1+q^{18n+12}} + \frac{J_6 J_{3,18}^2 J_{6,12}^2}{2J_{18}^3},
\end{aligned}$$

which can be recast as

$$G_0 - G_1 = -\frac{2qJ_{3,18}}{J_{9,18}}G_2 + \frac{J_6 J_{3,18}^2 J_{6,12}^2}{2J_{18}^3}. \quad (2.10)$$

Combining (2.10) into (2.8), we are led to the relation

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1+q^{6n}} = \left(1 - \frac{2qJ_{3,18}}{J_{9,18}}\right) G_2 + \frac{J_6 J_{3,18}^2 J_{6,12}^2}{2J_{18}^3}. \quad (2.11)$$

Applying the following identity [19, Lemma 3.1]

$$\frac{(q; q)_{\infty}}{(-q; q)_{\infty}} = \frac{(q^9; q^9)_{\infty}}{(-q^9; q^9)_{\infty}} - 2q(q^3, q^{15}, q^{18}; q^{18})_{\infty} = J_{9,18} - 2qJ_{3,18} \quad (2.12)$$

to (2.11), we get

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1+q^{6n}} = \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \cdot \frac{G_2}{J_{9,18}} + \frac{J_6 J_{3,18}^2 J_{6,12}^2}{2J_{18}^3}.$$

To complete the proof of Theorem 1.1, it remains to show that

$$\frac{J_6 J_{3,18}^2 J_{6,12}^2}{J_{18}^3} = \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \left\{ \frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^6 J_{9,18}} + q \frac{2J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^5 J_{9,18}^2} + q^2 \frac{4J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^4 J_{9,18}^3} \right\}. \quad (2.13)$$

In view of (2.12), the right hand side of (2.13) can be rewritten as

$$(J_{9,18} - 2qJ_{3,18}) \left\{ \frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^6 J_{9,18}} + q \frac{2J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^5 J_{9,18}^2} + q^2 \frac{4J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^4 J_{9,18}^3} \right\},$$

which equals

$$\begin{aligned}
&\left( \frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^6} - 8q^3 \frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^3 J_{9,18}^3} \right) + q \left( \frac{2J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^5 J_{9,18}} - \frac{2J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^5 J_{9,18}} \right) \\
&+ q^2 \left( \frac{4J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^4 J_{9,18}^2} - \frac{4J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^4 J_{9,18}^2} \right) \\
&= \frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^6} - 8q^3 \frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^3 J_{9,18}^3}.
\end{aligned}$$

Hence, (2.13) is equivalent to

$$\frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^6} - 8q^3 \frac{J_{18}^9 J_{6,12}^2}{J_6^3 J_{3,18}^3 J_{9,18}^3} = \frac{J_6 J_{3,18}^2 J_{6,12}^2}{J_{18}^3}, \quad (2.14)$$

which can be simplified to

$$J_{9,18}^3 - 8q^3 J_{3,18}^3 = \frac{J_6^4 J_{3,18}^8 J_{9,18}^3}{J_{18}^{12}}. \quad (2.15)$$

The above identity can be deduced from the following identity [4] by setting  $q = q^9$ ,  $x = -q^3$ ,  $y = q^3$  and  $z = -1$ :

$$j(x, q)^2 j(yz, q) j(yz^{-1}, q) = j(y, q)^2 j(xz, q) j(xz^{-1}, q) - yz^{-1} j(z, q)^2 j(xy, q) j(xy^{-1}, q).$$

This completes the proof of Theorem 1.1. ■

### 3 Proofs of Theorem 1.2 and Theorem 1.3

To prove Theorems 1.2 and 1.3, we need the following expression for  $\overline{R2}(\exp(\pi i/5); q)$ .

**Lemma 3.1.** *We have*

$$\overline{R2}(\exp(\pi i/5); q) = F_1(q) + (\xi_{10}^2 - \xi_{10}^3) F_2(q), \quad (3.1)$$

where

$$F_1(q) = \frac{2(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{10n}}, \quad (3.2)$$

$$F_2(q) = \frac{2(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n} (q^{2n} - 1)}{1 + q^{10n}}. \quad (3.3)$$

*Proof.* With the substitution  $z = \xi_{10}$ , the right-hand side of (1.2) equals

$$\begin{aligned} & \frac{(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(1 - \xi_{10})(1 - \xi_{10}^{-1})(-1)^n q^{n^2+2n}}{(1 - \xi_{10} q^{2n})(1 - \xi_{10}^{-1} q^{2n})} \\ &= \frac{(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(1 - \xi_{10})(1 - \xi_{10}^{-1})(1 - \xi_{10}^3 q^{2n})(1 - \xi_{10}^{-3} q^{2n})(1 + q^{2n})(-1)^n q^{n^2+2n}}{(1 - \xi_{10} q^{2n})(1 - \xi_{10}^{-1} q^{2n})(1 - \xi_{10}^3 q^{2n})(1 - \xi_{10}^{-3} q^{2n})(1 + q^{2n})}. \end{aligned} \quad (3.4)$$

Since  $1 - \xi_{10} - \xi_{10}^{-1} - \xi_{10}^3 - \xi_{10}^{-3} = 0$ , we deduce that

$$(1 - \xi_{10} q^{2n})(1 - \xi_{10}^{-1} q^{2n})(1 - \xi_{10}^3 q^{2n})(1 - \xi_{10}^{-3} q^{2n})(1 + q^{2n}) = 1 + q^{10n}$$

and

$$\begin{aligned}
& (1 - \xi_{10})(1 - \xi_{10}^{-1})(1 - \xi_{10}^3 q^{2n})(1 - \xi_{10}^{-3} q^{2n})(1 + q^{2n}) \\
&= (1 - \xi_{10}^2 + \xi_{10}^3) + (-1 + \xi_{10}^{-1} + \xi_{10})q^{2n} + (-1 + \xi_{10}^{-1} + \xi_{10})q^{4n} + (1 - \xi_{10}^2 + \xi_{10}^3)q^{6n} \\
&= (1 - \xi_{10}^2 + \xi_{10}^3) + (\xi_{10}^2 - \xi_{10}^3)q^{2n} + (\xi_{10}^2 - \xi_{10}^3)q^{4n} + (1 - \xi_{10}^2 + \xi_{10}^3)q^{6n}.
\end{aligned}$$

Thus the right hand side of (3.4) equals

$$\begin{aligned}
& \overline{R2}(\exp(\pi i/5); q) \\
&= \frac{(-q; q)_\infty}{(q; q)_\infty} \left\{ \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}(1 + q^{6n})}{1 + q^{10n}} \right. \\
&\quad \left. + (\xi_{10}^2 - \xi_{10}^3) \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}(q^{2n} + q^{4n} - 1 - q^{6n})}{1 + q^{10n}} \right\} \\
&= \frac{(-q; q)_\infty}{(q; q)_\infty} \left\{ \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{10n}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^{-n} q^{n^2-8n}}{1 + q^{-10n}} \right\} \\
&\quad + (\xi_{10}^2 - \xi_{10}^3) \frac{(-q; q)_\infty}{(q; q)_\infty} \left\{ \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+4n}}{1 + q^{10n}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^{-n} q^{n^2-6n}}{1 + q^{-10n}} \right. \\
&\quad \left. - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{10n}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^{-n} q^{n^2-8n}}{1 + q^{-10n}} \right\} \\
&= \frac{2(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1 + q^{10n}} \\
&\quad - (\xi_{10}^2 - \xi_{10}^3) \frac{2(-q; q)_\infty}{(q; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}(1 - q^{2n})}{1 + q^{10n}}.
\end{aligned}$$

This completes the proof. ■

**Lemma 3.2.** *We have*

$$\sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + \overline{N}_2(1, 10, n) - \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n))q^n = F_1(q) \quad (3.5)$$

and

$$\sum_{n=0}^{\infty} (\overline{N}_2(1, 10, n) + \overline{N}_2(2, 10, n) - \overline{N}_2(3, 10, n) - \overline{N}_2(4, 10, n))q^n = F_2(q), \quad (3.6)$$

where  $F_1(q)$  and  $F_2(q)$  are given in (3.2) and (3.3).

*Proof.* Setting  $z = \xi_{10} = e^{\frac{\pi i}{5}}$ , we get

$$\begin{aligned}\overline{R2}(\exp(\pi i/5); q) &= \sum_{n=0}^{\infty} \sum_{m=-\infty}^{\infty} \overline{N}_2(m, n) \xi_{10}^m q^n \\ &= \sum_{n=0}^{\infty} \sum_{t=0}^9 \sum_{m=-\infty}^{\infty} \overline{N}_2(10m + t, n) \xi_{10}^t q^n \\ &= \sum_{n=0}^{\infty} \sum_{t=0}^9 \overline{N}_2(t, 10, n) \xi_{10}^t q^n.\end{aligned}$$

Utilizing (2.3) and the fact that  $\xi_{10}^5 = -1$ , we deduce that

$$\begin{aligned}\overline{R2}(\exp(\pi i/5); q) &= \sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + (\xi_{10} - \xi_{10}^4) \overline{N}_2(1, 10, n) + (\xi_{10}^2 - \xi_{10}^3) \overline{N}_2(2, 10, n) \\ &\quad + (\xi_{10}^3 - \xi_{10}^2) \overline{N}_2(3, 10, n) + (\xi_{10}^4 - \xi_{10}) \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n)) q^n.\end{aligned}$$

But  $1 - \xi_{10} + \xi_{10}^2 - \xi_{10}^3 + \xi_{10}^4 = 0$ , so we arrive at

$$\begin{aligned}\overline{R2}(\exp(\pi i/5); q) &= \sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + (1 + \xi_{10}^2 - \xi_{10}^3) \overline{N}_2(1, 10, n) + (\xi_{10}^2 - \xi_{10}^3) \overline{N}_2(2, 10, n) \\ &\quad + (\xi_{10}^3 - \xi_{10}^2) \overline{N}_2(3, 10, n) - (1 + \xi_{10}^2 - \xi_{10}^3) \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n)) q^n \\ &= \sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + \overline{N}_2(1, 10, n) - \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n)) q^n \\ &\quad + (\xi_{10}^2 - \xi_{10}^3) \sum_{n=0}^{\infty} (\overline{N}_2(1, 10, n) + \overline{N}_2(2, 10, n) - \overline{N}_2(3, 10, n) - \overline{N}_2(4, 10, n)) q^n.\end{aligned}\tag{3.7}$$

Combining (3.1) with (3.7), we have

$$\begin{aligned}F_1(q) - \sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + \overline{N}_2(1, 10, n) - \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n)) q^n \\ = (\xi_{10}^2 - \xi_{10}^3) \left( \sum_{n=0}^{\infty} (\overline{N}_2(1, 10, n) + \overline{N}_2(2, 10, n) - \overline{N}_2(3, 10, n) - \overline{N}_2(4, 10, n)) q^n - F_2(q) \right).\end{aligned}\tag{3.8}$$

Let us rewrite (3.8) as

$$A(q) = (\xi_{10}^2 - \xi_{10}^3) B(q).\tag{3.9}$$

where

$$A(q) = \sum_{n=0}^{\infty} a_n q^n = F_1(q) - \sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + \overline{N}_2(1, 10, n) - \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n)) q^n,$$

and

$$B(q) = \sum_{n=0}^{\infty} b_n q^n = \sum_{n=0}^{\infty} (\overline{N}_2(1, 10, n) + \overline{N}_2(2, 10, n) - \overline{N}_2(3, 10, n) - \overline{N}_2(4, 10, n)) q^n - F_2(q).$$

From (3.9) it follows that for  $n \geq 0$ ,

$$a_n = (\xi_{10}^2 - \xi_{10}^3) b_n.$$

It is known that the minimal polynomial over  $\mathbb{Q}$  of  $\xi_{10}$  is  $1 + \xi_{10} + \xi_{10}^2 + \xi_{10}^3 + \xi_{10}^4$ . This implies that  $\xi_{10}^2 - \xi_{10}^3$  is not a rational number. On the other hand,  $a_n$  and  $b_n$  are integers for  $n \geq 0$ . So we conclude that  $a_n = b_n = 0$  for  $n \geq 0$ . This completes the proof.  $\blacksquare$

The following lemmas will be used in the proofs of Theorems 1.2 and 1.3.

**Lemma 3.3.** *Let*

$$U_1 = \frac{J_{10,100}^4 J_{15,100}^2 J_{30,100}^2 J_{35,100}^3 J_{40,100}^2 J_{50,100}^2}{J_{100}^{12}},$$

$$U_2 = \frac{J_{5,100}^3 J_{10,100}^3 J_{15,100}^3 J_{20,100}^3 J_{30,100}^3 J_{35,100}^2 J_{40,100}^2 J_{45,100} J_{50,100}}{J_{100}^{12}}.$$

We have

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1+q^{10n}} = \frac{1}{2} U_1 - q^3 U_2 + \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \cdot \frac{q^{24}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+40}}. \quad (3.10)$$

*Proof.* First, we split the sum of the left side of (3.10) as five sums:

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1+q^{10n}} &= \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+10n}}{1+q^{50n}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+20n+3}}{1+q^{50n+10}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+30n+8}}{1+q^{50n+20}} \\ &\quad - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+40n+15}}{1+q^{50n+30}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+24}}{1+q^{50n+40}}. \end{aligned} \quad (3.11)$$

Let  $P_0, P_1, P_2, P_3$  and  $P_4$  denote the summations on the right hand side of (3.11), respectively. Substituting  $q, z^2$  and  $\zeta^2$  by  $q^{25}, -q^{40}$  and  $q^{40}$ , respectively, (2.9) becomes

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+10n}}{1+q^{50n}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+90n+80}}{1+q^{50n+80}}$$

$$\begin{aligned}
&= -\frac{2q(q^{20}, q^{30}; q^{50})_{\infty}(-q^{25}, q^{25})_{\infty}^2}{(-q^{10}, -q^{15}; q^{25})_{\infty}(q^{10}, q^{40}; q^{50})_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+24}}{1+q^{50n+40}} \\
&\quad + \frac{(q^{10}, q^{15}; q^{25})_{\infty}(q^{10}; q^{10})_{\infty}(q^{50}; q^{50})_{\infty}}{2(-q^{10}, -q^{15}; q^{25})_{\infty}(-q^{10}; q^{10})_{\infty}(-q^{50}; q^{50})_{\infty}}. \tag{3.12}
\end{aligned}$$

Replacing  $n$  by  $n - 1$  in second summation on the left side of (3.12), we get

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+10n}}{1+q^{50n}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+40n+15}}{1+q^{50n+30}} \\
&= -\frac{2qJ_{15,50}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+24}}{1+q^{50n+40}} + \frac{1}{2}U_1,
\end{aligned}$$

which can be expressed as

$$P_0 - P_3 = -\frac{2qJ_{15,50}}{J_{25,50}}P_4 + \frac{1}{2}U_1. \tag{3.13}$$

Similarly, substituting  $q$ ,  $z^2$  and  $\zeta^2$  by  $q^{25}$ ,  $-q^{40}$  and  $q^{30}$ , respectively, (2.9) takes the form

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+20n}}{1+q^{50n+10}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+80n+60}}{1+q^{50n+70}} \\
&= -\frac{2q^{25}(q^{10}, q^{40}; q^{50})_{\infty}(-q^{25}; q^{25})_{\infty}^2}{(q^{20}, q^{30}; q^{50})_{\infty}(-q^5, -q^{20}; q^{25})_{\infty}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+40}} \\
&\quad + \frac{(q^{10}, q^{15}; q^{25})_{\infty}(q^{10}; q^{10})_{\infty}(q^{100}; q^{100})_{\infty}}{(-q^5, -q^{20}; q^{25})_{\infty}(-q^{10}; q^{10})_{\infty}(-q^{10}, -q^{40}; q^{50})_{\infty}}.
\end{aligned}$$

Replacing  $n$  by  $n - 1$  in second summation on the left side of the above relation, we find that

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+20n}}{1+q^{50n+10}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+30n+5}}{1+q^{50n+20}} \\
&= -\frac{2qJ_{5,50}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+24}}{1+q^{50n+40}} + U_2,
\end{aligned}$$

that is,

$$P_2 - P_1 = \frac{2q^4J_{5,50}}{J_{25,50}}P_4 - q^3U_2. \tag{3.14}$$

Plugging (3.13) and (3.14) into (3.11), we obtain that

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1+q^{10n}} = \left(1 - \frac{2qJ_{15,50}}{J_{25,50}} + \frac{2q^4J_{5,50}}{J_{25,50}}\right) P_4 + \frac{1}{2}U_1 - q^3U_2. \tag{3.15}$$

Employing the following 5-dissection formula [19, Lemma 3.1]

$$\begin{aligned}\frac{(q; q)_\infty}{(-q; q)_\infty} &= \frac{(q^{25}; q^{25})_\infty}{(-q^{25}; q^{25})_\infty} - 2q(q^{15}, q^{35}, q^{50}; q^{50})_\infty + 2q^4(q^5, q^{45}, q^{50}; q^{50})_\infty \\ &= J_{25,50} - 2qJ_{15,50} + 2q^4J_{5,50}\end{aligned}\quad (3.16)$$

and using (3.15), we infer that

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+2n}}{1+q^{10n}} = \frac{(q; q)_\infty}{(-q; q)_\infty} \cdot \frac{P_4}{J_{25,50}} + \frac{1}{2}U_1 - q^3U_2.$$

This completes the proof. ■

**Lemma 3.4.** *Let*

$$\begin{aligned}V_1 &= \frac{J_{5,100}^2 J_{10,100} J_{20,100}^3 J_{30,100}^4 J_{45,100}^2 J_{50,100}^2}{J_{100}^{12}}, \\ V_2 &= \frac{J_{5,100} J_{10,100}^3 J_{15,100} J_{20,100}^2 J_{30,100}^3 J_{35,100} J_{40,100} J_{45,100} J_{50,100}}{J_{100}^{12}}.\end{aligned}$$

We have

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+4n}}{1+q^{10n}} = \frac{1}{2}V_1 - q^7V_2 - \frac{(q; q)_\infty}{(-q; q)_\infty} \cdot \frac{q^{21}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+30}}. \quad (3.17)$$

*Proof.* To prove (3.17), we rewrite the left side of (3.17) as follows:

$$\begin{aligned}\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+4n}}{1+q^{10n}} &= \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+20n}}{1+q^{50n}} - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+30n+5}}{1+q^{50n+10}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+40n+12}}{1+q^{50n+20}} \\ &\quad - \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+21}}{1+q^{50n+30}} + \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+60n+32}}{1+q^{50n+40}}.\end{aligned}\quad (3.18)$$

Let  $S_0, S_1, S_2, S_3$  and  $S_4$  denote the summations on the right hand side of (3.18), respectively. We claim that the following two relations hold:

$$S_0 - S_1 = -\frac{2q^4 J_{5,50}}{J_{25,50}} S_3 + \frac{1}{2}V_1, \quad (3.19)$$

$$S_2 + S_4 = \frac{2q J_{15,50}}{J_{25,50}} S_3 - q^7 V_2. \quad (3.20)$$

These two relations turn out to be specializations of the following identity due to Chan [5, Theorem 2.1]:

$$\frac{(a_1, q/a_1, \dots, a_r, q/a_r; q)_\infty (q; q)_\infty^2}{(b_1, q/b_1, \dots, b_s, q/b_s; q)_\infty}$$

$$\begin{aligned}
&= \frac{(a_1/b_1, qb_1/a_1, \dots, a_r/b_1, qb_1/a_r; q)_\infty}{(b_2/b_1, qb_1/b_2, \dots, b_s/b_1, qb_1/b_s; q)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^{(s-r)n} q^{(s-r)n(n+1)/2}}{1 - b_1 q^n} \left( \frac{a_1 \cdots a_r b_1^{s-r-1}}{b_2 \cdots b_s} \right)^n \\
&\quad + \text{idem}(b_1; b_2, \dots, b_s), \tag{3.21}
\end{aligned}$$

where  $\text{idem}(b_1; b_2, \dots, b_s)$  is defined by the relation

$$\begin{aligned}
&F(b_1, b_2, \dots, b_m) + \text{idem}(b_1; b_2, \dots, b_m) \\
&= F(b_1, b_2, \dots, b_m) + F(b_2, b_1, b_3, \dots, b_m) + \cdots + F(b_m, b_2, \dots, b_{m-1}, b_1). \tag{3.22}
\end{aligned}$$

To derive (3.19) from (3.21), we set  $r = 2, s = 3$  and replace  $q, a_1, a_2, b_1, b_2, b_3$  by  $q^{50}, q^{30}, q^5, -1, -q^{10}, -q^{30}$ , respectively, to get

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty} (-1)^n q^{25n^2} \left( \frac{q^{20n}}{1 + q^{50n}} - \frac{q^{30n+5}}{1 + q^{50n+10}} \right) \\
&= -\frac{q^4 (-1, q^{10}, -q^{25}, -q^{25}, q^{40}, -q^{50}; q^{50})_\infty}{(-q^5, q^{20}, -q^{20}, q^{30}, -q^{30}, -q^{45}; q^{50})_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+21}}{1 + q^{25n+30}} \\
&\quad + \frac{(q^5, q^{10}, q^{40}, q^{45}; q^{50})_\infty (q^{20}, q^{30}, q^{50}; q^{50})_\infty^2}{(-1, -q^5, -q^{10}, -q^{40}, -q^{45}, -q^{50}; q^{50})_\infty (-q^{20}, -q^{30}; q^{50})_\infty^2} \\
&= -\frac{2q^4 J_{5,50}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+21}}{1 + q^{25n+30}} + V_1,
\end{aligned}$$

which coincides with (3.19).

The identity (3.20) can be justified by using the identity (3.21), setting  $r = 2, s = 3$  and replacing  $q, a_1, a_2, b_1, b_2, b_3$  by  $q^{50}, q^{30}, q^{55}, -q^{20}, -q^{40}, -q^{30}$ , respectively, we find that

$$\begin{aligned}
&\sum_{n=-\infty}^{\infty} (-1)^n q^{25n^2} \left( \frac{q^{40n}}{1 + q^{50n+20}} + \frac{q^{60n+20}}{1 + q^{50n+40}} \right) \\
&= \frac{2(q^{20}, q^{30}; q^{50})_\infty (-q^{25}; q^{25})_\infty^2}{(-q^{10}, -q^{15}, -q^{35}, -q^{40}; q^{50})_\infty (q^{10}, q^{40}; q^{50})_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+10}}{1 + q^{50n+30}} \\
&\quad - \frac{q^{-5} (q^5, q^{10}, q^{40}, q^{45}; q^{50})_\infty (q^{20}, q^{30}, q^{50}; q^{50})_\infty^2}{(-q^{15}, -q^{35}; q^{50})_\infty (-q^{10}, -q^{20}, -q^{30}, -q^{40}; q^{50})_\infty^2} \\
&= \frac{2q J_{15,50}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n+9}}{1 + q^{50n+30}} - q^{-5} V_2,
\end{aligned}$$

which gives (3.20).

Now, we restate (3.17) as follows:

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+4n}}{1 + q^{10n}} = -\frac{(q; q)_\infty}{(-q; q)_\infty} \cdot \frac{q^{21}}{J_{25,50}} S_3 + \frac{1}{2} V_1 - q^7 V_2. \tag{3.23}$$



Substituting (3.19) and (3.20) into (3.18), we deduce that

$$\sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{n^2+4n}}{1+q^{10n}} = - \left( 1 - \frac{2qJ_{15,50}}{J_{25,50}} + \frac{2q^4J_{5,50}}{J_{25,50}} \right) S_3 + \frac{1}{2}V_1 - q^7V_2.$$

In view of the 5-dissection formula (3.16), the above relation leads to (3.23). This completes the proof.  $\blacksquare$

The following two lemmas are needed in the proofs of Theorems 1.2 and 1.3.

**Lemma 3.5.** *Assume that  $U_1$  and  $U_2$  are defined as in Lemma 3.3. Then the following identity holds:*

$$\begin{aligned} & \frac{1}{2}U_1 - q^3U_2 \\ &= \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \cdot \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} \{A_0(q^5) + qA_1(q^5) + q^2A_2(q^5) + q^3A_3(q^5) + q^4A_4(q^5)\}, \end{aligned} \quad (3.24)$$

where  $A_0(q)$ ,  $A_1(q)$ ,  $A_2(q)$ ,  $A_3(q)$ ,  $A_4(q)$  are defined as in Theorem 1.2.

**Lemma 3.6.** *Recall that  $U_1$  and  $U_2$  are defined in Lemma 3.3 and  $V_1$  and  $V_2$  are defined as in Lemma 3.4. The following identity holds:*

$$\begin{aligned} & \frac{1}{2}V_1 - \frac{1}{2}U_1 + q^3U_2 - q^7V_2 \\ &= \frac{(q; q)_{\infty}}{(-q; q)_{\infty}} \cdot \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} \{B_0(q^5) + qB_1(q^5) + q^2B_2(q^5) + q^3B_3(q^5) + q^4B_4(q^5)\}, \end{aligned} \quad (3.25)$$

where  $B_0(q)$ ,  $B_1(q)$ ,  $B_2(q)$ ,  $B_3(q)$ ,  $B_4(q)$  are defined as in Theorem 1.3.

To present the proofs of Lemmas 3.5 and 3.6, we shall use techniques in the theory of modular forms. Recall that the Dedekind  $\eta$ -function is defined by

$$\eta(\tau) = q^{\frac{1}{24}}(q; q)_{\infty},$$

where  $\tau \in \mathcal{H} = \{\tau \in \mathcal{C} : \text{Im}\tau > 0\}$  and  $q = e^{2\pi i\tau}$ , and the generalized Dedekind  $\eta$ -function is defined by

$$\eta_{\delta,g}(\tau) = q^{P(g/\delta)\delta/2} \prod_{\substack{n > 0 \\ n \equiv g \pmod{\delta}}} (1 - q^n) \prod_{\substack{n > 0 \\ n \equiv -g \pmod{\delta}}} (1 - q^n), \quad (3.26)$$

where  $g, \delta \in \mathbb{Z}^+$ ,  $0 < g < \delta$ ,  $P(t) = \{t\}^2 - \{t\} + \frac{1}{6}$  is the second Bernoulli function, and  $\{t\} = t - [t]$  is the fractional parts of  $t$ . In particular,

$$\eta_{\delta,0}(\tau) = q^{\frac{\delta}{12}}(q^{\delta}; q^{\delta})_{\infty}^2$$

and

$$\eta_{\delta, \frac{\delta}{2}}(\tau) = q^{-\frac{\delta}{24}}(q^{\frac{\delta}{2}}; q^{\delta})_{\infty}^2.$$

Let  $N$  be a fixed positive integer. A generalized Dedekind  $\eta$ -quotient of level  $N$  has the form

$$f(\tau) = \prod_{\substack{\delta|N \\ 0 < g < \delta}} \eta_{\delta, g}^{r_{\delta, g}}(\tau), \quad (3.27)$$

where

$$r_{\delta, g} \in \begin{cases} \frac{1}{2}\mathbb{Z}, & \text{if } g = 0 \text{ or } g = \delta/2, \\ \mathbb{Z}, & \text{otherwise.} \end{cases}$$

Suppose that  $f$  is a modular function with respect to the congruence subgroup  $\Gamma$  of  $\Gamma_0(1)$ . For  $A \in \Gamma_0(1)$ , there exists a cusp given by  $\zeta = A^{-1}\infty$ . If

$$f(A^{-1}\tau) = \sum_{m=m_0}^{\infty} b_m q^{m/N}$$

and  $b_{m_0} \neq 0$ , then we say that  $m_0$  is the order of  $f$  at  $\zeta$  with respect to  $\Gamma$ , and it is denoted by  $ORD(f, \zeta, \Gamma)$ .

Robins [24] found sufficient conditions for a generalized  $\eta$ -quotient to be a modular function on  $\Gamma_1(N)$ .

**Theorem 3.7.** *Let  $f(\tau)$  be a generalized  $\eta$ -quotient defined as in (3.27). If*

$$\sum_{\substack{\delta|N \\ 0 \leq g \leq \delta}} \delta P_2\left(\frac{g}{\delta}\right) r_{\delta, g} \equiv 0 \pmod{2},$$

and

$$\sum_{\substack{\delta|N \\ 0 \leq g \leq \delta}} \frac{N}{\delta} P_2(0) r_{\delta, g} \equiv 0 \pmod{2},$$

then  $f(\tau)$  is a modular function on  $\Gamma_1(N)$ .

The following theorem of Garvan and Liang [10] can be used to prove generalized  $\eta$ -quotient identities.

**Theorem 3.8.** *Let  $f_1(\tau), f_2(\tau), \dots, f_n(\tau)$  be generalized  $\eta$ -quotients that are modular functions on  $\Gamma_1(N)$ . Let  $\mathcal{S}_N$  be a set of inequivalent cusps for  $\Gamma_1(N)$ . Define the constant*

$$B = \sum_{\substack{s \in \mathcal{S}_N \\ s \neq i\infty}} \min(\{ORD(f_j, s, \Gamma_1(N)) : 1 \leq j \leq n\} \cup \{0\}). \quad (3.28)$$

Let

$$g(\tau) = \alpha_1 f_1(\tau) + \alpha_2 f_2(\tau) + \cdots + \alpha_n f_n(\tau) + 1, \quad (3.29)$$

where each  $\alpha_j \in \mathbb{C}$ . Then

$$g(\tau) \equiv 0$$

if and only if

$$ORD(g(\tau), i\infty, \Gamma_1(N)) > -B. \quad (3.30)$$

We are now in a position to prove Lemmas 3.5 and 3.6 with the aid of Theorem 3.8.

*Proof of Lemma 3.5.* The relation (3.24) can be stated as follows by the definitions of  $A_0(q)$ ,  $A_1(q)$ ,  $A_2(q)$ ,  $A_3(q)$ ,  $A_4(q)$  as given in (1.8), (1.9), (1.10), (1.11) and (1.12):

$$\begin{aligned} & \frac{1}{2} \frac{J_{10,100}^4 J_{15,100}^2 J_{30,100} J_{35,100}^2 J_{40,100}^3 J_{50,100}^2}{J_{100}^{12}} - q^3 \frac{J_{5,100} J_{10,100}^3 J_{15,100} J_{20,100} J_{30,100}^3 J_{35,100} J_{40,100}^2 J_{45,100} J_{50,100}}{J_{100}^{12}} \\ &= \frac{(q; q)_\infty}{(-q; q)_\infty} \cdot \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} \left\{ \frac{J_{10}^4 J_{15,100}^2 J_{25,100}^8 J_{35,100}^2}{2 J_{20,100}^4 J_{30,100}^3 J_{40,100} J_{100}^6} + \frac{2q^5 J_5^3 J_{50} J_{25,100} J_{100}^5}{J_{5,100}^2 J_{10,100} J_{40,100}^3 J_{45,100}^2} \right. \\ & \quad + \frac{8q^{10} J_5^2 J_{10}^4 J_{10,100} J_{15,100} J_{35,100}}{J_{20,100}^2 J_{30,100} J_{40,100} J_{50,100}^2 J_{100}} - \frac{16q^{10} J_5^2 J_{10,100} J_{15,100}^2 J_{30,100} J_{35,100}^2}{J_{20,100} J_{25,100}^2 J_{100}^3} \\ & \quad + q \left( \frac{J_{10}^4 J_{15,100}^3 J_{25,100}^6 J_{35,100}^3}{J_{20,100}^4 J_{30,100}^3 J_{40,100} J_{100}^6} + \frac{8q^5 J_5^2 J_{50}^2 J_{10,100} J_{15,100}^2 J_{35,100}^2 J_{40,100}}{J_{5,100} J_{20,100}^2 J_{30,100} J_{45,100} J_{100}^4} - \frac{8q^5 J_5^2 J_{10,100}^4 J_{30,100}^2 J_{40,100}^3}{J_{5,100}^2 J_{15,100} J_{35,100} J_{45,100}^2 J_{100}^3} \right. \\ & \quad \left. + \frac{4q^{10} J_5 J_{10}^2 J_{25,100}^2 J_{35,100}^3 J_{45,100}^2}{J_{20,100}^2 J_{40,100} J_{100}^4 J_{50}} - \frac{8q^{15} J_5^2 J_{5,100}^2 J_{20,100} J_{30,100} J_{45,100}^2 J_{50,100}}{J_{25,100}^2 J_{40,100}^2 J_{100}^3} \right) \\ & \quad + q^2 \left( \frac{2J_{10}^4 J_{15,100}^4 J_{25,100}^4 J_{35,100}^4}{J_{20,100}^4 J_{30,100}^3 J_{40,100} J_{100}^6} + \frac{2q^5 J_5^2 J_{10,100} J_{25,100}^4 J_{30,100}}{J_{15,100} J_{20,100} J_{35,100} J_{100}^3} - \frac{8q^5 J_5^2 J_{50}^2 J_{15,100}^3 J_{20,100} J_{30,100} J_{35,100}^3}{J_{5,100} J_{25,100}^2 J_{40,100}^2 J_{45,100} J_{100}^4} \right. \\ & \quad \left. + \frac{4q^{10} J_5^2 J_{50}^2 J_{5,100} J_{20,100} J_{30,100} J_{45,100}}{J_{40,100}^2 J_{100}^4} + \frac{8q^{10} J_5^2 J_{5,100} J_{10,100} J_{30,100} J_{45,100}}{J_{20,100} J_{100}^3} \right) \\ & \quad + q^3 \left( -\frac{J_5 J_{10}^2 J_{25,100}^7}{J_{50} J_{20,100}^2 J_{40,100} J_{100}^4} + \frac{4J_{10}^4 J_{15,100}^5 J_{25,100}^2 J_{35,100}^5}{J_{20,100}^4 J_{30,100}^3 J_{40,100} J_{100}^6} + \frac{2q^5 J_5^2 J_{50}^2 J_{10,100}^2 J_{25,100}^2 J_{40,100}}{J_{20,100}^2 J_{30,100} J_{100}^4} \right. \\ & \quad \left. - \frac{12q^5 J_5^2 J_{10,100} J_{25,100}^2 J_{30,100}}{J_{20,100} J_{100}^3} - \frac{16q^{10} J_5^3 J_{100}^6}{J_{20,100}^2 J_{25,100}^3 J_{40,100} J_{50}} + \frac{8q^{10} J_5^3 J_{50} J_{10,100} J_{100}^5}{J_{20,100}^3 J_{25,100}^3 J_{30,100}^2} \right) \\ & \quad + q^4 \left( -\frac{2J_5^2 J_{10,100} J_{25,100}^4 J_{30,100}}{J_{5,100} J_{20,100} J_{45,100} J_{100}^3} - \frac{J_5^2 J_{50}^2 J_{10,100}^2 J_{25,100}^4 J_{40,100}}{J_{5,100} J_{20,100}^2 J_{30,100} J_{45,100} J_{100}^4} + \frac{8J_{10}^4 J_{15,100}^6 J_{35,100}^6}{J_{20,100}^4 J_{30,100}^3 J_{40,100} J_{100}^6} \right. \\ & \quad \left. - \frac{16q^5 J_5^2 J_{10,100} J_{15,100} J_{30,100} J_{35,100}}{J_{20,100} J_{100}^3} - \frac{8q^5 J_5^3 J_{50} J_{100}^5 J_{10,100}}{J_{5,100} J_{20,100}^3 J_{25,100} J_{30,100}^2 J_{45,100}} \right. \\ & \quad \left. - \frac{16q^{15} J_5 J_{10}^2 J_{5,100}^4 J_{45,100}^4}{J_{50} J_{100}^4 J_{20,100}^2 J_{25,100} J_{40,100}} \right) \left. \right\}. \quad (3.31) \end{aligned}$$

Multiplying both sides by  $2J_{10,100}^{-4} J_{15,100}^{-2} J_{30,100}^{-1} J_{35,100}^{-2} J_{40,100}^{-3} J_{50,100}^{-2} J_{100}^{12}$ , the above relation becomes

$$\begin{aligned}
& 1 - \frac{2\eta_{100,5}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} = \frac{\eta_{1,0}(\tau)\eta_{50,0}(\tau)^{\frac{5}{2}}\eta_{50,25}(\tau)}{\eta_{2,0}(\tau)^{\frac{1}{2}}\eta_{10,0}(\tau)^3\eta_{10,5}(\tau)^6} \left( \eta_{100,25}(\tau)^8 \right. \\
& + \frac{4\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,20}(\tau)^3\eta_{100,25}(\tau)^3\eta_{100,30}(\tau)^2}{\eta_{100,10}(\tau)^2\eta_{100,40}(\tau)^3} + \frac{16\eta_{10,0}(\tau)\eta_{10,5}(\tau)\eta_{100,10}(\tau)\eta_{100,20}(\tau)^2\eta_{100,30}(\tau)^2}{\eta_{100,0}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,50}(\tau)^2} \\
& - \frac{32\eta_{10,5}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)^2\eta_{100,40}(\tau)\eta_{100,50}(\tau)} + 2\eta_{100,15}(\tau)\eta_{100,25}(\tau)^6\eta_{100,35}(\tau) + \frac{16\eta_{10,5}(\tau)}{\eta_{100,5}(\tau)\eta_{100,45}(\tau)} \\
& - \frac{16\eta_{10,0}(\tau)\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)}{\eta_{100,0}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,50}(\tau)^2} + \frac{8\eta_{100,5}(\tau)^3\eta_{100,20}(\tau)\eta_{100,25}(\tau)^4\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^3}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& - \frac{16\eta_{100,5}(\tau)^4\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^4}{\eta_{100,10}(\tau)^2\eta_{100,40}(\tau)^3} + 4\eta_{100,15}(\tau)^2\eta_{100,25}(\tau)^4\eta_{100,35}(\tau)^2 \\
& + \frac{4\eta_{100,5}(\tau)^2\eta_{100,20}(\tau)\eta_{100,25}(\tau)^6\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} - \frac{16\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,15}(\tau)^2\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^2\eta_{100,35}(\tau)^2}{\eta_{100,10}(\tau)^2\eta_{100,25}(\tau)\eta_{100,40}(\tau)^3} \\
& + \frac{8\eta_{100,5}(\tau)^3\eta_{100,20}(\tau)^3\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^3}{\eta_{100,10}(\tau)^2\eta_{100,40}(\tau)^3} + \frac{16\eta_{100,5}(\tau)^3\eta_{100,20}(\tau)\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^3}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& - \frac{2\eta_{100,5}(\tau)\eta_{100,20}(\tau)\eta_{100,25}(\tau)^8\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} + 8\eta_{100,15}(\tau)^3\eta_{100,25}(\tau)^2\eta_{100,35}(\tau)^3 \\
& + 4\eta_{100,5}(\tau)^2\eta_{100,25}(\tau)^4\eta_{100,45}(\tau)^2 - \frac{24\eta_{100,5}(\tau)^2\eta_{100,20}(\tau)\eta_{100,25}(\tau)^4\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& - \frac{32\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,5}(\tau)^2\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& + \frac{16\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,5}(\tau)^2\eta_{100,45}(\tau)^2}{\eta_{100,25}(\tau)} - \frac{4\eta_{100,5}(\tau)\eta_{100,20}(\tau)\eta_{100,25}(\tau)^6\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& - 2\eta_{100,5}(\tau)\eta_{100,25}(\tau)^6\eta_{100,45}(\tau) + 16\eta_{100,15}(\tau)^4\eta_{100,35}(\tau)^4 \\
& - \frac{32\eta_{10,5}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} - \frac{16\eta_{10,5}(\tau)}{\eta_{100,15}(\tau)\eta_{100,35}(\tau)} \\
& \left. - \frac{32\eta_{100,5}(\tau)^5\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^5}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \right).
\end{aligned}$$

By means of Theorem 3.7, it can be shown that each term of the above identity is a modular function with respect to  $\Gamma_1(100)$ . Applying the algorithm of Garvan and Liang [10], we find that the constant  $B$  in (3.28) is equal to  $-600$ . Thus, by Theorem 3.8, (3.31) holds as long as one can verify it holds for the coefficients of  $q^n$  for  $1 \leq n \leq 600$ . This task can be easily accomplished, and hence the proof is complete.  $\blacksquare$

*Proof of Lemma 3.6.* By the definitions of  $B_0(q)$ ,  $B_1(q)$ ,  $B_2(q)$ ,  $B_3(q)$ ,  $B_4(q)$  as given in

(1.16), (1.17), (1.18), (1.19) and (1.20), we see that (3.25) can be expressed as

$$\begin{aligned}
& -\frac{1}{2} \frac{J_{10,100}^4 J_{15,100}^2 J_{30,100} J_{35,100}^2 J_{40,100}^3 J_{50,100}^2}{J_{100}^{12}} + q^3 \frac{J_{5,100} J_{10,100}^3 J_{15,100} J_{20,100} J_{30,100}^3 J_{35,100} J_{40,100}^2 J_{45,100} J_{50,100}}{J_{100}^{12}} \\
& + \frac{1}{2} \frac{J_{5,100}^2 J_{10,100} J_{20,100}^3 J_{30,100}^4 J_{45,100}^2 J_{50,100}^2}{J_{100}^{12}} - q^7 \frac{J_{5,100} J_{10,100}^3 J_{15,100} J_{20,100}^2 J_{30,100}^3 J_{35,100} J_{40,100} J_{45,100} J_{50,100}}{J_{100}^{12}} \\
& = \frac{(q; q)_\infty}{(-q; q)_\infty} \cdot \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} \left\{ -\frac{q^5 J_5^5 J_{5,100} J_{25,100} J_{45,100}}{J_{50} J_{100} J_{20,100}^2 J_{30,100} J_{40,100}} + \frac{4q^5 J_5^4 J_{100}^2 J_{15,100} J_{35,100}}{J_{5,100} J_{20,100}^2 J_{40,100} J_{45,100} J_{50,100}} \right. \\
& - \frac{16q^{10} J_{10}^7 J_{5,10}^{\frac{1}{2}} J_{5,100} J_{10,100} J_{45,100}}{J_{100} J_{20,100} J_{25,100}^3} + \frac{16q^{10} J_5^2 J_{10,100} J_{15,100}^2 J_{30,100} J_{35,100}^2}{J_{100}^3 J_{20,100} J_{25,100}^2} - \frac{8q^{15} J_5 J_{10}^2 J_{5,100}^2 J_{25,100}^3 J_{45,100}^2}{J_{50} J_{100}^4 J_{20,100} J_{40,100}^2} \\
& - \frac{4q^{15} J_5^4 J_{100}^2 J_{5,100} J_{45,100}}{J_{15,100} J_{20,100} J_{35,100} J_{40,100}^2 J_{50,100}} - \frac{16q^{20} J_{5,100}^2 J_{10,100} J_{30,100} J_{45,100}^2 J_5^2}{J_{100}^3 J_{25,100}^2 J_{40,100}} \\
& + q \left( -\frac{2q^5 J_5^2 J_{50}^2 J_{10,100} J_{25,100}^2}{J_{100}^4 J_{20,100}} + \frac{8q^5 J_5^3 J_{10,100} J_{15,100}^3 J_{30,100} J_{35,100}^3}{J_{50} J_{20,100} J_{25,100} J_{100}^6} + \frac{4q^{10} J_{10}^4 J_{5,100}^2 J_{10,100} J_{30,100} J_{45,100}^2}{J_{100}^6 J_{20,100} J_{50,100}} \right. \\
& + \frac{2q^{10} J_{10}^4 J_{5,100} J_{10,100} J_{25,100}^2 J_{30,100} J_{45,100}}{J_{100}^6 J_{40,100} J_{50,100}} - \frac{16q^{10} J_5^3 J_{50} J_{100}^5}{J_{20,100}^2 J_{25,100}^3 J_{30,100} J_{40,100}} + \frac{12q^{15} J_5^2 J_{5,100}^2 J_{10,100} J_{30,100} J_{45,100}^2}{J_{100}^3 J_{20,100} J_{25,100}^2} \\
& \left. + \frac{14q^{15} J_5^2 J_{5,100} J_{10,100} J_{30,100} J_{45,100}}{J_{100}^3 J_{40,100}} - \frac{16q^{20} J_5^2 J_{50}^2 J_{5,100}^4 J_{10,100} J_{45,100}^4}{J_{100}^4 J_{15,100} J_{20,100} J_{35,100} J_{45,100}^4} \right) \\
& + q^2 \left( \frac{q^5 J_5^3 J_{10,100} J_{25,100}^5 J_{30,100}}{J_{50} J_{100}^6 J_{40,100}} + \frac{4q^5 J_5^2 J_{50}^2 J_{5,100}^2 J_{20,100}^2 J_{30,100}^3 J_{45,100}^2}{J_{100}^4 J_{10,100}^2 J_{15,100} J_{35,100} J_{40,100}^3} \right. \\
& - \frac{8q^5 J_{10}^4 J_{5,100} J_{15,100}^5 J_{35,100} J_{45,100}}{J_{100}^6 J_{20,100}^4 J_{30,100}^3 J_{40,100}} + \frac{12q^5 J_5 J_{10}^2 J_{15,100}^4 J_{35,100}^4}{J_{50} J_{100}^4 J_{25,100}^2 J_{20,100} J_{40,100}} + \frac{4q^{10} J_5 J_{50} J_{10,100}^6 J_{30,100}^2 J_{40,100}^5}{J_{25,100} J_{100}^{12}} \\
& - \frac{12q^{10} J_5^2 J_{10,100} J_{25,100}^2 J_{30,100}}{J_{100}^3 J_{40,100}} - \frac{16q^{10} J_5^2 J_{5,100} J_{10,100} J_{30,100} J_{45,100}}{J_{100}^3 J_{20,100}} - \frac{8q^{15} J_5 J_{10}^2 J_{50} J_{10,100} J_{30,100}}{J_{100}^5 J_{20,100} J_{25,100}} \left. \right) \\
& + q^3 \left( \frac{J_5^3 J_{10,100} J_{25,100}^5 J_{30,100}}{J_{50} J_{100}^6 J_{20,100}} - \frac{8q^5 J_5^2 J_{50}^2 J_{10,100}^3 J_{15,100}^2 J_{35,100}^2 J_{40,100}^2}{J_{100}^4 J_{5,100} J_{20,100}^3 J_{30,100}^2 J_{45,100}} + \frac{12q^5 J_5^2 J_{10,100} J_{25,100}^2 J_{30,100}}{J_{100}^3 J_{20,100}} \right. \\
& + \frac{4q^{10} J_{10}^4 J_{5,100}^6 J_{25,100}^2 J_{30,100} J_{45,100}^6}{J_{100}^6 J_{10,100}^4 J_{15,100} J_{35,100} J_{40,100}^5} - \frac{16q^{15} J_5^3 J_{100}^6 J_{15,100} J_{35,100}}{J_{50} J_{5,100} J_{25,100}^3 J_{40,100}^3 J_{45,100}} - \frac{16q^{20} J_5 J_{10}^2 J_{5,100}^4 J_{45,100}^4}{J_{50} J_{100}^4 J_{50,100} J_{25,100} J_{40,100}^2} \left. \right) \\
& + q^4 \left( \frac{2J_5^2 J_{10,100} J_{25,100}^4 J_{30,100}}{J_{100}^3 J_{5,100} J_{20,100} J_{45,100}} - \frac{2q^5 J_5^2 J_{50}^2 J_{10,100}^5 J_{25,100}^2 J_{30,100}^2 J_{40,100}}{J_{100}^4 J_{5,100}^3 J_{15,100} J_{35,100} J_{45,100}^3} + \frac{16q^{10} J_5^2 J_{50}^2 J_{5,100}^2 J_{10,100} J_{45,100}^2}{J_{100}^4 J_{20,100} J_{25,100}^2} \right. \\
& - \frac{16q^{10} J_5^2 J_{10,100} J_{15,100}^2 J_{30,100} J_{35,100}^2}{J_{100}^3 J_{25,100}^2 J_{40,100}} + \frac{8q^{15} J_5^2 J_{5,100}^2 J_{10,100} J_{30,100} J_{45,100}^2}{J_{100}^3 J_{15,100} J_{35,100} J_{40,100}} + \frac{16q^{15} J_5 J_{10}^2 J_{5,100}^4 J_{45,100}^4}{J_{50} J_{100}^4 J_{20,100}^2 J_{25,100} J_{40,100}} \\
& \left. + \frac{32q^{20} J_5^2 J_{10,100}^3 J_{15,100}^2 J_{35,100}^2}{J_{50}^2 J_{100}^2 J_{20,100} J_{25,100}^2} \right\}. \tag{3.32}
\end{aligned}$$

Multiplying both sides of above relation by  $2J_{10,100}^{-4} J_{15,100}^{-2} J_{30,100}^{-1} J_{35,100}^{-2} J_{40,100}^{-3} J_{50,100}^{-2} J_{100}^{12}$ , the

above relation takes the form

$$\begin{aligned}
& -1 + \frac{2\eta_{100,5}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& + \frac{\eta_{100,5}(\tau)^2\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^3\eta_{100,45}(\tau)^2}{\eta_{100,10}(\tau)^3\eta_{100,15}(\tau)^2\eta_{100,35}(\tau)^2\eta_{100,40}(\tau)^3} - \frac{2\eta_{100,5}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)\eta_{100,45}(\tau)}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& = \frac{\eta_{1,0}(\tau)\eta_{50,0}(\tau)^{\frac{5}{2}}\eta_{50,25}(\tau)}{\eta_{2,0}(\tau)^{\frac{1}{2}}\eta_{10,0}(\tau)^3\eta_{10,5}(\tau)^6} \left( -\frac{2\eta_{10,5}(\tau)^3\eta_{100,10}(\tau)\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^3\eta_{100,40}(\tau)}{\eta_{100,15}(\tau)^3\eta_{100,35}(\tau)^3} \right. \\
& + \frac{8\eta_{10,5}(\tau)^{\frac{3}{2}}\eta_{100,20}(\tau)^2\eta_{100,25}(\tau)\eta_{100,30}(\tau)^3}{\eta_{100,50}(\tau)} - \frac{32\eta_{10,5}(\tau)\eta_{100,10}(\tau)\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^3\eta_{100,40}(\tau)}{\eta_{100,15}(\tau)^3\eta_{100,25}(\tau)^4\eta_{100,35}(\tau)^3} \\
& + \frac{32\eta_{10,5}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)^2\eta_{100,40}(\tau)\eta_{100,50}(\tau)} - \frac{8\eta_{10,5}(\tau)^2\eta_{100,5}(\tau)\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^3\eta_{100,45}(\tau)}{\eta_{100,15}(\tau)^3\eta_{100,35}(\tau)^3\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& - \frac{16\eta_{100,5}(\tau)^3\eta_{100,20}(\tau)^2\eta_{100,25}(\tau)^4\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^3}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)^2\eta_{100,50}(\tau)} - \frac{32\eta_{10,0}(\tau)\eta_{100,5}(\tau)^4\eta_{100,45}(\tau)^4}{\eta_{100,0}(\tau)\eta_{100,10}(\tau)^3\eta_{100,40}(\tau)^4\eta_{100,50}(\tau)^2} \\
& - \frac{4\eta_{10,5}(\tau)\eta_{100,20}(\tau)\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)^2\eta_{100,35}(\tau)^2\eta_{100,40}(\tau)} + \frac{16\eta_{10,5}(\tau)^2\eta_{100,20}(\tau)^2\eta_{100,30}(\tau)^3}{\eta_{100,5}(\tau)\eta_{100,25}(\tau)^2\eta_{100,45}(\tau)\eta_{100,50}(\tau)} \\
& + \frac{4\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,10}(\tau)\eta_{100,20}(\tau)^4\eta_{100,25}(\tau)\eta_{100,30}(\tau)^4}{\eta_{100,15}(\tau)^3\eta_{100,35}(\tau)^3\eta_{100,50}(\tau)} + \frac{8\eta_{10,5}(\tau)\eta_{100,10}(\tau)\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^4\eta_{100,40}(\tau)}{\eta_{100,15}(\tau)^4\eta_{100,25}(\tau)^2\eta_{100,35}(\tau)^4\eta_{100,50}(\tau)} \\
& - \frac{32\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,5}(\tau)^2\eta_{100,20}(\tau)\eta_{100,30}(\tau)\eta_{100,45}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)\eta_{100,40}(\tau)} + \frac{24\eta_{100,5}(\tau)^4\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^4}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& + \frac{28\eta_{100,5}(\tau)^3\eta_{100,20}(\tau)^2\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^3}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)^2\eta_{100,50}(\tau)} - \frac{32\eta_{100,5}(\tau)^6\eta_{100,20}(\tau)\eta_{100,30}(\tau)\eta_{100,45}(\tau)^6}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)^2\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)} \\
& + \frac{2\eta_{10,5}(\tau)^{\frac{3}{2}}\eta_{100,20}(\tau)^3\eta_{100,25}(\tau)^5\eta_{100,30}(\tau)^3}{\eta_{100,15}(\tau)^2\eta_{100,35}(\tau)^2\eta_{100,40}(\tau)\eta_{100,50}(\tau)} + \frac{8\eta_{100,5}(\tau)^4\eta_{100,20}(\tau)^4\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)^4\eta_{100,45}(\tau)^4}{\eta_{100,10}(\tau)^4\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)^4} \\
& - \frac{16\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,15}(\tau)^2\eta_{100,35}(\tau)^2}{\eta_{100,25}(\tau)} + \frac{24\eta_{10,5}(\tau)^{\frac{1}{2}}\eta_{100,15}(\tau)^2\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2\eta_{100,35}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& + \frac{8\eta_{10,0}(\tau)\eta_{100,5}(\tau)\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,45}(\tau)}{\eta_{100,0}(\tau)\eta_{100,15}(\tau)\eta_{100,20}(\tau)\eta_{100,35}(\tau)\eta_{100,50}(\tau)^2} - \frac{24\eta_{100,5}(\tau)^2\eta_{100,20}(\tau)^2\eta_{100,25}(\tau)^4\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)^2\eta_{100,50}(\tau)} \\
& - \frac{32\eta_{100,5}(\tau)^3\eta_{100,20}(\tau)\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^3}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} - \frac{16\eta_{100,5}(\tau)\eta_{100,10}(\tau)^3\eta_{100,20}(\tau)^2\eta_{100,30}(\tau)^3\eta_{100,45}(\tau)}{\eta_{100,15}(\tau)\eta_{100,35}(\tau)} \\
& + \frac{2\eta_{10,5}(\tau)^{\frac{3}{2}}\eta_{100,20}(\tau)^2\eta_{100,25}(\tau)^5\eta_{100,30}(\tau)^3}{\eta_{100,15}(\tau)^2\eta_{100,35}(\tau)^2\eta_{100,50}(\tau)} - \frac{16\eta_{10,5}(\tau)\eta_{100,10}(\tau)\eta_{100,40}(\tau)}{\eta_{100,5}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)\eta_{100,45}(\tau)} \\
& + \frac{24\eta_{100,5}(\tau)^2\eta_{100,20}(\tau)\eta_{100,25}(\tau)^4\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} + \frac{8\eta_{100,5}(\tau)^6\eta_{100,20}(\tau)^4\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)^4\eta_{100,45}(\tau)^6}{\eta_{100,10}(\tau)^4\eta_{100,15}(\tau)^3\eta_{100,35}(\tau)^3\eta_{100,40}(\tau)^4} \\
& - \frac{32\eta_{10,5}(\tau)\eta_{100,20}(\tau)^3\eta_{100,30}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)^2\eta_{100,40}(\tau)^3\eta_{100,50}(\tau)} - \frac{32\eta_{100,5}(\tau)^5\eta_{100,20}(\tau)^2\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^5}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)^2\eta_{100,50}(\tau)} \\
& + \frac{4\eta_{100,5}(\tau)\eta_{100,20}(\tau)\eta_{100,25}(\tau)^6\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} - \frac{4\eta_{100,10}(\tau)^3\eta_{100,20}(\tau)^2\eta_{100,25}(\tau)^4\eta_{100,30}(\tau)^3}{\eta_{100,5}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,45}(\tau)}
\end{aligned}$$

$$\begin{aligned}
& + \frac{32\eta_{100,5}(\tau)^4\eta_{100,20}(\tau)\eta_{100,30}(\tau)\eta_{100,45}(\tau)^4}{\eta_{100,10}(\tau)\eta_{100,40}(\tau)} - \frac{32\eta_{10,5}(\tau)\eta_{100,20}(\tau)^2\eta_{100,30}(\tau)^2}{\eta_{100,10}(\tau)\eta_{100,25}(\tau)^2\eta_{100,40}(\tau)^2\eta_{100,50}(\tau)} \\
& + \frac{16\eta_{100,5}(\tau)^4\eta_{100,20}(\tau)^2\eta_{100,25}(\tau)^2\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^4}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)^2\eta_{100,50}(\tau)} + \frac{32\eta_{100,5}(\tau)^5\eta_{100,20}(\tau)\eta_{100,30}(\tau)^2\eta_{100,45}(\tau)^5}{\eta_{100,10}(\tau)\eta_{100,15}(\tau)\eta_{100,35}(\tau)\eta_{100,40}(\tau)\eta_{100,50}(\tau)} \\
& + \frac{64\eta_{10,5}(\tau)\eta_{100,10}(\tau)\eta_{100,20}(\tau)\eta_{100,30}(\tau)}{\eta_{100,25}(\tau)^2\eta_{100,40}(\tau)\eta_{100,50}(\tau)}.
\end{aligned}$$

In light of Theorem 3.7, it can be verified that each term of the above identity is a modular function with respect to  $\Gamma_1(100)$ . Employing the algorithm by Garvan and Liang [10], we deduce that the constant  $B$  in (3.28) is equal to  $-640$ . By Theorem 3.8, (3.32) can be justified if it holds for coefficients of  $q^n$  for  $1 \leq n \leq 640$ , which can be easily checked. This completes the proof.  $\blacksquare$

Theorems 1.2 and 1.3 are immediate consequences of Lemmas 3.5 and 3.6.

*Proof of Theorem 1.2.* The relation (3.5) in Lemma 3.2 states that

$$\begin{aligned}
& \sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + \overline{N}_2(1, 10, n) - \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n))q^n \\
& = \frac{2(-q; q)_{\infty}}{(q; q)_{\infty}} \left( \frac{1}{2}U_1 - q^2U_3 \right) + \frac{2q^{24}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1 + q^{50n+40}}. \tag{3.33}
\end{aligned}$$

Substituting (3.24) into (3.33), we obtain that

$$\begin{aligned}
& \sum_{n=0}^{\infty} (\overline{N}_2(0, 10, n) + \overline{N}_2(1, 10, n) - \overline{N}_2(4, 10, n) - \overline{N}_2(5, 10, n))q^n \\
& = \frac{2J_{50}^4 J_{25,50}}{J_{5,10}^6} (A_0(q^5) + qA_1(q^5) + q^2A_2(q^5) + q^3A_3(q^5)) \\
& \quad + 2q^4 \left( \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} A_4(q^5) + \frac{q^{20}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1 + q^{50n+40}} \right).
\end{aligned}$$

Comparing the coefficients  $q^{5n}$ ,  $q^{5n+1}$ ,  $q^{5n+2}$ ,  $q^{5n+3}$  and  $q^{5n+4}$  of above relation, we get (1.6) and (1.7). This completes the proof.  $\blacksquare$

*Proof of Theorem 1.3.* The relation (3.6) in Lemma 3.2 reads

$$\begin{aligned}
& \sum_{n=0}^{\infty} (\overline{N}_2(1, 10, n) + \overline{N}_2(2, 10, n) - \overline{N}_2(3, 10, n) - \overline{N}_2(4, 10, n))q^n \\
& = \frac{2(-q; q)_{\infty}}{(q; q)_{\infty}} \left( \frac{1}{2}V_1 - q^7V_2 \right) - \frac{2q^{21}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1 + q^{50n+30}}
\end{aligned}$$

$$\begin{aligned}
& - \left\{ \frac{2(-q; q)_\infty}{(q; q)_\infty} \left( \frac{1}{2}U_1 - q^3U_2 \right) + \frac{2q^{24}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+40}} \right\} \\
& = \frac{2(-q; q)_\infty}{(q; q)_\infty} \left( \frac{1}{2}V_1 - \frac{1}{2}U_1 + q^3U_2 - q^7V_2 \right) \\
& \quad - \frac{2q^{21}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+30}} - \frac{2q^{24}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+40}}. \tag{3.34}
\end{aligned}$$

Substituting (3.25) into (3.34), we obtain that

$$\begin{aligned}
& \sum_{n=0}^{\infty} (\bar{N}_2(1, 10, n) + \bar{N}_2(2, 10, n) - \bar{N}_2(3, 10, n) - \bar{N}_2(4, 10, n))q^n \\
& = \frac{2J_{50}^4 J_{25,50}}{J_{5,10}^6} B_0(q^5) + 2q \left( \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} B_1(q^5) - \frac{q^{20}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+30}} \right) \\
& \quad + 2q^2 \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} B_2(q^5) + 2q^3 \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} B_3(q^5) \\
& \quad + 2q^4 \left( \frac{J_{50}^4 J_{25,50}}{J_{5,10}^6} B_4(q^5) - \frac{q^{20}}{J_{25,50}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{25n^2+50n}}{1+q^{50n+40}} \right).
\end{aligned}$$

Comparing the coefficients  $q^{5n}$ ,  $q^{5n+1}$ ,  $q^{5n+2}$ ,  $q^{5n+3}$  and  $q^{5n+4}$  of above relation, we get (1.13), (1.14) and (1.15). This completes the proof.  $\blacksquare$

## 4 Proof of Theorem 1.4

Before presenting the proof of Theorem 1.4, we recall the following theorem due to Liaw [16].

**Theorem 4.1.** *If  $p$  and  $r$  are positive integers with  $p \geq 2$  and  $r < p$  and*

$$\sum_{n=0}^{\infty} b_{p,r}(n)q^n = \frac{(q^p; q^p)_\infty}{(q^r; q^p)_\infty (q^{p-r}; q^p)_\infty},$$

then  $b_{p,r}(n) \geq 0$  for all  $n$ .

We are now in a position to prove Theorem 1.4.

*Proof of Theorem 1.4.* We first prove (1.21), (1.22) and (1.25). From Theorem 1.1, we find that

$$\bar{R}_{0,2}(0, 6) = \frac{J_6^9 J_{2,4}^2}{J_{1,6}^6 J_2^3 J_{3,6}} = \frac{(q^6; q^6)_\infty^2}{(q, q^5; q^6)_\infty^2} \frac{(-q; q^2)_\infty}{(q^4; q^4)_\infty (q^3; q^6)_\infty (q, q^5; q^6)_\infty^3}. \tag{4.1}$$



By Theorem 4.1, we see that the coefficients of  $q^n$  in

$$\frac{(q^6; q^6)_\infty^2}{(q, q^5; q^6)_\infty^2}$$

is nonnegative for  $n \geq 0$ . This implies that

$$\overline{N}_2(0, 6, 3n) + \overline{N}_2(1, 6, 3n) \geq \overline{N}_2(2, 6, 3n) + \overline{N}_2(3, 6, 3n). \quad (4.2)$$

Since  $i \equiv s \pmod{m}$  if and only if  $i \equiv s \pmod{2m}$  or  $i \equiv m + s \pmod{2m}$ , we find that

$$\overline{N}_2(s, m, n) = \overline{N}_2(s, 2m, n) + \overline{N}_2(m + s, 2m, n), \quad (4.3)$$

and thus

$$\overline{N}_2(0, 3, 3n) = \overline{N}_2(0, 6, 3n) + \overline{N}_2(3, 6, 3n). \quad (4.4)$$

Moreover, by (2.3)

$$\overline{N}_2(m + s, 2m, n) = \overline{N}_2(m - s, 2m, n),$$

(4.3) can be rewritten as

$$\overline{N}_2(s, m, n) = \overline{N}_2(s, 2m, n) + \overline{N}_2(m - s, 2m, n). \quad (4.5)$$

In particular, we get

$$\overline{N}_2(1, 3, 3n) = \overline{N}_2(1, 6, 3n) + \overline{N}_2(2, 6, 3n). \quad (4.6)$$

By means of the following identity due to Lovejoy and Osburn [20, Theorem 1.1],

$$\sum_{n=0}^{\infty} (\overline{N}_2(0, 3, 3n) - \overline{N}_2(1, 3, 3n))q^n = -1 + \frac{J_6^3 J_{3,6}}{J_{1,6}^2 J_2},$$

together with (4.4) and (4.6), we obtain

$$\sum_{n=0}^{\infty} (\overline{N}_2(0, 6, 3n) + \overline{N}_2(3, 6, 3n) - \overline{N}_2(1, 6, 3n) - \overline{N}_2(2, 6, 3n))q^n = -1 + \frac{J_6^3 J_{3,6}}{J_{1,6}^2 J_2}. \quad (4.7)$$

Consequently,

$$\frac{J_6^3 J_{3,6}}{J_{1,6}^2 J_2} = \frac{(q^3; q^6)_\infty^2 (q^6; q^6)_\infty}{(q, q^5; q^6)_\infty^2 (q^2, q^4; q^6)_\infty} = \frac{(q^3; q^3)_\infty}{(q, q^2; q^3)_\infty} \frac{(q^3; q^6)_\infty}{(q, q^5; q^6)_\infty} = \frac{(q^3; q^3)_\infty}{(q, q^2; q^3)_\infty} \frac{(q^3; q^6)_\infty^2}{(q; q^2)_\infty}$$

and

$$\frac{(q^3; q^6)_\infty^2}{(q; q^2)_\infty} = \frac{(-q; q)_\infty}{(-q^3; q^3)_\infty^2} = \frac{(-q; q^3)_\infty (-q^2; q^3)_\infty}{(-q^3; q^3)_\infty} = \frac{1}{(q^6; q^6)_\infty} \sum_{n=-\infty}^{\infty} q^{(3n^2-n)/2}.$$

Therefore, (4.7) can be written as

$$\begin{aligned} & \sum_{n=0}^{\infty} (\overline{N}_2(0, 6, 3n) + \overline{N}_2(3, 6, 3n) - \overline{N}_2(1, 6, 3n) - \overline{N}_2(2, 6, 3n)) q^n \\ &= -1 + \frac{(q^3; q^3)_{\infty}}{(q, q^2; q^3)_{\infty}} \frac{1}{(q^6; q^6)_{\infty}} \sum_{n=-\infty}^{\infty} q^{(3n^2-n)/2}. \end{aligned}$$

By Theorem 4.1, we deduce that for  $n \geq 1$

$$\overline{N}_2(0, 6, 3n) + \overline{N}_2(3, 6, 3n) \geq \overline{N}_2(1, 6, 3n) + \overline{N}_2(2, 6, 3n), \quad (4.8)$$

so (1.25) is verified. Adding (4.2) to (4.8), we reach (1.21).

Subtracting (4.7) from (4.1), we obtain that

$$\begin{aligned} \sum_{n=0}^{\infty} (\overline{N}_2(1, 6, 3n) - \overline{N}_2(3, 6, 3n)) q^n &= \frac{1}{2} \cdot \frac{J_6^9 J_{2,4}^2}{J_{1,6}^6 J_2^3 J_{3,6}} + \frac{1}{2} - \frac{1}{2} \cdot \frac{J_6^3 J_{3,6}}{J_{1,6}^2 J_2} \\ &= \frac{1}{2} + \frac{1}{2} \cdot \frac{J_6^3 J_{3,6}}{J_{1,6}^2 J_2} \left( \frac{J_6^6 J_{2,4}^2}{J_{1,6}^4 J_2^2 J_{3,6}^2} - 1 \right) \\ &= \frac{1}{2} + \frac{1}{2} \cdot \frac{J_6^3 J_{3,6}}{J_{1,6}^2 J_2} \cdot ((-q; q^2)_{\infty}^2 (-q; q)_{\infty}^2 - 1), \end{aligned}$$

which yields (1.22).

To verify (1.23), we rewrite (1.4) as

$$\overline{R}_{0,2}(1, 6) = \frac{2J_6^9 J_{2,4}^2}{J_{1,6}^5 J_2^3 J_{3,6}^2} = \frac{2(q^6; q^6)_{\infty}^2}{(q, q^5; q^6)_{\infty}^2} \frac{(-q; q^2)_{\infty}}{(q^4; q^4)_{\infty} (q, q^5; q^6)_{\infty}^2 (q^3; q^6)_{\infty}^3}. \quad (4.9)$$

By Theorem 4.1, we see that the coefficients of  $q^n$  in

$$\frac{(q^6; q^6)_{\infty}^2}{(q, q^5; q^6)_{\infty}^2}$$

is nonnegative for  $n \geq 0$ , which implies that

$$\overline{N}_2(0, 6, 3n+1) + \overline{N}_2(1, 6, 3n+1) \geq \overline{N}_2(2, 6, 3n+1) + \overline{N}_2(3, 6, 3n+1). \quad (4.10)$$

Using the following identity due to Lovejoy and Osburn [20, Theorem 1.1]:

$$\sum_{n=0}^{\infty} (\overline{N}_2(0, 3, 3n+1) - \overline{N}_2(1, 3, 3n+1)) q^n = \frac{2J_6^3}{J_{1,6} J_2} = \frac{(q^6; q^6)_{\infty}}{(q^2, q^4; q^6)_{\infty}} \frac{2}{(q, q^5; q^6)_{\infty}},$$

together with the relations (4.3) and (4.5), we get

$$\sum_{n=0}^{\infty} (\overline{N}_2(0, 6, 3n+1) + \overline{N}_2(3, 6, 3n+1) - \overline{N}_2(1, 6, 3n+1) - \overline{N}_2(2, 6, 3n+1)) q^n$$

$$= \frac{(q^6; q^6)_\infty}{(q^2, q^4; q^6)_\infty} \frac{2}{(q, q^5; q^6)_\infty}. \quad (4.11)$$

By Theorem 4.1, we see that the coefficients of  $q^n$  in

$$\frac{(q^6; q^6)_\infty}{(q^2, q^4; q^6)_\infty}$$

is nonnegative for  $n \geq 0$ . Hence

$$\overline{N}_2(0, 6, 3n+1) + \overline{N}_2(3, 6, 3n+1) \geq \overline{N}_2(1, 6, 3n+1) + \overline{N}_2(2, 6, 3n+1), \quad (4.12)$$

as given in (1.26). Adding (4.10) to (1.26), we arrive at (1.23). Subtracting (4.11) from (4.9), we obtain that

$$\begin{aligned} \sum_{n=0}^{\infty} (\overline{N}_2(1, 6, 3n+1) - \overline{N}_2(3, 6, 3n+1)) q^n &= \frac{J_6^9 J_{2,4}^2}{J_{1,6}^5 J_2^3 J_{3,6}^2} - \frac{J_6^3}{J_{1,6} J_2} \\ &= \frac{J_6^3}{J_{1,6} J_2} \left( \frac{J_6^6 J_{2,4}^2}{J_{1,6}^4 J_2^2 J_{3,6}^2} - 1 \right) \\ &= \frac{J_6^3}{J_{1,6} J_2} \cdot ((-q; q^2)_\infty^2 (-q; q)_\infty^2 - 1). \end{aligned} \quad (4.13)$$

It is easily seen that the coefficients of  $q^n$  in (4.13) are nonnegative for  $n \geq 1$ , and this justifies (1.24).

It remains to prove (1.27). Using the following relation due to Lovejoy and Osburn [20, Theorem 1.1],

$$\begin{aligned} &\sum_{n=0}^{\infty} (\overline{N}_2(0, 3, 3n+2) - \overline{N}_2(1, 3, 3n+2)) q^n \\ &= \frac{4(q^6; q^6)_\infty^4}{(q^2; q^2)_\infty (q^3; q^3)_\infty^2} + \frac{6(-q^3; q^3)_\infty}{(q^3; q^3)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+6n+1}}{1 - q^{6n+2}} \\ &= \frac{(q^6; q^6)_\infty}{(q^2, q^4; q^6)_\infty} \frac{4}{(q^3; q^6)_\infty^2} + \frac{6(-q^3; q^3)_\infty}{(q^3; q^3)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+6n+1}}{1 - q^{6n+2}}, \end{aligned}$$

together with (4.4) and (4.6), we get

$$\begin{aligned} &\sum_{n=0}^{\infty} (\overline{N}_2(0, 6, 3n+2) + \overline{N}_2(3, 6, 3n+2) - \overline{N}_2(1, 6, 3n+2) - \overline{N}_2(2, 6, 3n+2)) q^n \\ &= \frac{(q^6; q^6)_\infty}{(q^2, q^4; q^6)_\infty} \frac{4}{(q^3; q^6)_\infty^2} + \frac{6(-q^3; q^3)_\infty}{(q^3; q^3)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+6n+1}}{1 - q^{6n+2}}. \end{aligned}$$

By Theorem 4.1, we see that the coefficient of  $q^n$  in

$$\frac{(q^6; q^6)_\infty}{(q^2, q^4; q^6)_\infty}$$

is nonnegative for  $n \geq 0$ .

We now consider the positivity of

$$\frac{(-q^3; q^3)_\infty}{(q^3; q^3)_\infty} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+6n+1}}{1 - q^{6n+2}}. \quad (4.14)$$

The sum in (4.14) equals

$$\begin{aligned} & \sum_{n=0}^{\infty} \frac{(-1)^n q^{3n^2+6n+1}}{1 - q^{6n+2}} + \sum_{n=0}^{\infty} \frac{(-1)^n q^{3n^2+6n+2}}{1 - q^{6n+4}} \\ &= \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+1}}{1 - q^{12n+2}} - \sum_{n=0}^{\infty} \frac{q^{12n^2+24n+10}}{1 - q^{12n+8}} + \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+2}}{1 - q^{12n+4}} - \sum_{n=0}^{\infty} \frac{q^{12n^2+24n+11}}{1 - q^{12n+10}} \\ &= \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+1}}{(1 - q^{12n+2})(1 - q^{12n+8})} \left( (1 - q^{12n+8})(1 - q^{12n+9}) + q^{24n+11}(1 - q^6) \right) \\ & \quad + \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+2}}{(1 - q^{12n+4})(1 - q^{12n+10})} \left( (1 - q^{12n+9})(1 - q^{12n+10}) + q^{24n+13}(1 - q^6) \right) \\ &= (1 - q^3) \left( \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+1}}{1 - q^{12n+2}} \frac{1 - q^{12n+9}}{1 - q^3} + \sum_{n=0}^{\infty} \frac{q^{12n^2+36n+12}(1 + q^3)}{(1 - q^{12n+2})(1 - q^{12n+8})} \right. \\ & \quad \left. + \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+2}}{1 - q^{12n+4}} \frac{1 - q^{12n+9}}{1 - q^3} + \sum_{n=0}^{\infty} \frac{q^{12n^2+36n+15}(1 + q^3)}{(1 - q^{12n+4})(1 - q^{12n+10})} \right) \\ &= (1 - q^3) \left( \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+1}}{1 - q^{12n+2}} \sum_{m=0}^{4n+2} q^{3m} + \sum_{n=0}^{\infty} \frac{q^{12n^2+36n+12}(1 + q^3)}{(1 - q^{12n+2})(1 - q^{12n+8})} \right. \\ & \quad \left. + \sum_{n=0}^{\infty} \frac{q^{12n^2+12n+2}}{1 - q^{12n+4}} \sum_{m=0}^{4n+2} q^{3m} + \sum_{n=0}^{\infty} \frac{q^{12n^2+36n+15}(1 + q^3)}{(1 - q^{12n+4})(1 - q^{12n+10})} \right). \quad (4.15) \end{aligned}$$

Noting that

$$\frac{(-q^3; q^3)_\infty}{(q^3; q^3)_\infty} (1 - q^3) = \frac{(-q^3; q^3)_\infty}{(q^6; q^3)_\infty}, \quad (4.16)$$

it follows from (4.15) and (4.16) that the coefficient of  $q^n$  in (4.14) is nonnegative for  $n \geq 0$ . This completes the proof.  $\blacksquare$

## 5 Proofs of Theorems 1.5 and 1.6

This section is devoted to the proofs of the relations between the  $M_2$ -rank differences of overpartitions and mock theta functions as stated in Theorems 1.5 and 1.6. It is known that mock theta functions can be expressed in terms of the Appell-Lerch sum  $m(x, q, z)$ . Recall that the Appell-Lerch sum is defined by

$$m(x, q, z) = \frac{1}{j(z; q)} \sum_{r=-\infty}^{\infty} \frac{(-1)^r q^{\binom{r}{2}} z^r}{1 - q^{r-1} x z}, \quad (5.1)$$

where  $x, z \in \mathbb{C}^*$  with neither  $z$  nor  $xz$  an integral power of  $q$ .

Hickerson and Mortenson [12] proved that

$$\rho(q) = q^{-1} m(q, q^6, -q), \quad (5.2)$$

$$\phi(q) = -q^{-1} m(q, q^{10}, q) - q^{-1} m(q, q^{10}, q^2), \quad (5.3)$$

$$\psi(q) = -m(q^3, q^{10}, q) - m(q^3, q^{10}, q^3). \quad (5.4)$$

We are ready to present the proofs of Theorems 1.5 and 1.6.

*Proof of Theorem 1.5.* From Theorem 1.1, we have

$$\overline{R}_{0,2}(2, 6) = \frac{4J_6^9 J_{2,4}^2}{J_{1,6}^4 J_2^3 J_{3,6}^3} + \frac{2q^2}{J_{3,6}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2+6n}}{1 + q^{6n+4}} \quad (5.5)$$

$$\begin{aligned} &= \frac{4J_6^9 J_{2,4}^2}{J_{1,6}^4 J_2^3 J_{3,6}^3} - \frac{2q^2}{J_{3,6}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{3n^2-3}}{1 + q^{6n-2}} \\ &= \frac{4J_6^9 J_{2,4}^2}{J_{1,6}^4 J_2^3 J_{3,6}^3} - 2q^{-1} m(-q, q^6, q^3). \end{aligned} \quad (5.6)$$

Replacing  $q$  by  $-q$  in (5.2), we deduce that

$$\rho(-q) = -q^{-1} m(-q, q^6, q). \quad (5.7)$$

Recall the following identity in Hickerson and Mortenson [12]:

$$m(x, q, z_1) - m(x, q, z_0) = \frac{z_0 J_1^3 j(z_1/z_0; q) j(xz_0 z_1; q)}{j(z_0; q) j(z_1; q) j(xz_0; q) j(xz_1; q)}. \quad (5.8)$$

Replacing  $q, z_0, z_1$  and  $x$  with  $q^6, q, q^3$  and  $-q$ , respectively, in (5.8), we get

$$m(-q, q^6, q^3) - m(-q, q^6, q) = \frac{q J_6^3 J_2 \overline{J}_{1,6}}{J_{1,6} J_{3,6} \overline{J}_2^2}. \quad (5.9)$$

Combining (5.7) and (5.9), we obtain

$$m(-q, q^6, q^3) = -q\rho(-q) + \frac{q J_6^3 J_2 \overline{J}_{1,6}}{J_{1,6} J_{3,6} \overline{J}_2^2}. \quad (5.10)$$

Substituting (5.10) into (5.6), we get (1.28). This completes the proof of Theorem 1.5. ■

We conclude with a proof of Theorem 1.6.

*Proof of Theorem 1.6.* To prove (1.29), we apply Theorem 1.2 to get

$$\begin{aligned}
\bar{R}_{0,4}(4, 10) &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} A_4(q) + \frac{2q^4}{J_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2+10n}}{1+q^{10n+8}} \\
&= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} A_4(q) - \frac{2}{qJ_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2}}{1+q^{10n-2}} \\
&= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} A_4(q) - 2q^{-1} m(-q^3, q^{10}, q^5), \tag{5.11}
\end{aligned}$$

where  $A_4(q)$  is defined as (1.12). Replacing  $q$  by  $-q$  in (5.4), we are led to

$$\psi(-q) = -m(-q^3, q^{10}, -q) - m(-q^3, q^{10}, -q^3). \tag{5.12}$$

Replacing  $q, z_0, z_1$  and  $x$  by  $q^{10}, -q, q^5$  and  $-q^3$  in (5.8), respectively, we find that

$$m(-q^3, q^{10}, q^5) - m(-q^3, q^{10}, -q) = -\frac{qJ_{10}^3 \bar{J}_{4,10} J_{1,10}}{J_{4,10} J_{5,10} \bar{J}_{1,10} \bar{J}_{2,10}}. \tag{5.13}$$

Replacing  $q, z_1, z_0$  and  $x$  in (5.8) by  $q^{10}, q^5, -q^3$  and  $-q^3$ , respectively, we see that

$$m(-q^3, q^{10}, q^5) - m(-q^3, q^{10}, -q^3) = \frac{q^2 J_{10}^3 J_{1,10} \bar{J}_{2,10}}{J_{4,10} J_{5,10} \bar{J}_{2,10} \bar{J}_{3,10}}. \tag{5.14}$$

Combining (5.13) and (5.14) gives

$$\begin{aligned}
m(-q^3, q^{10}, q^5) &= \frac{1}{2} m(-q^3, q^{10}, -q) + \frac{1}{2} m(-q^3, q^{10}, -q^3) \\
&\quad + \frac{q^2 J_{10}^3 J_{1,10} \bar{J}_{2,10}}{2J_{4,10} J_{5,10} \bar{J}_{2,10} \bar{J}_{3,10}} - \frac{qJ_{10}^3 \bar{J}_{4,10} J_{1,10}}{2J_{4,10} J_{5,10} \bar{J}_{1,10} \bar{J}_{2,10}}. \tag{5.15}
\end{aligned}$$

Utilizing (5.12) and (5.15),  $m(-q^3, q^{10}, q^5)$  can be written as

$$m(-q^3, q^{10}, q^5) = -\frac{1}{2} \psi(-q) + \frac{q^2 J_{10}^3 J_{1,10} \bar{J}_{2,10}}{2J_{4,10} J_{5,10} \bar{J}_{2,10} \bar{J}_{3,10}} - \frac{qJ_{10}^3 \bar{J}_{4,10} J_{1,10}}{2J_{4,10} J_{5,10} \bar{J}_{1,10} \bar{J}_{2,10}}. \tag{5.16}$$

Substituting (5.16) into (5.11), we implies (1.29).

We now consider (1.30). By Theorem 1.3, we find that

$$\begin{aligned}
\bar{R}_{1,3}(1, 10) &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_1(q) - \frac{2q^4}{J_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2+10n}}{1+q^{10n+6}} \\
&= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_1(q) + \frac{2}{qJ_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2}}{1+q^{10n-4}}
\end{aligned}$$

$$= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_1(q) + 2q^{-1} m(-q, q^{10}, q^5), \quad (5.17)$$

where  $B_1(q)$  is defined as in (1.17). When  $q$  is replaced by  $-q$ , (5.3) becomes

$$\phi(-q) = q^{-1} m(-q, q^{10}, -q) + q^{-1} m(-q, q^{10}, q^2). \quad (5.18)$$

Replacing  $q, z_0, z_1$  and  $x$  by  $q^{10}, -q, q^5$  and  $-q$  in (5.8), respectively, we get

$$m(-q, q^{10}, q^5) - m(-q, q^{10}, -q) = -\frac{qJ_{10}^3 J_{3,10} \bar{J}_{4,10}}{J_{2,10} J_{5,10} \bar{J}_{1,10} \bar{J}_{4,10}}. \quad (5.19)$$

Replacing  $q, z_0, z_1$  and  $x$  by  $q^{10}, q^2, q^5$  and  $-q$  in (5.8), we obtain that

$$m(-q, q^{10}, q^5) - m(-q, q^{10}, q^2) = \frac{q^2 J_{10}^3 J_{3,10} \bar{J}_{2,10}}{J_{2,10} J_{5,10} \bar{J}_{3,10} \bar{J}_{4,10}}. \quad (5.20)$$

It follows from (5.19) and (5.20) that

$$\begin{aligned} m(-q, q^{10}, q^5) &= \frac{1}{2} m(-q, q^{10}, -q) + \frac{1}{2} m(-q, q^{10}, q^2) \\ &\quad - \frac{qJ_{10}^3 J_{3,10} \bar{J}_{4,10}}{2J_{2,10} J_{5,10} \bar{J}_{1,10} \bar{J}_{4,10}} + \frac{q^2 J_{10}^3 J_{3,10} \bar{J}_{2,10}}{2J_{2,10} J_{5,10} \bar{J}_{3,10} \bar{J}_{4,10}}. \end{aligned} \quad (5.21)$$

In view of (5.18) and (5.21),  $m(-q, q^{10}, q^5)$  can be rewritten as

$$m(-q, q^{10}, q^5) = \frac{q}{2} \phi(-q) - \frac{qJ_{10}^3 J_{3,10} \bar{J}_{4,10}}{2J_{2,10} J_{5,10} \bar{J}_{1,10} \bar{J}_{4,10}} + \frac{q^2 J_{10}^3 J_{3,10} \bar{J}_{2,10}}{2J_{2,10} J_{5,10} \bar{J}_{3,10} \bar{J}_{4,10}}. \quad (5.22)$$

Substituting (5.22) into (5.17), we obtain (1.30).

Finally, we prove (1.31). By Theorem 1.3, we see that

$$\begin{aligned} \bar{R}_{1,3}(4, 10) &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_4(q) - \frac{2q^4}{J_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2+10n}}{1+q^{10n+8}} \\ &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_4(q) + \frac{2}{qJ_{5,10}} \sum_{n=-\infty}^{\infty} \frac{(-1)^n q^{5n^2}}{1+q^{10n-2}} \\ &= \frac{2J_{10}^4 J_{5,10}}{J_{1,2}^6} B_4(q) + 2q^{-1} m(-q^3, q^{10}, q^5), \end{aligned} \quad (5.23)$$

where  $B_4(q)$  is given in (1.20). Substituting (5.16) into (5.23), we arrive at (1.31). This completes the proof.  $\blacksquare$

**Acknowledgments.** This work was supported by the National Science Foundation of China.

## References

- [1] G.E. Andrews, The ranks and cranks of partitions moduli 2, 3, and 4, *J. Number Theory* 85 (2000) 74–84.
- [2] G.E. Andrews and F.G. Garvan, Ramanujan’s “lost” notebook VI: The mock theta conjectures, *Adv. Math.* 73 (2) (1989) 245–255.
- [3] A.O.L. Atkin and S.M. Hussain, Some properties of partitions. II, *Trans. Amer. Math. Soc.* 89 (1958) 184–200.
- [4] A.O.L. Atkin and P. Swinnerton-Dyer, Some properties of partitions, *Proc. Lond. Math. Soc.* 66 (1954) 84–106.
- [5] S.H. Chan, Generalized Lambert series identities, *Proc. London Math. Soc.* 91 (2005) 598–622.
- [6] Y.-S. Choi, Tenth order mock theta functions in Ramanujan’s lost notebook, *Invent. Math.* 136 (1999) 497–569.
- [7] S. Corteel and J. Lovejoy, Overpartitions, *Trans. Amer. Math. Soc.* 356 (2004) 1623–1635.
- [8] F.J. Dyson, Some guesses in the theory of partitions, *Eureka (Cambridge)* 8 (1944) 10–15.
- [9] F.G. Garvan, New combinatorial interpretations of Ramanujan’s partition congruences mod 5, 7, 11, *Trans. Amer. Math. Soc.* 305 (1988) 47–77.
- [10] F.G. Garvan and J. Liang, Automatic proof of theta-function identities, <http://qseries.org/fgarvan/qmaple/thetaids/auttheta.pdf> Aug. 2016.
- [11] D. Hickerson, On the seventh order mock theta functions, *Invent. Math.* 94 (1988) 661–677.
- [12] D. Hickerson and E. Mortenson, Hecke-type double sums, Appell-Lerch sums, and mock theta functions, I, *Proc. Lond. Math. Soc.* (3) 109 (2014) 382–422.
- [13] C. Jennings-Shaffer, Higher order SPT functions for overpartitions, overpartitions with smallest part even, and partitions with smallest part even and without repeated odd parts, *J. Number Theory* 149 (2015) 285–312.
- [14] R.P. Lewis, On the ranks of partitions modulo 9, *Bull. London Math. Soc.* 23 (1991) 417–421.
- [15] R.P. Lewis, The ranks of partitions modulo 2, *Discrete Math.* 167/168 (1997) 445–449.
- [16] W.-C. Liaw, Note on the monotonicity of coefficients for some  $q$ -series arising from Ramanujan’s lost notebook, *Ramanujan J.* 3 (4) (1999) 385–388.
- [17] J. Lovejoy, Rank and conjugation for the Frobenius representation of an overpartition, *Ann. Comb.* 9 (3) (2005) 321–334.
- [18] J. Lovejoy, Rank and conjugation for a second Frobenius representation of an overpartition, *Ann. Comb.* 12 (2008) 101–113.
- [19] J. Lovejoy and R. Osburn, Rank differences for overpartitions, *Quart. J. Math.* 59 (2008) 257–273.
- [20] J. Lovejoy and R. Osburn,  $M_2$ -rank differences for overpartitions, *Acta Arith.* 144 (2) (2010) 193–212.
- [21] R. Mao, Ranks of partitions modulo 10, *J. Number Theory* 133 (2013) 3678–3702.
- [22] R. Mao, The  $M_2$ -rank of partitions without repeated odd parts modulo 6 and 10. *Ramanujan J.* 37 (2) (2015) 391–419.



- [23] G.N. Watson, The final problem: an account of the mock theta functions, J. London Math. Soc. 11 (1936) 55–80.
- [24] S. Robins, Generalized Dedekind  $\eta$ -products, in: The Rademacher Legacy to Mathematics (University Park, PA, 1992), 119–128 Contemp. Math. 166 Amer. Math. Soc. Providence RI 1994.
- [25] N. Santa-Gadea, On the ranks of partitions mod 9 and 12, J. Number Theory 40 (1992) 130–145.