# ON THE SUPREMUM OF PRODUCTS OF SYMMETRIC STABLE PROCESSES

CHRISTOPHE PROFETA

Abstract. We study the asymptotics, for small and large values, of the supremum of a product of symmetric stable processes. We show in particular that the persistence exponent remains the same as for only one process, up to some logarithmic terms.

## 1. INTRODUCTION

For  $n \in \mathbb{N}$ , let  $(Z^{(i)}, 1 \le i \le n)$  be independent symmetric  $\alpha$ -stable Lévy processes with  $\alpha \in (0, 2]$ . In this short note, we are interested in the study of the random variable

$$
\mathcal{S}_n = \sup_{0 \le u \le 1} \prod_{i=1}^n Z_u^{(i)}.
$$

Yn

Except when  $n = 1$ , in which case the double Laplace transform of  $S_1$  is classically given by fluctuation theory (see for instance Bertoin [\[4,](#page-10-0) p.174]), it does not seem evident to compute explicitly the law of  $\mathcal{S}_n$ , and we shall rather study its asymptotics  $\mathbb{P}(\mathcal{S}_n \geq x)$  as  $x \to +\infty$  and  $\mathbb{P}(\mathcal{S}_n \leq \varepsilon)$  as  $\varepsilon \to 0$ .

Most of the paper is devoted to the computation of the limit as  $\varepsilon \to 0$ , which is known as a persistence problem, see the surveys [\[1,](#page-10-1) [5\]](#page-10-2). By scaling, this amounts to the study of the first entrance time of the *n*-dimensional stable process  $(Z^{(i)}, 1 \le i \le n)$  into the "hyperbolic" domain  $\mathcal{H}_n = \{(z_1, \ldots, z_n) \in \mathbb{R}^n, \prod_{i=1}^n z_i \geq 1\}$ :

$$
\mathbb{P}\left(\mathcal{S}_n \leq \varepsilon\right) = \mathbb{P}\left(R_n > \frac{1}{\varepsilon^{\frac{\alpha}{n}}}\right) \qquad \text{where} \quad R_n = \inf\left\{u \geq 0, \prod_{i=1}^n Z_u^{(i)} \geq 1\right\}.
$$

There are several papers in the literature dealing with entrance and exit times of symmetric sta-ble processes, mainly for three families of domains : cones and wedges (Bañuelos and Bogdan [\[2\]](#page-10-3), Méndez-Hernández [\[9\]](#page-10-4)), parabolic domains (Bañuelos and Bogdan [\[3\]](#page-10-5)) and unbounded convex do-mains (Méndez-Hernández [\[8\]](#page-10-6)). Here, since the domain  $\mathcal{H}_n$  is non-connected, not much is known regarding  $R_n$  and we shall tackle the problem directly by working with  $S_n$ .

We start with the Brownian case, i.e.  $\alpha = 2$ .

**Theorem 1.** Let  $(W^{(i)}, 1 \leq i \leq n)$  be independent Brownian motions. There exist two constants  $0 < \kappa_1 \leq \kappa_2 < +\infty$  such that the following estimates hold.

(1) Large deviations :

$$
\kappa_1 x^{-\frac{1}{n}} \exp\left(-\frac{n}{2}x^{\frac{2}{n}}\right) \le \mathbb{P}\left(\sup_{0\le u\le 1} \prod_{i=1}^n W_u^{(i)} \ge x\right) \le \kappa_2 x^{-\frac{1}{n}} \exp\left(-\frac{n}{2}x^{\frac{2}{n}}\right) \qquad (x \to +\infty)
$$

2010 Mathematics Subject Classification. 60G52, 60J65.

Key words and phrases. Persistence probability - Stable processes.

(2) Persistence probability :

$$
\kappa_1 \,\varepsilon \le \mathbb{P}\left(\sup_{0\le u\le 1}\prod_{i=1}^n W_u^{(i)} \le \varepsilon\right) \le \kappa_2 \,\varepsilon \left|\ln(\varepsilon)\right|^n \qquad (\varepsilon \to 0)
$$

In the non-Gaussian stable case, the situation is different.

<span id="page-1-0"></span>**Theorem 2.** Let  $(L^{(i)}, 1 \leq i \leq n)$  be independent symmetric  $\alpha$ -stable Lévy processes with  $\alpha \in (0, 2)$ . There exist two constants  $0 < \kappa_1 \leq \kappa_2 < +\infty$  such that the following estimates hold.

(1) Large deviations :

$$
\kappa_1 \frac{(\ln(x))^{n-1}}{x^{\alpha}} \le \mathbb{P}\left(\sup_{0 \le u \le 1} \prod_{i=1}^n L_u^{(i)} \ge x\right) \le \kappa_2 \frac{(\ln(x))^{n-1}}{x^{\alpha}} \qquad (x \to +\infty)
$$

(2) Persistence probability :

$$
\kappa_1 \,\varepsilon^{\alpha/2} \le \mathbb{P}\left(\sup_{0\le u\le 1} \prod_{i=1}^n L_u^{(i)} \le \varepsilon\right) \le \kappa_2 \,\varepsilon^{\alpha/2} \,|\ln(\varepsilon)| \qquad (\varepsilon \to 0)
$$

Remark 3. The presence of extra logarithmic terms in the persistence probability of Brownian motion is due to some additive phenomenons. Indeed, recall the estimates (see Bertoin [\[4,](#page-10-0) p.219]) :

<span id="page-1-1"></span>
$$
\mathbb{P}(|Z_1| \leq \varepsilon) \underset{\varepsilon \to 0}{\sim} k \varepsilon \quad \text{and} \quad \mathbb{P}\left(\sup_{0 \leq u \leq 1} Z_u \leq \varepsilon\right) \underset{\varepsilon \to 0}{\sim} c \varepsilon^{\frac{\alpha}{2}} \tag{1}
$$

for some positive constants k and c. When  $\alpha < 2$ , the second asymptotics is the leading one, while for  $\alpha = 2$ , they are of the same order, and some compensations appear, see Lemma [4.](#page-2-0) In fact, the heuristic below leads us to believe that the right asymptotics in the Brownian case should be  $\varepsilon \left|\ln(\varepsilon)\right|^{n-1}.$ 

The main part of the proof deals with the computation of an upper bound for the persistence probabilities. A simple approach would be to try to bound the quantity  $S_n$  by  $\prod_{i=1}^n Z_{\theta_1}^{(i)}$  $\theta_1^{(i)}$  where  $\theta_1$ is the value at which one of the Lévy processes, say  $Z^{(n)}$ , reach its maximum on [0, 1]. This yields of course two main difficulties.

- i) First, the product of the other processes  $\prod_{i=1}^{n-1} Z^{(i)}_{\theta_1}$  $\theta_1^{(i)}$  might not be positive. This can be however easily circumvented thanks to Slepian's inequality, since the processes are symmetric.
- ii) The second difficulty is less obvious and is due to the arcsine law for stable processes. There is a high probability that  $\theta_1$  will be close to 0, hence, although  $Z_{\theta_1}^{(n)}$  $\theta_1^{(n)}$  will be large, the remaining product  $\prod_{i=1}^{n-1} Z_{\theta_1}^{(i)}$  $\theta_1^{(i)}$  will also be close to zero, thus not providing us with a good upper bound.

The general idea of the proof will be to decompose the path of the processes  $(Z^{(i)})$  at some last passage times and then use a time-reversal argument, so as to find a value not to close to the origin at which  $Z^{(n)}$  is large enough.

The outline of the paper is as follows : the large deviation results are proved in Section 2, the persistence probabilities in Section 3, and finally Section 4 provides the proof of an intermediary lemma.

## 2. Large deviations

The proof of the large deviation results relies on the symmetry of the processes  $(Z^{(i)})$ , and on the fact that the asymptotics of both random variables  $|Z_1|$  and sup  $0 \le u \le 1$  $Z_u$  are similar. Indeed, on the one hand, the lower bound is easily given by :

$$
\mathbb{P}(\mathcal{S}_n \geq x) \geq \mathbb{P}\left(\prod_{i=1}^n Z_1^{(i)} \geq x\right) = \frac{1}{2} \mathbb{P}\left(\prod_{i=1}^n |Z_1^{(i)}| \geq x\right).
$$

On the other hand, still by symmetry,

$$
\mathbb{P}(\mathcal{S}_n \ge x) \le \mathbb{P}\left(\sup_{0 \le u \le 1} Z_u^{(1)} \sup_{0 \le s \le 1} \prod_{i=1}^{n-1} Z_s^{(i)} \ge x\right) + \mathbb{P}\left(\inf_{0 \le u \le 1} Z_u^{(1)} \inf_{0 \le s \le 1} \prod_{i=1}^{n-1} Z_s^{(i)} \ge x\right)
$$
  

$$
\le 2 \mathbb{P}\left(\sup_{0 \le u \le 1} Z_u^{(1)} \sup_{0 \le s \le 1} \prod_{i=1}^{n-1} Z_s^{(i)} \ge x\right)
$$
  

$$
\le 2^n \mathbb{P}\left(\prod_{i=1}^n \sup_{0 \le u \le 1} Z_u^{(i)} \ge x\right) \qquad \text{(by iteration)}.
$$

It remains thus to compute the involved quantities in both cases.

 $\rightarrow$  In the Brownian case, since sup  $0 \le u \le 1$  $W_u \stackrel{\text{(law)}}{=} |W_1|$ , we deduce that the asymptotics of  $\mathcal{S}_n$  is given by that of  $\prod_{i=1}^{n} |W_1^{(i)}|$  $\binom{n}{1}$ . Its Mellin transform reads, for  $\nu > -1$ :

$$
\mathbb{E}\left[\prod_{i=1}^{n}|W_1^{(i)}|^{\nu}\right] = \left(\frac{2^{\nu}}{\pi}\right)^{\frac{n}{2}}\left(\Gamma\left(\frac{1+\nu}{2}\right)\right)^n.
$$
\n(2)

The converse mapping theorem, see Janson [\[7,](#page-10-7) Theorem 6.1], yields :

$$
\mathbb{P}\left(\prod_{i=1}^n|W_1^{(i)}|\in dx\right)/dx\underset{x\rightarrow+\infty}{\sim}\kappa\,x^{\frac{1}{n}-1}\,e^{-\frac{n}{2}x^{\frac{2}{n}}}
$$

for some positive constant  $\kappa$ . The result then follows by integration, using the asymptotics of the incomplete Gamma function.

 $\rightarrow$  Next, when  $\alpha \in (0, 2)$ , it is known from Bertoin [\[4,](#page-10-0) p.221] that there exists  $k > 0$  such that

$$
\mathbb{P}(|L_1| \geq x) \underset{x \to +\infty}{\sim} \frac{2k}{x^{\alpha}} \quad \text{and} \quad \mathbb{P}\left(\sup_{0 \leq u \leq 1} L_u \geq x\right) \underset{x \to +\infty}{\sim} \frac{k}{x^{\alpha}}.
$$

Point 1. of Theorem [2](#page-1-0) is then consequence of the following lemma (see for instance Lemma 2 in Profeta-Simon [\[10\]](#page-10-8)):

<span id="page-2-0"></span>**Lemma 4.** Let X and Y be two independent positive random variables satisfying the asymptotics :

$$
\mathbb{P}(X \ge z) \underset{z \to +\infty}{\sim} a \frac{(\ln(z))^n}{z^{\nu}} \qquad and \qquad \mathbb{P}(Y \ge z) \underset{z \to +\infty}{\sim} b \frac{(\ln(z))^p}{z^{\mu}}
$$

where  $n, p \in \mathbb{N}$  and  $a, b, \nu, \mu$  are positive constants such that  $0 < \nu \leq \mu$ . Then there exists  $c > 0$ such that :

$$
\mathbb{P}(XY \ge z) \underset{z \to +\infty}{\sim} \begin{cases} c \, z^{-\nu} (\ln(z))^n & \text{if } \nu < \mu \\ c \, z^{-\nu} (\ln(z))^{n+p+1} & \text{if } \nu = \mu. \end{cases}
$$

### 3. Persistence probabilities

We now turn our attention to the persistence estimates and start with some notations. Let X be a symmetric stable process. We denote by  $\mathbb{P}_x$  the probability measure of X when started from  $x \in \mathbb{R}$ , with the usual convention that  $\mathbb{P} = \mathbb{P}_0$ . Let  $T_0$  be the first time that X takes a negative value :

$$
T_0 = \inf\{t \ge 0, X_t \le 0\}.
$$

We recall from Bertoin [\[4,](#page-10-0) p.219] that since X is symmetric, there exists  $c > 0$  such that

<span id="page-3-0"></span>
$$
\mathbb{P}_1(T_0 \ge t) \underset{t \to +\infty}{\sim} \frac{c}{\sqrt{t}}.\tag{3}
$$

Finally, let us introduce the last change of sign of X before time  $t > 0$ :

$$
g_t = \sup \{ 0 \le u \le t, X_u X_{u^-} \le 0 \}.
$$

This random time will be the key to the computation of the persistence probabilities.

**Remark 5.** In the following, when applying the Markov property,  $\widehat{X}$  will always denote an independent copy of X. Besides, we shall use the notations c and  $\kappa$  to denote positive constants that may change from line to line.

We first show that the asymptotics of the distribution of  $g_1$  is similar to that of the arcsine law.

<span id="page-3-1"></span>Lemma 6. There exists a positive constant c such that

$$
\mathbb{P}(g_1 \in dr) / dr \underset{r \to 0}{\sim} \frac{c}{\sqrt{r}}.
$$

*Proof.* We first have, using the symmetry of X and applying the Markov property with  $r \in (0,1)$ :

$$
\mathbb{P}(g_1 \leq r) = \mathbb{E}\left[\widehat{\mathbb{P}}_{|X_r|}\left(\widehat{T}_0 \geq 1-r\right)\right].
$$

By scaling, this is further equal to

$$
\mathbb{P}(g_1 \le r) = \mathbb{E}\left[\widehat{\mathbb{P}}_1\left(\widehat{T}_0 \ge \frac{1-r}{r|X_1|^{\alpha}}\right)\right].
$$

Recall now from Doney-Savov [\[6\]](#page-10-9) that under  $\mathbb{P}_1$ , the random variable  $T_0$  admits a continuous density h satisfying  $h(z)$ <sub>z→+∞</sub>  $\kappa z^{-3/2}$  for some constant  $\kappa > 0$ . Therefore, differentiating, we deduce that

$$
\mathbb{P}(g_1 \in dr)/dr = \frac{1}{r^2} \mathbb{E}\left[\frac{1}{|X_1|^\alpha} h\left(\frac{1-r}{r|X_1|^\alpha}\right)\right] \underset{r \to 0}{\sim} \frac{\kappa}{\sqrt{r}} \mathbb{E}\left[|X_1|^{\frac{\alpha}{2}}\right]
$$

which is the announced result.  $\Box$ 

# 3.1. Lower bound for the persistence probabilities. Observe first that by scaling

$$
\mathbb{P}(\mathcal{S}_n \le \varepsilon) = \mathbb{P}\left(\sup_{u \in [0,\varepsilon^{-\alpha/n}]} \prod_{i=1}^n Z_u^{(i)} \le 1\right)
$$
  
\n
$$
\ge \mathbb{P}\left(\sup_{u \in [0,\varepsilon^{-\alpha/n}]} \prod_{i=1}^n Z_u^{(i)} \le 1, \prod_{i=1}^n Z_1^{(i)} \le 0, \sup_{1 \le i \le n} g_{1/\varepsilon}^{(i)} \le 1\right)
$$
  
\n
$$
= \mathbb{P}\left(\sup_{u \in [0,1]} \prod_{i=1}^n Z_u^{(i)} \le 1, \prod_{i=1}^n Z_1^{(i)} \le 0, \sup_{1 \le i \le n} g_{1/\varepsilon}^{(i)} \le 1\right)
$$

where the last equality follows from the fact that, by definition of the  $(g^{(i)})$ , the product  $\prod_{i=1}^{n} Z^{(i)}$ remains negative after time 1. We now apply the Markov property at time 1 :

$$
\mathbb{P}(\mathcal{S}_n \le \varepsilon) \ge \mathbb{E}\left[\prod_{i=1}^n \widehat{\mathbb{P}}_{|Z_1^{(i)}|}\left(\widehat{T}_0^{(i)} \ge \frac{1}{\varepsilon^{\frac{\alpha}{n}}} - 1\right), \sup_{u \in [0,1]} \prod_{i=1}^n Z_u^{(i)} \le 1, \prod_{i=1}^n Z_1^{(i)} \le 0\right]
$$
  
 
$$
\ge \mathbb{E}\left[\prod_{i=1}^n \widehat{\mathbb{P}}_1\left(\widehat{T}_0^{(i)} \ge \frac{1}{\varepsilon^{\frac{\alpha}{n}} |Z_1^{(i)}|^{\alpha}}\right), \sup_{u \in [0,1]} \prod_{i=1}^n Z_u^{(i)} \le 1, \prod_{i=1}^n Z_1^{(i)} \le 0\right].
$$
 (4)

From [\(3\)](#page-3-0), there exists  $\kappa > 0$  such that for  $\delta > 0$  small enough

<span id="page-4-0"></span>**110** 

$$
\widehat{\mathbb{P}}_1\left(\widehat{T}_0^{(i)} \geq \frac{1}{\varepsilon^{\frac{\alpha}{n}} |Z_1^{(i)}|^\alpha} \right) 1\!\!1_{\{\varepsilon^{\frac{\alpha}{n}} |Z_1^{(i)}|^\alpha \leq \delta\}} \geq \kappa \, \varepsilon^{\frac{\alpha}{2n}} |Z_1^{(i)}|^{\frac{\alpha}{2}} 1\!\!1_{\{\varepsilon^{\frac{\alpha}{n}} |Z_1^{(i)}|^\alpha \leq \delta\}}.
$$

Plugging this inequality in [\(4\)](#page-4-0), we deduce that

$$
\mathbb{P}(\mathcal{S}_n \leq \varepsilon) \geq \kappa^n \varepsilon^{\frac{\alpha}{2}} \mathbb{E} \left[ \prod_{i=1}^n |Z_1^{(i)}|^{\frac{\alpha}{2}} 1_{\{\varepsilon^{\frac{\alpha}{n}} |Z_1^{(i)}|^{\alpha} \leq \delta\}}, \sup_{u \in [0,1]} \prod_{i=1}^n Z_u^{(i)} \leq 1, \prod_{i=1}^n Z_1^{(i)} \leq 0 \right]
$$

$$
\sim_{\varepsilon \to 0} \kappa^n \varepsilon^{\frac{\alpha}{2}} \mathbb{E} \left[ \prod_{i=1}^n |Z_1^{(i)}|^{\frac{\alpha}{2}}, \sup_{u \in [0,1]} \prod_{i=1}^n Z_u^{(i)} \leq 1, \prod_{i=1}^n Z_1^{(i)} \leq 0 \right]
$$

which gives the lower bound.

3.2. Upper bound for the persistence probabilities. Since all the processes  $(Z^{(i)})$  have the same law, we first have :

$$
\mathbb{P}\left(\mathcal{S}_n \leq \varepsilon\right) = n \, \mathbb{P}\left(\sup_{0 \leq u \leq 1} \prod_{i=1}^n Z_u^{(i)} \leq \varepsilon, \ g_1^{(n)} \geq \sup_{1 \leq i \leq n-1} g_1^{(i)}\right).
$$

To simplify the notation, we shall remove the superscript  $(n)$  and denote

$$
X = Z^{(n)}
$$
,  $g_1 = g_1^{(n)}$  and  $\xi_t = \sup_{1 \le i \le n-1} g_t^{(i)}$ .

This yields, with the usual convention that empty products equal 1,

$$
\mathbb{P}(\mathcal{S}_n \le \varepsilon) = n \mathbb{P}\left(\sup_{0 \le u \le 1} X_u \prod_{i=1}^{n-1} Z_u^{(i)} \le \varepsilon, \ g_1 \ge \xi_1\right)
$$

$$
\le n \mathbb{P}\left(\sup_{0 \le u < g_1} X_u \prod_{i=1}^{n-1} Z_u^{(i)} \le \varepsilon, \ g_1 \ge \xi_1\right)
$$

$$
= 2n \mathbb{P}\left(\sup_{0\leq u<1} X_{ug_1} \prod_{i=1}^{n-1} Z_{ug_1}^{(i)} \leq \varepsilon, \prod_{i=1}^{n-1} Z_{g_1}^{(i)} \geq 0, g_1 \geq \xi_1\right)
$$

where the last equality follows by symmetry. By scaling, we further obtain

$$
\mathbb{P}\left(\mathcal{S}_n \leq \varepsilon\right) \leq 2n \int_0^1 \mathbb{P}\left(\sup_{0 \leq u < 1} \frac{X_{ur}}{r^{1/\alpha}} \prod_{i=1}^{n-1} Z_u^{(i)} r^{\frac{n}{\alpha}} \leq \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \geq 0, \ 1 \geq \xi_{\frac{1}{r}} \middle| \ g_1 = r\right) \mathbb{P}(g_1 \in dr).
$$

We set  $X^{(x,t,y)}$  for the  $\alpha$ -stable bridge of length t starting from x and ending at y. Notice that when  $X = W$  is a Brownian motion, then  $g_1$  coincides with the last zero of W before time 1, so that  $W_{g_1} = 0$  a.s. and it is well-known that the process  $\left(\frac{W_{ug_1}}{\sqrt{g_1}}, 0 \le u \le 1\right)$  is a standard Brownian bridge, independent from  $q_1$ , see Bertoin [\[4,](#page-10-0) p.230]. We shall extend this result to the stable case in the following lemma, whose proof is postponed at the end of the paper.

<span id="page-5-0"></span>**Lemma 7.** We set by convention  $X_{0}$ - =  $X_{0}$ . Conditionally on the event  $\begin{cases} X_{g_1}^{-1/\alpha} & \text{if } 0 \leq \alpha \leq \alpha \end{cases}$  $g_1^{1/\alpha}$  $= a$  $\mathcal{L}$ , the process  $\bigg(X_{(ug_1)^-}$  $\frac{u_{g_1}}{g_1^{1/\alpha}}, 0 \le u \le 1$ 1 !

is independent from  $g_1$  and has the same law as the stable bridge  $(X_{u^-}^{(0,1,a)}, 0 \le u \le 1)$ .

Let us denote by  $\rho(da, dr)$  the law of the pair  $(q_1^{-1/\alpha} X_{q_1^-}, g_1)$ . Since the  $(Z^{(i)})$  are quasi-left continuous and independent from  $X$ , we deduce from Lemma [7](#page-5-0) that

<span id="page-5-2"></span>
$$
\mathbb{P}(\mathcal{S}_n \le \varepsilon) \le 2n \int_0^{+\infty} \int_0^1 \mathbb{P}\left(\sup_{0 \le u \le 1} X_{u^-}^{(0,1,a)} \prod_{i=1}^{n-1} Z_u^{(i)} r^{\frac{n}{\alpha}} \le \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \ge 0, 1 \ge \xi_{\frac{1}{r}}\right) \rho(da, dr)
$$
  
=  $2n \int_0^{+\infty} \int_0^1 \mathbb{P}\left(\sup_{0 \le u \le 1} X_u^{(a,1,0)} \prod_{i=1}^{n-1} Z_{1-u}^{(i)} r^{\frac{n}{\alpha}} \le \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \ge 0, 1 \ge \xi_{\frac{1}{r}}\right) \rho(da, dr)$  (5)

where the equality follows from the time-reversal property of stable bridges. We shall now decompose the right-hand side of this inequality according as  $\{a \leq 1\}$  or  $\{a > 1\}$ .

3.2.1. The case  $\{a \leq 1\}$ . We start with the term giving the main contribution. Let us denote by  $p_t$ the density of the random variable  $X_t$ , and recall that it is even, and decreasing on  $(0, +\infty)$ . Using the absolute continuity formula of the stable bridge, we get :

<span id="page-5-1"></span>
$$
\iint_0^1 \mathbb{P}\left(\sup_{0\leq u\leq 1} X_u^{(a,1,0)} \prod_{i=1}^{n-1} Z_{1-u}^{(i)} r^{\frac{n}{\alpha}} \leq \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \geq 0, 1 \geq \xi_{\frac{1}{r}}\right) \rho(da, dr)
$$
\n
$$
\leq \iint_0^1 \mathbb{E}\left[\frac{p_{\frac{1}{2}}(a + X_{\frac{1}{2}})}{p_1(a)}, \sup_{0\leq u\leq 1/2} (a + X_u) \prod_{i=1}^{n-1} Z_{1-u}^{(i)} r^{\frac{n}{\alpha}} \leq \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \geq 0, 1 \geq \xi_{\frac{1}{r}}\right] \rho(da, dr)
$$
\n
$$
\leq \frac{p_{\frac{1}{2}}(0)}{p_1(1)} \iint_0^1 \mathbb{P}\left(\sup_{0\leq u\leq 1/2} (a + X_u) \prod_{i=1}^{n-1} Z_{1-u}^{(i)} r^{\frac{n}{\alpha}} \leq \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \geq 0, 1 \geq \xi_{\frac{1}{r}}\right) \rho(da, dr).
$$
\n(6)

We now study the integrand in [\(6\)](#page-5-1). Recall that X admits the representation  $(B_{\tau_u}, u \ge 0)$  where B is a standard Brownian motion and  $\tau$  a stable subordinator with index  $\frac{\alpha}{2}$  independent from B. Let us consider the conditional expectation :

$$
\mathbb{P}\left(\sup_{0\leq u\leq 1/2} (a+B_{\eta_u}) \prod_{i=1}^{n-1} \omega_{1-u}^{(i)} r^{\frac{n}{\alpha}} \leq \varepsilon \middle| \begin{array}{l} Z^{(i)} = \omega^{(i)}, & \tau = \eta \\ 1 \leq i \leq n-1 \end{array} \right)
$$

where  $\eta$  and  $(\omega^{(i)}, 1 \leq i \leq n-1)$  are some fixed càdlàg paths. We apply Slepian's lemma with the Gaussian processes

$$
U_u = z \prod_{i=1}^{n-1} \omega_u^{(i)} + B_{\eta_u} \prod_{i=1}^{n-1} \omega_u^{(i)} \quad \text{and} \quad V_u = a \prod_{i=1}^{n-1} \omega_u^{(i)} + B_{\eta_u} \prod_{i=1}^{n-1} |\omega_u^{(i)}|
$$

which satisfy for every  $0 \le u \le s \le \frac{1}{2}$  $\frac{1}{2}$ ,

$$
\mathbb{E}[U_u] = \mathbb{E}[V_u], \qquad \mathbb{E}[U_u^2] = \mathbb{E}[V_u^2] \quad \text{and} \quad \mathbb{E}[U_u U_s] \le \mathbb{E}[V_u V_s].
$$

This yields, using the tower property of conditional expectations :

$$
\mathbb{P}\left(\sup_{0\leq u\leq 1/2} (a + X_u) \prod_{i=1}^{n-1} Z_{1-u}^{(i)} r^{\frac{n}{\alpha}} \leq \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \geq 0, 1 \geq \xi_{\frac{1}{r}}\right) \leq \mathbb{P}\left(\sup_{0\leq u\leq 1/2} \left(a \prod_{i=1}^{n-1} Z_{1-u}^{(i)} + X_u \prod_{i=1}^{n-1} |Z_{1-u}^{(i)}|\right) r^{\frac{n}{\alpha}} \leq \varepsilon, \prod_{i=1}^{n-1} Z_1^{(i)} \geq 0, 1 \geq \xi_{\frac{1}{r}}\right).
$$

Observe next that, by taking  $u = 0$ , this quantity is null as soon as  $a \prod_{i=1}^{n-1} Z_1^{(i)}$  $n_1^{(i)} r^{\frac{n}{\alpha}} \geq \varepsilon$ . Therefore, denoting  $\theta_{\frac{1}{2}} = \operatorname{Argmax}_{0 \leq u \leq 1/2}$  $0 \le u \le 1/2$  $X_u$ , we may replace the supremum by its value at  $\theta_{\frac{1}{2}}$  to get the bound

<span id="page-6-0"></span>
$$
\mathbb{P}\left(\left(a\prod_{i=1}^{n-1} Z_{1-\theta_{\frac{1}{2}}}^{(i)} + X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1} |Z_{1-\theta_{\frac{1}{2}}}^{(i)}|\right)r^{\frac{n}{\alpha}} \leq \varepsilon, \ \varepsilon \geq a\ r^{\frac{n}{\alpha}}\prod_{i=1}^{n-1} Z_{1}^{(i)} \geq 0, \ 1 \geq \xi_{\frac{1}{r}}\right).
$$
 (7)

We further decompose this integral according to the sign of  $\prod_{i=1}^{n-1} Z_{1-i}^{(i)}$  $1-\theta_{\frac{1}{2}}$ .

*i)* When 
$$
\prod_{i=1}^{n-1} Z_{1-\theta_{\frac{1}{2}}}^{(i)} \ge 0
$$
, the expression (7) is smaller than  
\n
$$
\mathbb{P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1} |Z_{1-\theta_{\frac{1}{2}}}^{(i)}|r^{\frac{n}{\alpha}} \le \varepsilon, 1 \ge \xi_{\frac{1}{r}}\right) =: I_n(r, \varepsilon).
$$
\n(8)

*ii*) When  $\prod_{i=1}^{n-1} Z_{1-i}^{(i)}$  $\frac{1-\theta_1}{1-\theta_1} \leq 0$ , the situation is slightly more complex. We have

$$
\mathbb{P}\left((X_{\theta_{\frac{1}{2}}}-a)\prod_{i=1}^{n-1}|Z_{1-\theta_{\frac{1}{2}}}^{(i)}|r^{\frac{n}{\alpha}} \leq \varepsilon, \ \varepsilon \geq a r^{\frac{n}{\alpha}} \prod_{i=1}^{n-1}Z_{1}^{(i)} \geq 0, \ 1 \geq \xi_{\frac{1}{r}}\right)
$$
\n
$$
\leq \mathbb{P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1}|Z_{1-\theta_{\frac{1}{2}}}^{(i)}|r^{\frac{n}{\alpha}} \leq \varepsilon \left(1 + \frac{\prod_{i=1}^{n-1}|Z_{1-\theta_{\frac{1}{2}}}^{(i)}}{\prod_{i=1}^{n-1}Z_{1}^{(i)}}\right), \prod_{i=1}^{n-1}Z_{1}^{(i)} \geq 0, \ 1 \geq \xi_{\frac{1}{r}}\right)
$$
\n
$$
\leq I_{n}(r, 2\varepsilon) + \mathbb{P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1}|Z_{1}^{(i)}|r^{\frac{n}{\alpha}} \leq 2\varepsilon, \ 1 \geq \xi_{\frac{1}{r}}\right) =: I_{n}(r, 2\varepsilon) + J_{n}(r, 2\varepsilon)
$$

where we have used in the last line the inequality :  $1_{\{x \le \varepsilon(a+b)\}} \le 1_{\{x \le 2a\varepsilon\}} + 1_{\{x \le 2b\varepsilon\}}$ . Going back to [\(6\)](#page-5-1), we are thus led to study the asymptotics of

$$
\int_0^1 (I_n(r,\varepsilon) + J_n(r,\varepsilon)) \mathbb{P}(g_1 \in dr).
$$

We start with  $I_n(r, \varepsilon)$  which will give the main contribution. From Lemma [6,](#page-3-1) we may choose  $\delta \in (0,1)$  small enough such that

<span id="page-7-2"></span>
$$
\forall r \le \delta, \qquad \mathbb{P}(g_1 \in dr) / dr \le \frac{c}{\sqrt{r}} \tag{9}
$$

for some constant  $c > 0$ . On the one hand, when  $r \ge \delta$ , we obtain since  $\theta_{\frac{1}{2}} \le \frac{1}{2}$  $\frac{1}{2}$ :

$$
\int_{\delta}^{1} I_n(r,\varepsilon) \mathbb{P}\left(g_1 \in dr\right) \leq \mathbb{P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1}|Z_1^{(i)}|\delta^{n/\alpha} \leq 2^{\frac{n-1}{\alpha}}\varepsilon\right) \underset{\varepsilon \to +\infty}{\sim} \begin{cases} \kappa \varepsilon^{\frac{\alpha}{2}} & \text{if } \alpha \in (0,2), \\ \kappa \varepsilon \left|\ln(\varepsilon)\right|^{n-1} & \text{if } \alpha = 2. \end{cases}
$$

On the other hand, when  $r \leq \delta$ , we deduce from the Markov property at time 1 that :

$$
I_n(r,\varepsilon) \leq \mathbb{E}\left[\prod_{i=1}^{n-1} \widehat{\mathbb{P}}_1^{(i)} \left(|Z_1^{(i)}|^\alpha \widehat{T}_0^{(i)} \geq \frac{1-\delta}{r}\right), X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1} |Z_{1-\theta_{\frac{1}{2}}}^{(i)}| r^{\frac{n}{\alpha}} \leq \varepsilon\right].
$$

Using the identity  $Z_1^{(i)}$ 1  $\stackrel{\text{(law)}}{=} Z_{1-}^{(i)}$  $1-\theta_{\frac{1}{2}}$ +  $\theta_{\frac{1}{2}}^{\frac{1}{\alpha}} Y_1^{(i)}$  where  $Y_1^{(i)}$  $Z_1^{(i)}$  is a copy of  $Z_1^{(i)}$  $1^{(i)}$ , independent from the processes  $(Z^{(i)})$  and  $(\widetilde{Z}^{(i)})$ , we then obtain the bound

<span id="page-7-0"></span>
$$
I_n(r,\varepsilon) \le \mathbb{E}\left[\prod_{i=1}^{n-1} \widehat{\mathbb{P}}_1^{(i)} \left( \left( |Z_{1-\theta_{\frac{1}{2}}}^{(i)}|^{\alpha} + |Y_1^{(i)}|^{\alpha} \right) \widehat{T}_0^{(i)} \ge \frac{1-\delta}{2r} \right), X_{\theta_{\frac{1}{2}}} \prod_{i=1}^{n-1} |Z_{1-\theta_{\frac{1}{2}}}^{(i)}|^{\frac{n}{\alpha}} \le 1 \right] \tag{10}
$$

where we have used the classic inequality  $|x+y|^{\alpha} \leq 2(|x|^{\alpha} + |y|^{\alpha})$  since  $\alpha \in (0,2]$ . We further assume that  $\delta$  is taken small enough so that, from Lemma [4](#page-2-0) and the asymptotics [\(3\)](#page-3-0), we have

<span id="page-7-1"></span>
$$
\forall r \leq \delta, \qquad \mathbb{E}\left[\widehat{\mathbb{P}}_1^{(i)}\left(\left(|Z_{1-\theta_{\frac{1}{2}}}^{(i)}|^{\alpha} \vee 1 + |Y_1^{(i)}|^{\alpha}\right)\widehat{T}_0^{(i)} \geq \frac{1-\delta}{2r}\right)\right] \leq \kappa\sqrt{r} \tag{11}
$$

for some positive constant  $\kappa$ , and where  $a \vee b = \max(a, b)$ . We shall now proceed by iteration.

*i*) If  $|Z_{1-\theta_{1}}^{(n-1)}|$  $\left| \frac{1}{1-\theta_1} \right| \geq 1$ , then, we may remove  $|Z_{1-\theta_1}^{(n-1)}|$  $\left( \frac{n-1}{1-\theta} \right)$  from the second product in [\(10\)](#page-7-0), and deduce from [\(11\)](#page-7-1) that  $I_n(r,\varepsilon)$  is smaller than

$$
\kappa \sqrt{r} \mathbb{E} \left[ \prod_{i=1}^{n-2} \widehat{\mathbb{P}}^{(i)}_1 \left( \left( |Z^{(i)}_{1-\theta_{\frac{1}{2}}} |^{\alpha} + |Y^{(i)}_1 |^{\alpha}\right) \widehat{T}^{(i)}_0 \geq \frac{1-\delta}{2r} \right), X_{\theta_{\frac{1}{2}}} \prod_{i=1}^{n-2} |Z^{(i)}_{1-\theta_{\frac{1}{2}}} | r^{\frac{n}{\alpha}} \leq 1 \right].
$$

*ii*) If  $|Z_{1-\theta_{1}}^{(n-1)}|$  $\left| \frac{(n-1)}{1-\theta_1} \right| \leq 1$ , then, we may replace  $|Z_{1-\theta_1}^{(n-1)}|$  $\frac{1-(n-1)}{1-\theta}$  by 1 in the first product in [\(10\)](#page-7-0), and deduce, still from [\(11\)](#page-7-1), that  $I_n(r,\varepsilon)$  is smaller than

$$
\kappa\sqrt{r}\,\mathbb{E}\left[\prod_{i=1}^{n-2}\widehat{\mathbb{P}}_1^{(i)}\left(\left(|Z_{1-\theta_{\frac{1}{2}}}^{(i)}|^\alpha+|Y_1^{(i)}|^\alpha\right)\widehat{T}_0^{(i)}\geq \frac{1-\delta}{2r}\right),\,X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1}|Z_{1-\theta_{\frac{1}{2}}}^{(i)}|r^\frac{n}{\alpha}\leq 1\right].
$$

Iterating the procedure, we obtain that  $I_n(r, \varepsilon)$  may be bounded by a sum of  $2^{n-1}$  terms :

$$
I_n(r,\varepsilon)\leq \kappa\,r^{\frac{n-1}{2}}\sum_{\Delta\subset \{1,\dots,n-1\}}{\mathbb P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i\in\Delta}|Z_{1-\theta_{\frac{1}{2}}}^{(i)}|r^{\frac{n}{\alpha}}\leq 1\right)
$$

where the sum is taken over all the subsets of  $\{1, 2, \ldots, n-1\}$  (including the empty set). The change of variable  $\varepsilon x = r^{\frac{n}{\alpha}}$  and the estimate [\(9\)](#page-7-2) yield then the upper bound

<span id="page-8-0"></span>
$$
\int_0^\delta I_n(r,\varepsilon) \mathbb{P}(g_1 \in dr) \le \kappa \varepsilon^{\frac{\alpha}{2}} \sum_{\Delta \subset \{1,\dots,n-1\}} \int_0^{\frac{\delta^{n/\alpha}}{\varepsilon}} \mathbb{P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i\in\Delta} |Z_1^{(i)}|x \le 2^{\frac{n-1}{\alpha}}\right) x^{\frac{\alpha}{2}-1} dx\tag{12}
$$

and it remains to study the asymptotics of the integrands. From Lemma [4](#page-2-0) and [\(1\)](#page-1-1), we deduce that  $\rightarrow$  when  $\alpha \in (0, 2)$  all the terms have the same contribution:

$$
\mathbb{P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i\in\Delta}|L_1^{(i)}|x\leq 2^{\frac{n-1}{\alpha}}\right)\underset{x\to+\infty}{\sim}c\left(\frac{1}{x}\right)^{\alpha/2}
$$

 $\rightarrow$  while, for  $\alpha = 2$ , they depend on the cardinal of  $\Delta$ :

$$
\mathbb{P}\left(X_{\theta_{\frac{1}{2}}}\prod_{i\in\Delta}|W_1^{(i)}|x\leq 2^{\frac{n-1}{2}}\right)\underset{x\to+\infty}{\sim}c\,\frac{(\ln(x))^{|\Delta|}}{x}.
$$

Plugging these expressions in [\(12\)](#page-8-0) finally gives the announced upper bound.

The study of the asymptotics of  $J_n(r,\varepsilon)$  follows the same pattern of proof, except that we do not need to introduce the random variables  $(Y_1^{(i)}$  $T_1^{(i)}$ ). Indeed, when  $r \geq \delta$ , we get the same asymptotics bound while for  $r \leq \delta$  we obtain, applying the Markov property :

<span id="page-8-1"></span>
$$
J_n(r,\varepsilon) \leq \mathbb{E}\left[\prod_{i=1}^{n-1} \widehat{\mathbb{P}}_1^{(i)} \left(|Z_1^{(i)}|^\alpha \widehat{T}_0^{(i)} \geq \frac{1-\delta}{r}\right), X_{\theta_{\frac{1}{2}}}\prod_{i=1}^{n-1} |Z_1^{(i)}| r^{\frac{n}{\alpha}} \leq 1\right].
$$

Using the decompositions  $|Z_1^{(i)}|$  $|Z_1^{(i)}| \le 1$  (resp.  $|Z_1^{(i)}|$  $|I_1^{(i)}| \geq 1$  and following the same steps as for  $I_n(r, \varepsilon)$ , we deduce that

$$
\int_0^\delta J_n(r,\varepsilon) \mathbb{P}(g_1 \in dr)
$$
\n
$$
\leq \kappa \varepsilon^{\frac{\alpha}{2}} \sum_{\Delta \subset \{1,\dots,n-1\}} \int_0^{\frac{\delta^{n/\alpha}}{\varepsilon}} \mathbb{E} \left[ \prod_{i \in \Delta} |Z_1^{(i)}|^{\frac{\alpha}{2}}, X_{\theta_{\frac{1}{2}}} \prod_{i \in \Delta} |Z_1^{(i)}| x \leq 2^{\frac{n-1}{\alpha}} \right] x^{\frac{\alpha}{2}-1} dx \tag{13}
$$

and, as  $\varepsilon \to 0$ , all the terms on the right-hand side have the same asymptotics :  $\varepsilon^{\frac{\alpha}{2}} |\ln(\varepsilon)|$ .

3.2.2. The case  $\{a \ge 1\}$ . Starting back from [\(5\)](#page-5-2), we first bound the supremum by its value at  $u = 0$ :

$$
\int_{1}^{+\infty} \int_{0}^{1} \mathbb{P}\left(\sup_{0 \le u \le 1} X_{u}^{(a,1,0)} \prod_{i=1}^{n-1} Z_{1-u}^{(i)} r^{\frac{n}{\alpha}} \le \varepsilon, \prod_{i=1}^{n-1} Z_{1}^{(i)} \ge 0, 1 \ge \xi_{\frac{1}{r}}\right) \rho(da, dr)
$$
  

$$
\le \int_{1}^{+\infty} \int_{0}^{1} \mathbb{P}\left(a \prod_{i=1}^{n-1} Z_{1}^{(i)} r^{\frac{n}{\alpha}} \le \varepsilon, \prod_{i=1}^{n-1} Z_{1}^{(i)} \ge 0, 1 \ge \xi_{\frac{1}{r}}\right) \rho(da, dr)
$$
  

$$
\le \int_{0}^{1} \mathbb{P}\left(\prod_{i=1}^{n-1} |Z_{1}^{(i)}| r^{\frac{n}{\alpha}} \le \varepsilon, 1 \ge \xi_{\frac{1}{r}}\right) \mathbb{P}(g_{1} \in dr).
$$

The study of this last expression will be similar to that of  $J_n(r, \varepsilon)$ , replacing  $X_{\theta_1}$  by 1. Indeed, on the one hand, we first deduce from Lemma [4,](#page-2-0) taking  $\delta$  small enough as before, that :

$$
\int_{\delta}^1 \mathbb{P}\left(\prod_{i=1}^{n-1} |Z_1^{(i)}| r^{\frac{n}{\alpha}} \leq \varepsilon, \ 1 \geq \xi_{\frac{1}{r}}\right) \mathbb{P}\left(g_1 \in dr\right) \leq \mathbb{P}\left(\prod_{i=1}^{n-1} |Z_1^{(i)}| \delta^{\frac{n}{\alpha}} \leq \varepsilon\right) \underset{\varepsilon \to 0}{\sim} \kappa \varepsilon \left|\ln(\varepsilon)\right|^{n-2}.
$$

On the other hand, for  $r \leq \delta$ , we deduce, as for [\(13\)](#page-8-1), that

$$
\int_0^{\delta} \mathbb{P}\left(\prod_{i=1}^{n-1} |Z_1^{(i)}| r^{\frac{n}{\alpha}} \leq \varepsilon, 1 \geq \xi_{\frac{1}{r}}\right) \mathbb{P}\left(g_1 \in dr\right)
$$
  

$$
\leq \kappa \varepsilon^{\frac{\alpha}{2}} \sum_{\Delta \subset \{1, \ldots, n-1\}} \int_0^{\frac{\delta^{n/\alpha}}{\varepsilon}} \mathbb{E}\left[\prod_{i \in \Delta} |Z_1^{(i)}|^{\frac{\alpha}{2}}, \prod_{i \in \Delta} |Z_1^{(i)}| x \leq 2^{\frac{n-1}{\alpha}}\right] x^{\frac{\alpha}{2}-1} dx.
$$

When  $\varepsilon \to 0$ , all the integrals on the right-hand side are finite, hence we obtain the asymptotics  $\varepsilon \frac{\alpha}{2}$ which is negligible.  $\Box$ 

# 4. Proof of Lemma [7](#page-5-0)

*Proof.* This lemma being classic for Brownian motion, we assume that  $\alpha \in (0, 2)$ . Let  $0 < s \le t \le 1$ and take F a positive functional. Let us denote by  $f(y; z, r)$  the probability density function of  $(X_{T_0}, T_0)$  when  $X_0 = y$ . By symmetry and time reversal, we first have

$$
\mathbb{E}\left[F\left(\frac{X_{u^-}}{g_1^{1/\alpha}}, s \le u \le g_1\right)1_{\{g_1 \ge t\}}\right] = 2\int_0^{+\infty} \mathbb{E}^{(y,1,0)}\left[F\left(\frac{X_u}{(1-T_0)^{1/\alpha}}, T_0 \le u \le 1-s\right)1_{\{T_0 \le 1-t\}}\right]p_1(y)dy.
$$

The absolute continuity formula for the stable bridge as well as the Markov property then yield

$$
2\int_0^{+\infty} \mathbb{E}_y \left[ p_s(X_{1-s}) F\left(\frac{X_u}{(1-T_0)^{1/\alpha}}, T_0 \le u \le 1-s \right) 1_{\{T_0 \le 1-t\}} \right] dy
$$
  
= 
$$
2\int_0^{+\infty} \int_0^0 \int_{-\infty}^{1-t} \mathbb{E}_z \left[ p_s(X_{1-s-r}) F\left(\frac{X_u}{(1-r)^{1/\alpha}}, 0 \le u \le 1-s-r \right) \right] f(y; z, r) dy dz dr
$$

Next, by scaling and using that  $t^{1/\alpha} p_t(z) = p_1 \left( \frac{z}{t^{1/\alpha}} \right)$  $rac{z}{t^{1/\alpha}}\bigg),$ 

$$
2\int_{0}^{+\infty} \int_{-\infty}^{0} \int_{-\infty}^{1-t} \mathbb{E}_{\frac{z}{(1-r)^{1/\alpha}}} \left[ p_s((1-r)^{1/\alpha} X_{1-\frac{s}{1-r}}) F\left(X_{\frac{u}{1-r}}, 0 \le u \le 1-s-r\right) \right] f(y; z, r) dy dz dr
$$
  
\n
$$
= 2\int_{0}^{+\infty} \int_{-\infty}^{0} \int_{0}^{1-t} \mathbb{E}_{a} \left[ p_{\frac{s}{1-r}}(X_{1-\frac{s}{1-r}}) F\left(X_{u}, 0 \le u \le 1-\frac{s}{1-r}\right) \right] f(y; a(1-r)^{1/\alpha}, r) dy dadr
$$
  
\n
$$
= 2\int_{0}^{+\infty} \int_{-\infty}^{0} \int_{0}^{1-t} \mathbb{E}^{(a,1,0)} \left[ F\left(X_{u}, 0 \le u \le 1-\frac{s}{1-r}\right) \right] p_1(a) f(y; a(1-r)^{1/\alpha}, r) dy dadr
$$
  
\n
$$
= 2\int_{-\infty}^{0} da \mathbb{E}^{(0,1,a)} \left[ F\left(X_{u-}, \frac{s}{1-r} \le u \le 1\right) \right] p_1(a) \int_{0}^{+\infty} \int_{0}^{1-t} f(y; a(1-r)^{1/\alpha}, r) dy dr
$$

where in the second line we have used the change of variable  $z = a(1 - r)^{1/\alpha}$ . Letting  $s \to 0$ , we finally deduce that

$$
\mathbb{E}\left[F\left(\frac{X_{u^{-}}}{g_{1}^{1/\alpha}},0\leq u\leq g_{1}\right)1_{\{g_{1}\geq t\}}\right]
$$
  
=  $2\int_{-\infty}^{0}da\,\mathbb{E}^{(0,1,a)}\left[F\left(X_{u^{-}},0\leq u\leq 1\right)\right]p_{1}(a)\int_{0}^{+\infty}\int_{0}^{1-t}f(y;a(1-r)^{1/\alpha},r)dydr$ 

which proves Lemma [7.](#page-5-0)

 $\Box$ 

## **REFERENCES**

- <span id="page-10-3"></span><span id="page-10-1"></span>[1] F. Aurzada and T. Simon. Persistence probabilities and exponents. In : Lévy Matters V, 183-224, Springer, 2015.
- <span id="page-10-5"></span>[2] R. Bañuelos and K. Bogdan. Symmetric stable processes in cones. Potential Anal. 21 (3), 263-288, 2004.
- [3] R. Bañuelos and K. Bogdan. Symmetric stable processes in parabola-shaped regions. Proc. Amer. Math. Soc. 133 (12), 3581-3587, 2005.
- <span id="page-10-2"></span><span id="page-10-0"></span>[4] J. Bertoin. *Lévy processes*. Cambridge University Press, Cambridge, 1996.
- [5] A. J. Bray, S. N. Majumdar and G. Schehr. Persistence and first-passage properties in non-equilibrium systems. Adv. Physics 62 (3), 225-361, 2013.
- <span id="page-10-9"></span>[6] R.A. Doney and M. S. Savov. The asymptotic behavior of densities related to the supremum of a stable process. Ann. Probab. 38 (1), 316-326, 2010.
- <span id="page-10-7"></span><span id="page-10-6"></span>[7] S. Janson. Moments of Gamma type and the Brownian supremum process area. Prob. Surveys 7, 1-52, 2010.
- [8] P.J. Méndez-Hernández. Exit times of symmetric  $\alpha$ -stable processes from unbounded convex domains. Elect. J. Probab. 12, 100-121, 2007.
- <span id="page-10-4"></span>[9] P.J. Méndez-Hernández. Exit times from cones in  $\mathbb{R}^n$  of symmetric stable processes. Illinois J. Math. 46 (1), 155-163, 2002.
- <span id="page-10-8"></span>[10] C. Profeta and T. Simon. Windings of the stable Kolmogorov process. ALEA XII, 115-127, 2015.

LABORATOIRE DE MATHÉMATIQUES ET MODÉLISATION D'ÉVRY, UNIVERSITÉ D'EVRY-VAL D'ESSONNE, BÂTIMENT IBGBI, 23 BOULEVARD DE FRANCE, 91037 EVRY CEDEX, FRANCE. Email: christophe.profeta@univ-evry.fr