# KERNEL AND WAVELET DENSITY ESTIMATORS ON MANIFOLDS AND MORE GENERAL METRIC SPACES

G. CLEANTHOUS, A. G. GEORGIADIS, G. KERKYACHARIAN, P. PETRUSHEV, AND D. PICARD

ABSTRACT. We consider the problem of estimating the density of observations taking values in classical or nonclassical spaces such as manifolds and more general metric spaces. Our setting is quite general but also sufficiently rich in allowing the development of smooth functional calculus with well localized spectral kernels, Besov regularity spaces, and wavelet type systems. Kernel and both linear and nonlinear wavelet density estimators are introduced and studied. Convergence rates for these estimators are established, which are analogous to the existing results in the classical setting of real-valued variables.

## 1. INTRODUCTION

A great deal of efforts is nowadays invested in solving statistical problems, where the data are located in quite complex domains such as matrix spaces or surfaces (manifolds). A seminal example in this direction is the case of spherical data. Developments in this domain have been motivated by a number of important applications. We only mention here some of the statistical challenges posed by astrophysical data: denoising of signals, testing stationarity, rotation invariance or gaussianity of signals, investigating the fundamental properties of the cosmic microwave background (CMB), impainting of the CMB in zones on the sphere obstructed by other radiations, producing cosmological maps, exploring clusters of galaxies or point sources, investigating the true nature of ultra high energy cosmic rays (UHECR). We refer the reader to the overview by Starck, Murtagh, and Fadili [27] of the use of various wavelet tools in this domain as well as the work of some of the authors in this direction [1] and [22].

Dealing with complex data requires the development of more sophisticated tools and statistical methods than the existing tools and methods. In particular, these tools should capture the natural topology and geometry of the application domain.

Our contribution will be essentially theoretical, however, our statements will be illustrated by examples issued from different fields of applications.

Our purpose in this article is to study the *density estimation problem*, namely, one observes  $X_1, \ldots, X_n$  that are i.i.d. random variables defined on a space  $\mathcal{M}$  and the problem is to find a good estimation to the common density function.

Date: February 4, 2019.

<sup>2010</sup> Mathematics Subject Classification. Primary 62G07, 58J35; Secondary 43A85, 42B35. Key words and phrases. kernel density estimators, wavelet density estimators, adaptive density

estimators, non-parametric estimators, Ahlfors regularity, heat kernel, Besov spaces.

We thank the anonymous referees for a very thorough reading of an earlier version which enabled us to substantially improve the paper.

Corresponding author: Galatia Cleanthous, E-mail: galatia.cleanthous@newcastle.ac.uk.

### 2 G. CLEANTHOUS, A. GEORGIADIS, G. KERKYACHARIAN, P. PETRUSHEV, D. PICARD

This problem has a long history in mathematical statistics especially when the set  $\mathcal{M}$  is  $\mathbb{R}^d$  or a cube in  $\mathbb{R}^d$  (see e.g. the monograph [29] and the references herein). Here we will consider very general spaces  $\mathcal{M}$  such as Riemannian manifolds or spaces of matrices or graphs and prove that with some assumptions, we can build an estimation theory with estimation procedures, regularity sets and upper bounds evaluations quite parallel to what has been neatly done in  $\mathbb{R}^d$ . In particular we intend to develop kernel methods with upper bounds and oracle properties as well as wavelet thresholding estimators with adaptative behavior.

If we want to roughly summarize the basic assumptions that will be made in this work, let us mention that some of them are concerning the basic dimensional structure of the set (doubling conditions), whereas others are devoted to construct an environment where regularity spaces can be defined as well as kernels or wavelets can be constructed.

This setting is quite general but at the same time is sufficiently rich in allowing the development of smooth functional calculus with well localized spectral kernels, Besov regularity spaces, and wavelet type systems. Naturally, the classical setting on  $\mathbb{R}^d$  and the one on the sphere are contained in this general framework, but also various other settings are covered. In particular, spaces of matrices, Riemannian manifolds, convex subsets of (non-compact) Riemannian manifolds are covered.

As will be shown in this general setting, a regularity scale and a general nonparametric density estimation theory can be developed in full generality just as in the standard case of  $[0,1]^d$  or  $\mathbb{R}^d$ . This undertaking requires the development of new techniques and methods that break new ground in the density estimation problem. Our *main contributions* are as follows:

(a) In a general setting described below, we introduce kernel density estimators sufficiently concentrated to establish oracle inequalities and  $\mathbb{L}^{p}$ -error rates of convergence for probability density functions lying in *Besov spaces*.

(b) We also develop linear wavelet density estimators and obtain  $\mathbb{L}^{p}$ -error estimates for probability density functions in general *Besov* smoothness spaces.

(c) We establish  $\mathbb{L}^{p}$ -error estimates on *nonlinear wavelet density estimators with* hard thresholding in our general geometric setting. We obtain such estimates for probability density functions in general Besov spaces.

To put the results from this article in perspective we next compare them with the results in [2]. The geometric settings in both articles are comparable and the two papers study adaptive methods. In [2] different standard statistical models (regression, white noise model, density estimation) are considered in a Bayesian framework. The methods are different (because we do not consider here Bayesian estimators) and the results are also different (since, again, we are not interested here in a concentration result of the posterior distribution). It is noteworthy that the results in the so called *dense case* exhibit the same rates of convergence. It is also important to observe the wide adaptation properties of the thresholding estimates here which allow to obtain minimax rates of convergence in the so called *sparse case*, which was not possible in [2].

The organization of this article is as follows: In Section 2, we describe our general setting of a doubling measure metric space in the presence of a self-adjoint operator whose heat kernel has Gaussian localization and the Markov property. We provide motivation and inspiration for our developments and we present some first examples, both elementary and more involved. In Section 3, we review some basic

facts related to our setting such as smooth functional calculus, the construction of wavelet frames, Besov spaces, and other background. This section can be read quickly by a reader more motivated by the introduction of estimation procedures. We develop kernel density estimators in Section 4 and establish  $\mathbb{L}^p$ -error estimates for probability density functions in general Besov spaces. We also introduce and study linear wavelet density estimators. In Section 5, we introduce and study adaptive wavelet threshold density estimators. We establish  $\mathbb{L}^p$ -error estimates for probability density functions in Besov spaces. Section 7 is an appendix, where we place the proofs of some claims from previous sections.

Notation: Throughout  $\mathbb{1}_E$  will denote the indicator function of the set E and  $\|\cdot\|_p := \|\cdot\|_{\mathbb{L}^p(\mathcal{M},\mu)}$ . We denote by c, c' positive constants that may vary at every occurrence. Most of these constants will depend on some parameters that may be indicated in parentheses. We will also denote by  $c_0, c_1, \ldots$  as well as  $c_\star, c_\diamond$  constants that will remain unchanged throughout. The relation  $a \sim b$  means that there exists a constant c > 1 such that  $c^{-1}a \leq b \leq ca$ . We will also use the notation  $a \wedge b := \min\{a, b\}, a \vee b := \max\{a, b\}$  and  $C^k(\mathbb{R}_+), k \in \mathbb{N} \cup \{\infty\}$ , will stand for the set of all functions with continuous derivatives of order up to k on  $\mathbb{R}_+ := [0, \infty)$ .

## 2. Setting and motivation

We assume that  $(\mathcal{M}, \rho, \mu)$  is a metric measure space equipped with a distance  $\rho$  and a positive Radon measure  $\mu$ .

Let  $X_1, \ldots, X_n$  be independent identically distributed (i.i.d.) random variables on  $\mathcal{M}$  with common probability having a density function (pdf) f with respect to the measure  $\mu$ . Our purpose is to estimate the density f. To an estimator  $\hat{f}_n$  of f, we associate its risk:

$$R_n(\hat{f}, f, p) = \mathbb{E}_f \left( \int_{\mathcal{M}} |\hat{f}_n(x) - f(x)|^p \mu(dx) \right)^{\frac{1}{p}} = \mathbb{E}_f \|\hat{f}_n - f\|_p, \quad 1 \le p < \infty$$

as well as its  $\mathbb{L}_{\infty}$  risk:

$$R_n(\hat{f}, f, \infty) = \mathbb{E}_f\left(\operatorname{ess\,sup}_{x \in \mathcal{M}} |\hat{f}_n(x) - f(x)|\right) = \mathbb{E}_f ||\hat{f}_n - f||_{\infty}.$$

We will operate in the following setting. Most of the material can be found in an extended form in the papers [4, 17]. Note that, depending on the results we are going to establish, some of the following conditions will be assumed, others will not.

2.1. Doubling and non-collapsing conditions. The following conditions are concerning properties related to 'dimensional' structure of  $\mathcal{M}$ .

**C1.** We assume that the metric space  $(\mathcal{M}, \rho, \mu)$  satisfies the so called *doubling* volume condition:

(2.1) 
$$\mu(B(x,2r)) \le c_0 \mu(B(x,r)) \quad \text{for } x \in \mathcal{M} \text{ and } r > 0,$$

where  $B(x,r) := \{y \in \mathcal{M} : \rho(x,y) < r\}$  and  $c_0 > 1$  is a constant. The above implies that there exist constants  $c'_0 \ge 1$  and d > 0 such that

(2.2) 
$$\mu(B(x,\lambda r)) \le c'_0 \lambda^d \mu(B(x,r)) \quad \text{for } x \in \mathcal{M}, \, r > 0, \text{ and } \lambda > 1,$$

The least d such that (2.2) holds is the so called homogeneous dimension of  $(\mathcal{M}, \rho, \mu)$ From now on we will use the notation  $|E| := \mu(E)$  for  $E \subset \mathcal{M}$ .

#### 4 G. CLEANTHOUS, A. GEORGIADIS, G. KERKYACHARIAN, P. PETRUSHEV, D. PICARD

In developing adaptive density estimators in Section 5 we will additionally assume that  $(\mathcal{M}, \rho, \mu)$  is a compact measure space with  $\mu(\mathcal{M}) < \infty$  satisfying the following condition:

**C1A.** Ahlfors regular volume condition: There exist constants  $c_1, c_2 > 0$  and d > 0 such that

(2.3) 
$$c_1 r^d \le |B(x,r)| \le c_2 r^d \text{ for } x \in \mathcal{M} \text{ and } 0 < r \le \text{diam}(\mathcal{M}).$$

Clearly, condition C1A implies conditions C1 and the following condition C2 as well, with d from (2.3) being the homogeneous dimension of  $(\mathcal{M}, \rho, \mu)$ .

These doubling conditions have been introduced in Harmonic Analysis in the 70's by R. Coifman and G. Weiss [3].

It is interesting already to notice that d will indeed play the role of a dimension in the statistical results as well. Condition **C1A** is obviously true for  $\mathcal{M} = \mathbb{R}^d$  with  $\mu$  the Lebesgue measure.

Also, the doubling condition is precisely related to the *metric entropy* using the following lemma whose elementary proof can be found for instance in [2, Proposition 1]. For  $\epsilon > 0$ , we define, as usual, the covering number  $N(\epsilon, \mathcal{M})$  as the smallest number of balls of radius  $\epsilon$  covering  $\mathcal{M}$ .

**Lemma 2.1.** Under the condition C1A and if  $\mathcal{M}$  is compact, there exist constants c' > 0, c'' > 0 and  $\epsilon_0 > 0$  such that

(2.4) 
$$\frac{1}{c'} \left(\frac{1}{\epsilon}\right)^d \le N(\epsilon, \mathcal{M}) \le \frac{2^d}{c''} \left(\frac{1}{\epsilon}\right)^d,$$

for all  $0 < \epsilon \leq \epsilon_0$ .

**C2.** Non-collapsing condition: There exists a constant  $c_3 > 0$  such that

(2.5) 
$$\inf_{x \in \mathcal{M}} |B(x,1)| \ge c_3 > 0$$

This condition is not necessarily very restrictive. For instance, it is satisfied if  $\mathcal{M}$  is compact. It is satisfied for  $\mathbb{R}^d$  if  $\mu$  is the Lebesgue measure, but untrue for  $\mathbb{R}$  if  $\mu$  is a Gaussian measure.

2.2. **Smooth operator.** Here comes an important assumption which may seem strange to the reader at first glance. Before entering into the specificity of the set of assumptions described below, let us explain some motivations.

One rather standard method in density estimation is the kernel estimation method i.e. considering a family of functions indexed by  $\delta > 0$ :  $K_{\delta} : \mathcal{M} \times \mathcal{M} \longrightarrow \mathbb{R}$  an associated kernel density estimator is defined by

(2.6) 
$$\widehat{K}_{\delta}(x) := \frac{1}{n} \sum_{i=1}^{n} K_{\delta}(X_i, x), \quad x \in \mathcal{M}.$$

In  $\mathbb{R}^d$ , an important family is the family of translation kernels  $K_{\delta}(x, y) = [\frac{1}{\delta}]^d G(\frac{x-y}{\delta})$ , where G is a function  $\mathbb{R}^d \longrightarrow \mathbb{R}$ . When  $\mathcal{M}$  is a more involved set such as a manifold or a set of graphs, of matrices, the simple operations of translation and dilation may not be meaningful. Hence, even finding a family of kernels to start with might be a difficulty. It will be shown in Section 4 that the following assumptions provide quite 'naturally' a family of kernels.

When dealing with a kernel estimation method, it is standard to consider two quantities :

$$b_{\delta}(f) := \|\mathbb{E}_f \widehat{K}_{\delta} - f\|_p, \quad \|\xi_f\|_p := \|\widehat{K}_{\delta} - \mathbb{E}_f \widehat{K}_{\delta}\|_p.$$

The analysis of the second term (stochastic term)  $\|\xi_f\|_p$ , can be reduced via Rosenthal inequalities to proper bounds on norms of  $K_{\delta}(\cdot, \cdot)$  and f (see the Lemmas 4.7, 4.8), where in particular the assumptions of the previous subsection are also important).

The analysis of the first term  $b_{\delta}(f)$  is linked to the approximation properties of the family  $\mathbb{E}_f \hat{K}_{\delta}$ . One can stop at this level and precisely express the performance of an estimator in terms of  $\|\mathbb{E}_f \hat{K}_{\delta} - f\|_p$ . This is the purpose of oracle inequalities (see Theorem 4.3).

However, it might seem more convincing if one can relate the rate of approximation of the family  $\|\mathbb{E}_f \hat{K}_{\delta} - f\|_p$  to regularity properties of the function f. It is standardly proved (see e.g. [14]), that in  $\mathbb{R}^d$  if K is a translation family with mild properties on K, then polynomial rates of approximation are obtained for functions with Besov regularity.

Hence, an important issue becomes finding spaces of regularity associated to a possibly complex set  $\mathcal{M}$ . On a compact metric space  $(M, \rho)$  one can always define the scale of *s*-Lipschitz spaces defined by the following norm

(2.7) 
$$\|f\|_{Lip_s} := \|f\|_{\infty} + \sup_{x \neq y} \frac{|f(x) - f(y)|}{\rho(x, y)^s}, \quad 0 < s \le 1.$$

In Euclidian spaces a function can be much more regular than Lipschitz, for instance differentiable at different orders, or belong to some Sobolev or Besov spaces.

When  $\mathcal{M}$  is a set where there is no obvious notion of differentiability, one can make the observation that in  $\mathbb{R}^d$  or Riemannian manifolds, regularity properties can also be expressed via the associated Laplacian. The Laplacian itself is an operator of order 2, but its square root is of order 1 and can be interpreted as a substitute for derivation.

We will use this analogy to introduce an operator L playing the role of a Laplacian. However, conditions are needed to ensure that this analogy makes sense and can lead to a scale of spaces with suitable properties (which for instance, for small regularities correspond to Lipschitz spaces). This is why we adopt the setting introduced in [4, 17]. This setting is rich enough to develop a Littlewood-Paley theory in almost complete analogy with the classical case on  $\mathbb{R}^d$ , see [4, 17]. In particular, it allows to develop Besov spaces  $B_{pq}^s$  with all sets of indices. At the same time this framework is sufficiently general to cover a number of interesting cases as will be shown in what follows.

Our main assumption is that the space  $(\mathcal{M}, \rho, \mu)$  is complemented by an essentially self-adjoint non-negative operator L on  $\mathbb{L}^2(\mathcal{M}, \mu)$ , mapping real-valued to real-valued functions, such that the associated semigroup  $P_t = e^{-tL}$  consists of integral operators with the (heat) kernel  $p_t(x, y)$  obeying the following conditions: **C3.** Gaussian localization: There exist constants  $c_4, c_5 > 0$  such that

(2.8) 
$$|p_t(x,y)| \le \frac{c_4 \exp\left(-\frac{c_5 \rho^2(x,y)}{t}\right)}{\left(|B(x,\sqrt{t})||B(y,\sqrt{t})|\right)^{1/2}} \text{ for } x, y \in \mathcal{M}, t > 0.$$

## 6 G. CLEANTHOUS, A. GEORGIADIS, G. KERKYACHARIAN, P. PETRUSHEV, D. PICARD

C4. Hölder continuity: There exists a constant  $\alpha > 0$  such that

(2.9) 
$$|p_t(x,y) - p_t(x,y')| \le c_4 \left(\frac{\rho(y,y')}{\sqrt{t}}\right)^{\alpha} \frac{\exp\left(-\frac{c_5\rho^2(x,y)}{t}\right)}{\left(|B(x,\sqrt{t})||B(y,\sqrt{t})|\right)^{1/2}}$$

for  $x, y, y' \in \mathcal{M}$  and t > 0, whenever  $\rho(y, y') \le \sqrt{t}$ . C5. Markov property:

(2.10) 
$$\int_{\mathcal{M}} p_t(x, y) d\mu(y) = 1 \quad \text{for } x \in \mathcal{M} \text{ and } t > 0.$$

Above  $c_0, c_1, c_2, c_3, c_4, c_5, d, \alpha > 0$  are structural constants. These technical assumptions express that fact that the Heat kernel associated with the operator L 'behaves' as the standard Heat kernel of  $\mathbb{R}^d$ .

2.3. **Typical examples.** Here we present some examples of setups that are covered by the setting described above. We will use these examples in what follows to illustrate our theory. More involved examples will be given Section 6.

2.3.1. Classical case on  $\mathcal{M} = \mathbb{R}^d$ . Here  $d\mu$  is the Lebesgue measure and  $\rho$  is the Euclidean distance on  $\mathbb{R}^d$ . In this case we consider the operator

$$-L(f)(x) = \sum_{j=1}^{d} \partial_i^2 f(x) = \operatorname{div}(\nabla f)(x)$$

defined on the space  $\mathcal{D}(\mathbb{R}^d)$  of  $C^{\infty}$  functions with compact support. As is well known the operator L is positive essentially self-adjoint and has a unique extension to a positive self-adjoint operator. The associate semigroup  $e^{t\Delta}$  is given by the operator with the Gaussian kernel:  $p_t(x, y) = (4\pi t)^{-\frac{d}{2}} \exp\left(-\frac{|x-y|^2}{4t}\right)$ .

2.3.2. Periodic case on  $\mathcal{M} = [-1, 1]$ . Here  $d\mu$  is the Lebesgue measure and  $\rho$  is the Euclidean distance on the circle. The operator is L(f) = -f'' defined on the set on infinitely differentiable periodic functions. It has eigenvalues  $k^2\pi^2$  for  $k \in \mathbb{N}_0$  and eigenspaces

$$\ker(L) = \mathcal{H}_0 = \operatorname{span}\left\{\frac{1}{\sqrt{2}}\right\}, \quad \ker(L - k^2 \pi^2) = \mathcal{H}_k = \operatorname{span}\left\{\cos k\pi x, \sin k\pi x\right\}.$$

2.3.3. Non-periodic case on  $\mathcal{M} = [-1, 1]$  with Jacobi weight. (This example is further developed in Subsection 6.1.) Note that this example can arise when dealing with data issued from a density which itself has received a folding treatment such as in the Wicksell problem ([15, 18]). Now, the measure is

$$d\mu_{\alpha,\beta}(x) = W_{\alpha,\beta}(x)dx = (1-x)^{\alpha}(1+x)^{\beta}dx, \quad \alpha,\beta > -1$$

the distance  $\rho$  is the Euclidean distance, and L is the Jacobi operator

$$-L(f) = \frac{1}{W_{\alpha,\beta}(x)} D_x \left( (1-x^2) W_{\alpha,\beta}(x) D_x \right) f.$$

Conditions C1-C5 are satisfied, but not the Ahlfors condition C1A, unless  $\alpha = \beta = -\frac{1}{2}$ . The discrete spectral decomposition of L is given by one dimensional spectral spaces:

$$\mathbb{L}^{2}(M,\mu_{\alpha,\beta}) = \bigoplus E_{\lambda_{k}^{\alpha,\beta}}, \quad E_{\lambda_{k}^{\alpha,\beta}} = \ker(L-\lambda_{k}^{\alpha,\beta}I_{d}) = \operatorname{span}\left\{p_{k}^{\alpha,\beta}(x)\right\},$$

where  $p_k^{\alpha,\beta}(x)$  is the *k*th degree Jacobi polynomial and  $\lambda_k^{\alpha,\beta} = k(k + \alpha + \beta + 1)$ .

2.3.4. Riemannian manifold  $\mathcal{M}$  without boundary. If  $\mathcal{M}$  is a Riemannian manifold, then the Laplace operator  $\Delta_M$  is well defined on M (see [13]) and we consider

$$L = -\Delta_M.$$

If  $\mathcal{M}$  is compact, then conditions **C1-C5** are verified, including the Ahlfors condition **C1A**. Furthermore, there exists an associated discrete spectral decomposition with finite dimensional spectral eigenspaces of L:

$$\mathbb{L}^{2}(\mathcal{M},\mu) = \bigoplus E_{\lambda_{k}}, \quad E_{\lambda_{k}} = \ker(L - \lambda_{k}I_{d}), \quad \lambda_{0} = 0 < \lambda_{1} < \lambda_{1} < \cdots.$$

2.3.5. Unit sphere  $\mathcal{M} = \mathbb{S}^{d-1}$  in  $\mathbb{R}^d$ ,  $d \geq 3$ . This is the most famous Riemannian manifold with the induced structure from  $\mathbb{R}^d$ . Here  $d\mu$  is the Lebesgue measure on  $\mathbb{S}^{d-1}$ ,  $\rho$  is the geodesic distance on  $\mathbb{S}^{d-1}$ :

$$\rho(\xi,\eta) = \arccos(\langle \xi,\eta \rangle_{\mathbb{R}^d}),$$

and  $L := -\Delta_0$  with  $\Delta_0$  being the Laplace-Beltrami operator on  $\mathbb{S}^{d-1}$ . The spectral decomposition of the operator L can be described as follows:

$$\mathbb{L}^{2}(\mathbb{S}^{d-1},\mu) = \bigoplus E_{\lambda_{k}}, \quad E_{\lambda_{k}} = ker(L-\lambda_{k}I_{d}), \quad \lambda_{k} = k(k+d-2).$$

Here  $E_{\lambda_k}$  is the restriction to  $\mathbb{S}^{d-1}$  of harmonic homogeneous polynomials of degree k (spherical harmonics), see [28]. We have  $\dim(E_{\lambda_k}) = \binom{d-1}{d+k-1} - \binom{d-1}{d+k-3}$ .

2.3.6. Lie group of matrices:  $\mathcal{M} = SU(2)$ . This example is interesting in astrophysical problems, especially in the measures associated to the CMB, where instead of only measuring the intensity of the radiation we also measure spins. By definition

$$SU(2) := \left\{ \left( \begin{array}{cc} a & b \\ -\overline{b} & \overline{a} \end{array} 
ight), \quad a,b \in \mathbb{C}, |a|^2 + |b|^2 = 1 
ight\}.$$

Thus

$$q \in SU(2) \leftrightarrow q \in M(2, \mathbb{C}), \quad q^{-1} = -q^*, \quad \det(q) = 1.$$

This a compact group which topologically is the sphere  $\mathbb{S}^3 \subset \mathbb{R}^4$ . So, if

$$x = \begin{pmatrix} x_1 + ix_2 & x_3 + ix_4 \\ -(x_3 - ix_4) & x_1 - ix_2 \end{pmatrix}; \ y = \begin{pmatrix} y_1 + iy_2 & y_3 + iy_4 \\ -(y_3 - iy_4) & y_1 - iy_2 \end{pmatrix}$$
$$\|y\|^2 = \sum y_i^2 = 1 = \|x\|^2 = \sum x_i^2, \text{ then}$$

with  $||y||^2 = \sum_i y_i^2 = 1 = ||x||^2 = \sum_i x_i^2$ , then

$$\langle x, y \rangle_4 = \sum_i x_i y_i = \frac{1}{2} Tr[xy^*].$$

Thus

$$\rho_{SU(2)}(x,y) = \arccos \frac{1}{2} Tr[xy^*])$$

and for any  $q, x, y \in SU(2)$ 

$$v = \rho_{SU(2)}(qx, qy) = \rho_{SU(2)}(xq, yq) = \rho_{SU(2)}(x, y)$$

The eigenvalues of  $L = -\Delta$  are  $\lambda_k = k(k+2)$  and the dimension of the respective eigenspaces  $E_{\lambda_k}$  is  $(k+1)^2$ .

**Remark 2.2.** Looking at some of these examples an important question already arises: how to choose in a given problem the distance  $\rho$  as well as the dominating measure before even choosing the operator L and a class of regularity? In  $\mathbb{R}^d$ , most often the euclidean distance and the Lebesgue measure seems more or less unavoidable. In some other cases it might not be so obvious.

Let us take for instance the simple case of  $\mathcal{M}$  being an interval [-1, 1]. The cases of the ball, the simplex (see Section 6) and more generally sets with boundaries give rise in fact to identical discussions therefore we will focus on the case of the interval.

So, if  $\mathcal{M} = [-1, 1]$ , a possible choice -and probably the most standard one in statistical examples could be taking  $\rho$  as the euclidean distance and  $\mu$  as the Lebesgue measure. Then the usual translation kernels are available as well as the standard wavelet bases. However "something" -which generally is often swept under the carpet or not really detailed- has to be "done" about the boundary points  $\{-1, 1\}$ . Often special regularity conditions are assumed about these boundary points such as f(-1) = f(1) = 0 (subsection 2.3.2), which de facto lead to different methods for representing the functions to be estimated.

Let us now look at the choices (again for the interval [-1,1] Subsection 2.3.3) that are made in the "Jacobi" case. The distance  $\rho(x, y) = |\arccos x - \arccos y|$ , suggests a one-to-one correspondence with the semi-circle. The measure  $\mu$  ( $d\mu_{\alpha,\beta}(x) =$  $(1-x)^{\alpha}(1+x)^{\beta}dx$ ,  $\alpha, \beta > -1$ ) suggests that the points in the middle of the interval (say  $[-\frac{1}{2}, +\frac{1}{2}]$ , where the measure behaves as the Lebesgue measure) will not be weighted in the same way as the points near the boundary. And in some cases, this makes perfect sense: for instance if one needs to give a hard weight on these points because they require special attention, or at the contrary a small one.

Apart from these considerations, there are in fact two measures in the family  $\mu_{\alpha,\beta}$ , that are undeniable in the case  $\mathcal{M} = [-1,1]$  equipped with the distance  $\rho(x,y) = |\arccos x - \arccos y|$ . The first one is the Lebesgue measure (because Lebesgue is always undeniable), corresponding to  $\alpha = \beta = 0$ . The second one is  $\mu_{-\frac{1}{2},-\frac{1}{2}}$ , because in that case there is a one-to-one identification between  $(\mathcal{M},\rho,\mu_{-\frac{1}{2},-\frac{1}{2}})$ and the semi circle equipped with the euclidean distance and Lebesgue measure.

If we look more precisely into these two choices, we see that for the last case, all the required conditions including the Ahlfors one are satisfied, and the dimension d = 1, which is intuitively expected. Let us now observe that the case of the Lebesgue measure  $\mu_{0,0}$  would lead to a larger dimension d = 2.

#### 3. Background

In this section we collect some basic technical facts and results related to the setting described in Section 2 that will be needed for the development of density estimators. Most of them can be found in [4, 11, 17].

3.1. Functional calculus. A key trait of our setting is that it allows to develop a smooth functional calculus. If we recall that the operator L has been introduced as a substitute for Laplacian, we also have to recall that in  $\mathbb{R}^d$ , regularity properties of the functions are most often expressed in terms of Fourier transforms which are corresponding to spectral decompositions of the Laplacian. Hence there is no surprise that we will consider the spectral decomposition of L and define an associated functional calculus.

Let  $E_{\lambda}$ ,  $\lambda \geq 0$ , be the spectral resolution associated with the operator L in our setting. As L is non-negative, essentially self-adjoint and maps real-valued to real-valued functions, then for any real-valued, measurable, and bounded function h on  $\mathbb{R}_+$ 

(3.1) 
$$h(L) := \int_0^\infty h(\lambda) dE_{\lambda},$$

is well defined on  $\mathbb{L}^2(\mathcal{M})$ . The operator h(L), called *spectral multiplier*, is bounded on  $\mathbb{L}^2(\mathcal{M})$ , self-adjoint, and maps real-valued to real-valued functions [30]. We will be interested in integral spectral multiplier operators h(L). If h(L)(x, y) is the kernel of such an operator, it is real-valued and symmetric. From condition **C4** of our setting we know that  $e^{-tL}$  is an integral operator whose (heat) kernel  $p_t(x, y)$ is symmetric and real-valued:  $p_t(y, x) = p_t(x, y) \in \mathbb{R}$ .

3.1.1. *Examples.* Let us revisit some of the examples given in Subsection 2.3:

(a) Let  $\mathcal{M} = [-1, 1]$  be in the periodic case (Subsection 2.3.2). It is readily seen that the projection operators are:

$$P_{\mathcal{H}_0}(x,y) = \frac{1}{2}, \quad P_{\mathcal{H}_k}(x,y) = \cos k\pi (x-y).$$

Hence, formally,

$$h(L)(x,y) = \frac{1}{2}h(0) + \sum_{k \ge 1} h(k^2 \pi^2) \cos k\pi (x-y), \quad x,y \in [-1,1].$$

(b) If  $\mathcal{M}$  is a Riemanian manifold (Subsection 2.3.4), then h(L) is a kernel operator with kernel

$$h(L)(x,y) = \sum_{k} h(\lambda_k) P_k(x,y)$$

with  $P_k(x,y) = \sum_i v_i^{\lambda_k}(x) \overline{v_i^{\lambda_k}(y)}$ , where  $v_i^{\lambda_k}(x), i = 1, \dots, \dim(E_{\lambda_k})$  is an orthonormal basis of  $E_{\lambda_k}$ .

(c) In the case of the sphere (Subsection 2.3.5), the orthogonal projector operator  $P_{E_{\lambda_k}} : \mathbb{L}^2(\mathbb{S}^{d-1}) \mapsto E_{\lambda_k}$  is a kernel operator with kernel of the form

$$P_{E_{\lambda_k}}(\xi,\eta) = L_k(\langle \xi,\eta \rangle_{\mathbb{R}^d}), \text{ where } L_k(x) = \frac{1}{|\mathbb{S}^{d-1}|} \Big(1 + \frac{k}{\nu}\Big) C_k^{\nu}(x), \ \nu = \frac{d-2}{2}.$$

Here  $C_k^{\nu}(x)$  is the Gegenbauer polynomials of degree k. Usually, the polynomials  $\{C_k^{\nu}(x)\}$  are defined by the generating function

$$\frac{1}{(1-2rx+r^2)^{\nu}} = \sum_{k\geq 0} r^k C_k^{\nu}(x), \quad |r|<1, \ |x|<1.$$

Hence, formally

$$h(L)(\xi,\eta) = \sum_{k} h(k(k+d-2))L_k(\langle \xi,\eta \rangle_{\mathbb{R}^d}), \quad \xi,\eta \in \mathbb{S}^{d-1}.$$

(d) In the case of SU(2) (Subsection 2.3.6), the orthogonal projector operator  $P_{E_{\lambda_k}} : \mathbb{L}^2(SU(2)) \mapsto E_{\lambda_k}$  is the operator with kernel

$$P_{E_{\lambda_k}}(f)(\xi,\eta) = L_k\left(\frac{1}{2}Tr[\xi\eta^*]\right)$$

where  $L_k(x) = \frac{1}{|\mathbb{S}^3|} (1+k) C_k^1(x)$ . Hence, formally

$$h(L)(\xi,\eta) = \sum_{k} h(k(k+2)) L_k(\frac{1}{2} Tr[\xi\eta^*]), \quad \xi,\eta \in SU(2).$$

Our further development will heavily depend on the following result from the smooth functional calculus induced by the heat kernel, developed in [17, Theorem 3.4]. It asserts the localization properties of general spectral multipliers of the form  $g(\delta\sqrt{L})$  (corresponding to functions of the form  $h(u) = g(\delta\sqrt{u})$  in (3.1)). Again the appearance of the square root is by analogy with the Laplacian, which is an operator of degree 2. It is also interesting to remark that (3.2) is valid in  $\mathbb{R}^d$  when  $g(\delta\sqrt{L})$  is replaced by  $[\frac{1}{\delta}]^d G(\frac{x-y}{\delta})$ , where G is a bounded compactly supported function  $\mathbb{R}^d \longrightarrow \mathbb{R}$  for instance. This result is a building block for the properties of the kernel estimators defined in the sequel.

**Theorem 3.1.** Let  $g \in C^N(\mathbb{R})$ , N > d, be even, real-valued, and  $\operatorname{supp} g \subset [-R, R]$ , R > 0. Then  $g(\delta\sqrt{L})$ ,  $\delta > 0$ , is an integral operator with kernel  $g(\delta\sqrt{L})(x, y)$  satisfying

(3.2) 
$$\left| g(\delta\sqrt{L})(x,y) \right| \le c|B(x,\delta)|^{-1} \left( 1 + \delta^{-1}\rho(x,y) \right)^{-N+\frac{d}{2}}, \, \forall \, x, \, y \in \mathcal{M},$$

where c > 0 is a constant depending on  $||g||_{\infty}$ ,  $||g^{(N)}||_{\infty}$ , N, R and the constants  $c_0, c_4, c_5$  from our setting.

Furthermore, for any  $\delta > 0$  and  $x \in \mathcal{M}$ 

(3.3) 
$$\int_{\mathcal{M}} g(\delta\sqrt{L})(x,y)d\mu(y) = g(0).$$

3.2. Geometric properties. Conditions C1 and C2 yield

(3.4) 
$$|B(x,r)| \ge (c_3/c_0)r^d, \quad x \in \mathcal{M}, \ 0 < r \le 1.$$

To compare the volumes of balls with different centers  $x, y \in \mathcal{M}$  and the same radius r we will use the inequality

(3.5) 
$$|B(x,r)| \le c_0 \left(1 + \frac{\rho(x,y)}{r}\right)^d |B(y,r)|, \quad x,y \in \mathcal{M}, \ r > 0.$$

As  $B(x,r) \subset B(y,\rho(y,x)+r)$  the above inequality is immediate from (2.2).

We will also need the following simple inequality (see [4, Lemma 2.3]): If  $\tau>d,$  then for any  $\delta>0$ 

(3.6) 
$$\int_{\mathcal{M}} \left( 1 + \delta^{-1} \rho(x, y) \right)^{-\tau} d\mu(y) \le c |B(x, \delta)|, \quad x \in \mathcal{M},$$

where  $c = (2^{-d} - 2^{-\tau})^{-1}$ .

3.3. Spectral spaces. We recall the definition of the spectral spaces  $\Sigma_{\lambda}^{p}$ ,  $1 \leq p \leq \infty$ , from [4]. Denote by  $C_{0}^{\infty}(\mathbb{R})$  the set of all even real-valued compactly supported functions. We define

$$\Sigma_{\lambda}^{p} := \left\{ f \in \mathbb{L}^{p}(\mathcal{M}) : \theta(\sqrt{L})f = f \text{ for all } \theta \in C_{0}^{\infty}(\mathbb{R}), \ \theta \equiv 1 \text{ on } [0, \lambda] \right\}, \ \lambda > 0.$$

We will need the following proposition (Nikolski type inequality):

**Proposition 3.2.** Let  $1 \le p \le q \le \infty$ . If  $g \in \Sigma_{\lambda}^{p}$ ,  $\lambda \ge 1$ , then  $g \in \Sigma_{\lambda}^{q}$  and

(3.7) 
$$||g||_q \le c_\star \lambda^{d(1/p-1/q)} ||g||_p,$$

where the constant  $c_{\star} > 1$  is independent of p and q.

This proposition was established in [4, Proposition 3.12] (see also [17, Proposition 3.11]). We present its proof in the appendix because we need to control the constant  $c_{\star}$ .

3.4. Wavelets. In the setting of this article, wavelet type frames for Besov and Triebel-Lizorkin spaces are developed in [17]. Here, we review the construction of the frames from [17] and their basic properties. Indeed, in this setting the 'wavelets' do not form an orthonormal basis but a frame. In this case, the construction of a 'dual wavelet system' is necessary to get a representation of type (3.14).

This construction is inspired by to the Littlewood-Paley construction of the standard wavelets introduced by [7],[8], [9].

The construction of frames involves a "dilation" constant b > 1 whose role is played by 2 in the wavelet theory on  $\mathbb{R}$ .

The construction starts with the selection of a function  $\Psi_0 \in C^{\infty}(\mathbb{R}_+)$  with the properties:  $\Psi_0(\lambda) = 1$  for  $\lambda \in [0, 1], 0 \leq \Psi_0(\lambda) \leq 1$ , and  $\operatorname{supp} \Psi_0 \subset [0, b]$ . Denote  $\Psi(\lambda) := \Psi_0(\lambda) - \Psi_0(b\lambda)$  and set  $\Psi_j(\lambda) := \Psi(b^{-j}\lambda), j \in \mathbb{N}$ . From this it readily follows that

(3.8) 
$$\sum_{j=0}^{J} \Psi_j(\lambda) = \Psi_0(b^{-J}\lambda), \quad \lambda \in \mathbb{R}_+.$$

For  $j \ge 0$  we let  $\mathcal{X}_j \subset \mathcal{M}$  be a maximal  $\delta_j$ -net on  $\mathcal{M}$  with  $\delta_j := c_6 b^{-j}$ . It is easy to see that for any  $j \ge 0$  there exists a disjoint partition  $\{A_{j\xi}\}_{\xi \in \mathcal{X}_j}$  of  $\mathcal{M}$  consisting of measurable sets such that

$$B(\xi, \delta_j/2) \subset A_{j\xi} \subset B(\xi, \delta_j), \quad \xi \in \mathcal{X}_j.$$

Here  $c_6 > 0$  is a sufficiently small constant (see [17]).

**Lemma 3.3.** If  $\mathcal{M}$  is compact, then there exists a constant  $c_7 > 0$  such that

(3.9) 
$$\operatorname{card}(\mathcal{X}_j) \le c_7 b^{jd}, \quad j \ge 0$$

*Proof.* Assume  $\mathcal{M}$  is compact and let  $\mathcal{X}_{\delta}$  be a maximal  $\delta$ -net on  $\mathcal{M}, \delta > 0$ . Then

$$\sum_{\xi \in \mathcal{X}_{\delta}} \int_{B(\xi, \delta/2)} \frac{d\mu(x)}{\mu(B(x, \delta))} \leq \int_{\mathcal{M}} \frac{d\mu(x)}{\mu(B(x, \delta))} \leq \sum_{\xi \in \mathcal{X}_{\delta}} \int_{B(\xi, \delta)} \frac{d\mu(x)}{\mu(B(x, \delta))}$$

Therefore, using (2.2) we get

(3.10) 
$$\frac{1}{c_0 4^d} \operatorname{card}(\mathcal{X}_{\delta}) \le \int_M \frac{d\mu(x)}{\mu(B(x,\delta))} \le c_0 2^d \operatorname{card}(\mathcal{X}_{\delta}).$$

Since  $\mathcal{M}$  is compact we have  $\mu(\mathcal{M}) < \infty$  and  $B(x, D) = \mathcal{M}$  for  $x \in \mathcal{M}$ , where D is the diameter of  $\mathcal{M}$ , which is finite (see [4]). Using again (2.2) we get  $\mu(M) = \mu(B(x, D)) \leq c_0(\frac{D}{\delta})^d \mu(B(x, \delta) \text{ for } x \in M$ . Hence

$$\frac{1}{c_0 4^d} \operatorname{card}(\mathcal{X}_{\delta}) \le \int_M \frac{d\mu(x)}{\mu(B(x,\delta))} \le c_0 (D/\delta)^d,$$

which implies (3.9).

The *j*th level frame elements  $\psi_{j\xi}$  are defined by

(3.11) 
$$\psi_{j\xi}(x) := |A_{j\xi}|^{1/2} \Psi_j(\sqrt{L})(x,\xi), \quad \xi \in \mathcal{X}_j.$$

We will also use the more compact notation  $\psi_{\xi} := \psi_{j\xi}$  for  $\xi \in \mathcal{X}_j$ .

Let  $\mathcal{X} := \bigcup_{j \ge 0} \mathcal{X}_j$ , where equal points from different sets  $\mathcal{X}_j$  will be regarded as distinct elements of  $\mathcal{X}$ , so  $\mathcal{X}$  can be used as an index set. Then  $\{\psi_{\xi}\}_{\xi \in \mathcal{X}}$  is Frame #1.

The construction of a dual frame  $\{\tilde{\psi}_{\xi}\}_{\xi\in\mathcal{X}} = \bigcup_j \{\tilde{\psi}_{j\xi}\}_{\xi\in\mathcal{X}_j}$  is much more involved; we refer the reader to §4.3 in [17] for the details.

By construction, the two frames satisfy

(3.12) 
$$\Psi_j(\sqrt{L})(x,y) = \sum_{\xi \in \mathcal{X}_j} \psi_{j\xi}(y) \tilde{\psi}_{j\xi}(x), \quad j \ge 0.$$

A basic result from [17] asserts that for any  $f \in \mathbb{L}^p(\mathcal{M}, d\mu), 1 \leq p < \infty$ ,

(3.13) 
$$f = \sum_{j \ge 0} \Psi_j(\sqrt{L}) f \quad \text{(convergence in } \mathbb{L}^p)$$

and the same holds in  $\mathbb{L}^{\infty}$  if f is uniformly continuous and bounded (UCB) on  $\mathcal{M}$ . As a consequence, for any  $f \in \mathbb{L}^p(\mathcal{M}, d\mu)$ ,  $1 \leq p \leq \infty$ , ( $\mathbb{L}^{\infty} = \text{UCB}$ ) we have

(3.14) 
$$f = \sum_{j=0}^{\infty} \sum_{\xi \in \mathcal{X}_j} \langle f, \tilde{\psi}_{j\xi} \rangle \psi_{j\xi} \quad \text{(convergence in } \mathbb{L}^p\text{)}.$$

Furthermore, frame decomposition results are established in [17] for Besov and Triebel-Lizorkin spaces with full range of indices.

**Properties of frames in the Ahlfors regularity case.** We next present some properties of the frame elements in the case when condition **C1A** is stipulated (see [17]).

1. Localization: For every  $k \in \mathbb{N}$ , there exists a constant c(k) > 0 such that

(3.15) 
$$|\psi_{j\xi}(x)|, \ |\tilde{\psi}_{j\xi}(x)| \le c(k)b^{jd/2} (1+b^j\rho(x,\xi))^{-k}, \ x \in \mathcal{M}.$$

2. Norm estimation: For  $1 \le p \le \infty$ 

(3.16) 
$$c_{\diamond}^{-1}b^{jd(\frac{1}{2}-\frac{1}{p})} \leq \|\psi_{j\xi}\|_{p}, \ \|\tilde{\psi}_{j\xi}\|_{p} \leq c_{\diamond}b^{jd(\frac{1}{2}-\frac{1}{p})}, \ \xi \in \mathcal{X}_{j}, \ j \geq 0.$$

3. For  $1 \leq p \leq \infty$ 

(3.17) 
$$\left\|\sum_{\xi\in\mathcal{X}_j}\lambda_{\xi}\psi_{j\xi}\right\|_p \le c_{\diamond}b^{jd(\frac{1}{2}-\frac{1}{p})} \left(\sum_{\xi\in\mathcal{X}_j}|\lambda_{\xi}|^p\right)^{1/p}, \quad j\ge 0,$$

with the usual modification when  $p = \infty$ . Above the constant  $c_{\diamond} > 1$  depends only on  $p, b, \Psi_0$ , and the structural constants of the setting.

3.5. **Besov spaces.** We will deal with probability density functions (pdf's) in Besov spaces associated to the operator L in our setting. These spaces are developed in [4, 17]. Definition 3.4 coincides in  $\mathbb{R}^d$  with one the definitions of usual Besov spaces with L replaced by Laplacian ( $-\Delta$  in fact to get a positive operator).

Here we present some basic facts about Besov spaces that will be needed later on.

Let  $\Phi_0, \Phi \in C^{\infty}(\mathbb{R}_+)$  be real-valued functions satisfying the conditions:

(3.18) 
$$\operatorname{supp} \Phi_0 \subset [0, b], \ \Phi_0(\lambda) = 1 \text{ for } \lambda \in [0, 1], \ \Phi_0(\lambda) \ge c > 0 \text{ for } \lambda \in [0, b^{3/4}],$$

(3.19) 
$$\operatorname{supp} \Phi \subset [b^{-1}, b], \ \Phi(\lambda) \ge c > 0 \text{ for } \lambda \in [b^{-3/4}, b^{3/4}].$$

Set  $\Phi_j(\lambda) := \Phi(b^{-j}\lambda)$ , for  $j \ge 1$ .

**Definition 3.4.** Let s > 0,  $1 \le p \le \infty$ , and  $0 < q \le \infty$ . The Besov space  $B_{pq}^s = B_{pq}^s(\mathcal{M}, L)$  is defined as the set of all functions  $f \in \mathbb{L}^p(\mathcal{M}, \mu)$  such that

(3.20) 
$$||f||_{B^s_{pq}} := \left(\sum_{j\geq 0} \left(b^{sj} ||\Phi_j(\sqrt{L})f||_p\right)^q\right)^{1/q} < \infty,$$

where the  $\ell^q$ -norm is replaced by the sup-norm if  $q = \infty$ .

Note that as shown in [17] the above definition of the Besov spaces  $B_{pq}^s$  is independent of the particular choice of  $\Phi_0$ ,  $\Phi$  satisfying (3.18)-(3.19). For example with  $\Psi_i$  from the definition of the frame elements in §3.4 we have

(3.21) 
$$\|f\|_{B^s_{pq}} \sim \left(\sum_{j\geq 0} \left(b^{sj} \|\Psi_j(\sqrt{L})f\|_p\right)^q\right)^{1/q}$$

with the usual modification when  $q = \infty$ . The following useful inequality follows readily from above

(3.22) 
$$\|\Psi_j(\sqrt{L})f\|_p \le cb^{-sj} \|f\|_{B^s_{pq}}, \quad f \in B^s_{pq}, \ j \ge 0.$$

As in  $\mathbb{R}^d$ , we will need some embedding results involving Besov spaces. Recall the definition of embeddings: Let X and Y be two (quasi-)normed spaces. We say that X is continuously embedded in Y and write  $X \hookrightarrow Y$  if  $X \subset Y$  and for each  $f \in X$  we have  $||f||_Y \leq c||f||_X$ , where c > 0 is a constant independent of f.

**Proposition 3.5.** (i) If  $1 \le q \le r \le \infty$ ,  $0 < \tau \le \infty$ , s > 0, and  $\mu(\mathfrak{M}) < \infty$ , then  $B^s_{r\tau} \hookrightarrow B^s_{q\tau}$ .

(ii) If 
$$1 \le r \le q \le \infty$$
,  $0 < \tau \le \infty$  and  $s > d(\frac{1}{r} - \frac{1}{q})$ , then  $B^s_{r\tau} \hookrightarrow B^{s-d(\frac{1}{r} - \frac{1}{q})}_{q\tau}$ .  
(iii) If  $1 \le r \le \infty$ ,  $0 < \tau \le \infty$ , and  $s > d/r$ , then  $B^s_{r\tau} \hookrightarrow \mathbb{L}^{\infty}$ .  
(iv) If  $1 \le p \le r \le \infty$ ,  $0 < \tau \le \infty$ ,  $s > 0$ , and  $\mu(\mathcal{M}) < \infty$ , then  $B^s_{r\tau} \hookrightarrow \mathbb{L}^p$ .

To streamline our presentation we defer the proof of this proposition to the appendix.

**Besov spaces in the Ahlfors regularity case.** For the development of adaptive density estimators in Section 5 we will need some additional facts from the theory of Besov spaces when condition **C1A** is assumed. We first introduce the Besov bodies.

**Definition 3.6.** Assume s > 0,  $1 \le p \le \infty$ ,  $0 < q \le \infty$ , and let  $\mathcal{X} := \bigcup_{j \ge 0} \mathcal{X}_j$  be from the definition of the frames in §3.4. The Besov body  $\mathfrak{b}_{pq}^s = \mathfrak{b}_{pq}^s(\mathcal{X})$  is defined as the set of all sequences  $\{a_{\xi}\}_{\xi \in \mathcal{X}}$  of real (or complex) numbers such that

(3.23) 
$$||a||_{\mathfrak{b}_{pq}^{s}} := \Big(\sum_{j\geq 0} b^{jsq} \Big(\sum_{\xi\in\mathcal{X}_{j}} \left[b^{-jd(\frac{1}{p}-\frac{1}{2})}|a_{\xi}|\right]^{p}\Big)^{q/p}\Big)^{1/q} < \infty,$$

where the  $\ell^q$ -norm is replaced by the sup-norm if  $q = \infty$ .

One of the principle results in [17] asserts that the Besov spaces  $B_{pq}^s$  can be completely characterized in terms of the Besov bodies  $\mathfrak{b}_{pq}^s$  of the frame coefficients of the respective functions. To be specific, denote

(3.24) 
$$\beta_{j\xi}(f) := \langle f, \psi_{j\xi} \rangle, \quad \xi \in \mathcal{X}_j, \ j \ge 0.$$

We will also use the more compact notation:  $\beta_{\xi}(f) := \beta_{j\xi}(f)$  for  $\xi \in \mathcal{X}_j$ . In the current setting, assume s > 0,  $1 \le p \le \infty$ , and  $0 < q \le \infty$ . In light of [17, Theorem 6.10]  $f \in B_{pq}^s$  if and only if  $\{\beta_{\xi}(f)\} \in \mathfrak{b}_{pq}^s$  with equivalent norms:

(3.25) 
$$\|f\|_{B^s_{pq}} \sim \|\{\beta_{\xi}(f)\}\|_{\mathfrak{b}^s_{pq}}$$

This implies that if  $f \in B_{pq}^s$  for some s > 0,  $p \ge 1$ , and  $0 < q \le \infty$ , then

(3.26) 
$$\left(\sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}(f)|^p\right)^{1/p} \le c b^{-j(s+d(\frac{1}{2}-\frac{1}{p}))} ||f||_{B^s_{pq}}, \quad j \ge 0$$

where c = c(s, p, q) > 0.

By (3.17) and (3.26) it follows that, if  $f \in B^s_{pq}$  for some  $s > 0, p \ge 1$ , and  $0 < q \le \infty$ , then

(3.27) 
$$\left\|\sum_{\xi\in\mathcal{X}_j}\beta_{j\xi}(f)\psi_{j\xi}\right\|_p \le cb^{-sj}\|f\|_{B^s_{pq}}, \quad j\ge 0.$$

## 4. Kernel density estimators on the metric measure space $\mathcal{M}$

Our goal in this section is to introduce and study kernel density estimators (kde's) on a metric measure space  $(\mathcal{M}, \rho, \mu)$  in the general setting described in Section 2. More precisely, in this section, we assume that conditions **C1–C5** are satisfied, and do not necessarily assume the Ahlfors regular volume condition **C1A**.

To explain our construction of kernel estimators we begin by considering the classical example of the periodic case on  $\mathcal{M} = [-1, 1]$ , presented in Subsection 2.3.2. It will be commonly admitted among nonparametric statisticians to define an estimator of the form

$$\hat{f}_T(x) = \frac{1}{2} + \frac{1}{n} \sum_{i=1}^n \sum_{1 \le k \le T} \cos k\pi (x - X_i).$$

It falls into the category of orthogonal series estimator. It is well known that these estimators have nice  $\mathbb{L}^2$  properties but can drastically fail in  $\mathbb{L}^p$ ,  $p \neq 2$ , or locally.

In our setting we will replace  $\hat{f}_T$  by a 'smoothed version':

(4.1) 
$$\hat{K}_{\delta}(x) = \frac{1}{2}K(0) + \frac{1}{n}\sum_{i=1}^{n}\sum_{k\geq 1}K(\delta k)\cos k\pi(x-X_i) =: \frac{1}{n}\sum_{i=1}^{n}K_{\delta}(x,X_i),$$

where K is a smooth function  $\mathbb{R}_+ \longrightarrow \mathbb{R}$  eventually vanishing at infinity, and  $K_{\delta}(x,y) = \frac{1}{2}K(0) + \sum_{k>1} K(\delta k) \cos k\pi (x-y).$ 

In analogy to this case, replacing the circle by  $\mathcal{M}$  and the Laplacian by the operator -L we can naturally introduce kde's on  $\mathcal{M}$  by means of the machinery of spectral multipliers.

Let  $K : \mathbb{R}_+ \to \mathbb{R}$  be a bounded and measurable function. As we already alluded to in §3.1, if  $E_{\lambda}$ ,  $\lambda \ge 0$ , is the spectral resolution associated with the operator L, then the operator

(4.2) 
$$K(\sqrt{L}) := \int_0^\infty K(\sqrt{\lambda}) dE_\lambda$$

is well defined on  $\mathbb{L}^2(\mathcal{M})$ . Furthermore, the operator  $K(\sqrt{L})$  is self-adjoint and bounded on  $\mathbb{L}^2(\mathcal{M})$  with norm  $\|K(\sqrt{L})\|_{2\to 2} := \sup_{h, \|h\|_2 \leq 1} \|H(h)\|_2 \leq \|K\|_{\infty}$ .

We are interested in multiplier operators  $K(\sqrt{L})$  that are integral operators. In this case, since the function K is real-valued its *kernel*  $K(\sqrt{L})(x, y)$  is real-valued and symmetric. We will use Theorem 3.1 to define a family of multiplier operators whose kernels are suitable for the construction of kernel density estimators on  $\mathcal{M}$ .

We now introduce the kde's in the general setting of this article.

**Definition 4.1.** Let  $X_1, \ldots, X_n$  be i.i.d. random variables on  $\mathcal{M}$  in our setting. Let  $K(\delta\sqrt{L})(x,y)$  with  $0 < \delta \leq 1$  (the bandwidth) be the kernel of the integral operator  $K(\delta\sqrt{L})$ , where  $K : \mathbb{R}_+ \to \mathbb{R}$ . The associated kernel density estimator is defined by

(4.3) 
$$\widehat{K}_{\delta}(x) := \frac{1}{n} \sum_{i=1}^{n} K(\delta \sqrt{L})(X_i, x), \quad x \in \mathcal{M}.$$

**Remarks 4.2.** - The analogy with the Torus again explains why we take  $\sqrt{L}$  instead of L: in the torus, case the eigenvalues are  $\pi^2 k^2$ . It is mostly a comfort choice. Replacing  $\sqrt{L}$  by L is possible but would lead to a different regularity scale. - Again the analogy with the Torus case could lead to a choice of the form K to be the indicator function of the interval [0,1] for instance. This choice would induce  $\mathbb{L}^2$  properties, but not  $\mathbb{L}^p$  because this function is not smooth enough to get the concentration inequalities of Theorem 3.1.

- If  $K(\lambda) = e^{-\lambda^2}$ , then  $K(\delta\sqrt{L})(x, y) = p_{\delta^2}(x, y)$  (the "heat kernel") can be used to define a kernel density estimator. This choice relies to the Bayesian estimator provided in [2].

- This type of kernel estimators although constructed upon orthogonal projectors, because of the smoothing function K will finally have properties which are comparable to translation kernel estimators in  $\mathbb{R}^d$ . In  $\mathbb{R}^d$  some properties such as a number of moment annulation (see for instance [29]) to get a correct biais are required which will be here replaced by the vanishing properties at infinity of the function K and the smoothness of the function K.

4.1. Upper bound estimates for kernel density estimators. We will especially study kernel density estimators induced by compactly supported  $C^{\infty}$  multipliers, often called Littlewood-Paley functions. In fact other type of kernels among the family of multipliers could lead to quite similar results. The Littlewood-Paley are especially powerful technically to obtain upper-bounds. More explicitly, let  $\Phi$ be an even  $C^{\infty}(\mathbb{R})$  real-valued function with the following properties:

(4.4)  $\operatorname{supp} \Phi \subset [-1,1] \text{ and } \Phi(\lambda) = 1 \text{ for } \lambda \in [-1/2,1/2].$ 

By Theorem 3.1 it follows that  $\Phi(\delta\sqrt{L})$  is an integral operator with well localized symmetric kernel  $\Phi(\delta\sqrt{L})(x, y)$  and the Markov property:

(4.5) 
$$\int_{\mathcal{M}} \Phi(\delta\sqrt{L})(x,y)d\mu(y) = 1$$

As before we assume that  $X_1, \ldots, X_n$   $(n \ge 2)$  are i.i.d. random variables with values on  $\mathcal{M}$  and common probability density function (pdf) f with respect to the measure  $\mu$  on the space  $\mathcal{M}$ . Let  $X_i \sim X$ . We will denote by  $\mathbb{E} = \mathbb{E}_f$  the expectation with respect to the probability measure  $\mathbb{P} = \mathbb{P}_f$ . We are interested in the kernel

density estimator

(4.6) 
$$\widehat{\Phi}_{\delta}(x) = \widehat{\Phi}_{\delta}(x, X_1, \dots, X_n) := \frac{1}{n} \sum_{i=1}^n \Phi(\delta \sqrt{L})(x, X_i), \ \forall \ x \in \mathcal{M}.$$

4.2. Examples of Littlewood Paley kernel estimates. Let us take some examples issued from section 3.1.1. We already discussed the case [-1, 1] as a Torus in the introduction this section.

- For [-1, 1] in the Jacobi framework, (section 2.3.3), we get the following estimator:

$$\widehat{\Phi}_{\delta}(x) = \frac{1}{n} \sum_{i=1}^{n} \sum_{k} \Phi(\delta \sqrt{k(k+\alpha+\beta+1)}) p_k^{\alpha,\beta}(x) p_k^{\alpha,\beta}(X_i), \ \forall \ x \in [-1,1]$$

where  $p_k^{\alpha,\beta}(x)$  is the normalized Jacobi polynomial ( $\int_{-1}^1 |p_k^{\alpha,\beta}(x)|^2 (1-x)^{\alpha} (1+x)^{\beta} dx = 1$ ).

- For the sphere, (section 2.3.5), we get the following estimator

$$\widehat{\Phi}_{\delta}(x) = \frac{1}{n} \sum_{i=1}^{n} \sum_{k} \Phi(\delta \sqrt{k(k+d-2)}) L_k(\langle x, X_i \rangle_{\mathbb{R}^d}), \ \forall \ x \in \ \mathbb{S}^{d-1}$$

where

$$L_k(x) = \frac{1}{|\mathbb{S}^{d-1}|} \left(1 + \frac{k}{\nu}\right) C_k^{\nu}(x), \ \nu = \frac{d-2}{2}$$

- For SU(2), (section 2.3.6), we get the following estimator,

$$\widehat{\Phi}_{\delta}(x) = \frac{1}{n} \sum_{i=1}^{n} \sum_{k} \Phi(\delta \sqrt{k(k+2)}) L_k \left(\frac{1}{2} Tr[X_i x^*]\right), \ \forall \ x \in \ SU(2)$$

with  $L_k(x) = \frac{1}{|\mathbb{S}^3|} (1+k) C_k^1(x).$ 

4.3. Upper bound results. We next study the approximation of pdf's f by such kernel estimators. We first establish oracle inequalities:

**Theorem 4.3.** Assume  $1 \le p \le \infty$  and let  $\Phi$  be a Littlewood-Paley function as above. In the setting described above and with  $\widehat{\Phi}_{\delta}$  from (4.6) we have: (i) If  $2 \le p < \infty$ , then

$$\mathbb{E}\|\widehat{\Phi}_{\delta} - f\|_{p} \leq \frac{c(p)}{(n\delta^{d})^{1-\frac{1}{p}}} + \frac{c(p)}{(n\delta^{d})^{\frac{1}{2}}} \|f\|_{\frac{p}{2}}^{\frac{1}{2}} + \|\Phi(\delta\sqrt{L})f - f\|_{p}, \quad 0 < \delta \leq 1.$$

(ii) If  $1 \le p < 2$  and  $\operatorname{supp}(f) \subset B(x_0, R)$  for some  $x_0 \in \mathcal{M}$  and R > 0, then

$$\mathbb{E}\|\widehat{\Phi}_{\delta} - f\|_{p} \le \frac{c(p)}{(n\delta^{d})^{\frac{1}{2}}} |B(x_{0}, R)|^{\frac{1}{p} - \frac{1}{2}} + \|\Phi(\delta\sqrt{L})f - f\|_{p}, \quad 0 < \delta \le 1.$$

(iii) There exists a constant c such that for any  $q \ge 2$  and  $0 < \delta \le 1$  we have

$$\mathbb{E}\|\widehat{\Phi}_{\delta} - f\|_{\infty} \le c\delta^{-\frac{d}{q}} \left(\frac{q}{(n\delta^{d})^{1-\frac{1}{q}}} + \frac{q^{1/2}}{(n\delta^{d})^{\frac{1}{2}}} \|f\|_{\infty}^{\frac{1}{2}-\frac{1}{q}}\right) + \|\Phi(\delta\sqrt{L})f - f\|_{\infty}.$$

We next estimate the rates of  $\mathbb{L}^p\text{-approximation}$  of pdf's f lying in Besov space balls by kernel estimators. Denote

(4.7) 
$$B_{p\tau}^{s}(m) := \left\{ f \text{ is } pdf : ||f||_{B_{p\tau}^{s}} \le m \right\}$$

$$(4.8) \quad B^s_{p\tau}(m, x_0, R) := \left\{ f \in B^s_{p\tau}(m) : \operatorname{supp} f \subset B(x_0, R) \right\}, \quad x_0 \in \mathcal{M}, \ m, R > 0.$$

Here is our main result on the properties of these estimators, for density functions in Besov spaces, when the risk and the regularity classes are defined with the same norm.

**Theorem 4.4.** Assume s > 0,  $1 \le p \le \infty$ ,  $0 < \tau \le \infty$ , m > 0, and let  $\Phi$  be a Littlewood-Paley function as above. In the setting described above and with  $\widehat{\Phi}_{\delta}$  from (4.6) we have:

(i) If  $2 \le p < \infty$  and  $\delta = n^{-\frac{1}{2s+d}}$ , then

(4.9) 
$$\sup_{f \in B^s_{p\tau}(m)} \mathbb{E} \|\widehat{\Phi}_{\delta} - f\|_p \le cn^{-\frac{s}{2s+d}},$$

where  $c = c(p, s, m, \tau) > 0$ .

(*ii*) If 
$$1 \le p < 2$$
,  $x_0 \in \mathcal{M}$ ,  $R > 0$ , and  $\delta = n^{-\frac{1}{2s+d}}$ , then

(4.10) 
$$\sup_{f \in B_{p\tau}^s(m,x_0,R)} \mathbb{E} \|\widehat{\Phi}_{\delta} - f\|_p \le cn^{-\frac{s}{2s+d}},$$

where  $c = c(p, s, m, \tau, x_0, R) > 0.$ (*iii*) If  $\delta = \left(\frac{\log n}{n}\right)^{\frac{1}{2s+d}}$ , then (4.11)  $\sup_{f \in B_{\infty\tau}^s(m)} \mathbb{E} \|\widehat{\Phi}_{\delta} - f\|_{\infty} \le c \left(\frac{\log n}{n}\right)^{\frac{s}{2s+d}},$ 

where  $c = c(s, m, \tau) > 0$ .

**Remarks 4.5.** - Note that we do not claim that the above rates are necessarily minimax, although they show similarities with the results established in  $\mathbb{R}^d$ . The length of the paper does not allow to investigate the full lower bounds results. Let us just mention that if we add in this setting the Ahlfors condition C1A, then the lower bounds can be obtained in accordance with the upper bounds using a proof which is a direct adaptation of the one given in the case of the sphere in [1]. In the case where the Ahlfors condition is not valid, the problem is more complex since not only the regularity might be non-homogeneous due to Besov conditions but also the dimension itself may vary spatially. In this case, the upper bounds might not be optimal.

- It is interesting to compare the obtained upper bounds for  $\mathcal{M} = [-1, 1]$  in the different cases (Torus or Jacobi). In the Torus case, no surprise: the rate is the usual one with dimension d = 1. In the Jacobi case, the dimension is  $d = 1 + (2\alpha + 1)_+ \vee (2\beta + 1)_+$ , which in particular in the case  $\alpha = \beta = 0$  (corresponding to  $\mu$  the Lebesgue measure), gives a slower rate than the usual one. This is due to the fact that the 'boundaries' are playing a role so the spaces of approximation are not the same. The case  $\alpha = \beta = -\frac{1}{2}$ , which was corresponding to perfect identification with the semi-circle (see Remark 2.2) provides the expected rate with dimension d = 1.

We next compile some additional facts we need about kernels in the setting of this article and then carry out the proof of Theorems 4.3 and 4.4.

4.4. Spectral multiplier integral operators. The operator  $\Phi(\delta\sqrt{L})$  and its symmetric kernel  $\Phi(\delta\sqrt{L})(x,y)$  from above have a number of useful properties that we describe and prove next.

(a) For any k > d there exists a constant  $c_k > 0$  such that

(4.12) 
$$|\Phi(\delta\sqrt{L})(x,y)| \le c(k)|B(x,\delta)|^{-1} (1+\delta^{-1}\rho(x,y))^{-k}, \quad x,y \in \mathcal{M}, \ 0 < \delta \le 1,$$

where the constant c(k) > 0 depends only on k,  $\Phi$ , and constant from the setting in Section 2. This inequality follows immediately from Theorem 3.1.

(b) For any  $1 \le p \le \infty$ 

(4.13) 
$$\|\Phi(\delta\sqrt{L})(x,\cdot)\|_p \le c|B(x,\delta)|^{\frac{1}{p}-1} \le c_\star \delta^{-d(1-\frac{1}{p})}, \quad x \in \mathcal{M}, \ 0 < \delta \le 1,$$

where the constant  $c_{\star} > 0$  is independent of p. This estimate follows readily by (4.12), (3.6), and (3.4), just as the estimates in (7.3).

(c) Let X be a random variable on  $\mathcal{M}$  and  $X \sim f(u)d\mu(u)$ . Then

(4.14) 
$$\mathbb{E}(\Phi(\delta\sqrt{L})(x,X)) = \int_{\mathcal{M}} \Phi(\delta\sqrt{L})(x,u)f(u)d\mu(u) = \Phi(\delta\sqrt{L})f(x), \quad x \in \mathcal{M}.$$

This is a well known property of expected values.

We next estimate the bias term of the risk.

**Proposition 4.6.** Let s > 0,  $1 \le p \le \infty$ ,  $0 < q \le \infty$ . If  $f \in B^s_{pq}$ , then  $f \in \mathbb{L}^p$  and

(4.15) 
$$\|\Phi(\delta\sqrt{L})f - f\|_p \le c\delta^s \|f\|_{B^s_{pq}}, \quad 0 < \delta \le 1,$$

where c = c(s, p, q) > 0.

This statement is quite standard. For completeness we give its proof in the appendix.

We will also need the following two lemmas:

**Lemma 4.7.** Let  $2 \le p < \infty$  and  $0 < \delta \le 1$ . Then for any pdf f on  $\mathfrak{M}$  we have

(4.16) 
$$\left(\int_{\mathcal{M}} \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^p f(u) d\mu(u) d\mu(x)\right)^{1/p} \le c_\star \delta^{-d(1-1/p)}$$

and

(4.17) 
$$\left( \int_{\mathcal{M}} \left( \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^2 f(u) d\mu(u) \right)^{p/2} d\mu(x) \right)^{1/p} \le c_\star \delta^{-d/2} \|f\|_{p/2}^{1/2},$$

where  $c_{\star} > 0$  is the constant from (4.13);  $c_{\star}$  is independent of p.

*Proof.* Denote by  $S_1$  the quantity on the left-hand side in (4.16). To estimate  $S_1$  we use Fubini's theorem, (4.13), and the fact that  $\int_{\mathcal{M}} f(u)d\mu(u) = 1$ . We obtain

(4.18) 
$$S_1 = \left(\int_{\mathcal{M}} \|\Phi(\delta\sqrt{L})(\cdot, u)\|_p^p f(u) d\mu(u)\right)^{\frac{1}{p}} \le c_\star \delta^{-d(1-1/p)}$$

which confirms (4.16).

Let  $S_2$  denote the quantity on the left-hand side in (4.17) and consider the integral operator T with kernel

$$T(x,y) := |\Phi(\delta\sqrt{L})(x,y)|^2.$$

By (4.13) it follows that  $||T(x,\cdot)||_1 = ||T(\cdot,y)||_1 \le c_\star^2 \delta^{-d}$ . Therefore, by Schur's lemma (see e.g. [10, Theorem 6.36]) we obtain

(4.19) 
$$S_2^2 = \|Tf\|_{p/2} \le c_\star^2 \delta^{-d} \|f\|_{p/2}$$

and inequality (4.17) follows.

**Lemma 4.8.** Let  $1 \le p < 2$ . Then there exists a constant c = c(p) > 0 such that for any  $\delta > 0$  and any pdf f supported in a ball  $B(x_0, R)$  with  $x_0 \in \mathcal{M}$  and  $R \ge \delta/2$  we have

(4.20) 
$$\left(\int_{\mathcal{M}} \left(\int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|f(u)d\mu(u)\right)^{p/2} d\mu(x)\right)^{\frac{1}{p}} \le c|B(x_0,R)|^{\frac{1}{p}-\frac{1}{2}}.$$

*Proof.* We split the region of integration  $\mathcal{M}$  into two:  $B(x_0, 2R)$  and  $\mathcal{M} \setminus B(x_0, 2R)$ .

Since  $1 \le p < 2$ , there exists  $1 < r < \infty$  such that  $\frac{p}{2} + \frac{1}{r} = 1$ . Applying Hölder's inequality we obtain

$$I := \int_{B(x_0,2R)} \left( \int_{B(x_0,R)} |\Phi(\delta\sqrt{L})(x,u)| f(u) d\mu(u) \right)^{p/2} d\mu(x) \\ \le \left( \int_{B(x_0,2R)} \int_{B(x_0,R)} |\Phi(\delta\sqrt{L})(x,u)| f(u) d\mu(u) d\mu(x) \right)^{p/2} |B(x_0,2R)|^{1/r}.$$

We now use Fubini's theorem, (4.13) with p = 1, the fact that  $\int_{\mathcal{M}} f(u)d\mu(u) = 1$ , and the doubling property (2.1) to obtain (4.21)

$$I \leq \left(\int_{B(x_0,R)} \|\Phi(\delta\sqrt{L})(\cdot,u)\|_1 f(u) d\mu(u)\right)^{p/2} |B(x_0,2R)|^{1/r} \leq c |B(x_0,R)|^{1-\frac{p}{2}}.$$

To estimate the integral over  $\mathcal{M} \setminus B(x_0, 2R)$  we observe that if  $u \in B(x_0, R)$  and  $x \in \mathcal{M} \setminus B(x_0, 2R)$ , then  $\rho(x, x_0) \leq \rho(x, u) + \rho(u, x_0) < \rho(x, u) + R < 2\rho(x, u)$ . Then by (3.5) and (4.12) we get

$$\begin{aligned} |\Phi(\delta\sqrt{L})(x,u)| &\leq c|B(x,\delta)|^{-1} \left(1 + \delta^{-1}\rho(x,u)\right)^{-k} \\ &\leq c|B(x_0,\delta)|^{-1} \left(1 + \delta^{-1}\rho(x,x_0)\right)^{-k+d}. \end{aligned}$$

Choose k > d(1 + 2/p). Then using (3.6) we obtain

$$\int_{\mathcal{M}\setminus B(x_0,2R)} \left( \int_{B(x_0,R)} |\Phi(\delta\sqrt{L})(x,u)| f(u) d\mu(u) \right)^{p/2} d\mu(x) 
(4.22) \leq c |B(x_0,\delta)|^{-p/2} \int_{\mathcal{M}} \left( 1 + \delta^{-1} \rho(x,x_0) \right)^{-(k-d)p/2} d\mu(x) \left( \int_{\mathcal{M}} f(u) d\mu(u) \right)^{p/2} 
\leq c |B(x_0,\delta)|^{1-p/2} \leq c |B(x_0,2R)|^{1-p/2} \leq c' |B(x_0,R)|^{1-p/2},$$

where we used that  $R \ge \delta/2$ , p < 2, and  $\int_{\mathcal{M}} f(u)d\mu(u) = 1$ . Inequality (4.20) follows by (4.21) and (4.22).

4.5. **Proof of Theorem 4.3 and Theorem 4.4.** We will only prove Theorem 4.3. Theorem 4.4 follows readily.

By the triangle inequality we obtain the standard decomposition of the risk as the sum of stochastic and bias terms:

(4.23) 
$$\mathbb{E}\|\widehat{\Phi}_{\delta} - f\|_{p} \leq \mathbb{E}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{p} + \|\Phi(\delta\sqrt{L})f - f\|_{p}.$$

For the estimation of the bias term  $\|\Phi(\delta\sqrt{L})f - f\|_p$  we will use estimate (4.15). We next focus on the estimation of the stochastic term  $\mathbb{E}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_p$ . In the case  $1 \leq p < \infty$ , using Jensen's inequality, we get

$$\mathbb{E}(\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{p}) \leq \left(\mathbb{E}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{p}^{p}\right)^{\frac{1}{p}}$$

$$(4.24) \qquad = \left(\int_{\mathcal{M}} \mathbb{E}\left|\frac{1}{n}\sum_{i=1}^{n}\Phi(\delta\sqrt{L})(x,X_{i}) - \Phi(\delta\sqrt{L})f(x)\right|^{p}d\mu(x)\right)^{\frac{1}{p}}$$

(i) Assume the pdf  $f \in B^s_{p\tau}(m)$  and let  $X \sim X_i$ . We first prove estimate (4.9) for p = 2. Clearly

$$\mathbb{E} \left| \frac{1}{n} \sum_{i=1}^{n} \Phi(\delta \sqrt{L})(x, X_i) - \Phi(\delta \sqrt{L}) f(x) \right|^2 \le \frac{1}{n} \mathbb{E} [\Phi(\delta \sqrt{L})(x, X)]^2$$
$$= \frac{1}{n} \int_{\mathcal{M}} |\Phi(\delta \sqrt{L})(x, u)|^2 f(u) d\mu(u).$$

This coupled with (4.24) yields

(4.25) 
$$\mathbb{E} \| \overline{\Phi}_{\delta} - \Phi(\delta \sqrt{L}) f \|_{2}$$

$$\leq \frac{1}{n^{1/2}} \Big( \int_{\mathcal{M}} \int_{\mathcal{M}} |\Phi(\delta \sqrt{L})(x, u)|^{2} f(u) d\mu(u) d\mu(x) \Big)^{\frac{1}{2}} \leq \frac{c}{(n\delta^{d})^{1/2}},$$

where we used (4.16) with p = 2. Combining (4.23), (4.15), and (4.25) we get

$$\mathbb{E}\|\widehat{\Phi}_{\delta} - f\|_2 \le \frac{c}{(n\delta^d)^{1/2}} + cm\delta^s.$$

With  $\delta = n^{-\frac{1}{2s+d}}$ , i.e.  $\delta^s = \frac{1}{(n\delta^d)^{1/2}}$ , this yields (4.9) when p = 2.

Let 2 . We will use the following version of Rosenthal's inequality that $can be derived for instance from [14], p. 245, inequality (C.5) with <math>\tau = \frac{P}{2} + 1 \le p + 1$ : If  $Y_1, \ldots, Y_n$  are i.i.d. random variables and  $Y_i \sim Y$ , then

(4.26) 
$$\mathbb{E} \left| \frac{1}{n} \sum_{i=1}^{n} Y_i - \mathbb{E}Y \right|^p \le \frac{(p+1)^p}{n^{p-1}} \mathbb{E} |Y|^p + \frac{p(p+1)^{p/2} e^{p/2+1}}{n^{p/2}} \left( \mathbb{E} |Y|^2 \right)^{p/2}.$$

We get

$$\mathbb{E} \left| \frac{1}{n} \sum_{i=1}^{n} \Phi(\delta \sqrt{L})(x, X_i) - \Phi(\delta \sqrt{L}) f(x) \right|^p \\ \leq \frac{c}{n^{p-1}} \mathbb{E} \left| \Phi(\delta \sqrt{L})(x, X) \right|^p + \frac{c}{n^{p/2}} \left( \mathbb{E} \left| \Phi(\delta \sqrt{L})(x, X) \right|^2 \right)^{p/2}$$

This and (4.24) imply

$$\mathbb{E}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{p} \leq \frac{c}{n^{1-1/p}} \Big(\int_{\mathcal{M}} \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^{p} f(u)d\mu(u)d\mu(x)\Big)^{1/p} + \frac{c}{n^{1/2}} \Big(\int_{\mathcal{M}} \Big(\int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^{2} f(u)d\mu(u)\Big)^{p/2} d\mu(x)\Big)^{1/p} = \frac{c}{(n\delta^{d})^{1/2}} + c(n\delta^{d})^{-1} \|f\|_{p/2},$$

where we used (4.16) and (4.16). Since  $1 \leq \frac{p}{2} < p$  and  $||f||_1 = 1$ , we obtain by interpolation

$$(4.28) ||f||_{\frac{p}{2}} \le ||f||_{1}^{\frac{1}{p-1}} ||f||_{p}^{\frac{p-2}{p-1}} = ||f||_{p}^{\frac{p-2}{p-1}} \le c||f||_{B_{p_{\tau}}^{s}}^{\frac{p-2}{p-1}} \le cm^{\frac{p-2}{p-1}}.$$

Here we also used Proposition 3.5 (iv).

Combining (4.27)-(4.28) with (4.23) and (4.15), and taking into account that  $\delta = n^{-\frac{1}{2s+d}}$ , i.e.  $\delta^s = \frac{1}{(n\delta^d)^{1/2}}$  we arrive at

(4.29) 
$$\mathbb{E}\|\widehat{\Phi}_{\delta} - f\|_{p} \leq \frac{c}{(n\delta^{d})^{1/2}} + cm\delta^{s} \leq c'n^{-\frac{s}{2s+d}}.$$

The proof of part (i) of the theorem is complete.

(ii) Let  $1 \leq p < 2$  and  $f \in B^s_{p\tau}(m, x_0, R)$ . We use Jensen's inequality and the fact that  $|\Phi(\delta\sqrt{L})(x, u)| \leq c|B(x, \delta)|^{-1} \leq c'\delta^{-d}$ , using (4.12) and (3.4), to obtain

$$\begin{split} \mathbb{E} \Big| \frac{1}{n} \sum_{i=1}^{n} \Phi(\delta\sqrt{L})(x, X_{i}) - \Phi(\delta\sqrt{L})f(x) \Big|^{p} \\ &\leq \left( \mathbb{E} \Big| \frac{1}{n} \sum_{i=1}^{n} \Phi(\delta\sqrt{L})(x, X_{i}) - \Phi(\delta\sqrt{L})f(x) \Big|^{2} \right)^{\frac{p}{2}} \\ &\leq \frac{1}{n^{\frac{p}{2}}} \left( \mathbb{E} |\Phi(\delta\sqrt{L})(x, X)|^{2} \right)^{\frac{p}{2}} \\ &= \frac{1}{n^{\frac{p}{2}}} \Big( \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x, u)|^{2}f(u)d\mu(u) \Big)^{\frac{p}{2}} \\ &\leq \frac{c}{(n\delta^{d})^{\frac{p}{2}}} \Big( \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x, u)|f(u)d\mu(u) \Big)^{\frac{p}{2}}. \end{split}$$

This and (4.24) lead to

$$\mathbb{E}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{p} \leq \frac{c}{(n\delta^{d})^{\frac{1}{2}}} \Big(\int_{\mathcal{M}} \Big(\int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|f(u)d\mu(u)\Big)^{\frac{p}{2}} d\mu(x)\Big)^{\frac{1}{p}}.$$

We now invoke Lemma 4.8 to obtain

$$\mathbb{E}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{p} \le \frac{c}{(n\delta^{d})^{\frac{1}{2}}}|B(x_{0},R)|^{\frac{1}{p}-\frac{1}{2}} \le \frac{c'}{(n\delta^{d})^{\frac{1}{2}}}.$$

Using this and (4.15) we complete the proof of (4.10) just as above in (4.29).

(iii) Assume the pdf  $f \in B^s_{\infty\tau}(m)$  and let q > 2 be arbitrary. Since by construction supp  $\Phi \subset [-1,1]$ , the function  $\widehat{\Phi}_{\delta}(x) - \Phi(\delta\sqrt{L})f(x)$  belongs to the spectral space  $\Sigma_{1/\delta}$ . Then by Proposition 3.2

$$\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{\infty} \le c_{\star}\delta^{-\frac{d}{q}}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{q}$$

where the constant  $c_{\star} > 1$  is independent of q. This along with Jensen's inequality and Fubini's theorem lead to

(4.30) 
$$\mathbb{E}\|\widehat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{\infty} \le c_{\star}\delta^{-\frac{d}{q}} \Big(\int_{\mathcal{M}} \mathbb{E}|\widehat{\Phi}_{\delta}(x,\cdot) - \Phi(\delta\sqrt{L})f(x)|^{q}d\mu(x)\Big)^{\frac{1}{q}}$$

We now apply Rosenthal's inequality (4.26) to obtain

$$\begin{split} \mathbb{E}|\widehat{\Phi}_{\delta}(x) &- \Phi(\delta\sqrt{L})f(x)|^{q} \\ &\leq \frac{(q+1)^{q}}{n^{q-1}} \mathbb{E}|\Phi(\delta\sqrt{L})(x,X)|^{q} + \frac{q(q+1)^{\frac{q}{2}}e^{\frac{q}{2}+1}}{n^{\frac{q}{2}}} \big(\mathbb{E}|\Phi(\delta\sqrt{L})(x,X)|^{2}\big)^{\frac{q}{2}} \\ &= \frac{(q+1)^{q}}{n^{q-1}} \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^{q} f(u) d\mu(u) \\ &+ \frac{q(q+1)^{\frac{q}{2}}e^{\frac{q}{2}+1}}{n^{\frac{q}{2}}} \Big(\int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^{2} f(u) d\mu(u)\Big)^{\frac{q}{2}}. \end{split}$$

This coupled with (4.30) and the fact that 1/q < 1 imply

$$\mathbb{E} \| \hat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f \|_{\infty} \leq c_{\star} \delta^{-\frac{d}{q}} \frac{q+1}{n^{1-\frac{1}{q}}} \Big( \int_{\mathcal{M}} \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^{q} f(u)d\mu(u)d\mu(x) \Big)^{\frac{1}{q}} \\ (4.31) + c_{\star} \delta^{-\frac{d}{q}} \frac{q^{\frac{1}{q}}(q+1)^{\frac{1}{2}} e^{\frac{1}{2}+\frac{1}{q}}}{n^{\frac{1}{2}}} \Big( \int_{\mathcal{M}} \Big( \int_{\mathcal{M}} |\Phi(\delta\sqrt{L})(x,u)|^{2} f(u)d\mu(u) \Big)^{\frac{q}{2}} d\mu(x) \Big)^{\frac{1}{q}} \\ \leq c_{\star} \delta^{-\frac{d}{q}} \Big( \frac{2c_{\star}q}{(n\delta^{d})^{1-\frac{1}{q}}} + \frac{e^{2}c_{\star}q^{1/2}}{(n\delta^{d})^{1/2}} \|f\|_{q/2}^{1/2} \Big),$$

where we used (4.16), (4.17), and the inequality  $q^{\frac{1}{q}}(q+1)^{\frac{1}{2}}e^{\frac{1}{2}+\frac{1}{q}} \leq e^2q^{1/2}$ , (q > 2). Observe that the constant  $c_{\star}$  above is from (4.13) and is independent of q.

By Proposition 3.5 (iii) it follows that  $f \in \mathbb{L}^{\infty}$  and since  $||f||_1 = 1$  we obtain

$$\|f\|_{q/2} \le \|f\|_{\infty}^{1-2/q} \|f\|_1 \le \left(c\|f\|_{B_{\infty\tau}^s}\right)^{1-2/q} \le (cm)^{1-2/q} \le cm+1.$$

Let  $n \ge e^2$  and choose  $q := \log n$ . By assumption  $\delta = \left(\frac{\log n}{n}\right)^{1/(2s+d)}$ . Now, it is easy to see that  $n^{1/q} = e, \ \delta^{-d/q} \le e, \ \delta^s = \frac{q^{1/2}}{(n\delta^d)^{1/2}} = \left(\frac{\log n}{n}\right)^{s/(2s+d)}$ , and

$$\frac{q}{(n\delta^d)^{1-1/q}} \le \frac{q}{(n\delta^d)^{3/4}} \le \frac{\log n}{n^{\frac{3s/2}{2s+d}}} \le c \Big(\frac{\log n}{n}\Big)^{s/(2s+d)} \quad \text{if} \quad n \ge e^2.$$

Putting all of the above together we obtain

(4.32) 
$$\mathbb{E}\|\hat{\Phi}_{\delta} - \Phi(\delta\sqrt{L})f\|_{\infty} \le c \left(\frac{\log n}{n}\right)^{s/(2s+d)}$$

If  $2 \le n < e^2$ , then estimate (4.32) follows readily from (4.31) with q = 2.

As before we use (4.32) and (4.15) to obtain (4.11). The proof of Theorem 4.4 is complete.

A closer examination of the above proof shows that the oracle inequalities from Theorem 4.3 are valid.  $\hfill \Box$ 

4.6. Linear wavelet density estimators. In this section we establish  $\mathbb{L}^{p}$ -error estimates for linear wavelet density estimators. Let  $\{\psi_{j\xi}\}, \{\tilde{\psi}_{j\xi}\}$  be the pair of dual frames described in Subsection 3.4. We adhere to the notation from Section 3.4.

For any  $j \ge 0$  and  $\xi \in \mathcal{X}_j$  we define the *empirical coefficient estimators* by

(4.33) 
$$\hat{\beta}_{j\xi} := \frac{1}{n} \sum_{i=1}^{n} \tilde{\psi}_{j\xi}(X_i).$$

Using this we define the *linear wavelet density estimator* by

(4.34) 
$$f^*(x) = \sum_{j=0}^J \sum_{\xi \in \mathcal{X}_j} \hat{\beta}_{j\xi} \psi_{j\xi}(x), \ x \in \mathcal{M},$$

where the parameter  $J = J(n) \in \mathbb{N}$  is selected so that the factor  $b^{-J}$  de facto behaves as a bandwidth. More precisely, we define J as the unique positive integer such that

(4.35) 
$$b^J \le n^{1/(2s+d)} < b^{J+1}.$$

It is easy to see that  $f^*$  can be written in the following way

(4.36) 
$$f^*(x) = \frac{1}{n} \sum_{i=1}^n \sum_{j=0}^J \sum_{\xi \in \mathcal{X}_j} \psi_{j\xi}(x) \tilde{\psi}_{j\xi}(X_i)$$
$$= \frac{1}{n} \sum_{i=1}^n \sum_{j=0}^J \Psi_j(\sqrt{L})(X_i, x) = \frac{1}{n} \sum_{i=1}^n \Psi_0(b^{-J}\sqrt{L})(X_i, x).$$

where we used (3.12) and (3.8).

Thus, this linear wavelet estimator is in fact a particular case of the linear estimators investigated in the previous subsection. This enables us to state the following upper bound theorem, which is an immediate consequence of Theorem 4.4.

**Theorem 4.9.** Let s > 0,  $0 < \tau \le \infty$ , m > 0,  $x_0 \in \mathcal{M}$  and R > 0. (i) If  $2 \le p < \infty$  and J is as in (4.35), then

(4.37) 
$$\sup_{f \in B^*_{a,c}(m)} \mathbb{E} \| f^* - f \|_p \le c n^{-s/(2s+d)},$$

where  $c = c(p, \tau, s, m) > 0$ .

(ii) If  $1 \le p < 2$  and J is as in (4.35), then

(4.38) 
$$\sup_{f \in B_{p\tau}^s(m,x_0,R)} \mathbb{E} \| f^* - f \|_p \le c n^{-s/(2s+d)},$$

where  $c = c(p, \tau, s, m, x_0, R) > 0$ .

(iii) If J is the unique integer satisfying  $b^{J} \leq \left(\frac{n}{\log n}\right)^{1/(2s+d)} < b^{J+1}$ , then

(4.39) 
$$\sup_{f \in B_{\infty\tau}^s(m)} \mathbb{E} \| f^* - f \|_{\infty} \le c \Big( \frac{\log n}{n} \Big)^{\frac{s}{2s+d}}$$

where  $c = c(\tau, s, m) > 0$ .

#### 5. Adaptive wavelet density estimation by thresholding

If we want to parallel the achievements obtained in density estimation theory in -say-  $[0,1]^d$ , one important feature is lacking: adaptation (i.e. obtaining -up to logarithmic factors- optimal rates of convergence without knowing the regularity). There are various techniques for this. For instance, Lepski's method (see [23, 12]) could be applied to our kernel estimators.

We choose to develop here nonlinear wavelet estimators, where we apply hard thresholding. This method has been developed in the classical case of  $\mathbb{R}$  in [6] and on the sphere in [14]. We will operate in the general setting described in Section 2. Unlike the case of the kernel or linear wavelet density estimates considered in the previous section, here we assume that the space  $\mathcal{M}$  is compact ( $\mu(\mathcal{M}) < \infty$ ) and all conditions C1–C5 (including the Ahlfors regularity condition C1A) are satisfied, see Section 2.

As before we assume that  $X_1, \ldots, X_n$   $(n \ge 2)$  are i.i.d. random variables with values on  $\mathcal{M}$  and with a common density function f with respect to the measure  $\mu$  on  $\mathcal{M}$ . Let  $X_j \sim X$ . We denote by  $\mathbb{E} = \mathbb{E}_f$  the expectation with respect to the probability measure  $\mathbb{P} = \mathbb{P}_f$ .

In addition, we assume that f is bounded. Denote

(5.1) 
$$A := \max\{\|f\|_{\infty}, 4\}$$
 and set  $\kappa := c_{\diamond}(8A)^{1/2}$ ,

where  $c_{\diamond} > 1$  is the constant from the norm bounds of the frame elements in (3.16).

We will utilize the pair of frames  $\{\psi_{j\xi}\}, \{\overline{\psi}_{j\xi}\}\$  described in §3.4. We adhere to the notation from §3.4. Recall that any  $f \in \mathbb{L}^p(\mathcal{M}, d\mu)$  has the frame decomposition

(5.2) 
$$f = \sum_{j=0}^{\infty} \sum_{\xi \in \mathcal{X}_j} \beta_{j\xi}(f) \psi_{j\xi}, \quad \beta_{j\xi}(f) := \langle f, \tilde{\psi}_{j\xi} \rangle \quad \text{(convergence in } \mathbb{L}^p\text{)}.$$

Assuming the pdf f fixed, we will use the abbreviated notation  $\beta_{j\xi} := \beta_{j\xi}(f)$ . We introduce two parameters depending on n:

(5.3) 
$$\lambda_n := \kappa \left(\frac{\log n}{n}\right)^{1/2}$$

and  $J_n$  uniquely defined by the following inequalities

(5.4) 
$$b^{J_n} \le \left(\frac{n}{\log n}\right)^{1/d} < b^{J_n+1}.$$

As in  $\S4.6$  we introduce the *empirical coefficient estimators* 

(5.5) 
$$\hat{\beta}_{j\xi} := \frac{1}{n} \sum_{i=1}^{n} \tilde{\psi}_{j\xi}(X_i), \quad j \ge 0, \ \xi \in \mathcal{X}_j.$$

We now define the hard threshold coefficient estimators by

(5.6) 
$$\hat{\beta}_{j\xi}^* := \hat{\beta}_{j\xi} \mathbb{1}_{\{|\hat{\beta}_{j\xi}| > 2\lambda_n\}}, \quad j \ge 0, \ \xi \in \mathcal{X}_j.$$

Then the wavelet threshold density estimator is defined by

(5.7) 
$$\hat{f}_n(x) := \sum_{0 \le j \le J_n} \sum_{\xi \in \mathcal{X}_j} \hat{\beta}_{j\xi}^* \psi_{j\xi}(x), \quad x \in \mathcal{M}$$

**Remark 5.1.** Note that the density estimator  $\hat{f}_n$  of the pdf f depends only on the number n of observations, the geometric constant  $c_{\diamond}$ , and the  $\mathbb{L}^{\infty}$ -norm of f.

We now state our main result on the adaptive wavelet threshold estimator defined above.

**Theorem 5.2.** Let  $1 \le r \le \infty$ ,  $0 < \tau \le \infty$ ,  $1 \le p < \infty$ , s > d/r, and m > 0. Then there exists a constant  $c = c(r, \tau, p, s, m) > 0$  such that in the setting described above and with  $\hat{f}_n$  from (5.7) we have: (i)

(5.8)  $\sup_{f \in B^s_{r\tau}(m)} \mathbb{E} \| \hat{f}_n - f \|_{\infty} \le c \left( \frac{\log n}{n} \right)^{\frac{s - \frac{d}{r}}{2[s - d(\frac{1}{r} - \frac{1}{2})]}}.$ 

(ii) In the regular case  $s \ge \frac{dp}{2} \left(\frac{1}{r} - \frac{1}{p}\right)$ 

(5.9) 
$$\sup_{f \in B^s_{r\tau}(m)} \mathbb{E} \|\hat{f}_n - f\|_p \le c \log n \left(\frac{\log n}{n}\right)^{\frac{2s+d}{2s+d}}$$

(iii) In the sparse case  $s < \frac{dp}{2} \left( \frac{1}{r} - \frac{1}{p} \right)$ 

(5.10) 
$$\sup_{f \in B_{r\tau}^s(m)} \mathbb{E} \| \hat{f}_n - f \|_p \le c \log n \left( \frac{\log n}{n} \right)^{\frac{s - d(\frac{1}{r} - \frac{1}{p})}{2[s - d(\frac{1}{r} - \frac{1}{2})]}}.$$

Remark 5.3. Several observations are in order:

(a) The assumption s > d/r leads to  $||f||_{\infty} \leq c||f||_{B^s_{r\tau}} \leq cm$ , by Proposition 3.5 (iii). In addition in the sparse case it implies p > 2.

(b) The geometry of the setting is represented by the dimension d. Note that the exponents of  $\frac{\log n}{n}$  are the same as in the case of the sphere [1].

(c) In the regular case (modulo the logarithmic terms) we have the same rate of convergence  $n^{-s/(2s+d)}$  as in the case of the linear wavelet estimator.

(d) Just as in the case of kernel density estimators (see Remark 4.5) we note that since we assume here all conditions C1-C5 (including C1A) it would not be a problem to obtain lower bounds matching up to logarithmic terms the rates established above by a direct adaptation of the proof of the lower bounds in the case of the sphere from [1].

5.1. **Preparation for the proof of Theorem 5.2.** We first recall the classical *Bernstein inequality* (see e.g. [25]): Let  $Y_1, \ldots, Y_n$  be independent random variables such that  $\mathbb{E}Y_i = 0$ ,  $\mathbb{E}Y_i^2 \leq \sigma^2$ , and  $|Y_i| \leq M$ ,  $i = 1, \ldots, n$ . Then for any v > 0

$$(5.11) \quad \mathbb{P}\Big(\Big|\frac{1}{n}\sum_{i=1}^{n}Y_i\Big| \ge v\Big) \le 2\exp\Big(-\frac{nv^2}{2(\sigma^2 + Mv/3)}\Big) \le 2\exp\Big(-\frac{nv^2}{4\sigma^2} \wedge \frac{3nv}{4M}\Big)$$
$$= 2\exp\Big(-\frac{nv^2}{2\sigma^2}\Big)\mathbb{1}_{\{v \le \frac{3\sigma^2}{M}\}} + 2\exp\Big(-\frac{3nv}{4M}\Big)\mathbb{1}_{\{v > \frac{3\sigma^2}{M}\}}.$$

We next use Rosenthal's inequalities (4.26) to derive several useful estimates in the current setting. Clearly,

$$\beta_{j\xi} := \int_{\mathcal{M}} \tilde{\psi}_{j\xi}(x) f(x) d\mu(x) = \mathbb{E}(\tilde{\psi}_{j\xi}(X))$$

and using (3.16) we obtain

(5.12) 
$$\mathbb{E}|\tilde{\psi}_{j\xi}(X) - \beta_{j\xi}|^2 \le \mathbb{E}|\tilde{\psi}_{j\xi}(X)|^2 = \int_{\mathcal{M}} f(x)|\tilde{\psi}_{j\xi}(x)|^2 d\mu(x) \le c_{\diamond}^2 A$$

and

(5.13) 
$$|\tilde{\psi}_{j\xi}(X) - \beta_{j\xi}| \le 2 \|\tilde{\psi}_{j\xi}\|_{\infty} \le 2c_{\diamond} b^{jd/2}.$$

Also

$$\mathbb{E}|\tilde{\psi}_{j\xi}(X) - \beta_{j\xi}|^p \le 2^p \mathbb{E}|\tilde{\psi}_{j\xi}(X)|^p = 2^p \int_{\mathcal{M}} f(x)|\tilde{\psi}_{j\xi}(x)|^p dx \mu(x) \le (2c_\diamond)^p A b^{jd(\frac{p}{2}-1)}.$$

Therefore, using (5.11)

(5.14) 
$$\mathbb{P}\left(|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda\right) \le 2\exp\left(-\frac{n\lambda^2}{4c_{\diamond}^2 A} \wedge \frac{3n\lambda}{8c_{\diamond} b^{jd/2}}\right).$$

From this with the notation

(5.15) 
$$\mu_j := \frac{3}{2} c_\diamond A b^{-jd/2}$$

we obtain

(5.16) 
$$\mathbb{P}\left(|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda\right) \le 2 \exp\left(-\frac{n\lambda^2}{4c_{\diamond}^2 A}\right) \quad \text{if} \quad 0 \le \lambda \le \mu_j,$$

and

(5.17) 
$$\mathbb{P}(|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda) \le 2 \exp\left(-\frac{3n\lambda}{8c_{\diamond}b^{jd/2}}\right) \quad \text{if} \quad \lambda \ge \mu_j.$$

In particular, if  $0 \leq j \leq J_n$ , then

(5.18) 
$$\mathbb{P}(|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n) \le \frac{2}{n^2}$$

Now, by (4.26), for any  $p \ge 2$  there exists c = c(p) > 0 such that

(5.19) 
$$\mathbb{E}|\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \le c \Big(\frac{c_{\diamond}^2 A}{n}\Big)^{p/2} \Big(1 + 2^p \Big(\frac{b^{jd}}{nA}\Big)^{p/2-1}\Big).$$

Moreover, for any  $j \geq 0$  such that  $b^{jd} \leq nA$ 

(5.20) 
$$\mathbb{E}|\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \le c \left(\frac{c_{\diamond}^2 A}{n}\right)^{p/2}.$$

On the other hand, by Jensen inequality, for any 0

(5.21) 
$$\mathbb{E}|\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \le \left(\frac{cc_0^2 A}{n}\right)^{p/2}.$$

The following two lemmas will be instrumental in the proof of Theorem 5.2.

**Lemma 5.4.** For any  $n \ge 2, 0 \le j \le n$ , and  $\xi \in \mathcal{X}_j$  we have

$$(5.22) \quad I_{j\xi} := \int_0^\infty \mathbb{P}\Big(|\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbbm{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} > \lambda\Big) d\lambda \le cA^{1/2} \Big(\frac{\log n}{n}\Big)^{1/2} \frac{1}{n^2}.$$
  
where  $c = c(c_\diamond) > 0.$ 

*Proof.* We will use the following well known inequality:

(5.23) 
$$\int_{a}^{\infty} e^{-K\frac{\lambda^{2}}{2}} d\lambda \leq \frac{e^{-\frac{Ka^{2}}{2}}}{K^{1/2}} \Big( (\pi/2)^{1/2} \wedge \frac{1}{aK^{1/2}} \Big), \quad K, a > 0$$

We split the integral in (5.22) into three:

$$I_{j\xi} = \left(\int_0^{\lambda_n} + \int_{\lambda_n}^{\mu_j} + \int_{\mu_j}^{\infty}\right) \mathbb{P}\left(\left|\hat{\beta}_{j\xi} - \beta_{j\xi}\right| \mathbb{1}_{\left\{\left|\hat{\beta}_{j\xi} - \beta_{j\xi}\right| > \lambda_n\right\}} > \lambda\right) d\lambda =: S_1 + S_2 + S_3.$$

From the definitions in (5.3) and (5.15) it readily follows that  $\mu_j \ge \lambda_n$ .

To estimate  $S_1$  we use (5.16) and the definitions of  $\lambda_n$ ,  $\kappa$  in (5.3), (5.1). We get

$$S_{1} \leq \lambda_{n} \mathbb{P}(|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_{n}) \leq \lambda_{n} 2 \exp\left(-\frac{n\lambda_{n}^{2}}{4c_{\diamond}^{2}A}\right)$$
$$\leq 2\kappa \left(\frac{\log n}{n}\right)^{1/2} n^{-\frac{\kappa^{2}}{4c_{\diamond}^{2}A}} \leq 2\kappa \left(\frac{\log n}{n}\right)^{1/2} n^{-2},$$

where we used that  $\frac{\kappa^2}{4c^2A} = 2$ . For  $S_2$  we use inequality (5.23) and again (5.16) to obtain

$$S_{2} \leq \int_{\lambda_{n}}^{\infty} 2 \exp\left(-\frac{n\lambda^{2}}{4c_{\diamond}^{2}A}\right) d\lambda \leq 2(\pi/2)^{1/2} \left(\frac{2c_{\diamond}^{2}A}{n}\right)^{1/2} \exp\left(-\frac{n\lambda_{n}^{2}}{4c_{\diamond}^{2}A}\right) \\ \leq 6c_{\diamond}A^{1/2}n^{-1/2}n^{-\frac{\kappa^{2}}{4c_{\diamond}^{2}A}} = 6c_{\diamond}A^{1/2}n^{-5/2}.$$

We now estimate  $S_3$ . Using (5.17) we obtain

$$S_{3} \leq \int_{\mu_{j}}^{\infty} 2 \exp\left(-\frac{3n\lambda}{8c_{\diamond}b^{jd/2}}\right) d\lambda = \frac{16c_{\diamond}b^{j\frac{d}{2}}}{3n} \exp\left(-\frac{3n\mu_{j}}{8c_{\diamond}b^{jd/2}}\right)$$
$$= \frac{16c_{\diamond}b^{j\frac{d}{2}}}{3n} \exp\left(-\frac{9A}{16}nb^{-jd}\right) \leq \frac{6c_{\diamond}}{(n\log n)^{1/2}}n^{-\frac{9A}{16}} \leq \frac{6c_{\diamond}}{(n\log n)^{1/2}}n^{-9/4},$$

where for the second equality we used the definition of  $\mu_j$  in (5.15), for the former inequality we used that  $b^{-jd} \ge b^{-J_nd} = \frac{\log n}{n}$ , and for the last inequality that  $A \ge 4$ . Putting the above estimates for  $S_1, S_2$ , and  $S_3$  together we arrive at

$$I_{j\xi} \le 2\kappa \Big(\frac{\log n}{n}\Big)^{1/2} n^{-2} + 6c_{\diamond} A^{1/2} n^{-5/2} + \frac{6c_{\diamond} n^{-9/4}}{(n\log n)^{1/2}} \le cA^{1/2} \Big(\frac{\log n}{n}\Big)^{1/2} n^{-2}$$
  
claimed.

as claimed.

**Lemma 5.5.** Let  $\mathcal{F}$  be a finite family of functions  $\phi : \mathbb{R} \to \mathbb{R}$  with  $N := \operatorname{card}(\mathcal{F})$ and let X be a random variable. Assume

(5.24) 
$$\sup_{\phi \in \mathcal{F}} \|\phi\|_{\infty} \le M/2, \quad and \quad \sup_{\phi \in \mathcal{F}} \mathbb{E}(\phi^2(X)) \le \sigma^2.$$

Let  $X_1, \ldots, X_n$  be i.i.d. random variables and  $X_i \sim X$ . Then

(5.25) 
$$\mathbb{E}\left[\sup_{\phi\in\mathcal{F}}\left|\frac{1}{n}\sum_{i=1}^{n}\left(\phi(X_{i})-\mathbb{E}(\phi(X))\right)\right|\right] \leq 4\sigma\left(\frac{\log(2N)}{n}\right)^{1/2}+4M\frac{\log(2N)}{n}.$$

*Proof.* By Bernstein's inequality (5.11) it follows that for any  $\phi \in \mathcal{F}$ 

(5.26) 
$$\mathbb{P}\Big(\Big|\frac{1}{n}\sum_{i=1}^{n}\phi(X_i) - \mathbb{E}[\phi(X)]\Big| \ge \lambda\Big) \le 2\exp\Big(-\frac{n\lambda^2}{4\sigma^2} \wedge \frac{3n\lambda}{4M}\Big)$$

Let  $\lambda_0 := \frac{3\sigma^2}{M}$  and  $Z_\phi := |\frac{1}{n} \sum_{i=1}^n (\phi(X_i) - \mathbb{E}[\phi(X)])|$ . Then

$$\sup_{\phi \in \mathcal{F}} Z_{\phi} = \sup_{\phi \in \mathcal{F}} \{ Z_{\phi} \mathbb{1}_{\{ Z_{\phi} \le \lambda_0 \}} + Z_{\phi} \mathbb{1}_{\{ Z_{\phi} > \lambda_0 \}} \} \le \sup_{\phi \in \mathcal{F}} Z_{\phi} \mathbb{1}_{\{ Z_{\phi} \le \lambda_0 \}} + \sup_{\phi \in \mathcal{F}} Z_{\phi} \mathbb{1}_{\{ Z_{\phi} > \lambda_0 \}}$$

From (5.26) it follows that for any  $\lambda > 0$ 

$$\mathbb{P}(Z_{\phi}\mathbb{1}_{\{Z_{\phi} \leq \lambda_{0}\}} > \lambda) \leq 2e^{-\frac{n\lambda^{2}}{4\sigma^{2}}} \quad \text{and} \quad \mathbb{P}(Z_{\phi}\mathbb{1}_{\{Z_{\phi} > \lambda_{0}\}} > \lambda) \leq 2e^{-\frac{3n\lambda}{4M}}.$$

We use these inequalities and (5.23) to obtain

$$\begin{split} \mathbb{E}\Big[\sup_{\phi\in\mathcal{F}}Z_{\phi}\Big] &\leq \mathbb{E}\Big[\sup_{\phi\in\mathcal{F}}Z_{\phi}\mathbb{1}_{\{Z_{\phi}\leq\lambda_{0}\}}\Big] + \mathbb{E}\Big[\sup_{\phi\in\mathcal{F}}Z_{\phi}\mathbb{1}_{\{Z_{\phi}>\lambda_{0}\}}\Big] \\ &= \int_{0}^{\infty}\mathbb{P}\Big(\sup_{\phi\in\mathcal{F}}Z_{\phi}\mathbb{1}_{\{Z_{\phi}\leq\lambda_{0}\}}>\lambda\Big)d\lambda + \int_{0}^{\infty}\mathbb{P}\Big(\sup_{\phi\in\mathcal{F}}Z_{\phi}\mathbb{1}_{\{Z_{\phi}>\lambda_{0}\}}>\lambda\Big)d\lambda \\ &\leq a + \sum_{\phi\in\mathcal{F}}\int_{a}^{\infty}\mathbb{P}\Big(Z_{\phi}\mathbb{1}_{\{Z_{\phi}\leq\lambda_{0}\}}>\lambda\Big)d\lambda + b + \sum_{\phi\in\mathcal{F}}\int_{b}^{\infty}\mathbb{P}\Big(Z_{\phi}\mathbb{1}_{\{Z_{\phi}>\lambda_{0}\}}>\lambda\Big)d\lambda \\ &\leq a + N\int_{a}^{\infty}2e^{-\frac{n\lambda^{2}}{4\sigma^{2}}}d\lambda + b + N\int_{b}^{\infty}2e^{-\frac{3n\lambda}{4M}}d\lambda \\ &\leq a + 2N\frac{2\sigma^{2}}{na}e^{-\frac{na^{2}}{4\sigma^{2}}} + b + 2N\frac{4M}{3n}e^{-\frac{3nb}{4M}}. \end{split}$$

We now optimize with respect to a and b by taking  $2Ne^{-\frac{na^2}{4\sigma^2}} = 1$  and  $2Ne^{-\frac{3nb}{4M}} = 1$ . We obtain

$$\begin{split} \mathbb{E}\Big[\sup_{\phi\in\mathcal{F}}Z_{\phi}\Big] &\leq a + \frac{2\sigma^2}{na} + b + \frac{4M}{3n} \\ &= 2\sigma\Big(\frac{\log(2N)}{n}\Big)^{1/2} + \frac{\sigma}{(n\log(2N))^{1/2}} + \frac{4M}{3n}\log(2N) + \frac{4M}{3n} \\ &= \sigma\Big(\frac{\log(2N)}{n}\Big)^{1/2}\Big(2 + \frac{1}{\log(2N)}\Big) + \frac{4M}{n}\frac{1 + \log(2N)}{3} \\ &\leq 4\sigma\Big(\frac{\log(2N)}{n}\Big)^{1/2} + \frac{4M}{n}\log(2N), \end{split}$$
 where we used that  $2\log(2N) \geq \log 4 \geq 1$ . The proof of (5.25) is complete.

**Remark 5.6.** Note that by assumption (5.24) it follows that for any  $\phi \in \mathcal{F}$ 

(5.27) 
$$|\phi(X) - \mathbb{E}(\phi(X))| \le M \quad and \quad \mathbb{E}|\phi(X) - \mathbb{E}(\phi(X))|^2 \le \sigma^2.$$

5.2. Proof of Theorem 5.2. We will carry out this proof in several steps.

First, assuming that  $f \in \mathbb{L}^p$ ,  $1 \le p \le \infty$ , we use (3.13)-(3.14) and (5.7) to write

$$\hat{f}_n - f = \sum_{0 \le j \le J_n} \sum_{\xi \in \mathcal{X}_j} \left( \hat{\beta}_{j\xi} \mathbb{1}_{\{\hat{\beta}_{j\xi} > 2\lambda_n\}} - \beta_{j\xi} \right) \psi_{j\xi} + \sum_{j > J_n} \Psi_j(\sqrt{L}) f$$

which implies an estimate on the risk as a sum of stochastic and bias terms:

$$\mathbb{E}\|\hat{f}_n - f\|_p \le \sum_{0 \le j \le J_n} \mathbb{E}\left\|\sum_{\xi \in \mathcal{X}_j} (\hat{\beta}_{j,\xi}^* - \beta_{j\xi})\psi_{j,\xi}\right\|_p + \left\|\sum_{j > J_n} \Psi_j(\sqrt{L})f\right\|_p.$$

5.2.1. Estimation of the bias term. By the triangle inequality we get

$$\left\|\sum_{j>J_n}\Psi_j(\sqrt{L})f\right\|_p \le \sum_{j>J_n} \|\Psi_j(\sqrt{L})f\|_p.$$

Two cases are to be considered here. Let  $r \leq p \leq \infty$  and set  $s_1 := s - d(\frac{1}{r} - \frac{1}{p}) > 0$ . Then using Proposition 3.5 (ii), (3.22), and (5.4) we obtain

$$\begin{split} \sum_{j>J_n} \|\Psi_j(\sqrt{L})f\|_p &\leq \|f\|_{B_{p\tau}^{s_1}} \sum_{j>J_n} b^{-js_1} \\ &\leq c \|f\|_{B_{r\tau}^s} b^{-J_n s_1} \leq c \|f\|_{B_{r\tau}^s} \Big(\frac{\log n}{n}\Big)^{\frac{s}{d} - (\frac{1}{r} - \frac{1}{p})}. \end{split}$$

Let  $p < r \leq \infty$ . By Hölder's inequality (using  $\mu(M) < \infty$ ) and (3.22) we get

$$\sum_{j>J_n} \|\Psi_j(\sqrt{L})f\|_p \le c \sum_{j>J_n} \|\Psi_j(\sqrt{L})f\|_r \le c \|f\|_{B^s_{r\tau}} \sum_{j\ge J_n} b^{-js} \le c \|f\|_{B^s_{r\tau}} \left(\frac{\log n}{n}\right)^{\frac{s}{d}}.$$

Therefore, we have the following estimate for the bias

(5.28) 
$$\left\| \sum_{j>J_n} \Psi_j(\sqrt{L}) f \right\|_p \le c \|f\|_{B^s_{r\tau}} \left(\frac{\log n}{n}\right)^{\frac{s}{d} - (\frac{1}{r} - \frac{1}{p})_+}.$$

We next show that the rate from above is negligible compared to the rates in (5.8)-(5.10). First, if  $p = \infty$ , we have to verify that

$$\frac{s}{d} - \frac{1}{r} > \frac{s - \frac{d}{r}}{2(s - d(\frac{1}{r} - \frac{1}{2}))} = \left(\frac{s}{d} - \frac{1}{r}\right)\frac{d}{2(s - \frac{d}{r}) + d}$$

But this is obvious as  $s > \frac{d}{r}$ . If  $s \le \frac{dp}{2}(\frac{1}{r} - \frac{1}{p})$ , we have 2 and <math>r < p as  $s > \frac{d}{r}$ . We have to verify that

$$\frac{s}{d} - \left(\frac{1}{r} - \frac{1}{p}\right) > \frac{s - d(\frac{1}{r} - \frac{1}{p})}{2(s - d(\frac{1}{r} - \frac{1}{2}))} = \left(\frac{s}{d} - \left(\frac{1}{r} - \frac{1}{p}\right)\right) \frac{d}{2(s - \frac{d}{r}) + d}$$

which is obvious. If  $s > \frac{dp}{2}(\frac{1}{r} - \frac{1}{p})$  and  $p \le r$ , we have to show that  $\frac{s}{d} > \frac{s}{2s+d}$ , which again is

obvious. If  $s > \frac{dp}{2}(\frac{1}{r} - \frac{1}{p}) > 0$  (hence r < p) we have to verify that

$$\frac{s}{d} - \left(\frac{1}{r} - \frac{1}{p}\right) > \frac{s}{2s+d} \quad \Longleftrightarrow \quad \frac{s}{d} - \frac{s}{2s+d} > \frac{1}{r} - \frac{1}{p}.$$

In fact, we have

$$s > \frac{dp}{2} \left(\frac{1}{r} - \frac{1}{p}\right) \lor \frac{d}{r} =: a(p, r).$$

As the function  $s \mapsto \frac{s}{d} - \frac{s}{2s+d}$  is strictly increasing, we just need to show that  $\frac{a(p,r)}{d} - \frac{a(p,r)}{2s+d} \ge (\frac{1}{r} - \frac{1}{p})$ . However,

$$\frac{dp}{2}\left(\frac{1}{r} - \frac{1}{p}\right) \le \frac{d}{r} \quad \Longleftrightarrow \quad p \le r+2 \quad \Longleftrightarrow \quad \frac{d}{r} - \frac{d}{2s+d} \ge \frac{1}{r} - \frac{1}{p}$$

and

$$r+2 \frac{1}{r} - \frac{1}{p},$$

as it can be easily verified.

From above it follows that the rate in (5.28) is faster than the rates in (5.8)-(5.10).

5.2.2. Evaluation of the stochastic term. Note that by (3.17) we have (5.29)

$$\mathbb{E}\Big\|\sum_{\xi\in\mathcal{X}_j}(\hat{\beta}_{j\xi}^*-\beta_{j\xi})\psi_{j\xi}\Big\|_p \le cb^{jd(\frac{1}{2}-\frac{1}{p})}\mathbb{E}\Big[\Big(\sum_{\xi\in\mathcal{X}_j}|\hat{\beta}_{j\xi}^*-\beta_{j\xi}|^p\Big)^{1/p}\Big], \quad 1\le p<\infty,$$

and this holds with the usual modification when  $p = \infty$ .

## 30 G. CLEANTHOUS, A. GEORGIADIS, G. KERKYACHARIAN, P. PETRUSHEV, D. PICARD

To estimate the stochastic term we will use the following representation

$$\begin{split} \beta_{j\xi}^* - \beta_{j\xi} &= (\beta_{j\xi} - \beta_{j\xi}) \mathbb{1}_{\{|\hat{\beta}_{j\xi}| > 2\lambda_n\}} - \beta_{j\xi} \mathbb{1}_{\{|\hat{\beta}_{j\xi}| \le 2\lambda_n\}} \\ &= (\hat{\beta}_{j\xi} - \beta_{j\xi}) \mathbb{1}_{\{|\hat{\beta}_{j\xi}| > 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| \le \lambda_n\}} \\ &+ (\hat{\beta}_{j\xi} - \beta_{j\xi}) \mathbb{1}_{\{|\hat{\beta}_{j\xi}| > 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \\ &- \beta_{j\xi} \mathbb{1}_{\{|\hat{\beta}_{j\xi}| \le 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| \le 3\lambda_n\}} \\ &- \beta_{j\xi} \mathbb{1}_{\{|\hat{\beta}_{j\xi}| \le 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| > 3\lambda_n\}}. \end{split}$$

In the case when  $1 \le p < \infty$ , we use this and (5.29) to write

(5.30) 
$$\sum_{0 \le j \le J_n} \mathbb{E} \left\| \sum_{\xi \in \mathcal{X}_j} \left( \hat{\beta}_{j\xi} \mathbb{1}_{\{ | \hat{\beta}_{j\xi} | > 2\lambda_n \}} - \beta_{j\xi} \right) \psi_{j\xi} \right\|_p \le c \left( I + II + III + IV \right),$$

where

$$\begin{split} I &:= \sum_{0 \le j \le J_n} b^{-jd(\frac{1}{p} - \frac{1}{2})} \mathbb{E} \Big[ \Big( \sum_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \Big] \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big)^{\frac{1}{p}}, \\ II &:= \sum_{0 \le j \le J_n} b^{-jd(\frac{1}{p} - \frac{1}{2})} \mathbb{E} \Big[ \Big( \sum_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \mathbb{1}_{\{|\hat{\beta}_{j\xi}| > 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \Big)^{\frac{1}{p}} \Big], \\ III &:= \sum_{0 \le j \le J_n} b^{-jd(\frac{1}{p} - \frac{1}{2})} \mathbb{E} \Big[ \Big( \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^p \mathbb{1}_{\{|\hat{\beta}_{j\xi}| \le 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| \le 3\lambda_n\}} \Big)^{\frac{1}{p}} \Big] \\ IV &:= \sum_{0 \le j \le J_n} b^{-jd(\frac{1}{p} - \frac{1}{2})} \mathbb{E} \Big[ \Big( \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^p \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big)^{\frac{1}{p}} \Big]. \end{split}$$

In the case  $p = \infty$  we have

(5.31) 
$$\sum_{0 \le j \le J_n} \mathbb{E} \left\| \sum_{\xi \in \mathcal{X}_j} \left( \hat{\beta}_{j\xi} \mathbb{1}_{\{ |\hat{\beta}_{j\xi}| > 2\lambda_n \}} - \beta_{j\xi} \right) \psi_{j\xi} \right\|_{\infty} \le c \left( I' + II' + III' + IV' \right),$$

where

$$I' := \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} (|\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big],$$
  

$$II' := \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi}| > 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \Big],$$
  

$$III' := \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi}| \le 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j\xi}| \le 3\lambda_n\}} \Big],$$
  

$$IV' := \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big].$$

Estimation of I and I'. As card $(\mathcal{X}_j) \leq cb^{jd}$  by (3.9), we derive

$$I \leq \sum_{0 \leq j \leq J_n} b^{-jd(\frac{1}{p} - \frac{1}{2})} (\operatorname{card}(\mathcal{X}_j))^{\frac{1}{p}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big]$$
$$\leq c \sum_{0 \leq j \leq J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big] = cI'.$$

Now, in light of Lemma 5.4 we obtain

$$\begin{split} I &\leq cI' = c \sum_{0 \leq j \leq J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} (|\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big] \\ &\leq c \sum_{0 \leq j \leq J_n} b^{j\frac{d}{2}} \sum_{\xi \in \mathcal{X}_j} \mathbb{E} \Big[ |\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} \Big] \\ &= c \sum_{0 \leq j \leq J_n} b^{j\frac{d}{2}} \sum_{\xi \in \mathcal{X}_j} \int_0^\infty \mathbb{P} \Big( |\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n\}} > \lambda \Big) d\lambda \\ &\leq c \Big( \frac{\log n}{n} \Big)^{1/2} \frac{1}{n^2} \sum_{0 \leq j \leq J_n} b^{3j\frac{d}{2}} \leq c \Big( \frac{\log n}{n} \Big)^{1/2} \frac{1}{n^2} b^{3J_n\frac{d}{2}} = \frac{c}{n\log n}. \end{split}$$

Therefore,

(5.32) 
$$I \le cI' \le \frac{c}{n\log n},$$

and hence the terms I and I' are negligible compared to the rates in (5.8)-(5.10). Estimation of IV and IV'. Observe that

$$IV' = \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi} - \beta_{j,\xi}| > \lambda_n\}} \Big]$$
$$\leq \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \sup_{\xi \in \mathcal{X}_j} |\beta_{j\xi}| \sum_{\xi \in \mathcal{X}_j} \mathbb{P}(|\hat{\beta}_{j\xi} - \beta_{j\xi}| > \lambda_n) =: IV''.$$

On the other hand, Jensen's inequality and the fact that  $\operatorname{card}(\mathcal{X}_j) \leq c b^{jd}$  yield

$$IV \leq \sum_{0 \leq j \leq J_n} b^{j\frac{d}{2}} b^{-j\frac{d}{p}} \left( \mathbb{E} \left[ \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^p \mathbb{1}_{\{|\hat{\beta}_{j,\xi} - \beta_{j,\xi}| > \lambda_n\}} \right] \right)^{\frac{1}{p}}$$
$$\leq \sum_{0 \leq j \leq J_n} b^{j\frac{d}{2}} b^{-j\frac{d}{p}} \mathbb{E} \left[ \sup_{\xi \in \mathcal{X}_j} |\beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j,\xi} - \beta_{j,\xi}| > \lambda_n\}} \right] (\operatorname{card}(\mathcal{X}_j)^{\frac{1}{p}}$$
$$\leq c \sum_{0 \leq j \leq J_n} b^{j\frac{d}{2}} \mathbb{E} \left[ \sup_{\xi \in \mathcal{X}_j} |\beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j,\xi} - \beta_{j,\xi}| > \lambda_n\}} \right] \leq c I V''.$$

Now, as  $\mathbb{P}(|\hat{\beta}_{j,\xi} - \beta_{j,\xi}| > \lambda_n) \le 2n^{-2}, \ 0 \le j \le J_n$ , by (5.18), and (5.33)  $|\beta_{j\xi}| \le cb^{-j(s+d(\frac{1}{2}-\frac{1}{r}))} \|f\|_{B^s_{r\tau}}, \quad \xi \in \mathcal{X}_j$ 

by (3.26), we derive

$$\begin{split} IV'' &\leq cn^{-2} \sum_{0 \leq j \leq J_n} b^{jd/2} \sup_{\xi \in \mathcal{X}_j} |\beta_{j\xi}| \operatorname{card}(\mathcal{X}_j) \\ &\leq cn^{-2} \|f\|_{B^s_{r\tau}} \sum_{0 \leq j \leq J_n} b^{j\frac{3d}{2}} b^{-js} b^{jd(\frac{1}{r} - \frac{1}{2})} \\ &\leq cn^{-2} \|f\|_{B^s_{r\tau}} b^{J_n d} \sum_{j=0}^{\infty} b^{-j(s - \frac{d}{r})} \\ &\leq cn^{-2} \|f\|_{B^s_{r\tau}} \frac{n}{\log n} = c \|f\|_{B^s_{r\tau}} \frac{1}{n \log n}. \end{split}$$

Above we also used the fact that s > d/r. Therefore,

$$(5.34) IV, IV' \le \frac{c}{n\log n},$$

and hence the terms IV and IV' are also negligible.

Estimation of II' and III'. We first estimate III'. Using (5.33) we get

$$III' \leq \sum_{0 \leq j \leq J_n} \sup_{\xi \in \mathcal{X}_j} (b^{j\frac{d}{2}} |\beta_{j\xi}|) \wedge (3b^{j\frac{d}{2}} \lambda_n)$$
$$\leq c \sum_{0 \leq j \leq J_n} (b^{-j(s-\frac{d}{r})} ||f||_{B^s_{r\tau}}) \wedge (3b^{j\frac{d}{2}} \lambda_n)$$

We now introduce a new parameter  $J_n^\prime$  by the identity:

$$b^{-J'_n(s-\frac{d}{r})} \|f\|_{B^s_{r\tau}} = b^{J'_n \frac{d}{2}} \lambda_n$$

Hence,

$$b^{J'_n(s-d(\frac{1}{r}-\frac{1}{2}))} = \frac{\|f\|_{B^s_{r\tau}}}{\lambda_n} = \frac{\|f\|_{B^s_{r\tau}}}{\left(\log n\right)^{\frac{1}{2}}}.$$

Then we obtain, taking into account that s > d/r,

$$\begin{split} III' &\leq c\lambda_n \sum_{0 \leq j < J'_n} b^{j\frac{d}{2}} + c \|f\|_{B^s_{r\tau}} \sum_{J'_n \leq j < J_n} b^{-j(s-\frac{d}{r})} \\ &\leq c\lambda_n b^{J'_n\frac{d}{2}} + c(s,r) \|f\|_{B^s_{r\tau}} b^{-J'_n(s-\frac{d}{r})} \\ &\leq c \|f\|_{B^s_{r\tau}} b^{-J'_n(s-\frac{d}{r})} \leq c \|f\|_{B^s_{r\tau}}^{\frac{d}{s-d(\frac{1}{r}-\frac{1}{2})}} \left(\kappa \sqrt{\frac{\log n}{n}}\right)^{\frac{s-\frac{d}{r}}{s-d(\frac{1}{r}-\frac{1}{2})}} \\ &\leq c(s,r,\kappa) \|f\|_{B^s_{r\tau}}^{\frac{d}{2(s-d(\frac{1}{r}-\frac{1}{2}))}} \left(\frac{\log n}{n}\right)^{\frac{s-\frac{d}{r}}{2(s-d(\frac{1}{r}-\frac{1}{2}))}}. \end{split}$$

Therefore,

(5.35) 
$$III' \le c \left(\frac{\log n}{n}\right)^{\frac{s - \frac{d}{r}}{2(s - d(\frac{1}{r} - \frac{1}{2}))}}.$$

To estimate II' we first observe that

$$II' = \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\hat{\beta}_{j\xi}| > 2\lambda_n\}} \mathbb{1}_{\{|\beta_{j,\xi}| > \lambda_n\}} \Big]$$
$$\leq \sum_{0 \le j \le J_n} b^{j\frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}| \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \Big].$$

By (5.33) we have

$$\kappa \left(\frac{\log n}{n}\right)^{\frac{1}{2}} = \lambda_n < |\beta_{j\xi}| \le c \|f\|_{B^s_{\tau\tau}} b^{-j(s-d(\frac{1}{r}-\frac{1}{2}))}, \quad 0 \le j \le J_n.$$

Therefore, necessarily  $0 \leq j \leq \tilde{J}_n$ , where

$$b^{\tilde{J}_n} \sim \left(\frac{\|f\|_{B^s_{r\tau}}}{\kappa}\right)^{\frac{1}{s-d(\frac{1}{r}-\frac{1}{2})}} \left(\frac{n}{\log n}\right)^{\frac{1}{2(s-d(\frac{1}{r}-\frac{1}{2}))}}$$

and consequently

$$II' \leq \sum_{0 \leq j < \tilde{J}_n} b^{j \frac{d}{2}} \mathbb{E} \Big[ \sup_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j,\xi} - \beta_{j\xi}| \Big].$$

We next utilize Lemma 5.5. Consider the family  $\mathcal{F} := \{\tilde{\psi}_{j\xi} : \xi \in \mathcal{X}_j\}$  with  $\operatorname{card}(\mathcal{F}) \leq cb^{jd}$  and by (3.16)

$$\|\tilde{\psi}_{j\xi}\|_{\infty} \le c_{\diamond} b^{jd/2} =: M/2 \quad \text{and} \quad \mathbb{E}|\tilde{\psi}_{j\xi}(X)|^2 \le c_{\diamond} \|f\|_{\infty} =: \sigma^2.$$

Then applying Lemma 5.5 we get

$$\mathbb{E}\Big[\sup_{\xi \in \mathcal{X}_{j}} |\hat{\beta}_{j,\xi} - \beta_{j\xi}|\Big] \le c \|f\|_{\infty}^{\frac{1}{2}} \Big(\frac{\log(b^{jd})}{n}\Big)^{\frac{1}{2}} + cb^{j\frac{d}{2}} \frac{\log(b^{jd})}{n} \\ \le c \|f\|_{\infty}^{\frac{1}{2}} \Big(\frac{j}{n}\Big)^{\frac{1}{2}} + cb^{j\frac{d}{2}} \frac{j}{n}.$$

Since also  $\tilde{J}_n \leq c(s,r) \log n$ , we get

$$II' \le c \sum_{0 \le j \le \tilde{J}_n} b^{j\frac{d}{2}} \Big( \|f\|_{\infty}^{\frac{1}{2}} \Big( \frac{j}{n} \Big)^{\frac{1}{2}} + b^{j\frac{d}{2}} \frac{j}{n} \Big) \le c \|f\|_{\infty}^{\frac{1}{2}} \Big( b^{d\tilde{J}_n} \frac{\tilde{J}_n}{n} \Big)^{\frac{1}{2}} + c b^{d\tilde{J}_n} \frac{\tilde{J}_n}{n}.$$

But

$$\begin{split} b^{d\tilde{J}_n} \frac{\tilde{J}_n}{n} &\leq c \|f\|_{B^s_{r\tau}}^{\frac{d}{s-d(\frac{1}{r}-\frac{1}{2})}} \left(\frac{n}{\log n}\right)^{\frac{d}{2(s-d(\frac{1}{r}-\frac{1}{2}))}} \frac{\log n}{n} \\ &= c \|f\|_{B^s_{r\tau}}^{\frac{d}{s-d(\frac{1}{r}-\frac{1}{2})}} \left(\frac{\log n}{n}\right)^{\frac{(s-\frac{d}{r})}{(s-d(\frac{1}{r}-\frac{1}{2}))}}. \end{split}$$

Thus, because s > d/r we conclude, using Proposition 3.5 (ii),

$$II' \le c \left( \|f\|_{B^s_{r\tau}}^{1/2} \|f\|_{B^s_{r\tau}}^{\frac{d}{2(s-d(\frac{1}{r}-\frac{1}{2}))}} + \|f\|_{B^s_{r\tau}}^{\frac{d}{s-d(\frac{1}{r}-\frac{1}{2})}} \right) \left(\frac{\log n}{n}\right)^{\frac{s-\frac{d}{r}}{2(s-d(\frac{1}{r}-\frac{1}{2}))}}.$$

Therefore,

(5.36) 
$$II' \le c \left(\frac{\log n}{n}\right)^{\frac{s-\frac{d}{r}}{2(s-d(\frac{1}{r}-\frac{1}{2}))}}.$$

Estimation of II and III. As  $p\geq 1$  Jensen's inequality implies

$$II \leq \sum_{0 \leq j \leq J_n} b^{jd(\frac{1}{2} - \frac{1}{p})} \mathbb{E} \left[ \left( \sum_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \right)^{\frac{1}{p}} \right]$$
$$\leq \sum_{0 \leq j \leq J_n} b^{jd(\frac{1}{2} - \frac{1}{p})} \left( \mathbb{E} \left[ \sum_{\xi \in \mathcal{X}_j} |\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \right] \right)^{\frac{1}{p}}$$
$$= \sum_{0 \leq j \leq J_n} \left( b^{jd(\frac{p}{2} - 1)} \sum_{\xi \in \mathcal{X}_j} \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \mathbb{E} \left[ |\hat{\beta}_{j\xi} - \beta_{j\xi}|^p \right] \right)^{\frac{1}{p}}.$$

By (5.20) and (5.21) it follows that  $\mathbb{E}[|\hat{\beta}_{j,k} - \beta_{j,k}|^p] \le c \left(\frac{A}{n}\right)^{p/2}$  and hence

$$II \le c \left(\frac{A}{n}\right)^{\frac{1}{2}} \sum_{0 \le j \le J_n} \left( b^{jd(\frac{p}{2}-1)} \sum_{\xi \in \mathcal{X}_j} \mathbb{1}_{\{|\beta_{j\xi}| > \lambda_n\}} \right)^{\frac{1}{p}}.$$

Let q < p. Using the Bienayme-Chebyshev inequality we get

(5.37) 
$$II \le c \left(\frac{A}{n}\right)^{\frac{1}{2}} \sum_{0 \le j \le J_n} \left( b^{jd(\frac{p}{2}-1)} \frac{1}{\lambda_n^q} \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^q \right)^{\frac{1}{p}} \\ = c (\log n)^{-1/2} \lambda_n^{1-\frac{q}{p}} \sum_{0 \le j \le J_n} \left( b^{jd(\frac{p}{2}-1)} \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^q \right)^{\frac{1}{p}}.$$

For the term III we have

$$III \leq \sum_{0 \leq j \leq J_n} b^{jd(\frac{1}{2} - \frac{1}{p})} \mathbb{E} \left[ \left( \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^p \mathbb{1}_{\{|\beta_{j\xi}| \leq 3\lambda_n\}} \right)^{\frac{1}{p}} \right]$$
$$= \sum_{0 \leq j \leq J_n} \left( b^{jd(\frac{p}{2} - 1)} \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^p \mathbb{1}_{\{|\beta_{j\xi}| \leq 3\lambda_n\}} \right)^{\frac{1}{p}}.$$

Assuming that  $\lambda > 0, 0 < q < p$ , and  $\sigma$  is a positive measure on a measure space X, we have

$$\begin{split} \int_X \mathbb{1}_{\{|f| \le \lambda\}} |f|^p d\sigma &\leq \int_X (|f| \land \lambda)^p d\sigma = \int_0^\lambda p x^{p-1} \sigma(|f| > x) dx \\ &\leq \int_0^\lambda p x^{p-1} \frac{\|f\|_q^q}{x^q} dx = \frac{p}{p-q} \|f\|_q^q \lambda^{p-q}. \end{split}$$

Therefore, if q < p we have

$$III \leq \sum_{0 \leq j \leq J_n} \left( b^{jd(\frac{p}{2}-1)} \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^p \mathbb{1}_{\{|\beta_{j\xi}| \leq 3\lambda_n\}} \right)^{\frac{1}{p}}$$
$$\leq c(p,q) \sum_{0 \leq j \leq J_n} \left( \lambda_n^{p-q} b^{jd(\frac{p}{2}-1)} \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^q \right)^{\frac{1}{p}}$$
$$\leq c\lambda_n^{1-\frac{q}{p}} \sum_{0 \leq j \leq J_n} \left( b^{jd(\frac{p}{2}-1)} \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^q \right)^{\frac{1}{p}}.$$

This coupled with (5.37) yields

(5.38) 
$$II + III \le c\lambda_n^{1-\frac{q}{p}} \sum_{0 \le j \le J_n} \left( b^{jd(\frac{p}{2}-1)} \sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}|^q \right)^{\frac{1}{p}}, \quad 0 < q < p.$$

Assuming 0 < q < p we set  $s' := \frac{d(p-q)}{2q}$ . By (3.26) it follows that

$$\sum_{\xi \in \mathcal{X}_j} |\beta_{j\xi}(f)|^q \le c b^{-jq(s'+d(\frac{1}{2}-\frac{1}{q}))} ||f||^q_{B^{s'}_{q\tau}}, \quad j \ge 0$$

4

We combine this with (5.38) and the fact that  $J_n \leq c \log n$  to obtain

(5.39) 
$$II + III \leq c\lambda_n^{1-\frac{q}{p}} \|f\|_{B_{q\tau}^{s'}}^{\frac{q}{p}} \sum_{0 \leq j \leq J_n} \left( b^{jd(\frac{p}{2}-1)} b^{-jq(s'+d(\frac{1}{2}-\frac{1}{q}))} \right)^{\frac{1}{p}} \\ \leq c\lambda_n^{1-\frac{q}{p}} \|f\|_{B_{q\tau}^{s'}}^{\frac{q}{p}} J_n \leq c \log n \left(\kappa \frac{\log n}{n}\right)^{\frac{p-q}{2p}} \|f\|_{B_{q\tau}^{s'}}^{\frac{q}{p}}.$$

Thus, in what follows we have to show that  $\|f\|_{B^{s'}_{q\tau}} \leq c \|f\|_{B^s_{r\tau}}$  for a suitable q < p.

5.2.3. Proof of Theorem 5.2 (i). Let  $1 \le r \le \infty$ ,  $0 < \tau \le \infty$ , s > d/r, and m > 0. Assume  $f \in B^s_{r\tau}$  and  $\|f\|_{B^s_{r\tau}} \le m$ . Then combining (5.31), (5.32), (5.34), (5.35), (5.36), and (5.28) we obtain

$$\mathbb{E}\|\hat{f}_n - f\|_{\infty} \le c(m) \Big(\frac{\log n}{n}\Big)^{\frac{s - \frac{d}{r}}{2(s - d(\frac{1}{r} - \frac{1}{2}))}},$$

which confirms (5.8).

5.2.4. Proof of Theorem 5.2 (ii). Let  $1 \le r \le \infty$ ,  $0 < \tau \le \infty$ ,  $1 \le p < \infty$ , s > d/r, and m > 0. Assume  $f \in B^s_{r\tau}$ ,  $s \ge \frac{dp}{2} \left(\frac{1}{r} - \frac{1}{p}\right)$ , and  $\|f\|_{B^s_{r\tau}} \le m$ . Choose s' = s. Then using that from above

$$\frac{d(p-q)}{2q} = s' = s \ge \frac{dp}{2} \left(\frac{1}{r} - \frac{1}{p}\right)$$

we obtain

$$q = \frac{dp}{2s+d} < p$$
,  $\frac{p-q}{2p} = \frac{s}{2s+d}$  and  $q \le r$ .

From Proposition 3.5 (i) and (iii) it follows that

$$\|f\|_{B^{s'}_{q\tau}} = \|f\|_{B^s_{q\tau}} \le c \|f\|_{B^s_{r\tau}} \le cm \quad \text{and} \quad \|f\|_\infty \le c \|f\|_{B^s_{r\tau}} \le cm.$$

and from (5.39) we conclude

$$II + III \le c \log n \left( \kappa \frac{\log n}{n} \right)^{\frac{s}{2s+d}} \|f\|_{B^s_{r\tau}}^{\frac{d}{2s+d}} \le c(m) \log n \left( \frac{\log n}{n} \right)^{\frac{s}{2s+d}}.$$

This estimate along with (5.30), (5.32), (5.34), and (5.28) imply (5.9).

5.2.5. Proof of Theorem 5.2 (iii). Under the hypotheses of Theorem 5.2 let  $f \in B^s_{r\tau}$ ,  $s < \frac{dp}{2} \left(\frac{1}{r} - \frac{1}{p}\right)$ , and  $\|f\|_{B^s_{r\tau}} \leq m$ . In this case we choose  $s' := s - d\left(\frac{1}{r} - \frac{1}{p}\right) > 0$ . As above  $s' = \frac{d(p-q)}{2q}$  and hence

$$q = \frac{dp(\frac{1}{2} - \frac{1}{p})}{s - d(\frac{1}{r} - \frac{1}{2})} \ge r, \quad p > 2, \quad \text{and} \quad \frac{p - q}{2p} = \frac{s - d(\frac{1}{r} - \frac{1}{p})}{2(s - d(\frac{1}{r} - \frac{1}{2}))}.$$

By Proposition 3.5 (ii) it follows that  $||f||_{B^{s'}_{q\tau}} \leq c||f||_{B^s_{r\tau}}$  and hence from (5.39)

$$II + III \le c(m) \log n \left(\frac{\log n}{n}\right)^{\frac{s - d(\frac{1}{r} - \frac{1}{p})}{2(s - d(\frac{1}{r} - \frac{1}{2}))}}.$$

Combining this with (5.30), (5.32), (5.34), and (5.28) imply (5.10).

The proof of Theorem 5.2 is complete.

## 6. Examples of settings covered by our theory

In this section we present a number of examples of settings that are covered by the general framework from Section 2.

Clearly, the classical setup of  $\mathcal{M} = \mathbb{R}^d$  equipped with the Euclidean distance and Lebesgue measure, and with  $L = -\Delta$  the Laplacian obeys conditions **C1–C5**, together with **C1A** from Section 2. Therefore, our results on density estimators apply. They are compatible with the existing upper bound results. For instance, our result on adaptive wavelet estimators (Theorem 5.2) is compatible with the result in the classical setting on  $\mathbb{R}^d$  from [6].

The unit sphere  $\mathbb{S}^d$  in  $\mathbb{R}^{d+1}$  equipped with the standard (geodesic) distance, measure, and -L being the Laplace-Beltrami operator is another example of a setup that obeys conditions **C1–C5**, together with **C1A**. Our upper bound result on adaptive wavelet estimators (Theorem 5.2) is compatible with the upper bound estimate on the adaptive needlet estimator in [1].

A natural generalization of the above setup is the case of a compact Riemannian manifold  $\mathcal{M} \subset \mathbb{R}^m$  equipped with the natural Riemannian measure, geodesic

distance, and -L being the Laplace-Beltrami operator. Then conditions C1–C5, together with C1A from Section 2 are satisfied.

For other examples on Riemannian manifolds and Lie groups we refer the reader to [4] and the references therein.

We next describe the recently developed in [19] general setting of a subset of  $\mathbb{R}^n$  complemented by a differential operator L that is a realization in local coordinates of a weighted Laplace operator on suitable subset of a Riemannian manifold. In this setting conditions **C1–C5** and in some cases **C1A** from Section 2 are obeyed. This setting covers the weighted settings on the interval, ball, and simplex, which we will describe in more detail as well.

6.1. Convex subsets of Riemannian manifolds and counterparts on  $\mathbb{R}^n$ . We assume that  $V \subset \mathbb{R}^m$  is a connected open set in  $\mathbb{R}^m$  with the properties:  $X := \overline{V}$  is compact,  $\mathring{X} = V$ , and  $X \setminus V$  is of Lebesgue measure zero. Let L be a differential operator of the form

(6.1) 
$$L = \sum_{i,j=1}^{m} a_{ij}(x)\partial_i\partial_j + \sum_{j=1}^{m} b_j(x)\partial_j,$$

where  $a_{ij}$  and  $b_j$  are polynomials of degrees two and one, respectively. The underlying space is  $\mathbb{L}^2(V,\mu)$ , where  $d\mu(x) := \check{w}(x)dx$  with  $\check{w} \in C^{\infty}(V)$ ,  $\check{w} > 0$ , and  $\int_V \check{w}(x)dx < \infty$ .

On the other hand, we assume that there is (a closely related) counterpart to the above setup. Namely, we assume that  $(M, d, \nu)$  is an *m*-dimensional complete Riemannian manifold and  $M \subset \mathbb{R}^m$ , where the Riemannian metric is induced by the inner product on  $\mathbb{R}^m$ . We stipulate two conditions on  $(M, d, \nu)$ : (i) the volume doubling condition is valid, and (ii) the Poincaré inequality holds true (see [19]).

Further, we assume that  $(U, \varphi)$  is a chart on M, where U is a convex open relatively compact subset of M such that  $\varphi$  maps diffeomorphically U onto V, where  $V \subset \mathbb{R}^d$  is from above. We set  $\phi := \varphi^{-1}$  and  $Y := \overline{U}$ .

The key assumption is that the map  $\phi$  provides an one-to-one correspondence between the elements of the setting on X from above and the setting on Y. More precisely, it is assumed that the distance  $\rho(\cdot, \cdot)$  on X is induced by the geodesic distance  $d(\cdot, \cdot)$  on  $Y \subset M$ . The weighted measure  $d\mu(x) := \check{w}(x)dx$  on X is also induced by the respective wighted measure  $\nu_w$  on Y. Namely, assuming that g(x) = $(g_{ij}(x))$  is the Riemannian tensor it is stipulated that w > 0 is a  $C^{\infty}(U)$  weight function that is compatible with  $\check{w}$  from above in the following sense:

(6.2) 
$$\breve{w}(x) := w(\phi(x))\sqrt{\det g(x)}, \quad x \in V,$$

It is assumed that  $\nu_w = wd\nu$ . It is also assumed that the operator L from (6.1) is a realization in local coordinates (via the chart  $(U, \varphi)$ ) of the weighted Laplacian

$$\Delta_w f := \frac{1}{w} \operatorname{div}(w\nabla f) \quad \text{on } Y \subset M \quad (\text{see [19]}).$$

In addition, it is assumed that the volume doubling condition (2.1) on Y (and hence on X) is valid, and a natural regularity condition on the weighted functions and suitable Green's theorem are verified. See [19] for the details.

In [19] it is shown that under the above conditions the heat kernels associated to the operator L and weighted Laplacian  $\Delta_w$  have Gaussian localization just as in (2.8) and as a consequence the Hölder continuity (see (2.9)) is valid. Furthermore, these heat kernels have the Markov property (see (2.10)).

As a result, in the above described general setting conditions **C1–C5** in Section 2 are obeyed and our results on kernel and linear wavelet density estimators apply to the settings on  $X \subset \mathbb{R}^m$  and on  $Y \subset M$ . Furthermore, if the weight function  $w \equiv 1$ on Y, then the measures  $\mu$  on X and  $\nu_w$  on Y verify the Ahlfors condition (2.3), i.e. condition **C1A** is satisfied, and consequently our result on adaptive wavelet estimators (Theorem 5.2) applies.

We next show how the general theory described above is implemented in specific settings on [-1, 1], the ball and simplex.

## 6.2. Specific examples where our kernel and wavelet estimators work.

6.2.1. Jacobi operator on [-1, 1]. We consider the classical setting of  $\mathcal{M} = [-1, 1]$  equipped with the weighted measure

$$d\mu(x) := w(x)dx = (1-x)^{\alpha}(1+x)^{\beta}dx, \quad \alpha, \beta > -1,$$

and the distance  $\rho(x, y) := |\arccos x - \arccos y|$ , complemented with the classical Jacobi operator, defined by

$$Lf(x) := -\frac{\left[w(x)(1-x^2)f'(x)\right]'}{w(x)}$$

As is well known the Jacobi polynomials are eigenfunctions of this operator. Denote  $B(x,r) := \{y \in [-1,1] : \rho(x,y) < r\}$ . It is easy to see that (see e.g. [4])

(6.3) 
$$|B(x,r)| \sim r(1-x+r^2)^{\alpha+1/2}(1+x+r^2)^{\beta+1/2}, \quad x \in [-1,1], \ 0 < r \le \pi.$$

Hence, we have a doubling metric measure space with homogeneous dimension  $d = 1 + (2\alpha + 1)_+ \vee (2\beta + 1)_+$ . More importantly, as is shown in [4] the associated heat kernel has Gaussian localization, Hölder continuity, and the Markov property (see also [19]). Therefore, conditions **C1–C5** in Section 2 are obeyed.

Ahlfors space on [-1, 1]. In the above setting, if  $\alpha = \beta = -1/2$  then from (6.3)

$$|B(x,r)| \sim r, \quad x \in [-1,1], \ 0 < r \le \pi.$$

Therefore, condition C1A (see §2) is obeyed with d = 1.

6.2.2. Weighted unit ball. Consider the case when  $\mathcal{M}$  is  $\mathbb{B}^m := \{x \in \mathbb{R}^m : ||x|| < 1\}$  the unit ball in  $\mathbb{R}^m$  equipped with the measure

$$d\mu := (1 - \|x\|^2)^{\gamma - 1/2} dx, \quad \gamma > -1,$$

and the distance

$$\rho(x,y) := \arccos\left(\langle x,y\rangle + \sqrt{1 - \|x\|^2}\sqrt{1 - \|y\|^2}\right),$$

where  $\langle x, y \rangle$  is the inner product of  $x, y \in \mathbb{R}^m$  and  $||x|| := \sqrt{\langle x, x \rangle}$ . Denoting  $B(x, r) := \{y \in \mathbb{B}^m : \rho(x, y) < r\}$  it is easy to show (see [5]) that

(6.4) 
$$|B(x,r)| \sim r^m (1 - ||x||^2 + r^2)^{\gamma},$$

which implies that  $(\mathcal{M}, \mu, \rho)$  obeys the doubling condition (2.1) and non-collapsing condition (2.5) and it is of homogeneous dimension  $d = m + 2\gamma_+$ .

Consider the operator

$$L := -\sum_{i=1}^m (1 - x_i^2)\partial_i^2 + 2\sum_{1 \le i < j \le m} x_i x_j \partial_i \partial_j + (n + 2\gamma)\sum_{i=1}^m x_i \partial_i,$$

acting on sufficiently smooth functions on  $\mathbb{B}^m$ . This operator is essentially selfadjoint and positive (see [5] and also [19]). More importantly, its heat kernel has Gaussian localization, Hölder continuity, and the Markov property (see [19]). In fact, L is a realization of a weighted Laplace operator on the upper hemisphere of  $\mathbb{R}^{m+1}$  in local coordinates (see [19]). Consequently, conditions **C1–C5** in Section 2 are verified and this setting falls in the general framework from Section 2.

Ahlfors space on the unit ball. Assume that in the above setting  $\gamma = 0$ , i.e. the measure on  $\mathbb{B}^m$  is  $d\mu := (1 - ||x||^2)^{-1/2} dx$ . Then from (6.4)

$$|B(x,r)| \sim r^m, \quad x \in \mathbb{B}^m, \ 0 < r \le 1,$$

and hence condition C1A in Section 2 is obeyed with d = m. Thus this is another example of an Ahlfors space.

6.2.3. Weighted simplex. We now consider the simplex

$$\mathbb{T}^m := \left\{ x \in \mathbb{R}^d : x_1 > 0, \dots, x_m > 0, \ |x| < 1 \right\}, \quad |x| := x_1 + \dots + x_m,$$

in  $\mathbb{R}^m$  equipped with the measure

$$d\mu(x) = \prod_{i=1}^{m} x_i^{\kappa_i - 1/2} (1 - |x|)^{\kappa_{m+1} - 1/2} dx, \quad \kappa_i > -1/2,$$

and distance

$$\rho(x,y) = \arccos\Big(\sum_{i=1}^{m} \sqrt{x_i y_i} + \sqrt{1-|x|}\sqrt{1-|y|}\Big).$$

Similarly as before we use the notation:  $B(x,r) := \{y \in \mathbb{T}^m : \rho(x,y) < r\}$ . It is known (see [19]) that

(6.5) 
$$|B(x,r)| \sim r^m (1-|x|+r^2)^{\kappa_{m+1}} \prod_{i=1}^m (x_i+r^2)^{\kappa_i}$$

Hence, the doubling condition (2.1) and non-collapsing condition (2.5) are satisfied. Moreover  $d = m + 2((\kappa_1)_+ + \cdots + (\kappa_{m+1})_+)$ .

It is natural to consider the operator

$$L := -\sum_{i=1}^m x_i \partial_i^2 + \sum_{i=1}^m \sum_{j=1}^m x_i x_j \partial_i \partial_j - \sum_{i=1}^m \left(\kappa_i + \frac{1}{2} - \left(|\kappa| + \frac{m+1}{2}\right) x_i\right) \partial_i$$

with  $|\kappa| := \kappa_1 + \cdots + \kappa_{m+1}$ . In [19] (see also [20]) it is shown that this operator is essentially self-adjoint and positive. Furthermore, its heat kernel has Gaussian localization, Hölder continuity, and the Markov property. It is important to point out that the operator L is a realization of a weighted Laplacian on the sphere in the first octant in local coordinates (see [19]). Thus, conditions **C1–C5** in §2 are verified and this setup is covered by the general setting from Section 2. Ahlfors space on the simplex. If above  $\kappa_i = 0, i = 1, ..., m + 1$ , then the measure is given by  $d\mu(x) = \prod_{i=1}^m x_i^{-1/2} (1 - |x|)^{-1/2} dx$ , and (6.5) yields

$$|B(x,r)| \sim r^m, \quad x \in \mathbb{T}, \ 0 < r \le 1.$$

Therefore, condition C1A is again satisfied and this is also an example of an Ahlfors space with d = m where our result on adaptive wavelet estimators (Theorem 5.2) applies.

## 7. Appendix

7.1. **Proof of Proposition 3.2.** Let  $1 \le p < q \le \infty$  and  $g \in \Sigma_{\lambda}^{p}$ ,  $\lambda \ge 1$ . Let  $\varphi \in C_{0}^{\infty}(\mathbb{R})$  be an even function with the properties: supp  $\varphi \subset [-2, 2]$ ,  $0 \le \varphi \le 1$ , and  $\varphi(u) = 1$  for  $u \in [-1, 1]$ .

By Theorem 3.1  $\varphi(\delta\sqrt{L})$  is an integral operator whose kernel  $\varphi(\delta\sqrt{L})(x,y)$  is real-valued, symmetric, and such that for any k > d,  $x, y \in \mathcal{M}$ , and  $\delta > 0$ 

(7.1) 
$$|\varphi(\delta\sqrt{L})(x,y)| \le c(k)|B(x,\delta)|^{-1}(1+\delta^{-1}\rho(x,y))^{-k}.$$

where the constant c(k) > 0 depends only on k,  $\varphi$ , and the constants from the setting in Section 2.

Observe that since  $\varphi(u) = 1$  for  $u \in [-1, 1]$  and  $g \in \Sigma_{\lambda}^{p}$ , we have

(7.2) 
$$\varphi(\delta\sqrt{L})g = g \text{ if } \delta = 1/\lambda.$$

Set  $\delta := 1/\lambda$  and k := d+1. Define r > 1 from the identity 1/p - 1/q = 1 - 1/r. Using (7.1), (3.6), and (3.4) we obtain

$$\|\varphi(\delta\sqrt{L})(x,\cdot)\|_{r} \leq c(d+1)|B(x,\delta)|^{-1} \Big(\int_{\mathcal{M}} (1+\delta^{-1}\rho(x,y))^{-r(d+1)}d\mu(y)\Big)^{1/r}$$
(7.3) 
$$\leq c_{d}\tilde{c}|B(x,\delta)|^{1/r-1} \leq c_{d}\tilde{c}(c_{0}/c_{3})^{1-1/r}\delta^{d(1/r-1)} \leq c_{\star}\delta^{d(1/r-1)}.$$

Here  $\tilde{c} := (2^{-d} - 2^{-d-1})^{-1}$ ,  $c_d = c(d+1)$  is from (7.1), the constants  $c_0, c_3 > 0$  are from (2.1) and (2.5). We may assume that  $c_0/c_3 \ge 1$  and take  $c_{\star} = c_d \tilde{c}(c_0/c_3)$ , which is independent of p and q.

Thus, using the symmetry we have

$$\|\varphi(\delta\sqrt{L})(x,\cdot)\|_r = \|\varphi(\delta\sqrt{L})(\cdot,y)\|_r \le c_\star \delta^{d(1/r-1)} = c_\star \delta^{d(1/q-1/p)}$$

We now use the well known Theorem 6.36 from [10] to conclude that

$$\|g\|_{q} = \|\varphi(\delta\sqrt{L})g\|_{q} \le c_{\star}\delta^{d(1/q-1/p)}\|g\|_{p} = c_{\star}\lambda^{d(1/p-1/q)}\|g\|_{p}.$$

Here we also used (7.2). The proof is complete.

7.2. **Proof of Proposition 3.5.** (i) Let  $1 \le q \le r, 0 < \tau \le \infty, s > 0, \mu(\mathcal{M}) < \infty$ , and  $f \in B^s_{r\tau}$ . Let  $\Phi_j, j \ge 0$ , be the functions from the definition of Besov spaces, see Definition 3.4. Then by Hölder's inequality  $\|\Phi_j(\sqrt{L})f\|_q \le \mu(\mathcal{M})^{\frac{1}{q}-\frac{1}{r}} \|\Phi_j(\sqrt{L})f\|_r$ , which readily implies  $\|f\|_{B^s_{q\tau}} \le \mu(\mathcal{M})^{\frac{1}{q}-\frac{1}{r}} \|f\|_{B^s_{r\tau}}$  as claimed.

(ii) Let  $1 \leq r \leq q \leq \infty$ ,  $0 < \tau \leq \infty$ , and  $s > d(\frac{1}{r} - \frac{1}{q})$ . Assume  $f \in B^s_{r\tau}$ . Let  $\Phi_j$ ,  $j \geq 0$ , be the functions from the definition of Besov spaces. Clearly,  $\Phi_j \subset [b^{j-1}, b^{j+1}]$  and hence  $\Phi_j(\sqrt{L})f \in \Sigma_{b^{j+1}}$ . Applying Proposition 3.2 we obtain

(7.4) 
$$\|\Phi_j(\sqrt{L})f\|_q \le c b^{jd(\frac{1}{r} - \frac{1}{q})} \|\Phi_j(\sqrt{L})f\|_r.$$

From this it readily follows that  $\|f\|_{B^{s-d(\frac{1}{r}-\frac{1}{q})}_{q_{\tau}}} \leq c \|f\|_{B^{s}_{r_{\tau}}}$  as claimed.

(iii) Assume  $1 \le r \le \infty$ ,  $0 < \tau \le \infty$ , s > d/r, and  $f \in B^s_{r\tau}$ . Let the functions  $\Phi_j$ ,  $j \ge 0$ , in the definition of Besov spaces be selected so that  $\sum_{j=0}^{\infty} \Phi_j(\lambda) = 1$  for  $\lambda \ge 0$ . For example, the functions  $\Psi_j$ ,  $j \ge 0$ , from the definition of the frames in §3.4 have this property. Then as in (3.13) we have  $f = \sum_{j=0}^{\infty} \Phi_j(\sqrt{L})f$ . Just as in (7.4) by Proposition 3.2 we have

$$\|\Phi_j(\sqrt{L})f\|_{\infty} \le cb^{jd/r} \|\Phi_j(\sqrt{L})f\|_r$$

Also, evidently by the definition of Besov spaces  $\|\Phi_j(\sqrt{L})f\|_r \leq cb^{-js}\|f\|_{B^s_{r\tau}}$ . We use these two estimates and the fact that s > d/r to obtain

$$\begin{split} \|f\|_{\infty} &\leq \sum_{j=0}^{\infty} \|\Phi_{j}(\sqrt{L})f\|_{\infty} \leq c \sum_{j=0}^{\infty} b^{jd/r} \|\Phi_{j}(\sqrt{L})f\|_{r} \\ &\leq c \sum_{j=0}^{\infty} b^{-j(s-d/r)} \|f\|_{B^{s}_{r\tau}} \leq c \|f\|_{B^{s}_{r\tau}}. \end{split}$$

(iv) Let  $1 \le p \le r \le \infty$ ,  $0 < \tau \le \infty$ , s > 0, and  $\mu(\mathcal{M}) < \infty$ . Assume  $f \in B^s_{r\tau}$ . Let  $\Phi_j$ ,  $j \ge 0$ , be as in the proof of (iii) above. Since  $p \ge 1$  and s > 0 we have

$$\|f\|_{p} \leq \sum_{j=0}^{\infty} \|\Phi_{j}(\sqrt{L})f\|_{p} \leq c \sum_{j=0}^{\infty} b^{-js} \|f\|_{B_{p\tau}^{s}} \leq c \|f\|_{B_{r\tau}^{s}},$$

where for the last inequality we used (i).

7.3. **Proof of Proposition 4.6.** Let  $\check{\Phi}_0$  be just as the function  $\Phi$  in the definition of Besov spaces in §3.5 (see (3.18)), that is,  $\check{\Phi}_0 \in C^{\infty}(\mathbb{R}_+)$  is a real-valued function with the properties:  $\operatorname{supp}\check{\Phi}_0 \subset [0,b], \check{\Phi}_0(\lambda) = 1$  for  $\lambda \in [0,1]$ , and  $\check{\Phi}_0(\lambda) \ge c > 0$ for  $\lambda \in [0, b^{3/4}]$ . Set  $\check{\Phi}(\lambda) := \check{\Phi}_0(\lambda) - \check{\Phi}_0(b\lambda)$  and  $\check{\Phi}_j(\lambda) := \check{\Phi}(b^{-j}\lambda), j \ge 1$ . Clearly,  $\sum_{j\ge 0} \check{\Phi}_j(\lambda) = 1, \lambda \in [0,\infty)$ . Let  $0 < \delta \le (2b^2)^{-1}$ . The case when  $(2b^2)^{-1} < \delta \le 1$  is easier; we omit it.

Let  $0 < \delta \leq (2b^2)^{-1}$ . The case when  $(2b^2)^{-1} < \delta \leq 1$  is easier; we omit it. Choose  $\nu \geq 2$  and  $\ell \geq 1$  ( $\nu, \ell \in \mathbb{N}$ ) so that  $b^{\nu} \leq 1/(2\delta) < b^{\nu+1}$  and  $1/\delta \leq b^{\nu+\ell}$  $(\ell := \lfloor \log(2b)/\log b \rfloor + 1$  will do). From above and the properties of  $\Phi$  in (4.4) it readily follows that

$$1 - \Phi(\delta\lambda) = (1 - \Phi(\delta\lambda)) \sum_{j=\nu}^{\nu+\ell} \check{\Phi}(b^{-j}\lambda) + \sum_{j=\nu+\ell+1}^{\infty} \check{\Phi}(b^{-j}\lambda).$$

Set

(7.5) 
$$\Lambda(\lambda) := (1 - \Phi(\delta b^{\nu} \lambda)) \sum_{j=0}^{\ell} \breve{\Phi}(b^{-j} \lambda) \quad \text{and} \quad \Theta(\lambda) := \sum_{j=\nu-1}^{\nu+\ell+2} \breve{\Phi}(b^{-j} \lambda).$$

By the construction of  $\check{\Phi}_j$  it follows that  $\operatorname{supp} \Lambda \subset [1, b^{\ell+1}]$  and  $\Theta(\lambda) = 1$  for  $\lambda \in [b^{\nu}, b^{\nu+\ell+1}]$ , implying  $\Lambda(b^{-\nu}\lambda) = \Lambda(b^{-\nu}\lambda)\Theta(\lambda)$ . Therefore, we have

$$1 - \Phi(\delta\lambda) = \Lambda(b^{-\nu}\lambda) + \sum_{j=\nu+\ell+1}^{\infty} \breve{\Phi}(b^{-j}\lambda) = \Lambda(b^{-\nu}\lambda)\Theta(\lambda) + \sum_{j=\nu+\ell+1}^{\infty} \breve{\Phi}(b^{-j}\lambda).$$

From this we infer that for any  $f \in \mathbb{L}^p$ 

$$f - \Phi(\delta\sqrt{L})f = \Lambda(b^{-\nu}\sqrt{L})\Theta(\sqrt{L})f + \sum_{j=\nu+\ell+1}^{\infty} \check{\Phi}(b^{-j}\sqrt{L})f$$

and hence

(7.6) 
$$\|f - \Phi(\delta\sqrt{L})f\|_p \le \|\Lambda(b^{-\nu}\sqrt{L})\Theta(\sqrt{L})f\|_p + \sum_{j=\nu+\ell+1}^{\infty} \|\breve{\Phi}(b^{-j}\sqrt{L})f\|_p$$

From the definition of  $\Lambda(\lambda)$  in (7.5) it follow that  $\Lambda \in C^{\infty}(\mathbb{R}_+)$ , supp  $\Lambda \subset [1, b^{\ell+1}]$ , and  $\|\Lambda^{(r)}\|_{\infty} \leq c_r$  for  $r = 0, 1, \ldots$  with the constant  $c_r > 0$  depending only on r,  $\Phi$ ,  $\check{\Phi}$ , b. Now, we invoke Theorem 3.1 to conclude that  $\Lambda(b^{-\nu}\sqrt{L})$  is an integral operator with kernel  $\Lambda(b^{-\nu}\sqrt{L})(x, y)$  satisfying

$$|\Lambda(b^{-\nu}\sqrt{L})(x,y)| \le c(k)b^{d\nu} \left(1 + b^{\nu}\rho(x,y)\right)^k, \quad x,y \in \mathcal{M}, \ k > d.$$

As in (4.13) this with k = d+1 implies  $\|\Lambda(b^{-\nu}\sqrt{L})(\cdot, y)\|_1 = \|\Lambda(b^{-\nu}\sqrt{L})(x, \cdot)\|_1 \le c$ . and applying Schur's lemma it follows that  $\|\Lambda(b^{-\nu}\sqrt{L})\|_{p\to p} \le c < \infty$ . As a consequence, we get

$$\|\Lambda(b^{-\nu}\sqrt{L})\Theta(\sqrt{L})f\|_p \le c\|\Theta(\sqrt{L})f\|_p \le c\sum_{j=\nu-1}^{\nu+\ell+2} \|\breve{\Phi}(b^{-j}\sqrt{L})f\|_p$$

This coupled with (7.6) implies, when  $0 < q < \infty$ ,

$$\begin{split} \|f - \Phi(\delta\sqrt{L})f\|_{p} &\leq c \sum_{j=\nu-1}^{\infty} \|\check{\Phi}(b^{-j}\sqrt{L})f\|_{p} \leq c'b^{-s\nu} \Big(\sum_{j=\nu-1}^{\infty} \left(b^{sj}\|\check{\Phi}(b^{-j}\sqrt{L})f\|_{p}\right)^{q}\Big)^{1/q} \\ &\leq c\delta^{s} \Big(\sum_{j=0}^{\infty} \left(b^{sj}\|\check{\Phi}(b^{-j}\sqrt{L})f\|_{p}\right)^{q}\Big)^{1/q} \leq c\delta^{s}\|f\|_{B^{s}_{pq}}. \end{split}$$

For the second inequality above we use Hölder's inequality if q > 1 and the *q*-inequality if  $0 < q \leq 1$ ; for the last inequality we used the fact that the definition of the Besov space  $B_{pq}^s$  is independent of the particular choice of the functions  $\Phi_0$  and  $\Phi$  satisfying (3.18)-(3.19) in its definition, hance  $\check{\Phi}$  produces an equivalent norm.

The case when  $q = \infty$  is as easy; we omit it. The proof of Proposition 4.6 is complete.

#### References

- P. Baldi, G. Kerkyacharian, D. Marinucci, D. Picard, Adaptive density estimation for directional data using needlets, Ann. Statist. 37 (2009), no. 6A, 3362–3395.
- [2] I. Castillo, G. Kerkyacharian, D. Picard, Thomas Bayes' walk on manifolds, Probab. Theory Relat. Fields, 158 (2014), no. 3-4, 665–710.
- [3] R. Coifman, G. Weiss, Analyse Harmonique Non-commutative sur Certains Espaces Homogenes. Lecture Notes in Math. Vol. 242. Springer, Berlin 1971.
- [4] T. Coulhon, G. Kerkyacharian, P. Petrushev, Heat Kernel Generated Frames in the Setting of Dirichlet Spaces, J. Fourier Anal. Appl. 18 (2012), no. 5, 995–1066.
- [5] F. Dai, Y. Xu, Approximation theory and harmonic analysis on spheres and balls, Springer Monographs in Mathematics, Springer 2013.
- [6] D. L. Donoho, I. M. Johnstone, G. Kerkyacharian, D. Picard, Density estimation by wavelet thresholding, Ann. Statist. 24 (1996), no. 2, 508–539.
- [7] M. Frazier, B. Jawerth, Decomposition of Besov spaces, Indiana Univ. Math. J. 34 (1985), 777–799.
- [8] M. Frazier, B. Jawerth, A discrete transform and decomposition of distribution spaces, J. Funct. Anal. 93 (1990), 34–170.
- [9] M. Frazier, B. Jawerth, G. Weiss, Littlewood-Paley theory and the study of function spaces, CBMS No 79 (1991), AMS.

#### 42 G. CLEANTHOUS, A. GEORGIADIS, G. KERKYACHARIAN, P. PETRUSHEV, D. PICARD

- [10] G. Folland, Real Analysis: Modern techniques and their applications, 2nd edn. Wiley-Interscience, New York, 1999.
- [11] A. G. Georgiadis, G. Kerkyacharian, G. Kyriazis, P. Petrushev, Homogeneous Besov and Triebel-Lizorkin spaces of distributions associated to non-negative self-adjoint operators, J. Math. Anal. Appl. 449 (2017), no. 2, 1382–1412.
- [12] A. Goldenshluger, O. Lepski, On adaptive minimax density estimator on R<sup>d</sup>, Probab. Theory Relat. Fields, 159 (2014), 479–543.
- [13] A. Grigor'yan, Heat kernel and analysis on manifold, AMS/IP studies in advance mathematics, 47, AMS Providence, RI; International Press, Boston, MA, 2009.
- [14] W. Härdle, G. Kerkyacharian, D. Picard, A. Tsybakov, Wavelets, approximation, and statistical applications, Lecture Notes in Statistics, 129, Springer-Verlag, New York, 1998.
- [15] I. M. Johnstone and B. Silverman, Speed of estimation in positron emission tomography and related inverse problems, Ann. of Statist. 18 (1990), 251–280.
- [16] A. Juditsky, S. Lambert-Lacroix, On minimax density estimation on ℝ, Bernoulli, 10 (2004), no. 2, 187–220.
- [17] G. Kerkyacharian, P. Petrushev, Heat kernel based decomposition of spaces of distributions in the framework of Dirichlet spaces, Trans. Amer. Math. Soc. 367 (2015), 121–189.
- [18] G. Kerkyacharian, P. Petrushev, D. Picard, T. Willer, Needlet algorithms for estimation in inverse problems, Electron. J. Stat. 1:30–76 (electronic), 2007.
- [19] G. Kerkyacharian, P. Petrushev, Y. Xu, Gaussian bounds for the weighted heat kernels on the interval, ball and simplex, Constr. Approx. to appear. arXiv:1801.07325
- [20] G. Kerkyacharian, P. Petrushev, Y. Xu, Gaussian bounds for the heat kernels on the ball and simplex: Classical approach, Studia Math. to appear. arXiv:1801.07326
- [21] G. Kerkyacharian, D. Picard, Density estimation in Besov spaces, Statist. Probab. Lett. 13 (1992), 15–24.
- [22] G. Kerkyacharian, T. M. Pham Ngoc, D. Picard, Localized spherical deconvolution, Ann. Statist. 39 (2) (2011), 1042–1068
- [23] O. V. Lepskiĭ, Asymptotically minimax adaptive estimation. I. Upper bounds. Optimally adaptive estimates. Teor. Veroyatnost. i Primenen. 36(4) (1991), 645–659.
- [24] O. V. Lepski, E. Mammen, V. G. Spokoiny, Optimal spatial adaptation to inhomogeneous smoothness: an approach based on kernel estimates with variable bandwidth selectors, Ann. Statist. 25 (1997), no. 3, 929–947.
- [25] D.Pollard, Convergence of stochastic Processes, Springer, New York, 1984.
- [26] B. W. Silverman, Density estimation for statistics and data analysis, Monographs on Statistics and Applied Probability, Chapman & Hall, London, 1986.
- [27] J. L. Starck, F. Murtagh, J. M. Fadili, Sparse image and signal processing: wavelets, curvelets, morphological diversity, Cambridge University Press, Cambradge, 2010.
- [28] E. Stein, G. Weiss, Introduction to Fourier analysis on Euclidean spaces, Princeton Mathematical Series, No. 32. Princeton University Press, Princeton, N.J., 1971.
- [29] A. B. Tsybakov, Introduction to nonparametric estimation, Revised and extended from the 2004 French original. Translated by Vladimir Zaiats. Springer Series in Statistics, Springer, New York, 2009.
- [30] K. Yosida, Functional Analysis, Springer, Berlin, 1980.

Department of Mathematics, Statistics and Physics, Newcastle University, Newcastle Upon Tyne, NE1 7RU, UK

E-mail address: galatia.cleanthous@newcastle.ac.uk

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CYPRUS, 1678 NICOSIA, CYPRUS *E-mail address*: gathana@ucy.ac.cy

UNIVERSITY PARIS DIDEROT-PARIS 7, LPMA, PARIS, FRANCE *E-mail address*: kerk@math.univ-paris-diderot.fr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA *E-mail address*: pencho@math.sc.edu

UNIVERSITY PARIS DIDEROT-PARIS 7, LPMA, PARIS, FRANCE *E-mail address*: picard@math.univ-paris-diderot.fr