

Exact asymptotic formulae of the stationary distribution of a discrete-time 2d-QBD process: an example and additional proofs

Toshihisa Ozawa[†] and Masahiro Kobayashi^{††}

[†]Faculty of Business Administration, Komazawa University

^{††}Department of Mathematical Science, Tokai University

[†]1-23-1 Komazawa, Setagaya-ku, Tokyo 154-8525, Japan

E-mail: [†]toshi@komazawa-u.ac.jp, ^{††}m_kobayashi@tsc.u-tokai.ac.jp

Abstract

A discrete-time two-dimensional quasi-birth-and-death process (2d-QBD process), denoted by $\{\mathbf{Y}_n\} = \{(X_{1,n}, X_{2,n}, J_n)\}$, is a two-dimensional skip-free random walk $\{(X_{1,n}, X_{2,n})\}$ on \mathbb{Z}_+^2 with a supplemental process $\{J_n\}$ on a finite set S_0 . The supplemental process $\{J_n\}$ is called a phase process. The 2d-QBD process $\{\mathbf{Y}_n\}$ is a Markov chain in which the transition probabilities of the two-dimensional process $\{(X_{1,n}, X_{2,n})\}$ vary according to the state of the phase process $\{J_n\}$. This modulation is assumed to be space homogeneous except for the boundaries of \mathbb{Z}_+^2 . Under certain conditions, the directional exact asymptotic formulae of the stationary distribution of the 2d-QBD process have been obtained in Ref. [7]. In this paper, we give an example of 2d-QBD process and proofs of some lemmas and propositions appeared in Ref. [7].

Key words: quasi-birth-and-death process, stationary distribution, asymptotic property, matrix analytic method, two-dimensional reflecting random walk

Mathematical Subject Classification: 60J10, 60K25

1 Introduction

A discrete-time two-dimensional quasi-birth-and-death process (2d-QBD process), denoted by $\{\mathbf{Y}_n\} = \{(X_{1,n}, X_{2,n}, J_n)\}$, is a two-dimensional skip-free random walk $\{(X_{1,n}, X_{2,n})\}$ on \mathbb{Z}_+^2 with a supplemental process $\{J_n\}$ on a finite set S_0 (see Ref. [6]). The supplemental process $\{J_n\}$ is called a phase process. The 2d-QBD process $\{\mathbf{Y}_n\}$ is a Markov chain in which the transition probabilities of the two-dimensional process $\{(X_{1,n}, X_{2,n})\}$ vary according to the state of the phase process $\{J_n\}$. This modulation is assumed to be space homogeneous except for the boundaries of \mathbb{Z}_+^2 . Assume that the 2d-QBD process $\{\mathbf{Y}_n\}$ is irreducible, aperiodic and positive recurrent, and denote by $\boldsymbol{\nu} = (\nu_{k,l}, (k,l) \in \mathbb{Z}_+^2)$ its stationary distribution, where $\nu_{k,l} = (\nu_{k,l,j}, j \in S_0)$ and $\nu_{k,l,j} = \lim_{n \rightarrow \infty} \mathbb{P}(\mathbf{Y}_n = (k,l,j))$. In Ref. [7], we have obtained, under certain conditions, the directional exact asymptotic formulae $h_1(k)$ and $h_2(k)$ that satisfy, for some nonzero vector \mathbf{c}_1 and \mathbf{c}_2 ,

$$\lim_{k \rightarrow \infty} \frac{\nu_{k,0}}{h_1(k)} = \mathbf{c}_1, \quad \lim_{k \rightarrow \infty} \frac{\nu_{0,k}}{h_2(k)} = \mathbf{c}_2. \quad (1.1)$$

In this paper, we give an example of 2d-QBD process and proofs of some lemmas and propositions appeared in Ref. [7], in order to make that paper easy to understand.

The rest of the paper is organized as follows. In Section 2, we summarize main results of Ref. [7] and related topics. Numerical examples are presented in Section 3, where a single-server two-queue model is considered. Proofs of some lemmas and propositions in Ref. [7] are given in Section 4.

Notation. A set \mathbb{H} is defined as $\mathbb{H} = \{-1, 0, 1\}$ and \mathbb{H}_+ as $\mathbb{H}_+ = \{0, 1\}$. For $a, b \in \mathbb{R}_+$, $\mathbb{C}[a, b]$ and $\mathbb{C}(a, b)$ are defined as $\mathbb{C}[a, b] = \{z \in \mathbb{C} : a \leq |z| \leq b\}$ and $\mathbb{C}(a, b) = \{z \in \mathbb{C} : a \leq |z| < b\}$, respectively. $\mathbb{C}(a, b)$ and $\mathbb{C}(a, b)$ are analogously defined. For $r > 0$, $\varepsilon > 0$ and $\theta \in [0, \pi/2)$, $\tilde{\Delta}_r(\varepsilon, \theta)$ is defined as

$$\tilde{\Delta}_r(\varepsilon, \theta) = \{z \in \mathbb{C} : |z| < r + \varepsilon, z \neq r, |\arg(z - r)| > \theta\}.$$

For $r > 0$, we denote by “ $\tilde{\Delta}_r \ni z \rightarrow r$ ” that $\tilde{\Delta}_r(\varepsilon, \theta) \ni z \rightarrow r$ for some $\varepsilon > 0$ and some $\theta \in [0, \pi/2)$. For a matrix $A = (a_{ij})$, we denote by $[A]_{ij}$ the (i, j) -entry of A . The transpose of A is denoted by A^\top . We denote by $\text{spr}(A)$ the spectral radius of A , which is the maximum modulus of eigenvalue of A . We denote by $|A|$ the matrix each of whose entries is the modulus of the corresponding entry of A , i.e., $|A| = (|a_{ij}|)$. O is a matrix of 0's, $\mathbf{1}$ is a column vector of 1's and $\mathbf{0}$ is a column vector of 0's; their dimensions are determined in context. I is the identity matrix. For a $k \times l$ matrix $A = (\mathbf{a}_1 \ \mathbf{a}_2 \ \cdots \ \mathbf{a}_l)$, $\text{vec}(A)$ is a $kl \times 1$ vector defined as

$$\text{vec}(A) = \begin{pmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \vdots \\ \mathbf{a}_l \end{pmatrix}.$$

For matrices A , B and C , the identity $\text{vec}(ABC) = (C^\top \otimes A) \text{vec}(B)$ holds (see, for example, Horn and Johnson [3]).

2 Model description and main results of Ref. [7]

Here we summarize Section 2 of Ref. [7].

2.1 2d-QBD process

Let $S_0 = \{1, 2, \dots, s_0\}$ be a finite set, where s_0 is the number of elements of S_0 . A 2d-QBD process $\{\mathbf{Y}_n\} = \{(X_{1,n}, X_{2,n}, J_n)\}$ is a discrete-time Markov chain on the state space $\mathcal{S} = \mathbb{Z}_+^2 \times S_0$. The transition probability matrix P of $\{\mathbf{Y}_n\}$ is represented in block form as

$$P = \left(P_{(x_1, x_2), (x'_1, x'_2)}; (x_1, x_2), (x'_1, x'_2) \in \mathbb{Z}_+^2 \right),$$

where each block $P_{(x_1, x_2), (x'_1, x'_2)}$ is given as $P_{(x_1, x_2), (x'_1, x'_2)} = \left(p_{(x_1, x_2, j), (x'_1, x'_2, j')}; j, j' \in S_0 \right)$ and for $(x_1, x_2, j), (x'_1, x'_2, j') \in \mathcal{S}$, $p_{(x_1, x_2, j), (x'_1, x'_2, j')} = \mathbb{P}(\mathbf{Y}_1 = (x'_1, x'_2, j') | \mathbf{Y}_0 = (x_1, x_2, j))$. The block matrices are given in terms of $s_0 \times s_0$ non-negative matrices

$$A_{i,j}, i, j \in \mathbb{H}, \quad A_{i,j}^{(1)}, i \in \mathbb{H}, j \in \mathbb{H}_+, \quad A_{i,j}^{(2)}, i \in \mathbb{H}_+, j \in \mathbb{H}, \quad A_{i,j}^{(0)}, i, j \in \mathbb{H}_+,$$

as follows: for $(x_1, x_2), (x'_1, x'_2) \in \mathbb{Z}_+^2$,

$$P_{(x_1, x_2), (x'_1, x'_2)} = \begin{cases} A_{\Delta x_1, \Delta x_2}, & \text{if } x_1 \neq 0, x_2 \neq 0, \Delta x_1, \Delta x_2 \in \mathbb{H}, \\ A_{\Delta x_1, \Delta x_2}^{(1)}, & \text{if } x_1 \neq 0, x_2 = 0, \Delta x_1 \in \mathbb{H}, \Delta x_2 \in \mathbb{H}_+, \\ A_{\Delta x_1, \Delta x_2}^{(2)}, & \text{if } x_1 = 0, x_2 \neq 0, \Delta x_1 \in \mathbb{H}_+, \Delta x_2 \in \mathbb{H}, \\ A_{\Delta x_1, \Delta x_2}^{(0)}, & \text{if } x_1 = x_2 = 0, \Delta x_1, \Delta x_2 \in \mathbb{H}_+, \\ O, & \text{otherwise,} \end{cases}$$

where $\Delta x_1 = x'_1 - x_1$ and $\Delta x_2 = x'_2 - x_2$. Define matrices $A_{*,*}$, $A_{*,*}^{(1)}$, $A_{*,*}^{(2)}$ and $A_{*,*}^{(0)}$ as

$$A_{*,*} = \sum_{i,j \in \mathbb{H}} A_{i,j}, \quad A_{*,*}^{(1)} = \sum_{i \in \mathbb{H}, j \in \mathbb{H}_+} A_{i,j}^{(1)}, \quad A_{*,*}^{(2)} = \sum_{i \in \mathbb{H}_+, j \in \mathbb{H}} A_{i,j}^{(2)}, \quad A_{*,*}^{(0)} = \sum_{i,j \in \mathbb{H}_+} A_{i,j}^{(0)}.$$

All of these matrices are stochastic. We assume the following condition.

Assumption 2.1. *The Markov chain $\{\mathbf{Y}_n\}$ is irreducible and aperiodic.*

We consider three kinds of Markov chain generated from $\{\mathbf{Y}_n\}$ by removing one or two boundaries and denote them by $\{\tilde{\mathbf{Y}}_n\} = \{(\tilde{X}_{1,n}, \tilde{X}_{2,n}, \tilde{J}_n)\}$, $\{\tilde{\mathbf{Y}}_n^{(1)}\} = \{(\tilde{X}_{1,n}^{(1)}, \tilde{X}_{2,n}^{(1)}, \tilde{J}_n^{(1)})\}$ and $\{\tilde{\mathbf{Y}}_n^{(2)}\} = \{(\tilde{X}_{1,n}^{(2)}, \tilde{X}_{2,n}^{(2)}, \tilde{J}_n^{(2)})\}$, respectively. $\{\tilde{\mathbf{Y}}_n\}$ is a Markov chain on the state space $\mathbb{Z}^2 \times S_0$ and it is generated from $\{\mathbf{Y}_n\}$ by removing the boundaries on the x_1 and x_2 -axes. For $(x_1, x_2, j), (x'_1, x'_2, j') \in \mathbb{Z}^2 \times S_0$, denote by $\tilde{p}_{(x_1, x_2, j), (x'_1, x'_2, j')}$ the transition probability $\mathbb{P}(\tilde{\mathbf{Y}}_1 = (x'_1, x'_2, j') \mid \tilde{\mathbf{Y}}_0 = (x_1, x_2, j))$. The transition probability matrix \tilde{P} of $\{\tilde{\mathbf{Y}}_n\}$ is represented in block form as

$$\tilde{P} = \left(\tilde{P}_{(x_1, x_2), (x'_1, x'_2)}; (x_1, x_2), (x'_1, x'_2) \in \mathbb{Z}^2 \right),$$

where $\tilde{P}_{(x_1, x_2), (x'_1, x'_2)} = \left(\tilde{p}_{(x_1, x_2, j), (x'_1, x'_2, j')}; j, j' \in S_0 \right)$; each block $\tilde{P}_{(x_1, x_2), (x'_1, x'_2)}$ is given as

$$\tilde{P}_{(x_1, x_2), (x'_1, x'_2)} = \begin{cases} A_{\Delta x_1, \Delta x_2}, & \text{if } \Delta x_1, \Delta x_2 \in \mathbb{H}, \\ O, & \text{otherwise,} \end{cases}$$

where $\Delta x_1 = x'_1 - x_1$ and $\Delta x_2 = x'_2 - x_2$. The Markov chain $\{\tilde{\mathbf{Y}}_n\}$ is a Markov chain generated by $\{A_{k,l} : k, l \in \mathbb{H}\}$. $\{\tilde{\mathbf{Y}}_n^{(1)}\}$ is a Markov chain on the state space $\mathbb{Z} \times \mathbb{Z}_+ \times S_0$ and it is generated from $\{\mathbf{Y}_n\}$ by removing the boundary on the x_2 -axes. For $(x_1, x_2, j), (x'_1, x'_2, j') \in \mathbb{Z} \times \mathbb{Z}_+ \times S_0$, denote by $\tilde{p}_{(x_1, x_2, j), (x'_1, x'_2, j')}^{(1)}$ the transition probability $\mathbb{P}(\tilde{\mathbf{Y}}_1^{(1)} = (x'_1, x'_2, j') \mid \tilde{\mathbf{Y}}_0^{(1)} = (x_1, x_2, j))$. The transition probability matrix $\tilde{P}^{(1)}$ of $\{\tilde{\mathbf{Y}}_n^{(1)}\}$ is represented in block form as

$$\tilde{P}^{(1)} = \left(\tilde{P}_{(x_1, x_2), (x'_1, x'_2)}^{(1)}; (x_1, x_2), (x'_1, x'_2) \in \mathbb{Z} \times \mathbb{Z}_+ \right),$$

where $\tilde{P}_{(x_1, x_2), (x'_1, x'_2)}^{(1)} = \left(\tilde{p}_{(x_1, x_2, j), (x'_1, x'_2, j')}^{(1)}; j, j' \in S_0 \right)$; each block $\tilde{P}_{(x_1, x_2), (x'_1, x'_2)}^{(1)}$ is given as

$$\tilde{P}_{(x_1, x_2), (x'_1, x'_2)}^{(1)} = \begin{cases} A_{\Delta x_1, \Delta x_2}, & \text{if } x_2 \neq 0, \Delta x_1, \Delta x_2 \in \mathbb{H}, \\ A_{\Delta x_1, \Delta x_2}^{(1)}, & \text{if } x_2 = 0, \Delta x_1 \in \mathbb{H}, \Delta x_2 \in \mathbb{H}_+, \\ O, & \text{otherwise,} \end{cases}$$

where $\Delta x_1 = x'_1 - x_1$ and $\Delta x_2 = x'_2 - x_2$. $\{\tilde{\mathbf{Y}}_n^{(2)}\}$ is a Markov chain on the state space $\mathbb{Z}_+ \times \mathbb{Z} \times S_0$ and it is generated from $\{\mathbf{Y}_n\}$ by removing the boundary on the x_1 -axes. The transition probability matrix $\tilde{P}^{(2)}$ of $\{\tilde{\mathbf{Y}}_n^{(2)}\}$ is analogously given. The Markov chain $\{\tilde{\mathbf{Y}}_n^{(1)}\}$ is a Markov chain generated by $\{\{A_{k,l} : k, l \in \mathbb{H}\}, \{A_{k,l}^{(1)} : k \in \mathbb{H}, l \in \mathbb{H}_+\}\}$ and $\{\tilde{\mathbf{Y}}_n^{(2)}\}$ is that generated by $\{\{A_{k,l} : k, l \in \mathbb{H}\}, \{A_{k,l}^{(2)} : k \in \mathbb{H}_+, l \in \mathbb{H}\}\}$. Hereafter, we assume the following condition.

Assumption 2.2. $\{\tilde{\mathbf{Y}}_n\}$, $\{\tilde{\mathbf{Y}}_n^{(1)}\}$ and $\{\tilde{\mathbf{Y}}_n^{(2)}\}$ are irreducible and aperiodic.

Under Assumption 2.2, $A_{*,*}$ is irreducible and aperiodic. $A_{*,*}$ is, therefore, positive recurrent and ergodic since its dimension is finite. We denote by $\boldsymbol{\pi}_{*,*}$ the stationary distribution of $A_{*,*}$.

A 2d-QBD process $\{\mathbf{Y}_n\} = \{(X_{1,n}, X_{2,n}, J_n)\}$ can be represented as a QBD process with a countable phase space in two ways: One is $\{\mathbf{Y}_n^{(1)}\} = \{(X_{1,n}, (X_{2,n}, J_n))\}$, where $X_{1,n}$ is the level and $(X_{2,n}, J_n)$ the phase, and the other $\{\mathbf{Y}_n^{(2)}\} = \{(X_{2,n}, (X_{1,n}, J_n))\}$, where $X_{2,n}$ is the level and $(X_{1,n}, J_n)$ the phase (see Subsection 2.3 of Ref. [7]). Let $R^{(1)}$ and $R^{(2)}$ be the rate matrices of the QBD processes $\{\mathbf{Y}^{(1)}\}$ and $\{\mathbf{Y}^{(2)}\}$, respectively. Hereafter, we assume the following condition.

Assumption 2.3. The rate matrices $R^{(1)}$ and $R^{(2)}$ are irreducible.

2.2 Stationary condition

We define induced Markov chains and the mean drift vectors derived from the induced Markov chains (see Ref. [1]). Since the 2d-QBD process is a kind of two-dimensional reflecting random walk, there exist three induced Markov chains: $\mathcal{L}^{\{1,2\}}$, $\mathcal{L}^{\{1\}}$ and $\mathcal{L}^{\{2\}}$. $\mathcal{L}^{\{1,2\}}$ is the phase process $\{\tilde{J}_n\}$ of $\{\tilde{\mathbf{Y}}_n\}$ and it is a Markov chain governed by the transition probability matrix $A_{*,*}$. The mean drift vector $\mathbf{a}^{\{1,2\}} = (a_1^{\{1,2\}}, a_2^{\{1,2\}})$ derived from $\mathcal{L}^{\{1,2\}}$ is given as

$$a_1^{\{1,2\}} = \boldsymbol{\pi}_{*,*}(-A_{-1,*} + A_{1,*})\mathbf{1}, \quad a_2^{\{1,2\}} = \boldsymbol{\pi}_{*,*}(-A_{*, -1} + A_{*, 1})\mathbf{1},$$

where for $k \in \mathbb{H}$, $A_{k,*} = \sum_{l \in \mathbb{H}} A_{k,l}$ and $A_{*,k} = \sum_{l \in \mathbb{H}} A_{l,k}$. Define block tri-diagonal transition probability matrices $A_*^{(1)}$ and $A_*^{(2)}$ as

$$A_*^{(1)} = \begin{pmatrix} A_{*,0}^{(1)} & A_{*,1}^{(1)} & & & \\ A_{*, -1} & A_{*,0} & A_{*,1} & & \\ & A_{*, -1} & A_{*,0} & A_{*,1} & \\ & & \ddots & \ddots & \ddots \end{pmatrix}, \quad A_*^{(2)} = \begin{pmatrix} A_{0,*}^{(2)} & A_{1,*}^{(2)} & & & \\ A_{-1,*} & A_{0,*} & A_{1,*} & & \\ & A_{-1,*} & A_{0,*} & A_{1,*} & \\ & & \ddots & \ddots & \ddots \end{pmatrix},$$

where for $k \in \mathbb{H}_+$, $A_{*,k}^{(1)} = \sum_{l \in \mathbb{H}} A_{l,k}^{(1)}$ and $A_{k,*}^{(2)} = \sum_{l \in \mathbb{H}} A_{k,l}^{(2)}$. Under Assumption 2.2, $A_*^{(1)}$ and $A_*^{(2)}$ are irreducible and aperiodic. $\mathcal{L}^{\{1\}}$ (resp. $\mathcal{L}^{\{2\}}$) is a partial process $\{(\tilde{X}_{2,n}^{(1)}, \tilde{J}_n^{(1)})\}$ of $\{\tilde{\mathbf{Y}}_n^{(1)}\}$ (resp. $\{(\tilde{X}_{1,n}^{(2)}, \tilde{J}_n^{(2)})\}$ of $\{\tilde{\mathbf{Y}}_n^{(2)}\}$) and it is a Markov chain governed by $A_*^{(1)}$ (resp. $A_*^{(2)}$). Since $\mathcal{L}^{\{1\}}$ (resp. $\mathcal{L}^{\{2\}}$) is one-dimensional QBD process, it is positive recurrent if and only if $a_2^{\{1,2\}} < 0$ (resp. $a_1^{\{1,2\}} < 0$). Denote by $\boldsymbol{\pi}_*^{(1)} = (\boldsymbol{\pi}_{*,k}^{(1)}, k \in \mathbb{Z}_+)$ and $\boldsymbol{\pi}_*^{(2)} = (\boldsymbol{\pi}_{*,k}^{(2)}, k \in \mathbb{Z}_+)$ the stationary distributions

of $\mathcal{L}^{\{1\}}$ and $\mathcal{L}^{\{2\}}$, respectively, if they exist. Then, the mean drift vectors $\mathbf{a}^{\{1\}} = (a_1^{\{1\}}, a_2^{\{1\}})$ and $\mathbf{a}^{\{2\}} = (a_1^{\{2\}}, a_2^{\{2\}})$ derived from $\mathcal{L}^{\{1\}}$ and $\mathcal{L}^{\{2\}}$ are given as

$$\begin{aligned} a_1^{\{1\}} &= \boldsymbol{\pi}_{*,0}^{(1)}(-A_{-1,0}^{(1)} - A_{-1,1}^{(1)} + A_{1,0}^{(1)} + A_{1,1}^{(1)}) \mathbf{1} + \boldsymbol{\pi}_{*,1}^{(1)}(I - R_*^{(1)})^{-1}(-A_{-1,*} + A_{1,*}) \mathbf{1}, \\ a_2^{\{1\}} &= 0, \quad a_1^{\{2\}} = 0, \\ a_2^{\{2\}} &= \boldsymbol{\pi}_{*,0}^{(2)}(-A_{0,-1}^{(2)} - A_{1,-1}^{(2)} + A_{0,1}^{(2)} + A_{1,1}^{(2)}) \mathbf{1} + \boldsymbol{\pi}_{*,1}^{(2)}(I - R_*^{(2)})^{-1}(-A_{*, -1} + A_{*, 1}) \mathbf{1}, \end{aligned}$$

where $R_*^{(1)}$ and $R_*^{(2)}$ are the rate matrices of $A_*^{(1)}$ and $A_*^{(2)}$, respectively. The condition ensuring $\{\mathbf{Y}_n\}$ is positive recurrent or transient is given by Lemma 2.1 of Ref. [7]. In order for the 2d-QBD process $\{\mathbf{Y}_n\}$ to be positive recurrent, we assume the following condition.

Assumption 2.4. $a_1^{\{1,2\}}$ or $a_2^{\{1,2\}}$ is negative. If $a_1^{\{1,2\}} < 0$ and $a_2^{\{1,2\}} < 0$, then $a_1^{\{1\}} < 0$ and $a_2^{\{2\}} < 0$; if $a_1^{\{1,2\}} > 0$ and $a_2^{\{1,2\}} < 0$, then $a_1^{\{1\}} < 0$; if $a_1^{\{1,2\}} < 0$ and $a_2^{\{1,2\}} > 0$, then $a_2^{\{2\}} < 0$.

Denote by $\boldsymbol{\nu} = (\nu_{k,l,j}, (k,l,j) \in \mathbb{Z}_+^2 \times S_0)$ the stationary distribution of $\{\mathbf{Y}_n\}$. $\boldsymbol{\nu}$ is represented in block form as $\boldsymbol{\nu} = (\boldsymbol{\nu}_{k,l}, (k,l) \in \mathbb{Z}_+^2)$, where $\boldsymbol{\nu}_{k,l} = (\nu_{k,l,j}, j \in S_0)$.

2.3 Directional geometric decay rates

Let z_1 and z_2 be positive real numbers and define a matrix function $C(z_1, z_2)$ as

$$C(z_1, z_2) = \sum_{i,j \in \mathbb{H}} A_{i,j} z_1^i z_2^j = \sum_{j \in \mathbb{H}} A_{*,j}(z_1) z_2^j = \sum_{i \in \mathbb{H}} A_{i,*}(z_2) z_1^i,$$

where

$$A_{*,j}(z_1) = \sum_{i \in \mathbb{H}} A_{i,j} z_1^i, \quad A_{i,*}(z_2) = \sum_{j \in \mathbb{H}} A_{i,j} z_2^j.$$

The matrix function $C(z_1, z_2)$ is nonnegative and, under Assumption 2.2, it is irreducible and aperiodic. Let $\chi(z_1, z_2)$ be the Perron-Frobenius eigenvalue of $C(z_1, z_2)$, i.e., $\chi(z_1, z_2) = \text{spr}(C(z_1, z_2))$, and let $\mathbf{u}^C(z_1, z_2)$ and $\mathbf{v}^C(z_1, z_2)$ be the Perron-Frobenius left and right eigenvectors satisfying $\mathbf{u}^C(z_1, z_2) \mathbf{v}^C(z_1, z_2) = 1$. We have $\mathbf{u}^C(1, 1) = \boldsymbol{\pi}_{*,*}$ and $\mathbf{v}^C(1, 1) = \mathbf{1}$. Define a closed set $\bar{\Gamma}$ as

$$\bar{\Gamma} = \{(s_1, s_2) \in \mathbb{R}^2 : \chi(e^{s_1}, e^{s_2}) \leq 1\}.$$

Since $\chi(1, 1) = \chi(e^0, e^0) = 1$, $\bar{\Gamma}$ contains the point of $(0, 0)$ and thus it is not empty. By Proposition 2.3 and Lemma 2.2 of Ref. [7], $\bar{\Gamma}$ is a bounded convex set. Furthermore, by Lemma 2.3 of Ref. [7], the closed set $\bar{\Gamma}$ is not a singleton. For $i \in \{1, 2\}$, define the lower and upper extreme values of $\bar{\Gamma}$ with respect to s_i , denoted by \underline{s}_i^* and \bar{s}_i^* , as

$$\underline{s}_i^* = \min_{(s_1, s_2) \in \bar{\Gamma}} s_i, \quad \bar{s}_i^* = \max_{(s_1, s_2) \in \bar{\Gamma}} s_i,$$

where $-\infty < \underline{s}_i^* \leq 0$ and $0 \leq \bar{s}_i^* < \infty$. For $i \in \{1, 2\}$, define \underline{z}_i^* and \bar{z}_i^* as $\underline{z}_i^* = e^{\underline{s}_i^*}$ and $\bar{z}_i^* = e^{\bar{s}_i^*}$, respectively, where $0 < \underline{z}_i^* \leq 1$ and $1 \leq \bar{z}_i^* < \infty$. By Proposition 2.4 of Ref. [7], for each $z_1 \in (\underline{z}_1^*, \bar{z}_1^*)$ (resp. $z_2 \in (\underline{z}_2^*, \bar{z}_2^*)$), equation $\chi(z_1, z_2) = 1$ has just two different real solutions $\underline{z}_2(z_1)$ and $\bar{z}_2(z_1)$

(resp. $\zeta_1(z_2)$ and $\bar{\zeta}_1(z_2)$), where $\zeta_2(z_1) < \bar{\zeta}_2(z_1)$ (resp. $\zeta_1(z_2) < \bar{\zeta}_1(z_2)$). For $z_1 = \underline{z}_1^*$ or \bar{z}_1^* (resp. $z_2 = \underline{z}_2^*$ or \bar{z}_2^*), it has just one real solution $\zeta_2(z_1) = \bar{\zeta}_2(z_1)$ (resp. $\zeta_1(z_2) = \bar{\zeta}_1(z_2)$). If $z_1 \notin [\underline{z}_1^*, \bar{z}_1^*]$ (resp. $z_2 \notin [\underline{z}_2^*, \bar{z}_2^*]$), it has no real solutions.

Consider the following matrix quadratic equations of X :

$$A_{*, -1}(z_1) + A_{*, 0}(z_1)X + A_{*, 1}(z_1)X^2 = X, \quad (2.1)$$

$$A_{-1, *}(z_2) + A_{0, *}(z_2)X + A_{1, *}(z_2)X^2 = X. \quad (2.2)$$

Denote by $G_1(z_1)$ and $G_2(z_2)$ the minimum nonnegative solutions to matrix equations (2.1) and (2.2), respectively. Furthermore, consider the following matrix quadratic equations of X :

$$X^2 A_{*, -1}(z_1) + X A_{*, 0}(z_1) + A_{*, 1}(z_1) = X, \quad (2.3)$$

$$X^2 A_{-1, *}(z_2) + X A_{0, *}(z_2) + A_{1, *}(z_2) = X. \quad (2.4)$$

Denote by $R_1(z_1)$ and $R_2(z_2)$ the minimum nonnegative solutions to matrix equations (2.3) and (2.4), respectively. By Lemma 2.4 of Ref. [7], for $z \in \mathbb{R}_+ \setminus \{0\}$, the minimum nonnegative solutions $G_1(z)$ and $R_1(z)$ (resp. $G_2(z)$ and $R_2(z)$) to matrix equations (2.1) and (2.3) (resp. equations (2.2) and (2.4)) exist if and only if $z \in [\underline{z}_1^*, \bar{z}_1^*]$ (resp. $z \in [\underline{z}_2^*, \bar{z}_2^*]$). Furthermore, by Proposition 2.5 of Ref. [7], we have, for $z_1 \in [\underline{z}_1^*, \bar{z}_1^*]$ and $z_2 \in [\underline{z}_2^*, \bar{z}_2^*]$,

$$\text{spr}(G_1(z_1)) = \zeta_2(z_1), \quad \text{spr}(R_1(z_1)) = \bar{\zeta}_2(z_1)^{-1}, \quad (2.5)$$

$$\text{spr}(G_2(z_2)) = \zeta_1(z_2), \quad \text{spr}(R_2(z_2)) = \bar{\zeta}_1(z_2)^{-1}. \quad (2.6)$$

Define matrix functions $C_1(z_1, X)$ and $C_2(X, z_2)$ as

$$C_1(z_1, X) = A_{*, 0}^{(1)}(z_1) + A_{*, 1}^{(1)}(z_1)X, \quad C_2(X, z_2) = A_{0, *}^{(2)}(z_2) + A_{1, *}^{(2)}(z_2)X, \quad (2.7)$$

where X is an $s_0 \times s_0$ matrix and, for $i \in \mathbb{H}_+$,

$$A_{*, i}^{(1)}(z_1) = \sum_{j \in \mathbb{H}} A_{j, i}^{(1)} z_1^j, \quad A_{i, *}^{(2)}(z_2) = \sum_{j \in \mathbb{H}} A_{i, j}^{(2)} z_2^j.$$

Define $\psi_1(z_1)$ and $\psi_2(z_2)$ as

$$\psi_1(z_1) = \text{spr}(C_1(z_1, G_1(z_1))), \quad \psi_2(z_2) = \text{spr}(C_2(G_2(z_2), z_2)).$$

Next, we introduce points $(\theta_1^{(c)}, \theta_2^{(c)})$ and $(\eta_1^{(c)}, \eta_2^{(c)})$ on the closed curve $\chi(e^{s_1}, e^{s_2}) = 1$, as follows (see Fig. 1):

$$\theta_1^{(c)} = \max\{s_1 \in [0, \bar{s}_1^*] : \psi_1(e^{s_1}) \leq 1\}, \quad \theta_2^{(c)} = \log \zeta_2(e^{\theta_1^{(c)}}),$$

$$\eta_2^{(c)} = \max\{s_2 \in [0, \bar{s}_2^*] : \psi_2(e^{s_2}) \leq 1\}, \quad \eta_1^{(c)} = \log \zeta_1(e^{\eta_2^{(c)}}).$$

Furthermore, define $\bar{\theta}_2^{(c)}$ and $\bar{\eta}_1^{(c)}$ as

$$\bar{\theta}_2^{(c)} = \log \bar{\zeta}_2(e^{\theta_1^{(c)}}), \quad \bar{\eta}_1^{(c)} = \log \bar{\zeta}_1(e^{\eta_2^{(c)}}).$$




Figure 1: Configuration of $(\theta_1^{(c)}, \theta_2^{(c)})$ and $(\eta_1^{(c)}, \eta_2^{(c)})$

By Lemma 2.5 of Ref. [7], we see that $\theta_1^{(c)}$ and $\eta_2^{(c)}$ are always positive. Since $\bar{\Gamma}$ is convex, we can classify the possible configuration of points $(\theta_1^{(c)}, \theta_2^{(c)})$ and $(\eta_1^{(c)}, \eta_2^{(c)})$ in the following manner (see Fig. 1).

$$\begin{aligned} \text{Type I: } & \eta_1^{(c)} < \theta_1^{(c)} \text{ and } \theta_2^{(c)} < \eta_2^{(c)}, & \text{Type II: } & \eta_1^{(c)} < \theta_1^{(c)} \text{ and } \eta_2^{(c)} \leq \theta_2^{(c)}, \\ \text{Type III: } & \theta_1^{(c)} \leq \eta_1^{(c)} \text{ and } \theta_2^{(c)} < \eta_2^{(c)}. \end{aligned}$$

Define the directional decay rates, ξ_1 and ξ_2 , of the stationary distribution of the 2d-QBD process as

$$\begin{aligned} \xi_1 &= - \lim_{n \rightarrow \infty} \frac{1}{n} \log \nu_{n,j,k} \quad \text{for any } j \text{ and } k, \\ \xi_2 &= - \lim_{n \rightarrow \infty} \frac{1}{n} \log \nu_{i,n,k} \quad \text{for any } i \text{ and } k. \end{aligned}$$

By Lemma 2.6 of Ref. [7], ξ_1 and ξ_2 are given as follows.

$$(\xi_1, \xi_2) = \begin{cases} (\theta_1^{(c)}, \eta_2^{(c)}), & \text{Type I,} \\ (\bar{\eta}_1^{(c)}, \eta_2^{(c)}), & \text{Type II,} \\ (\theta_1^{(c)}, \bar{\theta}_2^{(c)}), & \text{Type III.} \end{cases} \quad (2.8)$$

The directional geometric decay rates r_1 in x_1 -coordinate and r_2 in x_2 -coordinate are given as $r_1 = e^{\xi_1}$ and $r_2 = e^{\xi_2}$.

2.4 Main results of Ref. [7]

We assume the following technical condition.

Assumption 2.5. *All the eigenvalues of $G_1(r_1)$ are distinct. Also, those of $G_2(r_2)$ are distinct.*

The exact asymptotic formulae $h_1(k)$ in x_1 -coordinate and $h_2(k)$ in x_2 -coordinate are given as follows.

Theorem 2.1 (Theorem 2.1 of Ref. [7]). *Under Assumptions 2.1 through 2.5, in the case of Type I, the exact asymptotic formulae are given as*

$$h_1(k) = \begin{cases} r_1^{-k}, & \psi_1(\bar{z}_1^*) > 1, \\ k^{-\frac{1}{2}(2l_1-1)}(\bar{z}_1^*)^{-k}, & \psi_1(\bar{z}_1^*) = 1, \\ k^{-\frac{1}{2}(2l_1+1)}(\bar{z}_1^*)^{-k}, & \psi_1(\bar{z}_1^*) < 1, \end{cases} \quad h_2(k) = \begin{cases} r_2^{-k}, & \psi_2(\bar{z}_2^*) > 1, \\ k^{-\frac{1}{2}(2l_2-1)}(\bar{z}_2^*)^{-k}, & \psi_2(\bar{z}_2^*) = 1, \\ k^{-\frac{1}{2}(2l_2+1)}(\bar{z}_2^*)^{-k}, & \psi_2(\bar{z}_2^*) < 1, \end{cases} \quad (2.9)$$

where l_1 and l_2 are some positive integers. In the case of Type II, they are given as

$$h_1(k) = \begin{cases} r_1^{-k}, & \eta_2^{(c)} < \theta_2^{(c)}, \\ k r_1^{-k}, & \eta_2^{(c)} = \theta_2^{(c)} \text{ and } \psi_1(\bar{z}_1^*) > 1, \\ (\bar{z}_1^*)^{-k}, & \eta_2^{(c)} = \theta_2^{(c)} \text{ and } \psi_1(\bar{z}_1^*) = 1, \\ k^{-\frac{1}{2}}(\bar{z}_1^*)^{-k}, & \eta_2^{(c)} = \theta_2^{(c)} \text{ and } \psi_1(\bar{z}_1^*) < 1, \end{cases} \quad h_2(k) = r_2^{-k}. \quad (2.10)$$

In the case of Type III, they are given as

$$h_1(k) = r_1^{-k}, \quad h_2(k) = \begin{cases} r_2^{-k}, & \theta_1^{(c)} < \eta_1^{(c)}, \\ k r_2^{-k}, & \theta_1^{(c)} = \eta_1^{(c)} \text{ and } \psi_2(\bar{z}_2^*) > 1, \\ (\bar{z}_2^*)^{-k}, & \theta_1^{(c)} = \eta_1^{(c)} \text{ and } \psi_2(\bar{z}_2^*) = 1, \\ k^{-\frac{1}{2}}(\bar{z}_2^*)^{-k}, & \theta_1^{(c)} = \eta_1^{(c)} \text{ and } \psi_2(\bar{z}_2^*) < 1. \end{cases} \quad (2.11)$$

3 An example

We consider the same queueing model as that used in Ozawa [6]. It is a single-server two-queue model in which the server visits the queues alternatively, serves one queue (denoted by Q_1) according to a 1-limited service and the other queue (denoted by Q_2) according to an exhaustive-type K -limited service (see Fig. 2). Customers in class 1 arrive at Q_1 according to a Poisson process with intensity λ_1 , those in class 2 arrive at Q_2 according to another Poisson process with intensity λ_2 , and they are mutually independent. Service times of class-1 customers are exponentially distributed with mean $1/\mu_1$, those of class-2 customers are also exponentially distributed with mean $1/\mu_2$, and they are mutually independent. The arrival processes and service times are also mutually independent. We define ρ_i , $i = 1, 2$, λ and ρ as $\rho_i = \lambda_i/\mu_i$, $\lambda = \lambda_1 + \lambda_2$ and $\rho = \rho_1 + \rho_2$, respectively. We refer to this model as a $(1, K)$ -limited service model.




Figure 2: Single server two-queue model

Let $X_1(t)$ be the number of customers at Q_1 at time t , $X_2(t)$ that of customers at Q_2 and $J(t) \in S_0 = \{0, 1, \dots, K\}$ the server state. When $X_1(t) = X_2(t) = 0$, $J(t)$ takes one of the states in S_0 at random; When $X_1(t) \geq 1$ and $X_2(t) = 0$, it also takes one of the states in S_0 at random; When $X_1(t) = 0$ and $X_2(t) \geq 1$, it takes the state of 0 or 1 at random if the server is serving the K -th customer in class 2 during a visit of the server at Q_2 and takes the state of $j \in \{2, \dots, K\}$ if the server is serving the $(K - j + 1)$ -th customer in class 2; When $X_1(t) \geq 1$ and $X_2(t) \geq 1$, it takes the state of 0 if the server is serving a class-1 customer and takes the state of $j \in \{1, \dots, K\}$ if the server is serving the $(K - j + 1)$ -th customer in class 2 during a visit of the server at Q_2 . The process $\{(X_1(t), X_2(t), J(t))\}$ becomes a continuous-time 2d-QBD process on the state space $\mathcal{S} = \mathbb{Z}_+^2 \times S_0$. By uniformization with parameter $\nu = \lambda + \mu_1 + \mu_2$, we obtain the corresponding discrete-time 2d-QBD process, $\{(X_{1,n}, X_{2,n}, J_n)\}$, governed by the following $(K+1) \times (K+1)$ block matrices:

$$A_{1,0} = \frac{\lambda_1}{\nu} I, \quad A_{0,1} = \frac{\lambda_2}{\nu} I, \quad A_{1,1} = A_{1,-1} = A_{-1,1} = A_{-1,-1} = O,$$

$$A_{-1,0} = \frac{\mu_1}{\nu} \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 0 & \cdots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 0 & 0 \end{pmatrix}, \quad A_{0,-1} = \frac{\mu_2}{\nu} \begin{pmatrix} 0 & & & \\ 1 & 0 & & \\ & \ddots & \ddots & \\ & & & 1 & 0 \end{pmatrix},$$

$$A_{0,0} = I + \frac{1}{\nu} \begin{pmatrix} -(\lambda + \mu_1) & & & \\ & -(\lambda + \mu_2) & & \\ & & \ddots & \\ & & & -(\lambda + \mu_2) \end{pmatrix},$$

$$A_{1,0}^{(1)} = \frac{\lambda_1}{\nu} I, \quad A_{0,0}^{(1)} = \left(1 - \frac{(\lambda + \mu_1)}{\nu}\right) I, \quad A_{1,1}^{(1)} = A_{-1,1}^{(1)} = O,$$

$$A_{0,1}^{(1)} = \frac{\lambda_2}{\nu} \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & \cdots & 0 \end{pmatrix}, \quad A_{-1,0}^{(1)} = \frac{\mu_1}{\nu(K+1)} \begin{pmatrix} 1 & \cdots & 1 \\ \vdots & \ddots & \vdots \\ 1 & \cdots & 1 \end{pmatrix},$$

$$A_{0,1}^{(2)} = \frac{\lambda_2}{\nu} I, \quad A_{0,0}^{(2)} = \left(1 - \frac{(\lambda + \mu_2)}{\nu}\right) I, \quad A_{1,1}^{(2)} = A_{1,-1}^{(2)} = O,$$

$$A_{1,0}^{(2)} = \frac{\lambda_1}{\nu} \begin{pmatrix} 0 & 1 & 0 & & \\ 0 & 1 & 0 & & \\ 0 & 0 & 1 & & \\ & & \ddots & \ddots & \\ & & & 0 & 1 \end{pmatrix}, \quad A_{0,-1}^{(2)} = \frac{\mu_2}{\nu} \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1/2 & 1/2 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ & & & \ddots & \ddots & \\ 0 & 0 & 0 & \cdots & 1 & 0 \end{pmatrix},$$

$$A_{1,0}^{(0)} = \frac{\lambda_1}{\nu} I, \quad A_{1,1}^{(0)} = O, \quad A_{0,0}^{(0)} = \left(1 - \frac{\lambda}{\nu}\right) I, \quad A_{0,1}^{(0)} = \frac{\lambda_2}{\nu} \begin{pmatrix} 0 & \cdots & 0 & 1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 0 & 1 \\ 0 & \cdots & 0 & 1 \end{pmatrix}.$$

Here we note that, because of the form of $A_{-1,0}$, only the $(K+1)$ -th column of $G_2(w)$ is nonzero for all $w \in [z_2^*, \bar{z}_2^*]$ and the value of 0 is always the eigenvalue of $G_2(w)$ with algebraic multiplicity K . Hence, $G_2(r_2)$ does not satisfy Assumption 2.5 but, in this case, we can give $G_2(w)$ in a specific form. $G_2(w)$ as well as $G_1(z)$ can analogously be defined for a complex variable (see Section 4 of Ref. [7]). For $z, w \in \mathbb{C}$, define a matrix function $L(z, w)$ as

$$L(z, w) = zw(C(z, w) - I) = zA_{*, -1}(z) + z(A_{*, 0}(z) - I)w + zA_{*, 1}(z)w^2, \quad (3.1)$$

and denote by $\phi(z, w)$ the determinant of $L(z, w)$, i.e., $\phi(z, w) = \det L(z, w)$. $L(z, w)$ is a $(K+1) \times (K+1)$ matrix polynomial in w with degree 2 and every entry of $L(z, w)$ is a polynomial in z and w . $\phi(z, w)$ is also a polynomial in z and w . For $w \in \mathbb{C}[z_2^*, \bar{z}_2^*]$, let $\alpha(w)$ be the solution to $\phi(z, w) = 0$ that corresponds to $\zeta_1(w)$ when $w \in [z_2^*, \bar{z}_2^*]$. Let $\mathbf{v}(w)$ be the column vector satisfying $L(\alpha(w), w)\mathbf{v}(w) = \mathbf{0}$. This $\mathbf{v}(w)$ is the right eigenvector of $G_2(w)$ with respect to the eigenvalue $\alpha(w)$. If $[\mathbf{v}(w)]_{K+1} \neq 0$, $G_2(w)$ is given as

$$G_2(w) = \begin{pmatrix} \mathbf{0} & \cdots & \mathbf{0} & \frac{\alpha(w)}{[\mathbf{v}(w)]_{K+1}} \mathbf{v}(w) \end{pmatrix}, \quad (3.2)$$

and it is decomposed as

$$G_2(w) = V_2(w)J_2(w)V_2(w)^{-1}, \quad (3.3)$$

where

$$J_2(w) = \text{diag}(0, \dots, 0, \alpha(w)), \quad (3.4)$$

$$V_2(w) = \begin{pmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \cdots & \mathbf{e}_K & \frac{1}{[\mathbf{v}(w)]_{K+1}} \mathbf{v}(w) \end{pmatrix}, \quad (3.5)$$

$$V_2(w)^{-1} = \begin{pmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \cdots & \mathbf{e}_K & 2\mathbf{e}_{K+1} - \frac{1}{[\mathbf{v}(w)]_{K+1}} \mathbf{v}(w) \end{pmatrix} \quad (3.6)$$

and, for $i \in \{1, 2, \dots, K+1\}$, \mathbf{e}_i is the i -th unit vector of dimension $K+1$. For any $w' \in \mathbb{C}$, if $\alpha(w)$ is analytic at $w = w'$, $G_2(w)$ is also entry-wise analytic at $w = w'$. Hence, the results of Lemma 4.7 of Ref. [7] hold for this $G_2(w)$. Furthermore, because of the decomposition of $G_2(w)$ above, the results in Section 5 of Ref. [7] also hold for $\varphi_2(w) = \sum_{j=1}^{\infty} \nu_{0,j} w^j$. Therefore, in the $(1, K)$ -limited service model, the results of Theorem 2.1 hold for the exact asymptotic formula $h_2(k)$.

In Tables 1 and 2, we show numerical examples of the geometric decay rates r_1 and r_2 in the $(1, K)$ -limited service model, where the value of K is varied. We have numerically verified that $G_1(r_1)$ satisfies Assumption 2.5 for all the numerical examples. In both the tables, the type of configuration of points $(\theta_1^{(c)}, \theta_2^{(c)})$ and $(\eta_1^{(c)}, \eta_2^{(c)})$ is Type I for all the values of K . We see from the tables that if $K \geq 6$, the exact asymptotic function $h_1(k)$ is geometric and if $K < 6$, it is not. On the other hand, in the case of Table 1, the exact asymptotic function $h_2(k)$ is not geometric for any value of K ; in the case of Table 2, if $K \leq 2$, $h_2(k)$ is geometric and $K > 2$, it is not.

Table 1: Directional geometric asymptotic rates ($\lambda_1 = 0.3, \lambda_2 = 0.3, \mu_1 = 1, \mu_2 = 1, \rho = 0.6$)

K	Type	$a_1^{\{1,2\}}$	$a_2^{\{1,2\}}$	$\psi(\bar{z}_1^*) - 1$	$\psi(\bar{z}_2^*) - 1$	r_1	r_2
1	I	-0.0769	-0.0769	-	-	1.968	1.968
2	I	-0.0128	-0.141	-	-	1.706	2.698
3	I	0.0192	-0.173	-	-	1.674	3.171
4	I	0.0385	-0.192	-	-	1.668	3.464
5	I	0.0513	-0.205	-	-	1.667	3.659
6	I	0.0604	-0.214	+	-	1.667	3.797
7	I	0.0673	-0.221	+	-	1.667	3.899
8	I	0.0726	-0.226	+	-	1.667	3.978
10	I	0.0804	-0.234	+	-	1.667	4.091
20	I	0.0971	-0.251	+	-	1.667	4.308
50	I	0.108	-0.262	+	-	1.667	4.421

Table 2: Directional geometric asymptotic rates ($\lambda_1 = 0.24, \lambda_2 = 0.7, \mu_1 = 1.2, \mu_2 = 1, \rho = 0.9$)

K	Type	$a_1^{\{1,2\}}$	$a_2^{\{1,2\}}$	$\psi(\bar{z}_1^*) - 1$	$\psi(\bar{z}_2^*) - 1$	r_1	r_2
1	I	-0.0973	0.0492	-	+	3.646	1.116
2	I	-0.0360	-0.00187	-	+	1.844	1.116
3	I	-0.00665	-0.0263	-	-	1.183	1.135
4	I	0.0105	-0.0406	-	-	1.102	1.273
5	I	0.0218	-0.0500	-	-	1.095	1.392
6	I	0.0298	-0.0567	+	-	1.095	1.481
7	I	0.0358	-0.0617	+	-	1.095	1.549
8	I	0.0404	-0.0655	+	-	1.095	1.603
10	I	0.0470	-0.0710	+	-	1.095	1.682
20	I	0.0611	-0.0828	+	-	1.095	1.859
50	I	0.0702	-0.0903	+	-	1.095	1.971

Remark 3.1. Consider a general 2d-QBD process. The results of Theorem 2.1 also hold for the case where several columns of $G_1(z)$ or $G_2(w)$ are zero. Here we only consider the case where several columns of $G_2(w)$ are zero. For simplicity, assume that, for some $k_0 \in \{1, 2, \dots, s_0 - 1\}$ and for any $w \in [\underline{z}_2^*, \bar{z}_2^*]$, the first k_0 columns of $G_2(w)$ are always zero. Further assume that the value of 0 is the eigenvalue of $G_2(r_2)$ with algebraic multiplicity k_0 and the other $s_0 - k_0$ eigenvalues of $G_2(r_2)$ are nonzero and distinct. $(1, K)$ -limited service models in which the Poisson arrival processes are replaced with Markovian arrival processes are like this.

For $w \in \mathbb{C}[\underline{z}_2^*, \bar{z}_2^*]$, let $\alpha_1(w), \alpha_2(w), \dots, \alpha_{s_0}(w)$ be the solutions to $\phi(z, w) = 0$, counting multiplicity, that satisfy $|\alpha_k(w)| \leq \zeta_1(w)$, $k = 1, 2, \dots, s_0$. By Lemma 4.3 of Ref. [7], these solutions are the eigenvalues of $G_2(w)$. Without loss of generality, we assume that $\alpha_k(w) \equiv 0$ for $k \in \{1, 2, \dots, k_0\}$ and $\alpha_{s_0}(w)$ corresponds to $\zeta_1(w)$ when $w \in [\underline{z}_2^*, \bar{z}_2^*]$. For $k \in \{k_0 + 1, k_0 + 2, \dots, s_0\}$, let $\mathbf{v}_k(w)$ be the column vector satisfying $L(\alpha_k(w), w)\mathbf{v}_k(w) = \mathbf{0}$. Each $\mathbf{v}_k(w)$ is the right eigenvector of $G_2(w)$ with respect to the eigenvalue $\alpha_k(w)$. Define $s_0 \times s_0$ matrix functions $J_2(w)$ and $V_2(w)$ as

$$J_2(w) = \text{diag}(0, \dots, 0, \alpha_{k_0+1}(w), \alpha_{k_0+2}(w), \dots, \alpha_{s_0}(w)), \quad (3.7)$$

$$V_2(w) = (\mathbf{e}_1 \quad \mathbf{e}_2 \quad \cdots \quad \mathbf{e}_{k_0} \quad \mathbf{v}_{k_0+1}(w) \quad \mathbf{v}_{k_0+2}(w) \quad \cdots \quad \mathbf{v}_{s_0}(w)), \quad (3.8)$$

If $\alpha_{k_0+1}(w), \alpha_{k_0+2}(w), \dots, \alpha_{s_0}(w)$ are distinct, $V_2(w)$ is nonsingular and $G_2(w)$ is given as

$$G_2(w) = V_2(w)J_2(w)V_2(w)^{-1}. \quad (3.9)$$

Therefore, the results of Lemma 4.7 of Ref. [7] hold for $G_2(w)$ and those in Section 5 of Ref. [7] hold for $\varphi_2(w)$. As a result, the exact asymptotic formula $h_2(k)$ is given by Theorem 2.1.

4 Additional proofs for Ref. [7]

In this section, we give proofs of some lemmas and propositions appeared in Ref. [7] as well as derivation of some coefficient vectors also appeared in Section 5 of Ref. [7].

4.1 Proof of Lemma 2.2 of Ref. [7]

Lemma 2.2 of Ref. [7]. Under Assumption 2.2, $\bar{\Gamma}$ is bounded.

Proof. For $n \geq 1$, $j \in S_0$ and $(s_1, s_2) \in \mathbb{R}^2$, $C(e^{s_1}, e^{s_2})^n$ satisfies

$$[C(e^{s_1}, e^{s_2})^n]_{jj} = \sum_{\mathbf{k}_{(n)} \in \mathbb{H}^n} \sum_{\mathbf{l}_{(n)} \in \mathbb{H}^n} [A_{k_1, l_1} A_{k_2, l_2} \cdots A_{k_n, l_n}]_{jj} e^{s_1 \sum_{p=1}^n k_p + s_2 \sum_{p=1}^n l_p}, \quad (4.1)$$

where $\mathbf{k}_{(n)} = (k_1, k_2, \dots, k_n)$ and $\mathbf{l}_{(n)} = (l_1, l_2, \dots, l_n)$. Consider the Markov chain $\{\tilde{\mathbf{Y}}_n\} = \{(\tilde{X}_{1,n}, \tilde{X}_{2,n}, \tilde{J}_n)\}$ governed by $\{A_{k,l} : k, l \in \mathbb{H}\}$ (see Subsection 2.1) and assume that $\{\tilde{\mathbf{Y}}_n\}$ starts from the state $(0, 0, j)$. Since $\{\tilde{\mathbf{Y}}_n\}$ is irreducible and aperiodic, there exists $n_0 \geq 1$ such that, for every $j \in S_0$, $\mathbb{P}(\tilde{\mathbf{Y}}_{n_0} = (1, 0, j) | \tilde{\mathbf{Y}}_0 = (0, 0, j)) > 0$, $\mathbb{P}(\tilde{\mathbf{Y}}_{n_0} = (0, 1, j) | \tilde{\mathbf{Y}}_0 = (0, 0, j)) > 0$, $\mathbb{P}(\tilde{\mathbf{Y}}_{n_0} = (-1, 0, j) | \tilde{\mathbf{Y}}_0 = (0, 0, j)) > 0$ and $\mathbb{P}(\tilde{\mathbf{Y}}_{n_0} = (0, -1, j) | \tilde{\mathbf{Y}}_0 = (0, 0, j)) > 0$. This implies that, for

some $\mathbf{k}_{1,(n_0)}^{(j)}$, $\mathbf{l}_{1,(n_0)}^{(j)}$, $\mathbf{k}_{2,(n_0)}^{(j)}$, $\mathbf{l}_{2,(n_0)}^{(j)}$, $\mathbf{k}_{3,(n_0)}^{(j)}$, $\mathbf{l}_{3,(n_0)}^{(j)}$, $\mathbf{k}_{4,(n_0)}^{(j)}$ and $\mathbf{l}_{4,(n_0)}^{(j)}$ in \mathbb{H}^{n_0} , we have

$$\begin{aligned} b_{1,j} &= [A_{k_{1,1}^{(j)}, l_{1,1}^{(j)}} A_{k_{1,2}^{(j)}, l_{1,2}^{(j)}} \cdots A_{k_{1,n_0}^{(j)}, l_{1,n_0}^{(j)}}]_{jj} > 0, & \sum_{p=1}^{n_0} k_{1,p}^{(j)} &= 1, & \sum_{p=1}^{n_0} l_{1,p}^{(j)} &= 0, \\ b_{2,j} &= [A_{k_{2,1}^{(j)}, l_{2,1}^{(j)}} A_{k_{2,2}^{(j)}, l_{2,2}^{(j)}} \cdots A_{k_{2,n_0}^{(j)}, l_{2,n_0}^{(j)}}]_{jj} > 0, & \sum_{p=1}^{n_0} k_{2,p}^{(j)} &= 0, & \sum_{p=1}^{n_0} l_{2,p}^{(j)} &= 1, \\ b_{3,j} &= [A_{k_{3,1}^{(j)}, l_{3,1}^{(j)}} A_{k_{3,2}^{(j)}, l_{3,2}^{(j)}} \cdots A_{k_{3,n_0}^{(j)}, l_{3,n_0}^{(j)}}]_{jj} > 0, & \sum_{p=1}^{n_0} k_{3,p}^{(j)} &= -1, & \sum_{p=1}^{n_0} l_{3,p}^{(j)} &= 0, \\ b_{4,j} &= [A_{k_{4,1}^{(j)}, l_{4,1}^{(j)}} A_{k_{4,2}^{(j)}, l_{4,2}^{(j)}} \cdots A_{k_{4,n_0}^{(j)}, l_{4,n_0}^{(j)}}]_{jj} > 0, & \sum_{p=1}^{n_0} k_{4,p}^{(j)} &= 0, & \sum_{p=1}^{n_0} l_{4,p}^{(j)} &= -1. \end{aligned}$$

Hence, the sum of any row of $C(e^{s_1}, e^{s_2})^{n_0}$ is greater than or equal to $b^*(e^{s_1} + e^{s_2} + e^{-s_1} + e^{-s_2})$, where

$$b^* = \min_{j \in S_0} \min_{1 \leq i \leq 4} b_{i,j} > 0,$$

and we obtain

$$\chi(e^{s_1}, e^{s_2}) = \text{spr}(C(e^{s_1}, e^{s_2})) = \text{spr}(C(e^{s_1}, e^{s_2})^{n_0})^{\frac{1}{n_0}} \geq (b^*(e^{s_1} + e^{s_2} + e^{-s_1} + e^{-s_2}))^{\frac{1}{n_0}}, \quad (4.2)$$

where we use the fact that, for a nonnegative square matrix $A = (a_{i,j})$, $\text{spr}(A) \geq \min_i \sum_j a_{i,j}$ (see, for example, Theorem 8.1.22 of Horn and Johnson [2]). This means that $\chi(e^{s_1}, e^{s_2})$ is unbounded in any direction, and since $\chi(e^{s_1}, e^{s_2})$ is convex in (s_1, s_2) , we see that Γ is a bounded set. \square

4.2 Proof of Lemma 2.5 of Ref. [7]

Lemma 2.5 of Ref. [7]. *If $a_2^{\{1,2\}} < 0$, then for some positive constant c_1 , $\psi'_1(1) = c_1 a_1^{\{1\}}$. If $a_1^{\{1,2\}} < 0$, then for some positive constant c_2 , $\psi'_2(1) = c_2 a_1^{\{2\}}$.*

Proof. We prove only the first claim. The second claim can be proved analogously. Assume $a_2^{\{1,2\}} < 0$. Then, the induced Markov chain $\mathcal{L}^{\{1\}}$ defined in Subsection 2.2 is positive recurrent and $G_1(1)$ is stochastic. For $z_1 \in [z_1^*, \bar{z}_1^*]$, $C_1(z_1, G_1(z_1))$ is a nonnegative matrix and $\psi_1(z_1)$ is its Perron-Frobenius eigenvalue. Let $\mathbf{u}^{C_1}(z_1)$ and $\mathbf{v}^{C_1}(z_1)$ be the Perron-Frobenius left and right eigenvectors of $C_1(z_1, G_1(z_1))$, satisfying $\mathbf{u}^{C_1}(z_1) \mathbf{v}^{C_1}(z_1) = 1$. Since $\mathcal{L}^{\{1\}}$ is positive recurrent, we have $\mathbf{u}^{C_1}(1) = c_1 \boldsymbol{\pi}_{*,0}^{(1)}$ and $\mathbf{v}^{C_1}(1) = \mathbf{1}$, where $c_1 = (\boldsymbol{\pi}_{*,0}^{(1)} \mathbf{1})^{-1}$ is the normalizing constant. The derivative of $\psi_1(z_1)$, $\psi'_1(z_1)$, is given as

$$\psi'_1(z_1) = \mathbf{u}^{C_1}(z_1) \left(\frac{d}{dz} C_1(z_1, G_1(z_1)) \right) \mathbf{v}^{C_1}(z_1), \quad (4.3)$$

$$\frac{d}{dz_1} C_1(z_1, G_1(z_1)) = -z_1^{-2} A_{-1,0}^{(1)} + A_{1,0}^{(1)} + (-z_1^{-2} A_{-1,1}^{(1)} + A_{1,1}^{(1)}) G_1(z_1) + A_{*,1}^{(1)}(z_1) G'_1(z_1). \quad (4.4)$$

Since $G_1(z_1)$ is a solution to equation (2.1) and $G_1(1)$ is stochastic, the derivative of $G_1(z_1)$, $G'_1(z_1)$, satisfies

$$(-A_{-1,*} + A_{1,*}) \mathbf{1} + (A_{*,0} + A_{*,1} G_1(1) + A_{*,1}) G'_1(1) \mathbf{1} = G'_1(1) \mathbf{1}. \quad (4.5)$$

Define a matrix N_1 as

$$N_1 = (I - (A_{*,0} + A_{*,1}G_1(1)))^{-1}. \quad (4.6)$$

This N_1 is well-defined since $G_1(1)$ is the G-matrix of the transition probability matrix $A_*^{(1)}$ (see Proposition 3.5 of Ozawa [6]). Furthermore, N_1 satisfies $G_1(1) = N_1A_{*, -1}$ and $R_*^{(1)} = A_{*,1}N_1$. From equation (4.5), we, therefore, obtain

$$G'_1(1)\mathbf{1} = N_1(-A_{-1,*} + A_{1,*})\mathbf{1} + N_1A_{*,1}G'_1(1)\mathbf{1}, \quad (4.7)$$

and this leads us to

$$G'_1(1)\mathbf{1} = N_1 \sum_{k=0}^{\infty} (R_*^{(1)})^k (-A_{-1,*} + A_{1,*})\mathbf{1}. \quad (4.8)$$

As a result, we obtain

$$\psi'_1(1) = c_1 \boldsymbol{\pi}_{*,0}^{(1)} \left(-A_{-1,0}^{(1)} + A_{1,0}^{(1)} - A_{-1,1}^{(1)} + A_{1,1}^{(1)} \right) \mathbf{1} + c_1 \boldsymbol{\pi}_{*,0}^{(1)} A_{*,1}^{(1)} G'_1(1)\mathbf{1} = c_1 a_1^{\{1\}}, \quad (4.9)$$

where we use the fact that $\boldsymbol{\pi}_{*,1}^{(1)} = \boldsymbol{\pi}_{*,0}^{(1)} A_{*,1}^{(1)} N_1$, and this completes the proof. \square

4.3 Proof of Lemma 3.2 of Ref. [7]

Lemma 3.2 of Ref. [7]. For $n \in \mathbb{Z}_+$, let \mathbf{a}_n be a $1 \times m$ complex vector and define a vector series $\phi(w)$ as $\phi(w) = \sum_{n=0}^{\infty} \mathbf{a}_n w^n$. Furthermore, for an $m \times m$ complex matrix X , define $\phi(X)$ as $\phi(X) = \sum_{n=0}^{\infty} \mathbf{a}_n X^n$. Assume that $\phi(w)$ converges absolutely for all $w \in \mathbb{C}$ such that $|w| < r$ for some $r > 0$. Then, if $\text{spr}(X) < r$, $\phi(X)$ also converges absolutely.

Proof. We denote by $J_k(\lambda)$ the k -dimensional Jordan block of eigenvalue λ . Note that the n -th power of $J_k(\lambda)$ is given by

$$J_k(\lambda)^n = \begin{pmatrix} {}_n C_0 \lambda^n & {}_n C_1 \lambda^{n-1} & \cdots & {}_n C_{k-1} \lambda^{n-k+1} \\ & \ddots & \ddots & \\ & & {}_n C_0 \lambda^n & {}_n C_1 \lambda^{n-1} \\ & & & {}_n C_0 \lambda^n \end{pmatrix}. \quad (4.10)$$

Without loss of generality, the Jordan canonical form J of X is represented as

$$J = T^{-1}XT = \begin{pmatrix} J_{k_1}(\lambda_1) & & & \\ & J_{k_2}(\lambda_2) & & \\ & & \ddots & \\ & & & J_{k_l}(\lambda_l) \end{pmatrix}, \quad (4.11)$$

where T is a nonsingular matrix, $\lambda_1, \lambda_2, \dots, \lambda_l$ are the eigenvalues of X and k_1, k_2, \dots, k_l are positive integers satisfying $k_1 + \dots + k_l = m$. We have

$$\begin{aligned} \left(\sum_{n=0}^{\infty} |\mathbf{a}_n X^n| \right)^\top &\leq \sum_{n=0}^{\infty} |T^{-1}|^\top |J|^n |T|^\top |\mathbf{a}_n|^\top \\ &= \sum_{n=0}^{\infty} \left(\mathbf{1}^\top \otimes |T^{-1}|^\top \right) (\text{diag}(|\mathbf{a}_n|) \otimes |J|^n) \text{vec}(|T|^\top) \\ &= \left(\mathbf{1}^\top \otimes (|T^{-1}|)^\top \right) \text{diag} \left(\sum_{n=0}^{\infty} [|\mathbf{a}_n|]_j |J|^n, j = 1, 2, \dots, s_0 \right) \text{vec}(|T|^\top), \end{aligned} \quad (4.12)$$

where we use the identity $\text{vec}(ABC) = (C^\top \otimes A)\text{vec}(B)$ for matrices A, B and C . Note that

$$\sum_{n=0}^{\infty} [|\mathbf{a}_n|]_j |J|^n = \text{diag} \left(\sum_{n=0}^{\infty} [|\mathbf{a}_n|]_j J_{k_s} (|\lambda_s|)^n, s = 1, 2, \dots, l \right), \quad (4.13)$$

and we have, for $t, u \in \{1, 2, \dots, k_s\}$ such that $t \leq u$,

$$\begin{aligned} &\sum_{n=0}^{\infty} [[|\mathbf{a}_n|]_j J_{k_s} (|\lambda_s|)^n]_{t,u} \\ &= \sum_{n=0}^{k_s-2} [[|\mathbf{a}_n|]_j J_{k_s} (|\lambda_s|)^n]_{t,u} + \sum_{n=k_s-1}^{\infty} \frac{n!}{(u-t)!(n-u+t)!} [|\mathbf{a}_n|]_j |\lambda_s|^{n-u+t} \\ &\leq \sum_{n=0}^{k_s-2} [[|\mathbf{a}_n|]_j J_{k_s} (|\lambda_s|)^n]_{t,u} + \sum_{n=u-t}^{\infty} \frac{n!}{(n-u+t)!} [|\mathbf{a}_n|]_j |\lambda_s|^{n-u+t} \\ &= \sum_{n=0}^{k_s-2} [[|\mathbf{a}_n|]_j J_{k_s} (|\lambda_s|)^n]_{t,u} + \frac{d^{u-t}}{dw^{u-t}} [\phi_{abs}(w)]_j \Big|_{w=|\lambda_s|}, \end{aligned} \quad (4.14)$$

where $\phi_{abs}(w) = \sum_{n=0}^{\infty} |\mathbf{a}_n| w^n$. For $w \in \mathbb{C}$ such that $|w| < r$, since $\phi(w)$ is absolutely convergent, $\phi_{abs}(w)$ is also absolutely convergent and analytic. Hence, $\phi_{abs}(w)$ is differentiable any number of times and we know that the second term on the last line of formula (4.14) is finite since $|\lambda_s| \leq \text{spr}(X) < r$; this completes the proof. \square

4.4 Proof of Lemma 4.1 of Ref. [7]

For $z \in \mathbb{C}$, we define $G_1(z)$ in a manner similar to that used for defining so-called *G-matrices* of QBD process (see Neuts [5]). For $z \in \mathbb{C}$, we also define $R_1(z)$ and $N_1(z)$. For the purpose, we use the following sets of index sequences: for $n \geq 1$ and for $m \geq 1$,

$$\begin{aligned} \mathcal{I}_n &= \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : \sum_{l=1}^k i_l \geq 0 \text{ for } k \in \{1, 2, \dots, n-1\} \text{ and } \sum_{l=1}^n i_l = 0 \right\}, \\ \mathcal{I}_{D,m,n} &= \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : \sum_{l=1}^k i_l \geq -m+1 \text{ for } k \in \{1, 2, \dots, n-1\} \text{ and } \sum_{l=1}^n i_l = -m \right\}, \\ \mathcal{I}_{U,m,n} &= \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : \sum_{l=1}^k i_l \geq 1 \text{ for } k \in \{1, 2, \dots, n-1\} \text{ and } \sum_{l=1}^n i_l = m \right\}, \end{aligned}$$

where $\mathbf{i}_{(n)} = (i_1, i_2, \dots, i_n)$. For $n \geq 1$, define $Q_{11}^{(n)}(z)$, $D_1^{(n)}(z)$ and $U_1^{(n)}(z)$ as

$$\begin{aligned} Q_{11}^{(n)}(z) &= \sum_{\mathbf{i}_{(n)} \in \mathcal{I}_n} A_{*,i_1}(z) A_{*,i_2}(z) \cdots A_{*,i_n}(z), \\ D_1^{(n)}(z) &= \sum_{\mathbf{i}_{(n)} \in \mathcal{I}_{D,1,n}} A_{*,i_1}(z) A_{*,i_2}(z) \cdots A_{*,i_n}(z), \\ U_1^{(n)}(z) &= \sum_{\mathbf{i}_{(n)} \in \mathcal{I}_{U,1,n}} A_{*,i_1}(z) A_{*,i_2}(z) \cdots A_{*,i_n}(z), \end{aligned}$$

and define $N_1(z)$, $R_1(z)$ and $G_1(z)$ as

$$N_1(z) = \sum_{n=0}^{\infty} Q_{11}^{(n)}(z), \quad G_1(z) = \sum_{n=1}^{\infty} D_1^{(n)}(z), \quad R_1(z) = \sum_{n=1}^{\infty} U_1^{(n)}(z),$$

where $Q_{11}^{(0)}(z) = I$. $N_1(z)$, $G_1(z)$ and $R_1(z)$ satisfy the following properties.

Lemma 4.1 of Ref. [7]. *For $z \in \mathbb{C}[\underline{z}_1^*, \bar{z}_1^*]$, the following statements hold.*

(i) $N_1(z)$, $G_1(z)$ and $R_1(z)$ converge absolutely and satisfy

$$|N(z)| \leq N(|z|), \quad |G_1(z)| \leq G_1(|z|), \quad |R_1(z)| \leq R_1(|z|). \quad (4.15)$$

(ii) $G_1(z)$ and $R_1(z)$ are represented in terms of $N_1(z)$, as follows.

$$G_1(z) = N_1(z)A_{*,-1}(z), \quad R_1(z) = A_{*,1}(z)N_1(z). \quad (4.16)$$

(iii) $G_1(z)$ and $R_1(z)$ satisfy the following matrix quadratic equations:

$$A_{*,-1}(z) + A_{*,0}(z)G_1(z) + A_{*,1}(z)G_1(z)^2 = G_1(z), \quad (4.17)$$

$$R_1(z)^2 A_{*,-1}(z) + R_1(z)A_{*,0}(z) + A_{*,1}(z) = R_1(z). \quad (4.18)$$

(iv) Define $H_1(z)$ as $H_1(z) = A_{*,0}(z) + A_{*,1}(z)N_1(z)A_{*,-1}(z)$, then $N_1(z)$ satisfies

$$(I - H_1(z))N_1(z) = I. \quad (4.19)$$

(v) For nonzero $w \in \mathbb{C}$, $I - C(z, w)$ satisfies the following factorization (see, for example, Ref. [4]).

$$I - C(z, w) = (w^{-1}I - R_1(z))(I - H_1(z))(wI - G_1(z)). \quad (4.20)$$

Proof. (i) First, we note that if z is a positive real number in $[\underline{z}_1^*, \bar{z}_1^*]$, then $N_1(z)$, $R_1(z)$ and $G_1(z)$ are finite. Let z be a complex number satisfying $|z| \in [\underline{z}_1^*, \bar{z}_1^*]$. Since $A_{i,j}$, $i, j \in \mathbb{H}$, are nonnegative, we have for $j \in \mathbb{H}$,

$$|A_{*,j}(z)| = \left| \sum_{i \in \mathbb{H}} A_{i,j} z^i \right| \leq \sum_{i \in \mathbb{H}} A_{i,j} |z|^i = A_{*,j}(|z|),$$

and for $n \geq 0$,

$$|Q_{11}^{(n)}(z)| = \left| \sum_{\mathbf{i}_{(n)} \in \mathcal{I}_n} A_{*,i_1}(z) A_{*,i_2}(z) \cdots A_{*,i_n}(z) \right| \leq Q_{11}^{(n)}(|z|).$$

From this and the fact that $N_1(|z|) = \sum_{n=0}^{\infty} Q_{11}^{(n)}(|z|)$ is finite (convergent), we see that $N_1(z) = \sum_{n=0}^{\infty} Q_{11}^{(n)}(z)$ converges absolutely and obtain $|N_1(z)| \leq N_1(|z|)$. Analogously, we see that both $G_1(z)$ and $R_1(z)$ also converge absolutely and satisfy expression (4.15).

(ii) Since, for $n \geq 1$, $\mathcal{I}_{D,1,n}$ and $\mathcal{I}_{U,1,n}$ satisfy

$$\begin{aligned} \mathcal{I}_{D,1,n} &= \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : \sum_{l=1}^k i_l \geq 0 \text{ for } k \in \{1, 2, \dots, n-2\}, \sum_{l=1}^{n-1} i_l = 0 \text{ and } i_n = -1 \right\} \\ &= \{(\mathbf{i}_{(n-1)}, -1) : \mathbf{i}_{(n-1)} \in \mathcal{I}_{n-1}\}, \\ \mathcal{I}_{U,1,n} &= \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : i_1 = 1, \sum_{l=2}^k i_l \geq 0 \text{ for } k \in \{2, \dots, n-1\} \text{ and } \sum_{l=2}^n i_l = 0 \right\} \\ &= \{(1, \mathbf{i}_{(n-1)}) : \mathbf{i}_{(n-1)} \in \mathcal{I}_{n-1}\}, \end{aligned}$$

where $\mathbf{i}_{(n)} = (i_1, i_2, \dots, i_n)$, we have

$$\begin{aligned} G_1(z) &= \sum_{n=1}^{\infty} Q_{11}^{(n-1)}(z) A_{*, -1}(z) = N_1(z) A_{*, -1}(z), \\ R_1(z) &= \sum_{n=1}^{\infty} A_{*, 1}(z) Q_{11}^{(n-1)}(z) = A_{*, 1}(z) N_1(z). \end{aligned}$$

(iii) We prove only equation (4.17) since equation (4.18) can analogously be proved. For $n \geq 3$, $\mathcal{I}_{D,1,n}$ satisfies

$$\begin{aligned} \mathcal{I}_{D,1,n} &= \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : i_1 = 0, \sum_{l=2}^k i_l \geq 0 \text{ for } k \in \{2, \dots, n-1\}, \sum_{l=2}^n i_l = -1 \right\} \\ &\quad \cup \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : i_1 = 1, \sum_{l=2}^k i_l \geq -1 \text{ for } k \in \{2, \dots, n-1\}, \sum_{l=2}^n i_l = -2 \right\} \\ &= \{(0, \mathbf{i}_{(n-1)}) : \mathbf{i}_{(n-1)} \in \mathcal{I}_{D,1,n-1}\} \cup \{(1, \mathbf{i}_{(n-1)}) : \mathbf{i}_{(n-1)} \in \mathcal{I}_{D,2,n-1}\}, \end{aligned}$$

and $\mathcal{I}_{D,2,n}$ satisfies

$$\begin{aligned} \mathcal{I}_{D,2,n} &= \bigcup_{m=1}^{n-1} \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : \sum_{l=1}^k i_l \geq 0 \text{ for } k \in \{1, 2, \dots, m-1\}, \sum_{l=1}^m i_l = -1, \right. \\ &\quad \left. \sum_{l=m+1}^k i_l \geq 0 \text{ for } k \in \{m+1, m+2, \dots, n-1\} \text{ and } \sum_{l=m+1}^n i_l = -1 \right\} \\ &= \bigcup_{m=1}^{n-1} \{(\mathbf{i}_{(m)}, \mathbf{i}_{(n-m)}) : \mathbf{i}_{(m)} \in \mathcal{I}_{D,1,m} \text{ and } \mathbf{i}_{(n-m)} \in \mathcal{I}_{D,1,m-n}\}. \end{aligned}$$

Hence, we have, for $n \geq 3$,

$$\begin{aligned} D_1^{(n)}(z) &= A_{*,0}(z)D_1^{(n-1)}(z) + A_{*,1}(z) \sum_{\mathbf{i}_{(n-1)} \in \mathcal{I}_{D,2,n-1}} A_{*,i_1}(z)A_{*,i_2}(z) \cdots A_{*,i_{n-1}}(z) \\ &= A_{*,0}(z)D_1^{(n-1)}(z) + A_{*,1}(z) \sum_{m=1}^{n-2} D_1^{(m)}(z)D_1^{(n-m-1)}(z), \end{aligned}$$

and obtain

$$\begin{aligned} G_1(z) &= D_1^{(1)}(z) + \sum_{n=2}^{\infty} A_{*,0}(z)D_1^{(n-1)}(z) + A_{*,1}(z) \sum_{n=3}^{\infty} \sum_{m=1}^{n-2} D_1^{(m)}(z)D_1^{(n-m-1)}(z) \\ &= A_{*,-1}(z) + A_{*,0}(z)G_1(z) + A_{*,1}(z)G_1(z)^2, \end{aligned}$$

where we use the fact that $D_1^{(1)}(z) = A_{*,-1}(z)$ and $D_1^{(2)}(z) = A_{*,0}(z)D_1^{(1)}(z) = A_{*,0}(z)A_{*,-1}(z)$.

(iv) For $n \geq 1$, \mathcal{I}_n satisfies

$$\begin{aligned} \mathcal{I}_n &= \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : i_1 = 0, \sum_{l=2}^k i_l \geq 0 \text{ for } k \in \{2, \dots, n-1\}, \sum_{l=2}^n i_l = 0 \right\} \\ &\quad \cup \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : i_1 = 1, \sum_{l=2}^k i_l \geq -1 \text{ for } k \in \{2, \dots, n-1\}, \sum_{l=2}^n i_l = -1 \right\} \\ &= \{(0, \mathbf{i}_{(n-1)}) : \mathbf{i}_{(n-1)} \in \mathcal{I}_{n-1}\} \\ &\quad \cup \left(\bigcup_{m=2}^n \left\{ \mathbf{i}_{(n)} \in \mathbb{H}^n : i_1 = 1, \sum_{l=2}^k i_l \geq 0 \text{ for } k \in \{2, \dots, m-1\}, \sum_{l=2}^m i_l = -1, \right. \right. \\ &\quad \left. \left. \sum_{l=m+1}^k i_l \geq 0 \text{ for } k \in \{m+1, \dots, n-1\}, \sum_{l=m+1}^n i_l = 0 \right\} \right) \\ &= \{(0, \mathbf{i}_{(n-1)}) : \mathbf{i}_{(n-1)} \in \mathcal{I}_{n-1}\} \\ &\quad \cup \left(\bigcup_{m=2}^n \{(1, \mathbf{i}_{(m-1)}, \mathbf{i}_{(n-m)}) : \mathbf{i}_{(m-1)} \in \mathcal{I}_{D,1,m-1}, \mathbf{i}_{(n-m)} \in \mathcal{I}_{n-m}\} \right), \end{aligned}$$

where $\mathcal{I}_0 = \emptyset$. Hence, we have, for $n \geq 1$,

$$Q_{1,1}^{(n)}(z) = A_{*,0}(z)Q_{1,1}^{(n-1)}(z) + \sum_{m=2}^n A_{*,1}(z)D_1^{(m-1)}(z)Q_{1,1}^{(n-m)}(z),$$

and this leads us to

$$N_1(z) = I + A_{*,0}(z)N_1(z) + A_{*,1}(z)G_1(z)N_1(z).$$

From this equation, we immediately obtain equation (4.19).

(v) Substituting $N_1(z)A_{*,-1}(z)$, $A_{*,1}(z)N_1(z)$ and $A_{*,0}(z) + A_{*,1}(z)N_1(z)A_{*,-1}(z)$ for $G_1(z)$, $R_1(z)$ and $H_1(z)$, respectively, in the right hand side of equation (4.20), we obtain the left hand side of the equation via straightforward calculation. \square

4.5 Proof of Proposition 4.1 of Ref. [7]

Proposition 4.1 of Ref. [7]. *Under Assumption 2.2, for $z, w \in \mathbb{C}$ such that $z \neq 0$ and $w \neq 0$, if $|z| \neq z$ or $|w| \neq w$, then $\text{spr}(C(z, w)) < \text{spr}(C(|z|, |w|))$.*

Proof. Set $z = re^{i\theta}$ and $w = r'e^{i\theta'}$, where $r, r' > 0$, $\theta, \theta' \in [0, 2\pi)$ and $i = \sqrt{-1}$. For $n \geq 1$ and $j \in S_0$, $C(z, w)^n$ satisfies

$$\begin{aligned} \left| [C(z, w)^n]_{jj} \right| &= \left| \sum_{\mathbf{k}_{(n)} \in \mathbb{H}^n} \sum_{\mathbf{l}_{(n)} \in \mathbb{H}^n} [A_{k_1, l_1} A_{k_2, l_2} \cdots A_{k_n, l_n}]_{jj} \right. \\ &\quad \left. \cdot r^{\sum_{p=1}^n k_p} (r')^{\sum_{p=1}^n l_p} e^{i(\theta \sum_{p=1}^n k_p + \theta' \sum_{p=1}^n l_p)} \right| \\ &\leq \sum_{\mathbf{k}_{(n)} \in \mathbb{H}^n} \sum_{\mathbf{l}_{(n)} \in \mathbb{H}^n} [A_{k_1, l_1} A_{k_2, l_2} \cdots A_{k_n, l_n}]_{jj} r^{\sum_{p=1}^n k_p} (r')^{\sum_{p=1}^n l_p} \\ &= [C(|z|, |w|)^n]_{jj}, \end{aligned} \tag{4.21}$$

where $\mathbf{k}_{(n)} = (k_1, k_2, \dots, k_n)$ and $\mathbf{l}_{(n)} = (l_1, l_2, \dots, l_n)$. In this formula, equality holds only when, for every $\mathbf{k}_{(n)}, \mathbf{l}_{(n)} \in \mathbb{H}^n$ such that $[A_{k_1, l_1} A_{k_2, l_2} \cdots A_{k_n, l_n}]_{jj} \neq 0$, $e^{i(\theta \sum_{p=1}^n k_p + \theta' \sum_{p=1}^n l_p)}$ takes some common value. Consider the Markov chain $\{\tilde{\mathbf{Y}}_n\} = \{(\tilde{X}_{1,n}, \tilde{X}_{2,n}, \tilde{J}_n)\}$ generated by $\{A_{k,l} : k, l \in \mathbb{H}\}$ (see Subsection 2.1) and assume that $\{\tilde{\mathbf{Y}}_n\}$ starts from the state $(0, 0, j)$. Since $\{\tilde{\mathbf{Y}}_n\}$ is irreducible and aperiodic, there exists $n_0 \geq 1$ such that $\mathbb{P}(\tilde{\mathbf{Y}}_{n_0} = (0, 0, j) \mid \tilde{\mathbf{Y}}_0 = (0, 0, j)) > 0$, $\mathbb{P}(\tilde{\mathbf{Y}}_{n_0} = (1, 0, j) \mid \tilde{\mathbf{Y}}_0 = (0, 0, j)) > 0$ and $\mathbb{P}(\tilde{\mathbf{Y}}_{n_0} = (0, 1, j) \mid \tilde{\mathbf{Y}}_0 = (0, 0, j)) > 0$. This implies that, for some $\mathbf{k}_{(n_0)}, \mathbf{l}_{(n_0)}, \mathbf{k}'_{(n_0)}, \mathbf{l}'_{(n_0)}, \mathbf{k}''_{(n_0)}$ and $\mathbf{l}''_{(n_0)}$ in \mathbb{H}^{n_0} , we have

$$\begin{aligned} [A_{k_1, l_1} A_{k_2, l_2} \cdots A_{k_{n_0}, l_{n_0}}]_{jj} &> 0, & \sum_{p=1}^{n_0} k_p &= 0, & \sum_{p=1}^{n_0} l_p &= 0, \\ [A_{k'_1, l'_1} A_{k'_2, l'_2} \cdots A_{k'_{n_0}, l'_{n_0}}]_{jj} &> 0, & \sum_{p=1}^{n_0} k'_p &= 1, & \sum_{p=1}^{n_0} l'_p &= 0, \\ [A_{k''_1, l''_1} A_{k''_2, l''_2} \cdots A_{k''_{n_0}, l''_{n_0}}]_{jj} &> 0, & \sum_{p=1}^{n_0} k''_p &= 0, & \sum_{p=1}^{n_0} l''_p &= 1. \end{aligned}$$

Hence, we see that if equality holds in formula (4.21), then $e^{i\theta} = e^{i\theta'} = 1$ and both θ and θ' must be zero. Therefore, if $\theta \neq 0$ or $\theta' \neq 0$, then we have

$$\left| [C(z, w)^{n_0}]_{jj} \right| = \left| [C(re^{i\theta}, r'e^{i\theta'})^{n_0}]_{jj} \right| < [C(r, r')^{n_0}]_{jj} = [C(|z|, |w|)^{n_0}]_{jj},$$

and obtain

$$\text{spr}(C(z, w))^{n_0} \leq \text{spr}(|C(z, w)^{n_0}|) < \text{spr}(C(|z|, |w|)^{n_0}) = \text{spr}(C(|z|, |w|))^{n_0}, \tag{4.22}$$

where we use Theorem 1.5 of Seneta [8] and the fact that $C(|z|, |w|)$ is irreducible. Obviously, this implies that $\text{spr}(C(z, w)) < \text{spr}(C(|z|, |w|))$. \square

4.6 Proof of Propositions 5.1 and 5.2 of Ref. [7]

For $w \in \mathbb{C}$, $\varphi_2(w)$ is defined as $\varphi_2(w) = \sum_{j=1}^{\infty} \nu_{0,j} w^j$. By Proposition 3.1 of Ref. [7], the radius of convergence of $\varphi_2(w)$ is given by r_2 . For a scalar z and matrix X , define vector functions $\varphi_2(X)$ and $\varphi_2^{\hat{C}_2}(z, X)$ as

$$\varphi_2(X) = \sum_{j=1}^{\infty} \nu_{0,j} X^j, \quad \varphi_2^{\hat{C}_2}(z, X) = \sum_{j=1}^{\infty} \nu_{0,j} \hat{C}_2(z, X) X^{j-1},$$

where

$$\hat{C}_2(z, X) = \sum_{j \in \mathbb{H}} A_{*,j}^{(2)}(z) X^{j+1}, \quad A_{*,j}^{(2)}(z) = \sum_{i \in \mathbb{H}_+} A_{i,j}^{(2)} z^i.$$

Proposition 5.1 of Ref. [7]. *If $G_1(z)$ is entry-wise analytic and satisfies $\text{spr}(G_1(z)) < r_2$ in a region $\Omega \subset \mathbb{C}$, then $\varphi_2(G_1(z))$ and $\varphi_2^{\hat{C}_2}(z, G_1(z))$ are also entry-wise analytic in the same region Ω .*

Proof. Let $X = (x_{k,l})$ be an $s_0 \times s_0$ complex matrix. By Lemma 3.2 of Ref. [7], if $\text{spr}(X) < r_2$, then $\varphi_2(X)$ converges absolutely. This means that each element of $\varphi_2(X)$ is an absolutely convergent series with s_0^2 valuables $x_{11}, x_{12}, \dots, x_{s_0, s_0}$ and it is analytic in the region $\{X = (x_{k,l}) \in \mathbb{C}^{s_0^2} : \text{spr}(X) < r_2\}$ as a function of s_0^2 complex valuables. Therefore, if $G_1(z)$ is entry-wise analytic in Ω and $\text{spr}(G_1(z)) < r_2$, we see that $\varphi_2(G_1(z)) = (\varphi_2 \circ G_1)(z)$ is also entry-wise analytic in Ω . For $z \in \mathbb{C}$ such that $|z| < r_2$, we have

$$\sum_{j=1}^{\infty} \left| \nu_{0,j} \hat{C}_2(z, G_1(z)) z^{j-1} \right| \leq |z|^{-1} \sum_{j=1}^{\infty} \nu_{0,j} |z|^j \hat{C}_2(|z|, G_1(|z|)) < \infty.$$

Hence, by Lemma 3.2 of Ref. [7], if $\text{spr}(X) < r_2$, $\sum_{j=1}^{\infty} \nu_{0,j} \hat{C}_2(z, G_1(z)) X^{j-1}$ converges absolutely, and we see that if $G_1(z)$ is entry-wise analytic in Ω and $\text{spr}(G_1(z)) < r_2$, then $\varphi_2^{\hat{C}_2}(z, G_1(z))$ is entry-wise analytic in Ω . \square

Proposition 5.2 of Ref. [7]. *Under Assumption 2.2, for any $z \in \mathbb{C}(\underline{z}_1^*, \bar{z}_1^*)$ such that $z \neq |z|$, we have*

$$\text{spr}(C_1(z, G_1(z))) < \text{spr}(C_1(|z|, G_1(|z|))). \quad (4.23)$$

Proof. Set $z = re^{i\theta}$, where $r > 0$, $\theta \in [0, 2\pi)$ and $i = \sqrt{-1}$. By the definition of $G_1(z)$, we have

$$C_1(z, G_1(z)) = \sum_{n=1}^{\infty} \sum_{\mathbf{k}_{(n+1)} \in \mathbb{H}^{n+1}} \sum_{\mathbf{l}_{(n)} \in \mathcal{D}_{D,1,n}} A_{k_1,1}^{(1)} A_{k_2,l_1} A_{k_3,l_2} \cdots A_{k_{n+1},l_n} \cdot r^{\sum_{p=1}^{n+1} k_p} e^{i(\theta \sum_{p=1}^{n+1} k_p)} + \sum_{k \in \mathbb{H}} A_{k,0}^{(1)} r^k e^{i\theta k}, \quad (4.24)$$

where $\mathbf{k}_{(n+1)} = (k_1, k_2, \dots, k_{n+1})$ and $\mathbf{l}_{(n)} = (l_1, l_2, \dots, l_n)$. Consider the Markov chain $\{\tilde{\mathbf{Y}}_n^{(1)}\} = \{(\tilde{X}_{1,n}^{(1)}, \tilde{X}_{2,n}^{(1)}, \tilde{J}_n^{(1)})\}$ generated by $\{\{A_{k,l} : k, l \in \mathbb{H}\}, \{A_{k,l}^{(1)} : k \in \mathbb{H}, l \in \mathbb{H}_+\}\}$ (see Subsection 2.1), and assume that $\{\tilde{\mathbf{Y}}_n^{(1)}\}$ starts from a state in $\{0\} \times \{0\} \times S_0$. For $n_0 \in \mathbb{N}$, let τ_{n_0} be the time when the Markov chain enters a state in $\mathbb{Z} \times \{0\} \times S_0$ for the n_0 -th time. Then, the term $\sum_{p=1}^{n+1} k_p$ (resp. k) in expression (4.24) indicates that $\tilde{X}_{1,\tau_1}^{(1)} = \sum_{p=1}^{n+1} k_p$ (resp. $\tilde{X}_{1,\tau_1}^{(1)} = k$). This point analogously holds for $\tilde{X}_{1,\tau_{n_0}}^{(1)}$ when $n_0 > 1$. Hence, $C_1(z, G_1(z))^{n_0}$ can be represented as

$$C_1(z, G_1(z))^{n_0} = \sum_{k \in \mathbb{Z}} \tilde{D}_k r^k e^{i\theta k}, \quad (4.25)$$

where k indicates that $\tilde{X}_{1,\tau_{n_0}}^{(1)} = k$ and the (j, j') -entry of nonnegative square matrix \tilde{D}_k is given as $[\tilde{D}_k]_{jj'} = \mathbb{P}(\tilde{\mathbf{Y}}_{\tau_{n_0}}^{(1)} = (k, 0, j') \mid \tilde{\mathbf{Y}}_0^{(1)} = (0, 0, j))$. We have

$$|C_1(z, G_1(z))^{n_0}| = \left| \sum_{k \in \mathbb{Z}} \tilde{D}_k r^k e^{i\theta k} \right| \leq C_1(|z|, G_1(|z|))^{n_0}, \quad (4.26)$$

and equality holds only when, for each $j, j' \in S_0$ and for every $k \in \mathbb{Z}$ such that $[\tilde{D}_k]_{jj'} > 0$, $e^{i\theta k}$ takes some common value. Under Assumption 2.2, $\{\tilde{\mathbf{Y}}_n^{(1)}\}$ is irreducible and aperiodic, and for any $j, j' \in S_0$, there exists $n_0 \geq 1$ such that $\mathbb{P}(\tilde{\mathbf{Y}}_{\tau_{n_0}}^{(1)} = (0, 0, j') \mid \tilde{\mathbf{Y}}_0^{(1)} = (0, 0, j)) > 0$ and $\mathbb{P}(\tilde{\mathbf{Y}}_{\tau_{n_0}}^{(1)} = (1, 0, j') \mid \tilde{\mathbf{Y}}_0^{(1)} = (0, 0, j)) > 0$. This implies that $[\tilde{D}_0]_{jj'} > 0$ and $[\tilde{D}_1]_{jj'} > 0$. Hence, if $\theta \neq 0$ ($z \neq |z|$), then we have

$$|[C_1(z, G_1(z))^{n_0}]_{jj'}| < [C_1(|z|, G_1(|z|))^{n_0}]_{jj'},$$

and obtain

$$\text{spr}(C_1(z, G_1(z))^{n_0}) \leq \text{spr}(|C_1(z, G_1(z))^{n_0}|) < \text{spr}(C_1(|z|, G_1(|z|))^{n_0}), \quad (4.27)$$

where we use Theorem 1.5 of Seneta [8] and the fact that $C_1(|z|, G_1(|z|))$ is irreducible. Obviously, this implies that $\text{spr}(C_1(z, G_1(z))) < \text{spr}(C_1(|z|, G_1(|z|)))$ when $z \neq |z|$. \square

4.7 Proof of Propositions 5.6 and 5.7 of Ref. [7]

Let $\alpha_j(z)$, $j = 1, 2, \dots, s_0$, be the eigenvalues of $G_1(z)$, counting multiplicity, and without loss of generality, let $\alpha_{s_0}(z)$ be the eigenvalue corresponding to $\zeta_2(z)$ when $z \in [\underline{z}_1^*, \bar{z}_1^*]$. For $j \in \{1, 2, \dots, s_0\}$, let $\mathbf{v}_j(z)$ be the right eigenvector of $G_1(z)$ with respect to the eigenvalue $\alpha_j(z)$. Define matrices $J_1(z)$ and $V_1(z)$ as $J_1(z) = \text{diag}(\alpha_1(z), \alpha_2(z), \dots, \alpha_{s_0}(z))$ and $V_1(z) = (\mathbf{v}_1(z) \ \mathbf{v}_2(z) \ \cdots \ \mathbf{v}_{s_0}(z))$, respectively. Under Assumption 2.5, $V_1(z)$ is nonsingular in a neighborhood of $z = r_1$, and $G_1(z)$ is factorized as $G_1(z) = V_1(z)J_1(z)V_1(z)^{-1}$ (Jordan decomposition). We represent $V_1(z)^{-1}$ as

$$V_1(z)^{-1} = \begin{pmatrix} \mathbf{u}_1(z) \\ \mathbf{u}_2(z) \\ \vdots \\ \mathbf{u}_{s_0}(z) \end{pmatrix},$$

where each $\mathbf{u}_j(z)$ is a $1 \times s_0$ vector. For $j \in \{1, 2, \dots, s_0\}$, $\mathbf{u}_j(z)$ is the left eigenvector of $G_1(z)$ with respect to the eigenvalue $\alpha_j(z)$ and satisfies $\mathbf{u}_j(z)\mathbf{v}_k(z) = \delta_{jk}$ for $k \in \{1, 2, \dots, s_0\}$, where δ_{jk} is the Kronecker delta. Under Assumption 2.5, if $r_1 < \bar{z}_1^*$, $J_1(z)$ and $V_1(z)$ are entry-wise analytic in a neighborhood of $z = r_1$; if $r_1 = \bar{z}_1^*$, $J_1(z)$ and $V_1(z)$ except for $\alpha_{s_0}(z)$ and $\mathbf{v}_{s_0}(z)$ are also entry-wise analytic in a neighborhood of $z = \bar{z}_1^*$, where $\alpha_{s_0}(z)$ and $\mathbf{v}_{s_0}(z)$ are given in terms of a function $\tilde{\alpha}_{s_0}(\zeta)$ and vector function $\tilde{\mathbf{v}}_{s_0}(\zeta)$ analytic in a neighborhood of the origin as $\alpha_{s_0}(z) = \tilde{\alpha}_{s_0}((\bar{z}_1^* - z)^{\frac{1}{2}})$ and $\mathbf{v}_{s_0}(z) = \tilde{\mathbf{v}}_{s_0}((\bar{z}_1^* - z)^{\frac{1}{2}})$ (see Section 4 of Ref. [7]).

Define $\tilde{J}_1(\zeta)$ and $\tilde{V}_1(\zeta)$ as

$$\begin{aligned}\tilde{J}_1(\zeta) &= \text{diag}(\alpha_1(\bar{z}_1^* - \zeta^2), \dots, \alpha_{s_0-1}(\bar{z}_1^* - \zeta^2), \tilde{\alpha}_{s_0}(\zeta)), \\ \tilde{V}_1(\zeta) &= (\mathbf{v}_1(\bar{z}_1^* - \zeta^2) \quad \cdots \quad \mathbf{v}_{s_0-1}(\bar{z}_1^* - \zeta^2) \quad \tilde{\mathbf{v}}_{s_0}(\zeta)).\end{aligned}$$

Under Assumption 2.5, $\tilde{J}_1(\zeta)$ and $\tilde{V}_1(\zeta)$ are entry-wise analytic in a neighborhood of the origin and $\tilde{V}_1(\zeta)$ is nonsingular. Therefore, $\tilde{G}_1(\zeta)$ defined as $\tilde{G}_1(\zeta) = \tilde{V}_1(\zeta)\tilde{J}_1(\zeta)\tilde{V}_1(\zeta)^{-1}$ is entry-wise analytic in a neighborhood of the origin and satisfies

$$G_1(z) = \tilde{G}_1((\bar{z}_1^* - z)^{\frac{1}{2}}). \quad (4.28)$$

Since $\tilde{\alpha}_{s_0}(\zeta)$ and $\tilde{G}_1(\zeta)$ are analytic in a neighborhood of the origin, their Taylor series are represented as

$$\tilde{\alpha}_{s_0}(\zeta) = \sum_{k=0}^{\infty} \tilde{\alpha}_{s_0,k} \zeta^k, \quad \tilde{G}_1(\zeta) = \sum_{k=0}^{\infty} \tilde{G}_{1,k} \zeta^k, \quad (4.29)$$

where $\tilde{\alpha}_{s_0,0} = \zeta_2(\bar{z}_1^*)$ and $\tilde{G}_{1,0} = G_1(\bar{z}_1^*)$. The Puiseux series for $G_1(z)$ at $z = \bar{z}_1^*$ is represented as

$$G_1(z) = \sum_{k=0}^{\infty} \tilde{G}_{1,k} (\bar{z}_1^* - z)^{\frac{k}{2}}.$$

In the following proposition and its proof, $\tilde{\alpha}_{s_0,1}$, $\mathbf{u}_{s_0}(z)$ and $\mathbf{v}_{s_0}(z)$ are denoted by $\tilde{\alpha}_{s_0,1}^{G_1}$, $\mathbf{u}_{s_0}^{G_1}(z)$ and $\mathbf{v}_{s_0}^{G_1}(z)$, respectively, in order to explicitly indicate that they are a coefficient and vector with respect to $G_1(z)$. This notation is also used in Proposition 5.7 of Ref. [7] and Proposition 4.1.

Proposition 5.6 of Ref. [7].

$$\lim_{\substack{\Delta_{\bar{z}_1^*} \ni z \rightarrow \bar{z}_1^* \\ \bar{z}_1^* - z \rightarrow \bar{z}_1^*}} \frac{G_1(\bar{z}_1^*) - G_1(z)}{(\bar{z}_1^* - z)^{\frac{1}{2}}} = -\tilde{G}_{1,1} = -\tilde{\alpha}_{s_0,1}^{G_1} N_1(\bar{z}_1^*) \mathbf{v}^{R_1}(\bar{z}_1^*) \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*) \geq \mathbf{0}^\top, \neq \mathbf{0}^\top, \quad (4.30)$$

where $\mathbf{v}^{R_1}(\bar{z}_1^*)$ is the right eigenvector of $R_1(\bar{z}_1^*)$ with respect to the eigenvalue $\bar{\zeta}_2(\bar{z}_1^*)^{-1}$, satisfying $\mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*) N_1(\bar{z}_1^*) \mathbf{v}^{R_1}(\bar{z}_1^*) = 1$.

Proof. $\tilde{G}_{1,1}$ is given by $\tilde{G}_{1,1} = (d/d\zeta) \tilde{G}_1(\zeta) |_{\zeta=0}$. Differentiating both sides of $\tilde{G}_1(\zeta) = \tilde{V}_1(\zeta)\tilde{J}_1(\zeta)\tilde{V}_1(\zeta)^{-1}$ and setting $\zeta = 0$, we obtain

$$\tilde{G}_{1,1} = \tilde{\alpha}_{s_0,1}^{G_1} \mathbf{v}^\dagger \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*), \quad (4.31)$$

where $\mathbf{v}^\dagger = \mathbf{v}_{s_0}^{G_1}(\bar{z}_1^*) + (\tilde{\alpha}_{s_0,1}^{G_1})^{-1}(\zeta_2(\bar{z}_1^*)I - G_1(\bar{z}_1^*))\tilde{\mathbf{v}}_{s_0,1}$ and $\tilde{\mathbf{v}}_{s_0,1} = (d/d\zeta)\tilde{\mathbf{v}}_{s_0}(\zeta)|_{\zeta=0}$. In the derivation of equation (4.31), we use the following identity:

$$\frac{d}{d\zeta}\tilde{V}_1(\zeta)^{-1} = -\tilde{V}_1(\zeta)^{-1}\left(\frac{d}{d\zeta}\tilde{V}_1(\zeta)\right)\tilde{V}_1(\zeta)^{-1}.$$

Note that $\mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*)\mathbf{v}^\dagger = 1$ and $\tilde{\alpha}_{s_0,1}^{G_1}$ is an eigenvalue of $\tilde{G}_{1,1}$. By equation (4.17), $\tilde{G}_1(\zeta)$ satisfies

$$\tilde{G}_1(\zeta) = A_{*, -1}(\bar{z}_1^* - \zeta^2) + A_{*, 0}(\bar{z}_1^* - \zeta^2)\tilde{G}_1(\zeta) + A_{*, 1}(\bar{z}_1^* - \zeta^2)\tilde{G}_1(\zeta)^2. \quad (4.32)$$

Differentiating both sides of equation (4.32) and setting $\zeta = 0$, we obtain

$$\tilde{G}_{1,1} = A^\dagger \tilde{G}_{1,1}, \quad (4.33)$$

where $A^\dagger = A_{*, 0}(\bar{z}_1^*) + \zeta_2(\bar{z}_1^*)A_{*, 1}(\bar{z}_1^*) + A_{*, 1}(\bar{z}_1^*)G_1(\bar{z}_1^*)$. In the derivation of equation (4.33), we use the fact that $\tilde{G}_{1,1}G_1(\bar{z}_1^*) = \zeta_2(\bar{z}_1^*)\tilde{G}_{1,1}$. Multiplying both sides of equation (4.33) by $\mathbf{v}_{s_0}^{G_1}(\bar{z}_1^*)$ from the right, we obtain $A^\dagger\mathbf{v}^\dagger = \mathbf{v}^\dagger$. Hence, \mathbf{v}^\dagger is the right eigenvector of A^\dagger with respect to the eigenvalue 1. From equation (4.20), we obtain

$$I - A^\dagger = (\bar{\zeta}_2(\bar{z}_1^*)^{-1}I - R_1(\bar{z}_1^*))(I - H_1(\bar{z}_1^*)), \quad (4.34)$$

and from equation (4.19), we know that $N_1(\bar{z}_1^*) = (I - H_1(\bar{z}_1^*))^{-1}$. Hence, $N_1(\bar{z}_1^*)\mathbf{v}^{R_1}(\bar{z}_1^*)$ is the right eigenvector of A^\dagger with respect to the eigenvalue 1, and we see that \mathbf{v}^\dagger can be given by $\mathbf{v}^\dagger = N_1(\bar{z}_1^*)\mathbf{v}^{R_1}(\bar{z}_1^*)$. Since $\mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*)$ is the left eigenvector of $\tilde{G}_{1,1}$ with respect to the eigenvalue $\tilde{\alpha}_{s_0,1}^{G_1}$, $\mathbf{v}^{R_1}(\bar{z}_1^*)$ must satisfy $\mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*)N_1(\bar{z}_1^*)\mathbf{v}^{R_1}(\bar{z}_1^*) = 1$. Since $\tilde{\alpha}_{s_0,1}^{G_1}$ is negative, nonnegativity of $-\tilde{G}_{1,1}$ is obvious. Since $N_1(\bar{z}_1^*) \geq I$, we have $N_1(\bar{z}_1^*)\mathbf{v}^{R_1}(\bar{z}_1^*) \geq \mathbf{v}^{R_1}(\bar{z}_1^*) \neq \mathbf{0}$ and $-\tilde{G}_{1,1}$ is nonzero. \square

From the results in Section 5 of Ref. [7], we know that $\varphi_2(G_1(z))$ is represented as

$$\varphi_2(G_1(z)) = (\varphi_2(\alpha_1(z))\mathbf{v}_1(z) \quad \varphi_2(\alpha_2(z))\mathbf{v}_2(z) \quad \cdots \quad \varphi_2(\alpha_{s_0}(z))\mathbf{v}_{s_0}(z))V(z)^{-1}. \quad (4.35)$$

Assume $r_1 < \bar{z}_1^*$ and $\text{spr}(G_1(r_1)) = r_2 < \bar{z}_2^*$. Then, the Laurent series for $\varphi_2(G_1(z))$ at $z = r_1$ is represented as

$$\varphi_2(G_1(z)) = \sum_{k=-1}^{\infty} \varphi_{2,k}^{G_1}(r_1 - z)^k. \quad (4.36)$$

Assume $r_1 = \bar{z}_1^*$ and define $\tilde{\varphi}_2(\tilde{G}_1(\zeta))$ as

$$\begin{aligned} \tilde{\varphi}_2(\tilde{G}_1(\zeta)) &= \left(\varphi_2(\alpha_1(\bar{z}_1^* - \zeta^2))\mathbf{v}_1(\bar{z}_1^* - \zeta^2) \quad \cdots \right. \\ &\quad \left. \varphi_2(\alpha_{s_0-1}(\bar{z}_1^* - \zeta^2))\mathbf{v}_{s_0-1}(\bar{z}_1^* - \zeta^2) \quad \varphi_2(\tilde{\alpha}_{s_0}(\zeta))\tilde{\mathbf{v}}_{s_0}(\zeta) \right) \tilde{V}(\zeta)^{-1}. \end{aligned} \quad (4.37)$$

If $\text{spr}(\tilde{G}_1(0)) = \text{spr}(G_1(\bar{z}_1^*)) < r_2$, then the Puiseux series for $\varphi_2(G_1(z))$ at $z = \bar{z}_1^*$ is represented as

$$\varphi_2(G_1(z)) = \tilde{\varphi}_2(\tilde{G}_1((\bar{z}_1^* - z)^{\frac{1}{2}})) = \sum_{k=0}^{\infty} \tilde{\varphi}_{2,k}^{\tilde{G}_1}(\bar{z}_1^* - z)^{\frac{k}{2}}, \quad (4.38)$$

where $\tilde{\varphi}_{2,0}^{\tilde{G}_1} = \varphi_2(G_1(\tilde{z}_1^*))$. If $\text{spr}(\tilde{G}_1(0)) = \text{spr}(G_1(\tilde{z}_1^*)) = r_2$, then the Puiseux series for $\varphi_2(G_1(z))$ at $z = \tilde{z}_1^*$ is represented as

$$\varphi_2(G_1(z)) = \tilde{\varphi}_2(\tilde{G}_1((\tilde{z}_1^* - z)^{\frac{1}{2}})) = \sum_{k=-1}^{\infty} \tilde{\varphi}_{2,k}^{\tilde{G}_1} (\tilde{z}_1^* - z)^{\frac{k}{2}}. \quad (4.39)$$

In the following proposition, $\mathbf{u}^{C_2}(w)$ is the left eigenvector of $C_2(G_2(w), w)$ with respect to the eigenvalue $\psi_2(w)$. For the definition of $c_{pole}^{\varphi_2}$, see Corollary 5.1 of Ref. [7].

Proposition 5.7 of Ref. [7]. *In the case of Type I, if $r_1 = \tilde{z}_1^*$, then $\text{spr}(G_1(\tilde{z}_1^*)) < r_2$ and $\tilde{\varphi}_{2,1}^{\tilde{G}_1}$ in expression (4.38) is given by*

$$\tilde{\varphi}_{2,1}^{\tilde{G}_1} = \sum_{k=1}^{\infty} \nu_{0,k} \sum_{l=1}^k \zeta_2(\tilde{z}_1^*)^{k-l} G_1(\tilde{z}_1^*)^{l-1} \tilde{G}_{1,1} \leq \mathbf{0}^\top, \neq \mathbf{0}^\top. \quad (4.40)$$

In the case of Type II, if $r_1 < \tilde{z}_1^$, then $\text{spr}(G_1(r_1)) = r_2 < \tilde{z}_2^*$ and $\varphi_{2,-1}^{G_1}$ in expression (4.36) is given by*

$$\varphi_{2,-1}^{G_1} = \zeta_{2,z}(r_1)^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(r_1) \mathbf{u}_{s_0}^{G_1}(r_1) \geq \mathbf{0}^\top, \neq \mathbf{0}^\top, \quad (4.41)$$

where $\zeta_{2,z}(z) = (d/dz)\zeta_2(z)$; *if $r_1 = \tilde{z}_1^*$, then $\text{spr}(G_1(\tilde{z}_1^*)) = r_2 < \tilde{z}_2^*$ and $\tilde{\varphi}_{2,-1}^{\tilde{G}_1}$ in expression (4.39) is given by*

$$\tilde{\varphi}_{2,-1}^{\tilde{G}_1} = (-\tilde{\alpha}_{s_0,1}^{G_1})^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(\tilde{z}_1^*) \mathbf{u}_{s_0}^{G_1}(\tilde{z}_1^*) \geq \mathbf{0}^\top, \neq \mathbf{0}^\top. \quad (4.42)$$

Before proving Proposition 5.7 of Ref. [7], we give the following proposition.

Proposition 4.1. *In the case of Type II, if $r_1 < \tilde{z}_1^*$, then $\zeta_2(r_1) = r_2 < \tilde{z}_2^*$ and*

$$\lim_{z \rightarrow r_1} (r_1 - z) \varphi_2(\alpha_{s_0}(z)) = \zeta_{2,z}(r_1)^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) > \mathbf{0}^\top; \quad (4.43)$$

if $r_1 = \tilde{z}_1^$, then $\zeta_2(\tilde{z}_1^*) = r_2 < \tilde{z}_2^*$ and*

$$\lim_{\tilde{\Delta}_{\tilde{z}_1^*} \ni z \rightarrow \tilde{z}_1^*} (\tilde{z}_1^* - z)^{\frac{1}{2}} \varphi_2(\alpha_{s_0}(z)) = (-\tilde{\alpha}_{s_0,1}^{G_1})^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) > \mathbf{0}^\top. \quad (4.44)$$

Proof. In the case of Type II, we have $\alpha_{s_0}(r_1) = \zeta_2(e^{\tilde{\eta}_1^{(c)}}) = e^{\eta_2^{(c)}} = r_2 < \tilde{z}_2^*$, and the point $w = r_2$ is a pole of $\varphi_2(w)$ with order one. If $r_1 < \tilde{z}_1^*$, $\alpha_{s_0}(z)$ is analytic at $z = r_1$ and, by Corollary 5.1 of Ref. [7], we have

$$\begin{aligned} \lim_{z \rightarrow r_1} (r_1 - z) \varphi_2(\alpha_{s_0}(z)) &= \lim_{z \rightarrow r_1} (r_2 - \alpha_{s_0}(z)) \varphi_2(\alpha_{s_0}(z)) \frac{r_1 - z}{r_2 - \alpha_{s_0}(z)} \\ &= c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) / \zeta_{2,z}(r_1), \end{aligned} \quad (4.45)$$

and if $r_1 = \bar{z}_1^*$,

$$\begin{aligned} \lim_{\tilde{\Delta}_{\bar{z}_1^*} \ni z \rightarrow \bar{z}_1^*} (\bar{z}_1^* - z)^{\frac{1}{2}} \varphi_2(\alpha_{s_0}(z)) &= \lim_{\tilde{\Delta}_{\bar{z}_1^*} \ni z \rightarrow \bar{z}_1^*} (r_2 - \alpha_{s_0}(z)) \varphi_2(\alpha_{s_0}(z)) \left(-\frac{(\bar{z}_1^* - z)^{\frac{1}{2}}}{\alpha_{s_0}(z) - r_2} \right) \\ &= c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) / (-\tilde{\alpha}_{s_0,1}^{G_1}), \end{aligned} \quad (4.46)$$

where we use Proposition 5.5 of Ref. [7]. \square

Proof of Proposition 5.7 of Ref. [7]. In the case of Type I, we always have $\text{spr}(G_1(r_1)) = \zeta_2(r_1) < r_2$. Assume $r_1 = \bar{z}_1^*$. By the definition of $\tilde{\varphi}_2(\tilde{G}_1(\zeta))$, we have $\tilde{\varphi}_2(\tilde{G}_1(\zeta)) = \varphi_2(\tilde{G}_1(\zeta))$. Therefore, by Proposition 5.1 of Ref. [7], since $\tilde{G}_1(0) = \text{spr}(G_1(\bar{z}_1^*)) < r_2$, $\tilde{\varphi}_2(\tilde{G}_1(\zeta))$ is given in a form of absolutely convergent series as

$$\tilde{\varphi}_2(\tilde{G}_1(\zeta)) = \sum_{k=0}^{\infty} \nu_{0,k} \tilde{G}_1(\zeta)^k. \quad (4.47)$$

This $\tilde{\varphi}_2(\tilde{G}_1(\zeta))$ is entry-wise analytic at $\zeta = 0$ and we have

$$\begin{aligned} \tilde{\varphi}_{2,1}^{\tilde{G}_1} &= \frac{d}{d\zeta} \tilde{\varphi}_2(\tilde{G}_1(\zeta)) \Big|_{\zeta=0} = \sum_{k=1}^{\infty} \nu_{0,k} \sum_{l=1}^k G_1(\bar{z}_1^*)^{l-1} \tilde{G}_{1,1} G_1(\bar{z}_1^*)^{k-l} \\ &= \sum_{k=1}^{\infty} \nu_{0,k} \sum_{l=1}^k \zeta_2(\bar{z}_1^*)^{k-l} G_1(\bar{z}_1^*)^{l-1} \tilde{G}_{1,1}, \end{aligned} \quad (4.48)$$

where we use the fact that

$$\tilde{G}_{1,1} G_1(\bar{z}_1^*) = \tilde{\alpha}_{s_0,1}^{G_1} N_1(\bar{z}_1^*) \mathbf{v}^{R_1}(\bar{z}_1^*) \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*) G_1(\bar{z}_1^*) = \zeta_2(\bar{z}_1^*) \tilde{G}_{1,1}.$$

Since $\tilde{G}_{1,1}$ is nonzero and nonpositive, $\tilde{\varphi}_{2,1}^{\tilde{G}_1}$ is also nonzero and nonpositive. In the case of Type II, we have $\text{spr}(G_1(r_1)) = \zeta_2(r_1) = r_2 < \bar{z}_2^*$ and $\varphi_2(w)$ has a pole at $w = \zeta_2(r_1)$. Hence, if $r_1 < \bar{z}_1^*$, we obtain from equation (4.35) and Proposition 4.1 that

$$\begin{aligned} \varphi_{2,-1}^{G_1} &= \lim_{z \rightarrow r_1} (r_1 - z) \varphi_2(G_1(z)) \\ &= \left(\mathbf{0} \quad \cdots \quad \mathbf{0} \quad \zeta_{2,z}(r_1)^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(r_1) \right) V_1(r_1)^{-1} \\ &= \zeta_{2,z}(r_1)^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(r_1) \mathbf{u}_{s_0}^{G_1}(r_1), \end{aligned} \quad (4.49)$$

where $\mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(r_1) > 0$; if $r_1 = \bar{z}_1^*$, we also obtain from equation (4.37) and Proposition 4.1 that

$$\begin{aligned} \tilde{\varphi}_{2,-1}^{\tilde{G}_1} &= \lim_{\tilde{\Delta}_{\bar{z}_1^*} \ni z \rightarrow \bar{z}_1^*} (\bar{z}_1^* - z)^{\frac{1}{2}} \varphi_2(G_1(z)) \\ &= \left(\mathbf{0} \quad \cdots \quad \mathbf{0} \quad (-\tilde{\alpha}_{s_0}^{G_1})^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(\bar{z}_1^*) \right) V_1(\bar{z}_1^*)^{-1} \\ &= (-\tilde{\alpha}_{s_0}^{G_1})^{-1} c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(\bar{z}_1^*) \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*), \end{aligned} \quad (4.50)$$

where $\mathbf{u}^{C_2}(r_2) \mathbf{v}_{s_0}^{G_1}(\bar{z}_1^*) > 0$. \square

(2) If $\psi_1(\bar{z}_1^*) < 1$, then the point $z = \bar{z}_1^*$ is a branch point of $\varphi_1(z)$ with order one and its Puiseux series is represented as

$$\varphi_1(z) = \tilde{\varphi}_1((\bar{z}_1^* - z)^{\frac{1}{2}}) = \sum_{k=0}^{\infty} \tilde{\varphi}_{1,k}^I (\bar{z}_1^* - z)^{\frac{k}{2}}, \quad (4.54)$$

where $\tilde{\varphi}_{1,0}^I = \varphi_1(\bar{z}_1^*)$ and

$$\begin{aligned} \tilde{\varphi}_{1,1}^I &= \left(\zeta_2(\bar{z}_1^*)^{-1} \varphi_2(\zeta_2(\bar{z}_1^*)) (A_{*,0}^{(2)}(\bar{z}_1^*) + A_{*,1}^{(2)}(\bar{z}_1^*) (\zeta_2(\bar{z}_1^*) I + G_1(\bar{z}_1^*))) \right. \\ &\quad + \sum_{k=1}^{\infty} \nu_{0,k} \hat{C}_2(\bar{z}_1^*, G_1(\bar{z}_1^*)) \sum_{l=1}^{k-1} \zeta_2(\bar{z}_1^*)^{k-l-1} G_1(\bar{z}_1^*)^{l-1} \\ &\quad - \sum_{k=1}^{\infty} \nu_{0,k} \sum_{l=1}^k \zeta_2(\bar{z}_1^*)^{k-l} G_1(\bar{z}_1^*)^{l-1} + \nu_{0,0} A_{*,1}^{(0)}(\bar{z}_1^*) \\ &\quad \left. + \varphi_1(\bar{z}_1^*) A_{*,1}^{(1)}(\bar{z}_1^*) \tilde{G}_{1,1} (I - C_1(\bar{z}_1^*, G_1(\bar{z}_1^*)))^{-1} \right) \end{aligned} \quad (4.55)$$

Derivation of $\tilde{\varphi}_{1,-1}^I$ and $\tilde{\varphi}_{1,1}^I$:

(1) $\tilde{\varphi}_{1,-1}^I$ of formula (4.53). From Propositions 5.9 and 5.11 of Ref. [7], we obtain

$$\tilde{\varphi}_{1,-1}^I = \lim_{\tilde{\Delta}_{\bar{z}_1^*} \ni z \rightarrow \bar{z}_1^*} (\bar{z}_1^* - z)^{\frac{1}{2}} \frac{\mathbf{g}_1(z) \operatorname{adj}(I - C_1(z, G_1(z)))}{\tilde{f}_1(1, (\bar{z}_1^* - z)^{\frac{1}{2}})} = \frac{\mathbf{g}_1(\bar{z}_1^*) \mathbf{v}^{C_1(\bar{z}_1^*)} \mathbf{u}^{C_1(\bar{z}_1^*)}}{-\tilde{\lambda}_{\zeta}^{C_1}(0)}.$$

(2) $\tilde{\varphi}_{1,1}^I$ of formula (4.55). From Propositions 5.7 and 5.8 of Ref. [7], we obtain

$$\begin{aligned} \tilde{\varphi}_{1,1}^I &= \frac{d}{d\zeta} \tilde{\mathbf{g}}_1(\zeta) (I - C_1(\bar{z}_1^* - \zeta^2, \tilde{G}_1(\zeta)))^{-1} \Big|_{\zeta=0} \\ &= (\tilde{\varphi}_{2,1}^{\hat{C}_2} - \tilde{\varphi}_{2,1}^{\tilde{G}_1} + \nu_{0,0} A_{*,1}^{(0)}(\bar{z}_1^*) \tilde{G}_{1,1}) (I - C_1(\bar{z}_1^*, G_1(\bar{z}_1^*)))^{-1} \\ &\quad + \mathbf{g}_1(\bar{z}_1^*) (I - C_1(\bar{z}_1^*, G_1(\bar{z}_1^*)))^{-1} A_{*,1}^{(1)}(\bar{z}_1^*) \tilde{G}_{1,1} (I - C_1(\bar{z}_1^*, G_1(\bar{z}_1^*)))^{-1}, \end{aligned}$$

and this leads us to formula (4.55).

4.10 Derivation of the coefficient vectors in Lemma 5.5 of Ref. [7]

For notation undefined in this subsection, see Section 5 of Ref. [7].

Lemma 5.5 of Ref. [7]. (1) If $\eta_2^{(c)} < \theta_2^{(c)}$, then the point $z = r_1$ is a pole of $\varphi_1(z)$ with order one and its Laurent series is represented as

$$\varphi_1(z) = \sum_{k=-1}^{\infty} \varphi_{1,k}^{II} (r_1 - z)^k, \quad (4.56)$$

where

$$\varphi_{1,-1}^{II} = \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) N_2(r_2) \mathbf{v}^{R_2}(r_2)}{\zeta_{2,z}(r_1)} \mathbf{u}_{s_0}^{G_1}(r_1) (I - C_1(r_1, G_1(r_1)))^{-1} > \mathbf{0}^\top \quad (4.57)$$

and $\mathbf{v}^{R_2}(r_2)$ is the right eigenvector of $R_2(r_2)$ with respect to the eigenvalue r_1^{-1} .

(2) If $\eta_2^{(c)} = \theta_2^{(c)}$ and $\psi_1(\bar{z}_1^*) > 1$, then the point $z = r_1$ is a pole of $\varphi_1(z)$ with order two and its Laurent series is represented as

$$\varphi_1(z) = \sum_{k=-2}^{\infty} \varphi_{1,k}^{II} (r_1 - z)^k, \quad (4.58)$$

where

$$\varphi_{1,-2}^{II} = \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) N_2(r_2) \mathbf{v}^{R_2}(r_2) \mathbf{u}_{s_0}^{G_1}(r_1) \mathbf{v}^{C_1}(r_1)}{\zeta_{2,z}(r_1) \psi_{1,z}(r_1)} \mathbf{u}^{C_1}(r_1) > \mathbf{0}^\top. \quad (4.59)$$

(3) If $\eta_2^{(c)} = \theta_2^{(c)}$ and $\psi_1(\bar{z}_1^*) = 1$, then the point $z = \bar{z}_1^*$ is a branch point of $\varphi_1(z)$ with order one and it is also a pole with order two. The Puiseux series for $\varphi_1(z)$ at $z = \bar{z}_1^*$ is represented as

$$\varphi_1(z) = \tilde{\varphi}_1((\bar{z}_1^* - z)^{\frac{1}{2}}) = \sum_{k=-2}^{\infty} \tilde{\varphi}_{1,k}^{II} (\bar{z}_1^* - z)^{\frac{k}{2}}, \quad (4.60)$$

where

$$\tilde{\varphi}_{1,-2}^{II} = \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) N_2(r_2) \mathbf{v}^{R_2}(r_2) \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*) \mathbf{v}^{C_1}(\bar{z}_1^*)}{\tilde{\alpha}_{s_0,1}^{G_1} \tilde{\lambda}_{\zeta}^{C_1}(0)} \mathbf{u}^{C_1}(\bar{z}_1^*) > \mathbf{0}^\top. \quad (4.61)$$

(4) If $\eta_2^{(c)} = \theta_2^{(c)}$ and $\psi_1(\bar{z}_1^*) < 1$, then the point $z = \bar{z}_1^*$ is a branch point of $\varphi_1(z)$ with order one and it is also a pole with order one. The Puiseux series for $\varphi_1(z)$ at $z = \bar{z}_1^*$ is represented as

$$\varphi_1(z) = \tilde{\varphi}_1((\bar{z}_1^* - z)^{\frac{1}{2}}) = \sum_{k=-1}^{\infty} \tilde{\varphi}_{1,k}^{II} (\bar{z}_1^* - z)^{\frac{k}{2}}, \quad (4.62)$$

where

$$\tilde{\varphi}_{1,-1}^{II} = \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) N_2(r_2) \mathbf{v}^{R_2}(r_2)}{-\tilde{\alpha}_{s_0,1}^{G_1}} \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*) (I - C_1(\bar{z}_1^*, G_1(\bar{z}_1^*)))^{-1} > \mathbf{0}^\top. \quad (4.63)$$

Derivation of $\varphi_{1,-1}^{II}$, $\varphi_{1,-2}^{II}$, $\tilde{\varphi}_{1,-2}^{II}$ and $\tilde{\varphi}_{1,-1}^{II}$:

Before deriving expressions for the coefficient vectors, we give the following proposition.

Proposition 4.2. *In the case of Type II, we have $\text{spr}(G_1(r_1)) = r_2$, $\text{spr}(R_2(r_2)) = r_1^{-1}$ and*

$$\mathbf{v}_{s_0}^{G_1}(r_1) = (r_1 I - G_2(r_2))^{-1} N_2(r_2) \mathbf{v}^{R_2}(r_2). \quad (4.64)$$

Proof. In the case of Type II, we have $\text{spr}(G_1(r_1)) = \zeta_2(r_1) = r_2$, $\text{spr}(R_1(r_1)) = \bar{\zeta}_2(r_1)^{-1} \leq r_2^{-1}$, $\text{spr}(G_2(r_2)) = \zeta_1(r_2) < r_1$ and $\text{spr}(R_2(r_2)) = \bar{\zeta}_1(r_2)^{-1} = r_1^{-1}$. Furthermore, by Lemma 4.1 and Remark 4.1 of Ref. [7], we have

$$\begin{aligned} I - C(r_1, r_2) &= (r_2^{-1}I - R_1(r_1))(I - H_1(r_1))(r_2I - G_1(r_1)) \\ &= (r_1^{-1}I - R_2(r_2))(I - H_2(r_2))(r_1I - G_2(r_2)). \end{aligned} \quad (4.65)$$

Multiplying both the sides of this equation by $\mathbf{v}_{s_0}^{G_1}(r_1)$ from the right, we obtain

$$(r_1^{-1}I - R_2(r_2))(I - H_2(r_2))(r_1I - G_2(r_2))\mathbf{v}_{s_0}^{G_1}(r_1) = \mathbf{0}. \quad (4.66)$$

Since both $(I - H_2(r_2))$ and $(r_1I - G_2(r_2))$ are nonsingular, we obtain

$$\mathbf{v}^{R_2}(r_2) = (I - H_2(r_2))(r_1I - G_2(r_2))\mathbf{v}_{s_0}^{G_1}(r_1) \neq \mathbf{0},$$

and this leads us to expression (4.64), where we use the fact that $N_2(r_2) = (I - H_2(r_2))^{-1}$. \square

(1) $\varphi_{1,-1}^{II}$ of formula (4.57). Note that we have

$$\begin{aligned} \mathbf{u}^{C_2}(r_2)(C_2(r_1, r_2) - I) &= \mathbf{u}^{C_2}(r_2)(C_2(r_1, r_2) - C_2(G_2(r_2), r_2)) \\ &= \mathbf{u}^{C_2}(r_2)A_{1,*}^{(2)}(r_2)(r_1I - G_2(r_2)). \end{aligned} \quad (4.67)$$

Hence, from Propositions 5.7 and 5.8 of Ref. [7], we obtain

$$\begin{aligned} \varphi_{1,-1}^{II} &= \lim_{z \rightarrow r_1} (r_1 - z) \mathbf{g}_1(z) (I - C_1(z, G_1(z)))^{-1} \\ &= (\varphi_{2,-1}^{\hat{C}_2} - \varphi_{2,-1}^{\hat{G}_1})(I - C_1(r_1, G_1(r_1)))^{-1} \\ &= \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) (r_1I - G_2(r_2)) \mathbf{v}_{s_0}^{G_1}(r_1) \mathbf{u}_{s_0}^{G_1}(r_1) (I - C_1(r_1, G_1(r_1)))^{-1}}{\zeta_{2,z}(r_1)}. \end{aligned}$$

This and Proposition 4.2 lead us to formula (4.57).

(2) $\varphi_{1,-2}^{II}$ of formula (4.59). From Propositions 5.3, 5.4, 5.7 and 5.8 of Ref. [7], we obtain

$$\begin{aligned} \varphi_{1,-2}^{II} &= \lim_{z \rightarrow r_1} (r_1 - z)^2 \frac{\mathbf{g}_1(z) \text{adj}(I - C_1(z, G_1(z)))}{f_1(1, z)} \\ &= \frac{(\varphi_{2,-1}^{\hat{C}_2} - \varphi_{2,-1}^{\hat{G}_1}) \mathbf{v}^{C_1}(r_1) \mathbf{u}^{C_1}(r_1)}{\psi_{1,z}(r_1)} \\ &= \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) (r_1I - G_2(r_2)) \mathbf{v}_{s_0}^{G_1}(r_1) \mathbf{u}_{s_0}^{G_1}(r_1) \mathbf{v}^{C_1}(r_1) \mathbf{u}^{C_1}(r_1)}{\zeta_{2,z}(r_1) \psi_{1,z}(r_1)}. \end{aligned}$$

This and Proposition 4.2 lead us to formula (4.59).

(3) $\tilde{\varphi}_{1,-2}^{II}$ of formula (4.61). From Propositions 5.7, 5.8, 5.9 and 5.11 of Ref. [7], we obtain

$$\begin{aligned}\tilde{\varphi}_{1,-2}^{II} &= \lim_{\tilde{\Delta}_{\bar{z}_1^*} \ni z \rightarrow \bar{z}_1^*} (\bar{z}_1^* - z) \frac{\tilde{\mathbf{g}}_1((\bar{z}_1^* - z)^{\frac{1}{2}}) \text{adj}(I - C_1(z, \tilde{G}_1((\bar{z}_1^* - z)^{\frac{1}{2}})))}{\tilde{f}_1(1, (\bar{z}_1^* - z)^{\frac{1}{2}})} \\ &= \frac{(\tilde{\varphi}_{2,-1}^{\hat{C}_2} - \tilde{\varphi}_{2,-1}^{\hat{G}_1}) \mathbf{v}^{C_1}(\bar{z}_1^*) \mathbf{u}^{C_1}(\bar{z}_1^*)}{-\tilde{\lambda}_\zeta^{C_1}(0)} \\ &= \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) (\bar{z}_1^* I - G_2(r_2)) \mathbf{v}_{s_0}^{G_1}(\bar{z}_1^*) \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*) \mathbf{v}^{C_1}(\bar{z}_1^*) \mathbf{u}^{C_1}(\bar{z}_1^*)}{\tilde{\alpha}_{s_0,1}^{G_1} \tilde{\lambda}_\zeta^{C_1}(0)}.\end{aligned}$$

This and Proposition 4.2 lead us to formula (4.61).

(4) $\tilde{\varphi}_{1,-1}^{II}$ of formula (4.63). From Propositions 5.7 and 5.8 of Ref. [7], we obtain

$$\begin{aligned}\tilde{\varphi}_{1,-1}^{II} &= \lim_{\tilde{\Delta}_{\bar{z}_1^*} \ni z \rightarrow \bar{z}_1^*} (\bar{z}_1^* - z)^{\frac{1}{2}} \tilde{\mathbf{g}}_1((\bar{z}_1^* - z)^{\frac{1}{2}}) (I - C_1(z, \tilde{G}_1((\bar{z}_1^* - z)^{\frac{1}{2}})))^{-1} \\ &= (\tilde{\varphi}_{2,-1}^{\hat{C}_2} - \tilde{\varphi}_{2,-1}^{\hat{G}_1}) (I - C_1(\bar{z}_1^*, G_1(\bar{z}_1^*)))^{-1} \\ &= \frac{c_{pole}^{\varphi_2} \mathbf{u}^{C_2}(r_2) A_{1,*}^{(2)}(r_2) (\bar{z}_1^* I - G_2(r_2)) \mathbf{v}_{s_0}^{G_1}(\bar{z}_1^*) \mathbf{u}_{s_0}^{G_1}(\bar{z}_1^*) (I - C_1(\bar{z}_1^*, G_1(\bar{z}_1^*)))^{-1}}{-\tilde{\alpha}_{s_0,1}^{G_1}}.\end{aligned}$$

This and Proposition 4.2 lead us to formula (4.63).

References

- [1] Fayolle, G., Malyshev, V.A., and Menshikov, M.V.: Topics in the Constructive Theory of Countable Markov Chains. Cambridge University Press, Cambridge (1995)
- [2] Horn, R.A. and Johnson, C.R.: Matrix Analysis. Cambridge University Press, Cambridge (1985)
- [3] Horn, R.A. and Johnson, C.R.: Topics in Matrix Analysis. Cambridge University Press, Cambridge (1991)
- [4] Miyazawa, M. and Zhao, Y.Q.: The stationary tail asymptotics in the GI/G/1 type queue with countably many background states. Advances in Applied Probability 36(4), 1231–1251 (2004)
- [5] Neuts, M.F.: Matrix-Geometric Solutions in Stochastic Models. Dover Publications, New York (1994)
- [6] Ozawa, T.: Asymptotics for the stationary distribution in a discrete-time two-dimensional quasi-birth-and-death process. Queueing Systems 74, 109–149 (2013)
- [7] Ozawa, T. and Kobayashi, M.: Exact asymptotic formulae of the stationary distribution of a discrete-time two dimensional QBD process. Queueing Systems (2018) (DOI:10.1007/s11134-018-9586-x)
- [8] Seneta, E.: Non-negative Matrices and Markov Chains, revised printing. Springer-Verlag, New York (2006)