

PARALLACTIC DISTANCES AND PROPER MOTIONS OF VIRTUALLY ALL STARS IN THE σ ORIONIS CLUSTER OR: HOW I LEARNED TO GET THE MOST OUT OF TOPCAT AND LOVE *Gaia* DR2

José A. Caballero¹

¹*Centro de Astrobiología (CSIC-INTA), ESAC, camino bajo del castillo s/n, E-28692 Villanueva de la Cañada, Madrid, Spain*

Keywords: astronomical data bases: miscellaneous — stars: massive — stars: low-mass — Galaxy: open clusters and associations (individual: σ Orionis)

The young σ Orionis open cluster is a cornerstone for studying X-ray emission, discs, jets, accretion, photometric variability, magnetism, abundances, and mass function of OB-type, Herbig Ae/Be and T Tauri stars, brown dwarfs, and objects beyond the deuterium burning mass limit (Garrison 1967; Caballero 2008a).

The *Gaia* Data Release 2 (DR2) was made public at noon (CEST) on 25 April 2018. Among many other things, it contains five-parameter solutions (α , δ , $\mu_\alpha \cos \delta$, μ_δ , π) for 1.332 million sources (Gaia Collaboration et al. 2018). By the evening of that date, and with the help of the Virtual Observatory tools TOPCAT and Aladin (Bonnarel et al. 2000; Taylor 2005), I had compiled more useful astro-photometric data on σ Orionis stars than all observations carried out in the previous five decades. Here I share useful new data that will facilitate further studies in the cluster.

First, I downloaded all *Gaia* DR2 data available on the 10,742 optical sources at less than 30 arcmin to the eponymous σ Ori star. It is located exactly at the center of the cluster, which is composed by a dense core that extends from the center to ~ 20 arcmin and a rarified halo at larger separations (Caballero 2008b). Because of abnormal *Gaia* $B_P - R_P$ colors and expected $G - K_s$ colors of known cluster low-mass stars and brown dwarfs with unequivocal features of extreme youth, I downloaded the 2MASS (Skrutskie et al. 2006) data of 5,742 near-infrared sources in the survey area. A total of 5,524 (96.2%) 2MASS sources have a *Gaia* counterpart at less than 1.0 arcsec. I also cross-matched all these sources with the Mayrit catalog of candidate and confirmed stars and brown dwarfs in σ Orionis (cf. Caballero 2008a). In spite of being over ten years old and suffering from known flaws (see corrections in Caballero et al. 2017), it is still the most precise and comprehensible catalog of stars and massive brown dwarfs in the cluster.

With TOPCAT, I plotted the color-magnitude diagram G vs. $G - K_s$ shown in Fig. 1. Next, I sketched two straight lines (not shown), redward of which all σ Orionis cluster members and candidates in the Mayrit catalog lie except for the background source 2MASS J05390697–0212168. The 816 *Gaia* DR2-2MASS sources redward of the lines are available in a comma-separated values file available as data behind Figure 1. They consist of true cluster members and stars that conservatively follow the photometric cluster sequence but are located closer or further or do not share the very low cluster proper motion (σ Orionis is in the antapex).

Of the 816 *Gaia* DR2-2MASS sources, 329 were tabulated in the Mayrit catalog (i.e. all except eight). Of the Mayrit stars and brown dwarfs, 281 have *Gaia* DR2 total proper motions $\mu < 15 \text{ mas a}^{-1}$ and parallaxes $1.5 \text{ mas} < \pi < 3.7 \text{ mas}$ and remain as good cluster member candidates. The 48 new outliers are stars either in the far background (43 stars including StHa 50) or near foreground (three stars: S Ori J053724.5–021856, HD 294297, S Ori 20), or that are well characterized T Tauri stars in the cluster but their parallaxes are affected by close multiplicity (two stars: Mayrit 873229 AB, Mayrit 359179 AB; Caballero et al. 2017). The mean and standard deviation of the parallaxes of the resulting 281 *Gaia* DR2-2MASS-Mayrit stars and brown dwarfs are $\bar{\pi} = 2.56 \text{ mas}$ and $\sigma_\pi = 0.29 \text{ mas}$, from where one can derive $d = 391 \pm 44 \text{ pc}$, identical within generous uncertainties to the value of $387.5 \pm 1.3 \text{ pc}$ derived by Schaefer et al. (2016) with interferometry.

Project AYA2016-79425-C3-2-P funded by MINECO/FEDER.




Figure 1. Color-magnitude diagram G vs. $G - K_s$ of all 2MASS-*Gaia* DR2 sources in a 30 arcmin-radius circular area centered on the eponymous σ Ori star, which illuminates the mane of Horsehead Nebula. Red circles: σ Orionis cluster member candidates in the Mayrit catalog (Caballero 2008a); blue dots: other sources that conservatively follow the cluster sequence (this work); grey tiny dots: other sources in the cluster fore- and background. The cluster sequence goes from O9.5 ($M \approx 17 M_{\odot}$) to M6–7 below the hydrogen burning limit ($M \approx 0.06\text{--}0.05 M_{\odot}$), i.e. covers almost three orders of magnitude in mass and over six orders of magnitude in optical flux. The virtual observatory-compliant .csv file for reproducing this figure, or investigating by yourself membership of stars in σ Orionis is available at RNAAS and at <http://exoterrae.eu/pub/soT.csv>.

REFERENCES

- Bonnarel, F., Fernique, P., Bienaymé, O., et al. 2000, A&AS, 143, 33
- Caballero, J. A. 2008a, A&A, 478, 667
- Caballero, J. A. 2008b, MNRAS, 383, 375
- Caballero, J. A., Novalbos, I., Tobal, T., & Miret, F. X. 2017, AN, in press (eprint arXiv:1707.00436)
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, A&A, in press (eprint arXiv:1804.09365)
- Garrison, R. F. 1967, PASP, 79, 433
- Schaefer, G. H., Hummel, C. A., Gies, D. R., et al. 2016, AJ, 152, 213
- Simón-Díaz, S., Caballero, J. A., Lorenzo, J., et al. 2015, ApJ, 799, 169
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163
- Taylor, M. B. 2005, Astronomical Data Analysis Software and Systems XIV, 347, 29