# ROTA–BAXTER OPERATORS AND POST-LIE ALGEBRA STRUCTURES ON SEMISIMPLE LIE ALGEBRAS

DIETRICH BURDE AND VSEVOLOD GUBAREV

ABSTRACT. Rota–Baxter operators R of weight 1 on  $\mathfrak n$  are in bijective correspondence to post-Lie algebra structures on pairs  $(\mathfrak{g}, \mathfrak{n})$ , where  $\mathfrak{n}$  is complete. We use such Rota–Baxter operators to study the existence and classification of post-Lie algebra structures on pairs of Lie algebras  $(\mathfrak{g}, \mathfrak{n})$ , where  $\mathfrak{n}$  is semisimple. We show that for semisimple  $\mathfrak{g}$  and  $\mathfrak{n}$ , with  $\mathfrak{g}$  or  $\mathfrak{n}$  simple, the existence of a post-Lie algebra structure on such a pair  $(\mathfrak{g}, \mathfrak{n})$  implies that  $\mathfrak{g}$  and  $\mathfrak{n}$  are isomorphic, and hence both simple. If  $\mathfrak n$  is semisimple, but  $\mathfrak g$  is not, it becomes much harder to classify post-Lie algebra structures on  $(\mathfrak{g}, \mathfrak{n})$ , or even to determine the Lie algebras  $\mathfrak{g}$  which can arise. Here only the case  $\mathfrak{n} = \mathfrak{sl}_2(\mathbb{C})$  was studied. In this paper we determine all Lie algebras g such that there exists a post-Lie algebra structure on  $(\mathfrak{g}, \mathfrak{n})$  with  $\mathfrak{n} = \mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{sl}_2(\mathbb{C})$ .

### 1. Introduction

Rota–Baxter operators were introduced by G. Baxter [\[3\]](#page-16-0) in 1960 as a formal generalization of integration by parts for solving an analytic formula in probability theory. Such operators  $R: A \rightarrow A$  are defined on an algebra A by the identity

$$
R(x)R(y) = R(R(x)y + xR(y) + \lambda xy)
$$

for all  $x, y \in A$ , where  $\lambda$  is a scalar, called the *weight* of R. These operators were then further investigated, by G.-C. Rota [\[31\]](#page-17-0), Atkinson [\[1\]](#page-16-1), Cartier [\[17\]](#page-16-2) and others. In the 1980s these operators were studied in integrable systems in the context of classical and modified Yang–Baxter equations [\[34,](#page-17-1) [4\]](#page-16-3). Since the late 1990s, the study of Rota–Baxter operators has made great progress in many areas, both in theory and in applications [\[26,](#page-16-4) [2,](#page-16-5) [23,](#page-16-6) [21,](#page-16-7) [22,](#page-16-8) [5,](#page-16-9) [20\]](#page-16-10).

Post-Lie algebras and post-Lie algebra structures also arise in many areas, e.g., in differential geometry and the study of geometric structures on Lie groups. Here post-Lie algebras arise as a natural common generalization of pre-Lie algebras [\[24,](#page-16-11) [27,](#page-17-2) [33,](#page-17-3) [6,](#page-16-12) [7,](#page-16-13) [8\]](#page-16-14) and LR-algebras [\[9,](#page-16-15) [10\]](#page-16-16), in the context of nil-affine actions of Lie groups, see [\[11\]](#page-16-17). A detailed account of the differential geometric context of post-Lie algebras is also given in [\[19\]](#page-16-18). On the other hand, post-Lie algebras have been introduced by Vallette [\[35\]](#page-17-4) in connection with the homology of partition posets and the study of Koszul operads. They have been studied by several authors in various contexts, e.g., for algebraic operad triples [\[29\]](#page-17-5), in connection with modified Yang– Baxter equations, Rota–Baxter operators, universal enveloping algebras, double Lie algebras, R-matrices, isospectral flows, Lie-Butcher series and many other topics [\[2,](#page-16-5) [19,](#page-16-18) [20\]](#page-16-10). There are several results on the existence and classification of post-Lie algebra structures, in particular on commutative post-Lie algebra structures [\[13,](#page-16-19) [14,](#page-16-20) [15\]](#page-16-21).

2000 Mathematics Subject Classification. Primary 17B20, 17D25.

Date: June 27, 2019.

Key words and phrases. Post-Lie algebra, Rota–Baxter operator.

### 2 D. BURDE AND V. GUBAREV

It is well-known [\[2\]](#page-16-5) that Rota–Baxter operators R of weight 1 on  $\mathfrak n$  are in bijective correspondence to post-Lie algebra structures on pairs  $(\mathfrak{g}, \mathfrak{n})$ , where  $\mathfrak{n}$  is complete. In fact, RB-operators always yield PA-structures. So it is possible (and desirable) to use results on RB-operators for the existence and classification of post-Lie algebra structures.

The paper is organized as follows. In section 2 we give basic definitions of RB-operators and PA-structures on pairs of Lie algebras. We summarize several useful results. For a complete Lie algebra **n** there is a bijection between PA-structures on  $(\mathfrak{g}, \mathfrak{n})$  and RB-operators of weight 1 on n. The PA-structure is given by  $x \cdot y = \{R(x), y\}$ . Here we study the kernels of R and  $R + id$ . If  $\mathfrak g$  and  $\mathfrak n$  are not isomorphic, then both R and  $R + id$  have a non-trivial kernel. Moreover, if one of g or n is not solvable, then at least one of  $\ker(R)$  and  $\ker(R + id)$  is non-trivial.

In section 3 we complete the classification of PA-structures on pairs of semisimple Lie algebras  $(\mathfrak{g}, \mathfrak{n})$ , where either  $\mathfrak{g}$  or  $\mathfrak{n}$  is simple. We already have shown the following in [\[11\]](#page-16-17). If  $\mathfrak{g}$  is simple, and there exists a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$ , then also  $\mathfrak{n}$  is simple, and we have  $\mathfrak{g} \cong \mathfrak{n}$  with  $x \cdot y = 0$ or  $x \cdot y = [x, y]$ . Here we deal now with the case that **n** is simple. Again it follows that **g** and n are isomorphic. The proof via RB-operators uses results of Koszul [\[28\]](#page-17-6) and Onishchik [\[30\]](#page-17-7). We also show a result concerning semisimple decompositions of Lie algebras. Suppose that  $\mathfrak{g} = \mathfrak{s}_1 + \mathfrak{s}_2$  is the vector space sum of two semisimple subalgebras of  $\mathfrak{g}$ . Then  $\mathfrak{g}$  is semisimple. As a corollary we show that the existence of a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  for  $\mathfrak{g}$  semisimple and  $\mathfrak{n}$ complete implies that  $\mathfrak n$  is semisimple.

In section 4 we determine all Lie algebras  $\mathfrak g$  which can arise by PA-structures on  $(\mathfrak g,\mathfrak n)$  with  $\mathfrak{n} = \mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{sl}_2(\mathbb{C})$ . This turns out to be much more complicated than the case  $\mathfrak{n} = \mathfrak{sl}_2(\mathbb{C})$ , which we have done in [\[11\]](#page-16-17). By Theorem 3.3 of [\[12\]](#page-16-22),  $\mathfrak{g}$  cannot be solvable unimodular. On the other hand, the result we obtain shows that there are more restrictions than that.

## 2. Preliminaries

Let A be a nonassociative algebra over a field K in the sense of Schafer [\[32\]](#page-17-8), with K-bilinear product  $A \times A \to A$ ,  $(a, b) \mapsto ab$ . We will assume that K is an arbitrary field of characteristic zero, if not said otherwise.

**Definition 2.1.** Let  $\lambda \in K$ . A linear operator  $R: A \rightarrow A$  satisfying the identity

(1) 
$$
R(x)R(y) = R(R(x)y + xR(y) + \lambda xy)
$$

for all x, y ∈ A is called a *Rota–Baxter operator on* A *of weight* λ, or just *RB-operator*.

Two obvious examples are given by  $R = 0$  and  $R = \lambda$  id, for an arbitrary nonassociative algebra. These are called the *trivial* RB-operators. The following elementary lemma was shown in [\[23\]](#page-16-6), Proposition 1.1.12.

**Lemma 2.2.** Let R be an RB-operator on A of weight  $\lambda$ . Then  $-R - \lambda$  id *is an RB*-operator *on* A *of weight*  $\lambda$ *, and*  $\lambda^{-1}R$  *is an RB-operator on* A *of weight* 1 *for all*  $\lambda \neq 0$ *.* 

It is also easy to verify the following results.

<span id="page-1-0"></span>**Proposition 2.3.** [\[5\]](#page-16-9) *Let* R *be an RB-operator on* A *of weight*  $\lambda$  *and*  $\psi \in \text{Aut}(A)$ *. Then*  $R^{(\psi)} = \psi^{-1} R \psi$  is an RB-operator on A of weight  $\lambda$ .

Proposition 2.4. [\[23\]](#page-16-6) *Let* B *be a countable direct sum of an algebra* A*. Then the operator* R *defined on* B *by*

$$
R((a_1, a_2, \ldots, a_n, \ldots)) = (0, a_1, a_1 + a_2, a_1 + a_2 + a_3, \ldots)
$$

*is an RB-operator on* B *of weight* 1*.*

<span id="page-2-0"></span>**Proposition 2.5.** Let  $B = A \oplus A$  and  $\psi \in \text{Aut}(A)$ . Then the operator R defined on B by

(2) 
$$
R((a_1, a_2)) = (0, \psi(a_1))
$$

*is an RB-operator on* B *of weight* 1*. Furthermore the operator* R *defined on* B *by*

(3) 
$$
R((a_1, a_2)) = (-a_1, -\psi(a_1))
$$

*is an RB-operator on* B *of weight* 1*.*

*Proof.* Let  $x = (a_1, a_2)$  and  $y = (b_1, b_2)$ . Then we have

$$
R(R(x)y + xR(y) + \lambda xy) = R((0, \psi(a_1)b_2 + (0, a_2\psi(b_1)) + (a_1b_1, a_2b_2))
$$
  
= (0, \psi(a\_1b\_1))  
= (0, \psi(a\_1)\psi(b\_1))  
= R(x)R(y).

The second claim follows similarly.

<span id="page-2-2"></span>**Proposition 2.6.** [\[26\]](#page-16-4) *Let*  $A = A_1 \oplus A_2$ ,  $R_1$  *be an RB-operator of weight*  $\lambda$  *on*  $A_1$ ,  $R_2$  *be an RB-operator of weight*  $\lambda$  *on*  $A_2$ . Then the operator  $R: A \rightarrow A$  defined by  $R((a_1, a_2))$  =  $(R_1(a_1), R_2(a_2))$  *is an RB-operator of weight*  $\lambda$  *on A*.

<span id="page-2-1"></span>**Proposition 2.7.** [\[23\]](#page-16-6) *Let*  $A = A_1 \dot{+} A_2$  *be the direct vector space sum of two subalgebras. Then the operator* R *defined on* A *by*

$$
(4) \t R(a_1 + a_2) = -\lambda a_2
$$

*for*  $a_1 \in A_1$  *and*  $a_2 \in A_2$  *is an RB-operator on* A *of weight*  $\lambda$ *.* 

We call such an operator *split*, with subalgebras  $A_1$  and  $A_2$ . Note that the set of all split RB-operators on A is in bijective correspondence with all decompositions  $A = A_1 \dot{+} A_2$  as a direct sum of subalgebras.

**Lemma 2.8.** [\[5\]](#page-16-9) Let R be an RB-operator of nonzero weight  $\lambda$  on an algebra A. Then R is *split if and only if*  $R(R + \lambda id) = 0$ *.* 

**Lemma 2.9.** Let  $A = A_+ + A_0 + A_+$  be a direct vector space sum of subalgebras of A. Suppose *that* R *is an RB-operator of weight*  $\lambda$  *on*  $A_0$ ,  $A_0$  *is an*  $(R + id)(A_0)$ *-module and*  $A_+$  *is an* R(A0)*-module. Define an operator* P *on* A *by*

(5) 
$$
P_{|A_{-}} = 0, P_{|A_{0}} = R, P_{|A_{+}} = -\lambda \operatorname{id}.
$$

*Then*  $P$  *is an RB-operator on*  $A$  *of weight*  $\lambda$ *.* 

**Definition 2.10.** Let P be an RB-operator on A defined as above such that not both  $A_-\$  and A<sup>+</sup> are zero. Then P is called *triangular-split*.

We also recall the definition of post-Lie algebra structures on a pair of Lie algebras  $(\mathfrak{g}, \mathfrak{n})$ over  $K$ , see [\[11\]](#page-16-17).

**Definition 2.11.** Let  $\mathfrak{g} = (V, [ , ])$  and  $\mathfrak{n} = (V, { , })$  be two Lie brackets on a vector space V over K. A *post-Lie algebra structure*, or *PA-structure* on the pair (g, n) is a K-bilinear product  $x \cdot y$  satisfying the identities:

<span id="page-3-5"></span>
$$
(6) \qquad \qquad x \cdot y - y \cdot x = [x, y] - \{x, y\}
$$

<span id="page-3-1"></span>(7) 
$$
[x, y] \cdot z = x \cdot (y \cdot z) - y \cdot (x \cdot z)
$$

<span id="page-3-0"></span>(8) 
$$
x \cdot \{y, z\} = \{x \cdot y, z\} + \{y, x \cdot z\}
$$

for all  $x, y, z \in V$ .

Define by  $L(x)(y) = x \cdot y$  the left multiplication operator of the algebra  $A = (V, \cdot)$ . By [\(8\)](#page-3-0), all  $L(x)$  are derivations of the Lie algebra  $(V, \{,\})$ . Moreover, by [\(7\)](#page-3-1), the left multiplication

$$
L: \mathfrak{g} \to \text{Der}(\mathfrak{n}) \subseteq \text{End}(V), \ x \mapsto L(x)
$$

is a linear representation of g.

If n is abelian, then a post-Lie algebra structure on (g, n) corresponds to a *pre-Lie algebra structure* on **g**. In other words, if  $\{x, y\} = 0$  for all  $x, y \in V$ , then the conditions reduce to

$$
x \cdot y - y \cdot x = [x, y],
$$
  
\n
$$
[x, y] \cdot z = x \cdot (y \cdot z) - y \cdot (x \cdot z),
$$

i.e.,  $x \cdot y$  is a *pre-Lie algebra structure* on the Lie algebra  $\mathfrak{g}$ , see [\[11\]](#page-16-17).

**Definition 2.12.** Let  $x \cdot y$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$ . If there exists a  $\varphi \in \text{End}(V)$  such that

$$
x \cdot y = \{\varphi(x), y\}
$$

for all  $x, y \in V$ , then  $x \cdot y$  is called an *inner* PA-structure on  $(\mathfrak{g}, \mathfrak{n})$ .

The following result is proved in [\[2\]](#page-16-5), Corollary 5.6.

<span id="page-3-2"></span>Proposition 2.13. *Let* (n, {, }, R) *be a Lie algebra together with a Rota–Baxter operator* R *of weight* 1*, i.e., a linear operator satisfying*

$$
{R(x), R(y)} = R({R(x), y} + {x, R(y)} + {x, y})
$$

*for all*  $x, y \in V$ *. Then* 

$$
x \cdot y = \{R(x), y\}
$$

*defines an inner PA-structure on* (g, n)*, where the Lie bracket of* g *is given by*

<span id="page-3-4"></span>(9) 
$$
[x, y] = \{R(x), y\} - \{R(y), x\} + \{x, y\}.
$$

Note that ker(R) is a subalgebra of **n**. For  $x, y \in \text{ker}(R)$  we have  $R(\lbrace x, y \rbrace) = 0$ . Recall that a Lie algebra is called *complete*, if it has trivial center and only inner derivations.

Proposition 2.14. *Let* n *be a Lie algebra with trivial center. Then any inner PA-structure on* (g, n) *arises by a Rota–Baxter operator of weight* 1*. Furthermore, if* n *is complete, then every PA-structure on* (g, n) *is inner.*

*Proof.* The first claim follows from Proposition 2.10 in [\[11\]](#page-16-17). By Lemma 2.9 in [\[11\]](#page-16-17) every PAstructure on  $(\mathfrak{g}, \mathfrak{n})$  with complete Lie algebra  $\mathfrak{n}$  is inner. The result can also be derived from the proof of Theorem 5.10 in [\[2\]](#page-16-5).  $\square$ 

<span id="page-3-3"></span>Corollary 2.15. *Let* n *be a complete Lie algebra. Then there is bijection between PA-structures on* (g, n) *and RB-operators of weight* 1 *on* n*.*

#### PA-STRUCTURES 5

As we have seen, any inner PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  with  $Z(\mathfrak{n}) = 0$  arises by a Rota–Baxter operator of weight 1. For Lie algebra n with non-trivial center this need not be true.

**Example 2.16.** Let  $(e_1, e_2, e_3)$  be a basis of V and  $\mathfrak{n} = \mathfrak{r}_2(K) \oplus K$  with  $\{e_1, e_2\} = e_2$ . Then

$$
\varphi = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ \alpha & \beta & \gamma \end{pmatrix}
$$

*defines an inner PA-structure on*  $(\mathfrak{g}, \mathfrak{n})$  *by*  $x \cdot y = \{\varphi(x), y\}$  *with*  $\mathfrak{g} = \mathfrak{n}$ *, i.e., with*  $[e_1, e_2] = e_2$ *. But*  $\varphi$  *is not always a Rota–Baxter operator of weight* 1 *for* **n**. It is easy to see that this is the *case if and only if*  $\beta = 0$ *.* 

Proposition 2.17. *Let* x · y *be an inner PA-structure arising from an RB-operator* R *on* n *of weight* 1*. Then* R *is also an RB-operator of weight* 1 *on* g*, i.e., it satisfies*

$$
[R(x), R[y)] = R([R(x), y] + [x, R(y)] + [x, y])
$$

*for all*  $x, y \in V$ *.* 

*Proof.* Because of  $R([x, y]) = \{R(x), R(y)\}\$ and the definition of  $[x, y]$  we have

$$
R([R(x), y] + [x, R(y)] + [x, y]) = \{R(R(x)), R(y)\} + \{R(x), R(R(y))\} + \{R(x), R(y)\}
$$
  
= [R(x), R(y)]

for all  $x, y \in V$ .

**Corollary 2.18.** Let  $x \cdot y = \{R(x), y\}$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  defined by an RB-operator R *of weight* 1 *on* n*. Denote by* g<sup>i</sup> *be the Lie algebra structure on* V *defined by*

$$
[x, y]_0 = \{x, y\},
$$
  

$$
[x, y]_{i+1} = [R(x), y]_i - [R(y), x]_i + [x, y]_i,
$$

*for all*  $i \geq 0$ *. Then* R *defines a PA-structure on each pair*  $(\mathfrak{g}_{i+1}, \mathfrak{g}_i)$ *.* 

We have  $[x, y]_1 = [x, y]$ , and both R and  $R + id$  are Lie algebra homomorphisms from  $\mathfrak{g}_{i+1}$ to  $\mathfrak{g}_i$ , see Proposition 7 in [\[34\]](#page-17-1). Hence we obtain a composition of homomorphisms

$$
\mathfrak{g}_i \xrightarrow[R+\text{id}]{R} \mathfrak{g}_{i-1} \xrightarrow[R+\text{id}]{R} \cdots \xrightarrow[R+\text{id}]{R} \mathfrak{g}_0
$$

So the kernels  $\ker(R^i)$  and  $\ker((R + id)^i)$  are ideals in  $\mathfrak{g}_j$  for all  $1 \leq i \leq j$ .

For a Lie algebra  $\mathfrak{g}$ , denote by  $\mathfrak{g}^{(i)}$  the derived ideals defined by  $\mathfrak{g}^{(1)} = \mathfrak{g}$  and  $\mathfrak{g}^{(i+1)} = [\mathfrak{g}^{(i)}, \mathfrak{g}^{(i)}]$ for  $i \geq 1$ . An immediate consequence of Proposition 2.[13](#page-3-2) is the following observation.

**Proposition 2.19.** Let  $x \cdot y = \{R(x), y\}$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  defined by an RB-operator R *of weight* 1 *on*  $\mathfrak{n}$ *. Then we have* dim  $\mathfrak{g}^{(i)} \leq \dim \mathfrak{n}^{(i)}$  *for all*  $i \geq 1$ *.* 

**Corollary 2.20.** Let  $x \cdot y$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$ , where  $\mathfrak{n}$  is complete. Then we have  $\dim \mathfrak{g}^{(i)} \leq \dim \mathfrak{n}^{(i)}$  for all  $i \geq 1$ . In particular, if  $\mathfrak n$  is solvable, so is  $\mathfrak g$ , and if  $\mathfrak g$  is perfect, so is n*.*

*Proof.* By Corollary 2.[15](#page-3-3) this follows from the proposition. □

<span id="page-4-0"></span>**Proposition 2.21.** Let  $x \cdot y = \{R(x), y\}$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  defined by an RB-operator R *of weight* 1 *on* n*. Then the following holds.*

- (1) If  $\mathfrak g$  *and*  $\mathfrak n$  *are not isomorphic, then both*  $R$  *and*  $R$  + id *have a non-trivial kernel.*
- (2) If either  $\mathfrak g$  or  $\mathfrak n$  *is not solvable, then at least one of the operators* R and  $R + id$  has a *non-trivial kernel.*

*Proof.* For (1), assume that ker(R) = 0. Then  $R: \mathfrak{g} \to \mathfrak{n}$  is invertible, hence an isomorphism. This is a contradiction. The same is true for  $R+id$ . For (2) assume that  $\ker(R) = \ker(R+id) =$ 0. Then R and R + id are isomorphisms from  $\mathfrak{g}$  to  $\mathfrak{n}$ , and  $\mathfrak{g} \cong \mathfrak{n}$ . Then we can apply a result of Jacobson [\[25\]](#page-16-23) to the automorphism  $\psi := (R + id) \circ R^{-1}$  of n, because n is not solvable. We obtain a nonzero fixed point  $x \in \mathfrak{n}$ , so that

$$
0 = \psi(x) - x = (R + id)R^{-1}(x) - x = R^{-1}(x).
$$

Since R is bijective,  $x = 0$ , a contradiction.

<span id="page-5-0"></span>Corollary 2.22. *Let* n *be a simple Lie algebra and* R *be an invertible RB-operator of nonzero weight*  $\lambda$  *on* **n***. Then we have*  $R = -\lambda$  id.

*Proof.* By rescaling we may assume that R has weight 1. We obtain a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$ by Proposition 2.[13,](#page-3-2) with Lie bracket [\(9\)](#page-3-4) on  $\mathfrak{g}$ . Since  $\mathfrak{n}$  is not solvable, either R or  $R + id$  have a nontrivial kernel. But  $\ker(R) = 0$  by assumption, so that  $\ker(R + id)$  is a nontrivial ideal of **n**. Hence we have  $R + id = 0$ .

### 3. PA-structures on pairs of semisimple Lie algebras

We will assume that all algebras in this section are finite-dimensional. Let  $x \cdot y$  be a PAstructure on  $(\mathfrak{g}, \mathfrak{n})$  over  $\mathbb{C}$ , where  $\mathfrak{g}$  is simple and  $\mathfrak{n}$  is semisimple. Then  $\mathfrak{n}$  is also simple, and both  $\mathfrak g$  and  $\mathfrak n$  are isomorphic, see Proposition 4.9 in [\[11\]](#page-16-17). We have a similar result for  $\mathfrak n$  simple and g semisimple. However, its proof is more difficult than the first one.

**Theorem 3.1.** Let  $x \cdot y$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  over  $\mathbb{C}$ , where  $\mathfrak{n}$  is simple and  $\mathfrak{g}$  is semisim*ple. Then* g *is also simple, and both* g *and* n *are isomorphic.*

*Proof.* By Corollary 2.[15](#page-3-3) we have  $x \cdot y = \{R(x), y\}$  for an RB-operator R of weight 1 on n. Assume that g and n are not isomorphic. By Proposition 2.[21](#page-4-0) (2) both ker(R) and ker(R+id) are proper nonzero ideals of  $\mathfrak{g}$ , with ker(R) ∩ ker(R + id) = 0. So we have

$$
\mathfrak{g} = \ker(R) \oplus \ker(R + id) \oplus \mathfrak{s}
$$

with a semisimple ideal  $\mathfrak{s}$ . We have  $\mathfrak{n} = \text{im}(R) + \text{im}(R+\text{id})$  because of  $x = R(-x) + (R+\text{id})(x)$ for all  $x \in \mathfrak{n}$ , and

$$
\operatorname{im}(R) \cong \mathfrak{g}/\ker(R) \cong \ker(R + \operatorname{id}) \oplus \mathfrak{s},
$$
  

$$
\operatorname{im}(R + \operatorname{id}) \cong \mathfrak{g}/\ker(R + \operatorname{id}) \cong \ker(R) \oplus \mathfrak{s}.
$$

This yields a semisimple decomposition

$$
\mathfrak{n} = (\ker(R + id) \oplus \mathfrak{s}) + (\ker(R) \oplus \mathfrak{s}).
$$

Suppose that  $\mathfrak s$  is nonzero. Then both summands are not simple. This is a contradiction to Theorem 4.2 in Onishchik's paper [\[30\]](#page-17-7), which says that at least one summand in a semisimple decomposition of a simple Lie algebra must be simple. Hence we obtain  $\mathfrak{s} = 0$ ,  $\text{im}(R) =$  $\ker(R + id), \, \text{im}(R + id) = \text{im}(R)$  and

$$
\mathfrak{n} = \text{im}(R) + \text{im}(R + \text{id}).
$$

#### PA-STRUCTURES 7

Then the main result of Koszul's note [\[28\]](#page-17-6) implies that  $\mathfrak{n} = \text{im}(R) \oplus \text{im}(R + \text{id})$ , which is a contradiction to the simplicity of **n**. Hence  $\mathfrak{g}$  and **n** are isomorphic.

If  $\mathfrak g$  is semisimple with only two simple summands, we can prove the same result for any field K of characteristic zero.

**Proposition 3.2.** Let  $x \cdot y$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$ , where  $\mathfrak{n}$  is semisimple, and  $\mathfrak{g} = \mathfrak{s}_1 \oplus \mathfrak{s}_2$ *is the direct sum of two simple ideals of* g*. Then* g *and* n *are isomorphic.*

The proof is the same as before. The only argument where we needed the complex numbers, was the result of [\[30\]](#page-17-7), which we do not need here.

Let  $\mathfrak{n} = \mathfrak{s}_1 \oplus \mathfrak{s}_2$  be a direct sum of two simple isomorphic ideals  $\mathfrak{s}_1$  and  $\mathfrak{s}_2$ . We would like to find all RB-operators of weight 1 on  $\mathfrak n$  such that  $\mathfrak g$  with bracket [\(9\)](#page-3-4) is isomorphic to  $\mathfrak n$ .

**Proposition 3.3.** All PA-structures on  $(\mathfrak{g}, \mathfrak{n})$  with  $\mathfrak{g} \cong \mathfrak{n} = \mathfrak{s}_1 \oplus \mathfrak{s}_2$ , where  $\mathfrak{s}_1$  and  $\mathfrak{s}_2$  simple *isomorphic ideals of* n*, arise by the trivial RB-operators or by one of the following RB-operators* R *on*  $\mathfrak{n}$ *, and*  $\psi \in \text{Aut}(\mathfrak{n})$ *,* 

$$
R((s_1, s_2)) = (-s_1, -\psi(s_1)),
$$
  
\n
$$
R((s_1, s_2)) = (0, \psi(s_1)),
$$
  
\n
$$
R((s_1, s_2)) = (-s_1, 0)),
$$

*up to permuting the factors and application of*  $\varphi(R) = -R - id$  *to these operators.* 

*Proof.* By Proposition [2](#page-2-0).5 and Proposition 2.[7](#page-2-1) the given operators are RB-operators of weight 1 on **n**, because R is. By Proposition 2.[21](#page-4-0) at least one of  $\text{ker}(R)$  and  $\text{ker}(R + id)$  is nonzero. Suppose first that both ker(R) and ker(R+id) are zero. Then we have  $\mathfrak{g} = \text{ker}(R) \oplus \text{ker}(R+\text{id})$ and  $\mathfrak{n} = \text{ker}(R) + \text{ker}(R + id)$ . It is easy to see that  $\text{ker}(R)$  coincides with  $\mathfrak{s}_1$  or  $\mathfrak{s}_2$  by using the Theorem of Koszul [\[28\]](#page-17-6). Applying  $\varphi$  if necessary, we can assume that ker(R) =  $\mathfrak{s}_2$ . Then again by Koszul's result we have  $R((s_1, s_2)) = (\psi_1(s_1), \psi_2(s_1))$  or  $R((s_1, s_2)) = (\psi_1(s_1), 0)$  for some  $\psi_1, \psi_2 \in \text{Aut}(\mathfrak{n})$ . Since  $\text{im}(R) = \text{ker}(R + \text{id})$  we either have  $R((s_1, s_2)) = (-s_1, -\psi(s_1))$ or  $R((s_1, s_2)) = (-s_1, 0).$ 

In the second case, one of the kernels is zero. Applying  $\varphi$  if necessary, we may assume that  $\ker(R + id) = 0$  and  $\ker(R) = \mathfrak{s}_1$ . Then  $\mathfrak{g}/\ker(R)$  is a simple Lie algebra, and  $-R - id$  is an invertible RB-operator of weight 1 on  $\mathfrak{g}/\text{ker}(R)$ . By Corollary 2.[22](#page-5-0) we obtain  $-R - \text{id} =$ - id, hence  $R = 0$  on  $\mathfrak{g}/\text{ker}(R)$ . This implies  $R^2 = 0$  on  $\mathfrak{g}$ . The projections of im $(R)$  to  $\mathfrak{s}_1$ and  $\mathfrak{s}_2$  are either zero or an isomorphism on one factor. So we have  $R((s, 0)) = (0, \psi(s))$  or  $R((s, 0)) = (\psi_1(s), \psi_2(s))$  for some automorphisms  $\psi, \psi_1, \psi_2$ . But the second operator does not satisfy  $R^2 = 0$ , and hence is impossible. Therefore we are done.

<span id="page-6-0"></span>**Proposition 3.4.** Let  $x \cdot y = \{R(x), y\}$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  defined by an RB-operator R of weight 1 on  $\mathfrak{n}$ . Let  $\mathfrak{n}_1 = \ker(R^n)$ ,  $\mathfrak{n}_2 = \ker(R + id)^n$ ,  $\mathfrak{n}_3 = \text{im}(R^n) \cap \text{im}((R + id)^n)$  for  $n = \dim(V)$ *. Then*  $\mathfrak{n} = \mathfrak{n}_1 + \mathfrak{n}_2 + \mathfrak{n}_3$  *with*  $\{\mathfrak{n}_1, \mathfrak{n}_3\} \subseteq \mathfrak{n}_1$ ,  $\{\mathfrak{n}_2, \mathfrak{n}_3\} \subseteq \mathfrak{n}_2$ *, and*  $\mathfrak{n}_3$  *is solvable.* 

*Proof.* We first show by induction that  $\ker(R^i)$  is a subalgebra of **n**, and that

$$
\{\ker(R^i), \operatorname{im}((R+\operatorname{id})^i)\} \subseteq \ker(R^i)
$$

for all  $i \geq 1$ . The case  $i = 1$  goes as follows. We already know that  $\ker(R)$  is a subalgebra of n. So we have to show that  $\{\ker(R), \operatorname{im}(R + id)\}\subseteq \ker(R)$ . Let  $x \in \ker(R)$  and  $y \in \mathfrak{n}$ . Then by [\(6\)](#page-3-5) we have

$$
\{x, (R + id)(y)\} = \{x, R(y)\} + \{x, y\}
$$

$$
= [x, y] + \{y, R(x)\}
$$

$$
= [x, y],
$$

which is in ker(R), since this is an ideal in g. For the induction step  $i \mapsto i + 1$  consider the iteration of the Lie bracket [\(9\)](#page-3-4) for all  $i \geq 0$ , given by

$$
[x, y]_i = [x, y]_{i+1} - [R(x), y]_i - [x, R(y)]_i
$$

for all  $i \geq 0$ . Then

$$
\begin{aligned} \{x, y\} &= [x, y]_1 - [R(x), y]_0 - [x, R(y)]_0 \\ &= [x, y]_2 - [R^2(x), y]_0 - 2[R(x), y]_0 - 2[R(x), R(y)]_0 - 2[x, R(y)]_0 - [x, R^2(y)]_0 \end{aligned}
$$

and so on. Define a degree of a term  $[R^l(x), R^k(y)]_m$  by  $l + k + m$ , and let  $x, y \in \text{ker}(R^{i+1})$ . We can iterate the brackets, until the degree of every summand on the right-hand side will be greater than 3*i*, so that all summands either have a term  $R^l(x)$  with  $l > i$ , or a term  $R^k(y)$  with  $k > i$ , or all summands lie in  $[\ker(R^{i+1}), \ker(R^{i+1})]_{i+1}$ . By induction hypothesis, such terms will vanish for  $l > i$  or  $k > i$ , and since ker $(R^{i+1})$  is an ideal in  $\mathfrak{g}_{i+1}$ , we have  $\{x, y\} \in \text{ker}(R^{i+1})$ , so that ker( $R^{i+1}$ ) is a subalgebra of **n**. The induction step for the second claim follows similarly.

Since the image of a subalgebra under the action of an RB-operator is a subalgebra,  $n_1$ ,  $n_2$  and their intersection  $n_3$  are subalgebras of n. We want to show that  $n = n_1 + n_2 + n_3$ . Because of  $\ker(R^n) \cap \text{im}(R^n) = 0$  we have  $\mathfrak{n} = \ker(R^n) + \text{im}(R^n)$ . In the same way we have  $\mathfrak{n} = \text{ker}((R + \text{id})^n) \dot{+} \text{im}((R + \text{id})^n)$ . We obtain

$$
\operatorname{im}(R^n) \cap \operatorname{ker}((R+\operatorname{id})^n) + \operatorname{im}(R^n) \cap \operatorname{im}((R+\operatorname{id})^n) \subseteq \operatorname{im}(R^n).
$$

We claim that  $\ker((R + id)^n) \subseteq \text{im}(R^n)$ , so that we have equality above. Indeed, for  $x \in$  $\ker((R + id)^n)$  we have by the binomial formula

$$
x + {n \choose n-1} R(x) + \dots + {n \choose 1} R^{n-1}(x) = -R^{n}(x) \in \text{im}(R^{n}).
$$

Applying  $R^{n-1}$  we obtain  $R^{n-1}(x) \in \text{im}(R^n)$  and

$$
x + nR(x) + \dots + \binom{n}{2} R^{n-2}(x) \in \text{im}(R^n).
$$

Iterating this we obtain  $x \in \text{im}(R^n)$ . This yields

$$
\begin{aligned} \mathfrak{n} &= \ker(R^n) \dot{+} \operatorname{im}(R^n) \\ &= \ker(R^n) \dot{+} \ker((R + \mathrm{id})^n) \dot{+} \operatorname{im}(R^n) \cap \operatorname{im}((R + \mathrm{id})^n) \\ &= \mathfrak{n}_1 \dot{+} \mathfrak{n}_2 \dot{+} \mathfrak{n}_3. \end{aligned}
$$

On  $\mathfrak{n}_3$  both operators R and  $R + id$  are invertible. By Proposition 2.[21](#page-4-0) part (2) it follows that  $\mathfrak{n}_3$  is solvable.

<span id="page-7-0"></span>**Corollary 3.5.** The decomposition  $\mathfrak{n} = \mathfrak{n}_1 + \mathfrak{n}_2 + \mathfrak{n}_3$  induces a decomposition  $\mathfrak{g}_i = \mathfrak{n}_1 + \mathfrak{n}_2 + \mathfrak{n}_3$ *for each*  $i \geq 1$  *with the same properties as in the Proposition. The Lie algebras*  $(\mathfrak{n}_j, [,]_i)$  *and*  $(\mathfrak{n}_j, [,]_0)$  *are isomorphic for*  $j = 1, 2, 3$ *.* 

*Proof.* Since R and  $R + id$  are RB-operators on all  $\mathfrak{g}_i$ , we obtain the same decomposition with the same subalgebras. Note that  $R + id$  is invertible on  $\mathfrak{n}_1$ , R is invertible on  $\mathfrak{n}_2$  and both are invertible on  $\mathfrak{n}_3$ . In order to show that  $(\mathfrak{n}_1, [,]_i$  is isomorphic to  $(\mathfrak{n}_1, [,]_0$ , we consider a chain of isomorphisms

$$
(\mathfrak{n}_1, [,]_n) \xrightarrow{R+\mathrm{id}} (\mathfrak{n}_1, [,]_{n-1}) \xrightarrow{R+\mathrm{id}} \cdots \xrightarrow{R+\mathrm{id}} (\mathfrak{n}_1, [,]_0).
$$

In a similar way we can deal with  $\mathfrak{n}_2$  and  $\mathfrak{n}_3$ .

*Note that Proposition* 3.6 *is not correct. Hence the proof of Proposition* 3.7 *and* 3.8 *is invalid. However, the statement of both results is true and we have given a new proof of it in our paper* [\[16\]](#page-16-24) *on decompositions of algebras and post-associative algebra structures.*

<span id="page-8-0"></span>**Proposition 3.6.** Let  $g = s_1 + s_2$  be the vector space sum of two complex semisimple subalgebras *of* g*. Then* g *is semisimple.*

*Proof.* Suppose that the claim is not true and let  $\mathfrak{g}$  be a counterexample of minimal dimension. Then g contains a nonzero abelian ideal a. Then we obtain

$$
\mathfrak{g}/\mathfrak{a}=\mathfrak{s}_1/(\mathfrak{s}_1\cap\mathfrak{a})+\mathfrak{s}_2/(\mathfrak{s}_2\cap\mathfrak{a}).
$$

Since  $\mathfrak{s}_1 \cap \mathfrak{a}$  is an abelian ideal  $\mathfrak{s}_1$ , it must be zero, i.e.,  $\mathfrak{s}_1 \cap \mathfrak{a} = 0$ . In the same way we have  $s_2 \cap \mathfrak{a} = 0$ . Hence we obtain a semisimple decomposition of  $\mathfrak{g}/\mathfrak{a}$  with  $\dim(\mathfrak{g}/\mathfrak{a}) < \dim(\mathfrak{g})$ . If  $\mathfrak{g}/\mathfrak{a}$ is semisimple, this is a contradiction to the minimality of the counterexample g. Otherwise we may assume that  $\mathfrak g$  has 1-dimensional solvable radical. Then  $\mathfrak g$  is reductive, and by Theorem 3.2 of [\[30\]](#page-17-7), there are no semisimple decompositions of a complex reductive non-semisimple Lie algebra. Hence we are done.

**Proposition 3.7.** Let  $x \cdot y = \{R(x), y\}$  be a PA-structure on  $(\mathfrak{g}, \mathfrak{n})$  over  $\mathbb{C}$ , where  $\mathfrak{n}$  is simple, *defined by an RB-operator* R *of weight* 1 *on*  $\mathfrak{n}$ *, with associated Lie algebras*  $\mathfrak{g}_i$  *for*  $i = 1, \ldots, n =$  $\dim(V)$ *. Assume that*  $\mathfrak{g}_0 = \mathfrak{n}$  *and*  $\mathfrak{g}_n$  *are semisimple. Then all*  $\mathfrak{g}_i$  *are isomorphic to* **n**.

*Proof.* Since  $\mathfrak{n}_1$  and  $\mathfrak{n}_2$  are kernels of homomorphisms, they are ideals in  $\mathfrak{g}_n$ . The quotient  $\mathfrak{g}_n/(\mathfrak{n}_1 + \mathfrak{n}_2) \cong \mathfrak{n}_3$  is semisimple and solvable by Proposition [3](#page-6-0).4. Hence  $\mathfrak{n}_3 = 0$ , and we obtain  $\mathfrak{g}_n = \text{ker}(R^n) \oplus \text{ker}((R + \text{id})^n)$ . Because of Corollary [3](#page-7-0).5 we have the decomposition  $\mathfrak{g}_i = \text{ker}(R^n) + \text{ker}((R + id)^n)$  for all  $i < n$ , where all Lie algebras  $(\text{ker}(R^n), [,]_i)$  are isomorphic, and all Lie algebras  $(\ker((R+id)^n), [,]_i)$  are isomorphic. By Proposition [3](#page-8-0).6 all  $\mathfrak{g}_i$  are semisimple. By Koszul's result [\[28\]](#page-17-6), all  $\mathfrak{g}_i$  are isomorphic.

Proposition 3.8. *Suppose that there is a post-Lie algebra structure on* (g, n) *over* C*, where* g *is semisimple and* n *is complete. Then* n *must be semisimple.*

*Proof.* By Corollary 2.[15](#page-3-3) the PA-structure is given by  $x \cdot y = \{R(x), y\}$ , where R is an RBoperator of weight 1 on n. If at least one of ker(R) and ker(R+id) is trivial, we obtain  $\mathfrak{g} \cong \mathfrak{n}$ by Proposition 2.[21,](#page-4-0) part (1). Otherwise  $\mathfrak{n} = \text{im}(R) + \text{im}(R + \text{id})$  is the sum of two nonzero semisimple subalgebras. By Proposition [3](#page-8-0).6  $\boldsymbol{\mathfrak{n}}$  is semisimple.

4. PA-STRUCTURES ON  $(\mathfrak{g}, \mathfrak{n})$  with  $\mathfrak{n} = \mathfrak{sl}_2(\mathbb{C}) \times \mathfrak{sl}_2(\mathbb{C})$ 

In [\[11\]](#page-16-17), Proposition 4.7 we have shown that PA-structures with  $\mathfrak{n} = \mathfrak{sl}_2(\mathbb{C})$  exist on  $(\mathfrak{g}, \mathfrak{n})$ if and only if g is isomorphic to  $\mathfrak{sl}_2(\mathbb{C})$ , or to one of the solvable non-unimodular Lie algebras  $\mathfrak{r}_{3,\lambda}(\mathbb{C})$  for  $\lambda \in \mathbb{C} \setminus \{-1\}$ . In this section we want to show an analogous result for  $\mathfrak{n} =$  $\mathfrak{sl}_2(\mathbb{C}) \times \mathfrak{sl}_2(\mathbb{C})$ . Here we will use RB-operators on **n** and an explicit classification by Douglas

and Repka [\[18\]](#page-16-25) of all subalgebras of n. This classification is up to inner automorphisms, but we will only need the subalgebras up to isomorphisms. Let us fix a basis  $(X_1, Y_1, H_1, X_2, Y_2, H_2)$ of **n** consisting of the following  $4 \times 4$  matrices.

 $X_1 = E_{12}$ ,  $Y_1 = E_{21}$ ,  $H_1 = E_{11} - E_{22}$ ,  $X_2 = E_{34}$ ,  $Y_2 = E_{43}$ ,  $H_2 = E_{33} - E_{44}$ . We use the following table.

Table 1: Complex 3-dimensional Lie algebras

	Lie brackets
$\mathfrak{n}_3(\mathbb{C})$	$ e_1, e_2  = e_3$
$\mathfrak{r}_2(\mathbb{C})\oplus\mathbb{C}$	$ e_1, e_2  = e_2$
$\mathfrak{r}_3(\mathbb{C})$	$[e_1, e_2] = e_2, [e_1, e_3] = e_2 + e_3$
$\mathfrak{r}_{3,\lambda}(\mathbb{C}), \lambda \neq 0$	$[e_1, e_2] = e_2, [e_1, e_3] = \lambda e_3$
$\mathfrak{sl}_2(\mathbb{C})$	$[e_1, e_2] = e_3, [e_1, e_3] = -2e_1, [e_2, e_3] = 2e_2$

Among the family  $\mathfrak{r}_{3,\lambda}(\mathbb{C}), \lambda \neq 0$  there are still isomorphisms. In fact,  $\mathfrak{r}_{3,\lambda}(\mathbb{C}) \cong \mathfrak{r}_{3,\mu}(\mathbb{C})$  if and only if  $\mu = \lambda^{-1}$  or  $\mu = \lambda$ . The list of subalgebras h of n is given as follows. We first list the solvable subalgebras, then the semisimple ones and the subalgebras with a non-trivial Levi decomposition.

Table 2: Solvable subalgebras

$\dim(\mathfrak{h})$	Representative	Isomorphism type
	$X_1$ , $\langle H_1 \rangle$ , $\langle X_1 + X_2 \rangle$ , $\langle X_1 + H_2 \rangle$ , $\langle H_1 + aH_2 \rangle$ , $a \in \mathbb{C}^*$	
	$\langle X_1, X_2 \rangle$ , $\langle X_1, H_2 \rangle$ , $\langle H_1, H_2 \rangle$	
	$\langle X_1 + X_2, H_1 + H_2 \rangle, \, \langle X_1, H_1 + X_2 \rangle, \, \langle X_1, H_1 + aH_2 \rangle, \, a \in \mathbb{C}$	$\mathfrak{r}_2(\mathbb{C})$
	$\langle X_1, X_2, H_1 + \lambda H_2 \rangle, \lambda \neq 0$	$\mathfrak{r}_{3,\lambda}(\mathbb{C}), \lambda \neq 0$
	$\langle X_1, H_1, H_2 \rangle, \, \langle X_1, H_1, X_2 \rangle$	$\oplus \mathbb{C}$ $\mathfrak{r}_2(\mathbb{C})$
	$\langle X_1, H_1, X_2, H_2 \rangle$	$\mathfrak{r}_2(\mathbb{C})\oplus\mathfrak{r}_2(\mathbb{C})$

Table 3: Semisimple subalgebras and Levi decomposable subalgebras

$\dim(\mathfrak{h})$	Representative	Isomorphism type
	$\langle X_1, Y_1, H_1 \rangle, \, \langle X_1 + X_2, Y_1 + Y_2, H_1 + H_2 \rangle$	$\mathfrak{sl}_2(\mathbb{C})$
	$\langle X_1, Y_1, H_1, H_2 \rangle, \, \langle X_1, Y_1, H_1, X_2 \rangle$	$A \oplus C$ $\mathfrak{sl}_2(\mathbb{C})$
	$\langle X_1, Y_1, H_1, X_2, H_2 \rangle$	$\mathfrak{sl}_2(\mathbb{C})\oplus\mathfrak{r}_2(\mathbb{C})$

**Theorem 4.1.** Suppose that there exists a post-Lie algebra structure on  $(\mathfrak{g}, \mathfrak{n})$ , where  $\mathfrak{n} =$ sl2(C) ⊕ sl2(C)*. Then* g *is isomorphic to one of the following Lie algebras, and all these possibilities do occur:*

(1) 
$$
\mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{sl}_2(\mathbb{C})
$$
.  
\n(2)  $\mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{sl}_2(\mathbb{C})$ .  
\n(3)  $\mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathfrak{r}_{3,\mu}(\mathbb{C}), \lambda \neq -1$ .  
\n(4)  $\mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_{2,\mu}(\mathbb{C}) \oplus \mathfrak{r}_{2,\mu}(\mathbb{C})$ .  
\n(5)  $\mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})$ .  
\n(6)  $\mathbb{C} \oplus ((\mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathbb{C}) \times \mathbb{C}) = \langle x_1, \ldots, x_6 \rangle$  and Lie brackets, for  $\alpha \neq 0, \beta \neq 0, -1$ .  
\n $[x_1, x_2] = x_1, [x_3, x_6] = x_3, [x_4, x_6] = \alpha x_4, [x_5, x_6] = \beta x_5$ .  
\n(6)  $\mathbb{C} \oplus ((\mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathbb{C}) \times \mathbb{C}) = \langle x_1, \ldots, x_6 \rangle$  and Lie brackets, for  $\lambda \neq 0, \alpha \neq 0, -1$ ,  
\n $[x_2, x_4] = x_2, [x_3, x_4] = \lambda x_3, [x_3, x_6] = x_3, [x_5, x_6] = \alpha x_5$ .  
\n(7)  $(\mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathbb{C}^2) \times \mathbb{C} = \langle x_1, \ldots, x_6 \rangle$  and Lie brackets, for  $\lambda \neq 0, \alpha_1, \alpha_2 \neq 0$ , and  
\n $(\lambda, \alpha_1, \alpha_2) \neq (-1, \alpha_1, -\alpha_1 - 1)$ ,  
\n $[x_1, x_3] = x_1, [x_2,$ 

$$
\alpha_3 - \alpha_1 \alpha_2 \neq 0, \, \alpha_1 \neq 0, 1.
$$

*Proof.* By Corollary 2.[15](#page-3-3) it is enough to consider the RB-operators R of weight 1 on n. Then  $\ker(R)$  and  $\ker(R + id)$  are ideals in g. If R is trivial, or one of the kernels is trivial, then we have  $\mathfrak{g} \cong \mathfrak{n}$ , which is type (1). So we assume that R is non-trivial, both ker(R) and ker(R+id) are non-zero, and  $\dim(\ker(R)) \ge \dim(\ker(R + id))$ . Then, for  $\mathfrak{n} \not\cong \mathfrak{g}$ , either  $\mathfrak{g}$  has a non-trivial Levi decomposition, or g is solvable.

*Case 1:* Assume that g has a non-trivial Levi decomposition, i.e., that  $g \cong \mathfrak{sl}_2(\mathbb{C}) \ltimes \mathfrak{r}$ . We claim that  $\mathfrak{sl}_2(\mathbb{C})$  is a direct summand of  $\mathfrak{g}$ , i.e.,  $\mathfrak{g} \cong \mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{r}$ , and that  $\mathfrak{r}$  is not isomorphic to  $\mathfrak{r}_3(\mathbb{C})$ . Then we can argue as follows. Because of Remark 2.12 of [\[12\]](#page-16-22), g cannot be unimodular, except for  $\mathfrak{g} \cong \mathfrak{n}$ . Thus  $\mathfrak{r}$  cannot be unimodular, so that  $\mathfrak{g}$  is isomorphic to  $\mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{r}_{3,\lambda}(\mathbb{C})$  with  $\lambda \neq -1$ . On the other hand, all such algebras do arise by Proposition 2.[6](#page-2-2) and Proposition 4.7 of [\[11\]](#page-16-17).

*Case 1a:* Suppose that  $\mathfrak{sl}_2(\mathbb{C})$  is not contained in ker $(R)$ , ker $(R + id)$  as a subalgebra. Then  $\dim(\ker(R + id)) = 1$  and  $\dim(\ker(R)) \in \{1,2\}$ . Let us assume, both have dimension 1. The other case goes similarly. Then we have  $\mathfrak{r} = \langle x_1, x_2, x_3 \rangle$ , ker(R) =  $\langle x_1 \rangle$  and ker(R + id) =  $\langle x_2 \rangle$ . Furthermore  $\text{im}(R) \cong \mathfrak{sl}_2(\mathbb{C}) \ltimes \langle x_2, x_3 \rangle$  and  $\text{im}(R + \text{id}) \cong \mathfrak{sl}_2(\mathbb{C}) \ltimes \langle x_1, x_3 \rangle$  are 5-dimensional subalgebras of **n**. By table 3,  $\mathfrak{sl}_2(\mathbb{C})$  is a direct summand of them. This implies that  $\mathfrak{sl}_2(\mathbb{C})$  is also a direct summand in g. Since both ker(R) and ker(R + id) are ideals in  $\mathfrak{r}$ , we can exclude that **r** is isomorphic to  $\mathfrak{r}_3(\mathbb{C})$ , and we are done.

*Case 1b:*  $\mathfrak{sl}_2(\mathbb{C})$  is contained in one of ker(R), ker(R + id). Without loss of generality we may assume that  $\mathfrak{sl}_2(\mathbb{C}) \subseteq \ker(R)$ . If  $\ker(R) = \mathfrak{sl}_2(\mathbb{C})$ , then  $\mathfrak{sl}_2(\mathbb{C})$  is an ideal of  $\mathfrak{g}$ , and we have  $\mathfrak{g} \cong \mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{r}$ , where  $\mathfrak{r} \cong \text{im}(R) \leq \mathfrak{n}$  is not isomorphic to  $\mathfrak{r}_3(\mathbb{C})$  by table 2, and we are done. Thus we may assume that  $\dim(\ker(R)) \geq 4$ . If R splits with subalgebras  $\ker(R)$  and  $\ker(R+i d)$ ,

### 12 D. BURDE AND V. GUBAREV

then  $\mathfrak{g} \cong \text{ker}(R) \oplus \text{ker}(R + id)$ , and  $\dim(\text{ker}(R)) + \dim(\text{ker}(R + id)) = 6$ . By table 3,  $\mathfrak{sl}_2(\mathbb{C})$ is a direct summand of ker(R), and hence of  $\mathfrak{g}$ . So we have again  $\mathfrak{g} \cong \mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{r}$ , and  $\mathfrak{r}$  is not isomorphic to  $\mathfrak{r}_3(\mathbb{C})$ . If R is not split, it remains to consider the case dim(ker(R)) = 4 and  $\dim(\ker(R + id)) = 1$ . We have  $\mathfrak{r} = \langle x, y, z \rangle$  with  $\ker(R) = \mathfrak{sl}_2(\mathbb{C}) \oplus \langle x \rangle$ ,  $\ker(R + id) = \langle y \rangle$ and  $[y, \mathfrak{sl}_2(\mathbb{C})] = 0$ . Assume that  $[z, \mathfrak{sl}_2(\mathbb{C})] \neq 0$ . Then  $\mathfrak{sl}_2(\mathbb{C})$  is not a direct summand of the 5-dimensional subalgebra im $(R + id)$  of **n**, which is a contradiction to table 3. Thus we have  $\mathfrak{g} \cong \mathfrak{sl}_2(\mathbb{C}) \oplus \mathfrak{r}$ . Since  $\mathfrak{r}$  has two disjoint 1-dimensional ideals  $\langle x \rangle$  and  $\langle y \rangle$ , it is not isomorphic to  $\mathfrak{r}_3(\mathbb{C})$ .

*Case 2:* Assume that  $\mathfrak g$  is solvable. Then  $\text{im}(R)$  and  $\text{im}(R + id)$  are solvable subalgebras of  $\mathfrak n$ of dimension at most 4 by table 2. So we have  $\dim(\ker(R)) \geq \dim(\ker(R + id)) \geq 2$ . Thus we have the following four cases:

- $(2a)$  dim(ker $(R)$ ) = 4, dim(ker $(R + id)$ ) = 2, (2b) dim(ker(R)) = 3, dim(ker(R + id)) = 3,  $(2c)$  dim(ker(R)) = 3, dim(ker(R + id)) = 2,
- (2d)  $\dim(\ker(R)) = 2$ ,  $\dim(\ker(R + id)) = 2$ .

For the cases (2a) and (2b), R is split since the dimensions add up to 6. Then  $\mathfrak g$  is a direct sum of two solvable subalgebras, which are both isomorphic to subalgebras of  $\mathfrak n$ . So we have  $\mathfrak{n} = \ker(R) + \ker(R + id)$  and  $\mathfrak{g} = \ker(R) \oplus \ker(R + id)$ .

*Case 2a:* Since we have only  $\mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})$  as 4-dimensional solvable subalgebra of **n**, we have  $\mathfrak{g} \cong \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C}) \oplus \mathbb{C}^2$ , which is of type (3) for  $(\lambda, \mu) = (0, 0)$ , or  $\mathfrak{g} \cong \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})$ , which is of type  $(4)$ . Both cases can arise. For the first one we will show this in case  $(2b)$ . For the second, it follows from Proposition 2.[7](#page-2-1) with  $\mathfrak{n} = \langle X_1, H_1, X_2, H_2 \rangle \dot{+} \langle Y_1, Y_2 + H_1 \rangle$ .

*Case 2b:* We have  $\mathfrak{g} \cong \mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathfrak{r}_{3,\mu}(\mathbb{C})$ . The case  $(\lambda, \mu) = (-1, -1)$  cannot arise by Theorem 3.3 of [\[11\]](#page-16-17). The cases  $(\lambda, \mu) = (-1, \mu)$  for  $\mu \neq -1$  arise by Proposition 2.[7](#page-2-1) with

$$
\mathfrak{n} = \langle X_1, X_2, H_1 - H_2 \rangle \dot{+} \langle Y_1, Y_2, H_1 + \mu H_2 \rangle.
$$

The other cases with  $\lambda, \mu \neq -1$  arise by Proposition [2](#page-2-2).6 and Proposition 4.7 of [\[11\]](#page-16-17).

*Case 2c:* Here **g** is isomorphic to  $(\mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})) \rtimes \mathbb{C}$  or  $(\mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathbb{C}^2) \rtimes \mathbb{C}$ . In the first case,  $\mathfrak{r}_2(\mathbb{C}) \rtimes \mathbb{C} \cong \text{im}(R)$  is a solvable subalgebra of **n**, hence isomorphic to  $\mathfrak{r}_{3,\nu}(\mathbb{C})$  by table 2. So  $\mathbb{C}$  acts trivially on  $\mathfrak{r}_2(\mathbb{C})$ , and  $\text{im}(R+\text{id}) \cong \mathfrak{r}_{3,\lambda}(\mathbb{C}) \rtimes \mathbb{C} \cong \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})$ . Then  $\mathfrak{g} \cong \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})$ , which we have already considered in Case (2a). For  $(\mathfrak{r}_{3,\lambda}(\mathbb{C}) \oplus \mathbb{C}^2) \rtimes \mathbb{C}$  we need to distinguish  $\lambda = 0$  and  $\lambda \neq 0$ .

*Case [2](#page-1-0)c,*  $\lambda = 0$ : By Proposition 2.3 we may assume that  $\text{im}(R + \text{id}) = \langle X_1, H_1, X_2, H_2 \rangle$ . Since  $\ker(R)$  is an ideal of im(R+id) isomorphic to  $\mathfrak{r}_2(\mathbb{C}) \oplus \mathbb{C}$ , we have  $\ker(R) = \langle X_1, H_1, X_2 \rangle$ . Let us consider the characteristic polynomial  $\chi_R$  of the linear operator R acting on n. By assumption on the kernels,  $\chi_R(t) = t^3(t+1)^2(t-\rho)$ .

*Case 2c,*  $\lambda = 0$ ,  $\rho \neq 0, -1$ : Then  $R(x_6) = \rho x_6$  for  $x_6 = H_2 + \alpha H_1 + \beta X_1 + \gamma X_2$ . Since ker(R+id) is an abelian 2-dimensional subalgebra of n, we have

$$
\ker(R + id) = \langle Y_1 + \nu_1 X_1 + \nu_2 H_1, Y_2 + \nu_3 X_2 + \nu_4 H_2 \rangle.
$$

We want to compute  $[x, y]$  for  $x = x_6$  and  $y \in \text{ker}(R + id)$ . By Proposition 2.[13](#page-3-2) we have, using  $R(x_6) = \rho x_6$ 

$$
[x, y] = \{R(x), y\} - \{R(y), x\} + \{x, y\}
$$
  
=  $\{R(x), y\}$   
=  $\rho\{x, y\}.$ 

For  $x_6 = H_2 + \alpha H_1 + \beta X_1 + \gamma X_2$  and  $y \in \text{ker}(R + \text{id})$  this yields, using the Lie brackets of n in the standard basis  $\{X_1, Y_1, H_1, X_2, Y_2, H_2\},\$ 

<span id="page-12-0"></span>(10) 
$$
[x_6, Y_1 + \nu_1 X_1 + \nu_2 H_1] = \rho((2\alpha\nu_1 - 2\beta\nu_2)X_1 - 2\alpha Y_1 + \beta H_1),
$$

<span id="page-12-1"></span>(11) 
$$
[x_6, Y_2 + \nu_3 X_2 + \nu_4 H_2] = \rho((2\nu_3 - 2\gamma\nu_4)X_2 - 2Y_2 + \gamma H_2).
$$

Since ker( $R + id$ ) is an ideal in g and  $\rho \neq 0$ , both vectors lie again in ker( $R + id$ ). Comparing coefficients for the basis vectors we obtain

$$
\beta = -2\alpha\nu_2, \ \alpha(\nu_1 + \nu_2^2) = 0, \ \gamma = -2\nu_4, \ \nu_3 = -\nu_4^2.
$$

Suppose that  $\alpha = 0$ . Then  $x_6 = H_2 - 2\nu_4 X_2$  and  $\langle X_1, H_1 \rangle \cong \mathfrak{r}_2(\mathbb{C})$  is a direct summand of  $\mathfrak{g}$ . Therefore  $\mathfrak{g} \cong \mathfrak{r}_2(\mathbb{C}) \oplus \mathbb{C} \oplus \mathfrak{r}_{3,\mu}(\mathbb{C})$  with  $\mathbb{C} = \langle Y_1+\nu_1X_1+\nu_2H_1 \rangle$ ,  $\mathfrak{r}_{3,\mu}(\mathbb{C}) = \langle X_2, H_2-2\nu_4X_2, Y_2+$  $\nu_4 H_2 - \nu_4^2 X_2$ ,  $\mu = -(\rho + 1)/\rho$ , which we have already considered above. Hence we may assume that  $\alpha \neq 0$  and  $\nu_1 = -\nu_2^2$ . Consider a new basis for **g** (note that we redefine  $x_6$ ) given by

$$
(x_1, \ldots, x_6) = (X_1, -\frac{1}{2}H_1 + \nu_2 X_1, X_2, Y_1 + \nu_2 H_1 - \nu_2^2 X_1, Y_2 + \nu_4 H_2 - \nu_4^2 X_2,
$$
  

$$
\frac{1}{2\rho}(H_2 + \alpha H_1 - 2\alpha \nu_2 X_1 - 2\nu_4 X_2)),
$$

with Lie brackets

$$
[x_1, x_2] = x_1, [x_1, x_6] = -\frac{\rho+1}{\rho} \alpha x_1, [x_3, x_6] = -\frac{\rho+1}{\rho} x_3, [x_4, x_6] = \alpha x_4, [x_5, x_6] = x_6.
$$

This algebra is of type (5), if we replace  $x_6$  by  $x_6 + \frac{\alpha(\rho+1)}{a}$  $\frac{p+1}{p}x_2$ . It arises for the triangular-split RB-operator R with  $A_-=\text{ker}(R)=\langle x_1, x_2, x_3\rangle$ ,  $\text{ker}(R'+\text{id}) = \langle x_4, x_5\rangle$  and  $A_0=\langle x_6\rangle$ , where  $x_6 = H_2 - 2\nu_4 X_2$ , with the action  $R(x_6) = \rho x_6$ .

*Case 2c,*  $\lambda = 0$ ,  $\rho = -1$ : We may assume that there exists  $x_6 = Y_2 + v$  such that  $(R + id)(x_6) =$  $\mu(H_2 + \alpha H_1 + \beta X_1 + \gamma X_2)$  for some non-zero  $\mu$  and some  $\alpha, \beta, \gamma \in \mathbb{C}$ . Since ker $(R + id)$  is an abelian subalgebra we obtain  $\alpha = \beta = 0$  and ker $(R + id) = \langle H_2 + \gamma X_2, Y_1 + \nu_1 X_1 + \nu_2 H_1 \rangle$ . Then we may choose  $x_6 = Y_2 + \kappa X_2 + \nu_3 H_1 + \nu_4 X_1$ . Then

$$
[x_6, H_2 + \gamma X_2] = \{R(x_6), H_2 + \gamma X_2\}
$$
  
=  $\{(R + id)(x_6) - x_6, H_2 + \gamma X_2\}$   
=  $-\{Y_2 + \kappa X_2, H_2 + \gamma X_2\}$   
=  $-2Y_2 + 2\kappa X_2 + \gamma H_2$ .

This is not contained in ker( $R + id$ ), which is a contradiction to the fact that ker( $R + id$ ) is an ideal.

*Case 2c,*  $\lambda = 0$ ,  $\rho = 0$ : Then we have  $R(H_2) = \alpha H_1 + \beta X_1 + \gamma X_2 \neq 0$  and ker $(R + id)$  $\{Y_1+\nu_1X_1+\nu_2H_1, Y_2+\nu_3X_2+\nu_4H_2\}$ . Since  $[H_2, Y_2+\nu_1X_1+\nu_2H_1] = \{\gamma X_2, Y_2+\nu_1X_2+\nu_2H_2\}$  is in ker(R+id), we obtain  $\gamma = 0$ . Since  $[H_2, Y_1 + \nu_1 X_1 + \nu_2 H_2] = {\alpha H_1 + \beta X_1, Y_1 + \nu_1 X_1 + \nu_2 H_1}$ 

is in ker( $R + id$ ), we obtain  $\alpha(\nu_1 + \nu_2^2) = 0$  and  $\beta = -2\alpha\nu_2$ . Since  $R(H_2) \neq 0$  we have  $\alpha \neq 0$ ,  $\nu_1 = -\nu_2^2$  and  $R(H_2) = \alpha H_1 - 2\alpha \nu_2 X_1$ . Consider a new basis for **g** given by

$$
(x_1, \ldots, x_6) = (X_1, -\frac{1}{2}H_1 + \nu_2 X_1, X_2, Y_1 + \nu_2 H_1 - \nu_2^2 X_1, Y_2 + \nu_3 X_2 + \nu_4 H_2, -\frac{1}{2}H_2),
$$

with Lie brackets

$$
[x_1, x_2] = x_1, [x_1, x_6] = \alpha x_1, [x_3, x_6] = x_3, [x_4, x_6] = -\alpha x_4.
$$

This algebra is of type (3), if we replace  $x_6$  by  $x_6 - \alpha x_2$ .

*Case 2c,*  $\lambda \neq 0$ : Then we have ker(R) =  $\langle X_1, X_2, -\frac{1}{2} \rangle$  $\frac{1}{2}(H_1 + \lambda H_2)$ . We again have  $\chi_R(t) =$  $t^3(t+1)^2(t-\rho)$ , where we distinguish the cases  $\rho \neq 0, -1, \rho = -1$  and  $\rho = 0$ .

*Case 2c,*  $\lambda \neq 0$ ,  $\rho \neq 0$ , -1*:* Then we may assume that  $R(x_6) = \rho x_6$  for  $x_6 = H_2 + \alpha H_1 + \beta X_1 + \beta X_2$  $\gamma X_2$ . As ker(R + id) is abelian, we have ker(R + id) =  $\langle Y_1 + \nu_1 X_1 + \nu_2 H_1, Y_2 + \nu_3 X_2 + \nu_4 H_2 \rangle$ . Since  $V = \text{ker}(R) \oplus \text{ker}(R + \text{id}) \oplus \langle x_6 \rangle$ , the two elements  $H_1 + \lambda H_2$  and  $H_2 + \alpha H_1$  need to be linearly independent, i.e.,  $1 - \alpha \lambda \neq 0$ . By [\(10\)](#page-12-0) and [\(11\)](#page-12-1) we obtain  $\gamma = -2\nu_4$ ,  $\beta = -2\alpha\nu_2$ ,  $\nu_3 = -\nu_4^2$  and  $\alpha(\nu_1 + \nu_2^2)$ . Suppose that  $\alpha = 0$ . Then  $x_6 = H_2 - 2\nu_4 X_2$ . Consider a new basis for g given by

$$
(x_1,\ldots,x_6)=(Y_1+\nu_1X_1+\nu_2H_1,X_1,X_2,-\frac{1}{2}(H_1+\lambda H_2),Y_2-\nu_4^2X_2+\nu_4H_2,-\frac{1}{2(\rho+1)}H_2),
$$

with Lie brackets

$$
[x_2, x_4] = x_2, [x_3, x_4] = \lambda x_3, [x_3, x_6] = x_3, [x_4, x_6] = -\lambda \nu_4 x_3, [x_5, x_6] = -\frac{\rho}{1+\rho} x_5.
$$

This is an algebra of type (6), if we replace  $x_4$  by  $x_4 + \lambda \nu_4 x_3$ . Now we assume that  $\alpha \neq 0$ . Consider a new basis for **g** given by

$$
(x_1, \ldots, x_6) = (X_1, X_2, -\frac{1}{2}(H_1 + \lambda H_2), Y_2 - \nu_4^2 X_2 + \nu_4 H_2, Y_1 - \nu_2^2 X_1 + \nu_2 H_1,
$$
  

$$
-\frac{1}{2(\rho + 1)}(H_2 - 2\nu_4 X_2 + \alpha (H_1 - 2\nu_2 X_1))),
$$

with Lie brackets

$$
[x_1, x_3] = x_1, [x_2, x_3] = \lambda x_2, [x_1, x_6] = \alpha x_1, [x_2, x_6] = x_2,
$$
  

$$
[x_3, x_6] = -\alpha \nu_2 x_1 - \lambda \nu_4 x_2, [x_4, x_6] = \delta x_4, [x_5, x_6] = \alpha \delta x_5,
$$

where  $\delta = -\frac{\rho}{\rho+1}$ . Replacing  $x_6$  by  $\frac{1}{\delta}(x_6 - \alpha \nu_2 x_1 - \nu_4 x_2 - \alpha x_3)$  we obtain the Lie brackets

$$
[x_1, x_3] = x_1, [x_2, x_3] = \lambda x_2, [x_2, x_6] = \alpha' x_2, [x_4, x_6] = x_4, [x_5, x_6] = \alpha x_5,
$$

where

$$
\alpha' = \frac{1 - \alpha \lambda}{\delta} = \frac{(\rho + 1)(\alpha \lambda - 1)}{\rho}.
$$

Note that  $\alpha' \neq 0$  and  $\alpha' \neq \alpha \lambda - 1$  by assumption. In other words,  $\alpha \neq \frac{\alpha'+1}{\lambda}$  $\frac{+1}{\lambda}$ . Consider a new basis for g given by

$$
(x'_1, \ldots, x'_6) = (x_2, x_1, \frac{1}{\lambda}x_3, x_4, x_5, x_6 - \frac{\alpha'}{\lambda}x_3),
$$

with Lie brackets

$$
[x'_1, x'_3] = x'_1, [x'_2, x'_3] = \lambda' x'_2, [x'_2, x'_6] = -\alpha' \lambda' x'_2, [x'_4, x'_6] = x'_4, [x'_5, x'_6] = \alpha x'_5,
$$

where  $\lambda' = \frac{1}{\lambda}$ <sup>1</sup>/<sub>λ</sub>. This is of type (7). Since  $\mathfrak{r}_{3,\lambda}(\mathbb{C}) \cong \mathfrak{r}_{3,\lambda'}(\mathbb{C})$ , one may check that we do not only have  $\alpha \neq \frac{\alpha'+1}{\lambda}$  $\frac{\hat{+1}}{\lambda}$ , but also  $\alpha \neq \lambda - \alpha'$ . For  $\frac{\alpha'+1}{\lambda}$  $\frac{+1}{\lambda} \neq \lambda - \alpha'$  we obtain no restriction for  $\alpha$ . However, for  $\frac{\alpha'+1}{\lambda} = \lambda - \alpha'$  we obtain  $\lambda = -1$  or  $\lambda = \alpha'+1$ , which excludes both  $(\lambda, \alpha', \alpha) = (-1, \alpha', -\alpha'-1)$ and  $(\lambda, \alpha', \alpha) = (\lambda, \lambda - 1, 1)$ . Rewriting this in the parameters of the Lie brackets from type (7), we obtain all cases except for  $(\lambda, \alpha', \alpha) = (\lambda, \lambda - 1, 1)$  with  $\lambda \neq -1$ . These PA-structures arise by a triangular-split RB-operator with  $A_-=\text{ker}(R)$ ,  $A_+=\text{ker}(R+\text{id})$  and  $A_0=\langle x_6\rangle$ with the action  $R(x_6) = \rho x_6, \ \rho \neq 0, -1.$ 

*Case 2c,*  $\lambda \neq 0, \rho = -1$ *:* This leads to a contradiction in the same way as case 2c with  $\lambda = 0, \rho = -1.$ 

*Case 2c,*  $\lambda \neq 0, \rho = 0$ *:* We have  $R(H_2) = \alpha X_1 + \beta X_2 + \gamma (H_1 + \lambda H_2)$  and ker $(R + id) = \langle x_4, x_5 \rangle$ with  $x_4 = Y_1 + \nu_1 X_1 + \nu_2 H_1$ ,  $x_5 = Y_2 + \nu_3 X_2 + \nu_4 H_2$ . Similarly to [\(10\)](#page-12-0), [\(11\)](#page-12-1) we obtain  $R(H_2) = \gamma(H_1 - 2\nu_2 X_1) + \gamma \lambda(H_2 - 2\nu_4 X_2)$ . This implies that  $\gamma \neq 0$  and  $x_4 = Y_1 - \nu_2^2 X_1 + \nu_2 H_1$ ,  $x_5 = Y_2 - \nu_4^2 X_2 + \nu_4 H_2$ . By setting  $x_1 = X_1$ ,  $x_2 = X_2$ ,  $x_3 = -\frac{1}{2}$  $\frac{1}{2}(H_1 + \lambda H_2)$  and  $x_6 = \frac{1}{24}$  $\frac{1}{2\gamma}H_2$  we obtain a new basis for g with Lie brackets

$$
[x_1, x_3] = x_1, [x_2, x_3] = \lambda x_2, [x_1, x_6] = -x_1, [x_2, x_6] = \delta x_2,
$$
  

$$
[x_3, x_6] = \nu_2 x_1 + \lambda^2 \nu_4 x_2, [x_4, x_6] = x_4, [x_5, x_6] = \lambda x_5,
$$

where  $\delta = -\frac{1+\lambda\gamma}{\gamma}$  $\frac{\partial^2 \phi}{\partial x^2}$  with  $\delta \neq -\lambda$ . Replacing  $x_6$  by  $x_6 + \nu_2 x_1 + \lambda \nu_4 x_2 + x_3$  we obtain the brackets

$$
[x_1, x_3] = x_1, [x_2, x_3] = \lambda x_2, [x_2, x_6] = \alpha_1 x_2, [x_4, x_6] = x_4, [x_5, x_6] = \lambda x_5
$$

with  $\alpha_1 = \delta + \lambda = -\frac{1}{\gamma}$  $\frac{1}{\gamma}$ . This is of type (7) with  $\alpha_2 = \lambda$ . It arises by the triangular-split RB-operator with  $A_ - = \langle x_1, x_2 \rangle$ ,  $A_+ = \langle x_4, x_5 \rangle$  and  $A_0 = \langle u, v \rangle$ , with  $u = \frac{1}{\gamma}$  $\frac{1}{\gamma}(H_2 - 2\nu_4 X_2)$  and  $v = H_1 - 2\nu_2X_1 + \lambda(H_2 - 2\nu_4X_2)$ , and the action  $R(u) = v$ ,  $R(v) = 0$ .

*Case 2d:* Suppose that one of the kernels  $ker(R)$  and  $ker(R+id)$  is non-abelian. Without loss of generality, let us assume that ker(R)  $\cong$  r<sub>2</sub>( $\mathbb{C}$ ). Write  $\mathfrak{g} \cong$  (ker(R)  $\oplus$  ker(R+id))  $\ltimes \langle a, b \rangle$ . Then  $\ker(R) \ltimes \langle a \rangle$  is a 3-dimensional solvable subalgebra of im( $R + id$ ). By table 2 we see that it is isomorphic to  $\mathfrak{r}_2(\mathbb{C}) \oplus \mathbb{C}$ . In this case there exist nonzero  $a' \in \ker(R) \oplus \langle a \rangle$  and  $b' \in \ker(R) \oplus \langle b \rangle$ such that  $[a',\ker(R)] = [b',\ker(R)] = 0$ . Then  $\mathfrak{g} \cong \ker(R) \oplus (\ker(R + id) \oplus \langle a',b' \rangle)$  with  $\ker(R) \cong \mathfrak{r}_2(\mathbb{C})$ , and  $\ker(R + id) \oplus \langle a', b' \rangle \cong \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})$  by Table 2. Hence we obtain  $\mathfrak{g} \cong \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C}) \oplus \mathfrak{r}_2(\mathbb{C})$ , which is of type (4).

So we may assume that  $\ker(R) \cong \ker(\tilde{R} + id) \cong \mathbb{C}^2$ . Then the characteristic polynomial of R has the form  $\chi_R(t) = t^2(t+1)^2(t-\rho_1)(t-\rho_2)$ .

*Case 2d,*  $\rho_1, \rho_2 \neq 0, -1$ *:* Suppose first that either  $\rho_1 \neq \rho_2$ , or that  $\rho_1 = \rho_2$  and the eigenspace is 2-dimensional. Then by Proposition [3](#page-6-0).4,  $\mathfrak{n} = \ker(R) + \ker(R + id) + \langle x_5', x_6' \rangle$  with linearly independent eigenvectors  $x'_5, x'_6$  corresponding to the eigenvalues  $\rho_1$  and  $\rho_2$ . Since ker(R) is an abelian ideal in  $\text{im}(R + \text{id}) = \langle X_1, H_1, X_2, H_2 \rangle$ , we may assume that  $\text{ker}(R) = \langle X_1, X_2 \rangle$ and  $[x'_5, x'_6] = 0$ . The decomposition  $\mathfrak{n} = \ker(R + id) + \text{im}(R + id)$  shows that  $\ker(R + id)$ has a basis  $x_3 = Y_1 + \alpha H_1 + \nu_3 X_1$ ,  $x_4 = Y_2 + \beta H_2 + \nu_4 X_2$ . Since  $[x'_5, x'_6] = 0$ , we have  $x'_5 = H_1 + \nu_1 X_1 + \xi_1 (H_2 + \nu_2 X_2), x'_6 = H_2 + \nu_2 X_2 + \xi_2 (H_1 + \nu_1 X_1)$  with  $\xi_1 \xi_2 \neq 1$ . So we have by [\(10\)](#page-12-0) and [\(11\)](#page-12-1)  $x_3 = Y_1 - \frac{\nu_1}{2}H_1 - \frac{\nu_1^2}{4}X_1$ ,  $x_4 = Y_2 - \frac{\nu_2}{2}H_2 - \frac{\nu_2^2}{4}X_2$ . Consider a basis for **g** given by

$$
(x_1, \ldots, x_6) = (X_1, X_2, x_3, x_4, -\frac{1}{2(1+\rho_1)}x'_5, -\frac{1}{2(1+\rho_2)}x'_6),
$$

with Lie brackets

$$
[x_1, x_5] = x_1, [x_1, x_6] = \xi_2 x_1, [x_2, x_5] = \xi_1 x_2, [x_2, x_6] = x_2,
$$
  

$$
[x_3, x_5] = \gamma x_3, [x_3, x_6] = \delta \xi_2 x_3, [x_4, x_5] = \gamma \xi_1 x_4, [x_4, x_6] = \delta x_4,
$$

where  $\gamma = -\frac{\rho_1}{\rho_1+1}$ ,  $\delta = -\frac{\rho_2}{\rho_2+1}$  with  $\gamma, \delta \neq 0, -1$  and  $\xi_1\xi_2 \neq 1$ . This is type (8*a*). It arises by the triangular-split RB-operator R with  $A_ = \langle X_1, X_2 \rangle$ ,  $A_+ = \langle x_3, x_4 \rangle$  and  $A_0 = \langle x_5, x_6 \rangle$ , where R acts on  $A_0$  by  $R(x_5) = \rho_1 x_5$  and  $R(x_6) = \rho_2 x_6$ . Note that for  $\nu_2 = \xi_2 = 0$  and  $\xi_1 \neq 0$  we get type (7) without the restriction  $(\lambda, \alpha_1, \alpha_2) \neq (\lambda, \lambda - 1, 1)$  for  $\lambda \neq -1$ , which we had in Case 2c,  $\lambda \neq 0, \, \rho \neq 0, -1.$ 

Suppose now that  $\rho_2 = \rho_1 \neq 0, -1$ , and the eigenspace for  $\rho_1$  is 1-dimensional. Let  $R(x_5') = \rho_1 x_5'$ and  $R(x_6') = x_5' + \rho_1 x_6'$ . In the same way as before we have  $x_5' = H_1 + \nu_1 X_1 + \xi (H_2 + \nu_2 X_2)$ ,  $x'_6 = \kappa (H_2 + \nu_2 X_2)$  with  $\kappa \neq 0$  and  $x_3 = Y_1 - \frac{\nu_1}{2} H_1 - \frac{\nu_1^2}{4} X_1$ ,  $x_4 = Y_2 - \frac{\nu_2}{2} H_2 - \frac{\nu_2^2}{4} X_2$ . Consider a basis for g given by

$$
(x_1,...,x_6) = (X_1, X_2, x_3, x_4, -\frac{1}{2(1+\rho_1)}x'_5, -\frac{1}{2(1+\rho_1)}x'_6),
$$

with Lie brackets

$$
[x_1, x_5] = x_1, [x_1, x_6] = (\gamma + 1)x_1, [x_2, x_5] = \xi x_2, [x_2, x_6] = (\kappa + \xi + \gamma \xi)x_2,
$$
  

$$
[x_3, x_5] = \gamma x_3, [x_3, x_6] = -(\gamma + 1)x_3, [x_4, x_5] = \gamma \xi x_4, [x_4, x_6] = (\kappa \gamma - \xi - \gamma \xi)x_4,
$$

where  $\gamma = -\frac{\rho_1}{\rho_1+1} \neq 0, -1$  and  $\kappa \neq 0$ . This is type (8b). It arises by the triangular-split RB-operator R with  $A_ = \langle X_1, X_2 \rangle$ ,  $A_+ = \langle x_3, x_4 \rangle$  and  $A_0 = \langle x_5, x_6 \rangle$ , where R acts on  $A_0$  by  $R(x_5) = \rho_1 x_5$  and  $R(x_6) = x_5 + \rho_1 x_6$ .

*Case 2d,*  $\rho_1 = \rho_2 = 0$ : We have  $\mathfrak{g} = \text{ker}(R + \text{id}) + \text{im}(R + \text{id})$  and we can assume that  $\ker(R) = \langle X_1, X_2 \rangle$  and  $\ker(R + id) = \langle Y_1 + \nu_1 X_1 + \nu_2 H_1, Y_2 + \nu_3 X_2 + \nu_4 H_2 \rangle$ . Suppose first that  $R(v) = X_1$  and  $R(w) = X_2$  for some v, w. Then

$$
[Y_1 + \nu_1 X_1 + \nu_2 H_1, v] = \{ Y_1 + \nu_1 X_1 + \nu_2 H_1, X_1 \} = -H_1 + 2\nu_2 X_1 \in \text{ker}(R + \text{id}),
$$

which is a contradiction. Otherwise we see from the possible Jordan forms of  $R$  that there exist v, w with  $R(v) = \alpha X_1 + \beta X_2 \neq 0$  and  $R(w) = v$ . This leads to a contradiction in the same way. *Case 2d,*  $\rho_1 = 0, \rho_2 \neq 0, -1$ : This case is analagous to the second part of the case before.

*Case 2d,*  $\rho_1 = 0, \rho_2 = -1$ : As above we may assume that  $\text{im}(R + \text{id}) = \langle X_1, X_2, H_1, H_2 \rangle$  and  $\ker(R) = \langle X_1, X_2 \rangle$ , and  $\alpha H_1 + \beta H_2 + \gamma X_1 + \delta X_2 \in \ker(R+i\epsilon) \cap \text{im}(R+i\epsilon)$  for some  $\alpha, \beta, \gamma, \delta \in \mathbb{C}$ . Since ker(R+id) is abelian, we may assume that ker(R+id) =  $\langle H_1 + \nu_1 X_1, Y_2 + \nu_2 X_2 + \nu_3 H_2 \rangle$ for some  $\nu_1, \nu_2, \nu_3 \in \mathbb{C}$ . Let  $v \in \ker(R^2)$  such that  $R(v) = \nu_4 X_1 + \nu_5 X_2 \neq 0$ . Then

$$
[v, Y_2 + \nu_2 X_2 + \nu_3 H_2] = \{ \nu_4 X_1 + \nu_5 X_2, Y_2 + \nu_2 X_2 + \nu_3 H_2 \} = \nu_5 (H_2 - 2\nu_3 X_2) \in \text{ker}(R + \text{id})
$$

implies that  $\nu_5 = 0$ . By  $[v, H_1 + \nu_1 X_1] = \{ \nu_4 X_1, H_1 + \nu_1 X_1 \} = -2\nu_4 X_1 \in \text{ker}(R + \text{id})$  we obtain  $\nu_4 = 0$ , which is a contradiction to  $R(v) \neq 0$ .

*Remark* 4.2*.* The algebras from different types are non-isomorphic, except for algebras of type (8), which have intersections with type (3) and (7) for certain parameter choices.

#### PA-STRUCTURES 17

#### Acknowledgments

Dietrich Burde is supported by the Austrian Science Foundation FWF, grant P28079 and grant I3248. Vsevolod Gubarev acknowledges support by the Austrian Science Foundation FWF, grant P28079.

### **REFERENCES**

- <span id="page-16-5"></span><span id="page-16-1"></span>[1] F. V. Atkinson: Some aspects of Baxters functional equation. J. Math. Anal. Appl. 7 (1963), 1–30.
- <span id="page-16-0"></span>[2] C. Bai, L. Guo, X. Ni: Nonabelian generalized Lax pairs, the classical Yang-Baxter equation and PostLie algebras. Comm. Math. Phys. 297 (2010), no. 2, 553–596.
- [3] G. Baxter: An analytic problem whose solution follows from a simple algebraic identity. Pacific J. Math. 10 (1960), 731–742.
- <span id="page-16-3"></span>[4] A. A. Belavin, V. G. Drinfel'd: Solutions of the classical Yang-Baxter equation for simple Lie algebras. Funct. Anal. Appl. 16 (1982), no. 3, 159–180.
- <span id="page-16-12"></span><span id="page-16-9"></span>[5] P. Benito, V. Gubarev, A. Pozhidaev: Rota–Baxter operators on quadratic algebras. [arXiv:1801.07037](http://arxiv.org/abs/1801.07037) (2018), 23 pp.
- <span id="page-16-13"></span>[6] D. Burde: Affine structures on nilmanifolds. Internat. J. Math.  $7$  (1996), no. 5, 599–616.
- [7] D. Burde, K. Dekimpe, S. Deschamps: The Auslander conjecture for NIL-affine crystallographic groups. Math. Ann. 332 (2005), no. 1, 161–176.
- <span id="page-16-14"></span>[8] D. Burde: Left-symmetric algebras, or pre-Lie algebras in geometry and physics. Cent. Eur. J. Math. 4 (2006), no. 3, 323–357.
- <span id="page-16-16"></span><span id="page-16-15"></span>[9] D. Burde, K. Dekimpe and S. Deschamps: LR-algebras. Contemp. Math. 491 (2009), 125–140.
- <span id="page-16-17"></span>[10] D. Burde, K. Dekimpe, K. Vercammen: Complete LR-structures on solvable Lie algebras. J. Group Theory 13 (2010), no. 5, 703–719.
- [11] D. Burde, K. Dekimpe and K. Vercammen: Affine actions on Lie groups and post-Lie algebra structures. Linear Algebra Appl. s 437 (2012), no. 5, 1250–1263.
- <span id="page-16-22"></span>[12] D. Burde, K. Dekimpe: Post-Lie algebra structures and generalized derivations of semisimple Lie algebras. Mosc. Math. J. 13 (2013), Issue 1, 1–18.
- <span id="page-16-20"></span><span id="page-16-19"></span>[13] D. Burde, K. Dekimpe: Post-Lie algebra structures on pairs of Lie algebras. J. Algebra 464 (2016), 226–245.
- <span id="page-16-21"></span>[14] D. Burde, W. A. Moens: Commutative post-Lie algebra structures on Lie algebras. J. Algebra 467 (2016), 183–201.
- [15] D. Burde, W. A. Moens, K. Dekimpe: Commutative post-Lie algebra structures and linear equations for nilpotent Lie algebras. [arXiv:1711.01964](http://arxiv.org/abs/1711.01964) (2017), 1–14.
- <span id="page-16-24"></span>[16] D. Burde, V. Gubarev: Decompositions of algebras and post-associative algebra structures. [arXiv:1906.09854](http://arxiv.org/abs/1906.09854) (2019).
- <span id="page-16-25"></span><span id="page-16-2"></span>[17] P. Cartier: On the structure of free Baxter algebras. Advances Math. 9 (1972), 253–265.
- [18] A. Douglas, J. Repka: Subalgebras of the rank two semisimple Lie algebras. Linear Multilinear Algebra 66 (2018), no. 10, 2049–2075.
- <span id="page-16-18"></span>[19] K. Ebrahimi-Fard, A. Lundervold, I. Mencattini, H. Z. Munthe-Kaas: Post-Lie Algebras and Isospectral Flows. SIGMA Symmetry Integrability Geom. Methods Appl. 11 (2015), Paper 093, 16 pp.
- <span id="page-16-10"></span>[20] V. Gubarev: Universal enveloping Lie Rota–Baxter algebra of pre-Lie and post-Lie algebras. [arXiv:1708.06747](http://arxiv.org/abs/1708.06747) (2017), 1–13.
- <span id="page-16-7"></span>[21] V. Gubarev, P. Kolesnikov: Embedding of dendriform algebras into Rota–Baxter algebras. Cent. Eur. J. Math. 11 (2013), no. 2, 226–245.
- <span id="page-16-8"></span><span id="page-16-6"></span>[22] L. Guo, W. Keigher: Baxter algebras and shuffle products. Adv. Math. 150 (2000), 117-149.
- [23] L. Guo: An Introduction to Rota–Baxter Algebra. Surveys of Modern Mathematics, Vol. 4 (2012), Somerville: Intern. Press; Beijing: Higher education press, 226 pp.
- <span id="page-16-23"></span><span id="page-16-11"></span>[24] J. Helmstetter: Radical d'une algèbre symétrique a gauche. Ann. Inst. Fourier 29 (1979), 17–35.
- <span id="page-16-4"></span>[25] N. Jacobson: A note on automorphisms of Lie algebras. Pacific J. Math.  $12$  (1962), no. 1, 303–315.
- [26] X. X. Li, D. P. Hou, C. M. Bai: Rota–Baxter operators on pre-Lie algebras. J. Nonlinear Math. Phys. 14, no. 2 (2007), 269–289.

#### <span id="page-17-2"></span>18 D. BURDE AND V. GUBAREV

- <span id="page-17-6"></span>[27] H. Kim: Complete left-invariant affine structures on nilpotent Lie groups. J. Differential Geom. 24 (1986), no. 3, 373–394.
- <span id="page-17-5"></span>[28] J. L. Koszul: *Variante dun théoréme de H. Ozeki*. Osaka J. Math.  $15$  (1978), 547–551.
- <span id="page-17-7"></span>[29] J.-L. Loday: *Generalized bialgebras and triples of operads*. Astrisque No. **320** (2008), 116 pp.
- <span id="page-17-0"></span>[30] A. L. Onishchik: *Decompositions of reductive Lie groups*. Mat. Sbornik. **80 (122)** (1969), no. 4, 515–554.
- <span id="page-17-8"></span>[31] G.-C. Rota: *Baxter algebras and combinatorial identities I*. Bull. Amer. Math. Soc. **75** (1969), 325–329.
- <span id="page-17-3"></span>[32] R. D. Schafer: A introduction to nonassociative algebras. Dover Publications, New York (1995), 166 pp.
- <span id="page-17-1"></span>[33] D. Segal: The structure of complete left-symmetric algebras. Math. Ann. 293 (1992), 569–578.
- <span id="page-17-4"></span>[34] M. A. Semenov-Tyan-Shanskii: What is a classical R-matrix? Funktsional. Anal. i Prilozhen. 17 (1983), no. 4, 17–33.
- [35] B. Vallette: Homology of generalized partition posets. J. Pure Appl. Algebra 208 (2007), no. 2, 699–725.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, 1090 WIEN, AUS-**TRIA** 

E-mail address: dietrich.burde@univie.ac.at

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT WIEN, OSKAR-MORGENSTERN-PLATZ 1, 1090 WIEN, AUS-**TRIA** 

E-mail address: vsevolod.gubarev@univie.ac.at