

AN APPLICATION OF THE THEORY OF FI-ALGEBRAS TO GRAPH CONFIGURATION SPACES

ERIC RAMOS

ABSTRACT. Recent work of An, Drummond-Cole, and Knudsen [ADK], as well as the author [R], has shown that the homology groups of configuration spaces of graphs can be equipped with the structure of a finitely generated graded module over a polynomial ring. In this work we study this module structure in certain families of graphs using the language of FI-algebras recently explored by Nagel and Römer [NR]. As an application we prove that the syzygies of the modules in these families exhibit a range of stable behaviors.

1. INTRODUCTION

There has been a recent raising interest in studying *configuration spaces of graphs* due to their connections with robotics and topological motion planning [Fa, G]. In this paper, a *graph* is a 1-dimensional, compact simplicial complex. Its *n-stranded configuration space* is the topological space

$$U\mathcal{F}_n(G) := \{(x_1, \dots, x_n) \in G^n \mid x_i \neq x_j\} / \mathfrak{S}_n.$$

In this paper, we explore the homology groups of these configuration spaces from a relatively new perspective, following the work of An, Drummond-Cole, and Knudsen [ADK], as well as the author [R].

For each $q \geq 0$, and each graph G , the *total q-th homology group* is the graded abelian group

$$\mathcal{H}_q(G) := \bigoplus_{n \geq 0} H_q(U\mathcal{F}_n(G))$$

Writing A_G for the integral polynomial ring with variables indexed by the edges of G , the following theorem was proven in the case where G is a tree by the author [R], and in the general case by An, Drummond-Cole, and Knudsen.

Theorem (Ramos, [R]; An, Drummond-Cole, Knudsen, [ADK]). The graded abelian group $\mathcal{H}_q(G)$ can be equipped with the structure of a finitely generated, graded module over A_G .

It therefore becomes natural to ask whether one can use techniques from commutative algebra to deduce facts about these homology groups. In [R], the author showed that the total homology groups could be decomposed into sums of graded shifts of square-free monomial ideals, in the cases wherein G is a tree. Other than that work, however, very little has been done in this direction. In this work, we study the modules $\mathcal{H}_q(G)$ for specific families of graphs G , using recent work of White and the author [RW], and Lütgehetmann [L], as well as Nagel and Römer [NR].

In [RW], White and the author introduced the notion of a *finitely generated FI-graph* (see Definition 3.9). For the purposes of this introduction, we restrict to a specific natural class of examples of this structure. Fix

Key words and phrases. FI-modules, graph configuration spaces.

The first author was supported by NSF grant DMS-1704811.

two graphs, G, H , as well as a choice of vertex v_G, v_H in each. Then for each $n \geq 0$ we may define

$$G_n := G \bigvee^n H.$$

The sequence of graphs $\{G_n\}$ is notable for us, as it carries an action by the category FI of finite sets and injections. Indeed, if we set $[n] := \{1, \dots, n\}$ and let $f : [n] \hookrightarrow [r]$ be an injection, then we obtain an injective homomorphism of graphs (see Definition 2.1)

$$G(f) : G_n \rightarrow G_r$$

which fixes the vertices of G , while permuting the vertices of H according to f . This structure was first exploited in the study of configuration spaces by Lütgehetmann in [L]. In particular, the following is proven.

Theorem (Lütgehetmann, [L]). Let G_n be as above. Then for each $q, m \geq 0$, the functor

$$[n] \mapsto H_q(\mathcal{U}\mathcal{F}_m(G_n))$$

from FI to abelian groups is finitely generated in the sense of [CEF].

Therefore, if we fix the number of points being configured, but allow our graph to grow in a certain way, one obtains a finiteness result. On the other hand, we have also seen that if we fix the graph, and allow the number of points being configured to increase, we obtain a similar finiteness result. The main theorem of this work is that these two types of finiteness are compatible with one another in the appropriate sense.

Theorem A. Let G_n be as above. Then for each $q \geq 0$, the functor

$$[n] \mapsto \mathcal{H}_q(G_n)$$

from FI to graded abelian groups is finitely generated as a graded module over the functor

$$[n] \mapsto A_{G_n}$$

from FI to \mathbb{Z} -algebras, in the sense of [NR] (see also Definition 3.2).

Remark 1.1. As previously stated, the most general theorem involves studying a larger class of families of graphs (see Theorem 4.4). That being said, it is unclear whether the scope this theorem extends to all finitely generated FI-graphs (see Definition 3.9). This difficulty is fundamentally related to the fact that the FI-algebra

$$[n] \mapsto A_{G_n}$$

is not necessarily Noetherian in general. It is an interesting question to ask whether the conclusion of Theorem A will hold for all FI-graphs, even in the case where $G_n = K_n$ is the complete graph on n vertices.

Coming back to the goal of studying the commutative algebra of $\mathcal{H}_q(G)$, we note that Theorem A has concrete consequences related to the Betti numbers of these modules. If M is a finitely generated graded module over the polynomial ring $k[x_1, \dots, x_n]$, where k is a field, then it admits a minimal free resolution

$$0 \rightarrow F^{(n)} \rightarrow \dots \rightarrow F^{(0)} \rightarrow M \rightarrow 0$$

Tensoring with the residue field k , we define the j -th Betti number in homological degree p to be

$$\beta_{p,j}(M) := \dim_k(\mathrm{Tor}_p(M, k)_j).$$

Knowledge of the Betti numbers allows one to access the number of generators in each grade of the p -th syzygies. Our second main result is that the Betti numbers of the modules $\mathcal{H}_q(G_n)$ stabilize in a strong sense.

Theorem B. Let G_n be as above, and let k be a field. Then for each $p, j \geq 0$, the function

$$n \mapsto \beta_{p,j}(\mathcal{H}_q(G_n; k))$$

agrees with a polynomial for $n \gg 0$. Moreover, there exists a finite list of integers

$$j_0(\mathcal{H}_q(G_n; k), p) < \dots < j_t(\mathcal{H}_q(G_n; k), p)$$

such that for all $n \gg 0$

$$\beta_{n,p,j}(\mathcal{H}_q(G_n; k)) \neq 0 \iff j \in \{j_0(\mathcal{H}_q(G_n; k), p), \dots, j_t(\mathcal{H}_q(G_n; k), p)\}.$$

Remark 1.2. It has been known since at least [KP, Theorem 3.16] that the A_G -module $\mathcal{H}_1(G)$ is generated in degree ≤ 2 , for all graphs G . This result has also appeared in An, Drummond-Cole, and Knudsen [ADK, Proposition 5.6]. To the knowledge of the author, the present work is the first to provide some evidence of such uniform boundedness for the higher syzygies of \mathcal{H}_q , for $q \geq 2$.

2. GRAPH CONFIGURATION SPACES

Definition 2.1. A **graph** is a 1-dimensional compact simplicial complex. We call the 0-simplices of G its **vertices**, $V(G)$, while the 1-simplices are its **edges**, $E(G)$. If v is a vertex of G , then the **degree** of v , denoted $\mu(v)$, is the number of edges adjacent to v . A vertex is said to be **essential** if it has $\mu(v) \neq 2$. A **homomorphism of graphs** is a map between vertices

$$\phi : V(G) \rightarrow V(G')$$

which preserves adjacency.

If G is a graph, the **(unordered) n -strand configuration space of G** , denoted $UF_n(G)$, is given by

$$UF_n(G) := \{(x_1, \dots, x_n) \in G^n \mid x_i \neq x_j\} / \mathfrak{S}_n$$

where \mathfrak{S}_n acts by permuting coordinates in the natural way.

Our main concern will be developing a better understanding of the homology groups $H_q(UF_n(G))$. While much of the current literature on the subject has been concerned with their relation to $\pi_1(UF_n(G))$ (see [FS, KKP], for instance), in this work we will approach their study from a more recent algebraic perspective.

Definition 2.2. Writing $E(G)$ for the set of edges of G , We will use A_G to denote the polynomial ring

$$A_G := \mathbb{Z}[x_e \mid e \in E(G)]$$

In [R], the author showed that the total q -homology group, $\mathcal{H}_q(G) = \bigoplus_n H_q(UF_n(G))$ could be endowed with the structure of a finitely generated, graded module over A_G whenever G was a tree. Following this, An, Drummond-Cole, and Knudsen showed that this could be extended to all graphs [ADK]. Our interest in this paper will be to understand the syzygies of the module \mathcal{H}_q for a certain family of graphs, which we call edge-linear (see Definition 3.11). In particular, we prove facts about the Betti numbers, in the commutative algebraic sense, by applying the techniques of FI-algebras recently developed by Nagel and Römer [NR], building on the work of Sam [Sa, Sa2], Snowden [Sn], and others.

Definition 2.3. Let G be a graph. A **half-edge** of G is a pair $h = (v, e)$ where v is a vertex and e is an edge adjacent to v . We will use $v(h)$ and $e(h)$ to denote the vertex and edge associated to h , respectively. The set of half edges with $v(h) = v$ will be written $H(v)$.

For a vertex $v \in V(G)$, we set $Sw(v)$ to be the abelian group freely generated by the symbols

$$Sw(v) = \langle \emptyset, v, h \in H(v) \rangle .$$

We consider $Sw(v)$ to be a bigraded abelian group by setting

$$|v| = (0, 1), |\emptyset| = (0, 0), |h| = (1, 1).$$

The **Swiatkowski complex** associated to G is the differential bigraded A_G -module

$$Sw(G) = A_G \otimes \bigotimes_v Sw(v),$$

with

$$|e| = (0, 1)$$

and differential defined by

$$\partial(h) = e(h) - v(h), \partial(e) = \partial(v) = \partial(\emptyset) = 0.$$

Expressed more classically, we may think of the Swiatkowski complex in the following way. For each $q \geq 0$, write $Sw_q(G)$ to be the subgroup of $Sw(G)$ generated by pure tensors of grade (q, x) for some $x \geq 0$. Then $Sw_q(G)$ can be thought of as a graded module over A_G . The Swiatkowski complex is the complex of graded A_G -modules

$$Sw_*(G) : \dots \rightarrow Sw_q(G) \xrightarrow{\partial} Sw_{q-1}(G) \rightarrow \dots \rightarrow Sw_0(G) \rightarrow 0$$

where ∂ is as above. We will use these two perspectives interchangeably in what follows.

Remark 2.4. Because it will be relevant later, we note that the assignment $G \mapsto Sw(G)$ is functorial. In particular, given an injective homomorphism of graphs $\phi : G \rightarrow G'$ one obtains a morphism $Sw(G) \rightarrow Sw(G')$. The reader can check that this morphism preserves all the relevant structure of the complex including the bigrading, as well as the differential.

We will follow the notation of An, Drummond-Cole, and Knudsen regarding shifts in the grading of bigraded modules. If M is a differential bigraded module over some algebra A , then we will write $M\{1\}$ to denote the bigraded module with

$$M\{1\}_{(a,b)} = M_{(a,b-1)}.$$

Similarly, we will write $M[1]$ to denote the module with bigrading

$$M[1]_{(a,b)} = M_{(a-1,b)}.$$

The differentials of these modules follow the usual sign conventions for degree shifting.

The Swiatkowski complex was originally expressed in the above form by An, Drummond-Cole, and Knudsen in [ADK], though they credit work of Swiatkowski [Sw] for the origin of the idea. The following theorem is extremely relevant for the remainder of this paper.

Theorem 2.5 (An, Drummond-Cole, and Knudsen, Theorem 4.3 [ADK]). *There is an isomorphism of functors from the category of graphs and injective homomorphisms to bigraded abelian groups,*

$$\mathcal{H}_*(G) \cong H_*(Sw(G))$$

In particular, for any graph G the total q -homology group $\mathcal{H}_q(G)$ carries the structure of a finitely generated graded module over the polynomial ring A_G .

In the case where G is a tree, the structure of $\mathcal{H}_q(G)$ as a A_G -module was explored by the author [R]. It is shown in that work that $\mathcal{H}_q(G)$ decomposes as a direct sum of graded shifts of square-free monomial ideals. In the work [ADK], An, Drummond-Cole and Knudsen also explore $\mathcal{H}_q(G)$ in the case where G has maximum valency 3.

For the purpose of doing computations later in this work, we will need to present a simplification of the Swiatkowski complex.

Definition 2.6. Let G be a graph with no isolated vertices. For each vertex v of G we may enumerate the half-edges containing v by $\{h_i\}$. Define a bigraded subgroup of $S(v)$ by,

$$\tilde{S}(v) := \langle \emptyset, h_{i,j} \rangle_{>,j},$$

where $h_{i,j} = h_i - h_j$. Then the **reduced Swiatkowski complex** is the submodule of $Sw(G)$ defined by

$$\tilde{Sw}(G) := A_G \otimes \bigotimes_v \tilde{S}(v)$$

One observes that for any pair of half edges h_i, h_j containing some fixed vertex v , one has

$$\partial(h_{i,j}) = e(h_i) - e(h_j)$$

It follows that $\tilde{Sw}(G)$ is indeed a differential bigraded submodule of the Swiatkowski complex.

Proposition 2.7 (An, Drummond-Cole, Knudsen, Proposition 4.9 [ADK]). *If G does not contain any isolated vertices, then the inclusion*

$$\widetilde{Sw}(G) \hookrightarrow Sw(G)$$

is a quasi-isomorphism.

3. FI-ALGEBRAS AND EDGE-LINEAR FI-GRAPHS

Definition 3.1. For each integer $n \geq 0$, we will write $[n]$ to denote the finite set $[n] := \{1, \dots, n\}$. The category FI is that whose objects are the sets $[n]$ and whose morphisms are set injections.

If k is a commutative Noetherian ring, then an **FI-module over k** is a functor M from FI to the category of k -modules. We will often write M_n to denote $M([n])$.

The representation theory of FI has recently risen to popularity due in large part to the seminal works of authors such as Church, Ellenberg, Farb, and Nagpal [CEF, CEFN], Djament and Vespa [D, DV], Sam and Snowden [SS, SS2], and many others. In this work we will be primarily concerned with two constructions stemming from FI: FI-algebras and FI-graphs.

Definition 3.2. If k is a commutative Noetherian ring, an FI-algebra is a functor \mathbf{A} from FI to the category of k -algebras. Given an FI-algebra \mathbf{A} , an **A-module** is an FI-module M satisfying the following:

1. M_n is an \mathbf{A}_n -module for each n ;
2. if $f : [n] \hookrightarrow [m]$ is an injection of sets then the following diagram commutes for all $a \in A_n$

$$\begin{array}{ccc} M_n & \xrightarrow{M(f)} & M_m \\ a \downarrow & & A(f)(a) \downarrow \\ M_n & \xrightarrow{M(f)} & M_m \end{array}$$

An **A-module** M is said to be **finitely generated** so long as there is a finite collection of elements $\{v\} \subseteq \bigoplus M_n$ which no proper sub **A-module** contains. We say that a finitely generated **A-module** M is **Noetherian** if every sub **A-module** of M is also finitely generated. We say that \mathbf{A} is itself **Noetherian** if all finitely generated modules over \mathbf{A} are Noetherian.

Example 3.3. The simplest example of an FI-algebra is that whose points are defined by $\mathbf{A}_n = k$, and whose transition maps are all the identity. In this case **A-modules** are no different than FI-modules over k .

The second example of an FI-algebra, and the example which will be the fundamental object of study in this paper, is that given on points by $\mathbf{A}_n = k[x_1, \dots, x_n]$, and whose transition maps act on indices in the obvious way.

Note that for any fixed $d \geq 0$, and any FI-algebra \mathbf{A} , we may always define a new algebra given by the tensor power

$$(\mathbf{A}^{\otimes d})_n := (\mathbf{A}_n)^{\otimes d}$$

It is famously known that all finitely generated FI-modules over Noetherian rings are themselves Noetherian (see [CEFN] for one such proof). However, finitely generated modules over a general FI-algebra do not share this property. This is true even if we assume that \mathbf{A} is itself finitely generated. Despite this, one can prove the following.

Theorem 3.4 (Nagel and Römer, Corollary 7.9 [NR]). *If k a commutative Noetherian ring, the FI-algebra \mathbf{A} given by $\mathbf{A}_n = k[x_1, \dots, x_n]$ is Noetherian. More generally, for any fixed integer $d \geq 0$, $\mathbf{A}^{\otimes d}$ is Noetherian.*

Remark 3.5. It is still open whether the Noetherian property of FI-algebras is preserved by tensor products. The above special case was proven by Nagel and Römer in the provided source.

Theorem 3.4 has a variety of interesting consequences in the case where we keep track of the natural grading on \mathbf{A}_n .

Definition 3.6. We say that an FI-algebra \mathbf{A} over a field k is **standard graded** if every algebra \mathbf{A}_n is a graded k -algebra generated in degree 1 with $\mathbf{A}_{n,0} = k$, and every induced map $A(f)$ is a morphism of graded k -algebras. We say that an \mathbf{A} -module M is **graded** if each M_n is a graded module over \mathbf{A}_n , and each induced map $M(f)$ preserves this grading.

If M is a finitely generated graded \mathbf{A} -module, then for each $p, n \geq 0, j \in \mathbb{Z}$ we define $\beta_{n,p,j}(M)$ to be the Betti number

$$\beta_{n,p,j}(M) := \dim_k(\mathrm{Tor}_p^{\mathbf{A}_n}(M_n, k)_j).$$

Corollary 3.7 (Nagel and Römer, Theorem 7.7 [NR]). *Let \mathbf{A} be a finitely generated standard graded FI-algebra over a field k . Then for any integers $p, j \geq 0$, and any finitely generated graded \mathbf{A} -module M , the function*

$$n \mapsto \beta_{n,p,j}(M)$$

agrees with a polynomial for $n \gg 0$. Moreover, there exists a finite list of integers

$$j_0(M, p) < \dots < j_t(M, p)$$

such that for all $n \gg 0$

$$\beta_{n,p,j}(M_n) \neq 0 \iff j \in \{j_0(M, p), \dots, j_t(M, p)\}.$$

Remark 3.8. The first part of Corollary 3.7 does not explicitly appear in [NR], although it is a clear consequence of the proof of [NR, Theorem 7.7].

One way to think about Corollary 3.7 is that if M is a finitely generated graded \mathbf{A} -module, then in each fixed degree p the p -th syzygies eventually stabilize. One should note, however, that it is not the case that this stabilization is uniform across all p . Namely, one should not expect the regularity of M_n to be eventually independent of n .

For the remainder of the paper, we will reserve \mathbf{A} to mean the FI-algebra defined by $\mathbf{A}_n = \mathbb{Z}[x_1, \dots, x_n]$.

The second FI-structure which will be important to us are FI-graphs.

Definition 3.9. An FI-graph is a functor G_\bullet from FI to the category of graphs with graph homomorphisms. As with FI-algebras, we will often write G_n to denote $G_\bullet([n])$ and $G(f)$ to denote $G_\bullet(f)$. We say that G_\bullet is finitely generated if for $n \gg 0$, the vertex set $V(G_{n+1})$ is equal to $\cup_f G(f)(V(G_n))$, where the union is over injections $f : [n] \hookrightarrow [n+1]$.

We say that G_\bullet is torsion-free if the induced maps $G(f)$ are injective for all f .

Example 3.10. The most typical example of an FI-graph is the complete graph K_n . This is the graph whose vertices are labeled by $[n]$, and whose edges are labeled by pairs $\{i, j\}$ with $i \neq j$.

For the purposes of this paper a more relevant example is given as follows. Let (G, v_G) and (H, v_H) be any pair of graphs with a choice of vertex. Then we define

$$G_n := G \bigvee^n H.$$

For any injection, the induced map on G_\bullet is defined by fixing the copy of G , and permuting the copies of H . For example, if G is a single vertex, and H is an edge, then G_n is the star graph, which has a unique essential vertex of degree n . This example was first discussed in the context of FI-graphs by Lütgehetmann in [L], although he does not use the language of FI-graphs.

FI-graphs were first introduced by the author and Graham White in [RW]. Configuration spaces of FI-graphs are shown to exhibit a variety of stable behaviors in that work, extending results of Lütgehetmann from [L]. In this work we combine these stabilization results with the aforementioned A_G -action.

Definition 3.11. Let G_\bullet be a finitely generated FI-graph. Then it is proven in [RW] that the function

$$n \mapsto |E(G_n)|$$

agrees with a polynomial for all $n \gg 0$. We say that G_\bullet is **edge-linear** if it is finitely generated, and $|E(G_n)|$ agrees with a linear polynomial for all $n \gg 0$.

Example 3.12. The second example given in Example 3.10 is edge-linear. Moreover, for any fixed $m \geq 1$ the FI-graph defined by $G_n = K_{n,m}$, the complete bipartite graph on $(n+m)$ -vertices, is also edge-linear.

Both of these examples can be thought of as being construct via the following gluing procedure. Let G and G' be two graphs, each sharing a common subgraph H . Then for any $n \geq 0$, we obtain a new graph

$$G \bigsqcup_H^n G'$$

by gluing n copies of G' to G along H . This family of graphs may be given an FI-graph structure, where the symmetric group acts by permuting the copies of G' , while fixing G . Graphs of this kind were originally studied by Lütgehetmann in [L]. The Classification Theorem 3.13 essentially implies that all edge-linear FI-graphs are isomorphic to something of the form $G \bigsqcup_H^\bullet G'$, for some G, G', H .

Edge-linear FI-graphs will be useful in this work due to their connection to the FI-algebra \mathbf{A} . We explore this connection as a corollary to the following classification result.

Theorem 3.13. *Let G_\bullet be an edge-linear FI-graph. Then there exists a pair of graphs G, G' , with a common subgraph H , such that for all $n \gg 0$*

$$G_n \cong G \bigsqcup_H^n G'$$

Proof. We begin by observe that an edge-linear FI-graph must also have linear growth in its number of vertices. Indeed, this follows from the graph-theoretic inequality

$$|V(G)| \leq 2|E(G)|/\delta(G)$$

where $\delta(G)$ is the minimum degree of a vertex of G . Appealing to the classification theorem for FI-sets [RSW, Theorem A], we conclude that $V(G_n)$ is isomorphic to a disjoint union of some number of copies of the \mathfrak{S}_n -action on $[n]$, along with some number of invariant vertices.

The only \mathfrak{S}_n -equivariant relation on $[n]$ is the identity relation $\{(i, i) \mid i \in [n]\}$, and the anti-identity relation $\{(i, j) \mid i \neq j\}$. In the latter case, the number of edges does not grow linearly in n . Therefore, there are no edges between two vertices in the same \mathfrak{S}_n -orbits. If we write two distinct \mathfrak{S}_n -orbits of vertices as

$$\mathcal{O}_1 = \{(i, 1) \mid i \in [n]\}, \mathcal{O}_2 = \{(i, 2) \mid i \in [n]\}$$

then the same argument before shows that there cannot be edges between vertices of the form $(i, 1)$ and $(j, 2)$ if $i \neq j$. Finally, if there is an edge between $(i, 1)$ and $(i, 2)$ for some i , then there must be an edge between $(i, 1)$ and $(i, 2)$ for all i . This then leads into the construction of the desired graphs G, G' , and H : let \tilde{G} be the graph whose vertices are labeled by \mathfrak{S}_n -orbits of vertices of G_n , and whose edges indicate the existence of an edge in G_n between two elements of the orbits. We let G be the induced subgraph of \tilde{G} whose vertices

correspond to singleton orbits, let G' be any subgraph of \tilde{G} containing all the non-singleton orbits and all edges adjacent to these orbits, and let H be the intersection in \tilde{G} of G and G' . \square

Remark 3.14. It was revealed in the above proof that the choices of G' and H are not necessarily unique, though G is unique.

The asymptotic nature of Theorem 3.13 is necessary. For instance, one may define an FI-graph which agrees with the bipartite graph $K_{n,n}$ for $n \leq 10$, and collapses to a single edge for all larger n . This observation inspires the following definition.

Definition 3.15. Let G_\bullet be an edge-linear FI-graph, and let m be sufficiently large so that $G_n \cong G \sqcup_H^n G'$ for all $n \geq m$, and for some G, G', H . The **tail** of G_\bullet , G_\bullet^{\gg} , is the edge-linear FI-graph,

$$G_n^{\gg} := \begin{cases} \emptyset & \text{if } n < m \\ G_n & \text{otherwise.} \end{cases}$$

We also define an FI-algebra \mathbf{A}_{G_\bullet} by setting

$$\mathbf{A}_{G_\bullet, n} := A_{G \sqcup_H^n G'}$$

for all $n \geq 0$, where G, G' , and H are as above.

Remark 3.16. While the tail of an FI-graph depends on the choice of m , this choice will not matter in what follows.

Proposition 3.17. *Let G_\bullet be an edge-linear FI-graph. Then there is some $d \geq 0$ such that \mathbf{A}_{G_\bullet} is a quotient of*

$$\mathbf{A}^{\otimes d}.$$

In particular, \mathbf{A}_{G_\bullet} is Noetherian.

Proof. Appealing to Theorem 3.13, it follows that $E(G_n)$ has $a + b$ \mathfrak{S}_n -orbits: a orbits isomorphic to the action of \mathfrak{S}_n on $[n]$, and b singleton orbits. We conclude that \mathbf{A}_{G_\bullet} is isomorphic to a quotient of $\mathbf{A}^{\otimes(a+b)}$. Namely, the quotient which identifies the variables of the first a tensor factors with the elements in the non-singleton orbits of $E(G_n)$, and identifies the variables of the final b tensor factors with the b singleton orbits.

The second part of the proposition follows from the fact that any quotient of a Noetherian FI-algebra must be Noetherian, and Theorem 3.4. \square

The goal of the next section is to show that if G_\bullet is an edge-linear FI-graph, then the assignments

$$n \mapsto \mathcal{H}_q(G_\bullet)$$

defines a finitely generated module over \mathbf{A}_{G_\bullet} . To do this we will make critical use of the fact that \mathbf{A}_{G_\bullet} is Noetherian.

4. THE PROOFS OF THE MAIN THEOREMS

For the remainder of this section we fix an edge-linear FI-graph G_\bullet . **In fact, we substitute G_\bullet by its tail G_\bullet^{\gg} so that G_\bullet is torsion free.** We will also assume that G_n does not contain any isolated vertices for all n . In view of Corollary 3.7 and Proposition 3.17, it will suffice to show that the FI-module

$$\mathcal{H}_q(G_\bullet) := \bigoplus_{n \geq 0} H_q(U\mathcal{F}_n(G_\bullet))$$

is finitely generated and graded over the FI-algebra \mathbf{A}_{G_\bullet} . We accomplish this by proving that the collection of reduced Swiatkowski complexes $\{\widetilde{Sw}(G_n)\}_{n \geq 0}$ can be extended to a complex of graded \mathbf{A}_{G_\bullet} -modules, $\widetilde{Sw}(G_\bullet)$. We will then show that the terms in the complex $\widetilde{Sw}(G_\bullet)$ are finitely generated \mathbf{A}_{G_\bullet} -modules. The Noetherian property (Proposition 3.17) will imply that $\mathcal{H}_q(G_\bullet)$ is finitely generated, as desired.

Lemma 4.1. *For any injection of sets $f : [n] \hookrightarrow [r]$, and any $q \geq 0$ the following diagram commutes*

$$\begin{array}{ccc} Sw_q(G_r) & \xrightarrow{\partial} & Sw_{q-1}(G_r) \\ Sw(G(f)) \uparrow & & \uparrow Sw(G(f)) \\ Sw_q(G_n) & \xrightarrow{\partial} & Sw_{q-1}(G_n) \end{array}$$

In particular, the collection of Swiatkowski complexes $\{Sw(G_n)\}_{n \geq 0}$ can be extended to a complex of graded \mathbf{A}_{G_\bullet} -modules.

Proof. This follows from the definition of ∂ , as well as the definition of $Sw(G(f))$. Indeed, we have for any half edge $h = (v, e)$

$$\partial(Sw(G(f))(h)) = \partial(G(f)(e), G(f)(v)) = G(f)(e) - G(f)(v) = Sw(G(f))(\partial(h)).$$

□

One observes that, for any injection $f : [n] \hookrightarrow [r]$, $Sw(G(f))$ restricts to a map $\widetilde{Sw}(G_n) \rightarrow \widetilde{Sw}(G_r)$. The above lemma therefore shows that the reduced Swiatkowski complex can also be extended to a complex of graded \mathbf{A}_{G_\bullet} -modules.

Lemma 4.2. *For each $q \geq 0$, the \mathbf{A}_{G_\bullet} -module $\widetilde{Sw}_q(G_\bullet)$ is finitely generated.*

Remark 4.3. One should observe that this lemma is false if we insist on working with the usual Swiatkowski complex instead of the reduced one.

Proof. We begin by defining an FI-set, in the sense of [RSW], $X_{q,d}(G_\bullet)$. For each n , the elements of $X_{q,d}(G_\bullet)$ are tuples

$$(m, \tilde{h}_1, \dots, \tilde{h}_q)$$

where m is a monomial in A_{G_n} of degree d , \tilde{h}_i is a pair of distinct half-edges containing a common vertex of G_n , and \tilde{h}_i, \tilde{h}_j are associated to different vertices whenever $i \neq j$. It isn't hard to see that $X_{q,d}(G_\bullet)$ is a FI-subset of a product of finitely generated FI-sets. This implies that it is itself finitely generated (see [RSW, Proposition 4.2]).

Note that we may associate to each element of $X_{q,d}(G_n)$ a generating element of $\widetilde{Sw}_q(G_n)$. In fact, we have that

$$\widetilde{Sw}(G_\bullet) \cong \bigoplus_{d \geq 0} \mathbb{Z}X_{q,d}(G_\bullet)$$

where $\mathbb{Z}X_{q,d}(G_\bullet)$ is the FI-module \mathbb{Z} -linearization of $X_{q,d}(G_\bullet)$. This shows that $\widetilde{Sw}(G_\bullet)$ is finitely generated, as desired. □

As previously stated, these two lemmas and the Noetherian property immediately lead to the following consequence.

Theorem 4.4. *Let G_\bullet be an edge-linear FI-graph. For each $q \geq 0$, the \mathbf{A}_{G_\bullet} -module $\mathcal{H}_q(G_\bullet)$ is finitely generated.*

Remark 4.5. The two preceding lemma do not use edge-linearity of the FI-graph G_\bullet . Indeed, the only thing preventing one from concluding an analog of Theorem 4.4 for all FI-graphs is the failing of Noetherianity outside of the linear case. It is unclear whether one should expect such a finite generation result in these cases.

Base change from \mathbb{Z} to any field will preserve finite generation of FI-modules, and so we also obtain the following as a corollary to the above

Corollary 4.6. *Let G_\bullet be an edge-linear FI-graph, and let k be a field. Then for each $p, j \geq 0$, the function*

$$n \mapsto \beta_{p,j}(\mathcal{H}_q(G_n; k))$$

agrees with a polynomial for $n \gg 0$. Moreover, there exists a finite list of integers

$$j_0(\mathcal{H}_q(G_n; k), p) < \dots < j_t(\mathcal{H}_q(G_n; k), p)$$

such that for all $n \gg 0$

$$\beta_{n,p,j}(\mathcal{H}_q(G_n; k)) \neq 0 \iff j \in \{j_0(\mathcal{H}_q(G_n; k), p), \dots, j_t(\mathcal{H}_q(G_n; k), p)\}.$$

5. AN INDUCTIVE METHOD FOR COMPUTATION

In this section we outline an inductive method with which one can compute the \mathbf{A}_{G_\bullet} -module $\mathcal{H}_q(G_\bullet)$. This method extends a similar method originally described by An, Drummond-Cole, and Knudsen [ADK, Proposition 5.13]. We then conclude the section by performing some example computations

As with the previous section, we fix a edge-linear FI-graph G_\bullet , which we make torsion free by working with its tail.

Definition 5.1. The theory of finitely generated FI-graphs tells us that for $n \gg 0$ the \mathfrak{S}_n -orbits of vertices of G_n eventually stabilize (see, [RSW, Proposition 3.6]), in the sense that the transition maps of G_\bullet induce isomorphisms

$$V(G_n)/\mathfrak{S}_n \cong V(G_{n+1})/\mathfrak{S}_{n+1}.$$

Let $m \gg 0$ be in this stable range, and let v be a vertex of G_m which is fixed by \mathfrak{S}_m . We will often say that v is an **invariant vertex of G_\bullet** . Then all images of v under all transition maps of G_\bullet are also invariant under the action of the relevant symmetric group. By abuse of notation, we say that v is a vertex of G_n for all $n \geq m$.

The **blow up of G_\bullet at v** is an FI-graph $Bl_v(G_\bullet)$, defined as follows. For $n \geq m$, $Bl_v(G_n)$ is the graph with vertex set

$$V(Bl_v(G_n)) := (V(G_n) - \{v\}) \cup (\{v\} \times H(v)),$$

where $H(v)$ is the set of half-edges containing v . The edges of $Bl_v(G_n)$ are the edges of G_n , altered in the obvious way to account for the new vertices. We also set $Bl_v(G_n) = \emptyset$ for $n < m$. See [ADK, Figure 9] for a pictorial representation of the blow up.

Let $f : [n] \hookrightarrow [r]$ be an injection of sets. Then the transition map induced by f is defined as follows. On vertices not of the form $v \times h$, the transition map agrees with $G(f)$. Otherwise, we have

$$Bl_v(G_\bullet)(f)(v \times h) = v \times G(f)(h).$$

This is well defined, as v was an invariant vertex.

Remark 5.2. We will often talk of invariant vertices of an FI-graph, without making explicit reference to what degree they originate from. Because much of what we care above is asymptotic, this shouldn't cause confusion.

Following the techniques of [ADK], our goal will be to relate $\mathcal{H}_q(G_\bullet)$ to $\mathcal{H}_q(\text{Bl}_v(G_\bullet))$. We first observe that $\mathcal{H}_q(\text{Bl}_v(G_\bullet))$ is a \mathbf{A}_{G_\bullet} -module. Indeed, the construction of the blowup does not erase, or meaningfully alter edges.

Lemma 5.3. *Let v be an invariant vector of G_\bullet . Then there is an inclusion of complexes*

$$\widetilde{Sw}(\text{Bl}_v(G_\bullet)) \hookrightarrow \widetilde{Sw}(G_\bullet).$$

Proof. one can realize $\widetilde{Sw}(\text{Bl}_v(G_n))$ as the subcomplex of $\widetilde{Sw}(G_n)$ generated by pure tensors whose v -th term is \emptyset . The fact that this extends to the desired inclusion follows from the fact that v was chosen to be an invariant vector. \square

The above proof immediately points to how one can describe the cokernel of $\widetilde{Sw}(\text{Bl}_v(G_\bullet)) \hookrightarrow \widetilde{Sw}(G_\bullet)$. To accomplish this we will need some notation. If v is an invariant vector of G_\bullet , then we define an FI-module $\widetilde{H}(v)$ to be that for which

$$\widetilde{H}(v)_n = \mathbb{Z} \langle h_i - h_j \mid h_i, h_j \in H(v)_n \rangle,$$

where $H(v)_n$ is the set of half-edges containing v , thought of as a vertex of G_n . We now extend the above lemma as follows. Note that this result generalizes a lemma of An, Drummond-Cole, and Knudsen [ADK].

Lemma 5.4. *Let v be an invariant vector of G_\bullet . Then there is an exact sequence of complexes*

$$0 \rightarrow \widetilde{Sw}(\text{Bl}_v(G_\bullet)) \rightarrow \widetilde{Sw}(G_\bullet) \rightarrow \widetilde{H}(v) \otimes \widetilde{Sw}(\text{Bl}_v(G_\bullet))\{1\} \rightarrow 0$$

It therefore follows that computing $\mathcal{H}_q(G_\bullet)$ can often times be reduced to computing $\mathcal{H}_q(\text{Bl}_v(G_\bullet))$. This inductive approach becomes even more appealing when one considers how simple its base case is. Indeed, the classification theorem implies that after sufficiently many blow-ups one is left with some fixed number of disjoint edges, invariant under the action of the symmetric group, paired with n disjoint copies of some graph, which the symmetric group permutes.

Example 5.5. Take G_\bullet to be the star graph

$$G_n = K_{n,1}$$

This graph has a unique invariant vertex, and $\text{Bl}_v(G_n)$ is a disjoint union of n edges. It is fact that $U\mathcal{F}_n(G_n)$ only has homology in degrees 0 and 1 (see, for instance, [Sw]). We therefore have,

$$0 \rightarrow \mathcal{H}_1(G_\bullet) \rightarrow S^{(\bullet-1,1)} \otimes \mathcal{H}_0(\text{Bl}_v(G_\bullet))\{1\} \rightarrow \mathcal{H}_0(\text{Bl}_v(G_\bullet)) \rightarrow \mathcal{H}_0(G_\bullet) \rightarrow 0,$$

where $S^{(\bullet-1,1)}$ is the FI-module for which $S^{(n-1,1)}$ is the standard representation of \mathfrak{S}_n . Examining this exact sequence from right to left, we note that $\mathcal{H}_0(G_n) = \mathbb{Z}$ for all $n \geq 0$, while $\mathcal{H}_0(\text{Bl}_v(G_\bullet)) \cong \mathbf{A}$. The map

$$\mathcal{H}_0(\text{Bl}_v(G_\bullet)) \rightarrow \mathcal{H}_0(G_\bullet) \rightarrow 0$$

identifies all of the disjoint edges of $\text{Bl}_v(G_\bullet)$. In particular, the kernel of this map is the module \mathbf{m}_\bullet , defined on points by

$$\mathbf{m}_n = \langle x_i - x_j \mid i \neq j \rangle$$

On the other hand, the map

$$S^{(\bullet-1,1)} \otimes \mathbf{A}\{1\} \rightarrow \mathbf{m}_\bullet \rightarrow 0$$

is given by

$$(e_i - e_j) \otimes 1 \mapsto x_i - x_j.$$

The kernel of this map is the submodule of $S^{(\bullet-1,1)} \otimes \mathbf{A}\{1\}$ generated in degree 3 by

$$(e_1 - e_2) \otimes x_3 + (e_3 - e_1) \otimes x_2 + (e_2 - e_3) \otimes x_1$$

In particular, we conclude that all of the homology classes in $H_1(U\mathcal{F}_n(G_n))$ are determined by $H_1(U\mathcal{F}_2(G_3))$. This recovers [ADK, Lemma 5.5]. A straight forward computation also verifies that for all $d \gg 0$, the $\mathbb{Z}[x_1, \dots, x_d]$ -module

$$\mathcal{H}_1(G_d)$$

is generated in grade 2, with relations generated in grades ≤ 3 , verifying the second part of Theorem B for the first two Betti numbers.

REFERENCES

- [ADK] B. H. An, G. C. Drummond-Cole and B. Knudsen, *Subdivisional spaces and graph braid groups*, [arXiv:1708.02351](#).
- [CEF] T. Church, J. S. Ellenberg and B. Farb, *FI-modules and stability for representations of symmetric groups*, *Duke Math. J.* 164, no. 9 (2015), 1833-1910.
- [CEFN] T. Church, J. S. Ellenberg, B. Farb, and R. Nagpal, *FI-modules over Noetherian rings*, *Geom. Topol.* 18 (2014) 2951-2984.
- [CF] T. Church and B. Farb, *Representation theory and homological stability*, *Advances in Mathematics* (2013), 250-314.
- [D] A. Djament, *Des propriétés de finitude des foncteurs polynomiaux*, [arXiv:1308.4698](#).
- [DV] A. Djament, and C. Vespa, *Foncteurs faiblement polynomiaux*, [arXiv:1308.4106](#).
- [Fa] M. Farber, *Topological complexity of motion planning* *Discrete and Computational Geometry* 29 (2003), 211-221.
- [FS] D. Farley and L. Sabalka, *Discrete Morse theory and graph braid groups*, *Algebr. Geom. Topol.* 5 (2005), 1075-1109 (electronic). <http://www.users.miamioh.edu/farleyds/FS1.pdf>.
- [G] R. Ghrist, *Configuration spaces and braid groups on graphs in robotics*, *Knots, braids, and mapping class groups - papers dedicated to Joan S. Birman (New York, 1998)*, *AMS/IP Stud. Adv. Math.*, 24, Amer. Math. Soc., Providence, RI (2001), 2940. <https://www.math.upenn.edu/~ghrist/preprints/birman.pdf>.
- [L] D. Lütgehetmann, *Representation Stability for Configuration Spaces of Graphs*, [arXiv:1701.03490](#).
- [KP] K. H. Ko and H. W. Park, *Characteristics of graph braid groups*, *Discrete Comput Geom* (2012) 48: 915. [arXiv:1101.2648](#).
- [KKP] J. H. Kim, K. H. Ko, and H. W. Park, *Graph braid groups and right-angled Artin groups*, *Trans. Amer. Math. Soc.* 364 (2012), 309-360. [arXiv:0805.0082](#).
- [NR] U. Nagel and T. Römer, *FI- and OI-modules with varying coefficients*, [arXiv:1710.09247](#).
- [R] E. Ramos, *Stability phenomena in the homology of tree braid groups*, to appear, *Algebraic and Geometric Topology*, [arXiv:1609.05611](#).
- [RW] E. Ramos and G. White, *Families of nested graphs with compatible symmetric-group actions*, [arXiv:1711.07456](#).
- [RSW] E. Ramos, D. Speyer, and G. White, *FI-sets with relations*, [arXiv:1804.04238](#).
- [Sa] S. Sam, *Szygies of bounded rank symmetric tensors are generated in bounded degree*, *Math. Ann.*, to appear. [arXiv:1608.01722](#).
- [Sa2] S. Sam, Steven V Sam, *Ideals of bounded rank symmetric tensors are generated in bounded degree*, *Invent. Math.* 207 (2017), no. 1, 121, [arXiv:1510.04904](#).
- [Sn] A. Snowden, *Szygies of Segre embeddings and Δ -modules*, *Duke Math. J.* 162 (2013), no. 2, 225-277, [arXiv:1006.5248](#).
- [Sw] J. Swiatkowski, *Estimates for homological dimension of configuration spaces of graphs* *Colloquium Mathematicum* 89.1 (2001): 69-79.
- [SS] S. Sam and A. Snowden, *GL-equivariant modules over polynomial rings in infinitely many variables*, *Trans. Amer. Math. Soc.* 368 (2016), 1097-1158.
- [SS2] S. Sam, and A. Snowden, *Gröbner methods for representations of combinatorial categories*, *J. Amer. Math. Soc.* 30 (2017), 159-203. [arXiv:1409.1670](#).

UNIVERSITY OF MICHIGAN DEPARTMENT OF MATHEMATICS, 530 CHURCH ST., ANN ARBOR, MI 48109

E-mail address: egramos@umich.edu