

## LOCAL CURVATURE ESTIMATES FOR THE LAPLACIAN FLOW

YILI

ABSTRACT. In this paper we give local curvature estimates for the Laplacian flow on closed  $G_2$ -structures under the condition that the Ricci curvature is bounded along the flow. The main ingredient consists of the idea of Kotschwar-Munteanu-Wang [24] who gave local curvature estimates for the Ricci flow on complete manifolds and then provided a new elementary proof of Sesum's result [36], and the particular structure of the Laplacian flow on closed  $G_2$ -structures. As an immediate consequence, this estimates give a new proof of Lotay-Wei's [33] result which is an analogue of Sesum's theorem.

The second result is about an interesting evolution equation for the scalar curvature of the Laplacian flow of closed  $G_2$ -structures. Roughly speaking, we can prove that the time derivative of the scalar curvature  $R_t$  is equal to the Laplacian of  $R_t$ , plus an extra term which can be written as the difference of two nonnegative quantities.

## 1. INTRODUCTION

Let  $\mathcal{M}$  be a smooth 7-manifold. The Laplacian flow for closed  $G_2$ -structures on  $\mathcal{M}$  introduced by Bryant [1] is to study the torsion-free  $G_2$ -structures

$$(1.1) \quad \partial_t \varphi_t = \Delta_{\varphi_t} \varphi_t, \quad \varphi_0 = \varphi,$$

where  $\Delta_{\varphi_t} \varphi_t = dd_{\varphi_t}^* \varphi_t + d_{\varphi_t}^* d\varphi_t$  is the Hodge Laplacian of  $g_{\varphi_t}$  and  $\varphi$  is an initial closed  $G_2$ -structure. Since  $dd_{\varphi_t}^* \varphi_t = \partial_t d\Delta_{\varphi_t} \varphi_t = 0$ , we see that the flow (1.1) preserves the closedness of  $\varphi_t$ . For more background on  $G_2$ -structures, see Section 2. When  $\mathcal{M}$  is compact, the flow (1.1) can be viewed as the gradient flow for the Hitchin functional introduced by Hitchin [17]

$$(1.2) \quad \mathcal{H} : [\bar{\varphi}]_+ \longrightarrow \mathbb{R}^+, \quad \varphi \longmapsto \frac{1}{7} \int_{\mathcal{M}} \varphi \wedge \psi = \int_{\mathcal{M}} *_{\varphi} 1.$$

Here  $\bar{\varphi}$  is a closed  $G_2$ -structure on  $\mathcal{M}$  and  $[\bar{\varphi}]_+$  is the open subset of the cohomology class  $[\bar{\varphi}]$  consisting of  $G_2$ -structures. Any critical point of  $\mathcal{H}$  gives a torsion-free  $G_2$ -structure.

The study of Laplacian flows on some special 7-manifolds, Laplacian solitons, and other flows on  $G_2$ -structures can be found in [12, 13, 14, 15, 18, 23, 28, 34, 35, 37, 38].

Recently, Donaldson [6, 7, 8, 9] studied the co-associative Kovalev-Lefschetz fibrations  $G_2$ -manifolds and  $G_2$ -manifolds with boundary.

---

2010 *Mathematics Subject Classification*. Primary 53C44, 53C10.

*Key words and phrases*. Laplacian flow,  $G_2$ -structures, local curvature estimates.

**1.1. Notions and conventions.** To state the main results, we fix our notions used throughout this paper. Let  $\mathcal{M}$  be as before a smooth 7-manifold. The space of smooth functions and the space of smooth vector fields are denoted respectively by  $C^\infty(\mathcal{M})$  and  $\mathfrak{X}(\mathcal{M})$ . The space of  $k$ -tensors (i.e.,  $(0, k)$ -covariant tensor fields) and  $k$ -forms on  $\mathcal{M}$  are denoted, respectively, by  $\otimes^k(\mathcal{M}) = C^\infty(\otimes^k(T^*\mathcal{M}))$  and  $\wedge^k(\mathcal{M}) = C^\infty(\wedge^k(T^*\mathcal{M}))$ . For any  $k$ -tensor field  $T \in \otimes^k(\mathcal{M})$ , we locally have the expression  $T = T_{i_1 \dots i_k} dx^{i_1} \otimes \dots \otimes dx^{i_k} =: T_{i_1 \dots i_k} dx^{i_1 \otimes \dots \otimes i_k}$ . A  $k$ -form  $\alpha$  on  $\mathcal{M}$  can be written in the *standard form* as  $\alpha = \frac{1}{k!} \alpha_{i_1 \dots i_k} dx^{i_1} \wedge \dots \wedge dx^{i_k} =: \frac{1}{k!} \alpha_{i_1 \dots i_k} dx^{i_1 \wedge \dots \wedge i_k}$ , where  $\alpha_{i_1 \dots i_k}$  is fully skew-symmetric in its indices. Using the standard forms, if we take the interior product  $X \lrcorner \alpha$  of a  $k$ -form  $\alpha \in \wedge^k(\mathcal{M})$  with a vector field  $X \in \mathfrak{X}(\mathcal{M})$ , we obtain the  $(k-1)$ -form  $X \lrcorner \alpha = \frac{1}{(k-1)!} X^m \alpha_{m i_1 \dots i_{k-1}} dx^{i_1 \wedge \dots \wedge i_{k-1}}$  which is also in the standard form. In particular, consider the vector space  $\otimes^2(\mathcal{M})$  of 2-tensors. For any 2-tensor  $A = A_{ij} dx^{i \otimes j}$ , define  $A^\odot := \frac{1}{2}(A_{ij} + A_{ji}) dx^{i \otimes j} \equiv A_{ij}^\odot dx^{i \otimes j}$  and  $A^\wedge := \frac{1}{2}(A_{ij} - A_{ji}) dx^{i \otimes j} \equiv A_{ij}^\wedge dx^{i \otimes j}$ . Then  $A^\odot$  is an element of  $\odot^2(\mathcal{M})$ , the space of symmetric 2-tensors. Since  $dx^{i \wedge j} = dx^{i \otimes j} - dx^{j \otimes i}$ , it follows that  $A^\wedge = \frac{1}{2} A_{ij} dx^{i \wedge j}$ . Define  $\alpha^A := \frac{1}{2} \alpha_{ij}^A dx^{i \wedge j}$  with  $\alpha_{ij}^A := A_{ij}$ . Then we see that  $\alpha^A = A^\wedge \in \wedge^2(\mathcal{M})$  and  $\otimes^2(\mathcal{M}) = \odot^2(\mathcal{M}) \oplus \wedge^2(\mathcal{M})$ .

A given Riemannian metric  $g$  on  $\mathcal{M}$  determines two isomorphisms between vector fields and 1-forms:  $b_g : \mathfrak{X}(\mathcal{M}) \rightarrow \wedge^1(\mathcal{M})$  and  $\sharp_g : \wedge^1(\mathcal{M}) \rightarrow \mathfrak{X}(\mathcal{M})$ , where, for every vector field  $X = X^i \frac{\partial}{\partial x^i}$  and 1-form  $\alpha = \alpha_i dx^i$ ,  $b_g(X) = X^i g_{ij} dx^j \equiv X_j dx^j$  and  $\sharp_g(\alpha) = \alpha_i g^{ij} \frac{\partial}{\partial x^j} \equiv \alpha^j \frac{\partial}{\partial x^j}$ . Using these two natural maps, we can frequently raise or lower indices on tensors. The metric  $g$  also induces a metric on  $k$ -forms  $g(dx^{i_1 \wedge \dots \wedge i_k}, dx^{j_1 \wedge \dots \wedge j_k}) = \det(g(dx^{i_a}, dx^{j_b})) = \sum_{\sigma \in \mathfrak{S}_7} \text{sgn}(\sigma) g^{i_1 j_{\sigma(1)}} \dots g^{i_k j_{\sigma(k)}}$  where  $\mathfrak{S}_7$  is the group of permutations of seven letters and  $\text{sgn}(\sigma)$  denotes the sign ( $\pm 1$ ) of an element  $\sigma$  of  $\mathfrak{S}_7$ . The inner product  $\langle \cdot, \cdot \rangle_g$  of two  $k$ -forms  $\alpha, \beta \in \wedge^k(\mathcal{M})$  now is given by  $\langle \alpha, \beta \rangle_g = \frac{1}{k!} \alpha_{i_1 \dots i_k} \beta^{i_1 \dots i_k} = \frac{1}{k!} \alpha_{i_1 \dots i_k} \beta_{j_1 \dots j_k} g^{i_1 j_1} \dots g^{i_k j_k}$ .

Given two 2-tensors  $A, B \in \otimes^2(\mathcal{M})$ , with the forms  $A = A_{ij} dx^{i \otimes j}$  and  $B = B_{ij} dx^{i \otimes j}$ . Define  $\langle \langle A, B \rangle \rangle_g := A_{ij} B^{ij}$ . There are two special cases which will be used later:

- (1)  $\alpha = \frac{1}{2} \alpha_{ij} dx^{i \wedge j} \in \wedge^2(\mathcal{M})$  and  $B = B_{ij} dx^{i \otimes j} \in \otimes^2(\mathcal{M})$ . In this case,  $\alpha$  can be written as a 2-tensor  $A^\alpha = A_{ij}^\alpha dx^{i \otimes j}$  with  $A_{ij}^\alpha = \alpha_{ij}$ . Then  $\langle \langle \alpha, B \rangle \rangle_g := \langle \langle A^\alpha, B \rangle \rangle_g = \alpha_{ij} B^{ij}$ .
- (2)  $\alpha = \frac{1}{2} \alpha_{ij} dx^{i \wedge j}$  and  $\beta = \frac{1}{2} \beta_{ij} dx^{i \wedge j} \in \wedge^2(\mathcal{M})$ . In this case,  $\alpha, \beta$  can be both written as 2-tensors  $A^\alpha = A_{ij}^\alpha dx^{i \otimes j}$  and  $B^\beta = B_{ij}^\beta dx^{i \otimes j}$  with  $A_{ij}^\alpha = \alpha_{ij}$  and  $B_{ij}^\beta = \beta_{ij}$ . Then  $\langle \langle \alpha, \beta \rangle \rangle_g := \langle \langle A^\alpha, B^\beta \rangle \rangle_g = \alpha_{ij} \beta^{ij} = 2 \langle \alpha, \beta \rangle_g$ .

---

<sup>1</sup>In our convention, for any 2-form  $\alpha = \frac{1}{2} \alpha_{ij} dx^{i \wedge j}$ , we have

$$\alpha \left( \frac{\partial}{\partial x^k}, \frac{\partial}{\partial x^\ell} \right) = \frac{1}{2} \alpha_{ij} (dx^{i \otimes j} - dx^{j \otimes i}) \left( \frac{\partial}{\partial x^k}, \frac{\partial}{\partial x^\ell} \right) = \frac{1}{2} \alpha_{ij} (\delta_k^i \delta_\ell^j - \delta_k^j \delta_\ell^i) = \frac{1}{2} (\alpha_{k\ell} - \alpha_{\ell k}) = \alpha_{k\ell}$$

which justifies the notion  $\alpha_{k\ell}$  as  $\alpha(\partial/\partial x^k, \partial/\partial x^\ell)$ . In general, for any  $k$ -form  $\alpha = \frac{1}{k!} \alpha_{i_1 \dots i_k} dx^{i_1 \wedge \dots \wedge i_k}$  we have  $\alpha_{i_1 \dots i_k} = \alpha(\partial/\partial x^{i_1}, \dots, \partial/\partial x^{i_k})$ , because  $dx^{i_1 \wedge \dots \wedge i_k} = \sum_{\sigma \in \mathfrak{S}_k} \text{sgn}(\sigma) dx^{i_{\sigma(1)} \otimes \dots \otimes i_{\sigma(k)}}$ .

The norm of  $A \in \otimes^2(\mathcal{M})$  is defined by  $\|A\|_g^2 := \langle A, A \rangle_g = A_{ij}A^{ij}$ , while the norm of  $\alpha \in \wedge^k(\mathcal{M})$  is  $|\alpha|_g^2 := \langle \alpha, \alpha \rangle_g = \frac{1}{k!} \alpha_{i_1 \dots i_k} \alpha^{i_1 \dots i_k}$ . In particular,  $\|X\|_g^2 = X_i X^i = |b_g(X)|_g^2$  and  $\|\alpha\|_g^2 = 2|\alpha|_g^2$ , for any vector field  $X \in \mathfrak{X}(\mathcal{M})$  and 2-form  $\alpha$ .

The Levi-Civita connection associated to a given Riemannian metric  $g$  is denoted by  ${}^g\nabla$  or simply  $\nabla$ . Our convention on Riemann curvature tensor is  $R_{ijk}^m \frac{\partial}{\partial x^m} := \text{Rm}(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}) \frac{\partial}{\partial x^k} = (\nabla_i \nabla_j - \nabla_j \nabla_i) \frac{\partial}{\partial x^k}$  and  $R_{ijkl} := R_{ijk}^m g_{m\ell}$ . The Ricci curvature of  $g$  is given by  $R_{jk} := R_{ijk\ell} g^{i\ell}$ . We use  $dV_g$  and  $*_g$  to denote the volume form and Hodge star operator, respectively, on  $\mathcal{M}$  associated to a metric  $g$  and an orientation.

We use the standard notion  $A * B$  to denote some linear combination of contractions of the tensor product  $A \otimes B$  relative to the metric  $g_t$  associated the  $\varphi_t$ . In Theorem 1.4 and its proof, all universal constants  $c, C$  below depend only on the given real number  $p$ .

**1.2. Main results.** Applying De Turck's trick and Hamilton's Nash-Moser inverse function theorem, Bryant and Xu [2] proved the following local time existence for (1.1).

**Theorem 1.1. (Bryant-Xu [2])** *For a compact 7-manifold  $\mathcal{M}$ , the initial value problem (1.1) has a unique solution for a short time interval  $[0, T_{\max})$  with the maximal time  $T_{\max} \in (0, \infty]$  depending on  $\varphi$ .*

As in the Ricci flow, we can prove following results on the long time existence for the Laplacian flow (1.1).

**Theorem 1.2. (Lotay-Wei [33])** *Let  $\mathcal{M}$  be a compact 7-manifold and  $\varphi_t, t \in [0, T)$ , where  $T < \infty$ , be a solution to the flow (1.1) for closed  $G_2$ -structures with associated metric  $g_t = g_{\varphi_t}$  for each  $t$ .*

(a) *If the velocity of the flow satisfies*

$$\sup_{\mathcal{M} \times [0, T)} \|\Delta_t \varphi_t\|_t < \infty,$$

*then the solution  $\varphi_t$  can be extended past time  $T$ .*

(b) *If  $T = T_{\max}$ , then*

$$\lim_{t \rightarrow T_{\max}} \sup_{\mathcal{M}} \left( \|\text{Rm}_t\|_t^2 + \|\nabla_t \mathbf{T}_t\|_t^2 \right) = \infty.$$

*Here  $\mathbf{T}_t$  is the torsion of  $\varphi_t$  (see (2.14)).*

In this paper, we give a new elementary proof of Theorem 1.2, based on the idea of [24] and the structure of the equation (1.1).

**Theorem 1.3.** *Let  $\mathcal{M}$  be a compact 7-manifold and  $\varphi_t, t \in [0, T)$ , where  $T < \infty$ , be a solution to the flow (1.1) for closed  $G_2$ -structures with associated metric  $g_t = g_{\varphi_t}$  for each  $t$ . Suppose that*

$$K := \sup_{\mathcal{M} \times [0, T)} \|\text{Ric}_t\|_t < \infty, \quad \Lambda := \sup_{\mathcal{M}} \|\text{Rm}_0\|_0.$$

Then

$$\sup_{\mathcal{M} \times [0, T]} \|\text{Rm}_t\|_t < \infty,$$

where the bound depends only on  $n, K, T$  and  $\Lambda$ .

When  $\mathcal{M}$  is compact, the theorem immediately implies the part (a) in Theorem 1.2. Indeed, we shall show that (see (3.18) and (3.37))

$$\sup_{\mathcal{M} \times [0, T]} \|\Delta_t \varphi_t\|_t < \infty \iff \sup_{\mathcal{M} \times [0, T]} \|\text{Ric}_t\|_t < \infty.$$

In the compact case, Theorem 1.3 shows that, if the conclusion in part (a) does not hold, then  $T = T_{\max}$  and  $\sup_{\mathcal{M} \times [0, T_{\max})} \|\text{Rm}_t\|_t < \infty$  which implies  $\sup_{\mathcal{M} \times [0, T_{\max})} (\|\text{Rm}_t\|_t^2 + \|\nabla_t T_t\|_t^2) < \infty$ , since the norm  $\|\nabla_t T_t\|_t^2$  can be controlled by  $\|\text{Rm}_t\|_t^2$  (see (3.63)). However, by part (b) in Theorem 1.2, it is impossible. Therefore, the conclusion in part (a) is true.

As remarked in [24], to prove Theorem 1.3, it suffices to establish the following integral estimate.

**Theorem 1.4.** *Let  $\mathcal{M}$  be a smooth 7-manifold and  $\varphi_t, t \in [0, T)$ , where  $T < \infty$ , be a solution to the flow (1.1) for closed  $G_2$ -structures with associated metric  $g_t = g_{\varphi_t}$  for each  $t$ . Assume that there exist constants  $A, K > 0$  and a point  $x_0 \in \mathcal{M}$  such that the geodesic ball  $B_{g_0}(x_0, A/\sqrt{K})$  is compactly contained in  $\mathcal{M}$  and that*

$$|\text{Ric}_t|_t \leq K \quad \text{on } B_{g_0}\left(x_0, \frac{A}{\sqrt{K}}\right) \times [0, T].$$

Then, for any  $p \geq 5$ , there exists  $c = c(p) > 0$  so that

$$(1.3) \quad \int_{B_{g_0}(x_0, A/2\sqrt{K})} \|\text{Rm}_t\|_t^p dV_t \leq c(1+K)e^{cKT} \int_{B_{g_0}(x_0, A/\sqrt{K})} \|\text{Rm}_0\|_0^p dV_0 + cK^p \left(1 + A^{-2p}\right) e^{cKT} \text{vol}_t \left(B_{g_0}\left(x_0, \frac{A}{\sqrt{K}}\right)\right)$$

for all  $t \in [0, T]$ .

Now by the standard De Giorgi-Nash-Moser iteration (our manifold is compact and the Ricci curvature is uniformly bounded), under the condition in Theorem 1.4, we can prove

$$(1.4) \quad \|\text{Rm}_T\|_T(x_0) \leq d_1(d_2 + \Lambda_0),$$

where  $d_1, d_2$  are constants depending on  $K, T, A$ , and

$$\Lambda_0 := \sup_{B_{g_0}(x_0, A/\sqrt{K})} \|\text{Rm}_0\|_0.$$

Actually, this follows from the same argument in [24] by noting that

$$(1.5) \quad (\Delta_t - \partial_t) \|\text{Rm}_t\|_t \geq -c \|\text{Rm}_t\|_t^2.$$

To verify (1.5), we use (2.26), (3.61) and (3.65) to deduce that  $\|\nabla_t T_t\|_t \leq c \|\text{Rm}_t\|_t$  and

$$\|\nabla_t^2 T_t\|_t \leq c \|\nabla_t \text{Rm}_t\|_t + c \|\text{Rm}_t\|_t^{3/2}.$$

Then, by (3.31) and the Cauchy inequality

$$\begin{aligned} \|\nabla_t \mathbf{Rm}_t\|_t^2 &\leq -\frac{1}{2}(\partial_t - \Delta_t)\|\mathbf{Rm}_t\|_t^2 + c\|\mathbf{Rm}_t\|_t^3 + c\|\mathbf{Rm}_t\|_t^{3/2}\|\nabla_t \mathbf{Rm}_t\|_t \\ &\leq -\frac{1}{2}(\partial_t - \Delta_t)\|\mathbf{Rm}_t\|_t^2 + c\|\mathbf{Rm}_t\|_t^3 + \|\nabla_t \mathbf{Rm}_t\|_t^2 \end{aligned}$$

which implies (1.5). Now the estimate (1.4) yields Theorem 1.3.

The analogue of Theorem 1.2 in the Ricci flow was proved by Hamilton [16] (for part (b)) and Sesum [36] (for part (a)). It is an open question (due to Hamilton, see [3]) that the Ricci flow will exist as long as the scalar curvature remains bounded. For the Kähler-Ricci flow [39] or type-I Ricci flow [10], this question was settled. For the general case, some partial result on Hamilton's conjecture was carried out in [3].

For the Ricci-harmonic flow introduced by List [29, 30] (see also, [31, 32]), the analogue of Theorem 1.2 was proved in [29, 30] (see also, [31, 32]) and [4] (see [27] for another proof). The author [25, 26] extended Cao's result [3] to the Ricci-harmonic flow. The same Hamilton's conjecture was asked by the author in [25, 26].

We can ask the same question for the Laplacian flow on closed  $G_2$ -structures. In [33] (see Page 171, line -6 to -3, or Open Problem (3) in Page 230), Lotay and Wei asked that whether the Laplacian flow on closed  $G_2$ -structures will exist as long as the torsion tensor or scalar curvature remains bounded. Let  $g_t$  be the associated metric of  $\varphi_t$ . Then the evolution equation for  $g_t$  is given by

$$(1.6) \quad \partial_t g_{ij} = -2R_{ij} - \frac{4}{3}|T_t|_t^2 g_{ij} - 4T_i^k T_{kj}.$$

For the Laplacian flow on closed  $G_2$ -structures, the torsion  $T_t$  is actually a 2-form for each  $t$ , hence we use the norm  $|\cdot|_t$  in (1.6). The standard formula for the scalar curvature  $R_t$  gives (see (3.23))

$$(1.7) \quad \partial_t R_t = \Delta_t R_t + 2\|\mathbf{Ric}_t\|_t^2 - \frac{2}{3}R_t^2 + 4R_{ijkl}T^{ik}T^{j\ell} + 4(\nabla^j T^{ik})(\nabla_i T_{jk}).$$

Now the above mentioned open problem states that

$$\text{Is it true that } \lim_{t \rightarrow T_{\max}} R_t = -\infty?$$

The "minus infinity" comes from the fact that along the Laplacian flow on closed  $G_2$ -structures the scalar curvature is always nonpositive (see (2.26)). The following Proposition 1.5 is motivated to solve this problem, and starts from the basic evolution equation (1.7) where the last two terms on the right-hand side do not have good signature. However, using the closedness of  $\varphi_t$  (in particular, the identity (3.23)), we can prove the following interesting evolution equation for  $R_t$ .

**Proposition 1.5.** *Let  $\mathcal{M}$  be a smooth 7-manifold and  $\varphi_t$ ,  $t \in [0, T)$ , where  $T \in (0, \infty]$ , be a solution to the flow (1.1) for closed  $G_2$ -structures with associated metric  $g_t = g_{\varphi_t}$  for*

each  $t$ . Then the scalar curvature  $R_t$  satisfies

$$\begin{aligned}
\partial_t R_t &= \Delta_t R_t + \left\{ 2 \left\| R_{ij} + \frac{2}{3} |\mathbf{T}_t|_t^2 g_{ij} \right\|_t^2 + \frac{1}{2} \left\| R_{ijab} R^{ijmn} - \psi_{abmn} \right\|_t^2 \right. \\
&\quad + \frac{1}{2} \left\| 2\mathbf{T}_{ia} \mathbf{T}_{jb} R^{ijmn} - \psi_{abmn} \right\|_t^2 + \frac{1}{2} \left\| 2\widehat{\mathbf{T}}_{am} \widehat{\mathbf{T}}_{bn} - \psi_{abmn} \right\|_t^2 \\
(1.8) \quad &\quad + 2 \left\| \widehat{\mathbf{T}}_t \right\|_t^2 + 4 \left\| \nabla_t \mathbf{T}_t \right\|_t^2 \left. \right\} - \left\{ \|\mathrm{Rm}_t\|_t^2 + \frac{26}{9} R_t^2 + \frac{1}{2} \left\| R_{ijab} R^{ijmn} \right\|_t^2 \right. \\
&\quad \left. + 2 \left\| \mathbf{T}_{ia} \mathbf{T}_{jb} R^{ijmn} \right\|_t^2 + 2 \left\| \widehat{\mathbf{T}}_t \right\|_t^4 + 210 \right\}.
\end{aligned}$$

Here  $\widehat{\mathbf{T}}_{ij} = \mathbf{T}_i^k \mathbf{T}_{kj}$ .

Observe that the above well-arranged evolution equation can give us a weakly lower bound for  $R_t$ , which can not prove or disprove the conjecture of Lotay and Wei.

We give an outline of the current paper. We review the basic theory in Section 2 about  $G_2$ -structures,  $G_2$ -decompositions of 2-forms and 3-forms, and general flows on  $G_2$ -structures. In Section 3, we rewrite results in Section 2 for closed  $G_2$ -structures, and the local curvature estimates will be given in the last subsection.

**1.3. Acknowledgments.** The author is supported in part by the Fonds National de la Recherche Luxembourg (FNR) under the OPEN scheme (project GEOMREV O14/7628746).

The main result was carried out during the Young Geometric Analysts Forum 2018, 29th January – 2th February, in Tsinghua Sanya International Mathematics Forum.

The author, together with other six friends, thanks Yunhui Wu who personally provided us 14, the dimension of  $G_2$ , very fresh coconuts during the forum.

The author thanks Joel Fine, Brett Kotschwar, Chengjian Yao, Yong Wei, and Anton Thalmaier for useful discussion on the Laplacian flows and the earlier version of this paper. He also thanks Jason Lotay for his interested in this paper.

## 2. BASIC THEORY OF $G_2$ -STRUCTURES

In this section, we view some basic theory of  $G_2$ -structures, following [1, 19, 20, 21, 22, 33]. Let  $\{e_1, \dots, e_7\}$  denote the standard basis of  $\mathbb{R}^7$  and let  $\{e^1, \dots, e^7\}$  be its dual basis. Define the 3-form

$$\phi := e^{1\wedge 2\wedge 3} + e^{1\wedge 4\wedge 5} + e^{1\wedge 6\wedge 7} + e^{2\wedge 4\wedge 6} - e^{2\wedge 5\wedge 7} - e^{3\wedge 4\wedge 7} - e^{3\wedge 5\wedge 6},$$

where  $e^{i\wedge j\wedge k} := e^i \wedge e^j \wedge e^k$ . The subgroup  $G_2$ , which fixes  $\phi$ , of  $\mathrm{GL}(7, \mathbb{R})$  is the 14-dimensional Lie subgroup of  $\mathrm{SO}(7)$ , acts irreducibly on  $\mathbb{R}^7$ , and preserves the metric and orientation for which  $\{e_1, \dots, e_7\}$  is an oriented orthonormal basis. Note that  $G_2$  also preserves the 4-form

$$*_\phi \phi = e^{4\wedge 5\wedge 6\wedge 7} + e^{2\wedge 3\wedge 6\wedge 7} + e^{2\wedge 3\wedge 4\wedge 5} + e^{1\wedge 3\wedge 5\wedge 7} - e^{1\wedge 3\wedge 4\wedge 6} - e^{1\wedge 2\wedge 5\wedge 6} - e^{1\wedge 2\wedge 4\wedge 7}.$$

where the Hodge star operator  $*_\phi$  is determined by the metric and orientation.

For a smooth 7-manifold  $\mathcal{M}$  and a point  $x \in \mathcal{M}$ , define as in [33]

$$\Lambda_+^3(T_x^*\mathcal{M}) := \left\{ \varphi_x \in \Lambda^3(T_x^*\mathcal{M}) : \begin{array}{l} \mathbf{u}^*\varphi = \varphi_x \text{ for some invertible} \\ \text{map } \mathbf{u} \in \text{Hom}_{\mathbb{R}}(T_x\mathcal{M}, \mathbb{R}^7) \end{array} \right\}$$

and the bundle

$$\Lambda_+^3(T^*\mathcal{M}) := \bigsqcup_{x \in \mathcal{M}} \Lambda_+^3(T_x^*\mathcal{M}).$$

We call a section  $\varphi$  of  $\Lambda_+^3(T^*\mathcal{M})$  a *positive 3-form* on  $\mathcal{M}$  or a  $G_2$ -structure on  $\mathcal{M}$ , and denote the space of positive 3-forms by  $\Lambda_+^3(\mathcal{M})$ . The existence of  $G_2$ -structures is equivalent to the property that  $\mathcal{M}$  is oriented and spin, which is equivalent to the vanishing of the first and second Stiefel-Whitney classes. From the definition of  $G_2$ -structures, we see that any  $\varphi \in \Lambda_+^3(\mathcal{M})$  uniquely determines a Riemannian metric  $g_\varphi$  and an orientation  $dV_\varphi$ , hence the Hodge star operator  $*_\varphi$  and the associated 4-form

$$(2.1) \quad \psi := *_\varphi \varphi.$$

We also have the isomorphisms  $\flat_\varphi := \flat_{g_\varphi}$  and  $\sharp_\varphi := \sharp_{g_\varphi}$ . For a given  $G_2$ -structure  $\varphi \in \Lambda_+^3(\mathcal{M})$ , we denote by  $\langle \cdot, \cdot \rangle_\varphi$ ,  $\langle\langle \cdot, \cdot \rangle\rangle$ ,  $|\cdot|_\varphi$ ,  $\|\cdot\|_\varphi$ , the corresponding inner products  $\langle \cdot, \cdot \rangle_{g_\varphi}$ ,  $\langle\langle \cdot, \cdot \rangle\rangle_{g_\varphi}$  and norms  $|\cdot|_{g_\varphi}$ ,  $\|\cdot\|_{g_\varphi}$ .

Given a  $G_2$ -structure  $\varphi \in \Lambda_+^3(\mathcal{M})$ . We say that  $\varphi$  is *torsion-free* if  $\varphi$  is parallel with respect to the metric  $g_\varphi$ . Equivalently,  $\varphi$  is torsion-free if and only if  ${}^\varphi\nabla\varphi = 0$ , where  ${}^\varphi\nabla$  is the Levi-Civita connection of  $g_\varphi$ .

**Theorem 2.1. (Fernández-Gray [11])** *The  $G_2$ -structure  $\varphi$  is torsion-free if and only if  $\varphi$  is both closed (i.e.,  $d\varphi = 0$ ) and co-closed (i.e.,  $d*_\varphi\varphi = d\psi = 0$ ).*

When  $\mathcal{M}$  is compact, the above theorem says that a  $G_2$ -structure  $\varphi$  is torsion-free if and only if  $\varphi$  is harmonic with respect to the induces metric  $g_\varphi$ .

We say that a  $G_2$ -structure  $\varphi$  is *closed* (resp., *co-closed*) if  $d\varphi = 0$  (resp.,  $d\psi = 0$ ). Theorem 2.1 can be restated as that a  $G_2$ -structure is torsion-free if and only if it is both closed and co-closed.

**2.1.  $G_2$ -decompositions of  $\Lambda^2(\mathcal{M})$  and  $\Lambda^3(\mathcal{M})$ .** A  $G_2$ -structure  $\varphi$  induces splittings of the bundles  $\Lambda^k(T^*\mathcal{M})$ ,  $2 \leq k \leq 5$ , into direct summands, which we denote by  $\Lambda_\ell^k(T^*\mathcal{M}, \varphi)$  with  $\ell$  being the rank of the bundle. We let the space of sections of  $\Lambda_\ell^k(T^*\mathcal{M}, \varphi)$  by  $\Lambda_\ell^k(\mathcal{M}, \varphi)$ . Define the natural projections

$$(2.2) \quad \pi_\ell^k : \Lambda^k(\mathcal{M}) \longrightarrow \Lambda_\ell^k(\mathcal{M}, \varphi), \quad \alpha \longmapsto \pi_\ell^k(\alpha).$$

We mainly focus on the  $G_2$ -decompositions of  $\Lambda^2(\mathcal{M})$  and  $\Lambda^3(\mathcal{M})$ . Recall that

$$(2.3) \quad \Lambda^2(\mathcal{M}) = \Lambda_7^2(\mathcal{M}, \varphi) \oplus \Lambda_{14}^2(\mathcal{M}, \varphi),$$

$$(2.4) \quad \Lambda^3(\mathcal{M}) = \Lambda_1^3(\mathcal{M}, \varphi) \oplus \Lambda_7^3(\mathcal{M}, \varphi) \oplus \Lambda_{27}^3(\mathcal{M}, \varphi).$$

Here each component is determined by

$$\begin{aligned}\wedge_7^2(\mathcal{M}, \varphi) &= \{X \lrcorner \varphi : X \in \mathfrak{X}(\mathcal{M})\} = \{\beta \in \wedge^2(\mathcal{M}) : *_{\varphi}(\varphi \wedge \beta) = 2\beta\}, \\ \wedge_{14}^2(\mathcal{M}, \varphi) &= \{\beta \in \wedge^2(\mathcal{M}) : \psi \wedge \beta = 0\} = \{\beta \in \wedge^2(\mathcal{M}) : *_{\varphi}(\varphi \wedge \beta) = -\beta\}, \\ \wedge_1^3(\mathcal{M}, \varphi) &= \{f\varphi : f \in C^{\infty}(\mathcal{M})\}, \\ \wedge_7^3(\mathcal{M}, \varphi) &= \{*_\varphi(\varphi \wedge \alpha) : \alpha \in \wedge^1(\mathcal{M})\} = \{X \lrcorner \psi : X \in \mathfrak{X}(\mathcal{M})\}, \\ \wedge_{27}^3(\mathcal{M}, \varphi) &= \{\eta \in \wedge^3(\mathcal{M}) : \eta \wedge \varphi = \eta \wedge \psi = 0\}.\end{aligned}$$

For any 2-form  $\beta = \frac{1}{2}\beta_{ij}dx^i \wedge dx^j \in \wedge^2(\mathcal{M})$ , its two components  $\pi_7^2(\beta)$  and  $\pi_{14}^2(\beta)$  are determined by

$$(2.5) \quad \pi_7^2(\beta) = \frac{\beta + *_\varphi(\varphi \wedge \beta)}{3} = \frac{1}{2} \left( \frac{1}{3}\beta_{ab} + \frac{1}{6}\beta^{\ell m}\psi_{\ell mab} \right) dx^{ab},$$

$$(2.6) \quad \pi_{14}^2(\beta) = \frac{2\beta - *_\varphi(\varphi \wedge \beta)}{3} = \frac{1}{2} \left( \frac{2}{3}\beta_{ab} - \frac{1}{6}\beta^{\ell m}\psi_{\ell mab} \right) dx^{ab}.$$

To decompose 3-forms, recall two maps introduced by Bryant [1]

$$(2.7) \quad i_{\varphi} : \odot^2(\mathcal{M}) \longrightarrow \wedge^3(\mathcal{M}), \quad j_{\varphi} : \wedge^3(\mathcal{M}) \longrightarrow \odot^2(\mathcal{M}),$$

where

$$\begin{aligned}i_{\varphi}(h) &:= h_{ij}g^{j\ell}dx^i \wedge \left( \frac{\partial}{\partial x^{\ell}} \lrcorner \varphi \right) = \frac{1}{2}h_{i\ell}\varphi^{\ell}_{jk}dx^{ijk} \\ (2.8) \quad &= \frac{1}{6} \left( h_{i\ell}\varphi^{\ell}_{jk} + h_{j\ell}\varphi^{\ell}_{ik} + h_{k\ell}\varphi^{\ell}_{ij} \right) dx^{ijk}, \quad h = h_{ij}dx^{ij} \in \odot^2(\mathcal{M}),\end{aligned}$$

and

$$(2.9) \quad (j_{\varphi}(\eta)) (X, Y) := *_\varphi((X \lrcorner \varphi) \wedge (Y \lrcorner \varphi) \wedge \eta).$$

Then  $i_{\varphi}$  is injective and is isomorphic onto  $\wedge_1^3(\mathcal{M}, \varphi) \oplus \wedge_{27}^3(\mathcal{M}, \varphi)$ , and  $j_{\varphi}$  is an isomorphism between  $\wedge_1^3(\mathcal{M}, \varphi) \oplus \wedge_{27}^3(\mathcal{M}, \varphi)$  and  $\odot^2(\mathcal{M})$ . Moreover, for any 3-form  $\eta \in \wedge^3(\mathcal{M})$ , we have

$$(2.10) \quad \eta = i_{\varphi}(h) + X \lrcorner \psi$$

for some symmetric 2-tensor  $h \in \odot^2(\mathcal{M})$  and vector field  $X \in \mathfrak{X}(\mathcal{M})$ . Then

$$\begin{aligned}\eta &= h_i^{\ell}dx^i \wedge \left( \frac{\partial}{\partial x^{\ell}} \lrcorner \varphi \right) + X^{\ell} \left( \frac{\partial}{\partial x^{\ell}} \lrcorner \psi \right) = \frac{1}{2}h_i^{\ell}\varphi_{\ell jk}dx^{ijk} + \frac{1}{6}X^{\ell}\psi_{\ell jk}dx^{ijk} \\ &= \frac{1}{6} \left( 3h_i^{\ell}\varphi_{\ell jk} + X^{\ell}\psi_{\ell jk} \right) dx^{ijk} = \frac{1}{6}\eta_{ijk}dx^{ijk}.\end{aligned}$$

Write  $h$  as  $h_{ij} = \hat{h}_{ij} + \frac{1}{7}\text{tr}_{\varphi}(h)g_{\varphi}$ , where  $\hat{h} \in \odot_0^2(\mathcal{M})$  is the trace-free part of  $h$ , one has

$$(2.11) \quad \eta = \underbrace{\frac{3}{7}(\text{tr}_{\varphi}(h))\varphi}_{\pi_1^3(\eta)} + \underbrace{\frac{1}{2}\hat{h}_i^{\ell}\varphi_{\ell jk}dx^{ijk}}_{\pi_{27}^3(\eta)} + \underbrace{\frac{1}{6}X^{\ell}\psi_{\ell jk}dx^{ijk}}_{\pi_7^3(\eta)}.$$



**2.2. The torsion tensors of a  $G_2$ -structure.** By Hodge duality we obtain the  $G_2$ -decompositions of 4-forms  $\wedge^4(\mathcal{M}) = \wedge_1^4(\mathcal{M}, \varphi) \oplus \wedge_7^4(\mathcal{M}, \varphi) \oplus \wedge_{27}^4(\mathcal{M}, \varphi)$  and 5-forms  $\wedge^5(\mathcal{M}) = \wedge_7^5(\mathcal{M}, \varphi) \oplus \wedge_{14}^5(\mathcal{M}, \varphi)$ , respectively. By definition, we can find forms  $\tau_0 \in C^\infty(\mathcal{M})$ ,  $\tau_1, \tilde{\tau}_1 \in \wedge^1(\mathcal{M})$ ,  $\tau_2 \in \wedge_{14}^2(\mathcal{M}, \varphi)$ , and  $\tau_3 \in \wedge_{27}^3(\mathcal{M}, \varphi)$  such that

$$(2.12) \quad d\varphi = \tau_0\psi + 3\tau_1 \wedge \varphi + *_\varphi\tau_3, \quad d\psi = 4\tilde{\tau}_1 \wedge \psi - *_\varphi\tau_2.$$

Since  $\tau_2 \in \wedge_{14}^2(\mathcal{M}, \varphi)$ , it follows that  $\tau_2 \wedge \varphi = -*_\varphi\tau_2$ . Then (2.12) can be written as in the sense of Bryant [1]

$$(2.13) \quad d\varphi = \tau_0\psi + 3\tau_1 \wedge \varphi + *_\varphi\tau_3, \quad d\psi = 4\tilde{\tau}_1 \wedge \psi + \tau_2 \wedge \varphi.$$

It can be proved that  $\tau_1 = \tilde{\tau}_1$  (see [22]). We call  $\tau_0$  the *scalar torsion*,  $\tau_1$  the *vector torsion*,  $\tau_2$  the *Lie algebra torsion*, and  $\tau_3$  the *symmetric traceless torsion*. We also call  $\tau_\varphi := \{\tau_0, \tau_1, \tau_2, \tau_3\}$  the *intrinsic torsion forms* of the  $G_2$ -structure  $\varphi$ .

Recall that a  $G_2$ -structure  $\varphi$  is torsion-free if and only if  $d\varphi = d\psi = 0$  by Theorem 2.1. From (2.12) we see that  $\varphi$  is torsion-free if and only if the intrinsic torsion forms  $\tau_\varphi \equiv 0$ ; that is,  $\tau_0 = \tau_1 = \tau_2 = \tau_3 = 0$ .

**Lemma 2.2. (Fernández-Gray, [11])** *For any  $X \in \mathfrak{X}(\mathcal{M})$ , the 3-form  $\nabla_X\varphi$  lies in the space  $\wedge_7^3(\mathcal{M}, \varphi)$ . Therefore the covariant derivative  $\nabla\varphi \in \wedge^1(\mathcal{M}) \otimes \wedge_7^3(\mathcal{M})$ .*

Consequently, there exists a 2-tensor  $T = T_{ij}dx^{i\otimes j}$ , called the *full torsion tensor*, such that

$$(2.14) \quad \nabla_\ell\varphi = T_\ell{}^n\psi_{nabc}.$$

Equivalently,

$$(2.15) \quad T_{\ell m} = \frac{1}{24}(\nabla_\ell\varphi_{abc})\psi_m{}^{abc}.$$

Write

$$(2.16) \quad \tau_1 = (\tau_1)_i dx^i \in \wedge^1(\mathcal{M}),$$

$$(2.17) \quad \tau_2 = \frac{1}{2}(\tau_2)_{ab} dx^{ab} \in \wedge_{14}^2(\mathcal{M}),$$

$$(2.18) \quad \tau_3 = \frac{1}{2}(\tau_3)_{i^\ell} \varphi_{\ell ij} dx^{ijk} \in \wedge_{27}^3(\mathcal{M}, \varphi).$$

The associated 2-tensor  $\tau_3 := (\tau_3)_{ij} dx^{i\otimes j}$  of  $\tau_3$  lies in the space  $\odot_0^2(\mathcal{M})$ . With this convenience, the full torsion tensor  $T_{\ell m}$  is determined by

$$(2.19) \quad T_{\ell m} = \frac{\tau_0}{4}g_{\ell m} - (\tau_3)_{\ell m} - (\sharp_\varphi(\tau_1)\lrcorner\varphi)_{\ell m} - \frac{1}{2}(\tau_2)_{\ell m}$$

or as 2-tensors,

$$(2.20) \quad T = \frac{\tau_0}{4}g_\varphi - \tau_3 - \sharp_\varphi(\tau_1)\lrcorner\varphi - \frac{1}{2}\tau_2.$$

Here the 2-form  $\sharp_\varphi(\tau_1)\lrcorner\varphi$  is defined by

$$\sharp_\varphi(\tau_1)\lrcorner\varphi = \frac{1}{2}(\sharp_\varphi(\tau_1)\lrcorner\varphi) dx^{a\wedge b} = \frac{1}{2}\left((\tau_1)_k \varphi^k{}_{ab}\right) dx^{a\wedge b}.$$

As an application, this gives another proof of Theorem 2.1.

For fixed indices  $i$  and  $j$ , set

$$(2.21) \quad R_{ij|k\ell} := R_{ijk\ell} \text{ is skew-symmetric in } k \text{ and } \ell,$$

where

$$(2.22) \quad R_{ij|\bullet\bullet} := \frac{1}{2}R_{ij|k\ell}dx^{k\ell} = \frac{1}{2}R_{ijk\ell}dx^{k\ell} \in \wedge^2(\mathcal{M}).$$

Then, according to (2.5) and (2.6)

$$R_{ijk\ell} = R_{ij|k\ell} = \left( \pi_7^2(R_{ij|\bullet\bullet}) \right)_{k\ell} + \left( \pi_{14}^2(R_{ij|\bullet\bullet}) \right)_{k\ell},$$

where

$$\begin{aligned} \left( \pi_7^2(R_{ij|\bullet\bullet}) \right)_{k\ell} &= \frac{1}{3}R_{ij|k\ell} + \frac{1}{6}R_{ij|ab}\psi^{ab}{}_{k\ell} = \frac{1}{3}R_{ijk\ell} + \frac{1}{6}R_{ijab}\psi^{ab}{}_{k\ell}, \\ \left( \pi_{14}^2(R_{ij|\bullet\bullet}) \right)_{k\ell} &= \frac{2}{3}R_{ij|k\ell} - \frac{1}{6}R_{ij|ab}\psi^{ab}{}_{k\ell} = \frac{1}{3}R_{ijk\ell} - \frac{1}{6}R_{ijab}\psi^{ab}{}_{k\ell}. \end{aligned}$$

Karigiannis [22] (see also the equivalent formula obtained by Bryant in [1]) proved that the Ricci curvature is given by

$$\begin{aligned} R_{jk} &= R_{ijk\ell}g^{i\ell} = 3 \left( \pi_7^2(R_{ij|\bullet\bullet}) \right)_{k\ell} g^{i\ell} = \frac{3}{2} \left( \pi_{14}^2(R_{ij|\bullet\bullet}) \right)_{k\ell} g^{i\ell} \\ (2.23) \quad &= -(\nabla_i T_{jm} - \nabla_j T_{im}) \varphi^m{}_k{}^i - T_j{}^i T_{ik} + (\text{tr}_\varphi T) T_{jk} + T_{jb} T_{ia} \psi^{iab}{}_k, \\ &= -\nabla_i \left( T_j{}^n \varphi_{nk}{}^i \right) + \nabla_j \left( T_i{}^n \varphi_{nk}{}^i \right) - T_j{}^i T_{ik} + (\text{tr}_\varphi T) T_{jk} - T_{jb} T_{ia} \psi^{iab}{}_k. \end{aligned}$$

Cleyton and Ivanov [5] also derived a formula for the Ricci tensor for closed  $G_2$ -structures in terms of  $d_\varphi^* \varphi$ . Taking the trace of (2.23), we obtain Bryant's formula [1] for the scalar curvature

$$\begin{aligned} R &= -12\nabla^\ell(\tau_1)_\ell + \frac{21}{8}\tau_0^2 - \|\tau_3\|_\varphi^2 + 5\|\#_\varphi(\tau_1) \lrcorner \varphi\|_\varphi^2 - \frac{1}{4}\|\tau_2\|_\varphi^2, \\ (2.24) \quad &= -12\nabla^\ell(\tau_1)_\ell + \frac{21}{8}\tau_0^2 - \|\tau_3\|_\varphi^2 + 30|\tau_1|_\varphi^2 - \frac{1}{2}|\tau_2|_\varphi^2, \end{aligned}$$

For a closed  $G_2$ -structure, we have  $\tau_0 = \tau_1 = \tau_3 = 0$  and then  $R = -\frac{1}{4}\|\tau_2\|_\varphi^2 \leq 0$ . On the other hand, we have  $(\tau_2)_{ij} = -2T_{ij}$  by (2.20). Thus the full torsion tensor  $T$  is actually a 2-form

$$(2.25) \quad T = \frac{1}{2}T_{ij}dx^{ij} \in \wedge^2(\mathcal{M})$$

and the scalar curvature can be written in terms of  $T$

$$(2.26) \quad R = -\|T\|_\varphi^2 = -2|T|_\varphi^2 \leq 0.$$

Hence, for closed  $G_2$ -structures, scalar curvatures are always non-positive.

Finally, we mention a Bianchi type identity

$$(2.27) \quad \nabla_i T_{j\ell} - \nabla_j T_{i\ell} = -\frac{1}{2}R_{ijab}\varphi^{ab}{}_\ell - T_{ia} T_{jb} \varphi^{ab}{}_\ell = -\left( \frac{1}{2}R_{ijab} + T_{ia} T_{jb} \right) \varphi^{ab}{}_\ell.$$

The proof can be found in [22].

**2.3. General flows on  $G_2$ -structures.** For any family  $(\varphi_t)_t$  of  $G_2$ -structures, according to the decomposition (2.10), we can consider the general flow

$$(2.28) \quad \partial_t \varphi_t = i_{\varphi_t}(h_t) + X_t \lrcorner \psi_t$$

where  $h_t \in \odot^2(\mathcal{M})$  and  $X_t \in \mathfrak{X}(\mathcal{M})$ . The general flow (2.28) locally can be written as

$$(2.29) \quad \partial_t \varphi_{ijk} = h_i^\ell \varphi_{\ell jk} + h_j^\ell \varphi_{i \ell k} + h_k^\ell \varphi_{ij \ell} + X^\ell \psi_{\ell jk}.$$

We write for  $g_t$  and  $dV_t$  the metric and volume form associated to  $\varphi_t$ , respectively.

**Theorem 2.3.** *Under the general flow (2.28), we have*

$$(2.30) \quad \partial_t g_{ij} = 2h_{ij},$$

$$(2.31) \quad \partial_t g^{ij} = -2h^{ij},$$

$$(2.32) \quad \partial_t dV_t = (\text{tr}_t h_t) dV_t,$$

$$(2.33) \quad \partial_t T_{pq} = T_p^m h_{mq} - T_p^m X^k \varphi_{kmq} - (\nabla_k h_{ip}) \varphi^{ki}_q + \nabla_p X_q.$$

These evolution equations can be found in [22].

### 3. LAPLACIAN FLOWS ON CLOSED $G_2$ -STRUCTURES

We now consider the Laplacian flow for closed  $G_2$ -structures

$$(3.1) \quad \partial_t \varphi_t = \Delta_{\varphi_t} \varphi_t = \Delta_t \varphi_t, \quad \varphi_0 = \varphi,$$

where  $\Delta_{\varphi_t} \varphi_t = dd_{\varphi_t}^* \varphi_t + d_{\varphi_t}^* d\varphi_t$  is the Hodge Laplacian of  $g_{\varphi_t}$  and  $\varphi$  is an initial closed  $G_2$ -structure. The short time existence for (3.1) was proved by Bryant and Xu [2], see also Theorem 1.1.

A criterion for the long time existence for the Laplacian flow on compact manifolds was given in Theorem 1.2. In this section, we give a new elementary proof of Lotay-Wei's result in compact case.

**3.1. Basic theory of closed  $G_2$ -structures.** Let  $\wedge_{+,\bullet}^3(\mathcal{M}) \subset \wedge_+^3(\mathcal{M}, \varphi)$  be the set of all closed  $G_2$ -structures on  $\mathcal{M}$ . If  $\varphi \in \wedge_{+,\bullet}^3(\mathcal{M})$  is closed, i.e.,  $d\varphi = 0$ , then  $\tau_0, \tau_1, \tau_3$  are all zero, so the only nonzero torsion form is

$$(3.2) \quad \tau \equiv \tau_2 = \frac{1}{2}(\tau_2)_{ij} dx^{ij} = \frac{1}{2} \tau_{ij} dx^{ij}.$$

According to (2.20) and (2.25), we have  $T_{ij} = -\frac{1}{2} \tau_{ij}$  so that

$$(3.3) \quad T \equiv \frac{1}{2} T_{ij} dx^{ij} \quad \text{or equivalently} \quad T = -\frac{1}{2} \tau,$$

is a 2-form. Since  $d\psi = \tau \wedge \varphi = -*_\varphi \tau$ , we get  $d_\varphi^* \tau = *_\varphi d *_\varphi \tau = -*_\varphi d^2 \psi = 0$  which is given in local coordinates by

$$(3.4) \quad \nabla^i \tau_{ij} = 0$$

For a closed  $G_2$ -structure  $\varphi$ , according to (2.23), the Ricci curvature is given by (in this case  $T_{ij}$  is a 2-form)

$$R_{jk} = (\nabla_j T_{im} - \nabla_i T_{jm}) \varphi^m_k{}^i - T_j^i T_{ik} + T_{jb} T_{ia} \psi^{iab}{}_k.$$

Since  $\tau \in \wedge_{14}^2(\mathcal{M}, \varphi)$  and  $T_{ij} = -\frac{1}{2}\tau_{ij}$ , it follows from [33] (see page 179 – 180) that

$$(3.5) \quad (\nabla_j T_{im})\varphi^{m i} = 2T_j^\ell T_{\ell k}.$$

and therefore, for a closed  $G_2$ -structure  $\varphi$ , the Ricci curvature is given by

$$(3.6) \quad R_{jk} = -(\nabla_i T_{jm})\varphi_k^{im} - T_j^i T_{ik}.$$

Taking the trace of (3.6) yields (2.26). Moreover, the factor  $\nabla_i T_{jm}$  in (3.6) can be expressed as (see Proposition 2.4 in [33])

$$(3.7) \quad \begin{aligned} \nabla_i T_{jk} &= -\frac{1}{4}R_{ijmn}\varphi_k^{mn} - \frac{1}{4}R_{kjmn}\varphi_i^{mn} + \frac{1}{4}R_{ikmn}\varphi_j^{mn} \\ &\quad - \frac{1}{2}T_{im}T_{jn}\varphi_k^{mn} - \frac{1}{2}T_{km}T_{jn}\varphi_i^{mn} + \frac{1}{2}T_{im}T_{kn}\varphi_j^{mn}. \end{aligned}$$

If  $\varphi$  is a closed  $G_2$ -structure, Section 2.2 in [33] shows that  $\pi_7^3(\Delta_\varphi\varphi) = 0$  and hence, according to (2.10),

$$(3.8) \quad \Delta_\varphi\varphi = i_\varphi(h) \in \wedge_1^3(\mathcal{M}, \varphi) \oplus \wedge_{27}^3(\mathcal{M}, \varphi),$$

where

$$(3.9) \quad h_{ij} = \frac{1}{2}\nabla_m \tau_{ni}\varphi_j^{mn} - \frac{1}{6}|\tau|_\varphi^2 g_{ij} - \frac{1}{4}\tau_i^\ell \tau_{\ell j} = -R_{ij} - \frac{2}{3}|T|_\varphi^2 g_{ij} - 2T_i^k T_{kj}.$$

Here  $|T|_\varphi^2 = \frac{1}{2}T_{k\ell}T^{k\ell} = \frac{1}{2}\|T\|_\varphi^2$ .

**3.2. Evolution equations for closed  $G_2$ -structures.** Since the Laplacian flow (3.1) preserves the closedness of  $\varphi_t$ , it follows from (3.10) that we have

$$(3.10) \quad \Delta_{\varphi_t}\varphi_t = i_{\varphi_t}(h_t) \in \wedge_1^3(\mathcal{M}, \varphi_t) \oplus \wedge_{27}^3(\mathcal{M}, \varphi_t),$$

where

$$(3.11) \quad h_{ij} = -R_{ij} - \frac{2}{3}|T_t|_t^2 g_{ij} - 2T_i^k T_{kj}.$$

From Theorem 2.3, we see that the associated metric tensor  $g_t$  evolves by

$$(3.12) \quad \partial_t g_{ij} = 2h_{ij} = -2R_{ij} - \frac{4}{3}|T_t|_t^2 g_{ij} - 4T_i^k T_{kj}.$$

and the volume form  $dV_t$  evolves by

$$(3.13) \quad \partial_t dV_t = (\text{tr}_t h_t) dV_t = \left(-R - \frac{14}{3}|T_t|_t^2 + 4|T_t|_t^2\right) dV_t = \frac{4}{3}|T_t|_t^2 dV_t.$$

Hence, along the flow (3.1), the volume of  $g_t$  is nondecreasing.

Introduce the following notions

$$(3.14) \quad \blacksquare_t := \partial_t - \blacktriangle_t, \quad |\cdot|_t := |\cdot|_{\varphi_t}, \quad \Delta_t := \Delta_{\varphi_t},$$

where  $\blacktriangle_t := g^{ij}\nabla_i\nabla_j$  is the usual Laplacian of  $g_t$  and  $\Delta_t$  is the Hodge Laplacian of  $g_t$ , and also the 2-tensor  $S_{ic_t}$  with components

$$(3.15) \quad S_{ij} := R_{ij} + \frac{2}{3}|T_t|_t^2 g_{ij} + 2T_i^k T_{kj} = -h_{ij}.$$

Then the evolution equation (3.12) can be written as

$$(3.16) \quad \partial_t g_{ij} = -2S_{ij}.$$

Moreover, the trace of  $\text{Sic}_t$  is exactly the scalar curvature, up to a multiplying constant,

$$(3.17) \quad S_t := \text{tr}_t \text{Sic}_t = R_t + \frac{14}{3} |\mathbf{T}_t|_t^2 - 4 |\mathbf{T}_t|_t^2 = -\frac{4}{3} |\mathbf{T}_t|_t^2 = \frac{2}{3} R_t.$$

It was proved in [33] that

$$(3.18) \quad |\Delta_t \varphi_t|_t^2 = (\text{tr}_t h_t)^2 + 2 \|h_t\|_t^2 = \frac{16}{9} |\mathbf{T}_t|_t^4 + 2 \|\text{Sic}_t\|_t^2.$$

This identity together with (2.26) shows that the boundedness of  $\Delta_t \varphi_t$  is equivalent to the boundedness of  $\text{Ric}_t$ .

The evolution equation (2.33) implies that for the Laplacian flow on closed  $G_2$ -structures, the torsion  $T_{ij}$  evolves by evolves

$$(3.19) \quad \partial_t T_{ij} = T_i^k h_{kj} - (\nabla_m h_{ni}) \varphi_j^{mn}.$$

Furthermore, we can prove

**Proposition 3.1.** *Under the flow (3.1), we have*

$$(3.20) \quad \begin{aligned} \blacksquare_t T_{ij} &= 3R_j^k T_{ki} - R_i^k T_{kj} - \frac{1}{2} R_{ijmk} T^{mk} - \frac{1}{2} R_{mpi}{}^k R_{qk} \psi_j^{pqm} - \frac{2}{3} \varphi_j^i{}^m \nabla_m |\mathbf{T}_t|_t^2 \\ &+ \nabla_p T_{qi} \left( T^{pk} \varphi_{kj}{}^q - 2T^{qk} \varphi_{kj}{}^p \right) - \frac{2}{3} |\mathbf{T}_t|_t^2 T_{ij} - 4T_i^k T_k^m T_{mj}. \end{aligned}$$

*Proof.* See [33]. □

For a geometric flow  $\partial_t g_{ij} = \eta_{ij}$ , for some symmetric 2-tensor  $\eta_{ij}$ , we have

$$\begin{aligned} \partial R_{ijk}^\ell &= \frac{1}{2} g^{\ell p} \left( \nabla_i \nabla_j \eta_{kp} + \nabla_i \nabla_k \eta_{jp} - \nabla_i \nabla_p \eta_{jk} \right. \\ &\quad \left. - \nabla_j \nabla_i \eta_{kp} - \nabla_j \nabla_k \eta_{ip} + \nabla_j \nabla_p \eta_{ik} \right), \\ \partial_t R_{jk} &= \frac{1}{2} g^{pq} \left( \nabla_q \nabla_j \eta_{kp} + \nabla_q \nabla_k \eta_{jp} - \nabla_q \nabla_p \eta_{jk} - \nabla_j \nabla_k \eta_{qp} \right), \\ \partial_t R_t &= -\blacktriangle_t \text{tr}_t \eta_t + \text{div}_t(\text{div}_t \eta_t) - R_{ij} h^{ij}, \end{aligned}$$

where  $(\text{div}_t \eta_t)_j = \nabla^i \eta_{ij}$ . Applying those evolution equations to  $\eta_{ij} = -2R_{ij} - \frac{4}{3} |\mathbf{T}_t|_t^2 g_{ij} - 4T_i^k T_{kj} = -2S_{ij}$  we have

$$\begin{aligned} \text{tr}_t \eta_t &= -2R_t - \frac{28}{3} |\mathbf{T}_t|_t^2 + 8 |\mathbf{T}_t|_t^2 = \frac{8}{3} |\mathbf{T}_t|_t^2, \\ (\text{div}_t \eta_t)_j &= -2 \nabla^i R_{ij} - \frac{4}{3} \nabla_j |\mathbf{T}_t|_t^2 - 4 \nabla^i \widehat{T}_{ij} \\ &= -\nabla_j R_t - \frac{4}{3} \nabla_j |\mathbf{T}_t|_t^2 - 4 \nabla^i \widehat{T}_{ij}, \\ \text{div}_t(\text{div}_t \eta_t) &= \nabla^j (\text{div}_t \eta_t)_j = -\blacktriangle_t R_t - \frac{4}{3} \blacktriangle_t |\mathbf{T}_t|_t^2 - 4 \nabla^j \nabla^i \widehat{T}_{ij}, \end{aligned}$$

where the symmetric 2-tensor  $\widehat{T}$  is given by

$$(3.21) \quad \widehat{T}_{ij} := T_{ik} T^k{}_j.$$

Plugging those identities into the above evolution equation for  $R_t$ , we get

$$\begin{aligned}\partial_t R_t &= -4\blacktriangle_t |T_t|_t^2 - \blacktriangle_t R_t - 4\nabla^j \nabla^i \widehat{T}_{ij} - R^{ij} \left( -2R_{ij} - \frac{4}{3} |T_t|_t^2 g_{ij} - 4\widehat{T}_{ij} \right) \\ &= \blacktriangle_t R_t - 4\nabla^j \nabla^i \widehat{T}_{ij} + 2\|\text{Ric}_t\|_t^2 + \frac{4}{3} |T_t|_t^2 R_t + 4R^{ij} \widehat{T}_{ij}\end{aligned}$$

which implies

$$(3.22) \quad \blacksquare_t R_t = 2\|\text{Ric}_t\|_t^2 - \frac{2}{3} R_t^2 - 4\nabla^j \nabla^i \widehat{T}_{ij} + 4\langle \text{Ric}_t, \widehat{T} \rangle_t.$$

Observe that the last two terms on the right-hand side of (3.22) are not determined of their signs. In the following, we shall use the identity

$$(3.23) \quad \nabla^i T_{ij} = 0$$

follows from from (3.3) and (3.4), to simplify those two terms. Using the identity (3.23), the term  $\nabla^j \nabla^i \widehat{T}_{ij}$  can be simplified as follows.

$$\begin{aligned}\nabla^j \nabla^i \widehat{T}_{ij} &= \nabla^j \nabla^i (T_i^k T_{kj}) = \nabla^j \left[ (\nabla^i T_i^k) T_{kj} + T_i^k (\nabla^i T_{kj}) \right] \\ &= T^{ik} (\nabla_j \nabla_i T_k^j) - (\nabla^j T^{ik}) (\nabla_i T_{jk}).\end{aligned}$$

On the other hand, from the Ricci identity

$$\nabla_j \nabla_i T_k^j = \nabla_i \nabla_j T_k^j - R_{jikl} T_k^l - R_{jil} T_{kl} = R_{ijk\ell} T^{\ell j} + R_{i\ell} T_k^\ell,$$

we see that the evolution equation (3.22) is equivalent to

$$(3.24) \quad \blacksquare_t R_t = 2\|\text{Ric}_t\|_t^2 - \frac{2}{3} R_t^2 + 4R_{ijk\ell} T^{ik} T^{j\ell} + 4(\nabla^j T^{ik}) (\nabla_i T_{jk}).$$

From (3.15) and (3.21) we can rewrite the term  $\|\text{Ric}_t\|_t^2$  in (3.24) in terms of  $\text{Sic}_t$  according to the following relation:

$$\begin{aligned}\|\text{Sic}_t\|_t^2 &= \left( R_{ij} + \frac{2}{3} |T_t|_t^2 g_{ij} + 2\widehat{T}_{ij} \right) \left( R^{ij} + \frac{2}{3} |T_t|_t^2 g^{ij} + 2\widehat{T}^{ij} \right) \\ &= \|\text{Ric}_t\|_t^2 + \frac{4}{3} |T_t|_t^2 R_t + 4\langle \text{Ric}_t, \widehat{T}_t \rangle_t + \frac{28}{9} |T_t|_t^4 + \frac{8}{3} |T_t|_t^2 \text{tr}_t \widehat{T}_t + 4\|\widehat{T}_t\|_t^2 \\ &= \|\text{Ric}_t\|_t^2 - \frac{2}{3} R_t^2 + 4\langle \text{Ric}_t, \widehat{T}_t \rangle_t + \frac{7}{9} R_t^2 - \frac{4}{3} R_t^2 + 4\|\widehat{T}_t\|_t^2 \\ &= \|\text{Ric}_t\|_t^2 + 4\|\widehat{T}_t\|_t^2 + 4\langle \text{Ric}_t, \widehat{T}_t \rangle_t - \frac{11}{9} R_t^2,\end{aligned}$$

where we used  $\text{tr}_t \widehat{T}_t = g^{ij} T_{ik} T^k_j = T_{ik} T^{ki} = -2|T_t|_t^2$  and  $R_t = -2|T_t|_t^2$ . Replacing  $R_t$  by  $S_t$  according to the identity (3.17), we can rewrite (3.24) as

$$\begin{aligned}\blacksquare_t S_t &= \frac{4}{3} \|\text{Sic}_t\|_t^2 - \frac{16}{3} \|\widehat{T}_t\|_t^2 - \frac{16}{3} \langle \text{Ric}_t, \widehat{T}_t \rangle_t + \frac{32}{27} R_t^2 \\ &\quad + \frac{8}{3} R_{ijk\ell} T^{ik} T^{j\ell} + \frac{8}{3} (\nabla^j T^{ik}) (\nabla_i T_{jk}).\end{aligned}$$

Similarly, replacing  $\langle \text{Ric}_t, \widehat{T}_t \rangle_t$  by  $\langle \text{Sic}_t, \widehat{T}_t \rangle_t$  with respect to the identity

$$\langle \langle \text{Sic}_t, \widehat{T}_t \rangle \rangle = \left( R_{ij} + \frac{2}{3} |T_t|_t^2 g_{ij} + 2\widehat{T}_{ij} \right) \widehat{T}^{ij} = \langle \langle \text{Ric}_t, \widehat{T}_t \rangle \rangle_t - \frac{1}{3} R_t^2 + 2\|\widehat{T}_t\|_t^2,$$

we obtain the following evolution equation for  $S_t$ ,

$$(3.25) \quad \blacksquare_t S_t = \frac{4}{3} \left[ \left\| \text{Si}c_t - 2\widehat{T}_t \right\|_t^2 - S_t^2 \right] + \frac{8}{3} \left[ R_{ijkl} T^{ik} T^{j\ell} + (\nabla^j T^{ik})(\nabla_i T_{jk}) \right].$$

Next, we try to deal with the last bracket in (3.25), which contains two terms  $R_{ijkl} T^{ik} T^{j\ell}$  and  $(\nabla^j T^{ik})(\nabla_i T_{jk})$ . Using (2.27) and (3.7), the term  $(\nabla^j T^{ik})(\nabla_i T_{jk})$  is equal to

$$\begin{aligned} (\nabla^j T^{ik})(\nabla_i T_{jk}) &= \left[ \nabla^i T^{jk} + \left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \varphi^{kab} \right] \nabla_i T_{jk} \\ &= \|\nabla_t T_t\|_t^2 + \frac{1}{2} \left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left[ -\frac{1}{2} R_{ijmn} \varphi^{mn}{}_k \varphi^{kab} - \frac{1}{2} R_{kjmn} \varphi_i{}^{mn} \varphi^{kab} \right. \\ &\quad \left. + \frac{1}{2} R_{ikmn} \varphi_j{}^{mn} \varphi^{kab} - T_{im} T_{jn} \varphi^{mn}{}_k \varphi^{kab} - T_{km} T_{jn} \varphi_i{}^{mn} \varphi^{kab} + T_{im} T_{kn} \varphi_j{}^{mn} \varphi^{kab} \right]. \end{aligned}$$

By symmetry the term

$$\left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left( -\frac{1}{2} R_{kjmn} \varphi_i{}^{mn} \varphi^{kab} + \frac{1}{2} R_{ikmn} \varphi_j{}^{mn} \varphi^{kab} \right)$$

is equal to, interchanging  $i \leftrightarrow j$  and  $a \leftrightarrow b$  in the second term,

$$\left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left( -\frac{1}{2} R_{kjmn} \varphi_i{}^{mn} \varphi^{kab} \right) + \left( \frac{1}{2} R^{ji}{}_{ba} + T^j{}_b T^i{}_a \right) \left( \frac{1}{2} R_{jkmn} \varphi_i{}^{mn} \varphi^{kba} \right)$$

which is zero. Similarly, we have, by interchanging  $m \leftrightarrow n$  and then  $i \leftrightarrow j$ ,  $a \leftrightarrow b$  in the first term,

$$\begin{aligned} &\left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left( -T_{km} T_{jn} \varphi_i{}^{mn} \varphi^{kab} + T_{im} T_{kn} \varphi_j{}^{mn} \varphi^{kab} \right) \\ &= \left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left( -T_{kn} T_{jm} \varphi_i{}^{nm} \varphi^{kab} + T_{im} T_{kn} \varphi_j{}^{mn} \varphi^{kab} \right) \\ &= \left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left( -T_{kn} T_{im} \varphi_j{}^{nm} \varphi^{kba} + T_{im} T_{kn} \varphi_j{}^{mn} \varphi^{kab} \right) = 0. \end{aligned}$$

Therefore, using the identity  $\varphi_{ijk} \varphi^{kab} = g_{ia} g_{jb} - g_{ib} g_{ja} + \psi_{ijab}$  (see [22]), we arrive at

$$\begin{aligned} (\nabla^j T^{ik})(\nabla_i T_{jk}) &= \|\nabla_t T_t\|_t^2 - \frac{1}{2} \left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left( \frac{1}{2} R_{ij}{}^{mn} + T_i{}^m T_j{}^n \right) \varphi_{mnk} \varphi^{kab} \\ &= \|\nabla_t T_t\|_t^2 - \frac{1}{2} \left( \frac{1}{2} R^{ij}{}_{ab} + T^i{}_a T^j{}_b \right) \left( \frac{1}{2} R_{ij}{}^{mn} + T_i{}^m T_j{}^n \right) \left( \delta_m^a \delta_n^b - \delta_m^b \delta_n^a + \psi_{mn}{}^{ab} \right) \\ &= \|\nabla_t T_t\|_t^2 - \frac{1}{8} \left( R_{ijab} + 2T_{ia} T_{jb} \right) \left[ \left( R^{ijab} + 2T^{ia} T^{jb} \right) \right. \\ &\quad \left. - \left( R^{ijba} + 2T^{ib} T^{ja} \right) + \left( R^{ijmn} + 2T^{im} T^{jn} \right) \psi_{mn}{}^{ab} \right]. \end{aligned}$$

Since, by our convention,

$$\begin{aligned} \left( R_{ijab} + 2T_{ia} T_{jb} \right) \left( R^{ijab} + 2T^{ia} T^{jb} \right) &= \|\text{Rm}_t\|_t^2 + 4R_{ijab} T^{ia} T^{jb} + 4\|\widehat{T}_t\|_t^4, \\ \left( R_{ijab} + 2T_{ia} T_{jb} \right) \left( R^{ijba} + 2T^{ib} T^{ja} \right) &= -\|\text{Rm}_t\|_t^2 - 4R_{ijab} T^{ia} T^{jb} + 4\|\widehat{T}_t\|_t^2, \end{aligned}$$

it follows that

$$\begin{aligned} (\nabla^j \mathbf{T}^{ik})(\nabla_i \mathbf{T}_{jk}) &= \|\nabla_t \mathbf{T}_t\|_t^2 + \frac{1}{8} \left[ -2\|\mathbf{Rm}_t\|_t^2 - 8R_{ijab} \mathbf{T}^{ia} \mathbf{T}^{jb} - 4\|\mathbf{T}_t\|_t^4 \right. \\ &\quad \left. + 4\|\widehat{\mathbf{T}}_t\|_t^2 - (R_{ijab} + 2\mathbf{T}_{ia} \mathbf{T}_{jb}) (R^{ijmn} + 2\mathbf{T}^{im} \mathbf{T}^{jn}) \psi_{mn}{}^{ab} \right] \end{aligned}$$

and (3.25) can be written as

$$\begin{aligned} \blacksquare_t S_t &= \frac{4}{3} \left\| \text{Sic}_t - 2\widehat{\mathbf{T}}_t \right\|_t^2 + \frac{8}{3} \|\nabla_t \mathbf{T}_t\|_t^2 + \frac{4}{3} \|\widehat{\mathbf{T}}_t\|_t^2 - \frac{2}{3} \|\mathbf{Rm}_t\|_t^2 - \frac{13}{3} S_t^2 \\ (3.26) \quad &- \frac{1}{3} (R_{ijab} + 2\mathbf{T}_{ia} \mathbf{T}_{jb}) (R^{ijmn} + 2\mathbf{T}^{im} \mathbf{T}^{jn}) \psi_{mn}{}^{ab}. \end{aligned}$$

Finally, we deal with the last term  $J$  on the right-hand side of (3.26). From the identity  $\psi_{ijkl} \psi^{ijkl} = 168$ , we find that

$$\begin{aligned} J &:= -\frac{1}{3} (R_{ijab} + 2\mathbf{T}_{ia} \mathbf{T}_{jb}) (R^{ijmn} + 2\mathbf{T}^{im} \mathbf{T}^{jn}) \psi_{mn}{}^{ab} \\ &= \frac{1}{3} \left( -R_{ij}{}^{ab} R^{ijmn} \psi_{mnab} - 4\mathbf{T}_i{}^a \mathbf{T}_j{}^b R^{ijmn} \psi_{mnab} - 4\mathbf{T}_i{}^a \mathbf{T}^{im} \mathbf{T}^b{}_j \mathbf{T}^{jn} \psi_{mnab} \right) \\ &= \frac{1}{3} \left[ \left\| R_{ij}{}^{ab} R^{ijmn} - \frac{1}{2} \psi^{abmn} \right\|_t^2 - \left\| R_{ij}{}^{ab} R^{ijmn} \right\|_t^2 - \frac{168}{4} \right. \\ &\quad \left. + \left\| 2\mathbf{T}_i{}^a \mathbf{T}_j{}^b R^{ijmn} - \psi^{abmn} \right\|_t^2 - 4 \left\| \mathbf{T}_i{}^a \mathbf{T}_j{}^b R^{ijmn} \right\|_t^2 - 168 \right. \\ &\quad \left. + \left\| 2\widehat{\mathbf{T}}^{am} \widehat{\mathbf{T}}^{bn} - \psi^{mnab} \right\|_t^2 - 4\|\widehat{\mathbf{T}}_t\|_t^4 - 168 \right]. \end{aligned}$$

Plugging the expression for  $J$  into (3.26), we obtain

**Proposition 3.2.** *The scalar curvature  $R_t$  or  $S_t$  evolves by*

$$\begin{aligned} \blacksquare_t S_t &= \frac{4}{3} \left\| \text{Sic}_t - 2\widehat{\mathbf{T}}_t \right\|_t^2 + \frac{8}{3} \|\nabla_t \mathbf{T}_t\|_t^2 + \frac{1}{3} \left\| R_{ijab} R^{ij}{}_{mn} - \psi_{abmn} \right\|_t^2 + \frac{4}{3} \|\widehat{\mathbf{T}}_t\|_t^2 \\ (3.27) \quad &+ \frac{1}{3} \left\| 2\mathbf{T}_{ia} \mathbf{T}_{jb} R^{ij}{}_{mn} - \psi_{abmn} \right\|_t^2 + \frac{1}{3} \left\| 2\widehat{\mathbf{T}}_{am} \widehat{\mathbf{T}}_{bn} - \psi_{abmn} \right\|_t^2 - \frac{4}{3} \|\widehat{\mathbf{T}}_t\|_t^4 \\ &- \frac{2}{3} \|\mathbf{Rm}_t\|_t^2 - \frac{13}{3} S_t^2 - \frac{1}{3} \left\| R_{ijab} R^{ij}{}_{mn} \right\|_t^2 - \frac{4}{3} \left\| \mathbf{T}_{ia} \mathbf{T}_{jb} R^{ij}{}_{mn} \right\|_t^2 - 126. \end{aligned}$$

Since  $S_t = \frac{2}{3} R_t$ , it follows from the above theorem that (1.6) holds true.

Before giving local curvature estimates for Laplacian flow in the next subsection, we derive evolution equations for  $\text{Ric}_t$ ,  $\mathbf{Rm}_t$ , and  $\mathbf{T}_t$  in different forms. Using the Lichnerowicz Laplacian

$$\blacktriangle_{L,t} \eta_{jk} := \blacktriangle_t \eta_{jk} - R_j{}^p \eta_{pk} - R_k{}^p \eta_{jp} + 2R_{pj}{}^q \eta^{qp},$$

we see that the evolution equation for  $R_{ij}$  can be written as

$$\partial_t R_{jk} = -\frac{1}{2} \left[ \blacktriangle_{L,t} \eta_{jk} + \nabla_j \nabla_k \text{tr}_t \eta_t + \nabla_j (d_t^* \eta_t)_k + \nabla_k (d_t^* \eta_t)_j \right],$$



where  $(d_t^* \eta_t)_k := -\nabla^j \eta_{jk}$ . For  $\eta_{ij} = -2R_{ij} - \frac{4}{3} \|\mathbf{T}_t\|_t^2 g_{ij} - 4\mathbf{T}_i^k \mathbf{T}_{kj}$  we have proved  $\text{tr}_t \eta_t = \frac{8}{3} \|\mathbf{T}_t\|_t^2$  and  $(d_t^* \eta_t)_j = \nabla_j R_t + \frac{4}{3} \nabla_j \|\mathbf{T}_t\|_t^2 + 4\nabla^i \widehat{\mathbf{T}}_{ij}$  with  $\widehat{\mathbf{T}}_{ij} = \mathbf{T}_i^k \mathbf{T}_{kj}$ . Then

$$\begin{aligned} \partial_t R_{jk} &= \blacktriangle_{L,t} \left( R_{jk} + \frac{2}{3} \|\mathbf{T}_t\|_t^2 g_{jk} + 2\widehat{\mathbf{T}}_{jk} \right) - \frac{1}{2} \nabla_j \left( \nabla_k R_t + \frac{4}{3} \nabla_k \|\mathbf{T}_t\|_t^2 + 4\nabla^i \widehat{\mathbf{T}}_{ik} \right) \\ &\quad - \frac{4}{3} \nabla_j \nabla_k \|\mathbf{T}_t\|_t^2 - \frac{1}{2} \nabla_k \left( \nabla_j R_t + \frac{4}{3} \nabla_j \|\mathbf{T}_t\|_t^2 + 4\nabla^i \widehat{\mathbf{T}}_{ij} \right) \\ &= \blacktriangle_{L,t} \left( R_{jk} + \frac{2}{3} \|\mathbf{T}_t\|_t^2 g_{jk} + 2\widehat{\mathbf{T}}_{jk} \right) - 2\nabla_j \nabla^i \widehat{\mathbf{T}}_{ik} - 2\nabla_k \nabla^i \widehat{\mathbf{T}}_{ij} - \frac{2}{3} \nabla_j \nabla_k \|\mathbf{T}_t\|_t^2. \end{aligned}$$

But the first term is equal to

$$\begin{aligned} \blacktriangle_{L,t} \left( R_{jk} + \frac{2}{3} \|\mathbf{T}_t\|_t^2 g_{jk} + 2\widehat{\mathbf{T}}_{jk} \right) &= \blacktriangle_t R_{jk} - 2R_j^p R_{pk} + 2R_{pj} R_{pk} \\ &\quad + \left[ \frac{2}{3} \left( \blacktriangle_t \|\mathbf{T}_t\|_t^2 \right) g_{jk} + 2\blacktriangle_t \widehat{\mathbf{T}}_{jk} - 2R_j^p \widehat{\mathbf{T}}_{pk} - 2\widehat{\mathbf{T}}_j^p R_{pk} + 4R_{pj} R_{pk} \widehat{\mathbf{T}}^{pq} \right], \end{aligned}$$

we have

$$\begin{aligned} \blacksquare_t R_{ij} &= -2R_i^p R_{pj} + 2R_{pij} R^{pq} + \left[ \frac{2}{3} \left( \blacktriangle_t \|\mathbf{T}_t\|_t^2 \right) g_{ij} + 2\blacktriangle_t \widehat{\mathbf{T}}_{ij} - 2R_i^p \widehat{\mathbf{T}}_{pj} \right. \\ (3.28) \quad &\quad \left. - 2\widehat{\mathbf{T}}_i^p R_{pj} + 4R_{pij} \widehat{\mathbf{T}}^{pq} - 2\nabla_i \nabla^p \widehat{\mathbf{T}}_{pj} - 2\nabla_j \nabla^p \widehat{\mathbf{T}}_{pi} - \frac{2}{3} \nabla_i \nabla_j \|\mathbf{T}_t\|_t^2 \right]. \end{aligned}$$

Consequently, the norm of  $\text{Ric}_t$  satisfies

$$\begin{aligned} \blacksquare_t \|\text{Ric}_t\|_t^2 &= -2\|\nabla_t \text{Ric}_t\|_t^2 + 4R_{kij\ell} R^{k\ell} R^{ij} + \left[ \frac{4}{3} R_t \blacktriangle_t \|\mathbf{T}_t\|_t^2 + 8R^{kj\ell} \widehat{\mathbf{T}}_{k\ell} R^{ij} + \right. \\ (3.29) \quad &\quad \left. \frac{8}{3} \|\text{Ric}_t\|_t^2 \|\mathbf{T}_t\|_t^2 + 4R^{ij} \blacktriangle_t \widehat{\mathbf{T}}_{ij} - 8R^{ij} \nabla_i \nabla^k \widehat{\mathbf{T}}_{kj} - \frac{4}{3} R^{ij} \nabla_i \nabla_j \|\mathbf{T}_t\|_t^2 \right]. \end{aligned}$$

The general formula for  $R_{ijk}^\ell$  gives

$$\begin{aligned} \partial_t R_{ijk}^\ell &= -\nabla_i \nabla_k R_j^\ell - \nabla_j \nabla^\ell R_{ik} + \nabla_i \nabla^\ell R_{jk} + \nabla_j \nabla_k R_i^\ell + R_{ijk}^q R_q^\ell + R_{ij}^{\ell q} R_{kp} \\ &\quad + 2R_{ijk}^q \widehat{\mathbf{T}}_q^\ell + 2R_{ij}^{\ell q} \widehat{\mathbf{T}}_{kp} - \frac{2}{3} \left( \nabla_i \nabla_k \|\mathbf{T}_t\|_t^2 \right) g_j^\ell - \frac{2}{3} \left( \nabla_j \nabla^\ell \|\mathbf{T}_t\|_t^2 \right) g_{ik} \\ (3.30) \quad &\quad + \frac{2}{3} \left( \nabla_i \nabla^\ell \|\mathbf{T}_t\|_t^2 \right) g_{jk} + \frac{2}{3} \left( \nabla_j \nabla_k \|\mathbf{T}_t\|_t^2 \right) g_i^\ell \\ &\quad - 2\nabla_i \nabla_k \widehat{\mathbf{T}}_j^\ell - 2\nabla_j \nabla^\ell \widehat{\mathbf{T}}_{ik} + 2\nabla_i \nabla^\ell \widehat{\mathbf{T}}_{jk} + 2\nabla_j \nabla_k \widehat{\mathbf{T}}_i^\ell. \end{aligned}$$

Hence, the evolution equation for  $\|\text{Rm}_t\|_t^2$  is given by

$$\begin{aligned} \partial_t \|\text{Rm}_t\|_t^2 &= \nabla_t^2 \text{Ric}_t * \text{Rm}_t + \text{Ric}_t * \text{Rm}_t * \text{Rm}_t + \text{Rm}_t * \text{Rm}_t * \widehat{\mathbf{T}}_t \\ (3.31) \quad &\quad + \text{Ric}_t * \nabla_t^2 \|\mathbf{T}_t\|_t^2 + \text{Rm}_t * \nabla_t^2 \widehat{\mathbf{T}}_t + \frac{8}{3} \|\mathbf{T}_t\|_t^2 \|\text{Rm}_t\|_t^2. \end{aligned}$$

Moreover, it was proved in [33] that

$$\begin{aligned} \|\nabla_t \text{Rm}_t\|_t^2 &\leq -\frac{1}{2} \blacksquare_t \|\text{Rm}_t\|_t^2 + C_1 \|\text{Rm}_t\|_t^3 \\ (3.32) \quad &\quad + C_1 \|\text{Rm}_t\|_t^{3/2} \|\nabla_t^2 \mathbf{T}_t\|_t + C_1 \|\text{Rm}_t\|_t \|\nabla_t \mathbf{T}_t\|_t^2 \end{aligned}$$

where  $C_1$  is some universal constant, and

$$(3.33) \quad \blacksquare_t \mathbf{T}_t = \text{Rm}_t * \mathbf{T}_t + \text{Rm}_t * \mathbf{T}_t * \psi_t + \nabla_t \mathbf{T}_t * \mathbf{T}_t * \varphi_t + \mathbf{T}_t * \mathbf{T}_t * \mathbf{T}_t.$$

Squaring (3.33) gives

$$(3.34) \quad \begin{aligned} \|\nabla_t \mathbf{T}_t\|_t^2 &\leq -\frac{1}{2} \blacksquare_t \|\mathbf{T}_t\|_t^2 + C_2 \|\text{Rm}_t\|_t \|\mathbf{T}_t\|_t^2 \\ &\quad + C_2 \|\nabla_t \mathbf{T}_t\|_t \|\mathbf{T}_t\|_t^2 + C_2 \|\mathbf{T}_t\|_t^4 \end{aligned}$$

for another universal constant  $C_2$  which may differ from  $C_1$ . The Cauchy-Schwartz inequality shows  $2C_2 \|\nabla_t \mathbf{T}_t\|_t \|\mathbf{T}_t\|_t^2 \leq \|\nabla_t \mathbf{T}_t\|_t^2 + C_2^2 \|\mathbf{T}_t\|_t^4$ , so that the evolution inequality (3.34) becomes

$$(3.35) \quad \|\nabla_t \mathbf{T}_t\|_t^2 \leq -\blacksquare_t \|\mathbf{T}_t\|_t^2 + C_3 \|\text{Rm}_t\|_t \|\mathbf{T}_t\|_t^2 + C_3 \|\mathbf{T}_t\|_t^4.$$

Here  $C_3$  is a universal constant.

**3.3. Local curvature estimates.** In this section, we consider the Laplacian flow (3.1) on  $\mathcal{M} \times [0, T]$ , where  $T \in (0, T_{\max})$ . From now on we always omit the time subscripts from all considered quantities. From (3.15), (3.29), (3.31), (3.32), and (3.35) we have

$$\begin{aligned} \|\nabla \text{Ric}\|^2 &= -\frac{1}{2} \blacksquare \|\text{Ric}\|^2 + \text{Ric} * \text{Ric} * \text{Rm} - \frac{1}{3} (\blacktriangle R) R - \frac{2}{3} \|\text{Ric}\|^2 R \\ &\quad + 2 \langle \langle \text{Ric}, \blacktriangle \widehat{T} \rangle \rangle + \frac{1}{3} \langle \langle \text{Ric}, \nabla^2 R \rangle \rangle + \text{Ric} * \widehat{T} * \text{Rm} + \text{Ric} * \nabla^2 \widehat{T}, \\ \|\nabla \text{Rm}\|^2 &\leq -\frac{1}{2} \blacksquare \|\text{Rm}\|^2 + C \|\text{Rm}\|^3 + C \|\text{Rm}\|^{3/2} \|\nabla^2 \mathbf{T}\| + C \|\text{Rm}\| \|\nabla \mathbf{T}\|^2, \\ \partial_t \|\text{Rm}\|^2 &= \nabla^2 \text{Ric} * \text{Rm} + \text{Ric} * \text{Rm} * \text{Rm} + \text{Rm} * \text{Rm} * \widehat{T} \\ &\quad + \text{Ric} * \nabla^2 \|\mathbf{T}\|^2 + \text{Rm} * \nabla^2 \widehat{T} + \frac{4}{3} \|\mathbf{T}\|^2 \|\text{Rm}\|^2, \\ \|\nabla \mathbf{T}\|^2 &\leq -\blacksquare \|\mathbf{T}\|^2 + C \|\text{Rm}\| \|\mathbf{T}\|^2 + C \|\mathbf{T}\|^4, \\ \partial_t dV &= \frac{2}{3} \|\mathbf{T}\|^2 dV, \quad R = -\|\mathbf{T}\|^2. \end{aligned}$$

Choose an open domain  $\Omega$  of  $\mathcal{M}$  and assume that

$$(3.36) \quad \|\text{Ric}\| \leq K$$

on  $\Omega \times [0, T]$ , Then the torsion  $\mathbf{T}$  satisfies  $\|\mathbf{T}\| \lesssim K^{1/2}$  and metrics  $g_t$  are all equivalent to  $g_0$ . We also observe from (2.25) and (3.19) that

$$(3.37) \quad \|\text{Ric}\| \lesssim 1 \iff |\Delta \varphi| \lesssim 1$$

and the following simple fact

$$(3.38) \quad \partial_t \|A\|^2 = \frac{p}{2} \|A\|^{p-2} \partial_t \|A\|^2$$

for any tensor  $A$ .

Choose a Lipschitz function  $\eta$  with support in  $\Omega$  and consider the quantity

$$\frac{d}{dt} \int \|\text{Rm}\|^p \eta^{2p} dV, \quad \int := \int_{\mathcal{M}'}$$

where  $p \geq 5$ . As in [27], we introduce the following “good” quantities

$$\begin{aligned} A_1 &:= \int \|\text{Rm}\|^p \eta^{2p} dV, \quad A_2 := \int \|\text{Rm}\|^{p-1} \eta^{2p} dV, \\ A_3 &:= \int \|\text{Rm}\|^{p-1} \|\nabla \eta\|^2 \eta^{2p-1} dV, \quad A_4 := \int \|\text{Rm}\|^{p-1} \|\nabla \eta\|^2 \eta^{2p-2} dV \end{aligned}$$

and also “bad” quantities

$$B_1 := \frac{1}{K} \int \|\nabla \text{Ric}\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV, \quad B_2 := \int \|\nabla \text{Rm}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} dV.$$

We split the proof of Theorem 1.4 into four steps.

(a) In the first step, we can show that, see Lemma 3.3,

$$\begin{aligned} \frac{d}{dt} A_1 &\leq B_1 + cKB_2 + cKA_4 + cKA_1 + cK^2 A_2 \\ &\quad + c \int \left( -\blacksquare \|T\|^2 \right) \|\text{Rm}\|^{p-1} \eta^{2p} dV. \end{aligned}$$

(b) In the second step, we can prove that the term

$$c \int \left( -\blacksquare \|T\|^2 \right) \|\text{Rm}\|^{p-1} \eta^{2p} dV$$

is bounded from above by (see (3.47))

$$B_1 + cKB_2 + cK^2 A_2 + cKA_1 - \frac{d}{dt} \left[ \int c(-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right].$$

Observe that the above integral is nonnegative, since the scalar curvature  $R$  is nonpositive along the Laplacian flow on closed  $G_2$ -structures. Hence we obtain from the first step that, see Lemma 3.4,

$$\begin{aligned} \frac{d}{dt} A_1 &\leq 2B_1 + cKB_2 + cKA_4 + cKA_1 + cK^2 A_2 \\ &\quad - \frac{d}{dt} \left[ \int c(-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]. \end{aligned}$$

(c) In the next two steps, we estimate the bad terms  $B_1$  and  $B_2$ . In the third step,  $B_1$  is estimated by (see (3.57))

$$\begin{aligned} B_1 &\leq cKB_2 + cKA_4 + cKA_1 + cK^2 A_2 \\ &\quad - \frac{d}{dt} \left[ \frac{1}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]. \end{aligned}$$

Then the second step can be simplified as, see Lemma 3.5,

$$\begin{aligned} \frac{d}{dt} A_1 &\leq cKB_2 + cKA_4 + cKA_1 + cK^2 A_2 \\ &\quad - \frac{d}{dt} \left[ \frac{1}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]. \end{aligned}$$

(d) Finally, we estimate the term  $B_2$ . In this step we shall use the assumption that  $p \geq 5$ . Using the inequality  $\|\nabla T\| \lesssim \|\text{Rm}\|$  and  $\|\nabla^2 T\| \lesssim \|\nabla \text{Rm}\| + \|\text{Rm}\| \|T\| + \|\nabla T\| \|T\| + \|T\|^3$ , we can prove (see (3.67))

$$B_2 \leq cA_4 + cA_1 - \frac{d}{dt} \left[ \frac{1}{p-1} \int \|\text{Rm}\|^{p-1} \eta^{2p} dV \right].$$

Plugging it into the third step, we arrive at, see Lemma 3.6,

$$\begin{aligned} \frac{d}{dt}(A_1 + cKA_2) &\leq cK(A_1 + cKA_2) + cKA_4 \\ &\quad - \frac{d}{dt} \left[ \frac{c}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV \right. \\ &\quad \left. + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]. \end{aligned}$$

If we choose a geodesic ball  $\Omega := B_{g_0}(x_0, \rho/\sqrt{K})$  and a cut-off function  $\eta$  so that  $\|\nabla\phi\| \leq \sqrt{K}e^{cKT}/\rho$ , then the above inequality gives a proof of Theorem 1.4.

We are going to carry out the above mentioned four steps. From (3.39) and the above evolution equations, we have

$$\begin{aligned} \frac{d}{dt} \int \|\text{Rm}\|^p \eta^{2p} dV &= \int (\partial_t \|\text{Rm}\|^p) \eta^{2p} dV + \int \|\text{Rm}\|^p \eta^{2p} \partial_t dV \\ &= \int \frac{p}{2} \|\text{Rm}\|^{p-2} (\partial_t \|\text{Rm}\|^2) \eta^{2p} dV + \int \|\text{Rm}\|^p \eta^{2p} \left( -\frac{2}{3} R \right) dV \\ &= \int \frac{p}{2} \|\text{Rm}\|^{p-2} \left[ \begin{array}{l} \nabla^2 \text{Ric} * \text{Rm} + \text{Ric} * \text{Rm} * \text{Rm} \\ + \text{Rm} * \text{Rm} * \hat{T} + \text{Ric} * \nabla^2 \|\mathbf{T}\|^2 \\ + \text{Rm} * \nabla^2 \hat{T} + \frac{4}{3} \|\mathbf{T}\|^2 \|\text{Rm}\|^2 \end{array} \right] \eta^{2p} dV \\ (3.39) \quad &\quad - \frac{2}{3} \int R \|\text{Rm}\|^p \eta^{2p} dV \\ &\leq c \int \|\text{Rm}\|^{p-2} \left[ \nabla^2 \text{Ric} * \text{Rm} + K \|\text{Rm}\|^2 + K \|\text{Rm}\|^2 + \nabla^2 \|\mathbf{T}\|^2 * \text{Ric} \right. \\ &\quad \left. + \nabla^2 \hat{T} * \text{Rm} \right] \eta^{2p} dV + cK \int \|\text{Rm}\|^p \eta^{2p} dV \\ &\leq c \int \|\text{Rm}\|^{p-2} \left[ \nabla^2 \text{Ric} * \text{Rm} + \nabla^2 \|\mathbf{T}\|^2 * \text{Ric} + \nabla^2 \hat{T} * \text{Rm} \right] \eta^{2p} dV \\ &\quad + cK \int \|\text{Rm}\|^p \eta^{2p} dV. \end{aligned}$$

It was proved in [24] that the first integral in (3.39) is bounded by

$$\begin{aligned} c \int \|\text{Rm}\|^{p-2} (\nabla^2 \text{Ric} * \text{Rm}) \eta^{2p} dV &\leq \frac{1}{K} \int \|\nabla \text{Ric}\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV \\ (3.40) \quad &+ cK \int \|\nabla \text{Rm}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} dV + cK \int \|\text{Rm}\|^{p-1} \|\nabla \eta\|^2 \eta^{2p-2} dV. \end{aligned}$$

Since  $\|\mathbf{T}\|^2 = -R$ , the same inequality holds for the integral

$$c \int \|\text{Rm}\|^{p-2} (\nabla^2 \|\mathbf{T}\|^2 * \text{Ric}) \eta^{2p} dV.$$

To deal with the last term in the bracket of (3.39), we use the same argument of [24] to conclude

$$\begin{aligned} c \int \|\text{Rm}\|^{p-2} (\nabla^2 \hat{T} * \text{Rm}) \eta^{2p} dV &= c \int (\nabla \|\text{Rm}\|^{p-2} * \nabla \hat{T} * \text{Rm}) \eta^{2p} dV \\ &\quad + c \int (\|\text{Rm}\|^{p-2} * \nabla \hat{T} * \nabla \text{Rm}) \eta^{2p} dV \end{aligned}$$

$$\begin{aligned}
& + c \int \left( \|\mathbf{Rm}\|^{p-2} * \nabla \widehat{T} * \mathbf{Rm} * \nabla \eta \right) \eta^{2p-1} dV \\
\leq & c \int \|\mathbf{Rm}\|^{p-2} \|\nabla \mathbf{Rm}\| \|\nabla \widehat{T}\| \eta^{2p} dV + c \int \|\mathbf{Rm}\|^{p-2} \|\nabla \widehat{T}\| \|\nabla \mathbf{Rm}\| \eta^{2p} dV \\
& + c \int \|\mathbf{Rm}\|^{p-1} \|\nabla \widehat{T}\| \|\nabla \eta\| \eta^{2p-1} dV \\
\leq & c \int \|\mathbf{Rm}\|^{p-2} \|\nabla \mathbf{Rm}\| \|\nabla \widehat{T}\| \eta^{2p} dV + c \int \|\mathbf{Rm}\|^{p-1} \|\nabla \widehat{T}\| \|\nabla \eta\| \eta^{2p-1} dV.
\end{aligned}$$

According to the Cauchy-Schwartz inequality, the first and second integrals are bounded by

$$\begin{aligned}
& \int \|\mathbf{Rm}\|^{p-2} \|\nabla \mathbf{Rm}\| \|\nabla \widehat{T}\| \eta^{2p} dV \\
& = \int \left( \|\mathbf{Rm}\|^{\frac{p-3}{2}} \|\nabla \mathbf{Rm}\| \eta^p \right) \left( \|\mathbf{Rm}\|^{\frac{p-1}{2}} \|\nabla \widehat{T}\| \eta^p \right) dV \\
\leq & cK \int \|\nabla \mathbf{Rm}\|^2 \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV + \frac{1}{K} \int \|\nabla \widehat{T}\|^2 \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV
\end{aligned}$$

and

$$\begin{aligned}
& \int \|\mathbf{Rm}\|^{p-1} \|\nabla \widehat{T}\| \|\nabla \eta\| \eta^{2p-1} dV \\
& = \int \left( \|\mathbf{Rm}\|^{\frac{p-1}{2}} \|\nabla \widehat{T}\| \eta^p \right) \left( \|\mathbf{Rm}\|^{\frac{p-1}{2}} \|\nabla \eta\| \eta^{p-1} \right) dV \\
\leq & \frac{1}{K} \int \|\nabla \widehat{T}\|^2 \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV + cK \int \|\mathbf{Rm}\|^{p-1} \|\nabla \eta\|^2 \eta^{2p-2} dV.
\end{aligned}$$

Hence we obtain

$$\begin{aligned}
& c \int \|\mathbf{Rm}\|^{p-2} \left( \nabla^2 \widehat{T} * \mathbf{Rm} \right) \eta^{2p} dV \leq \frac{1}{K} \int \|\nabla \widehat{T}\|^2 \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \\
(3.41) \quad & + cK \int \|\nabla \mathbf{Rm}\|^2 \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV + cK \int \|\mathbf{Rm}\|^{p-1} \|\nabla \eta\|^2 \eta^{2p-2} dV.
\end{aligned}$$

Using  $\widehat{T} = T * T$  and  $R = -\|T\|^2$  yields

$$\begin{aligned}
& \frac{1}{K} \int \|\nabla \widehat{T}\|^2 \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \\
(3.42) \quad & \leq \frac{c}{K} \int \|\nabla T\|^2 \|T\|^2 \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \leq c \int \|\nabla T\|^2 \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \\
& \leq c \int \left( -\frac{1}{4} \blacksquare \|T\|^2 + c \|\mathbf{Rm}\| \|T\|^2 + c \|T\|^4 \right) \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \\
& = c \int \left( -\blacksquare \|T\|^2 \right) \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \\
& + cK \int \|\mathbf{Rm}\|^p \eta^{2p} dV + cK^2 \int \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV.
\end{aligned}$$

Hence, using (3.40), (3.41), and (3.42), we arrive at

**Lemma 3.3.** *One has*

$$\begin{aligned}
(3.43) \quad A'_1 \equiv \frac{d}{dt} A_1 & \leq B_1 + cKB_2 + cKA_4 + cKA_1 + cK^2 A_2 \\
& + c \int \left( -\blacksquare \|T\|^2 \right) \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV.
\end{aligned}$$

In the following computations, we are mainly going to estimate or simplify the bad terms  $B_1, B_2$ , and also the term involving  $-\blacksquare\|T\|^2$ . Integration by parts on the last integral in (3.43) and using  $R = -\|T\|^2$ , we obtain

$$\begin{aligned}
& c \int \left( -\blacksquare\|T\|^2 \right) \|\text{Rm}\|^{p-1} \eta^{2p} dV = c \int ((\partial_t - \Delta)R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \\
& = c \int (\partial_t R) \|\text{Rm}\|^{p-1} \eta^{2p} dV + c \int \left\langle \nabla R, \nabla \left( \|\text{Rm}\|^{p-1} \eta^{2p} \right) \right\rangle dV \\
& = \frac{d}{dt} \left( c \int R \|\text{Rm}\|^{p-1} \eta^{2p} dV \right) - c \int R \left( \partial_t \|\text{Rm}\|^{p-1} \right) \eta^{2p} dV \\
& \quad - c \int R \|\text{Rm}\|^{p-1} \eta^{2p} \partial_t dV + c \int \left\langle \nabla R, \|\text{Rm}\|^{p-3} \text{Rm} * \nabla \text{Rm} \right\rangle \eta^{2p} dV \\
& \quad + c \int \left\langle \nabla R, \|\text{Rm}\|^{p-1} \eta^{2p-1} \nabla \eta \right\rangle dV \\
& \leq c \int \|\text{Rm}\|^{p-2} \langle \nabla R, \nabla \text{Rm} \rangle \eta^{2p} dV + c \int \|\text{Rm}\|^{p-1} \|\nabla R\| \|\nabla \eta\| \eta^{2p-1} dV \\
& \quad + c \int R^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV - c \int R \left( \partial_t \|\text{Rm}\|^{p-1} \right) \eta^{2p} dV \\
& \quad + \frac{d}{dt} \left( c \int R \|\text{Rm}\|^{p-1} \eta^{2p} dV \right).
\end{aligned}$$

The first two integrals can be simplified by using the Cauchy-Schwarz inequality as follows:

$$\begin{aligned}
& c \int \|\text{Rm}\|^{p-2} \langle \nabla R, \nabla \text{Rm} \rangle \eta^{2p} dV \leq c \int \|\nabla \text{Ric}\| \|\nabla \text{Rm}\| \|\text{Rm}\|^{p-2} \eta^{2p} dV \\
& \leq c \int \left( \|\nabla \text{Rm}\| \|\text{Rm}\|^{\frac{p-3}{2}} \eta^p \right) \left( \|\nabla \text{Ric}\| \|\text{Rm}\|^{\frac{p-1}{2}} \eta^p \right) dV \leq \frac{1}{50} B_1 + cKB_2
\end{aligned}$$

and

$$\begin{aligned}
& c \int \|\text{Rm}\|^{p-1} \|\nabla R\| \|\nabla \eta\| \eta^{2p-1} dV \leq c \int \|\text{Rm}\|^{p-1} \|\nabla \text{Ric}\| \|\nabla \eta\| \eta^{2p-1} dV \\
& \leq c \int \left( \|\text{Rm}\|^{\frac{p-1}{2}} \|\nabla \eta\| \eta^{p-1} \right) \left( \|\text{Rm}\|^{\frac{p-1}{2}} \|\nabla \text{Ric}\| \eta^p \right) dV \leq \frac{1}{50} B_1 + cKA_4.
\end{aligned}$$

Therefore

$$\begin{aligned}
& c \int \left( -\blacksquare\|T\|^2 \right) \|\text{Rm}\|^{p-1} \eta^{2p} dV \leq \frac{2}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 \\
(3.44) \quad & + \frac{d}{dt} \left( c \int R \|\text{Rm}\|^{p-1} \eta^{2p} dV \right) - c \int R \left( \partial_t \|\text{Rm}\|^{p-1} \right) \eta^{2p} dV.
\end{aligned}$$

Now, the second integral in (3.44) is equal to

$$\begin{aligned}
& -c \int R \left( \partial_t \|\text{Rm}\|^{p-1} \right) \eta^{2p} dV = c \int (-R) \|\text{Rm}\|^{p-3} \left( \partial_t \|\text{Rm}\|^2 \right) \eta^{2p} dV \\
& = c \int (-R) \|\text{Rm}\|^{p-3} \left[ \nabla^2 \text{Ric} * \text{Rm} + \text{Ric} * \text{Rm} * \text{Rm} + \text{Rm} * \text{Rm} * \widehat{T} \right. \\
& \quad \left. + \text{Ric} * \nabla^2 \|T\|^2 + \text{Rm} * \nabla^2 \widehat{T} + \frac{4}{3} \|T\|^2 \|\text{Rm}\|^2 \right] \eta^{2p} dV \\
& \leq c \int (-R) \|\text{Rm}\|^{p-3} \left[ \nabla^2 \text{Ric} * \text{Rm} - \text{Ric} * \nabla^2 R + \nabla^2 \widehat{T} * \text{Rm} \right] \eta^{2p} dV + cK^2 A_2.
\end{aligned}$$

Using the identity, where  $p \geq 5$ ,

$$\nabla \|\text{Rm}\|^{p-3} = \frac{p-3}{2} \left( \|\text{Rm}\|^2 \right)^{\frac{p-3}{2}-1} \nabla \|\text{Rm}\|^2 = \|\text{Rm}\|^{p-5} \text{Rm} * \nabla \text{Rm}$$

we obtain

$$\begin{aligned} & c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p} (\nabla^2 \text{Ric} * \text{Rm}) dV = c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p} (\nabla \text{Ric} * \nabla \text{Rm}) dV \\ & \quad + c \int \left\{ \nabla \left[ (-R) \|\text{Rm}\|^{p-3} \phi^{2p} \right] * \nabla \text{Ric} * \text{Rm} \right\} dV \\ = & c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p} (\nabla \text{Ric} * \nabla \text{Rm}) dV + c \int \|\text{Rm}\|^{p-3} \eta^{2p} (\nabla R * \nabla \text{Ric} * \text{Rm}) dV \\ & \quad + c \int (-R) \eta^{2p} \left( \nabla \|\text{Rm}\|^{p-3} * \nabla \text{Ric} * \text{Rm} \right) dV \\ & \quad + c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p-1} (\nabla \phi * \nabla \text{Ric} * \text{Rm}) dV \\ \leq & c \int \|\text{Rm}\|^{p-2} \eta^{2p} \|\nabla \text{Ric}\| \|\nabla \text{Rm}\| dV + c \int \|\nabla \text{Ric}\| \|\nabla R\| \|\text{Rm}\|^{p-2} \eta^{2p} dV \\ & \quad + c \int \|\text{Rm}\|^{p-2} \|\nabla \text{Ric}\| \|\nabla \text{Rm}\| \eta^{2p} dV + c \int \|\text{Rm}\|^{p-1} \eta^{2p-1} \|\nabla \eta\| \|\nabla \text{Ric}\| dV \\ \leq & c \int \left( \|\nabla \text{Ric}\| \|\text{Rm}\|^{\frac{p-1}{2}} \eta^p \right) \left( \|\nabla \text{Rm}\| \|\text{Rm}\|^{\frac{p-3}{2}} \eta^p \right) dV \\ & \quad + c \int \left( \|\nabla \text{Ric}\| \|\text{Rm}\|^{\frac{p-1}{2}} \eta^p \right) \left( \|\nabla \phi\| \|\text{Rm}\|^{\frac{p-1}{2}} \eta^{p-1} \right) dV \leq \frac{1}{50} B_1 + cKB_2 + cKA_4. \end{aligned}$$

Similarly, we can prove

$$c \int (-R) \|\text{Rm}\|^{p-3} \left( -\text{Ric} * \nabla^2 R \right) \eta^{2p} dV \leq \frac{1}{50} B_1 + cKB_2 + cKA_4.$$

Using  $\nabla \hat{T} = \nabla T * T \leq c \|\nabla T\| \|T\| \leq cK^{1/2} \|\nabla T\|$  yields

$$\begin{aligned} & c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p} \left( \nabla^2 \hat{T} * \text{Rm} \right) dV = c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p} (\nabla \hat{T} * \nabla \text{Rm}) dV \\ & \quad + c \int \left\{ \nabla \left[ (-R) \|\text{Rm}\|^{p-3} \eta^{2p} \right] * \nabla \hat{T} * \text{Rm} \right\} dV \\ = & c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p} (\nabla \hat{T} * \nabla \text{Rm}) dV + c \int \|\text{Rm}\|^{p-3} \eta^{2p} (\nabla R * \nabla \hat{T} * \text{Rm}) dV \\ & \quad + c \int (-R) \eta^{2p} \left( \nabla \|\text{Rm}\|^{p-3} * \nabla \hat{T} * \text{Rm} \right) dV \\ & \quad + c \int (-R) \|\text{Rm}\|^{p-3} \eta^{2p-1} \left( \nabla \eta * \nabla \hat{T} * \text{Rm} \right) dV \\ \leq & c \int \left( \|\text{Rm}\|^{p-2} \eta^{2p} \|\nabla \text{Rm}\| + \|\text{Rm}\|^{p-1} \eta^{2p-1} \|\nabla \eta\| \right) \left( K^{1/2} \|\nabla T\| \right) dV \\ \leq & c \int \left( \|\nabla \text{Rm}\| \|\text{Rm}\|^{\frac{p-3}{2}} \eta \right) \left( \|\nabla T\| K^{1/2} \|\text{Rm}\|^{\frac{p-1}{2}} \eta^p \right) dV \\ & \quad + \int \left( \|\nabla \eta\| \|\text{Rm}\|^{\frac{p-1}{2}} \eta^{p-1} \right) \left( \|\nabla T\| K^{1/2} \|\text{Rm}\|^{\frac{p-1}{2}} \eta^p \right) dV \\ \leq & \epsilon c \int \|\nabla T\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV + \frac{cK}{\epsilon} B_2 + \frac{cK}{\epsilon} A_4. \end{aligned}$$

According to (3.44) we get

$$c \int \|\nabla T\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV$$

$$\begin{aligned}
&\leq c \int \left( -\blacksquare \|T\|^2 \right) \|\text{Rm}\|^{p-1} \eta^{2p} dV + cKA_1 + cK^2 A_2 \\
&\leq \frac{2}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 \\
&+ \frac{d}{dt} \left( c \int R \|\text{Rm}\|^{p-1} \eta^{2p} dV \right) - c \int R \left( \partial_t \|\text{Rm}\|^{p-1} \right) \eta^{2p} dV \\
&\leq \frac{2}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 \\
&+ \frac{d}{dt} \left( \int cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right) + c \int (-R) \|\text{Rm}\|^{p-3} \left( \partial_t \|\text{Rm}\|^2 \right) \eta^{2p} dV.
\end{aligned}$$

Hence

$$\begin{aligned}
c \int (-R) \|\text{Rm}\|^{p-3} \left( \partial_t \|\text{Rm}\|^2 \right) \eta^{2p} dV &\leq \frac{2}{50} B_1 + cKB_2 + cKA_4 + \frac{cK}{\epsilon} B_2 + \frac{cK}{\epsilon} A_4 \\
&+ \epsilon \left[ \frac{2}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 + \frac{d}{dt} \left( \int cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right) \right] \\
&+ \epsilon c \int (-R) \|\text{Rm}\|^{p-3} \left( \partial_t \|\text{Rm}\|^2 \right) \eta^{2p} dV.
\end{aligned}$$

Choosing  $\epsilon = \frac{1}{2}$  yields

$$\begin{aligned}
&\frac{c}{2} \int (-R) \|\text{Rm}\|^{p-3} \left( \partial_t \|\text{Rm}\|^2 \right) \eta^{2p} dV \\
&\leq \frac{3}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 + \frac{d}{dt} \left( \int cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right)
\end{aligned}$$

and

$$\begin{aligned}
&c \int \|\nabla T\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV \\
&\leq \frac{8}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 + \frac{d}{dt} \left( \int 2cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right).
\end{aligned}$$

Thus

$$\begin{aligned}
(3.45) \quad &c \int (-R) \|\text{Rm}\|^{p-3} \left( \partial_t \|\text{Rm}\|^2 \right) \eta^{2p} dV \leq \frac{3}{50} B_1 + cKB_2 \\
&+ cKA_4 + cK^2 A_2 + cKA_1 + \frac{d}{dt} \left( \int cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right)
\end{aligned}$$

and

$$\begin{aligned}
(3.46) \quad &c \int \|\nabla T\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV \leq \frac{8}{50} B_1 + cKB_2 \\
&+ cKA_4 + cK^2 A_2 + cKA_1 + \frac{d}{dt} \left( \int cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right)
\end{aligned}$$

and

$$\begin{aligned}
(3.47) \quad &c \int \left( -\blacksquare \|T\|^2 \right) \|\text{Rm}\|^{p-1} \eta^{2p} dV \leq \frac{5}{50} B_1 + cKB_2 \\
&+ cK^2 A_2 + cKA_1 + \frac{d}{dt} \left( \int cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right).
\end{aligned}$$

From (3.43) and (3.47) we arrive at



**Lemma 3.4.** *One has*

$$(3.48) \quad \begin{aligned} A'_1 &\leq 2B_1 + cKB_2 + cKA_4 + cK^2A_2 + cKA_1 \\ &\quad + \frac{d}{dt} \left( \int cR \|\text{Rm}\|^{p-1} \eta^{2p} dV \right). \end{aligned}$$

We next estimate  $B_1$  and  $B_2$ . Actually, we shall see that  $B_1$  can be estimated in terms of  $B_2$ . Hence the key step is to estimate  $B_2$ . For  $B_1$ , using

$$\begin{aligned} \|\nabla \text{Ric}\|^2 &= -\frac{1}{2} \blacksquare \|\text{Ric}\|^2 + \text{Ric} * \text{Ric} * \text{Rm} - \frac{1}{3} (\blacktriangle R) T - \frac{2}{3} R \|\text{Ric}\|^2 \\ &\quad + 2 \langle \text{Ric}, \blacktriangle \hat{T} \rangle + \frac{1}{3} \langle \text{Ric}, \nabla^2 R \rangle + \text{Ric} * \hat{T} * \text{Rm} + \text{Ric} * \nabla^2 \hat{T}. \end{aligned}$$

we obtain

$$(3.49) \quad \begin{aligned} B_1 &\leq \frac{1}{2K} \int \|\text{Rm}\|^{p-1} \eta^{2p} (\blacktriangle - \partial_t) \|\text{Ric}\|^2 dV + cKA_1 \\ &\quad + \frac{1}{3K} \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} \Delta R dV + \frac{2}{K} \int \langle \text{Ric}, \blacktriangle \hat{T} \rangle \|\text{Rm}\|^{p-1} \eta^{2p} dV \\ &\quad + \frac{1}{3K} \int \langle \text{Ric}, \nabla^2 R \rangle \|\text{Rm}\|^{p-1} \eta^{2p} dV + \frac{1}{K} \int \|\text{Rm}\|^{p-1} (\text{Ric} * \nabla^2 \hat{T}) \eta^{2p} dV. \end{aligned}$$

From the estimates  $\nabla \|\text{Ric}\|^2 \lesssim \|\text{Ric}\| \|\nabla \text{Ric}\|$ ,  $\nabla \|\text{Rm}\|^{p-1} \lesssim \|\text{Rm}\|^{p-2} \|\nabla \text{Rm}\|$ , and  $\partial_t \|\text{Rm}\|^{p-1} = \frac{p-1}{2} \|\text{Rm}\|^{p-3} \partial_t \|\text{Rm}\|^2$ , we have

$$\begin{aligned} &\int \|\text{Rm}\|^{p-1} \eta^{2p} (\blacktriangle - \partial_t) \|\text{Ric}\|^2 dV \\ &= \int \nabla \|\text{Ric}\|^2 * \nabla (\|\text{Rm}\|^{p-1} \eta^{2p}) dV - \int \|\text{Rm}\|^{p-1} \eta^{2p} (\partial_t \|\text{Ric}\|^2) dV \\ &= \int (\nabla \|\text{Ric}\|^2 * \nabla \|\text{Rm}\|^{p-1}) \eta^{2p} dV + \int (\nabla \|\text{Ric}\|^2 * \nabla \eta) \|\text{Rm}\|^{p-1} \eta^{2p-1} dV \\ &\quad - \frac{d}{dt} \left[ \int \|\text{Rm}\|^{p-1} \eta^{2p} \|\text{Ric}\|^2 dV \right] + \int (\partial_t \|\text{Rm}\|^{p-1}) \eta^{2p} \|\text{Ric}\|^2 dV \\ &\quad + \int \|\text{Rm}\|^{p-1} \eta^{2p} \|\text{Ric}\|^2 (\partial_t dV) \\ &\leq cK \int \|\nabla \text{Ric}\| \|\nabla \text{Rm}\| \|\text{Rm}\|^{p-2} \eta^{2p} dV + cK \int \|\nabla \text{Ric}\| \|\nabla \eta\| \|\text{Rm}\|^{p-1} \eta^{2p-1} dV \\ &\quad + c \int \|\text{Rm}\|^{p-3} (\partial_t \|\text{Rm}\|^2) \eta^{2p} \|\text{Ric}\|^2 dV + cK^2 A_1 \\ &\quad - \frac{d}{dt} \left[ \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV \right] \\ &\leq cK \left( \frac{1}{50c} B_1 + cKB_2 \right) + cK \left( \frac{1}{50c} B_1 + cKA_4 \right) + cK^2 A_1 \\ &\quad + c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} (\partial_t \|\text{Rm}\|^2) dV - \frac{d}{dt} \left[ \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV \right] \\ &\leq \frac{2}{50} KB_1 + cK^2 B_2 + cK^2 A_4 + cK^2 A_1 \\ &\quad + c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} (\partial_t \|\text{Rm}\|^2) dV - \frac{d}{dt} \left[ \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV \right]. \end{aligned}$$

Thus

$$(3.50) \quad \int \|\text{Rm}\|^{p-1} \eta^{2p} \|\text{Ric}\|^2 dV \leq \frac{2}{50} K B_1 + c K^2 B_2 + c K^2 A_4 + c K^2 A_1 \\ + c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( \partial_t \|\text{Rm}\|^2 \right) dV - \frac{d}{dt} \left[ \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV \right].$$

Consider the term

$$c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( \partial_t \|\text{Rm}\|^2 \right) dV = c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \\ \left[ \nabla^2 \text{Ric} * \text{Rm} + \text{Ric} * \text{Rm} * \text{Rm} + \text{Rm} * \text{Rm} * \widehat{T} + \text{Ric} * \nabla^2 \|\mathbf{T}\|^2 + \text{Rm} * \nabla^2 \widehat{T} \right. \\ \left. + \frac{4}{3} \|\mathbf{T}\|^2 \|\text{Rm}\|^2 \right] dV \leq c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left[ \nabla^2 \text{Ric} * \text{Rm} - \nabla^2 R * \text{Ric} \right. \\ \left. + \nabla^2 \widehat{T} * \text{Rm} \right] dV + c K^2 A_2.$$

The three terms in the bracket can be estimated as follows. Firstly

$$c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( \nabla^2 \text{Ric} * \text{Rm} \right) dV \\ = c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( \nabla \text{Ric} * \nabla \text{Rm} \right) dV \\ + c \int \left\{ \nabla \left[ \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \right] * \nabla \text{Ric} * \text{Rm} \right\} dV \\ = c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( \nabla \text{Ric} * \nabla \text{Rm} \right) dV \\ + c \int \|\text{Rm}\|^{p-3} \eta^{2p} \left( \nabla \|\text{Ric}\|^2 * \nabla \text{Ric} * \text{Rm} \right) dV \\ + c \int \|\text{Ric}\|^2 \eta^{2p} \left( \nabla \|\text{Rm}\|^{p-3} * \nabla \text{Ric} * \text{Rm} \right) dV \\ + c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p-1} \left( \nabla \eta * \nabla \text{Ric} * \text{Rm} \right) dV \\ \leq c K \int \|\text{Rm}\|^{p-2} \eta^{2p} \|\nabla \text{Ric}\| \|\nabla \text{Rm}\| dV + c K \int \|\text{Rm}\|^{p-1} \eta^{2p-1} \|\nabla \text{Ric}\| \|\nabla \eta\| dV \\ \leq c K \left( \epsilon B_1 + \frac{K}{\epsilon} B_2 \right) + c K \left( \epsilon B_1 + \frac{K}{\epsilon} A_4 \right) \leq \frac{1}{50} K B_1 + c K^2 B_2 + c K^2 A_4.$$

The same estimate holds for

$$c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( -\nabla^2 R * \text{Ric} \right) dV.$$

Finally,

$$c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( \nabla^2 \widehat{T} * \text{Rm} \right) dV = c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \\ \left( \nabla \widehat{T} * \nabla \text{Rm} \right) dV + c \int \left\{ \nabla \left( \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \right) * \nabla \widehat{T} * \text{Rm} \right\} dV \\ \leq c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} \left( K^{1/2} \|\nabla \mathbf{T}\| \|\nabla \text{Rm}\| \right) dV \\ + c \int \left( \nabla \|\text{Ric}\|^2 \right) \|\text{Rm}\|^{p-3} \eta^{2p} \|\nabla \widehat{T}\| \|\text{Rm}\| dV \\ + c \int \|\text{Rm}\|^2 \left( \nabla \|\text{Rm}\|^{p-3} \right) \eta^{2p} \|\nabla \widehat{T}\| \|\text{Rm}\| dV$$

$$\begin{aligned}
& + c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p-1} \|\nabla \eta\| \|\nabla \widehat{T}\| \|\text{Rm}\| dV \\
& \leq cK \int \|\text{Rm}\|^{p-2} \eta^{2p} \left( K^{1/2} \|\nabla T\| \|\nabla \text{Rm}\| \right) dV \\
& \quad + cK \int \|\text{Rm}\|^{p-1} \eta^{2p-1} \left( K^{1/2} \|\nabla \eta\| \|\nabla T\| \right) dV \\
& \leq K \left[ cKB_2 + \frac{cK}{\epsilon} A_4 + \epsilon c \int \|\nabla T\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV \right] \\
& \leq \frac{8}{50} KB_1 + cK^2 B_2 + cK^2 A_4 + cK^3 A_2 + cK^2 A_1 + \frac{d}{dt} \left[ cK \int R \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]
\end{aligned}$$

Therefore

$$\begin{aligned}
(3.51) \quad c \int \|\text{Ric}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} (\partial_t \|\text{Rm}\|^2) dV & \leq \frac{10}{50} KB_1 + cK^2 B_2 + cK^2 A_4 + cK^3 A_2 \\
& \quad + cK^2 A_1 + cK \frac{d}{dt} \left[ \int R \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]
\end{aligned}$$

and

$$\begin{aligned}
(3.52) \quad \frac{1}{2K} \int \|\text{Rm}\|^{p-1} \eta^{2p} (\blacktriangle - \partial_t) \|\text{Ric}\|^2 dV & \leq \frac{6}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 \\
& \quad - \frac{1}{K} \frac{d}{dt} \left[ \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV \right] + c \frac{d}{dt} \left[ \int R \|\text{Rm}\|^{p-1} \eta^{2p} dV \right] \\
& \leq \frac{6}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 \\
& \quad - \frac{d}{dt} \left[ \frac{1}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right].
\end{aligned}$$

In the following, we estimate the left four terms in (3.49). We start from terms involving the scalar curvature.

$$\begin{aligned}
(3.53) \quad \frac{1}{3K} \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} \Delta R dV & = -\frac{1}{3K} \int \nabla R \cdot \nabla \left[ (-R) \|\text{Rm}\|^{p-1} \eta^{2p} \right] dV \\
& = -\frac{1}{3K} \int \nabla R \cdot \left[ -\nabla R \|\text{Rm}\|^{p-1} \eta^{2p} + (-R) \nabla \|\text{Rm}\|^{p-1} \eta^{2p} \right. \\
& \quad \left. + 2p(-R) \|\text{Rm}\|^{p-1} \eta^{2p-1} \nabla \eta \right] dV \leq \frac{1}{3K} \int \|\nabla R\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV \\
& \quad + \frac{c}{K} \int (-R) \|\text{Rm}\|^{p-2} \|\nabla R\| \|\nabla \text{Rm}\| \eta^{2p} dV \\
& \quad + \frac{c}{K} \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p-1} \|\nabla R\| \|\nabla \eta\| dV \\
& \leq \frac{1}{3K} \int \|\nabla R\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV + \frac{1}{3K} \int \|\nabla R\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV + cKB_2 \\
& \quad + \frac{1}{3K} \int \|\nabla R\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV + cKA_4 \\
& \leq \frac{1}{K} \int \|\nabla R\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV + cKB_2 + cKA_4.
\end{aligned}$$

The another term involving the scalar curvature can be estimated by

$$\begin{aligned}
& \frac{1}{3K} \int \langle \langle \text{Ric}, \nabla^2 R \rangle \rangle ||\text{Rm}||^{p-1} \eta^{2p} dV = -\frac{1}{3K} \int \nabla^j R \nabla^i [R_{ij} ||\text{Rm}||^{p-1} \eta^{2p}] dV \\
& = -\frac{1}{3K} \int \nabla^j R \left[ \frac{1}{2} \nabla_j R ||\text{Rm}||^{p-1} \eta^{2p} + R_{ij} \nabla^i ||\text{Rm}||^{p-1} \eta^{2p} \right. \\
(3.54) \quad & \left. + R_{ij} ||\text{Rm}||^{p-1} 2p \eta^{2p-1} \nabla^i \eta \right] dV \leq -\frac{1}{6K} \int ||\nabla R||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV \\
& \quad + \frac{c}{K} \int ||\text{Ric}|| ||\nabla R|| ||\text{Rm}||^{p-2} ||\nabla \text{Rm}|| \eta^{2p} dV \\
& \quad + \frac{c}{K} \int ||\nabla R|| ||\text{Ric}|| ||\text{Rm}||^{p-1} \eta^{2p-1} ||\nabla \eta|| dV \\
& \leq -\frac{1}{6K} \int ||\nabla R||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV + \frac{1}{18K} \int ||\nabla R||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV + cKB_2 \\
& \quad + \frac{1}{18K} \int ||\nabla R||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV + cKA_4 \leq cKB_2 + cKA_4.
\end{aligned}$$

Using (3.46) we obtain

$$\begin{aligned}
& \frac{2}{K} \int \langle \langle \text{Ric}, \blacktriangle \hat{T} \rangle \rangle ||\text{Rm}||^{p-1} \eta^{2p} dV = \frac{1}{K} \int (\text{Ric} * \blacktriangle \hat{T}) ||\text{Rm}||^{p-1} \eta^{2p} dV \\
& = \frac{1}{K} \int (\nabla \text{Ric} * \nabla \hat{T}) ||\text{Rm}||^{p-1} \eta^{2p} dV + \frac{1}{K} \int \text{Ric} * \nabla \hat{T} * \nabla (||\text{Rm}||^{p-1} \eta^{2p}) dV \\
& \leq \frac{c}{K} \int ||\nabla \text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-1} \eta^{2p} dV + \frac{c}{K} \int ||\text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-2} ||\nabla \text{Rm}|| \eta^{2p} dV \\
& \quad + \frac{c}{K} \int ||\text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-1} \eta^{2p-1} ||\nabla \eta|| dV \\
(3.55) \quad & \leq \frac{1}{50} B_1 + c \int ||\nabla T||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV + cKB_2 \\
& \quad + c \int ||\nabla T||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV + cKA_4 + c \int ||\nabla T||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV \\
& \leq \frac{1}{50} B_1 + cKB_2 + cKA_4 + c \int ||\nabla T||^2 ||\text{Rm}||^{p-1} \eta^{2p} dV \\
& \leq \frac{9}{50} B_1 + cKB_2 + cKA_4 + cK^2 A_2 + cKA_1 + \frac{d}{dt} \left[ \int cR ||\text{Rm}||^{p-1} \eta^{2p} dV \right].
\end{aligned}$$

Similarly, we can prove

$$\begin{aligned}
& \frac{1}{K} \int (\text{Ric} * \nabla^2 \hat{T}) ||\text{Rm}||^{p-1} \eta^{2p} dV = \frac{1}{K} \int (\nabla \text{Ric} * \nabla \hat{T}) ||\text{Rm}||^{p-1} \eta^{2p} dV \\
& + \frac{1}{K} \int \text{Ric} * \nabla \hat{T} * \nabla (||\text{Rm}||^{p-1} \eta^{2p}) dV \leq \frac{1}{K} \int (\nabla \text{Ric} * \nabla \hat{T}) ||\text{Rm}||^{p-1} \eta^{2p} dV \\
& \quad + \frac{c}{K} \int ||\text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-2} ||\nabla \text{Rm}|| \eta^{2p} dV \\
(3.56) \quad & \quad + \frac{c}{K} \int ||\text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-1} \eta^{2p-1} ||\nabla \eta|| dV \\
& \leq \frac{c}{K} \int ||\nabla \text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-1} \eta^{2p} dV + \frac{c}{K} \int ||\text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-2} ||\nabla \text{Rm}|| \eta^{2p} dV \\
& \quad + \frac{c}{K} \int ||\text{Ric}|| ||\nabla \hat{T}|| ||\text{Rm}||^{p-1} \eta^{2p-1} ||\nabla \eta|| dV
\end{aligned}$$

$$\leq \frac{9}{50}B_1 + cKB_2 + cKA_4 + cK^2A_2 + cKA_1 + \frac{d}{dt} \left[ \int cR |\text{Rm}|^{p-1} \eta^{2p} dV \right].$$

Plugging (3.50) and (3.53) – (3.56) into (3.49), and using (3.46) and  $\|\nabla R\|^2 \leq cK\|\nabla T\|^2$ , we obtain

$$\begin{aligned} B_1 &\leq \frac{6}{50}B_1 + cKB_2 + cKA_4 + cK^2A_2 + cKA_1 \\ &\quad - \frac{d}{dt} \left[ \frac{1}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right] \\ &+ \frac{1}{K} \int \|\nabla R\|^2 \|\text{Rm}\|^{p-1} \eta^{2p} dV + \frac{18}{50}B_1 - \frac{d}{dt} \left[ c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right] \\ &\leq \frac{32}{50}B_1 + cKB_2 + cKA_4 + cK^2A_2 + cKA_1 \\ &\quad - \frac{d}{dt} \left[ \frac{1}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]. \end{aligned}$$

Thus

$$(3.57) \quad \begin{aligned} B_1 &\leq cKB_2 + cKA_4 + cK^2A_2 + cKA_1 \\ &\quad - \frac{d}{dt} \left[ \frac{1}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right] \end{aligned}$$

From (3.48) and (3.57), we can conclude that

**Lemma 3.5.** *One has*

$$(3.58) \quad \begin{aligned} A'_1 &\leq cKB_2 + cKA_4 + cK^2A_2 + cKA_1 \\ &\quad - \frac{d}{dt} \left[ \frac{c}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]. \end{aligned}$$

Observe that two terms in the bracket are both nonnegative, since  $R = -\|T\|^2 \leq 0$ .

Finally, we estimate the term  $B_2$ . Using the evolution inequality

$$\|\nabla \text{Rm}\|^2 \leq -\frac{1}{2} \blacksquare \|\text{Rm}\|^2 + c\|\text{Rm}\|^3 + c\|\nabla^2 T\| \|\text{Rm}\|^{3/2} + c\|\text{Rm}\| \|\nabla T\|^2$$

we obtain

$$(3.59) \quad \begin{aligned} B_2 &= \int \|\nabla \text{Rm}\|^2 \|\text{Rm}\|^{p-3} \eta^{2p} dV \leq \int \left[ -\frac{1}{2} \blacksquare \|\text{Rm}\|^2 + c\|\text{Rm}\|^3 \right. \\ &\quad \left. + c\|\nabla^2 T\| \|\text{Rm}\|^{3/2} + c\|\text{Rm}\| \|\nabla T\|^2 \right] \|\text{Rm}\|^{p-3} \eta^{2p} dV \\ &\leq -\frac{1}{2} \int \left( \blacksquare \|\text{Rm}\|^2 \right) \|\text{Rm}\|^{p-3} \eta^{2p} dV + cA_1 \\ &\quad + c \int \|\nabla^2 T\| \|\text{Rm}\|^{p-3/2} \eta^{2p} dV + c \int \|\nabla^2 T\|^2 \|\text{Rm}\|^{p-2} \eta^{2p} dV. \end{aligned}$$

For the first integral one has

$$-\frac{1}{2} \int \left( \blacksquare \|\text{Rm}\|^2 \right) \|\text{Rm}\|^{p-3} \eta^{2p} dV = \frac{1}{2} \int \left( \blacktriangle \|\text{Rm}\|^2 \right) \|\text{Rm}\|^{p-3} \eta^{2p} dV$$

$$\begin{aligned}
& -\frac{1}{2} \int \left( \partial_t \|\mathbf{Rm}\|^2 \right) \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV = -\frac{1}{2} \int \left( \partial_t \|\mathbf{Rm}\|^2 \right) \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV \\
& \quad - \frac{1}{2} \int \nabla \|\mathbf{Rm}\|^2 \left[ \left( \nabla \|\mathbf{Rm}\|^{p-3} \right) \eta^{2p} + \|\mathbf{Rm}\|^{p-3} \left( \nabla \eta^{2p} \right) \right] dV \\
& \quad = -\frac{p-3}{4} \int \left( \nabla \|\mathbf{Rm}\|^2 \right)^2 \|\mathbf{Rm}\|^{p-5} \eta^{2p} dV \\
& + c \int \|\mathbf{Rm}\|^{p-2} \|\nabla \mathbf{Rm}\| \|\nabla \eta\| \eta^{2p-1} dV - \frac{1}{2} \int \left( \partial_t \|\mathbf{Rm}\|^2 \right) \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV \\
& \leq \frac{1}{50} B_2 + cA_4 - \frac{1}{2} \int \left( \partial_t \|\mathbf{Rm}\|^2 \right) \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV.
\end{aligned}$$

Here we used the assumption that  $p \geq 5$ . On the other hand,

$$\begin{aligned}
& -\frac{1}{2} \int \left( \partial_t \|\mathbf{Rm}\|^2 \right) \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV = -\frac{1}{2} \frac{d}{dt} \left[ \int \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \right] \\
& \quad + \frac{1}{2} \int \|\mathbf{Rm}\|^2 \left( \partial_t \|\mathbf{Rm}\|^{p-3} \right) \eta^{2p} dV + \frac{1}{2} \int \|\mathbf{Rm}\|^{p-1} \eta^{2p} \left( \partial_t dV \right) \\
& \leq \frac{p-3}{4} \int \|\mathbf{Rm}\|^{p-3} \left( \partial_t \|\mathbf{Rm}\|^2 \right) \eta^{2p} dV + cA_1 - \frac{1}{2} \frac{d}{dt} \left[ \int \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \right]
\end{aligned}$$

so that

$$-\frac{1}{2} \int \left( \partial_t \|\mathbf{Rm}\|^2 \right) \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV \leq cA_1 - \frac{1}{p-1} \frac{d}{dt} \left[ \int \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \right].$$

Therefore

$$\begin{aligned}
& -\frac{1}{2} \int \left( \blacksquare \|\mathbf{Rm}\|^2 \right) \|\mathbf{Rm}\|^{p-3} \eta^{2p} dV \leq \frac{1}{50} B_2 + cA_4 + cA_1 \\
(3.60) \quad & - \frac{1}{p-1} \frac{d}{dt} \left[ \int \|\mathbf{Rm}\|^{p-1} \eta^{2p} dV \right].
\end{aligned}$$

To estimate the remainder two integrals, we recall from (3.9) that

$$(3.61) \quad \nabla \mathbf{T} = \mathbf{Rm} * \varphi + \mathbf{T} * \mathbf{T} * \varphi$$

and from (2.14) that

$$(3.62) \quad \nabla \varphi = \mathbf{T} * \psi.$$

From (3.61) we get

$$(3.63) \quad \|\nabla \mathbf{T}\| \leq c\|\mathbf{Rm}\| + c\|\mathbf{T}\|^2 \leq c\|\mathbf{Rm}\|.$$

In particular, the inequality (3.63) yields

$$(3.64) \quad \int \|\nabla \mathbf{T}\|^2 \|\mathbf{Rm}\|^{p-2} \eta^{2p} dV \leq c \int \|\mathbf{Rm}\|^p \eta^{2p} dV \leq cA_1.$$

Taking the derivative of (3.61) and using (3.62) we obtain

$$(3.65) \quad \nabla^2 \mathbf{T} = \nabla \mathbf{Rm} * \varphi + \mathbf{Rm} * \mathbf{T} * \psi + \nabla \mathbf{T} * \mathbf{T} * \varphi + \mathbf{T} * \mathbf{T} * \mathbf{T} * \psi.$$

The particular case  $\|\nabla^2 \mathbf{T}\| \leq c\|\nabla \mathbf{Rm}\| + c\|\mathbf{Rm}\| \|\mathbf{T}\| + c\|\nabla \mathbf{T}\| \|\mathbf{T}\| + c\|\mathbf{T}\|^3$  leads to

$$\begin{aligned}
& c \int \|\nabla^2 \mathbf{T}\| \|\mathbf{Rm}\|^{p-3/2} \eta^{2p} dV \leq c \int \left[ \|\nabla \mathbf{Rm}\| + \|\mathbf{Rm}\| \|\mathbf{T}\| + \|\nabla \mathbf{T}\| \|\mathbf{T}\| \right. \\
& \left. + \|\mathbf{T}\|^3 \right] \|\mathbf{Rm}\|^{p-3/2} \eta^{2p} dV \leq c \int \left( \|\nabla \mathbf{Rm}\| \|\mathbf{Rm}\|^{p-3/2} \eta^p \right) \left( \|\mathbf{Rm}\|^{p/2} \eta^p \right) dV
\end{aligned}$$

$$(3.66) \quad + c \int \|\text{Rm}\|^p \eta^{2p} dV \leq \frac{1}{50} B_2 + cA_1.$$

Plugging (3.60), (3.64), and (3.66) into (3.59) we arrive at

$$(3.67) \quad B_2 \leq cA_4 + cA_1 - \frac{d}{dt} \left[ \frac{1}{p-1} \int \|\text{Rm}\|^{p-1} \eta^{2p} dV \right].$$

Together with (3.58) and (3.67) we finally obtain

$$(3.68) \quad \begin{aligned} & (A_1 + cKA_2)' \leq cK(A_1 + cKA_2) + cKA_4 \\ & - \frac{d}{dt} \left[ \frac{c}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right]. \end{aligned}$$

Equivalently,

**Lemma 3.6.** *If  $\|\text{Ric}\| \leq K$  and  $p \geq 5$ , one has*

$$(3.69) \quad \begin{aligned} & \frac{d}{dt} \left[ A_1 + cKA_2 + \frac{c}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV \right] \\ & \leq cK(A_1 + cKA_2) + cKA_4. \end{aligned}$$

As in [24, 27], we choose the domain  $\Omega := B_{g_0}(x_0, \rho/\sqrt{K})$  and the function

$$\eta = \left( \frac{\rho/\sqrt{K} - d_{g(0)}(x_0, \cdot)}{\rho/\sqrt{K}} \right)_+.$$

Then, for all  $t \in [0, T]$ ,

$$e^{-cKt} g_0 \leq g(t) \leq e^{cKt} g_0, \quad \|\nabla_{g(t)} \phi\|_{g(t)} \leq e^{cKt} \|\nabla_{g_0} \phi\|_{g_0} \leq \frac{\sqrt{K} e^{cKt}}{\rho}.$$

The proof of Theorem 1.4. Define

$$(3.70) \quad \begin{aligned} U & := \int \|\text{Rm}\|^p \phi^{2p} dV + cK \int \|\text{Rm}\|^{p-1} \eta^{2p} dV \\ & + \frac{c}{K} \int \|\text{Rm}\|^{p-1} \|\text{Ric}\|^2 \eta^{2p} dV + c \int (-R) \|\text{Rm}\|^{p-1} \eta^{2p} dV. \end{aligned}$$

Then (3.69) yields

$$(3.71) \quad U' \leq cKU + cKA_4.$$

For  $A_4$ , using the Young inequality, we have

$$\begin{aligned} A_4 & = \int \|\text{Rm}\|^{p-1} \|\nabla \eta\|^2 \eta^{2p-2} dV \leq \int_{B_{g_0}(x_0, \rho/\sqrt{K})} \|\text{Rm}\|^{p-1} \eta^{2p-2} K \rho^{-2} e^{cKt} dV \\ & \leq \int_{B_{g_0}(x_0, \rho/\sqrt{K})} \left[ \frac{(\|\text{Rm}\|^{p-1} \eta^{2p-2})^{p/(p-1)}}{\frac{p}{p-1}} + \frac{(K \rho^{-2} e^{cKt})^p}{p} \right] dV \\ & \leq A_1 + K^p \rho^{-2p} p e^{cKt} \text{vol}_{g(t)} \left( B_{g_0} \left( x_0, \frac{\rho}{\sqrt{K}} \right) \right) \\ & \leq U + cK^p e^{cKt} \rho^{-2p} \text{vol}_{g(t)} \left( B_{g_0} \left( x_0, \frac{\rho}{\sqrt{K}} \right) \right). \end{aligned}$$

Thus

$$U' \leq cKU + cK^{p+1} e^{cKt} \rho^{-2-p} \text{vol}_{g(t)} \left( B_{g_0} \left( x_0, \frac{\rho}{\sqrt{K}} \right) \right).$$

As in the proof of [24], one can easily deduce from above that

$$(3.72) \quad \int_{B_{g_0}(x_0, \frac{\rho}{2\sqrt{K}})} \|\mathrm{Rm}_{g(t)}\|_{g(t)}^p dV_{g(t)} \leq c(1+K)e^{cKT} \int_{B_{g_0}(x_0, \frac{\rho}{\sqrt{K}})} \|\mathrm{Rm}_{g_0}\|_{g_0}^p dV_{g_0} \\ + cK^p \left(1 + \rho^{-2p}\right) e^{cKT} \mathrm{vol}_{g(t)} \left( B_{g_0} \left( x_0, \frac{\rho}{\sqrt{K}} \right) \right).$$

As an immediate consequence of the inequality (3.72) we give another proof of the part (a) in Theorem 1.2.

#### REFERENCES

- [1] Bryant, Robert L. *Some remarks on  $G_2$ -structures*, Proceedings of Gökova Geometry-Topology Conference 2005, 75–09, Gökova Geometry/Topology Conference (GGT), Gökova, 2006. MR2282011 (2007k: 53019)
- [2] Bryant, Robert L.; Xu, Feng. *Laplacian flow for closed  $G_2$ -structures: short time behavior*, arXiv: 1101.2004.
- [3] Cao, Xiaodong. *Curvature pinching estimate and singularities of the Ricci flow*, *Comm. Anal. Geom.*, **19**(2011), no. 5, 975–990. MR2886714
- [4] Cheng, Liang; Zhu, Anqiang. *On the extension of the harmonic Ricci flow*, *Geom. Dedicata*, **164**(2013), 179–185. MR3054623
- [5] Cleyton, R.; Ivanov, S. *On the geometry of closed  $G_2$ -structure*, *Commun. Math. Phys.*, **270**(2007), no. 1, 53–67. MR2276440 (2007m: 53025)
- [6] Donaldson, Simon. *Adiabatic limits of co-associative Kovalev-Lefschetz fibrations*, *Algebra, geometry, and physics in the 21st century*, 1–29, *Progr. Math.*, **324**, Birkhäuser/Springer, Cham, 2017. MR3702382
- [7] Donaldson, Simon. *Boundary value problems in dimension seven, four and three related to exceptional holonomy*, arXiv: 1708.01649.
- [8] Donaldson, Simon. *An elliptic boundary value problem for  $G_2$  structures*, arXiv: 1801.01806.
- [9] Donaldson, Simon. *Remarks on  $G_2$ -manifolds with boundary*, arXiv: 1802.09694.
- [10] Enders, Joerg; Müller Reto; Topping, Peter M. *On type-I singularities in Ricci flow*, *Comm. Anal. Geom.*, **19**(2011), no. 5, 905–922. MR2886712
- [11] Fernández, M.; Gray, A. *Riemannian manifolds with structure group  $G_2$* , *Ann. Mat. Pura Appl.*, (4)**132**(1982), 19–45. MR0696037 (84e: 53056)
- [12] Fernández, Marisa; Fino, Anna; Manero, Victor. *Laplacian flow of closed  $G_2$ -structures inducing nilsoltons*, *J. Geom. Anal.*, **26**(2016), no. 3, 1808–1837. MR3511459
- [13] Fine, Joel; Yao, Chengjian. *Hypersymplectic 4-manifolds, the  $G_2$ -Laplacian flow and extension assuming bounded scalar curvature*, arXiv: 1704.07620
- [14] Grigorian, Sergey. *Short-time behavior of a modified Laplacian coflow of  $G_2$ -structures*, *Adv. Math.*, **248**(2013), 378–415. MR3107516
- [15] Grigorian, Sergey. *Modified Laplacian coflow of  $G_2$ -structures on manifolds with symmetry*, *Differential Geom. Appl.*, **46**(2016), 39–78. MR3475531
- [16] Hamilton, Richard. *Three-manifolds with positive Ricci curvature*, *J. Differential Geom.*, **17**(1982), no. 2, 255–306. MR0664497 (84a: 53050)
- [17] Hitchin, N. *The geometry of three-forms in six dimensions*, *J. Differential Geom.*, **55**(2000), no. 3, 547–576. MR1863733 (2002m: 53070)
- [18] Huang, Hongnian; Wang, Yuanqi; Yao, Chengjian. *Cohomogeneity-one  $G_2$ -Laplacian flow on 7-torus*, to appear in *Journal of the London Mathematical Society*.
- [19] Karigiannis, Sprio. *Deformations of  $G_2$  and  $\mathrm{Spin}(7)$ -structures on manifolds*, Ph. D. Thesis, Harvard University, 2003.
- [20] Karigiannis, Sprio. *Deformations of  $G_2$  and  $\mathrm{Spin}(7)$ -structures on manifolds*, *Canad. J. Math.*, **57**(2005), no. 5, 1012–1055. MR2164593 (2006j: 53070)
- [21] Karigiannis, Spiro. *Some notes on  $G_2$  and  $\mathrm{Spin}(7)$  geometry*, *Recent Advances in Geometric Analysis*, 129–146, *Adv. Lect. Math. (ALM)*, **11**, Internat. Press, Somerville, MA, 2010. MR2648941 (2011f: 53094)
- [22] Karigiannis, Sprio. *Flows of  $G_2$  structures*, *I, Q. J. Math.*, **60**(2009), no. 4, 487–522. MR2559631 (2011g: 53046)



- [23] Karigiannis, Spiro; McKay, Ben; Tsui, Mao-Pei. *Soliton solutions for the Laplacian coflow of some  $G_2$ -structures with symmetry*, *Diff. Geom. Appl.*, **30**(2012), no. 4, 318–333. MR2926272
- [24] Kotschwar, Brett; Munteanu, Ovidiu; Wang, Jiaping. *A local curvature estimate for the Ricci flow*, *J. Funct. Anal.*, **271**(2016), no. 9, 2604–2630. MR3545226
- [25] Li, Yi. *Long time existence of Ricci-harmonic flow*, *Front. Math. China*, **11**(2016), no. 5, 1313–1334. MR3547931
- [26] Li, Yi. *Long time existence and bounded scalar curvature in the Ricci-harmonic flow*, *J. Differential Equations*, **265**(2018), 69–97. MR3782539
- [27] Li, Yi. *Local curvature estimates for Ricci-harmonic flow*, preprint, 2017. In preparation of the second revised version.
- [28] Lin, Christopher. *Laplacian solitons and  $G_2$ -geometry*, *J. Geom. Phys.*, **64**(2013), 111–119. MR3004019
- [29] List, Bernhard. *Evolution of an extended Ricci flow system*, PhD thesis, AEI Potsdam, 2005.
- [30] List, Bernhard. *Evolution of an extended Ricci flow system*, *Comm. Anal. Geom.*, **16**(2008), no. 5, 1007–1048. MR2471366 (2010i: 53126)
- [31] Müller, Reto. *The Ricci flow coupled with harmonic map flow*, PhD thesis, ETH Zürich, doi: 10.3929/ethz-a-005842361, 2009.
- [32] Müller, Reto. *Ricci flow coupled with harmonic map flow*, *Ann. Sci. Éc. Norm. Supér. (4)* **45**(2012), no. 1, 101–142. MR2961788
- [33] Lotay, Jason, D; Wei, Yong. *Laplacian flow for closed  $G_2$ -structures: Shi-type estimates, uniqueness and compactness*, *Geom. Funct. Anal.*, **27**(2017), no. 1, 165–233. MR3613456
- [34] Lotay, Jason, D; Wei, Yong. *Stability of torsion-free  $G_2$ -structures along the Laplacian flow*, to appear in *J. Differential Geom.*
- [35] Lotay, Jason, D; Wei, Yong. *Laplacian flow for closed  $G_2$ -structures: real analyticity*, to appear in *Comm. Anal. Geom.*
- [36] Sesum, Natasa. *Curvature tensor under the Ricci flow*, *Amer. J. Math.*, **127**(2005), no. 6, 1315–1324. MR2183526 (2006f:53097)
- [37] Weiss, Hartmut; Witt, Frederik. *A heat flow for special metrics*, *Adv. Math.*, **231**(2012), no. 6, 3288–3322. MR2980500
- [38] Weiss, Hartmut; Witt, Frederik. *Energy functionals and soliton equations for  $G_2$ -forms*, *Ann. Glob. Anal. Geom.*, **42**(2012), no. 4, 585–610. MR2995206
- [39] Zhang, Zhou. *Scalar curvature behavior for finite-time singularity of Kähler-Ricci flow*, *Michigan Math.*, **59**(2010), no. 2, 419–433. MR2677630 (2011j: 53128)

FACULTY OF SCIENCE, TECHNOLOGY AND COMMUNICATION (FSTC), MATHEMATIC RESEARCH UNIT, CAMPUS BELVAL, UNIVERSITE DU LUXEMBOURG, MAISON DU NOMBRE, 6, AVENUE DE LA FONTE, L-4364, ESCH-SUR-ALZETTE, GRAND-DUCHY OF LUXEMBOURG  
E-mail address: yilicms@gmail.com