

# Uniqueness of solution to scalar BSDEs with $L \exp\left(\mu\sqrt{2\log(1+L)}\right)$ -integrable terminal values

Rainer Buckdahn\*      Ying Hu†      Shanjian Tang‡

**Abstract.** In [4], the existence of the solution is proved for a scalar linearly growing backward stochastic differential equation (BSDE) if the terminal value is  $L \exp\left(\mu\sqrt{2\log(1+L)}\right)$ -integrable with the positive parameter  $\mu$  being bigger than a critical value  $\mu_0$ . In this note, we give the uniqueness result for the preceding BSDE.

**AMS Subject Classification:** 60H10

**Key Words** Backward stochastic differential equation,  $L \exp\left(\mu\sqrt{2\log(1+L)}\right)$  integrability, uniqueness.

## 1 Introduction

Let  $\{W_t, t \geq 0\}$  be a standard Brownian motion with values in  $\mathbb{R}^d$  defined on some complete probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , and  $\{\mathcal{F}_t, t \geq 0\}$  its natural filtration augmented by all  $\mathbb{P}$ -null sets of  $\mathcal{F}$ . Let us fix a nonnegative real number  $T > 0$ . The  $\sigma$ -field of predictable subsets of  $\Omega \times [0, T]$  is denoted by  $\mathcal{P}$ .

For any real  $p \geq 1$ , denote by  $L^p$  the set of all  $\mathcal{F}_T$ -measurable random variables  $\eta$  such that  $E|\eta|^p < \infty$ , by  $\mathcal{S}^p$  the set of (equivalent classes of) all real-valued, adapted and càdlàg processes  $\{Y_t, 0 \leq t \leq T\}$  such that

$$\|Y\|_{\mathcal{S}^p} := \mathbb{E} \left[ \sup_{0 \leq t \leq T} |Y_t|^p \right]^{1/p} < +\infty,$$

by  $\mathcal{L}^p$  the set of (equivalent classes of) all real-valued adapted processes  $\{Y_t, 0 \leq t \leq T\}$  such that

$$\|Y\|_{\mathcal{L}^p} := \mathbb{E} \left[ \int_0^T |Y_t|^p dt \right]^{1/p} < +\infty,$$

and by  $\mathcal{M}^p$  the set of (equivalent classes of) all predictable processes  $\{Z_t, 0 \leq t \leq T\}$  with values in  $\mathbb{R}^{1 \times d}$  such that

$$\|Z\|_{\mathcal{M}^p} := \mathbb{E} \left[ \left( \int_0^T |Z_t|^2 dt \right)^{p/2} \right]^{1/p} < +\infty.$$

---

\*Laboratoire de Mathématiques, Université de Bretagne Occidentale, 29285 Brest Cedex, France (Rainer.Buckdahn@univ-brest.fr).

†IRMAR, Université Rennes 1, Campus de Beaulieu, 35042 Rennes Cedex, France (ying.hu@univ-rennes1.fr) and School of Mathematical Sciences, Fudan University, Shanghai 200433, China. Partially supported by Lebesgue center of mathematics “Investissements d’avenir” program - ANR-11-LABX-0020-01, by ANR CAESARS - ANR-15-CE05-0024 and by ANR MFG - ANR-16-CE40-0015-01.

‡Department of Finance and Control Sciences, School of Mathematical Sciences, Fudan University, Shanghai 200433, China (sjtang@fudan.edu.cn). Partially supported by National Science Foundation of China (Grant No. 11631004) and Science and Technology Commission of Shanghai Municipality (Grant No. 14XD1400400).

Consider the following Backward Stochastic Differential Equation (BSDE):

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s, \quad t \in [0, T]. \quad (1.1)$$

Here,  $f$  (hereafter called the generator) is a real valued random function defined on the set  $\Omega \times [0, T] \times \mathbb{R} \times \mathbb{R}^{1 \times d}$ , measurable with respect to  $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R}^{1 \times d})$ , and continuous in the last two variables with the following linear growth:

$$|f(s, y, z) - f(s, 0, 0)| \leq \beta|y| + \gamma|z|, \quad (s, y, z) \in [0, T] \times \mathbb{R} \times \mathbb{R}^{1 \times d}$$

with  $f_0 := f(\cdot, 0, 0) \in \mathcal{L}^1$ ,  $\beta \geq 0$  and  $\gamma > 0$ .  $\xi$  is a real  $\mathcal{F}_T$ -measurable random variable, and hereafter called the terminal condition or terminal value.

**Definition 1.1** *By a solution to BSDE (1.1) we mean a pair  $\{(Y_t, Z_t), 0 \leq t \leq T\}$  of predictable processes with values in  $\mathbb{R} \times \mathbb{R}^{1 \times d}$  such that  $\mathbb{P}$ -a.s.,  $t \mapsto Y_t$  is continuous,  $t \mapsto Z_t$  belongs to  $L^2(0, T)$  and  $t \mapsto f(t, Y_t, Z_t)$  is integrable, and  $\mathbb{P}$ -a.s.  $(Y, Z)$  verifies (1.1).*

By BSDE  $(\xi, f)$ , we mean the BSDE with generator  $f$  and terminal condition  $\xi$ .

It is well known that for  $(\xi, f_0) \in L^p \times \mathcal{L}^p$  (with  $p > 1$ ), BSDE (1.1) admits a unique adapted solution  $(y, z)$  in the space  $\mathcal{S}^p \times \mathcal{M}^p$  if the generator  $f$  is uniformly Lipschitz in the pair of unknown variables. See e.g. [6, 3, 1] for more details. For  $(\xi, f_0) \in L^1 \times \mathcal{L}^1$ , one needs to restrict the generator  $f$  to grow sub-linearly with respect to  $z$ , i.e., with some  $q \in [0, 1)$ ,

$$|f(t, y, z) - f_0(t)| \leq \beta|y| + \gamma|z|^q, \quad (t, y, z) \in [0, T] \times \mathbb{R} \times \mathbb{R}^{1 \times d}$$

for BSDE (1.1) to have a unique adapted solution (see [1]) if the generator  $f$  is uniformly Lipschitz in the pair of unknown variables.

In [4], the existence of the solution is given for a scalar linearly growing BSDE (1.1) if the terminal value is  $L \exp\left(\mu\sqrt{2\log(1+L)}\right)$ -integrable with the positive parameter  $\mu$  being bigger than a critical value  $\mu_0 = \gamma\sqrt{T}$ , and the preceding integrability of the terminal value for a positive parameter  $\mu$  less than critical value  $\mu_0$  is shown to be not sufficient for the existence of a solution. In this note, we give the uniqueness result for the preceding BSDE under the preceding integrability of the terminal value for  $\mu > \mu_0$ .

We first establish some interesting properties of the function  $\psi(x, \mu) = x \exp\left(\mu\sqrt{2\log(1+x)}\right)$ . We observe that the obtained solution  $Y$  in [4] has the nice property:  $\psi(|Y|, a)$  belongs to the class  $(D)$  for some  $a > 0$ , which is used to prove the uniqueness of the solution by dividing the whole interval  $[0, T]$  into a finite number of sufficiently small subintervals.

## 2 Uniqueness

Define the function  $\psi$ :

$$\psi(x, \mu) := x \exp\left(\mu\sqrt{2\log(1+x)}\right), \quad (x, \mu) \in [0, +\infty) \times (0, +\infty).$$

We denote also  $\psi(\cdot, \mu)$  as  $\psi_\mu(\cdot)$

The following two lemmas can be found in Hu and Tang [4].

**Lemma 2.1** *For any  $x \in \mathbb{R}$  and  $y \geq 0$ , we have*

$$e^x y \leq e^{\frac{x^2}{2\mu^2}} + e^{2\mu^2} \psi(y, \mu). \quad (2.2)$$

**Lemma 2.2** Let  $\mu > \gamma\sqrt{T}$ . For any  $d$ -dimensional adapted process  $q$  with  $|q_t| \leq \gamma$  almost surely, for  $t \in [0, T]$ ,

$$\mathbb{E} \left[ e^{\frac{1}{2\mu^2} \left| \int_t^T q_s dW_s \right|^2} \middle| \mathcal{F}_t \right] \leq \frac{1}{\sqrt{1 - \frac{\gamma^2}{\mu^2}(T-t)}}. \quad (2.3)$$

**Proposition 2.3** We have the following assertions on  $\psi$ :

- (i) For  $\mu > 0$ ,  $\psi(\cdot, \mu)$  is convex.
- (ii) For  $c > 1$ , we have  $\psi_\mu(cx) \leq \psi_\mu(c)\psi_\mu(x)$  for any  $x \geq 0$ .
- (iii) For any triple  $(a, b, c)$  with  $a > 0, b > 0$  and  $c > 0$ , we have

$$\psi(\psi(x, a), b) \leq e^{\frac{ab^2}{c}} \psi(x, a + b + c).$$

*Proof.* The first assertion has been shown in [4]. It remains to show the Assertions (ii) and (iii). We prove Assertion (ii).

$$\begin{aligned} \psi_\mu(cx) &= cx \exp\left(\mu\sqrt{2\log(1+cx)}\right) \\ &\leq cx \exp\left(\mu\sqrt{2\log[(1+c)(1+x)]}\right) \\ &= cx \exp\left(\mu\sqrt{2\log(1+c) + 2\log(1+x)}\right) \\ &\leq cx \exp\left(\mu\sqrt{2\log(1+c)} + \mu\sqrt{2\log(1+x)}\right) \\ &= \psi_\mu(c)\psi_\mu(x). \end{aligned}$$

We now prove Assertion (iii). We have

$$\begin{aligned} &(\psi_b \circ \psi_a)(x) \\ &= \psi_a(x) \exp\left(b\sqrt{2\log(1+\psi_a(x))}\right) \\ &= x \exp\left(a\sqrt{2\log(1+x)}\right) \exp\left(b\sqrt{2\log\left(1+x e^{a\sqrt{2\log(1+x)}}\right)}\right) \\ &\leq x \exp\left(a\sqrt{2\log(1+x)}\right) \exp\left(b\sqrt{2\log\left((1+x)e^{a\sqrt{2\log(1+x)}}\right)}\right) \\ &= x \exp\left(a\sqrt{2\log(1+x)}\right) \exp\left(b\sqrt{2\log(1+x) + 2a\sqrt{2\log(1+x)}}\right). \end{aligned}$$

In view of the following elementary inequality:

$$2a\sqrt{2\log(1+x)} \leq \frac{a^2b^2}{c^2} + \frac{2c^2}{b^2} \log(1+x),$$

we have

$$\begin{aligned} &(\psi_b \circ \psi_a)(x) \\ &\leq x \exp\left(a\sqrt{2\log(1+x)}\right) \exp\left(b\sqrt{2\log(1+x) + \frac{a^2b^2}{c^2} + \frac{2c^2}{b^2} \log(1+x)}\right) \\ &\leq x \exp\left(a\sqrt{2\log(1+x)}\right) \exp\left(b\sqrt{2\log(1+x)} + b\sqrt{\frac{a^2b^2}{c^2}} + b\sqrt{\frac{2c^2}{b^2} \log(1+x)}\right). \end{aligned}$$

Therefore,

$$\begin{aligned}
& (\psi_b \circ \psi_a)(x) \\
& \leq x \exp\left(a\sqrt{2\log(1+x)}\right) \exp\left(b\sqrt{2\log(1+x)} + \frac{ab^2}{c} + c\sqrt{2\log(1+x)}\right) \\
& \leq xe^{\frac{ab^2}{c}} \exp\left((a+b+c)\sqrt{2\log(1+x)}\right).
\end{aligned}$$

□

Consider the following BSDE:

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s, \quad (2.4)$$

where  $f$  satisfies

$$|f(s, y, z) - f_0(s, 0, 0)| \leq \beta|y| + \gamma|z|, \quad (2.5)$$

with  $f_0 := f(\cdot, 0, 0) \in \mathcal{L}^1$ ,  $\beta \geq 0$  and  $\gamma > 0$ .

**Theorem 2.4** *Let  $f$  be a generator which is continuous with respect to  $(y, z)$  and verifies inequality (2.5), and  $\xi$  be a terminal condition. Let us suppose that there exists  $\mu > \gamma\sqrt{T}$  such that  $\psi(|\xi| + \int_0^T |f_0(t)| dt, \mu) \in L^1(\Omega, \mathbb{P})$ . Then BSDE (2.4) admits a solution  $(Y, Z)$  such that*

$$|Y_t| \leq \frac{1}{\sqrt{1 - \frac{\gamma^2}{\mu^2}(T-t)}} e^{\beta(T-t)} + e^{2\mu^2 + \beta(T-t)} \mathbb{E} \left[ \psi_\mu \left( |\xi| + \int_t^T |f_0(s)| ds \right) \middle| \mathcal{F}_t \right].$$

Furthermore, there exists  $a > 0$  such that  $\psi(Y, a)$  belongs to the class (D).

*Proof.* Let us fix  $n \in \mathbb{N}^*$  and  $p \in \mathbb{N}^*$ . Set

$$\xi^{n,p} := \xi^+ \wedge n - \xi^- \wedge p, \quad f_0^{n,p} := f_0^+ \wedge n - f_0^- \wedge p, \quad f^{n,p} := f - f_0 + f_0^{n,p}.$$

As the terminal value  $\xi^{n,p}$  and  $f^{n,p}(\cdot, 0, 0)$  are bounded (hence square-integrable) and  $f^{n,p}$  is a continuous generator with a linear growth, in view of the existence result in [5], the BSDE  $(\xi^{n,p}, f^{n,p})$  has a (unique) minimal solution  $(Y^{n,p}, Z^{n,p})$  in  $\mathcal{S}^2 \times \mathcal{M}^2$ . Set

$$\bar{f}^{n,p}(s, y, z) = |f_0^{n,p}(s)| + \beta y + \gamma|z|, \quad (s, y, z) \in [0, T] \times \mathbb{R} \times \mathbb{R}^{1 \times d}.$$

In view of Pardoux and Peng [6], the BSDE  $(|\xi^{n,p}|, \bar{f}^{n,p})$  has a unique solution  $(\bar{Y}^{n,p}, \bar{Z}^{n,p})$  in  $\mathcal{S}^2 \times \mathcal{M}^2$ .

By comparison theorem,

$$|Y_t^{n,p}| \leq \bar{Y}_t^{n,p}.$$

Letting  $q_s^{n,p} = \gamma \operatorname{sgn}(Z_s^{n,p})$  and

$$\mathbb{P}_{q^{n,p}} = \exp\left\{ \int_0^T q_s^{n,p} dW_s - \frac{1}{2} \int_0^T |q_s^{n,p}|^2 ds \right\} \mathbb{P},$$

we obtain,

$$\begin{aligned}
|Y_t^{n,p}| & \leq \bar{Y}_t^{n,p} \\
& = \mathbb{E}_{q^{n,p}} \left[ e^{\beta(T-t)} |\xi^{n,p}| \middle| \mathcal{F}_t \right] + \int_t^T e^{\beta(s-t)} |f_0^{n,p}(s)| ds \\
& \leq e^{\beta(T-t)} \mathbb{E}_{q^{n,p}} \left[ |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \middle| \mathcal{F}_t \right] \\
& \leq \frac{1}{\sqrt{1 - \frac{\gamma^2}{\mu^2}(T-t)}} e^{\beta(T-t)} + e^{2\mu^2 + \beta(T-t)} \mathbb{E} \left[ \psi_\mu \left( |\xi| + \int_t^T |f_0(s)| ds \right) \middle| \mathcal{F}_t \right].
\end{aligned}$$

Since  $Y^{n,p}$  is nondecreasing in  $n$  and non-increasing in  $p$ , then by the localization method in [2], there is some  $Z \in L^2(0, T; \mathbb{R}^{1 \times d})$  almost surely such that  $(Y := \inf_p \sup_n Y^{n,p}, Z)$  is an adapted solution. Therefore, we have for  $a > 0$ , using Jensen's inequality and the convexity of  $\psi_a(\cdot) := \psi(\cdot, a)$  together with Assertion (ii) of Proposition 2.3, we have

$$\begin{aligned} \psi_a(|Y_t^{n,p}|) &\leq \psi_a \left( e^{\beta(T-t)} \mathbb{E}_{q^{n,p}} \left[ |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \middle| \mathcal{F}_t \right] \right) \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \psi_a \left( \mathbb{E}_{q^{n,p}} \left[ |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \middle| \mathcal{F}_t \right] \right) \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \mathbb{E}_{q^{n,p}} \left[ \psi_a \left( |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \right) \middle| \mathcal{F}_t \right] \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \mathbb{E} \left[ \exp \left( \int_t^T q_s^{n,p} dW_s \right) \psi_a \left( |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \right) \middle| \mathcal{F}_t \right]. \end{aligned}$$

For  $b > \gamma\sqrt{T}$ , applying Lemma 2.1, we have

$$\begin{aligned} \psi_a(|Y_t^{n,p}|) &\leq \psi_a \left( e^{\beta(T-t)} \right) \mathbb{E} \left[ \exp \left( \int_t^T q_s^{n,p} dW_s \right) \psi_a \left( |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \right) \middle| \mathcal{F}_t \right] \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \left( \mathbb{E} \left[ \exp \left( \frac{1}{2b^2} \left( \int_t^T q_s^{n,p} dW_s \right)^2 \right) \middle| \mathcal{F}_t \right] \right. \\ &\quad \left. + e^{2b^2} \mathbb{E} \left[ \psi_b \circ \psi_a \left( |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \right) \middle| \mathcal{F}_t \right] \right). \end{aligned}$$

Using Lemma 2.2 and Assertion (iii) of Proposition 2.3, we have for any  $c > 0$ ,

$$\begin{aligned} &\psi_a(|Y_t^{n,p}|) \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \left( \frac{1}{\sqrt{1 - \frac{\gamma^2}{b^2}(T-t)}} + e^{2b^2 + \frac{ab^2}{c}} \mathbb{E} \left[ \psi_{a+b+c} \left( |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \right) \middle| \mathcal{F}_t \right] \right). \end{aligned}$$

For  $\mu > \gamma\sqrt{T}$ , we can choose  $a > 0, b > \gamma\sqrt{T}$ , and  $c > 0$  such that  $a + b + c = \mu$ . Then, we have

$$\begin{aligned} &\psi_a(|Y_t^{n,p}|) \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \left( \frac{1}{\sqrt{1 - \frac{\gamma^2}{b^2}(T-t)}} + e^{2b^2 + \frac{ab^2}{c}} \mathbb{E} \left[ \psi_\mu \left( |\xi^{n,p}| + \int_t^T |f_0^{n,p}(s)| ds \right) \middle| \mathcal{F}_t \right] \right) \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \left( \frac{1}{\sqrt{1 - \frac{\gamma^2}{b^2}(T-t)}} + e^{2b^2 + \frac{ab^2}{c}} \mathbb{E} \left[ \psi_\mu \left( |\xi| + \int_t^T |f_0(s)| ds \right) \middle| \mathcal{F}_t \right] \right). \end{aligned}$$

Letting first  $n \rightarrow \infty$  and then  $p \rightarrow \infty$ , we have

$$\begin{aligned} &\psi_a(|Y_t|) \\ &\leq \psi_a \left( e^{\beta(T-t)} \right) \left( \frac{1}{\sqrt{1 - \frac{\gamma^2}{b^2}(T-t)}} + e^{2b^2 + \frac{ab^2}{c}} \mathbb{E} \left[ \psi_\mu \left( |\xi| + \int_t^T |f_0(s)| ds \right) \middle| \mathcal{F}_t \right] \right) \\ &\leq \psi_a \left( e^{\beta T} \right) \left( \frac{1}{\sqrt{1 - \frac{\gamma^2 T}{b^2}}} + e^{2b^2 + \frac{ab^2}{c}} \mathbb{E} \left[ \psi_\mu \left( |\xi| + \int_0^T |f_0(s)| ds \right) \middle| \mathcal{F}_t \right] \right). \end{aligned}$$

Consequently, we have  $\psi_a(|Y|)$  belongs to the class (D). □

Now we state our main result of this note.

**Theorem 2.5** Assume that the generator  $f$  is uniformly Lipschitz in  $(y, z)$ , i.e., there are  $\beta > 0$  and  $\gamma > 0$  such that for all  $(y^i, z^i) \in R \times R^{1 \times d}$ ,  $i = 1, 2$ , we have

$$|f(t, y^1, z^1) - f(t, y^2, z^2)| \leq \beta|y^1 - y^2| + \gamma|z^1 - z^2|.$$

Furthermore, assume that there exists  $\mu > \gamma\sqrt{T}$  such that  $\psi(|\xi| + \int_0^T |f(t, 0, 0)| dt, \mu) \in L^1(\Omega, P)$ . Then, BSDE (2.4) admits a unique solution  $(Y, Z)$  such that  $\psi(Y, a)$  belongs to the class (D) for some  $a > 0$ .

*Proof.* The existence of an adapted solution has been proved in the preceding theorem. It remains to prove the uniqueness.

For  $i = 1, 2$ , let  $(Y^i, Z^i)$  be a solution of BSDE (2.4) such that  $\psi_{a^i}(Y^i)$  belongs to the class (D) for some  $a^i > 0$ . Define

$$a := a^1 \wedge a^2, \quad \delta Y := Y^1 - Y^2, \quad \delta Z := Z^1 - Z^2.$$

Then both  $\psi_a(Y^1)$  and  $\psi_a(Y^2)$  are in the class (D), since  $\psi(x, \mu)$  is nondecreasing in  $\mu$ , and the pair  $(\delta Y, \delta Z)$  satisfies the following equation

$$\delta Y_t = \int_t^T [f(s, Y_s^1, Z_s^1) - f(s, Y_s^2, Z_s^2)] ds - \int_t^T \delta Z_s dW_s, \quad t \in [0, T].$$

By a standard linearization we see that there exists an adapted pair of processes  $(u, v)$  such that  $|u_s| \leq \beta, |v_s| \leq \gamma$ , and  $f(s, Y_s^1, Z_s^1) - f(s, Y_s^2, Z_s^2) = u_s \delta Y_s + \delta Z_s v_s$ .

We define the stopping times

$$\tau_n := \inf\{t \geq 0 : |Y_t^1| + |Y_t^2| \geq n\} \wedge T, \quad n = 1, 2, \dots,$$

with the convention that  $\inf \emptyset = \infty$ . Since  $(\delta Y, \delta Z)$  satisfies the linear BSDE

$$\delta Y_t = \int_t^T (u_s \delta Y_s + \delta Z_s v_s) ds - \int_t^T \delta Z_s dW_s, \quad t \in [0, T],$$

we have the following formula

$$\delta Y_{t \wedge \tau_n} = \mathbb{E} \left[ e^{\int_{t \wedge \tau_n}^{\tau_n} u_s ds + \int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle - \frac{1}{2} \int_{t \wedge \tau_n}^{\tau_n} |v_s|^2 ds} \delta Y_{\tau_n} \mid \mathcal{F}_t \right].$$

Therefore,

$$\begin{aligned} |\delta Y_{t \wedge \tau_n}| &\leq \mathbb{E} \left[ e^{\int_{t \wedge \tau_n}^{\tau_n} u_s ds + \int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle} |\delta Y_{\tau_n}| \mid \mathcal{F}_t \right] \\ &\leq e^{\beta T} \mathbb{E} \left[ e^{\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle} |\delta Y_{\tau_n}| \mid \mathcal{F}_t \right]. \end{aligned} \quad (2.6)$$

Now we show that the family of random variables  $e^{\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle} |\delta Y_{\tau_n}|$  is uniformly integrable. For this note that, thanks to Lemma 2.1,

$$e^{\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle} |\delta Y_{\tau_n}| \leq e^{\frac{1}{2a^2} (\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle)^2} + e^{2a^2} \psi_a(|\delta Y_{\tau_n}|). \quad (2.7)$$

For  $t \in [T - \frac{a^2}{4\gamma^2}, T]$ , we have from Lemma 2.2,

$$\mathbb{E} \left[ \left| e^{\frac{1}{2a^2} (\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle)^2} \right|^2 \right] = \mathbb{E} \left[ e^{\frac{1}{a^2} (\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle)^2} \right] \leq \frac{1}{\sqrt{1 - \frac{2\gamma^2}{a^2} (T - t)}} \leq \sqrt{2},$$

and, thus, the family of random variables  $e^{\frac{1}{2a^2}(\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle)^2}$  is uniformly integrable.

On the other hand, since  $\psi_a$  is nondecreasing and convex, we have thanks to Proposition 2.3 (ii)

$$\begin{aligned} \psi_a(|\delta Y_{\tau_n}|) &\leq \psi_a(|Y_{\tau_n}^1| + |Y_{\tau_n}^2|) = \psi_a\left(\frac{1}{2} \times 2|Y_{\tau_n}^1| + \frac{1}{2} \times 2|Y_{\tau_n}^2|\right) \\ &\leq \frac{1}{2}\psi_a(2|Y_{\tau_n}^1|) + \frac{1}{2}\psi_a(2|Y_{\tau_n}^2|) \leq \frac{1}{2}\psi_a(2)[\psi_a(|Y_{\tau_n}^1|) + \psi_a(|Y_{\tau_n}^2|)]. \end{aligned}$$

From (2.7) it now follows that, for  $t \in [T - \frac{a^2}{4\gamma^2}, T]$ , the family of random variables  $e^{\int_{t \wedge \tau_n}^{\tau_n} \langle v_s, dW_s \rangle} |\delta Y_{\tau_n}|$  is uniformly integrable.

Finally, letting  $n \rightarrow \infty$  in inequality (2.6), we have  $\delta Y = 0$  on the interval  $[T - \frac{a^2}{4\gamma^2}, T]$ . It is then clear that  $\delta Z = 0$  on  $[T - \frac{a^2}{4\gamma^2}, T]$ . The uniqueness of the solution is obtained on the interval  $[T - \frac{a^2}{4\gamma^2}, T]$ . In an identical way, we have the uniqueness of the solution on the interval  $[T - \frac{a^2}{2\gamma^2}, T - \frac{a^2}{4\gamma^2}]$ . By a finite number of steps, we cover in this way the whole interval  $[0, T]$ , and we conclude the uniqueness of the solution on the interval  $[0, T]$ .  $\square$

## References

- [1] P. Briand, B. Delyon, Y. Hu, E. Pardoux and L. Stoica,  $L^p$  solutions of backward stochastic differential equations. *Stochastic Process. Appl.* **108** (2003), no. 1, 109–129.
- [2] P. Briand and Y. Hu, BSDE with quadratic growth and unbounded terminal value. *Probab. Theory Related Fields* **136** (2006), no. 4, 604–618.
- [3] N. El Karoui, S. Peng and M. C. Quenez, Backward stochastic differential equations in finance. *Math. Finance* **7** (1997), no. 1, 1–71.
- [4] Y. Hu and S. Tang, Existence of solution to scalar BSDEs with  $L \exp\left(\sqrt{\frac{2}{\lambda} \log(1+L)}\right)$ -integrable terminal values. *Electron. Commun. Probab.* **23** (2018), Paper No. 27, 11pp.
- [5] J. P. Lepeltier and J. San Martin, Backward stochastic differential equations with continuous coefficient. *Statist. Probab. Lett.* **32** (1997), 425–430.
- [6] E. Pardoux and S. Peng, Adapted solution of a backward stochastic differential equation. *Systems Control Lett.* **14** (1990), no. 1, 55–61.