

Some useful lemmas on the edge Szeged index

Shengjie He^{1*}

1. Department of Mathematics, Beijing Jiaotong University, Beijing, 100044, China

Abstract

The edge Szeged index of a graph G is defined as $Sz_e(G) = \sum_{uv \in E(G)} m_u(uv|G)m_v(uv|G)$, where $m_u(uv|G)$ (resp., $m_v(uv|G)$) is the number of edges whose distance to vertex u (resp., v) is smaller than the distance to vertex v (resp., u), respectively. Recently, Wang et al. [G. Wang, S. Li, D. Qi, H. Zhang, On the edge-Szeged index of unicyclic graphs with given diameter, Appl. Math. Comput. 336 (2018) 94-106] identified the minimum edge-Szeged index of the n -vertex unicyclic graphs with a given diameter. By using another approach, we characterize the graph with minimum edge Szeged index among all the unicyclic graphs with given order and diameter in this paper.

Keywords: Edge Szeged index; Szeged index; Wiener index; Unicyclic graph; Diameter.

2010 MSC: 05C40, 05C90

1 Introduction

Throughout this paper, all graphs we considered are finite, undirected, and simple. Let G be a connected graph with vertex set $V(G)$ and edge set $E(G)$. For a vertex $u \in V(G)$, the degree of u , denote by $d_G(u)$, is the number of vertices which are adjacent to u . Call a vertex u a pendent vertex of G , if $d_G(u) = 1$ and call an edge uv a pendent edge of G , if $d_G(u) = 1$ or $d_G(v) = 1$. For any two vertices $u, v \in V(G)$, let $d(u, v|G)$ denote the distance between u and v in G . The diameter of a graph G is the maximum distance between pairs of vertices of G . Denote by P_n , S_n and C_n a path, star and cycle on n vertices, respectively. Let \mathbb{N} represent the set of natural numbers. If a and b are two natural numbers with $a \leq b$, $[a, b]$ denote the set $\{n \in \mathbb{N} | a \leq n \leq b\}$.

The Wiener index is one of the oldest and the most thoroughly studied topological indices. The Wiener index of a graph G is defined as

$$W(G) = \sum_{\{u,v\} \subseteq V(G)} d(u, v|G).$$

For any edge $e = uv$ of G , $V(G)$ can be partitioned into three sets by comparing with the distance of the vertex in $V(G)$ to u and v , and the three sets are as follows:

$$\begin{aligned} N_u(e|G) &= \{w \in V(G) : d(u, w|G) < d(v, w|G)\}, \\ N_v(e|G) &= \{w \in V(G) : d(v, w|G) < d(u, w|G)\}, \\ N_0(e|G) &= \{w \in V(G) : d(u, w|G) = d(v, w|G)\}. \end{aligned}$$

*Corresponding author. Emails: he1046436120@126.com (Shengjie He)

The number of vertices of $N_u(e|G)$, $N_v(e|G)$ and $N_0(e|G)$ are denoted by $n_u(e|G)$, $n_v(e|G)$ and $n_0(e|G)$, respectively. If G is a tree, then the formula $W(G) = \sum_{e=uv \in E(G)} n_u(e|G)n_v(e|G)$ gives a long time known property of the Wiener index. According to this result, a new topological index, named by Szeged index, was introduced by Gutman [4], which is an extension of the Wiener index and defined by

$$Sz(G) = \sum_{e=uv \in E(G)} n_u(e|G)n_v(e|G).$$

For other results on the Wiener and Szeged indices, we refer to [3, 8, 16, 17].

An edge version of the Wiener index (resp., Szeged index) is also introduced, named edge Wiener index (resp., edge Szeged index). If $e = uv$ and $f = ab$ are two edges of G and w is a vertex of G , then the distance between e and w is defined as $d(e, w|G) = \min\{d(u, w|G), d(v, w|G)\}$ and the distance between e and f is defined as $d(e, f|G) = \min\{d(u, f|G), d(v, f|G)\}$. The edge Wiener index [2] is defined as follows:

$$W_e(G) = \sum_{\{e, f\} \subseteq E(G)} d(e, f|G).$$

For $e = uv \in E(G)$, let $M_u(e|G)$ be the set of edges whose distance to the vertex u is smaller than the distance to the vertex v , and $M_v(e|G)$ be the set of edges whose distance to the vertex v is smaller than the distance to the vertex u . Set $m_u(e|G) = |M_u(e|G)|$ and $m_v(e|G) = |M_v(e|G)|$. Gutman and Ashrafi [5] introduced an edge version of the Szeged index, named edge Szeged index. The edge Szeged index of G is defined as

$$Sz_e(G) = \sum_{uv \in E(G)} m_u(uv|G)m_v(uv|G).$$

In [5], some basic properties of the edge Szeged index were established by Gutman. It can be checked that the pendent edges make no contributions to the edge Szeged index of a graph. Nadjafi-Arani et al. [12] proved that for every connected graph G , $Sz_e(G) \geq W_e(G)$ with equality if and only if G is a tree. A unicyclic graph of order n is a connected graph with n edges. In [1], Cai and Zhou determined the n -vertex unicyclic graphs with the largest, the second largest, the smallest and the second smallest edge Szeged indices, respectively. Li [9] established a relation between the edge Szeged index and Szeged index of the unicyclic graphs.

Let $\mathcal{U}_{n,d}$ be the set of unicyclic graphs of order n with diameter d for $2 \leq d \leq n - 2$. Tan [14] and Shi [13] independently determined the graph in $\mathcal{U}_{n,d}$ with minimum Wiener index. Liu et al. [11] characterized the graph in $\mathcal{U}_{n,d}$ with minimum Szeged index. Recently, Wang et al. [15] identified the minimum edge-Szeged index of the n -vertex unicyclic graphs with a given diameter. By using another approach, we determine the graph in $\mathcal{U}_{n,d}$ with minimum edge Szeged index in this paper. In order to state our results, we need more notations.

Suppose that l_1 , l_2 and a are three nonnegative integers. Let $P' = P_{l_1+1}$ be a path on $l_1 + 1$ vertices with a terminal vertex u and $P'' = P_{l_2+1}$ be a path on $l_2 + 1$ vertices with a terminal vertex v . Let $S' = S_{a+1}$ be a star on $a + 1$ vertices with center vertex w . We denote by P_{l_1, l_2}^a the tree obtained from P' , P'' and S' by identifying u , v and w to u' , and call u' the root vertex of P_{l_1, l_2}^a (see Fig. 1). For convenience, we write $P_{l_1}^a$ for $P_{l_1, 0}^a$ (see Fig. 1). Note that $P_{0, 0}^a \cong S_{a+1}$.




Fig. 1 Graphs P_{l_1, l_2}^a and $P_{l_1}^a$

Let $C_3(P_{\lceil \frac{d-1}{2} \rceil}^{n-d-2}, P_{\lfloor \frac{d-1}{2} \rfloor+1}, S_1)$ be the graph obtained from the cycle $C_3 = v_1 v_2 v_3 v_1$ by identifying the root vertex of $P_{\lceil \frac{d-1}{2} \rceil}^{n-d-2}$ with v_1 , and identifying one end vertex of $P_{\lfloor \frac{d-1}{2} \rfloor+1}$ with v_2 (see Fig. 2).




Fig. 2 Graphs $C_3(P_{\lceil \frac{d-1}{2} \rceil}^{n-d-2}, P_{\lfloor \frac{d-1}{2} \rfloor+1}, S_1)$ and $T_{n, d, \lfloor \frac{d}{2} \rfloor}$

Let G be a unicyclic graph of order n with diameter d . Then $1 \leq d \leq n-2$. If $d=1$, then $G \cong C_3$. If $d=2$, then $G \in \{C_4, C_3(P_1^0, S_1, S_1)\}$ for $n=4$; $G \in \{C_5, C_3(P_1^1, S_1, S_1)\}$ for $n=5$ and $G = C_3(P_1^{n-4}, S_1, S_1)$ for $n \geq 6$. For $1 \leq d \leq 2$, it is not difficult to determine $Sz_e(G)$. Hence throughout the paper, we assume that $n \geq 6$ and $3 \leq d \leq n-2$. Our main result is the following theorem.

Theorem 1.1. *Among the unicyclic graphs of order $n \geq 6$ with diameter d for $3 \leq d \leq n-2$, $C_3(P_{\lceil \frac{d-1}{2} \rceil}^{n-d-2}, P_{\lfloor \frac{d-1}{2} \rfloor+1}, S_1)$ is the unique graph with minimum edge Szeged index.*

The rest of this paper is organized as follows. In Section 2, we present some useful lemmas. In Section 3, we establish some transformations of graphs which keep the diameter and the order of the graphs, but decrease the edge Szeged index of the graphs. In section 4, Theorem 1.1 is proved.

2 Useful lemmas

We first introduce a result useful for computing the edge Szeged index of a unicyclic graph. Let $g \geq 3$ be an integer, and let $C_g = v_1 v_2 \cdots v_g v_1$ be a cycle on g vertices. Let T_1, T_2, \dots, T_g be vertex-disjoint trees, and let the root vertex of T_i be u_i for $1 \leq i \leq g$. Denote by $C_{T_1, T_2, \dots, T_g}^{u_1, u_2, \dots, u_g}$ the unicyclic graph obtained from C_g by identifying the root vertex u_i of T_i with v_i for each $1 \leq i \leq g$. If there is no confusion, we write $C_g(T_1, T_2, \dots, T_g)$ for $C_{T_1, T_2, \dots, T_g}^{u_1, u_2, \dots, u_g}$. Any n -vertex unicyclic graph G with a g -cycle is of the form $C_g(T_1, T_2, \dots, T_g)$, where $\sum_{i=1}^g |T_i| = n = \sum_{i=1}^g |E(T_i)| + g$. For $v \in V(G)$, we define

$$D(v|G) = \sum_{w \in V(G)} d(v, w|G).$$

For an integer g , define

$$\delta(g) = \begin{cases} 1, & \text{if } g \text{ is odd;} \\ 0, & \text{if } g \text{ is even.} \end{cases}$$

Lemma 2.1. [9] Let $G = C_r(T_1, T_2, \dots, T_r)$ with $|V(G)| = n$. Then

$$Sz_e(G) = Sz(G) + \sum_{i=1}^r D(v_i|T_i) - n^2 + \begin{cases} nr & \text{if } r \text{ is odd;} \\ r & \text{if } r \text{ is even.} \end{cases}$$

The following are some lemmas which will be used in our proof.

Lemma 2.2. [7] Let G and G' be the graphs shown as in Fig. 3, where G consists of G_0 and G_1 with a common vertex u , and G' consists of G_0 and G_2 with a common vertex u . Then each of the followings holds:

(i) For any edge $e = w_1w_2 \in E(G_0)$ and $1 \leq i \leq 2$, we have

$$m_{w_i}(e|G) = m_{w_i}(e|G_0) + \tau(u)|E(G_1)|,$$

where

$$\tau(u) = \begin{cases} 1, & u \in N_{w_i}(e|G_0); \\ 0, & \text{otherwise.} \end{cases}$$

(ii) If $|E(G_1)| = |E(G_2)|$, then

$$\sum_{e=w_1w_2 \in E(G_0)} m_{w_1}(e|G)m_{w_2}(e|G) = \sum_{e=w_1w_2 \in E(G_0)} m_{w_1}(e|G')m_{w_2}(e|G').$$




Fig. 3. G and G' in Lemma 2.2

Lemma 2.3. [7] Let G be a graph of order n with a cycle $C_g = v_1v_2 \dots v_gv_1$. Assume that $G - E(C_g)$ has exactly g components G_1, G_2, \dots, G_g , where G_i is the component of $G - E(C_g)$ that contains v_i for $1 \leq i \leq g$. Let

$$G' = G - \cup_{i=2}^g \{wv_i : w \in N_{G_i}(v_i)\} + \cup_{i=2}^g \{wv_1 : w \in N_{G_i}(v_i)\}.$$

Then $Sz_e(G') \leq Sz_e(G)$ with equality if and only if C_g is an end-block, that is, $G \cong G'$.

Lemma 2.4. [5] If T is a tree with order n , then $Sz_e(T) = Sz(T) - (n - 1)^2$.

Note that if T is a tree, $W_e(T) = Sz_e(T)$ and $W(T) = Sz(T)$. The conclusion of Lemma 2.4 can also be written as $W_e(T) = W(T) - (n - 1)^2$.

Let $n \geq 3$ and $2 \leq d \leq n - 1$ be integers. Let $P_{d+1} = u_0u_1 \dots u_d$ be a path of order $d + 1$ and $T_{n,d, \lfloor \frac{d}{2} \rfloor}$ be the tree obtained from P_{d+1} by attaching $n - d - 1$ pendent vertices to $u_{\lfloor \frac{d}{2} \rfloor}$ (see Fig. 2). Note that $T_{n,d, \lfloor \frac{d}{2} \rfloor} \cong P_{\lfloor \frac{d}{2} \rfloor, \lceil \frac{d}{2} \rceil}^{n-d-1}$.

Lemma 2.5. [10] Among the trees of order n with diameter $2 \leq d \leq n - 2$, $T_{n,d, \lfloor \frac{d}{2} \rfloor}$ has minimum Wiener index.

Lemma 2.6. [11] Let $C_g = v_1v_2 \cdots v_gv_1$ and $d_{i,j} = d(v_i, v_j|C_g)$ for $i, j \in \{1, 2, \dots, g\}$.

(i) If g is an even number,

$$(d_{2,j} - d_{1,j} + 1, d_{g,j} - d_{1,j} + 1) = \begin{cases} (0, 2), & \text{if } 2 \leq j \leq \frac{g}{2}; \\ (0, 0), & \text{if } j = \frac{g}{2} + 1; \\ (2, 0), & \text{if } \frac{g}{2} + 2 \leq j \leq g - 1. \end{cases}$$

(ii) If g is an odd number,

$$(d_{2,j} - d_{1,j} + 1, d_{g,j} - d_{1,j} + 1) = \begin{cases} (0, 2), & \text{if } 2 \leq j \leq \frac{g-1}{2}; \\ (0, 1), & \text{if } j = \frac{g+1}{2}; \\ (1, 0), & \text{if } j = \frac{g+3}{2}; \\ (2, 0), & \text{if } \frac{g+5}{2} \leq j \leq g - 1. \end{cases}$$

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Nos. 11371052, 11771039), the Fundamental Research Funds for the Central Universities (Nos. 2016JBM071, 2016JBZ012) and the 111 Project of China (B16002).

References

- [1] X. Cai, B. Zhou, Edge Szeged index of unicyclic graphs, *MATCH Commun. Math. Comput. Chem.* 63 (2010) 133–144.
- [2] P. Dankelmann, I. Gutman, S. Mukwembi, H. C. Swart, The edge-Wiener index of a graph, *Discrete Math.* 309 (2009) 3452–3457.
- [3] A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: theory and applications, *Acta Appl. Math.* 66 (2001) 211–249.
- [4] I. Gutman, A formula for the Wiener number of trees and its extension to graphs containing cycles, *Graph Theory Notes N. Y.* 27 (1994) 9–15.
- [5] I. Gutman, A. R. Ashrafi, The edge version of the Szeged index, *Croat. Chem. Acta.* 81 (2) (2008) 263–266.
- [6] I. Gutman, L. Popović, P. V. Khadikar, S. Karmarkar, S. Joshi, M. Mandloi, Relations between Wiener and Szeged indices of monocyclic molecules, *MATCH Commun. Math. Comput. Chem.* 35 (1997) 91–103.
- [7] S. He, R.-X. Hao, A. Yu, On extremal cacti with respect to the edge Szeged index and edge-vertex Szeged index, arXiv:1711.02394v1 [math.CO] 7 Nov 2017.
- [8] P. Khadikar, P. Kale, N. Deshpande, S. Karmarkar, V. Agrawal, Szeged indices of hexagonal chains, *MATCH Commun. Math. Comput. Chem.* 43 (2000) 7–15.
- [9] J. Li, A relation between the edge Szeged index and the ordinary Szeged index, *MATCH Commun. Math. Comput. Chem.* 70 (2013) 621–625.
- [10] H. Liu, X. Pan, On the Wiener index of trees with fixed diameter, *MATCH Commun. Math. Comput. Chem.* 60 (2008) 85–94.
- [11] Y. Liu, A. Yu, M. Lu, R.-X. Hao, On the Szeged index of unicyclic graphs with given diameter, *Discrete Appl. Math.* 233 (2017) 118–130.

- [12] M. J. Nadjafi-Arani, H. Khodashenas, A. R. Ashrafi, Relationship between edge Szeged and edge Wiener indices of graphs, *Glas. Mat.* 47 (67) (2012) 21–29.
- [13] C. Ren, J. Shi, On the Wiener index of unicyclic graphs with fixed diameter, *J. East China Univ. Sci. Technol.* 39 (2013) 768–772.
- [14] S. Tan, The minimum Wiener index of unicyclic graphs with a fixed diameter, *J. Appl. Math. Comput.* (2016) 1–22.
- [15] G. Wang, S. Li, D. Qi, H. Zhang, On the edge-Szeged index of unicyclic graphs with given diameter, *Appl. Math. Comput.* 336 (2018) 94-106.
- [16] H. Zhang, S. Li, L. Zhao, On the further relation between the (revised) Szeged index and the Wiener index of graphs, *Discrete Appl. Math.* 206 (2016) 152–164.
- [17] B. Zhou, X. Cai, Z. Du, On Szeged indices of unicyclic graphs, *MATCH Commun. Math. Comput. Chem.* 63 (2010) 113–132.