

Octonions, exceptional Jordan algebra, and the role of the group F_4 in particle physics¹

Ivan Todorov and Svetla Drenska

Institute for Nuclear Research and Nuclear Energy
 Bulgarian Academy of Sciences
 Tsarigradsko Chaussee 72, BG-1784 Sofia, Bulgaria
 e-mail: ivbortodorov@gmail.com, sdren@abv.bg

Abstract

Normed division rings are reviewed in the more general framework of composition algebras that include the split (indefinite metric) case. The Jordan - von Neumann - Wigner classification of finite euclidean Jordan algebras is outlined with special attention to the 27 dimensional exceptional Jordan algebra \mathcal{J} . The automorphism group F_4 of \mathcal{J} and its maximal Borel - de Siebenthal subgroups $\frac{SU(3) \times SU(3)}{\mathbb{Z}_3}$ and Spin(9) are studied in detail and applied to the classification of fundamental fermions and gauge bosons. Their intersection in F_4 is demonstrated to coincide with the gauge group of the Standard Model of particle physics. The first generation's fundamental fermions form a basis of primitive idempotents in the euclidean extension of the Jordan subalgebra J_2^8 of \mathcal{J} .

Keywords: Automorphism groups, Clifford algebras, Exceptional Jordan algebra, Octonions, Standard Model.

¹Extended version of lectures presented by I.T. at the Institute for Nuclear Research and Nuclear Energy during the spring of 2017 (see [TD17]) and in May 2018 (published in: *Adv. Appl. Clifford Alg.* **28** (2018) 82).

Contents

1	Composition and Clifford algebras	2
1.1	Normed alternative algebras ([R08] Sect.1)	2
1.2	Relation to Clifford algebras. Classification.	6
1.3	Historical note	8
2	Octonions. Isometries and automorphisms	10
2.1	Eight dimensional alternative algebras	10
2.2	Isometry group of the (split) octonions. Triality	12
2.3	Automorphism group G_2 of the octonions.	16
2.4	Roots and weights of $\mathfrak{g}_2 \subset \mathfrak{B}_3 \subset \mathfrak{D}_4$	18
3	Jordan algebras and related groups	21
3.1	Classification of finite dimensional Jordan algebras	21
3.2	The Tits-Kantor-Koecher (TKK) construction	24
3.3	Automorphism groups of the exceptional Jordan algebras $\mathcal{H}_3(\mathbb{O}_{(s)})$ and their maximal subgroups	26
4	F_4 as a grand unified symmetry of the standard model	30
4.1	The Jordan subalgebra $JSpin_9$ of $\mathcal{H}_3(\mathbb{O})$	30
4.2	The representation $\underline{26}$ of F_4	34
4.3	The symmetry group of the standard model	37
5	Extended euclidean Jordan algebra; the 32 primitive idem- potents of the fermion states	38
5.1	Euclidean extension of J_2^8 . Fermion Fock space	39
5.2	Fermion state space in terms of primitive idempotents	40
5.3	Nonabelian superselection sectors	41
6	Concluding remarks	43
	References	48

Introduction

Division and Clifford algebras were introduced in 19th century with an eye for applications in geometry and physics (for a historical survey see the last chapter of [L]). Pascual Jordan introduced and studied his algebras in the 1930's in order to describe observables in quantum mechanics (for "a taste of Jordan algebras" see [McC] along with the original paper [JvNW]). Yet, the first serious applications of these somewhat exotic structures appeared (in mid-twentieth century) in pure mathematics: in the theory of exceptional Lie groups and symmetric spaces (cf. [F] as well as the later surveys [A, K, BS, R08]), in topology [ABS]. For an entertaining review on division algebras and quantum theory - see [B12]. Possible applications to particle physics were first advocated by Feza Gürsey and his students in the 1970's - see his lecture and his posthumous book (with C.-H.Tze) [G] and references therein). They continue in various guises to attract attention until these days, never becoming a mainstream activity. The present lectures are meant as a background for the ongoing work [DV, TD, D-V.T] centered on the exceptional 27 dimensional Jordan algebra $\mathcal{J} = \mathcal{H}_3(\mathbb{O})$. In Chapter 4 we elaborate on a possible application of the automorphism group F_4 of $\mathcal{J} = J_3^8$ as a novel "grand unified symmetry" of the Standard Model. In Chapter 5 we apply the notion of extended special euclidean Jordan algebra of [D-V.T] to the spin factor J_2^8 and put forward a fermion Fock space construction, inspired by [F18], of the first generation of leptons and quarks. Although such a proposal of an "exceptional finite quantum geometry" is still tentative, we feel that it is worth pursuing.² In any case, the mathematical background which is the main subject of these notes is sound and beautiful - and deserves to be known by particle theorists.

1 Composition and Clifford algebras

1.1 Normed alternative algebras ([R08] Sect.1)

A composition (or Hurwitz) algebra \mathcal{A} is a vector space over a field $\mathbb{K} = (\mathbb{R}, \mathbb{C}, \dots)$ equipped with a bilinear (not necessarily associative) product xy with a unit 1 ($1x = x1 = x$) and a nondegenerate quadratic form $N(x)$, the

²For related attempts to provide an algebraic counterpart of the Standard Model of particle physics see [D04, D10, D14, CGKK, F16, S18, F18] and references therein.

norm, satisfying

$$N(xy) = N(x)N(y), \quad N(\lambda x) = \lambda^2 N(x) \text{ for } x \in \mathcal{A}, \lambda \in \mathbb{K}. \quad (1.1)$$

The norm allows to define by polarization a *symmetric bilinear form* $\langle x, y \rangle$ setting:

$$2 \langle x, y \rangle = N(x + y) - N(x) - N(y) (= 2 \langle y, x \rangle). \quad (1.2)$$

(Nondegeneracy of N means that if $\langle x, y \rangle = 0$ for all $y \in \mathcal{A}$ then $x = 0$.) By repeated polarization of the identity $\langle xy, xy \rangle = \langle x, x \rangle \langle y, y \rangle$ one obtains

$$\langle ab, ac \rangle = N(a) \langle b, c \rangle = \langle ba, ca \rangle \quad (1.3)$$

$$\langle ac, bd \rangle + \langle ad, bc \rangle = 2 \langle a, b \rangle \langle c, d \rangle. \quad (1.4)$$

Setting in (1.4) $a = c = x$, $b = 1$, $d = y$ and using (1.3) we find:

$$\langle x^2 + N(x)1 - t(x)x, y \rangle = 0,$$

where $t(x) := 2 \langle x, 1 \rangle$ is by definition the *trace*, or, using the non-degeneracy of the form \langle, \rangle ,

$$x^2 - t(x)x + N(x)1 = 0, \quad t(x) = 2 \langle x, 1 \rangle. \quad (1.5)$$

Thus every $x \in \mathcal{A}$ satisfies a quadratic relation with coefficients the trace $t(x)$ and the norm $N(x)$ (a linear and a quadratic scalar functions) taking values in \mathbb{K} .

The trace functional (1.5) allows to introduce *Cayley conjugation*,

$$x \rightarrow x^* = t(x) - x, \quad t(x) = t(x)1 \in \mathcal{A} \quad (1.6)$$

an important tool in the study of composition algebras. It is an (orthogonal) reflection ($\langle x^*, y^* \rangle = \langle x, y \rangle$) that leaves the scalars $\mathbb{K}1$ invariant (in fact, $t(\lambda 1) = 2\lambda$ implying $(\lambda 1)^* = \lambda 1$ for $\lambda \in \mathbb{K}$). It is also an involution and an antihomomorphism:

$$(x^*)^* = x, \quad (xy)^* = y^* x^*. \quad (1.7)$$

Furthermore Eqs.(1.5) and (1.6) allow to express the trace and the norm as a sum and a product of x and x^* :

$$t(x) = x + x^*, \quad N(x) = xx^* = x^*x = N(x^*).$$

The relation (1.4) allows to deduce

$$\langle ax, y \rangle = \langle x, a^*y \rangle, \quad \langle xa, y \rangle = \langle x, ya^* \rangle. \quad (1.8)$$

From these identities it follows $\langle ab, 1 \rangle = \langle a, b^* \rangle = \langle ba, 1 \rangle$, hence, the trace is commutative:

$$t(ab) = \langle b, a^* \rangle = \langle a, b^* \rangle = t(ba). \quad (1.9)$$

Similarly, one proves that t is associative and symmetric under cyclic permutations.

$$t((ab)c) = t(a(bc)) =: t(abc) = t(cab) = t(bca). \quad (1.10)$$

Moreover, using the quadratic relation (1.5) and the above properties of the trace one proves the identities that define an *alternative algebra*:

$$x^2y = x(xy), \quad yx^2 = (yx)x \quad (1.11)$$

(see Sect.1 of [R08] for details). The conditions (1.11) guarantee that the *associator*

$$[x, y, z] = (xy)z - x(yz) \quad (1.12)$$

changes sign under odd permutations (and is hence preserved by even, cyclic, permutations). This implies, in particular, the *flexibility conditions*.

$$(xy)x = x(yx). \quad (1.13)$$

An unital alternative algebra with an involution $x \rightarrow x^*$ satisfying (1.7) is a composition algebra if the norm N and the trace t defined by (1.9) are scalars (i.e. belong to $\mathbb{K}(= \mathbb{K}1)$) and the norm is non-degenerate.

Given a finite dimensional composition algebra \mathcal{A} Cayley and Dickson have proposed a procedure to construct another composition algebra \mathcal{A}' with twice the dimension of \mathcal{A} . Each element x of \mathcal{A}' is written in the form

$$x = a + eb, \quad a, b \in \mathcal{A} \quad (1.14)$$

where e is a new "imaginary unit" such that

$$e^2 = -\mu, \mu \in \{1, -1\} (\mu^2 = 1). \quad (1.15)$$

Thus \mathcal{A} appears as a subalgebra of \mathcal{A}' . The product of two elements $x = a + eb, y = c + ed$ of \mathcal{A}' is defined as

$$xy = ac - \mu d\bar{b} + e(\bar{a}d + cb) \quad (1.16)$$

where $a \rightarrow \bar{a}$ is the Cayley conjugation in \mathcal{A} . (The order of the factors becomes important, when the product in \mathcal{A} is noncommutative.) The Cayley conjugation $x \rightarrow x^*$ and the norm $N(x)$ in \mathcal{A}' are defined by:

$$\begin{aligned} x^* &= (a + eb)^* = \bar{a} + \bar{b}e^* = \bar{a} - \bar{b}e = \bar{a} - eb \\ N(x) &= xx^* = a\bar{a} + \mu b\bar{b} = x^*x. \end{aligned} \quad (1.17)$$

Let us illustrate the meaning of (1.16) and (1.17) in the first simplest cases.

For $\mathcal{A} = \mathbb{R}$, $\bar{a} = a$, Eq.(1.16) coincides with the definition of complex numbers for $\mu = 1$ ($e = i$) and defines the split complex numbers for $\mu = -1$. Taking next $\mathcal{A} = \mathbb{C}$ and $\mu = 1$ we can identify \mathcal{A}' with a 2×2 matrix representations setting

$$\mathbf{a} = a_0 + e_1 a_1 = a_0 + i\sigma_3 a_1 = \begin{pmatrix} a & 0 \\ 0 & \bar{a} \end{pmatrix} \quad (a = a_0 + ia_1) \quad (1.18)$$

$$x = \mathbf{a} + e\mathbf{b}, \quad e = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \Rightarrow x = \begin{pmatrix} a & -\bar{b} \\ b & \bar{a} \end{pmatrix}, \quad \mathbf{b} = b_0 + e_1 b_1.$$

Anticipating Baez *Fano plane* [B] notation for the octonion imaginary units (see Appendix A) we shall set $e = e_4$, $e_4 e_1 = e_2$ ($= i\sigma_1$).

It is then easily checked that the multiplication law (1.16) reproduces the standard matrix multiplication, the Cayley conjugation $x \rightarrow x^*$ coincides with the hermitian conjugation of matrices, while the norm $N(x)$ in \mathcal{A}' is given by the determinant:

$$\mathbb{H} = \{x \in \mathbb{C}[2]; \quad xx^* = \det x (\geq 0)\}. \quad (1.19)$$

Similarly, starting with the split complex numbers, we can write

$$\mathbf{a}_s = \mathbf{a}_0 + \tilde{\mathbf{e}}_1 \mathbf{a}_1, \quad \tilde{\mathbf{e}}_1 = \sigma_3 \iff \mathbf{a}_s = \begin{pmatrix} a_s & 0 \\ 0 & \bar{a}_s \end{pmatrix} \quad (\mathbf{a}_s = \mathbf{a}_0 + \mathbf{a}_1, \quad \bar{\mathbf{a}}_s = \mathbf{a}_0 - \mathbf{a}_1).$$

and choosing the same e as above we can identify the *split quaternions* \mathbb{H}_s with real 2×2 matrices:

$$\mathbb{H}_s = \left\{ x = \begin{pmatrix} a_s & -\bar{b}_s \\ b_s & \bar{a}_s \end{pmatrix} \in \mathbb{R}[2], \quad x^* = \begin{pmatrix} \bar{a}_s & \bar{b}_s \\ -b_s & a_s \end{pmatrix}, \quad xx^* = \det x \right\} \quad (1.20)$$

its norm having signature $(2, 2)$.

The next step in Cayley-Dickson construction gives the octonions, which have a nonassociative (but alternative) multiplication and thus do not have matrix realization.

1.2 Relation to Clifford algebras. Classification.

Given a composition algebra \mathcal{A} we define subspace $\mathcal{A}_0 \subset \mathcal{A}$ of pure imaginary elements with respect to the Cayley conjugation (1.6):

$$\mathcal{A}_0 = \{y \in \mathcal{A}; \quad y^* = -y\}. \quad (1.21)$$

It is a subspace of co-dimension one, orthogonal to the unit 1 of \mathcal{A} . For any $x \in \mathcal{A}$ we define its imaginary part as

$$x_0 = \frac{1}{2}(x - x^*) = x - \langle x, 1 \rangle \Rightarrow \langle x_0, 1 \rangle = 0. \quad (1.22)$$

From the expression $N(x) = xx^*$ (1.8) and from the defining property (1.21) of imaginary elements it follows that

$$x_0 \in \mathcal{A}_0 \Rightarrow x_0^2 = -N(x_0). \quad (1.23)$$

In other words, if the composition algebra \mathcal{A} is n -dimensional then its $(n-1)$ -dimensional subalgebra \mathcal{A}_0 gives rise to a Clifford algebra. If the norm N is positive definite then ${}^3 \mathcal{A}_0 = Cl(0, n-1) = Cl_{(1-n)}$. In the case of *split* complex numbers, quaternions and octonions one encounters instead the algebras $Cl_1 \equiv Cl(1, 0)$, $Cl(2, 1)$ and $Cl(4, 3)$, respectively.

It turns out that the classification of the Clifford algebras $Cl_{(1-n)}$ implies the classification of normed division rings of dimension n . So we recall it in

n	$Cl_{(1-n)}$	Irreducible spinor	n	$Cl_{(1-n)}$	Irreducible spinor
1	\mathbb{R}	$S_1 = \mathbb{R}$	5	$\mathbb{H}[2]$	$S_5 = \mathbb{H}^2$
2	\mathbb{C}	$S_2 = \mathbb{C}$	6	$\mathbb{C}[4]$	$S_6 = \mathbb{C}^4$
3	\mathbb{H}	$S_3 = \mathbb{H}$	7	$\mathbb{R}[8]$	$S_7 = \mathbb{R}^8$
4	$\mathbb{H} \oplus \mathbb{H}$	$S_4^+ = \mathbb{H}, S_4^- = \mathbb{H}$	8	$\mathbb{R}[8] \oplus \mathbb{R}[8]$	$S_8^+ = \mathbb{R}^8, S_8^- = \mathbb{R}^8$

Table 1: Irreducible spinors in the Clifford algebras $Cl_{(1-n)}$.

Table 1. Here we use the notation $\mathbb{A}[n]$ for the algebra of $n \times n$ matrices with entries in the (associative) algebra \mathbb{A} . As discovered by Élie Cartan in 1908 $Cl_{(-\nu-8)} = Cl_{-\nu} \otimes \mathbb{R}[16]$ so that the above Table 1 suffices to reconstruct all Clifford algebras of type $Cl_{-\nu}$. We see that the (real) dimension of the irreducible representation of $Cl_{(1-n)}$ coincides with n for $n = 1, 2, 4, 8$ only, thus implying Hurwitz theorem (see [B] Theorem 1 and the subsequent discussion).

Proceeding to the split alternative composition algebras we note that the type of $Cl(p, q)$ only depends on the signature $p - q$ which is 1 (similar to -7 modulo 8); we have: $Cl(1, 0) = \mathbb{R} \oplus \mathbb{R}$, $Cl(2, 1) = \mathbb{R}[2] \oplus \mathbb{R}[2]$, $Cl(4, 3) = \mathbb{R}[8] \oplus \mathbb{R}[8]$, for all above cases

$$Cl(p, p - 1) \cong \mathbb{R}[2^{(p-1)}] \oplus \mathbb{R}[2^{(p-1)}]. \quad (1.24)$$

We note here the difference in the treatment of the representations of $Cl(p, p-1)$ in the cases $p = 1, 2$, in which we are dealing with real associative composition algebras \mathbb{C}_s and \mathbb{H}_s , and $p = 4$ of the split octonions. In the associative case we deal with the action of $Cl(p, p-1)$ on the direct sum $\mathbb{R}^n \oplus \mathbb{R}^n$, $n = 2^{(p-1)}$ (for $p = 1, 2$) while in the non-associative case it acts on the irreducible subspace \mathbb{R}^n ($n = 8$), thus again fitting the dimension of the corresponding alternative algebra.

Remark 1.1. The *spinors* S_n are here understood as quantities transforming under the lowest order faithful irreducible representation of the (compact) group $\text{Spin}(n)$ which consists of the norm one even elements of Cl_{-n} . In fact, the even part $Cl_0(p, q)$ of $Cl(p, q)$ is isomorphic, for $q > 0$, to $Cl(p, q-1)$. $\text{Spin}(n)$ is the double cover of the rotation group $\text{SO}(n)$. The group of all norm one elements $Cl_{(1-n)}$ is the double cover $\text{Pin}(n-1)$ of the full orthogonal group $\text{O}(n-1)$ and its irreducible representations are called "pinors" -

³We adopt the sign convention of [L], [G], [T]; the opposite sign convention, $Cl_{(n-1)}$ for the positive definite $N(x)$, is used e.g. in [B].

see [B], (Sect.2.3).

In summary, the alternative algebras are classified as follows ([R08] Proposition 1.6):

Theorem 1 *Let (\mathcal{A}, N) be a composition algebra. For $\mu = \pm 1$, denote by $\mathcal{A}(\mu)$ the algebra $\mathcal{A}(\mu) = \mathcal{A} \oplus e\mathcal{A}$ with $e^2 = -\mu$ and product (1.16). Then*

- $\mathcal{A}(\mu)$ is commutative iff $\mathcal{A} = \mathbb{K}$;
- $\mathcal{A}(\mu)$ is associative iff \mathcal{A} is associative and commutative;
- $\mathcal{A}(\mu)$ is alternative iff \mathcal{A} is associative.

Theorem 2 ([R08] Theorems (1.7)-(1.10)). *A composition algebra is, as a vector space, 1, 2, 4 or 8 dimensional. There are four composition algebras \mathcal{A}_j over \mathbb{C} of dimension 2^j , $j = 0, 1, 2, 3$. There are seven composition algebras over \mathbb{R} ; the division algebras \mathcal{A}_j^+ , ($j = 0, 1, 2, 3$) with $N(x) \geq 0$ and $x^{-1} = \frac{x^*}{N(x)}$ for $x \neq 0$, and the split algebras \mathcal{A}_j^s , $j = 1, 2, 3$ of signature $(2^{j-1}, 2^{j-1})$.*

All above algebras are unique up to isomorphism. The multiplication rule (1.16) varies in different expositions. Different conventions are related by algebra automorphisms. (Our notation differs from Roos [R08] only by the sign of μ , as we set $e^2 = -\mu$.) The only nontrivial automorphism of the algebra of complex numbers is the complex conjugation. The automorphism group of the (real) quaternions is $SO(3)$ realized by

$$x \rightarrow uxu^*, \quad u \in SU(2) \ (u^* = u^{-1}). \quad (1.25)$$

Similarly, the automorphism group of the split quaternions is $SO(2, 1)$:

$$\mathbb{H} \ni x \rightarrow gxg^{-1}, \quad g \in SL(2, \mathbb{R}). \quad (1.26)$$

We shall survey the octonions and their automorphisms in the next section.

1.3 Historical note

The simplest relation of type (1.1), the one applicable to the absolute value square of a product of complex numbers

$$(xu - yv)^2 + (xv + yu)^2 = (x^2 + y^2)(u^2 + v^2)$$

$(x, y, u, v \in \mathbb{R})$, was found by Diophantus of Alexandria around 250. A more general relation of this type,

$$(xu + Dyv)^2 - D(xv + yu)^2 = (x^2 - Dy^2)(u^2 - Dv^2)$$

occurs for special values of D in Indian mathematics (Brahmegupta 598) - see [B61] Sect.2. For D positive it applies to the *split complex numbers*. The geometric interpretation by Gauss comes much later. (The fact that complex numbers are useful and should be taken seriously is sometimes attributed to Gerolamo Cardano (1501-1576), whose book *Ars Magna* (The Great Art) contains the solution of the cubic equation. In fact, it was his contemporary, Bologna's mathematician Rafael Bombelli (1526-1572) who first thoroughly understood the complex numbers and described them in his *L'Algebra*, published in 1572.)

The multiplicativity of the norm of the quaternions was noted by Euler in 1748, a century before Hamilton discovered the algebra of quaternions in 1843 (when "in a famous act of a mathematical vandalism, he carved the equations $i^2 = j^2 = k^2 = ijk = -1$ into the stone of Brougham Bridge" [B] p. 145). The corresponding relation for the octonions was discovered by the Danish mathematician Degen in 1818 - again before the discovery of the octonions (which took place in late 1843 - in a letter to Hamilton by his college friend J.T.Graves). The first publication about octonions appears as an appendix to an otherwise erroneous paper of the English mathematician (at the time, lawyer) Arthur Cayley (1821-1895) in 1845 (see Introduction and references 17, 18 of [B])

The American algebraist and author of a three-volume *History of the Theory of Numbers*, Leonard E. Dickson (1874-1954) contributed to the construction of composition algebras in 1919 [Di]. The theorem that the only normed division algebras are \mathbb{R} , \mathbb{C} , \mathbb{H} and \mathbb{O} was proven by A. Hurwitz (1859-1919) in 1898. The extension of this result to alternative (including split) algebras belongs to M. Zorn (1906-1993) in 1930 and 1933. The fact that the only division algebras (without extra structure) have dimensions 1, 2, 4, 8 was established as late as in 1958 (independently by R. Bott and J. Milnor and by M. Kervaire).

2 Octonions. Isometries and automorphisms

2.1 Eight dimensional alternative algebras

The multiplication table of the imaginary octonions (see Appendix A) can be introduced by first selecting a quaternion subalgebra

$$\begin{aligned} e_j e_k &= \epsilon_{jkl} e_l - \delta_{jk}, \quad j, k, l = 1, 2, 4, \\ \epsilon_{124} = 1 &= \epsilon_{412} = \epsilon_{241} = 1 = -\epsilon_{214} = \dots \end{aligned} \quad (2.1)$$

The somewhat exotic labeling of the units (jumping over 3) is justified by the following memorable multiplication rules *mod*7:

$$\begin{aligned} e_i e_j = e_k &\Rightarrow e_{i+1} e_{j+1} = e_{k+1}, \quad e_{2i} e_{2j} = e_{2k} \equiv e_{2k(\text{mod}7)} \\ e_7 e_j &= e_{3j(\text{mod}7)}, \quad \text{for } j = 1, 2, 4 \text{ (} e_7 e_4 = e_5 \text{)}. \end{aligned} \quad (2.2)$$

A convenient complex isotropic basis for the vector representation of the isometry Lie algebra $so(8)$ of \mathbb{O} (which contains the automorphism algebra \mathfrak{g}_2 of the octonions) is given by:

$$\rho^\epsilon = \frac{1}{2}(1 + i\epsilon e_7), \quad \zeta_j^\epsilon = \rho^\epsilon e_j = \frac{1}{2}(e_j + i\epsilon e_{3j}) \quad j = 1, 2, 4, \quad \epsilon = \pm \quad (2.3)$$

(the imaginary unit i commutes with the octonion units e_a). The multiplication table of the octonion units is summarized by the following relations:

$$\begin{aligned} (\zeta_j^\epsilon)^2 &= 0 = \rho^+ \rho^-, \quad (\rho^\epsilon)^2 = \rho^\epsilon, \quad \rho^+ + \rho^- = 1, \quad \zeta_j^\epsilon \zeta_k^\epsilon = \epsilon_{jkl} \zeta_l^{-\epsilon} \\ \zeta_j^\epsilon \zeta_k^{-\epsilon} &= -\rho^{-\epsilon} \delta_{jk}, \quad \Rightarrow [\zeta_j^+, \zeta_k^-]_+ = \delta_{jk}, \quad j, k, l = 1, 2, 4. \end{aligned} \quad (2.4)$$

The idempotents ρ^\pm (which go back to Gürsey) are also exploited in [D10]. The last equation (2.4) coincides with the canonical anticommutation relations for fermionic creation and annihilation operators (cf. [CGKK]).

The *split octonions* $x_s \in \mathbb{O}_s$ with units \tilde{e}_a can be embedded in the algebra $\mathbb{C}\mathbb{O}$ of complexified octonions by setting $\tilde{e}_\mu = e_\mu$, $\mu = 0, 1, 2, 4$; $\tilde{e}_7 = ie_7$, $\tilde{e}_{3j} = ie_{3j(\text{mod}7)}$, so that

$$\begin{aligned} x_s &= \sum_{a=0}^7 x_s^a \tilde{e}_a \Rightarrow N(x_s) = x_s x_s^* \\ &= \sum_{\mu=0,1,2,4} (x_s^\mu)^2 - (x_s^7)^2 - (x_s^3)^2 - (x_s^6)^2 - (x_s^5)^2. \end{aligned} \quad (2.5)$$

The quark-lepton correspondence suggests the splitting of octonions into a direct sum,

$$\begin{aligned} \mathbb{O} &= \mathbb{C} \oplus \mathbb{C}^3; \quad x = a + \mathbf{z} \mathbf{e} = a + z^1 e_1 + z^2 e_2 + z^4 e_4 \quad (e_1 e_2 = e_4) \\ a &= x^0 + x^7 e_7, \quad z^j = x^j + x^{3j(\text{mod}7)} e_7 \quad (x^{12} \equiv x^5) \end{aligned} \quad (2.6)$$

thus e_7 playing the role of *imaginary unit within the real octonions*. The Cayley-Dickson construction corresponds to the splitting of \mathbb{O} into two quaternions:

$$\mathbb{O} = \mathbb{H} \oplus \mathbb{H} : x = u + e_7 v, \quad u = x^0 + x^j e_j, \quad v = x^7 + x^{3j} e_j. \quad (2.7)$$

One may speculate that upon complexification u and v could be identified with the "up-" and "down-", leptons and quarks: $u = (\nu; u^j, j = 1, 2, 4)$. $v = (e^-, d_j)$ j playing the role of a colour index, but we shall not pursue this line of thought. For $x_r = u_r + e_7 v_r, r = 1, 2$ the Cayley-Dickson formula (1.16) and its expression in terms of the complex variable a_r, z_r^j reads:

$$\begin{aligned} x_1 x_2 &= u_1 u_2 - v_2 v_1^* + e_7 (u_1^* v_2 + u_2 v_1) = \\ &= a_1 a_2 - \mathbf{z}_1 \bar{\mathbf{z}}_2 + (a_1 \mathbf{z}_2 + \bar{a}_2 \mathbf{z}_1 + \bar{\mathbf{z}}_1 \times \bar{\mathbf{z}}_2) \mathbf{e} \end{aligned} \quad (2.8)$$

where the star indicates quaternionic conjugation while the bar stands for a change of the sign of e_7 ($\bar{z}^j = x^j - e_7 x^{3j}$). The two representations (2.8) display the covariance of the product under the action of two subgroups of maximal rank of the automorphism group of the octonions. If p and q are two unit quaternions

$$\begin{aligned} p &= p^0 + p^j e_j, \quad q = q^0 + q^j e_j \\ pp^* &= N(p) = (p^0)^2 + \mathbf{p}^2 = 1 = qq^* \Leftrightarrow (p, q) \in SU(2) \times SU(2), \end{aligned} \quad (2.9)$$

it is easy to verify, using the first equation (2.8), that the transformation

$$\begin{aligned} (p, q) : x &= u + e_7 v, \rightarrow pup^* + e_7 pvq^*, \\ (p, q) &\in \frac{SU(2) \times SU(2)}{\mathbb{Z}_2^{diag}} \end{aligned} \quad (2.10)$$

where $\mathbb{Z}_2^{diag} = \{(p, q) = (1, 1), (-1, -1)\}$, is an automorphism of the octonion algebra. Similarly, if $U \in SU(3)$ acts on x (2.6) as

$$U : x = a + z^j e_j \rightarrow a + U_j^i z^j e_i \text{ then } U(x_1)U(x_2) = U(x_1 x_2). \quad (2.11)$$

The subgroups (2.10), (2.11) are the two closed connected subgroups of maximal rank of the compact group G_2 corresponding to the *Borel - de Siebenthal theory* [BdS] that plays a central role in [TD] as well as in Chapter 4 below.

2.2 Isometry group of the (split) octonions. Triality

The norm $N(x)$ (1.17) and the associated scalar product of the (split) octonions,

$$N_{(s)}(x) = \sum_{a=0}^7 \eta_a^{(s)} x_a^2, \quad \eta_a = 1 \quad \text{for all } a, \quad \text{for } x \in \mathbb{O}$$

$$\eta_0^s = \eta_1^s = \eta_2^s = \eta_4^s = 1 = -\eta_7^s = -\eta_3^s = -\eta_6^s = -\eta_5^s, \quad \text{for } x \in \mathbb{O}_s \quad (2.12)$$

is the (compact) orthogonal group $O(8)$ (respectively, the split orthogonal group $O(4, 4)$). As stressed in [B], while invariant quadratic forms are common, trilinear forms are rare. It is, therefore, noteworthy, that the trilinear form $t(xyz)$ (1.10) is invariant under the 2-fold cover $\text{Spin}(8)$ (respectively $\text{Spin}(4, 4)$) of the connected orthogonal group $\text{SO}(8)$ (respectively, $\text{SO}(4, 4)$). The existence of such trilinear invariant is related to the exceptional symmetry of the Lie algebra $\mathfrak{D}_4 = \mathfrak{so}(8)$ which is visualized by the symmetry of its Dynkin diagram, Figure 1.




Figure 1: The Dynkin diagram of the Lie algebra \mathfrak{D}_4 . The three outer nodes correspond to the two 8-dimensional chiral spinor representations S_8^\pm of $\text{Spin}(8)$ and to the vector representation of $\text{SO}(8)$, the central node corresponds to the 28-dimensional adjoint representation. The *triality* automorphisms implementing the symmetries of the diagram were discovered in 1925 by Élie Cartan [Ca].

Accordingly, the symmetric group S_3 which permutes the three external nodes of \mathfrak{D}_4 is the group of outer automorphisms of the Lie algebra $\mathfrak{so}(8)$. This is the *triality group* of D_4 . We proceed to describing the Lie algebra \mathfrak{D}_4 and to displaying the action of the outer automorphisms on it. To begin with \mathfrak{D}_4 is generated by left multiplications L_α (as well as by right multiplications R_α) by pure imaginary elements $\alpha \in \text{Im}\mathbb{O}$ acting on the algebra of octonions. If we set (following [M89], [Y])

$$\begin{aligned} L_\alpha x &= \alpha x, & R_\alpha x &= x\alpha, & T_\alpha &= L_\alpha + R_\alpha \\ \text{(i.e. } T_\alpha x &= \alpha x + x\alpha) & \text{for } \text{Re}\alpha &= \langle \alpha, 1 \rangle = 0 \end{aligned} \quad (2.13)$$

then each of the three actions annihilates the inner product:

$$\langle L_\alpha x, y \rangle + \langle x, L_\alpha y \rangle = 0 \quad \text{etc. since } \alpha^* = -\alpha.$$

Note that the product of operators $L_a L_b$, $a, b \in \mathbb{O}$ cannot be written in general as L_{ab} (or as left multiplication by any element of \mathbb{O}). But, as emphasized in the thesis [F16] one can work with composition of maps $L_a L_b$ which is associative while the product of octonions is not. In fact, the Lie algebra $\mathfrak{so}(8)$ is spanned by L_{e_i} and $[L_{e_j}, L_{e_k}]$ for $1 \leq i, j, k \leq 7$. The action of the group

S_3 of outer automorphism of \mathfrak{D}_4 is generated by an automorphism of order three ν and an involution π defined on the triple $(L_\alpha, R_\alpha, T_\alpha)$ by

$$\begin{aligned} \nu L_\alpha &= R_\alpha, & \nu R_\alpha (= \nu^2 L_\alpha) &= -T_\alpha & \Rightarrow & \nu T_\alpha = -L_\alpha, & \nu^3 &= id \\ \pi L_\alpha &= T_\alpha, & \pi R_\alpha &= -R_\alpha & \Rightarrow & \pi T_\alpha = L_\alpha, & \pi^2 &= id. \end{aligned} \quad (2.14)$$

We leave it as an exercise to the reader to verify that the product $\kappa = \pi\nu$ is an involution ($\kappa^2 = id$) satisfying

$$\kappa L_\alpha = -R_\alpha, \quad \kappa R_\alpha = -L_\alpha, \quad \kappa T_\alpha = -T_\alpha, \quad \nu\kappa = \pi, \quad \nu\pi = \pi\nu^2. \quad (2.15)$$

The involutive automorphism κ can be extended to an arbitrary element D of the Lie algebra \mathfrak{D}_4 with the formula ([Y] Sect.1.3).

$$(\kappa D)x = (Dx^*)^*, \quad \text{for all } D \in \mathfrak{D}_4 = \mathfrak{so}(8), \quad x \in \mathbb{O}. \quad (2.16)$$

We shall introduce, following⁴ [M89, Y], two bases, G_{ab} and F_{ab} , in \mathfrak{D}_4 , defined by

$$\begin{aligned} G_{ab}e_c &= \delta_{bc}e_a - \delta_{ac}e_b, & a, b, c &= 0, 1, \dots, 7, & e_0 &= 1 \\ F_{ab}x &= \frac{1}{2}e_a(e_b^*x) = -F_{ba}x & (F_{aa} &:= 0, \quad e_0^* = e_0) \end{aligned} \quad (2.17)$$

which satisfy the same commutation relations:

$$[X_{ab}, X_{cd}] = \delta_{bc}X_{ad} - \delta_{bd}X_{ac} + \delta_{ad}X_{bc} - \delta_{ac}X_{bd}, \quad X = G, F. \quad (2.18)$$

Using the identity $L_{e_k} = 2F_{k0}$ (for $k = 1, \dots, 7$) and Eq. (2.14) one verifies that $\pi(F_{k0}) = G_{k0}$. More generally, we have

$$\pi(G_{ab}) = F_{ab}, \quad \pi(F_{ab}) = G_{ab} \quad (2.19)$$

(cf. Lemma 1.3.3 of [Y]). We shall demonstrate in Appendix B that the involution π splits into seven 4-dimensional involutive transformations. Here, we display one of them which involves our choice of the Cartan subalgebra

⁴We note that the convention $e_1e_2 = e_3$ (rather than our $e_1e_2 = e_4$ which follows [B, CS]) is used in [M89, Y] so that our formulas relating G and F (Appendix B) differ from theirs.

of $so(8)$:

$$\begin{aligned}
F_7 &= X_7 G_7, \quad G_7 = \begin{pmatrix} G_{07} \\ G_{13} \\ G_{26} \\ G_{45} \end{pmatrix}, \quad F_7 = \begin{pmatrix} F_{07} \\ F_{13} \\ F_{26} \\ F_{45} \end{pmatrix}, \\
X_7 &= \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}, \tag{2.20}
\end{aligned}$$

a straightforward calculation gives $X_7^2 = \mathbb{I}$, $\det X_7 = -1$.

Theorem 3 (*Infinitesimal triality*) *For any $D \in \mathfrak{D}_4$ and $x, y \in \mathbb{O}$ the following identity holds:*

$$(Dx)y + x\nu(D)y = \pi(D)(xy). \tag{2.21}$$

For a given $D \neq 0$ the automorphisms ν and π , given by (2.14), (2.19) are uniquely determined from (2.21).

For $D = L_\alpha$ ($\operatorname{Re}\alpha = 0$) Theorem 3 follows from the definition (2.14). For a general $D \in \mathfrak{D}_4$ one uses the fact that it can be written as a linear combination of L_α and their commutators (see Theorem 1.3.6 of [Y]).

There exists a group theoretic counterpart of the principle of infinitesimal triality which we shall only sketch. (For a pedagogical exposition based on the concepts of Moufang loops and their isotopies - see Chapters 7 and 8 of [CS].)

The notion of left and right multiplication are well defined and preserve the norm for unit octonions; for instance $N(L_a x) = N(x)$ for $aa^* = 1$. In order to obtain a general $SO(8)$ action on the octonions one needs seven left (or seven right) multiplications ([CS] Sect. 8.4). The group theoretic counterpart of T_α is the bimultiplicaton $B_a x = axa$; for $a_t = \exp(t\alpha)$, $t \in \mathbb{R}$, $\alpha \in \operatorname{Im}\mathbb{O}$

$$\lim_{t \rightarrow 0} \frac{d}{dt} B_{a_t} x = T_\alpha x = \alpha x + x\alpha. \tag{2.22}$$

The automorphisms (2.14), (2.15) can be lifted to group automorphisms:

$$\begin{aligned}
\nu(L_a, R_a, B_a) &= (R_a, B_{a^*}, L_{a^*}), & \pi(L_a, R_a, B_a) &= (B_a, R_{a^*}, L_a) \\
\kappa(L_a, R_a, B_a) &= (R_{a^*}, B_{a^*}, L_{a^*}), & \kappa^2 &= \pi^2 = \nu^3 = 1.
\end{aligned} \tag{2.23}$$

Theorem 4 For each rotation $g \in \text{SO}(8)$ there are elements g^\pm of the double cover $\text{Spin}(8)$ of $\text{SO}(8)$ such that the normed trilinear form

$$t_8(x, y, z) = \frac{1}{2}t(x, y, z) = \langle xyz, 1 \rangle, \quad x, y, z \in \mathbb{O} \quad (2.24)$$

satisfy the invariance condition

$$t_8(gx, g^+y, g^-z) = t_8(x, y, z). \quad (2.25)$$

The elements g^\pm are determined from (2.25) up to a common sign: for each g there are exactly two pairs (g^+, g^-) and $(-g^+, -g^-)$ obeying (2.25). For $g = L_a$ ($aa^* = 1$) we can set ,

$$g^+ = \nu(L_a) = R_a, \quad g^- = \nu^2 L_a = B_{a^*}. \quad (2.26)$$

(The factor $\frac{1}{2}$ in (2.24) follows the definition of *normed triality* of [B] which demands $|t_8(x, y, z)|^2 \leq N(x)N(y)N(z)$.)

2.3 Automorphism group G_2 of the octonions.

As proven by Élie Cartan in 1914, the automorphism group of the octonions is the rank 2 exceptional Lie group G_2 which can thus be defined by

$$G_2 = \{g \in L(\mathbb{O}, \mathbb{R}) \mid (gx)(gy) = g(xy), \quad x, y \in \mathbb{O}\} \quad (2.27)$$

where $L(\mathbb{O}, \mathbb{R})$ is the group of non-singular linear transformations of the 8-dimensional real vector space \mathbb{O} . It follows from $e_0^2 = e_0 = 1$ and $e_k^* = -e_k$ for $k > 0$ that G_2 preserves the octonion unit and the Cayley conjugation and hence the norm $N(x)$:

$$g1 = 1, \quad (gx)^* = g(x^*), \quad N(gx) = N(x). \quad (2.28)$$

Thus G_2 is a subgroup of the isometry (orthogonal) group $O(\mathbb{O}) = O(8)$ of the 8-dimensional euclidean space of the octonions. In fact, it is a subgroup of the connected orthogonal group $\text{SO}(7)$ of the 7-dimensional space $\text{Im}\mathbb{O}$ of imaginary octonions, the Lie algebra $\mathfrak{so}(7)$ splitting as a vector space into

a direct sum of the Lie algebra \mathfrak{g}_2 and the vector representation $\underline{7}$ of $\mathfrak{so}(7)$. Thus the dimension of G_2 is $\binom{7}{2} - 7 = 14$.

$$\mathfrak{so}(7) \cong \mathfrak{g}_2 \oplus \underline{7} (= \mathfrak{g}_2 \oplus \mathbb{R}^7).$$

Moreover, the group G_2 acts transitively on the unit sphere \mathbb{S}^6 in \mathbb{R}^7 ; every point of \mathbb{S}^6 can be transformed, say, into e_7 by an automorphism of \mathbb{O} . The stabilizer of e_7 is the subgroup $SU(3)$ of G_2 , defined by (2.11):

$$(G_2)_{e_7} = SU(3) \Rightarrow \frac{G_2}{SU(3)} \cong \mathbb{S}^6. \quad (2.29)$$

It follows that the group G_2 is connected. The maximal subgroups of G_2 , whose action was defined by (2.10) and (2.11) (and which correspond to the Borel - de Siebenthal theory) can be characterized as follows. The *complex conjugation* γ (in the notation of [Y]): $e_7 \rightarrow -e_7$ belongs to the automorphism group G_2 of \mathbb{O} (corresponding, in fact, to the reflection of four imaginary units e_7 , $e_7e_1 = e_3$, $e_7e_2 = e_6$ and $e_7e_4 = e_5$) and has square one:

$$\gamma x = \gamma(u + e_7v) = u - e_7v, \quad \gamma^2 = 1. \quad (2.30)$$

The rank two (semisimple) subgroup (2.9), (2.10) of G_2 can be characterized as the commutant of γ in G_2 :

$$G_2^\gamma = \{g \in G_2 | \gamma g = g \gamma\} = \frac{SU(2) \times SU(2)}{\mathbb{Z}_2}. \quad (2.31)$$

Denote, on the other hand, by ω the generator of the center of $SU(3)$ acting on z by (2.11)

$$\omega x = a + \omega_7 z^j e_j, \quad \omega_7 = -\frac{1}{2} + \frac{\sqrt{3}}{2} e_7 \quad (\omega_7^3 = 1 = \omega^3). \quad (2.32)$$

Then the subgroup (2.11) of G_2 is characterized by

$$G_2^\omega := \{\omega \in G_2 | \omega g = g \omega\} = SU(3) (= (G_2)_{e_7}). \quad (2.33)$$

(One uses, in particular, the relation $\omega_7 \mathbf{z} e \omega_7 = \omega_7 \bar{\omega}_7 \mathbf{z} e = \mathbf{z} e$.)

2.4 Roots and weights of $\mathfrak{g}_2 \subset \mathfrak{B}_3 \subset \mathfrak{D}_4$

It is convenient (in particular, for the study of the Jordan algebra $JSpin_9$ in Sect. 3.3 below to view the Lie algebra \mathfrak{D}_4 as embedded into the Clifford algebra Cl_8). Indeed, its 16×16 matrix-generators \hat{P} , $a = 0, 1, \dots, 7$, satisfy the same anticommutation relations as the 2×2 hermitian octonionic matrices

$$\hat{e}_a = \begin{pmatrix} 0 & e_a \\ e_a^* & 0 \end{pmatrix}, \text{ i.e. } \hat{e}_0 = \begin{pmatrix} 0 & e_0 \\ e_0 & 0 \end{pmatrix}, (e_0 = 1), \hat{e}_k = \begin{pmatrix} 0 & e_k \\ -e_k & 0 \end{pmatrix} \quad (2.34)$$

$k = 1, \dots, 7$; we have

$$[\hat{e}_a, \hat{e}_b]_+ = 2\delta_{ab}\mathbb{1}_2 \quad \leftrightarrow \quad \left[\hat{P}_a, \hat{P}_b \right]_+ = 2\delta_{ab}\mathbb{1}_{16}, \quad a, b = 0, 1, \dots, 7. \quad (2.35)$$

(The symbol 1_n , the $n \times n$ unit matrix, will henceforth be omitted wherever this would not give rise to ambiguity.) The Clifford algebra gives also room for the symmetry generators. If we set

$$\hat{G}_{ab} = \begin{pmatrix} G_{ab} & 0 \\ 0 & G_{ab}^\kappa \end{pmatrix}, \quad \text{where} \quad G_{ab}^\kappa = \kappa(G_{ab}), \quad (2.36)$$

so that, in view of (2.16) and (2.17), $G_{ab}^\kappa \hat{e}_c^* = \delta_{bc}e_a^* - \delta_{ac}e_b^*$ we shall have

$$\hat{G}_{ab}\hat{e}_c = \delta_{bc}\hat{e}_a - \delta_{ac}\hat{e}_b, \quad a, b, c = 0, 1, \dots, 7. \quad (2.37)$$

Writing the \hat{P} , in analogy with (2.34), as

$$\begin{aligned} \hat{P}_0 &= \sigma_1 \otimes P_0 = \begin{pmatrix} 0 & P_0 \\ P_0 & 0 \end{pmatrix}, \quad \hat{P} = c \otimes P_k = \begin{pmatrix} 0 & P_k \\ -P_k & 0 \end{pmatrix}, \quad k = 1, \dots, 7, \\ P_0 &= \mathbb{1}_8, \quad c = i\sigma_2, \quad [P_j, P_k]_+ = -2\delta_{jk}, \quad P_1 P_2 \dots P_6 = P_7, \end{aligned} \quad (2.38)$$

we see that the generators G_{ab} of the Lie algebra \mathfrak{D}_4 are represented by

$$\frac{1}{2}\hat{P}_{ab} = \frac{1}{4}[\hat{P}_a, \hat{P}_b], \quad \left[\frac{1}{2}\hat{P}_{ab}, \hat{P}_c \right] = \delta_{bc}\hat{P}_a - \delta_{ac}\hat{P}_b, \quad a, b, c = 0, 1, \dots, 7. \quad (2.39)$$

We stress that the map $\hat{e}_a \rightarrow \hat{P}_a$ (and hence the map $\hat{e}_a \rightarrow \Gamma_a$) - unlike the representation $q_j \rightarrow -i\sigma_j$ of the imaginary quaternion units - only respects the anticommutators, the relation $L_{e_1} \dots L_{e_6} 1 = e_7$, and the $SO(\mathbb{O})$ symmetry properties of the octonions, not their commutators.

The four operators G_7 of (2.20) are diagonalized in the counterpart of the isotropic octonionic basis (2.3), (2.4) given by

$$\begin{aligned}\hat{P}_0^\epsilon &= \frac{1}{2}(\hat{P}_0 + i\epsilon\hat{P}_7), & \hat{P}_j^\epsilon &= \frac{1}{2}(\hat{P}_j + i\epsilon\hat{P}_{3j}), \quad j = 1, 2, 4, \quad (3 \times 4 = 5(\text{mod}7)) \\ \epsilon &= \pm, \text{ s.t. } (\hat{P}_\mu^\epsilon)^2 = 0, & \hat{P}_\mu^{-\epsilon}\hat{P}_\mu^\epsilon &=: \rho_\mu^\epsilon, \quad (\rho_\mu^\epsilon)^2 = \rho_\mu^\epsilon, \quad \rho_\mu^+ + \rho_\mu^- = \mathbb{I}_{16}.\end{aligned}\quad (2.40)$$

Their commutators define an orthogonal weight basis in \mathfrak{D}_4 (as well as in $\mathfrak{B}_4 = \mathfrak{so}(9)$) - see Sect. 4.1):

$$\Lambda_\mu = \frac{1}{2} [\hat{P}_\mu^-, \hat{P}_\mu^+] = \frac{1}{2}(\rho_\mu^+ - \rho_\mu^-), \quad \Lambda_0 = \frac{i}{2}\hat{P}_{07}, \quad \Lambda_j = \frac{i}{2}\hat{P}_{3j}, \quad j = 1, 2, 4, \quad (2.41)$$

which can be represented by real diagonal matrices (belonging actually to $\mathfrak{so}(4, 4) \subset Cl(4, 4)$). The operator Λ_μ are normalized to have eigenvalues ± 1 (and 0) under commutation:

$$[\Lambda_\mu, \hat{P}_\nu^\epsilon] = -\epsilon\delta_{\mu\nu}\hat{P}_\nu^\epsilon, \quad \mu, \nu = 0, 1, 2, 4, \quad \epsilon = \pm. \quad (2.42)$$

They also satisfy the orthogonality relation $\langle \Lambda_\mu, \Lambda_\nu \rangle := \frac{1}{4}tr(\Lambda_\mu, \Lambda_\nu) = \delta_{\mu\nu}$. A Chevalley-Cartan basis corresponding to the simple roots α_μ of \mathfrak{D}_4 is given by

$$\begin{aligned}\alpha_0 &\leftrightarrow H_0 = \Lambda_0 - \Lambda_1 = \rho_0^+ - \rho_1^+, & E_0 &= \hat{P}_0^- \hat{P}_1^+, & F_0 &= \hat{P}_1^- \hat{P}_0^+ \\ \alpha_1 &\leftrightarrow H_1 = \Lambda_1 - \Lambda_2 = \rho_1^+ - \rho_2^+, & E_1 &= \hat{P}_1^- \hat{P}_2^+, & F_1 &= \hat{P}_2^- \hat{P}_1^+ \\ \alpha_2 &\leftrightarrow H_2 = \Lambda_2 - \Lambda_4 = \rho_2^+ - \rho_4^+, & E_2 &= \hat{P}_2^- \hat{P}_4^+, & F_2 &= \hat{P}_4^- \hat{P}_2^+ \\ (H_4 =) H_{\alpha_4} &= \Lambda_2 + \Lambda_4 = \pm\rho_2^\pm \mp \rho_4^\mp, & E_{\alpha_4} &= \hat{P}_4^- \hat{P}_2^-, & F_{\alpha_4} &= \hat{P}_2^+ \hat{P}_4^+\end{aligned}\quad (2.43)$$

and satisfy the standard commutation relations

$$[H_\mu, E_\nu] = c_{\mu\nu}^D E_\nu, \quad [H_\mu, F_\nu] = -c_{\mu\nu}^D F_\nu, \quad [E_\mu, F_\nu] = \delta_{\mu\nu} H_\nu \quad (2.44)$$

where $c_{\mu\nu}^D$ is the \mathfrak{D}_4 Cartan matrix

$$(c_{\mu\nu}^D) = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & -1 \\ 0 & -1 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}.$$

As it follows from our discussion in Sect. 2.3 of the automorphism group of the octonions the derivations of \mathbb{O} span, in fact, a subalgebra of the 21-dimensional Lie algebra

$$\begin{aligned}\mathfrak{B}_3 &= \mathfrak{so}_7 = \{\mathfrak{D} \in \mathfrak{D}_4 : De_0 = 0\} = \{D \in \mathfrak{D}_4; \kappa(D) = D\} \\ &= \text{Span}\{G_{kl}; k, l = 1, \dots, 7\}.\end{aligned}\tag{2.45}$$

The first two simple roots α_1 and α_2 of \mathfrak{B}_3 coincide with those of \mathfrak{D}_4 , while the third, α_s , is a short root:

$$\{\text{Simple roots of } \mathfrak{B}_3\} = \{\alpha_1 = \Lambda_1 - \Lambda_2, \alpha_2 = \Lambda_2 - \Lambda_4, \alpha_s = \Lambda_4\}\tag{2.46}$$

the coroot corresponding to α_s being

$$\alpha_s^\vee = \frac{2\alpha_s}{(\alpha_s, \alpha_s)} = 2\Lambda_4$$

(so that $C_{2s}^B = (\alpha_2, \alpha_s^\vee) = -2$).

Comparing the defining relation for a derivation $D \in \mathfrak{g}_2$,

$$(Dx)y + xDy = D(xy)\tag{2.47}$$

with the infinitesimal triality relation (2.21) we conclude that $D \in \mathfrak{g}_2$ only if $\nu(D) = \pi(D) = D$. Taking into account the fact that $\kappa(D) = D$ in \mathfrak{B} and that $\nu = \pi\kappa$ (according to (2.15)) we arrive at the following

Proposition 2.3 *If $D \in \mathfrak{B}_3$ then each of the condition $\nu(D) = D$ and $\pi(D) = D$ implies the other and the resulting triality invariance is necessary and sufficient for D to belong to \mathfrak{g}_2 .*

Combining this result with the last equation (2.45) we deduce that a linear combination of G_{kl} belongs to \mathfrak{g}_2 if it is invariant under the involution π (2.19). In particular, taking (2.20) into account, we deduce that $\lambda G_{13} + \mu G_{26} + \nu G_{45} \in \mathfrak{g}_2$ iff $\lambda + \mu + \nu = 0$. In other words, the Cartan subalgebra \mathfrak{g}_2 is spanned by $G_{12} - G_{26}$ and $G_{26} - G_{45}$ whose representatives within Γ_{ab} are proportional to H_1 and H_2 of eq. (2.43). More generally, using Appendix B, we find that the following seven linear combinations of G_{lk} span \mathfrak{g}_2

$$\begin{aligned}\lambda G_{24} + \mu G_{37} + \nu G_{56}, & \quad \lambda G_{14} - \mu G_{35} + \nu G_{76}, \\ \lambda G_{17} + \mu G_{25} - \nu G_{46}, & \quad -\lambda G_{12} + \mu G_{36} + \nu G_{75}, \\ \lambda G_{16} - \mu G_{23} + \nu G_{47}, & \quad -\lambda G_{15} + \mu G_{27} + \nu G_{43}, \\ \lambda G_{13} + \mu G_{26} + \nu G_{45}, & \quad \text{with } \lambda + \mu + \nu = 0.\end{aligned}\tag{2.48}$$

3 Jordan algebras and related groups

3.1 Classification of finite dimensional Jordan algebras

Pascual Jordan (1902-1980) the "unsung hero among the creators of quantum theory" (in the words of Schweber, 1994) asked himself in 1932 a question you would expect of an idle mathematician: Can one construct an algebra of (hermitian) observables without introducing an auxiliary associative product? He arrived, after some experimenting with the *special Jordan product*

$$A \circ B = \frac{1}{2}(AB + BA) (= B \circ A), \quad (3.1)$$

at two axioms (Jordan, 1933)

$$(i) : A \circ B = B \circ A; \quad (ii) : A^2 \circ (B \circ A) = (A^2 \circ B) \circ A, \quad (3.2)$$

where $A^2 := (A \circ A)$. They imply, in particular, power associativity and

$$A^m \circ A^n = A^{m+n}, \quad m, n = 0, 1, 2, \dots \quad (3.3)$$

(Jordan algebras are assumed to contain a unit and $A^0 = 1$.) Being interested in extracting the properties of the algebra of hermitian matrices (or selfadjoint operators) for which $A^2 \geq 0$, Jordan adopted Artin's *formal reality condition*

$$A_1^2 + \dots + A_n^2 = 0 \implies A_1 = 0 = \dots = A_n. \quad (3.4)$$

(It is enough to demand (3.4) for $n=2$.) Algebras over the real numbers satisfying both (3.2) and (3.4) are now called *euclidean Jordan algebras*. In a fundamental paper of 1934 Jordan, von Neumann and Wigner [JvNW] classified all finite dimensional euclidean Jordan algebras. They split into a direct sum of *simple algebras*, which belong to four infinite families,

$$\mathcal{H}_n(\mathbb{R}), \quad \mathcal{H}_n(\mathbb{C}), \quad \mathcal{H}_n(\mathbb{H}), \quad JSpin_n, \quad (3.5)$$

and a single exceptional one

$$\mathcal{J}(= J_3^8) = \mathcal{H}_3(\mathbb{O}). \quad (3.6)$$

Here $\mathcal{H}_n(\mathbb{A})$ stands for the set of $n \times n$ hermitian matrices with entries in the division ring $\mathbb{A}(= \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O})$, equipped with the commutative product (3.1).

(One uses the same notation when \mathbb{A} is replaced by one of the alternative split composition rings, $\mathbb{C}_s, \mathbb{H}_s$ or \mathbb{O}_s albeit the resulting algebra is not euclidean in that case.) $JSpin_n$ is an algebra of elements $(\xi, x; \xi \in \mathbb{R}, x \in \mathbb{R}^n)$ where \mathbb{R}^n is equipped with the (real) euclidean scalar product $\langle x, y \rangle$ and the product in $JSpin_n$ is given by

$$(\xi, x)(\eta, y) = (\xi\eta + \langle x, y \rangle, \xi y + \eta x). \quad (3.7)$$

The first three algebras $\mathcal{H}_n(\mathbb{A})$ (3.5) are *special*: the matrix product AB in (3.1) is *associative*. The algebra $JSpin_n$ is also special as a Jordan subalgebra of the 2^n dimensional (associative) Clifford algebra Cl_n .

Remark 3.1. The Jordan algebras $\mathcal{H}_2(\mathbb{A})$ for $\mathbb{A} = \mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$ are isomorphic to $JSpin_n$ for $n = 2, 3, 5, 9$, respectively. In fact, more generally, let $x = x^0 + \mathbf{x}$ where $\mathbf{x} = \sum_{k=1}^{n-2} x^k e_k \in Cl_{2-n}$; then the two-by-two (Clifford valued) matrix:

$$X = \begin{pmatrix} \xi + x_{n-1} & x \\ x^* & \xi - x_{n-1} \end{pmatrix} \quad (3.8)$$

satisfies Eq. (1.5) with $N(X) = \det X$, $t(X) = \text{tr}(X) = 2\xi$ where the determinant of X has a (time-like) Minkowski space signature:

$$\det X = \xi^2 - x_{n-1}^2 - xx^*, \quad xx^* = x^*x = (x^0)^2 + (x^1)^2 + \dots + (x^{n-2})^2 \quad (3.9)$$

and is thus invariant under the Lorentz group $SO(n, 1)$. We leave it to the reader to verify that the multiplication law (3.7) can be obtained from Eq. (1.5) for the matrix (3.8) by polarization. (Cf. Sect. 1.2.)

On the other hand, the algebras $\mathcal{H}_n(\mathbb{O})$ for $n > 3$ are not Jordan since they violate condition (ii) of (3.2). The exceptional Jordan algebra $\mathcal{J} = \mathcal{H}_3(\mathbb{O})$ did not seem to be special but the authors of [JvNW] assigned the proof that the product $A \circ B$ of two elements of \mathcal{J} cannot be represented in the form (3.1) with an *associative* product AB to A. Adrian Albert (1905-1972) - a PhD student of L. Dickson. As a result, many authors, including [McC] call \mathcal{J} an *Albert algebra*.

With the realization (3.8) of the elements of $JSpin_n$ we see that each (simple) euclidean Jordan algebra is a matrix algebra of some kind and so has a well defined (matrix) trace. The trace ρ of the unit element of a Jordan algebra defines its *rank*; in particular, the rank of the algebra $JSpin_n$ for any n is $\rho(JSpin_n) = 2$. The real dimension δ of the non-diagonal elements of the

matrix representation of a Jordan algebra is called its *degree*. (For a concise survey of euclidean Jordan algebras and their two numerical characteristics, the rank and the degree - see [M] (Sect. 2); note that Meng denotes the $JSpin_n$ by $\Gamma(n)$ and calls them *Dirac type*.) The algebras listed in (3.5) have degrees $\delta = 1, 2, 4, n - 1$, respectively, while the Albert algebra (3.6) has degree 8. The dimension of an euclidean Jordan algebra $V = J_\rho^\delta$ of rank ρ and degree δ is $\dim J_\rho^\delta = \binom{\rho}{2}\delta + \rho$. The rank and the degree completely characterize the simple euclidean Jordan algebras.

We introduce the 1-dimensional projectors E_i and the hermitian octonionic matrices $F_i(x_i)$ writing down a general element of $\mathcal{H}_3(\mathbb{O})$ as

$$\begin{aligned} X &= \begin{pmatrix} \xi_1 & x_3 & x_2^* \\ x_3^* & \xi_2 & x_1 \\ x_2 & x_1^* & \xi_3 \end{pmatrix} \\ &= \xi_1 E_1 + \xi_2 E_2 + \xi_3 E_3 + F_1(x_1) + F_2(x_2) + F_3(x_3). \end{aligned} \quad (3.10)$$

We can then write the Jordan multiplication $X \circ Y$ setting

$$\begin{aligned} E_i \circ E_j &= \delta_{ij} E_j, \quad E_i \circ F_j(x) = \begin{cases} 0, & \text{if } i = j, \\ \frac{1}{2} F_j(x), & \text{if } i \neq j. \end{cases} \\ F_i(x) \circ F_i(y) &= \langle x, y \rangle (E_{i+1} + E_{i+2}), \\ F_i(x) \circ F_{i+1}(y) &= \frac{1}{2} F_{i+2}(y^* x^*), \end{aligned} \quad (3.11)$$

where the indices are counted mod 3: $E_4 \equiv E_1$, $F_5 \equiv F_2$, \dots . We define the trace, a symmetric bilinear inner product and a trilinear scalar product in J by

$$\begin{aligned} \text{tr} X &= \xi_1 + \xi_2 + \xi_3, \\ \langle X, Y \rangle &= \text{tr}(X \circ Y), \quad \text{tr}(X, Y, Z) = \langle X, Y \circ Z \rangle. \end{aligned} \quad (3.12)$$

The exceptional algebra \mathcal{J} also admits a (symmetric) *Freudenthal product*:

$$X \times Y = \frac{1}{2} [2X \circ Y - X \text{tr} Y - Y \text{tr} X + (\text{tr} X \text{tr} Y - \langle X, Y \rangle) E] \quad (3.13)$$

where E is the 3×3 unit matrix, $E = E_1 + E_2 + E_3$. Finally, we define a 3-linear form (X, Y, Z) and the determinant $\det X$ by

$$\begin{aligned} (X, Y, Z) &= \langle X, Y \times Z \rangle = \langle X \times Y, Z \rangle, \quad \det X = \frac{1}{3} (X, X, X) \\ &= \xi_1 \xi_2 \xi_3 + 2 \text{Re}(x_1 x_2 x_3) - \xi_1 x_1 x_1^* - \xi_2 x_2 x_2^* - \xi_3 x_3 x_3^*. \end{aligned} \quad (3.14)$$

The following identities hold:

$$\begin{aligned} X \times X \circ X &= (\det X)E \quad (\text{Hamilton-Cayley}) \\ (X \times X) \times (X \times X) &= (\det X)X. \end{aligned} \quad (3.15)$$

3.2 The Tits-Kantor-Koecher (TKK) construction

The symmetrized product $u \circ x$ (3.1) is not the only way to construct a hermitean operator out of two such operators u and x . The quadratic in u binary operation $P_u x = u x u$ also gives a hermitean result whenever u and x are hermitean. Its expression in terms of the Jordan product looks somewhat clumsy:

$$P_u = 2L_u^2 - L_{u^2}, \quad \text{—i.e.} \quad P_u x = u \circ (u \circ x) - u^2 \circ x \quad (3.16)$$

but, as emphasized by McCrimmon [McC], it can be advantageously taken as a basic notion defined axiomatically. Introduce first the polarized form of P :

$$S_{uv}w = \frac{1}{2}(P_{u+w} - P_u - P_w)v = \{uvw\} := u \circ (v \circ w) + w \circ (v \circ u) - (u \circ w) \circ v = \{wvu\}. \quad (3.17)$$

A *unital quadratic Jordan algebra* is a space together with a distinguished element 1 and a product $P_u(x)$ linear in x and quadratic in u , which is *unital* and satisfies the *Commuting Formula* and the *Fundamental Formula*:

$$P_1 = 1, \quad P_u S_{vu} = S_{uv} P_u, \quad P_{P_u(v)} = P_u P_v P_u. \quad (3.18)$$

The triple product $\{uvw\}$ is symmetric (according to the last equation (3.17)) and obeys the *5-linear relation*

$$\{xy\{uvw\}\} = \{\{xyu\}vw\} - \{u\{yxv\}w\} + \{uv\{xyw\}\}. \quad (3.19)$$

This identity can be read as a Lie algebra relation:

$$[S_{xy}, S_{uv}] = S_{\{xyu\}v} - S_{u\{yxv\}} \quad (3.20)$$

applied to an arbitrary element $w \in V$. It defines the *structure Lie algebra* \mathfrak{st} of V . The generators S_{uv} can be expressed in terms of the (left) multiplication operators L_x as follows:

$$S_u (= S_{u1}) = L_u, \quad S_{uv} = L_{uv} + [L_u, L_v]. \quad (3.21)$$

The *derivation algebra* $\mathfrak{der}(V)$ of V , spanned by the commutators $[L_u, L_v]$, appears as the maximal compact Lie subalgebra of $\mathfrak{str}(V)$.

The *conformal algebra* $\mathfrak{co}(v)$ is an extension of $\mathfrak{str}(V)$ defined, as a vector space, as

$$\mathfrak{co}(v) = V \dot{+} \mathfrak{str}(V) \dot{+} V^* \quad (3.22)$$

with the natural *TKK commutation relations* (under the action of the structure algebra, u and v in S_{uv} transforming as a vector and a covector, respectively - see Theorem 3.1 of [M]). Here there are three relevant examples (the reader can find a complete list in [M]):

$$\begin{aligned} \mathfrak{der}(JSpin_n) &= \mathfrak{so}(n), & \mathfrak{str}(JSpin_n) &= \mathfrak{so}(n, 1) \oplus \mathbb{R}, & \mathfrak{co}(JSpin_n) &= \mathfrak{so}(n+1, 2) \\ \mathfrak{der}(\mathcal{H}_n(\mathbb{C})) &= \mathfrak{su}(n), & \mathfrak{str}(\mathcal{H}_n(\mathbb{C})) &= \mathfrak{sl}(n, \mathbb{C}) \oplus \mathbb{R}, & \mathfrak{co}(\mathcal{H}_n(\mathbb{C})) &= \mathfrak{su}(n, n) \\ \mathfrak{der}(\mathcal{H}_3(\mathbb{O})) &= \mathfrak{f}_4, & \mathfrak{str}(\mathcal{H}_3(\mathbb{O})) &= \mathfrak{e}_{6(-26)} \oplus \mathbb{R}, & \mathfrak{co}(\mathcal{H}_3(\mathbb{O})) &= e_{7(-25)}. \end{aligned} \quad (3.23)$$

A *Jordan triple system* V with a triple product $V^{\times 3} \rightarrow V, \{uvw\}$, satisfying (i) $\{uvw\} = \{wvu\}$ and (3.19) is a generalization of a Jordan algebra (as every Jordan algebra generates a Jordan triple system). The same structure arises⁵ in any 3-graded Lie algebra $\mathfrak{g} = \mathfrak{g}_{-1} \dot{+} \mathfrak{g}_0 \dot{+} \mathfrak{g}_1$ with an involution τ exchanging $\mathfrak{g}_{\pm 1}$ (see [P]):

$$\{uvw\} = [[u, \tau(v)], w], \quad u, v, w \in \mathfrak{g}_1, \quad \tau(v) \in \mathfrak{g}_{-1}, \quad [u, \tau(v)] \in \mathfrak{g}_0. \quad (3.24)$$

Another important discovery, due to Koecher and his school, is the existence of a one-to-one correspondence between (simple) euclidean Jordan algebras V and (irreducible) symmetric cones $\Omega(V)$ (see [K], [FK]). To the four matrix algebras correspond the cones of positive definite matrices $\Omega_n(\mathbb{R}), \Omega_n(\mathbb{C}), \Omega_n(\mathbb{H}), \Omega_3(\mathbb{O})$ of rank $\rho := tr_V(1)$ equal to n or 3 , respectively. The positive cone of $JSpin_n$ coincides with the forward light cone:

$$\Omega_n(JSpin_n) = \{(\xi, x) \in JSpin_n \mid \xi > \sqrt{x_1^2 + \cdots + x_n^2}\}. \quad (3.25)$$

In all cases $\Omega(V)$ is spanned by (convex) linear combinations of squares of elements $x \in V$ with positive coefficients; equivalently, $\Omega(V)$ is the connected component of the unit element $1 \in V$ of the invertible elements of V . The

⁵This fact has been discovered by Isaiah Kantor (1964) - see the emotional essay [Z] by Efim Zelmanov.

cones $\Omega(V)$ are all selfdual and invariant under the structure group $Str(V)$. The conformal group $Co(V)$ can be defined as the automorphism group of the tube domain:

$$Co(V) = Aut\{V + i\Omega(V)\}. \quad (3.26)$$

3.3 Automorphism groups of the exceptional Jordan algebras $\mathcal{H}_3(\mathbb{O}_{(s)})$ and their maximal subgroups

Classical Lie groups appear as symmetries of classical symmetric spaces. For quite some time there was no such interpretation for the exceptional Lie groups. The situation only changed with the discovery of the exceptional Jordan algebra $\mathcal{H}_3(\mathbb{O})$ and its split octonions' cousin $\mathcal{H}_3(\mathbb{O}_s)$.

The automorphism group of $\mathcal{H}_3(\mathbb{O})$ is the rank four compact simple Lie group ⁶ F_4 . It leaves the unit element E invariant and is proven to preserve the trace (3.12) (see Lemma 2.2.1 in [Y]). The stabilizer of E_1 in F_4 is the double cover $Spin(9)$ of the rotation group in nine dimensions (which preserves X_0^2 (3.8)). Moreover, we have

$$\begin{aligned} F_4/Spin(9) &\simeq \mathbb{O}\mathbb{P}^2 \implies \\ \dim F_4 &= \dim Spin(9) + \dim \mathbb{O}^2 = 36 + 16 = 52. \end{aligned} \quad (3.27)$$

Building on our treatment of $D_4 \supset \mathfrak{g}_2$ of Sect. 2.4 we shall first construct the Cartan subalgebra of the Lie algebra \mathfrak{f}_4 of *derivation* (infinitesimal automorphisms) of $\mathcal{H}_3(\mathbb{O}_s)$. It is again spanned by the orthonormal weight basis Λ_μ , $\mu = 0, 1, 2, 4$ (that actually belongs to the real form $f_{4(4)} \supset \mathfrak{so}(4, 4)$), their restriction to \mathfrak{D}_4 being given by (2.41). The simple roots α and the corresponding coroots α^\vee of \mathfrak{f}_4 are given by:

$$\begin{aligned} \alpha_1 &= \Lambda_1 - \Lambda_2 = \alpha_1^\vee = H_1, & \alpha_2 &= \Lambda_2 - \Lambda_4 = \alpha_2^\vee = H_2 \\ (\alpha_4^{(s)})_{s_4} &= \Lambda_4, \quad s_4^\vee = 2\Lambda_4 = H_4^s, & s_0 &= \frac{1}{2}(\Lambda_0 - \Lambda_1 - \Lambda_2 - \Lambda_4) (= \alpha_0^{(s)}) \\ s_0^\vee &= \Lambda_0 - \Lambda_1 - \Lambda_2 - \Lambda_4 = H_0^s. \end{aligned} \quad (3.28)$$

⁶This was proven by Claude Chevalley and Richard Schafer in 1950. The result was prepared by Ruth Moufang's study in 1933 of the octonionic projective plane, then Jordan's construction in 1949 of $\mathbb{O}\mathbb{P}^2$ in terms of 1-dimensional projections in $\mathcal{H}_3(\mathbb{O})$ and Armand Borel's observation that F_4 is the isometry group of $\mathbb{O}\mathbb{P}^2$; for a review and references - see [B] (Sect. 4.2). Octonionic quantum mechanics in the Moufang plane was considered in [GPR].

The corresponding Cartan matrix reads:

$$(c_{ij}^f = \langle \alpha_i^\vee, \alpha_j \rangle) = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -2 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{pmatrix}. \quad (3.29)$$

The highest root θ coincides with that of the rank four simple subalgebra $\mathfrak{B}_4 \supset \mathfrak{D}_4$ (respectively, of the real forms $so(5, 4) \supset so(4, 4)$):

$$\theta = \Lambda_0 + \Lambda_1 (= 2\alpha_1 + 3\alpha_2 + 4\alpha_4 + 2\alpha_0). \quad (3.30)$$

The elements D of $so(8)$ act on X of \mathcal{J} (3.10) through their action on the octonions.

$$DX = F_1(Dx_1) + F_2(\kappa(D)x_2) + F_3(\pi(D)x_3), \quad (3.31)$$

where $D =: D_1$, $\kappa(D) =: D_2$, $\pi(D) =: D_3$, obey the principle of infinitesimal triality:

$$(D_1x)y + x(D_2y) = (D_3((xy)^*))^*. \quad (3.32)$$

For $D \in G_2$ we have $D_1 = D_2 = D_3 = D$.

The remaining 24 generators of \mathfrak{f}_4 (outside $so(8)$) can be identified with the skew-hermitian matrices $A_i(e_a)$, $i = 1, 2, 3$, $a = 0, 1, \dots, 7$

$$\begin{aligned} A_1(x) &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & x \\ 0 & -x^* & 0 \end{pmatrix}, \quad A_2(x) = \begin{pmatrix} 0 & 0 & -x^* \\ 0 & 0 & 0 \\ x & 0 & 0 \end{pmatrix}, \\ A_3(x) &= \begin{pmatrix} 0 & x & 0 \\ -x^* & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (3.33)$$

They act on \mathcal{J} through the commutators

$$\tilde{A}_i(e_a)X = \frac{1}{2}[A_i(e_a), X], \quad i = 1, 2, 3, \quad a = 0, 1, \dots, 7. \quad (3.34)$$

The Borel - de Siebenthal theory [BdS] describes the maximal rank closed connected subgroups of a compact Lie group. In order to reveal the physical meaning of the symmetry of $\mathcal{H}_3(\mathbb{O})$ we shall consider, elaborating on [TD],

those maximal rank subgroups of F_4 which contain the (unbroken) colour symmetry group $SU(3)_c \subset G_2 \subset F_4$. There are two such subgroups:

$$\frac{SU(3) \times SU(3)}{\mathbb{Z}_3} \quad \text{and} \quad \text{Spin}(9). \quad (3.35)$$

Postponing the study of the maximal subgroup $\text{Spin}(9)$ to Sect. 4.1 we shall display here the action of

$$F_4^\omega = \frac{SU(3) \times SU(3)}{\mathbb{Z}_3}, \quad \omega(a + z^j e_j) = a + \omega_7 z^j e_j \quad (\omega^3 = 1 = \omega_7^3) \quad (3.36)$$

$$(a = a^0 + a^7 e_7, \quad z^j e_j = z^1 e_1 + z^2 e_2 + z^4 e_4)$$

(cf. (2.32)) on the exceptional Jordan algebra. To do that we shall first extend the splitting of the octonions $\mathbb{O} = \mathbb{C} \oplus \mathbb{C}^3$ to a splitting of the exceptional Jordan algebra, $\mathcal{H}_3(\mathbb{O}) = \mathcal{H}_3(\mathbb{C}) \oplus \mathbb{C}[3]$:

$$(\mathcal{H}_3(\mathbb{O}) \ni) X(\xi, x) = \begin{pmatrix} \xi_1 & x_3 & x_2^* \\ x_3^* & \xi_2 & x_1 \\ x_2 & x_1^* & \xi_3 \end{pmatrix} = X(\xi, a) + X(0, \mathbf{z}\mathbf{e}) \quad (3.37)$$

where

$$\begin{aligned} X(\xi, a) &= \begin{pmatrix} \xi_1 & a_3 & \bar{a}_2 \\ \bar{a}_3 & \xi_2 & a_1 \\ a_2 & \bar{a}_1 & \xi_3 \end{pmatrix}, \\ a_r &= x_r^0 + x_r^7 e_7, \quad \bar{a}_r = x_r^0 - x_r^7 e_7, \quad r = 1, 2, 3; \\ X(0, \mathbf{z}\mathbf{e}) &= \begin{pmatrix} 0 & \mathbf{z}_3 \mathbf{e} & -\mathbf{z}_2 \mathbf{e} \\ -\mathbf{z}_3 \mathbf{e} & 0 & \mathbf{z}_1 \mathbf{e} \\ \mathbf{z}_2 \mathbf{e} & -\mathbf{z}_1 \mathbf{e} & 0 \end{pmatrix}, \\ \mathbf{z}_r \mathbf{e} &= z_r^1 e_1 + z_r^2 e_2 + z_r^4 e_4, \quad z_r^j = x_r^j + x_r^{3j(\text{mod}7)} e_7, \end{aligned} \quad (3.38)$$

(we have used the conjugation property $(\mathbf{z}\mathbf{e})^* = -\mathbf{z}\mathbf{e}$ of imaginary octonions). Multiplication mixes the two terms in the right hand side of (3.37). The Freudenthal product $X(\xi, x) \times Y(\eta, b)$ can be expressed in a nice compact way if we substitute the skew symmetric octonionic matrices $X(0, \mathbf{z}\mathbf{e}), X(0, \mathbf{w}\mathbf{e})$ by 3×3 complex matrices Z, W :

$$X(0, \mathbf{z}\mathbf{e}) \longleftrightarrow Z = (z_r^j, \quad r = 1, 2, 3, \quad j = 1, 2, 4) \in \mathbb{C}[3], \quad (3.39)$$

which transform naturally under the subgroup (3.36).

Indeed, using the fact that the matrices $X(0, \mathbf{ze})$ and $X(0, \mathbf{we})$ are traceless, their Fredenthal product (3.13) simplifies and we find:

$$\begin{aligned} X(\xi, a) \times X(0, \mathbf{we}) &= X(\xi, a) \circ X(0, \mathbf{we}) - \frac{\xi_1 + \xi_2 + \xi_3}{2} X(0, \mathbf{we}) \\ \implies X(\xi, a) \times W &= -\frac{1}{2} W X(\xi, a), \text{ for } W = (w_r^j); \end{aligned} \quad (3.40)$$

$$\begin{aligned} X(0, \mathbf{ze}) \times X(0, \mathbf{we}) &= X(0, \mathbf{ze}) \circ X(0, \mathbf{we}) - \frac{1}{2} \text{tr}(X(0, \mathbf{ze})X(0, \mathbf{we}))E \\ X(0, \mathbf{ze}) \times X(0, \mathbf{we}) &\leftrightarrow -\frac{1}{2}(W^*Z + Z^*W + \bar{Z} \times \bar{W}) \end{aligned} \quad (3.41)$$

where $Z \times W = (\epsilon_{rst}(\mathbf{z}_s \times \mathbf{w}_t)^j)$, so that

$$\begin{aligned} (X(\xi, a) + Z) \times (X(\eta, b) + W) &= X(\zeta, c) + V \\ X\zeta, c) &= X(\xi, a) \times X(\eta, b) - \frac{1}{2}(Z^*W - W^*Z) \\ V &= -\frac{1}{2}(WX(\xi, a) + ZX(\eta, b) + \bar{Z} \times \bar{W}). \end{aligned} \quad (3.42)$$

Thus, if we set $V = (v_r^j)$ we would have

$$\begin{aligned} 2\mathbf{v}_1 &= -\xi_1 \mathbf{w}_1 - \bar{a}_3 \mathbf{w}_2 - a_2 \mathbf{w}_3 - \bar{\mathbf{z}}_2 \times \bar{\mathbf{w}}_3 \\ 2\mathbf{v}_2 &= -a_3 \mathbf{w}_1 - \xi_2 \mathbf{w}_2 - \bar{a}_1 \mathbf{w}_3 - \bar{\mathbf{z}}_3 \times \bar{\mathbf{w}}_1 \\ 2\mathbf{v}_3 &= -\bar{a}_2 \mathbf{w}_1 - a_1 \mathbf{w}_2 - \xi_3 \mathbf{w}_3 - \bar{\mathbf{z}}_1 \times \bar{\mathbf{w}}_2. \end{aligned}$$

The inner product in \mathcal{J} is expressed in terms of the components $X(\xi, a)$ and Z as:

$$\begin{aligned} (X, Y) (= \text{tr} X \circ Y) &= (X(\xi, a), X(\eta, b)) + 2(Z, W) \\ \text{where } 2(Z, W) &= \text{Tr}(Z^*W + W^*Z) = 2 \sum_{r=1}^3 \sum_{j=1,2,4} (\bar{z}_r^j w_r^j + \bar{w}_r^j z_r^j). \end{aligned} \quad (3.43)$$

In the applications to the standard model of particle physics the (upper) index j of z ($j = 1, 2, 4$) labels quark's colour while $r \in \{1, 2, 3\}$ is a *flavour* index. The $SU(3)$ subgroup of G_2 , displayed in Sect. 2 acting on individual (imaginary) octonions is the colour group.

The subgroup F_4^ω (3.36) is defined as the commutant of the automorphism ω of order three in F_4 (see (2.32)):

$$\begin{aligned}\omega X(\xi, x) &= \begin{pmatrix} \xi_1 & \omega x_3 & (\omega x_2)^* \\ (\omega x_3)^* & \xi_2 & \omega x_1 \\ \omega x_2 & (\omega x_1)^* & \xi_3 \end{pmatrix}, \\ \omega(X(\xi, a) + Z) &= X(\xi, a) + \omega_7 Z.\end{aligned}\tag{3.44}$$

The automorphisms $g \in F_4^\omega$ that commute with ω (3.44) are given by pairs $g = (A, U) \in SU(3) \times SU(3)$ acting on $\mathcal{H}_3(\mathbb{O})$ by

$$(A, U)(X(\xi, a) + Z) = AX(\xi, a)A^* + UZA^*.\tag{3.45}$$

The central subgroup

$$\mathbb{Z}_3 = \{(1, 1), (\omega_7, \omega_7), (\omega_7^2, \omega_7^2)\} \in SU(3) \times SU(3)\tag{3.46}$$

acts trivially on $\mathcal{H}_3(\mathbb{O})$. We see that the unitary matrix U acts (in (3.44)) on the colour index j and hence belongs to the (unbroken) *colour group* $SU_c(3)$, while the action of A on the flavour indices will be made clear in Sect. 4.2 below.

4 F_4 as a grand unified symmetry of the standard model

This chapter provides a tentative application of the exceptional Jordan algebra to the standard model (SM) of particle physics. We first study, in Sect. 4.1, the special Jordan subalgebra $JSpin_9$ of \mathcal{J} and its automorphism group $Spin(9) \subset F_4$ singling out the 16-dimensional spinor representation of $Spin(9)$ and interpret it in terms of the fundamental fermionic doublets of the SM. Then, in Sect.4.2, we examine the full 26-dimensional representation $\underline{26}$ of F_4 , considering its restriction to both its maximal rank subgroups that contain the colour $SU(3)_c$ as subgroup. We demonstrate that $\underline{26}$ gives also room, to the electroweak gauge bosons.

4.1 The Jordan subalgebra $JSpin_9$ of $\mathcal{H}_3(\mathbb{O})$

The ten dimensional Jordan algebra $JSpin_9$ can be identified with the algebra of 2×2 hermitian octonionic matrices $\mathcal{H}_2(\mathbb{O})$ equipped with the Jordan

matrix product. It is generated by the 9-dimensional vector subspace $s\mathcal{H}_2(\mathbb{O})$ of traceless matrices of $\mathcal{H}_2(\mathbb{O})$ whose square is, in fact, a positive real scalar:

$$X = \begin{pmatrix} \xi & x \\ x^* & -\xi \end{pmatrix} \Rightarrow X^2 = (\xi^2 + x^*x)1, \quad x \in \mathbb{O}, \xi \in \mathbb{R}. \quad (4.1)$$

$JSpin_9$ is a (special) Jordan subalgebra of the (associative) matrix algebra $\mathbb{R}[2^4]$ that provides an irreducible representation of Cl_9 . Clearly, it is a subalgebra of $\mathcal{H}_3(\mathbb{O})$ -consisting of 3×3 matrices with vanishing first row and first column. Its automorphism group is the subgroup $Spin(9) \subset F_4$ which stabilizes the projector E_1 : $Spin(9) = (F_4)_{E_1} \subset F_4$.

We shall see that the spinor representation of $Spin_9$ can be interpreted as displaying the first generation of (left chiral) doublets of quarks and leptons

$$\begin{pmatrix} \nu_L & u_L^j \\ e_L^- & d_L^j \end{pmatrix} \quad (j \text{ is the colour index})$$

and their antiparticles. As demonstrated in [D-V.T] and elaborated in Sect. 4.2 below (Eqs.(4.17-21) the exceptional Jordan algebra gives room to a similar construction of the second and third generation fermions.

The 16-dimensional (real) spinor representation S_9 of $Spin(9)$ splits into a direct sum of the two 8-dimensional chiral representations S_8^+ and S_8^- of $Spin(8)$ that appear as eigenvectors of the Coxeter element ω_8 of Cl_8 :

$$S_9 = S_8^+ \oplus S_8^-, \quad \omega_8 S_8^\pm = \pm S_8^\pm. \quad (4.2)$$

We shall use the relation (2.38) ($\hat{P}_0 = \sigma_1 \otimes P_0$, $\hat{P}_a = c \otimes P_a$, $a = 1, \dots, 7$) with real P_j ($j = 1, 2, 4$) and iP_7, iP_3, iP_6, iP_5 .

$$\begin{aligned} P_1 &= \mathbb{1} \otimes \sigma_1 \otimes c, & P_2 &= \sigma_1 \otimes \sigma_3 \otimes c^*, & P_4 &= c \otimes \sigma_1 \otimes \sigma_1 \\ iP_7 &= \mathbb{1} \otimes \mathbb{1} \otimes \sigma_3, & iP_3 &= \sigma_3 \otimes \sigma_1 \otimes \sigma_1, & iP_6 &= -\sigma_1 \otimes \mathbb{1} \otimes \sigma_1, \\ iP_5 &= \sigma_1 \otimes c \otimes c, & \Rightarrow \omega_{-7} &:= P_1 P_2 P_3 P_4 P_5 P_6 P_7 = -\mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1}, \end{aligned} \quad (4.3)$$

$$\mathbb{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad c = i\sigma_2, \quad c^* = -c = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

The Coxeter element ω_8 of Cl_8 , appearing in (4.2) is given by:

$$\hat{P}_8 = \omega_8 = \hat{P}_0 \hat{P}_1 \dots \hat{P}_7 = \sigma_3 \otimes \omega_{-7} = -\sigma_3 \otimes \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1}. \quad (4.4)$$

Inserting the matrices in the tensor products from right to left, so that

$$\mathbb{I} \otimes \sigma_3 = \begin{pmatrix} \sigma_3 & 0 \\ 0 & \sigma_3 \end{pmatrix}, \quad \sigma_3 \otimes \mathbb{I} = \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix}, \quad \sigma_3 \otimes \sigma_3 = \begin{pmatrix} \sigma_3 & 0 \\ 0 & -\sigma_3 \end{pmatrix}$$

we can write the weight matrices Λ_μ (2.41) as:

$$\begin{aligned} \Lambda_0 &= \frac{i}{2} \hat{P}_{07} = -\frac{1}{2} \sigma_3 \otimes \mathbb{I} \otimes \mathbb{I} \otimes \sigma_3 = -\frac{1}{2} \text{diag}(\sigma_3, \sigma_3, \sigma_3, \sigma_3, -\sigma_3, -\sigma_3, -\sigma_3, -\sigma_3) \\ \Lambda_1 &= \frac{i}{2} \hat{P}_{13} = -\frac{1}{2} \mathbb{I} \otimes \sigma_3 \otimes \mathbb{I} \otimes \sigma_3 = -\frac{1}{2} \mathbb{I} \otimes \text{diag}(\sigma_3, \sigma_3, -\sigma_3, -\sigma_3) \\ \Lambda_2 &= \frac{i}{2} \hat{P}_{26} = -\frac{1}{2} \mathbb{I} \otimes \mathbb{I} \otimes \sigma_3 \otimes \sigma_3 = -\frac{1}{2} \mathbb{I} \otimes \text{diag}(\sigma_3, -\sigma_3, \sigma_3, -\sigma_3) \\ \Lambda_4 &= \frac{i}{2} \hat{P}_{45} = -\frac{1}{2} \mathbb{I} \otimes \sigma_3 \otimes \sigma_3 \otimes \sigma_3 = -\frac{1}{2} \mathbb{I} \otimes \text{diag}(\sigma_3, -\sigma_3, -\sigma_3, \sigma_3). \end{aligned} \quad (4.5)$$

They form a (commuting) orthonormal basis in the weight space:

$$\Lambda_\mu^2 = \frac{1}{4} 1_{16}, \quad [\Lambda_\mu, \Lambda_\nu] = 0, \quad \langle \Lambda_\mu, \Lambda_\nu \rangle := \frac{1}{4} \text{tr}(\Lambda_\mu \Lambda_\nu) = \delta_{\mu\nu}, \quad \mu, \nu = 0, 1, 2, 4.$$

The Lie algebra $\mathfrak{B}_4 (= \mathfrak{so}(9))$ of Spin(9) is spanned by the commutators $\hat{P}_{ab} = \frac{1}{2} [\hat{P}_a, \hat{P}_b]$, $a, b = 0, 1, \dots, 8$. The physical meaning of \mathfrak{B}_4 is best revealed by identifying its maximal (rank 4) subalgebra

$$\mathfrak{su}(2) \oplus \mathfrak{su}(4) \cong \mathfrak{so}(3) \oplus \mathfrak{so}(6) \in \mathfrak{so}(9). \quad (4.6)$$

that is part of the *Pati-Salam grand unified Lie algebra*

$$\mathfrak{g}_{PS} := \mathfrak{su}(2)_L \oplus \mathfrak{su}(2)_R \oplus \mathfrak{su}(4). \quad (4.7)$$

(See for a nice review [BH].) The SU(4) of Pati-Salam is designed to unify the quark colour group SU(3)_c with the lepton number. The colour Lie algebra $\mathfrak{su}(3)_c$ is identified with the commutant in $\mathfrak{su}(4)$ of the hypercharge $Y(\underline{4})$. In the defining 4-dimensional representation $\underline{4}$ of $\mathfrak{su}(4)$ it is given by the traceless diagonal matrix

$$Y(\underline{4}) = \begin{pmatrix} \frac{1}{3} & 0 & 0 & 0 \\ 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & \frac{1}{3} & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \quad (4.8)$$

With our splitting of the octonions $\mathbb{O} = \mathbb{C} + \mathbb{C}^3$ and the corresponding embedding of $SU(3)_c$ into G_2 (see (2.11)) the subalgebra $\mathfrak{so}(6)$ is spanned by \hat{P}_{jk} with $1 \leq j < k \leq 6$. The Lie subalgebra $\mathfrak{su}(3)_c$ appears as the intersection of \mathfrak{g}_2 and $\mathfrak{so}(6)$ in $\mathfrak{so}(7)$. As a consequence of (2.43) it is spanned by the following combinations of \hat{P}_{ab} :

$$\begin{aligned} \mathfrak{su}(3)_c = \text{Span}\{ & \hat{P}_{13} - \hat{P}_{26} (= -iH_1), \quad \hat{P}_{26} - \hat{P}_{45} (-iH_2), \\ & \hat{P}_{12} + \hat{P}_{36}, \quad \hat{P}_{24} + \hat{P}_{45}, \quad \hat{P}_{14} + \hat{P}_{35} \\ & \hat{P}_{16} + \hat{P}_{23}, \quad \hat{P}_{25} + \hat{P}_{46}, \quad \hat{P}_{15} + \hat{P}_{43}\}. \end{aligned} \quad (4.9)$$

The first two span the Cartan subalgebra of $\mathfrak{su}(3)_c$ - cf. (3.28). The next three span the maximal compact subalgebra $\mathfrak{so}(3)$ of the real form $\mathfrak{sl}(3, \mathbb{R}) \subset \mathfrak{so}(4, 4)$. The generator Y of the commutant of $\mathfrak{su}(3)_c$ in (the complexification of) $\mathfrak{su}(4)$ is given by

$$Y = \frac{2}{3} (\Lambda_1 + \Lambda_2 + \Lambda_4) = \frac{i}{3} (\hat{P}_{13} + \hat{P}_{26} + \hat{P}_{45}) \quad (\in \mathfrak{so}(3, 3)), \quad (4.10)$$

$$3Y = \mathbb{1}_4 \otimes \begin{pmatrix} -3\sigma_3 & 0 & 0 & 0 \\ 0 & \sigma_3 & 0 & 0 \\ 0 & 0 & \sigma_3 & 0 \\ 0 & 0 & 0 & \sigma_3 \end{pmatrix} = \mathbb{1}_4 \otimes \text{diag}(-3\sigma_3, \sigma_3, \sigma_3, \sigma_3).$$

It (commutes with and) is orthogonal to the Cartan $\{H_1, H_2\}$ of $\mathfrak{su}(3)_c$. The algebra $\mathfrak{su}(2)$ in the left hand side of (4.6) is then spanned by

$$\begin{aligned} I_3 := \Lambda_0 = \frac{i}{2} \hat{P}_{07}, \quad I_+ = \hat{P}_0^- \hat{P}_8, \quad I_- = \hat{P}_8 \hat{P}_0^+ \\ ([I_3, I_\pm] = \pm I_\pm, \quad [I_+, I_-] = 2I_3). \end{aligned} \quad (4.11)$$

The spinor representation $\underline{16}$ of $\text{Spin}(9)$ can now be associated with the doublet representation

$$(\underline{4}^*, \underline{2}, \underline{1}) \oplus (\underline{4}, \underline{2}, \underline{1}) \in \mathfrak{g}_{PS}. \quad (4.12)$$

‘ It is natural to identify Λ_0 which commutes with $\mathfrak{su}(4)$ and has eigenvalues $\pm \frac{1}{2}$ with the third component I_3 of the weak isospin, while Y that commutes with $\mathfrak{su}(3)_c$ and is orthogonal to I_3 should coincide with the weak

hypercharge. The spinor representation consists of two octets of the fermion (lepton-quark) left chiral doublets and antifermion right chiral doublets

$$\begin{aligned} & \begin{pmatrix} \nu_L \\ e_L^- \end{pmatrix}, \quad Y = -1, \quad \begin{pmatrix} u_L \\ d_L \end{pmatrix}, \quad Y = \frac{1}{3}, \quad I_3 = \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}, \\ & \begin{pmatrix} e_R^+ \\ \bar{\nu}_R \end{pmatrix}, \quad Y = 1, \quad \begin{pmatrix} \bar{d}_R \\ \bar{u}_R \end{pmatrix}, \quad Y = -\frac{1}{3}, \quad I_3 = \begin{pmatrix} \frac{1}{2} \\ -\frac{1}{2} \end{pmatrix}. \end{aligned} \quad (4.13)$$

These are the fundamental (anti) fermions that participate in the weak interactions. The missing fundamental fermion representations are the sixteen singlets with respect to the weak isospin, $I = 0 (= I_3)$.

4.2 The representation 26 of F_4

As the automorphism group F_4 of $\mathcal{J} = \mathcal{H}_3(\mathbb{O})$ preserves the unit element and the trace of \mathcal{J} it acts faithfully and irreducibly on the 26-dimensional subspace $\mathcal{J}_0 = s\mathcal{H}_3(\mathbb{O}) \subset \mathcal{J}$ of traceless 3×3 hermitian octonionic matrices. In fact, 26 is the lowest nontrivial (fundamental) representation (of highest weight Λ_0) of F_4 . Restricted to the maximal subgroup $\text{Spin}(9)$ of F_4 it splits into three irreducible components:

$$\underline{26} = \underline{16} + \underline{9} + \underline{1}. \quad (4.14)$$

We have identified in the preceding Sect. 4.1 the spinor representation as four isospin chiral doublets of quarks and leptons and their antiparticles. The 9-vector representation of $\mathfrak{so}(9)$ is spanned by the generators \hat{P} of $C\ell_9$. The matrices \hat{P}_μ^ϵ (2.40) and \hat{P}_8 diagonalize the adjoint action of the physical Cartan elements Y (4.10) and I_3 (4.11); we find:

$$\begin{aligned} [I_3, \hat{P}_0^\mp] &= \pm \hat{P}_0^\mp, & [I_3, \hat{P}_8] &= 0 = [I_3, \hat{P}_j^\epsilon], \quad (j = 1, 2, 4) \\ [Y, \hat{P}_0^\mp] &= 0 = [Y, \hat{P}_8], & [Y, \hat{P}_j^\mp] &= \pm \frac{2}{3} \hat{P}_j^\mp. \end{aligned} \quad (4.15)$$

This allows to consider \hat{P}_0^\mp as representatives of the W^\pm bosons while \hat{P}_8 appears as the neutral component, W^0 , of the isotopic triplet. Then the Z boson and the photon can be associated with appropriate mixtures of W^0 and the trivial representation of $\mathfrak{so}(9)$. The isotropic matrices \hat{P}_j^\pm , on the other hand, carry the quantum numbers ($I_3 = 0$, $Y = \mp \frac{2}{3}$) of the right handed

d -quarks d_R and the left handed d -antiquarks, \bar{d}_L , respectively ($j = 1, 2, 4$ being the colour index). Let us note that all eigenvalues of the pair (I_3, Y) appearing in the representation $\underline{26}$ of F_4 satisfy the SM constraint

$$I_3 + \frac{3}{2}Y \in \mathbb{Z}.$$

We recall that according to the identification of the 2^5 -dimensional fermion Fock space with the exterior algebra $\Lambda\mathbb{C}^5$ (see [BH]) all fundamental fermions (of the first generation) can be expressed as exterior products of d_R, e_R^+ , and $\bar{\nu}_R$ (with $\bar{\nu}_L$ identified with the Fock vacuum). In particular, all fundamental fermions can be written as wedge products of d_R, \bar{d}_L and the chiral spinors in (4.13):

$$\begin{aligned} e_L^+(I_3 = 0, Y = 2) &= e_R^+ \wedge \bar{\nu}_R \\ u_R(I_3 = 0, Y = \frac{4}{3}) &= e_L^+ \wedge d_R \\ \bar{u}_L^j(I_3 = 0, Y = -\frac{4}{3}) &= \epsilon^{jkl} d_R^k \wedge d_R^l \\ e_R^-(I_3 = 0, Y = -2) &= \sum_{j=1,2,4} \bar{u}_L^j \wedge d_R^j \\ \nu_R(I_3 = 0 = Y) &= \sum_j \bar{d}_L^j \wedge d_R^j, \quad (j, k, l = 1, 2, 4). \end{aligned} \quad (4.16)$$

(Here ϵ^{ijk} is the fully antisymmetric unit tensor with $\epsilon^{124} = 1$.) The isosinglets $\bar{\nu}_L, d_R, \bar{d}_L$ and the wedge products (4.16) together with the isotopic doublets (4.13) exhaust the 32 first generation fundamental fermions of the SM (cf. Sect. 5.3). We assume that right handed fermions have negative chirality so that wedge product with d_R or $\bar{\nu}_R$ changes the sign of chirality (i.e. transforms right to left and vice versa). If we associate the Higgs boson with the trace (or the unit element) of the exceptional Jordan algebra \mathcal{J} we see that all fundamental particles of the first family of the SM are generated either directly or as exterior products of elements of \mathcal{J} . The same assignment of quantum numbers to fundamental particles is obtained, as it should, if we consider the reduction of F_4 with respect to the other admissible maximal subgroup F_4^ω (3.36). On the way of demonstrating this we shall express the third component of the weak isospin I_3 and the hypercharge Y in terms of the Cartan elements of the flavour $SU(3)$. To begin with we introduce the

three Weyl units ϵ_i and their conjugate (=transposed) ϵ_i^* ($i = 1, 2, 3$) by

$$\epsilon_1^* = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad \epsilon_2^* = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \epsilon_3^* = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (4.17)$$

and set

$$\begin{aligned} 2I_3 &= [\epsilon_1^*, \epsilon_1] = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix} \\ 3Y &= [\epsilon_3^*, \epsilon_3] - [\epsilon_2^*, \epsilon_2] = \begin{pmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}. \end{aligned} \quad (4.18)$$

We see that $\epsilon_i^{(*)}$ are eigenvectors of I_3 and Y under their adjoint action corresponding to the (first generation) leptons and to the W^\pm bosons:

$$\begin{aligned} [I_3, \epsilon_1^*] &= \epsilon_1^*, \quad [Y, \epsilon_1^*] = 0, \quad \epsilon_1^* \leftrightarrow W^+ \\ [I_3, \epsilon_2^*] &= -\frac{1}{2}\epsilon_2^*, \quad [Y, \epsilon_2^*] = -\epsilon_2^* \quad \epsilon_2^* \leftrightarrow e_L^- \\ [I_3, \epsilon_3^*] &= \frac{1}{2}\epsilon_3^*, \quad [Y, \epsilon_3^*] = -\epsilon_3^* \Rightarrow \epsilon_3^* \leftrightarrow \nu_L \\ (\epsilon_1 &\leftrightarrow W^-, \quad \epsilon_2 \leftrightarrow e_R^+, \quad \epsilon_3 \leftrightarrow \bar{\nu}_R). \end{aligned} \quad (4.19)$$

The form of I_3 and Y depends on the generation, - i.e., on the choice of E_i which commutes with the corresponding subalgebra $J_2^8 \subset J_3^8$. We have:

$$\begin{aligned} 2I_3^{(i)} &= [\epsilon_i^*, \epsilon_i], \quad 3Y^{(i)} = [\epsilon_{i-1}^*, \epsilon_{i-1}] - [\epsilon_{i+1}^*, \epsilon_{i+1}] \\ &\text{for } i = 1, 2, 3 \pmod{3}. \end{aligned} \quad (4.20)$$

Correspondingly, the W^\pm boson for the i -th generation is associated with $\epsilon_i^{(*)}$. The generations are permuted by the element

$$\tau = \epsilon_1 + \epsilon_2 + \epsilon_3 \Rightarrow \tau \epsilon_i^{(*)} \tau^* = \epsilon_{i+1 \pmod{3}}^{(*)}, \quad \tau^3 = 1. \quad (4.21)$$

(where $\epsilon_i^{(*)}$ stands for either ϵ_i or ϵ_i^*). Coming back to the first generation, we note that the operators I_3 and Y act on the part $X(\xi, a)$ (3.38) of \mathcal{J}

by commutation and on the matrix Z (3.39) by right multiplication with $(-I_3, -Y)$. Their eigenvectors are given by

$$Z_1 = (z_1^j, 0, 0), \quad Z_2 = (0, z_2^j, 0), \quad Z_3 = (0, 0, z_3^j) \quad (4.22)$$

$$\begin{aligned} Z_1(-I_3) &= 0, & Z_2(-I_3) &= -\frac{1}{2}Z_2, & Z_3(-I_3) &= \frac{1}{2}Z_3 \\ Z_1(-Y) &= -\frac{2}{3}Z_1 & Z_{2,3}(-Y) &= \frac{1}{3}Z_{2,3}, \end{aligned} \quad (4.23)$$

while their conjugates have the same eigenvalues with opposite sign and the diagonal matrices $X(\xi, 0)$ correspond to $I_3 = Y = 0$ ($I = 0, 1$)⁷. Comparing with the quantum numbers of (4.13) we end up with the following correspondence

$$\begin{aligned} Z_1 &\rightarrow d_R, & Z_2 &\rightarrow d_L, & Z_3 &\rightarrow u_L, \\ (\bar{Z}_1 &\rightarrow \bar{d}_L, & \bar{Z}_2 &\rightarrow \bar{d}_R, & \bar{Z}_3 &\rightarrow \bar{u}_R). \end{aligned} \quad (4.24)$$

Thus the basic representation $\underline{26}$ of F_4 splits when restricted to $SU(3)_f \times SU(3)_c$ into the following irreducible components:

$$\underline{26} = \bar{\underline{3}} \otimes \underline{3} + \underline{3} \otimes \bar{\underline{3}} + \underline{8} \otimes \underline{1} \quad (4.25)$$

where the adjoint representation $\underline{8}$ of the flavour $SU(3) = SU(3)_f$ consists of two doublets (of leptons and antileptons) and of the three massive gauge bosons and the photon.

4.3 The symmetry group of the standard model

It has been observed by Baez and Huerta [BH] that the gauge group of the SM,

$$G_{SM} = S(U(2) \times U(3)) = \frac{SU(2) \times SU(3) \times U(1)}{\mathbb{Z}_6} \quad (4.26)$$

can be obtained as the intersection of the Georgi-Glashow and Pati-Salam *grand unified theory groups* $SU(5)$ and $(SU(4) \times SU(2) \times SU(2))/\mathbb{Z}_2$ viewed

⁷The element $X(\xi, 0) = \xi_1 \Lambda_3$ carries total isospin $I = 1$ while $\xi_2 \Lambda_8$ corresponds to $I = 0$, both having $I_3 = 0$.

as subgroups of Spin(10). Here we elaborate on the suggestion of [TD] that one can deduce the symmetry of the SM by applying the Borel - de Siebenthal theory to *admissible* maximal rank subgroups of F_4 - those that contain the exact colour symmetry $SU(3)_c$. Our analysis in this chapter demonstrates that the intersection of the two admissible maximal subgroups Spin(9) and $\frac{SU(3) \times SU(3)}{\mathbb{Z}_3}$ of F_4 is precisely the symmetry group G_{SM} (4.23) of the SM. The constraints on the $U(1)_Y$ term coming from factoring the 6-element central subgroup \mathbb{Z}_6 imply

$$I_3 + \frac{3}{2}Y (= Q + Y) \in \mathbb{Z} \quad (4.27)$$

(where $Q = I_3 + \frac{1}{2}Y$ is the electric charge); furthermore, if the central element of $SU(3)_c$ is represented by a nontrivial eigenvalue ω , i.e. if

$$\omega^2 + \omega + 1 = 0 \quad \text{then} \quad U_Y^2 + U_Y + 1 = 0 \quad \text{for} \quad U_Y = e^{2\pi i Y}. \quad (4.28)$$

The "colourless" Cartan elements I_3, Y complemented with the total isospin I , ($I_1^2 + I_2^2 + I_3^2 = I(I + 1)$) and the chirality γ (which has eigenvalue 1 for left and -1 for right chiral fermions) completely characterize all 32 first generation fermions as well as the four electroweak gauge bosons. The colour index j can be related to the eigenvalues of the $SU(3)_c$ Cartan matrices, but different "colours" are physically indistinguishable. We only need the chirality γ to separate $\nu_R(\gamma = -1)$ from $\bar{\nu}_L(\gamma = 1)$.

5 Extended euclidean Jordan algebra; the 32 primitive idempotents of the fermion states

The purpose of this section is to relate our treatment of fundamental fermions in Sect. 4 with the recent realization [D-V.T] that a natural euclidean extension of the special Jordan algebra $J_2^8 = JSpin_9$ plays the role of the quantum observable algebra of the first generation of the SM. We shall proceed in three steps. First, we recall the notion of an euclidean extension of a special Jordan algebra [D-V.T] as applied to J_2^8 . Secondly, we provide a fermion Fock space construction of the complexified associative envelope Cl_9 of J_2^8 , inspired by a similar construction by Furey [F18]. We construct 32 *primitive idempotents* [A95, T85/16] which diagonalize three commuting superselection rules and span two irreducible Weyl spinor representations of the Lie algebra $so(9, 1)$.

Finally, we recover the octet of weak doublets of Sect. 4.1 by identifying the weak isospin I_3 with the difference $I_3^L - I_3^R$ of the "left" and the "right" Pati-Salam isospins.

5.1 Euclidean extension of J_2^8 . Fermion Fock space

The associative envelope of the spin factor J_2^8 is given by the real Clifford algebra $Cl_9 = \mathbb{R}[16] \oplus \mathbb{R}[16]$ which is again a Jordan algebra (under anticommutation) but is not euclidean. As observed in [D-V.T] the Jordan subalgebra of hermitian matrices $\mathcal{H}_{16}(\mathbb{C}) \oplus \mathcal{H}_{16}(\mathbb{C})$ of the complexified Clifford algebra $Cl_9(\mathbb{C})$ is euclidean (and has the same real dimension 2×16^2 , as the above real associative envelope). It provides a *canonical euclidean extension* of the 10-dimensional Jordan algebra J_2^8 . The splitting of Cl_9 into two real matrix algebras corresponds to the splitting of the spinorial representation of $so(9,1)$ into two irreducible 16 dimensional chiral Majorana-Weyl spinors. Indeed the Lie algebra $so(9,1)$ belongs to the even part $Cl^0(9,1)$ of the Clifford algebra $Cl(9,1)$ which is isomorphic to Cl_9 . It is instructive to present (following [BH09]) the mapping Γ from J_2^8 to $Cl(9,1)$ through its action on $\underline{16}^+ \oplus \underline{16}^-$:

$$\Gamma(X)(\psi_+, \psi_-) = (\tilde{X}\psi_-, X\psi_+) \quad (5.1)$$

where \tilde{X} is the *trace reversal* $\tilde{X} = X - (tr X)1$.

$$a_\mu^* = \Gamma_\mu^-, \quad a_\mu = \Gamma_\mu^+, \quad \mu = 0, 1, 2, 4 \quad \Gamma_\mu^\pm = \sigma_1 \otimes \hat{P}_\mu^\pm \quad \text{see eq. (2.40)}$$

with another pair (a_8^*, a_8) with all five pairs satisfying

$$\begin{aligned} a_\mu + a_\mu^* &= \Gamma_\mu, \quad \mu = 0, 1, 2, 4, 8, \quad a_0 - a_0^* = i\Gamma_7, \\ a_j - a_j^* &= \Gamma_{3j(mod7)}, \quad a_8 - a_8^* = \Gamma_{-1} \end{aligned} \quad (5.2)$$

where $\Gamma_{-1}(= \Gamma(1)$ of (5.1)) is the (antihermitian) 10th generator of $Cl(9,1)$ satisfying $\Gamma_{-1}^2 = -1$ ($[\Gamma_{-1}, \Gamma_a]_+ = 0$, $a = 0, 1, \dots, 8$). In accord with our preceding conventions, we identify the electric charge Q , the hypercharge Y and the charge $B - L$ with the quadratic expressions:

$$\begin{aligned} Q &= \frac{1}{3}(a_1^*a_1 + a_2^*a_2 + a_4^*a_4) - a_8^*a_8 \\ Y &= \frac{2}{3}(a_1^*a_1 + a_2^*a_2 + a_4^*a_4) - a_0^*a_0 - a_8^*a_8 \\ B - L &= \frac{1}{3}([a_1^*, a_1] + [a_2^*, a_2] + [a_4^*, a_4]). \end{aligned} \quad (5.3)$$

5.2 Fermion state space in terms of primitive idempotents

The states of an euclidean Jordan algebra are given by idempotents - (hermitian) projection operators. The fermion creation and annihilation operators (5.1) allow to define five pairs of commuting idempotents

$$\pi_\mu = a_\mu a_\mu^*, \quad \bar{\pi}_\mu = a_\mu^* a_\mu, \quad \pi_\mu + \bar{\pi}_\mu = 1, \quad \mu = 0, 1, 2, 4, 8. \quad (5.4)$$

The canonical anticommutation relations imply the defining relation for idempotents:

$$\pi_\mu^2 = \pi_\mu, \quad (\bar{\pi}_\mu^2 = \bar{\pi}_\mu). \quad (5.5)$$

The products of π_μ are again idempotents. An idempotent is called *primitive* if it cannot be decomposed into a sum of (nonzero) mutually orthogonal idempotents. Any product of less than all five π_μ 's is not primitive. For instance, the product of the first four π_μ splits into the following two:

$$\pi_0 \pi_1 \pi_2 \pi_4 = \pi_0 \pi_1 \pi_2 \pi_4 \pi_8 + \pi_0 \pi_1 \pi_2 \pi_4 \bar{\pi}_8.$$

Examples of primitive idempotents are:

$$\Omega = \pi_0 \pi_1 \pi_2 \pi_4 \pi_8, \quad \bar{\Omega} = \bar{\pi}_0 \bar{\pi}_1 \bar{\pi}_2 \bar{\pi}_4 \bar{\pi}_8. \quad (5.6)$$

All mixed products of five factors $\pi_\mu, \bar{\pi}_\nu$ are also primitive. Thus, there are exactly $2^5 = 32$ primitive idempotents corresponding to the pure states of the 32 fermions of a given generation (in our case, the first). We postulate, following [T85/16], that the trace of each primitive idempotent is one. Then the trace of each π_μ (and each $\bar{\pi}_\nu$) is found to be 16 while the trace of the unit operator equals to *the rank, 32 of the extended Jordan algebra*. The chirality

$$\gamma = [a_0^*, a_0][a_1^*, a_1][a_2^*, a_2][a_4^*, a_4][a_8^*, a_8] \quad (5.7)$$

commutes with the generators $\Gamma_{ab} = \frac{1}{2}[\Gamma_a, \Gamma_b]$, $a, b = -1, 0, \dots, 8$ of $so(9, 1)$ and splits its 32 dimensional spinor representation into two irreducible chiral components. The right chiral representation $\underline{16}^-$ is obtained from the lowest

weight state Ω by acting on it with the ten raising operators $a_\mu^* a_\nu^*$ ($\mu < \nu$). The left chiral fermions $\underline{16}^+$ are obtained by acting on the highest weight vector $\bar{\Omega}$ by the ten lowering operators $a_\mu a_\nu$. We note that the operators $a_\mu a_\nu$ and their conjugate belong to the real form $so(5, 5)$ of the complexified Lie algebra $so(10)$. The 25-dimensional maximal rank subalgebra $u(3, 2)$ of $so(5, 5)$ with generators $\frac{1}{2}[a_\mu^*, a_\nu]$ splits each of the chiral 16-plets into three irreducible components:

$$\begin{aligned}
\underline{16}^- &= \underline{1}_R + \underline{10}_R + \underline{5}_R; & \underline{16}^+ &= \underline{1}_L + \underline{10}_L + \underline{5}_L, & \underline{1}_R &= \Omega, & \underline{1}_L &= \bar{\Omega}, \\
\underline{10}_R &= Span\{a_\mu^* a_\nu^* \Omega a_\nu a_\mu, \mu < \nu\}, & \underline{5}_R &= Span\{a_\mu \bar{\Omega} a_\mu^*, \mu = 0, 1, 2, 4, 8\} \\
\underline{10}_L &= Span\{a_\mu a_\nu \bar{\Omega} a_\nu^* a_\mu^*, \mu < \nu\}, & \underline{5}_L &= Span\{a_\mu^* \Omega a_\mu, \mu = 0, 1, 2, 4, 8\}.
\end{aligned} \tag{5.8}$$

They are familiar from the Georgi-Glashow $SU(5)$ Grand Unified Theory [BH, W].

5.3 Nonabelian superselection sectors

The leptons are given by conventional (1-dimensional) pure states characterized by the eigenvalues of the *superselected* charges (in the terminology introduced in [WWW]) $(Q, Y, B - L)$ (5.3). We have:

$$\begin{aligned}
(Q, Y, B - L)\Omega &= (0, 0, -1)\Omega \Rightarrow \Omega = |\nu_R \rangle \langle \nu_R| \\
(Q, Y, B - L)\bar{\Omega} &= (0, 0, 1)\bar{\Omega} \Rightarrow \bar{\Omega} = |\bar{\nu}_L \rangle \langle \bar{\nu}_L|;
\end{aligned} \tag{5.9}$$

similarly

$$\begin{aligned}
a_0^* \Omega a_0 &= |\nu_L \rangle \langle \nu_L| \Leftrightarrow (0, -1, -1) \\
a_8^* \Omega a_8 &= |e_L^- \rangle \langle e_L^-| \Leftrightarrow (-1, -1, -1) \\
a_0^* a_8^* \Omega a_8 a_0 &= |e_R^- \rangle \langle e_R^-| = a_1 a_2 a_4 \bar{\Omega} a_4^* a_2^* a_1^* \Leftrightarrow (-1, -2, -1) \\
a_0 \bar{\Omega} a_0^* &= |\bar{\nu}_R \rangle \langle \bar{\nu}_R| \Leftrightarrow (0, 1, 1), \\
a_8 \Omega a_8^* &= |e_R^+ \rangle \langle e_R^+| \Leftrightarrow (1, 1, 1) \\
a_1^* a_2^* a_4^* \Omega a_4 a_2 a_1 &= |e_L^+ \rangle \langle e_L^+| = a_0 a_8 \bar{\Omega} a_8^* a_0^* \Leftrightarrow (1, 2, 1).
\end{aligned} \tag{5.10}$$

It is not meaningful to characterize similarly the quark states since the "colour" is not observable (as the non-abelian gauge symmetry $SU(3)_c$ is

unbroken). We shall introduce instead $SU(3)_c$ -invariant 3-dimensional projection operators which characterize the observable quark states:

$$\begin{aligned}
a_j^* \Omega a_j &= a_1^* \Omega a_1 + a_2^* \Omega a_2 + a_4^* \Omega a_4 = |\bar{d}_L \rangle \langle \bar{d}_L| \Leftrightarrow \left(\frac{1}{3}, \frac{2}{3}, -\frac{1}{3} \right), \\
a_0^* a_j^* \Omega a_j a_0 &= |\bar{d}_R \rangle \langle \bar{d}_R| \Leftrightarrow \left(\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3} \right) \\
a_j^* a_8^* \Omega a_8 a_j &= |\bar{u}_R \rangle \langle \bar{u}_R| \Leftrightarrow \left(-\frac{2}{3}, -\frac{1}{3}, -\frac{1}{3} \right) \\
a_0^* a_j^* a_8^* \Omega a_8 a_j a_0 &= |\bar{u}_L \rangle \langle \bar{u}_L| \Leftrightarrow \left(-\frac{2}{3}, -\frac{4}{3}, -\frac{1}{3} \right) \\
a_j \bar{\Omega} a_j^* &= |d_R \rangle \langle d_R| \Leftrightarrow \left(-\frac{1}{3}, -\frac{2}{3}, \frac{1}{3} \right) \\
a_0 a_j \bar{\Omega} a_j^* a_0^* &= |d_L \rangle \langle d_L| \Leftrightarrow \left(-\frac{1}{3}, \frac{1}{3}, \frac{1}{3} \right) \\
a_j a_8 \bar{\Omega} a_8^* a_j^* &= |u_L \rangle \langle u_L| \Leftrightarrow \left(\frac{2}{3}, \frac{1}{3}, \frac{1}{3} \right) \\
a_0 a_j a_8 \bar{\Omega} a_8^* a_j^* a_0^* &= |u_R \rangle \langle u_R| \Leftrightarrow \left(\frac{2}{3}, \frac{4}{3}, \frac{1}{3} \right). \tag{5.11}
\end{aligned}$$

(Summation is understood throughout in the repeated index $j = 1, 2, 4$ - as displayed in the first formula.) We note that the antiparticle to a left chiral quark is a right chiral antiquark and vice versa. All three superselection charges (Q,Y,B-L) change sign going from a particle to its antiparticle.

The passage from the 32-dimensional representation of $so(9, 1)$ to the 16-dimensional one $so(9)$, displayed in Sect. 4.1, is similar to the passage from Lorentzian Weyl spinors to the Pauli spinors. To display it we introduce the third components of the "left" and "right" isospin of the Pati-Salam model:

$$2I_3^L = \frac{1}{2} ([a_0^*, a_0] - [a_8^*, a_8]), \quad 2I_3^R = -\frac{1}{2} ([a_0^*, a_0] + [a_8^*, a_8]) \tag{5.12}$$

and identify the I_3 of Sect. 4.1 with their difference

$$I_3 = I_3^L - I_3^R = \frac{1}{2} [a_0^*, a_0]. \tag{5.13}$$

Identifying in addition the hypercharge in the $\underline{16}$ of $so(9)$ with $B - L$ we reproduce the assignment of quantum numbers to the weakly interacting doublets of Sect. 4.1.

6 Concluding remarks

The idea that exceptional structures in mathematics should characterize the fundamental constituents of matter has been with us since the Ancient Greeks first contemplated the Platonic solids⁸. The octonions, the elements of the ultimate division algebra, have been linked to the Standard Model of particle physics starting with the paper of Günaydin and Gürsey [GG] which related them to the coloured quarks. Vigorous attempts to implement them in superstring theory [GNORS, CH] remained inconclusive. When the idea of a finite quantum geometry emerged [DKM, CL] (see also the recent contributions [CCS, BF, L18, CCS18] and references therein) it became natural to look for a role of special algebraic structures in such a context. The application of the exceptional Jordan algebra $\mathcal{J} = J_3^8$ to the SM, put forward in Chapters 4 and 5 is a continuation of [TD] connected with the recently introduced notion of an extended special euclidean Jordan algebra [D-V.T]. It was initiated by the paper [DV] of Michel Dubois-Violette (see also [CDD] where differential calculus and the theory of connections on Jordan algebras and Jordan modules is developed).

We find it remarkable that our version of the Borel - de Siebenthal theory [BdS] applied to the automorphism group F_4 of \mathcal{J} , yields unambiguously the gauge group G_{SM} of the SM and that the relevant irreducible representation $\underline{26}$ of F_4 combines in a single multiplet all quarks' and leptons' doublets and the d-quarks' isotopic singlets with the gauge bosons of the electroweak interactions. On the other hand, the extended observable algebra $\mathcal{H}_{16}(\mathbb{C}) \oplus \mathcal{H}_{16}(\mathbb{C})$ of J_2^8 gives room to the eight primitive idempotents corresponding to the leptons and antileptons of the SM and to the eight 3-dimensional projectors to the internal state space of quarks and antiquarks of the first generation.


In fact, the exceptional Jordan algebra is intimately related to all exceptional Lie groups (Sect. 3.2 - see also [BS, McC]). It will be interesting to reveal the role of the structure group $E_{6(-26)}$ and the conformal group $E_{7(-25)}$

⁸Theaetetus, a contemporary of Plato, gave the first mathematical description of the five regular solids. Plato in the dialog Timaeus (c. 360 B.C.) related the four classical elements, earth, air, water and fire to the four regular solids, cube, octahedron, icosahedron and tetrahedron, respectively, while, according to him "God used the dodecahedron for arranging the constellations on the whole heaven". In the 20th century special structures became part of the excitement with the A-D-E classification that includes the exceptional Lie groups - see [McKay] and references therein.

of \mathcal{J} in the physics of the SM. We intend to return to this problem in future work.

Acknowledgments. I.T. thanks Michel Dubois-Violette for discussions. Both authors thank Ludmil Hadjiivanov for his participation. I.T. acknowledges the hospitality of the IHES (Bures-sur-Yvette) and the NCCR SwissMAP (Geneva) where part of this work was done. S. Drenka's work has been supported by the Bulgarian National Science Fund, DFNI E02/6.

Appendix A. The Fano plane of imaginary octonions ([B])



$$\begin{aligned}
 e_1 &= (0, 0, 1), e_2 = (0, 1, 0) \Rightarrow e_1 e_2 = e_4 = (0, 1, 1) \\
 e_3 &= (1, 0, 0) \Rightarrow e_2 e_3 = e_5 = (1, 1, 0) \\
 e_1 e_5 &= e_6 = (1, 1, 1) \\
 e_4 e_5 &= e_7 = (1, 0, 1).
 \end{aligned}$$

Figure 2.

Projective plane in \mathbb{Z}_2^3 with seven points and seven lines.

The multiplication rules for the seven imaginary quaternionic units can be summarized by

$$e_a e_b = -\delta_{ab} + f_{abc} e_c \quad (\text{A.1})$$

where f_{abc} are fully antisymmetric and

$$f_{124} = f_{235} = f_{346} = f_{561} = f_{672} = f_{713} = 1. \quad (\text{A.2})$$

The relation (A.2) obey the rules

$$f_{ijk} = 1 \Rightarrow f_{i+1j+1k+1} = 1 = f_{2i2j2k} \quad (\text{A.3})$$

where indices are counted mod 7. Eqs.(A.2) can be recovered from anyone of them and the first relation Eqs.(A.3).

We have displayed on Figure 2. the points e_i as non-zero triples of homogeneous coordinates taking values 0 and 1 such that the product $e_i e_j$ (in clockwise order) is obtained by adding the coordinates (a, b, c) , $a, b, c \in \{0, 1\}$, modulo two.

Appendix B. Two bases of $so(8)$ related by the outer automorphism π .

The generators G_{ab} of $so(8)$ are given directly by their action on the octonion units (2.24):

$$G_{ab}e_b = e_a, \quad G_{ab}e_a = -e_b, \quad G_{ab}e_c = 0 \quad \text{for } a \neq c \neq b. \quad (\text{B.1})$$

The action of F_{ab} can also be deduced from definition (2.24) and multiplication rules:

$$\begin{aligned} F_{ab}e_b &= \frac{1}{2}e_a, & F_{ab}e_a &= -\frac{1}{2}e_b, & (a \neq b)F_{ab} &= -F_{ba}. \\ F_{j0}e_{2j} &= \frac{1}{2}e_{4j \bmod 7} (= -F_{0j}e_{2j}) & \text{for } j &= 1, 2, 4, \\ F_{j0}e_{3j} &= \frac{1}{2}e_7, & F_{70}e_7 &= \frac{1}{2}e_{3j}, & F_{07} &= \frac{1}{2}e_j, \\ F_{0j}e_{6j} &= \frac{1}{2}e_{5j}, & F_{j0}e_{5j} &= \frac{1}{2}e_{6j}, & [F_{j0}, F_{0k}] &= F_{jk}. \end{aligned} \quad (\text{B.2})$$

(All indices are counted $\bmod 7$.) From (B.1) and (B.2) we find

$$\begin{aligned} 2F_{0j} &= G_{0j} + G_{2j4j} + G_{3j7} + G_{5j6j} \\ 2F_{03j} &= G_{03j} - G_{J7} - G_{2j5j} + G_{4j6j}, \quad j = 1, 2, 4 \\ 2F_{07} &= G_{07} + G_{13} + G_{26} + G_{45}. \end{aligned} \quad (\text{B.3})$$

In particular, taking the skew symmetry of G_{ab} and the counting $\bmod 7$ into account we can write

$$\begin{aligned} 2F_{02} &= G_{02} - G_{14} + G_{35} - G_{76}, \\ 2F_{04} &= G_{04} + G_{12} - G_{36} - G_{75}, \\ 2F_{03} &= G_{03} - G_{17} - G_{25} + G_{46}, \\ 2F_{06} &= G_{06} + G_{15} - G_{27} - G_{43}, \\ 2F_{05} &= G_{05} - G_{16} + G_{23} - G_{47}. \end{aligned} \quad (\text{B.4})$$

Note that with the abc are ordering $(1, 2, 4, 3, 7, 5, 6)$ The first (positive) indices of G $(2, 3, 5; 1, 3, 7; 1, 2, 4)$ correspond to quaternionic triples: $e_2e_3 = e_5$, $e_1e_3 = e_7$, $e_1e_2 = e_4$. Setting

$$\begin{aligned} G_1 &= \begin{pmatrix} G_{01} \\ G_{24} \\ G_{37} \\ G_{56} \end{pmatrix}, G_2 = \begin{pmatrix} G_{02} \\ G_{14} \\ G_{35} \\ G_{76} \end{pmatrix}, G_4 = \begin{pmatrix} G_{04} \\ G_{12} \\ G_{36} \\ G_{75} \end{pmatrix}, G_3 = \begin{pmatrix} G_{03} \\ G_{17} \\ G_{25} \\ G_{46} \end{pmatrix}, \\ G_7 &= \begin{pmatrix} G_{07} \\ G_{13} \\ G_{26} \\ G_{45} \end{pmatrix}, G_5 = \begin{pmatrix} G_{05} \\ G_{16} \\ G_{23} \\ G_{47} \end{pmatrix}, G_6 = \begin{pmatrix} G_{06} \\ G_{15} \\ G_{27} \\ G_{43} \end{pmatrix}, \end{aligned}$$

and similarly for F_1, \dots, F_6 we find

$$F_a = X_a G_a, \quad a = 1, 2, 4, 3, 7, 5, 6, \text{ with}$$

$$\begin{aligned} X_1 &= X_7 = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}, X_2 = X_5 = \frac{1}{2} \begin{pmatrix} 1 & -1 & 1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & 1 & 1 & 1 \\ -1 & -1 & 1 & 1 \end{pmatrix}, \\ X_4 &= X_6 = \frac{1}{2} \begin{pmatrix} 1 & 1 & -1 & -1 \\ 1 & 1 & 1 & 1 \\ -1 & 1 & 1 & -1 \\ -1 & 1 & -1 & 1 \end{pmatrix}, X_3 = \frac{1}{2} \begin{pmatrix} 1 & -1 & -1 & 1 \\ -1 & 1 & -1 & 1 \\ -1 & -1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}. \end{aligned}$$

They all define involutive transformations:

$$X_k^2 = \mathbb{I}, \quad \det X_k = -1, \quad k = 1, 2, 3, 4. \quad (\text{B.5})$$

and satisfy

$$X_1 X_2 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} = -X_3 X_4, \quad X_1 X_3 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} = -X_2 X_4$$

$$X_1 X_4 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{pmatrix} \Rightarrow X_1 X_2 X_3 X_4 = -\mathbb{I}.$$

$$(X_i X_j = X_j X_i, \quad X_1 X_2 + X_3 X_4 = 0, \quad X_1 X_3 + X_2 X_4 = 0). \quad (\text{B.6})$$

References

- [A] J.F.Adams, *Lectures on Exceptional Lie Groups*, eds. Z. Mahmud, M. Mimira, Univ. Chicago Press, 1996.
- [ABS] M.F. Atiyah, R. Bott, A. Shapiro, Clifford modules, *Topology* **3**, Suppl. 1 (1964) 3-38.
- [A95] R.Ablamowicz, Construction of spinors via Witt decomposition and primitive idempotents: a review, Clifford algebras and spinor structures. Kluwer Academic Publishers, 1995.
- [B] J. Baez, The octonions, *Bull. Amer. Math. Soc.* **39**:2 (2002) 145-205, Errata, *ibid.* **42** (2005) 213; math/0105155v4 [math.RA].
- [B12] J.C. Baez, Division algebras and quantum theory, *Found. Phys.* **42** (2012) 819-855; arXiv:1101.5690v3 [quant-ph].
- [BH] J.C. Baez, J. Huerta, The algebra of grand unified theory, *Bull. Amer. Math. Soc.* **47**:3 (2010) 483-552; arXiv:0904.1556v2 [hep-th].
- [BH09] J.C. Baez, J. Huerta, Division algebras and supersymmetry I, in: *Superstrings, Geometry, Topology, and C*-algebras*, eds. R. Doran, G. Friedman and J. Rosenberg, Proc. Symp. Pure Math. 81, AMS, Providence, 2010, pp. 65-80; arXiv:0909.0551v3 [hep-th].
- [BS] C.H. Barton, A. Sudbery, Magic squares and matrix models of Lie algebras, *Adv. in Math.* **180** (2003) 596-647; math/0203010v2.
- [B61] F. van der Blij, History of the octaves, *Simon Stevin, Wis. en Naturkundig Tijdschrift* 34e Jaargang Aflevering III (Februari 1961) 106-125.
- [BdS] A. Borel, J. de Siebenthal, Les sous-groupes fermés de rang maximum des groupes de Lie clos, *Commentarii Mathematici Helvetici* **23** (1949) 200-221.
- [BF] L. Boyle, S. Farnsworth, A new algebraic standard model of particle physics, arXiv:1604.00847 [hep-th]; -. Rethinking Connes' approach to the standard model of particle physics via non-commutative geometry, *New J. Phys.* **17** (2015) 023021; arXiv:1408.5367 [hep-th].

- [CDD] A. Carotenuto, L. Dabrowski, M. Dubois-Violette, Differential calculus on Jordan algebras and Jordan modules, arXiv:1803.08373 [math.QA].
- [Ca] Élie Cartan, Le principe de dualité et la théorie des groupes simple et semisimple, *Bull.Sci.Math.* **49** (1925) 361-374.
- [CGKK] S. Catto, Y. Gürcan, A. Khalfan, L. Kurt, Unifying ancient and modern geometries through octonions, *J. Phys. Conference Series* **670** (2016) 012016.
- [CCS] A.H. Chamseddine, A. Connes, W.D. van Suijlekom, Grand unification in the spectral Pati-Salam model, *JHEP* **1511** (2015) 011; arXiv:1507.08161 [hep-ph].
- [CCS18] A.H. Chamseddine, A. Connes, W.D. van Suijlekom, Entropy and the spectral action, arXiv:1809.02944 [hep-th].
- [CL] A. Connes and J. Lott, Particle models and noncommutative geometry. *Nucl. Phys. Proc. Suppl.*, **B 18** (1990) 29–47 .
- [CS] J.H. Conway, D.A.Smith, *On Quaternions and Octonions: Their Geometry, Arithmetic and Symmetry* A.K. Peters (2003) 1-143. J.C. Baez, *Bull. (New Series) Amer. Math. Soc.* **42**(2005) 229-243
- [CH] E. Corrigan, T.J. Hollowood, The exceptional Jordan algebra and the superstring, *Commun. Math. Phys.* **122** (1989) 393-410.
- [Di] L.E. Dickson, On quaternions and their generalization and the history of the eight square theorem, *Annals of Math.* **20** (1919) 155-171.
- [D04] G.M. Dixon, Division algebras: family replication, *J.Math. Phys.* **45** (2004) 3878.
- [D10] G.M. Dixon, Division algebras; spinors; idempotents; the algebraic structure of reality, arXiv:1012.1304 [hep.th].
- [D14] G.M. Dixon, Seeable matter; unseeable antimatter, *Comment. Math. Univ. Carolin.* **55**:3 (2014) 381-386; arXiv:1407.4818 [physics.gen-ph].
- [DV] M. Dubois-Violette, Exceptional quantum geometry and particle physics, *Nucl. Phys. B* **912** (2016) 426-444; arXiv:1604.01247 [hep-th].

- [D-V.T] M. Dubois-Violette, I. Todorov. Exceptional quantum geometry and particle physics II, arXiv:1808.08110 [hep-th].
- [DKM] M. Dubois-Violette, R. Kerner, and J. Madore, Gauge bosons in a non-commutative geometry. *Phys. Lett.*, **B 217** (1989) 485–488.
- [FK] J. Faraut, A. Koranyi, *Analysis on Symmetric Cones*, Oxford Univ. Press, 1994 (xii + 382 pp.); K.I. Gross, Book Review *Bull. AMS* **35** (1998) 77-86.
- [F] H. Freudenthal, Lie groups in the foundation of geometry, *Adv. Math.* **1** (1964) 145-190.
- [F16] C. Furey, Standard model physics from an algebra? arXiv:1611.09182 [hep-th].
- [F18] C. Furey, $SU(3)_c \times SU(2)_L \times U(1)_Y (\times U(1)_X)$ as a symmetry of the division algebra ladder operators, *Eur. Phys. J.* **C78** (2018) 375, 12 pages; arXiv:1806.00612 [hep-th].
- [GNORS] P. Goddard, W. Nahm, D.I. Olive, H. Ruegg, A. Schwimmer, Fermions and octonions, *Commun. Math. Phys.* **112** (1987) 385-408.
- [GG] M. Günaydin, F. Gürsey, Quark structure and octonions, *J. Math. Phys.* **14**:11 (1973) 1651-1667.
- [GPR] M. Günaydin, C. Pirron, H. Ruegg, Moufang plane and octonionic quantum mechanics, *Commun. Math. Phys.* **61** (1978) 69-85.
- [G] F. Gürsey, Octonionic structures in particle physics, *Group Theoretical Methods in Physics*, Lecture Notes in Physics **94**, Springer, 1979, pp. 508-521; F. Gürsey, C.-H. Tze, *The Role of Division, Jordan and Related Algebras in Particle Physics*, World Scientific, 1996 (460 pages).
- [JvNW] P. Jordan, J. v. Neumann, E. Wigner, On an algebraic generalization of the quantum mechanical formalism, *Ann. of Math. (2)* **36** (1934) 29-64.
- [K] M. Koecher, *The Minnesota Notes on Jordan Algebras and Their Applications*, Lecture Notes in Math. **1710** ed. by A. Krieg, S. Walcher, Springer, Berlin et al. 1999.

- [L18] F. Lizzi, Noncommutative geometry and particle physics, *Proc. of Sci. Corfu Summer Institute 2017*; arXiv:1805.00411 [hep-th].
- [L] P. Loumesto, *Clifford Algebras and Spinors*, second edition, London Math. Soc. Lecture Notes Series **286**, Cambridge Univ. Press, 2001.
- [McC] K. McCrimmon, *A Taste of Jordan Algebras*, Springer, 2004 (xxv + 564 pages); Jordan algebras and their applications, *Bull. Amer. Math. Soc.* **84**:4 (1978) 612-627,
- [McKay] John McKay, A rapid introduction to ADE theory, <http://math.ucr.edu/home/baez/ADE.html>
- [M] G. Meng, Euclidean Jordan algebras, hidden actions, and J-Kepler problems, *J. Math. Phys.* **52** (2011) 112104; arXiv:0911.2977v3 [math-ph].
- [M89] Shingo Murakami, Differential Geometry and Topology. Exceptional Simple Lie Groups and Related Topics in Recent Differential Geometry, in: *Differential Geometry and Topology*, (Jiang B., Peng Ch.-K., Hou Z., eds.), Lecture Notes in Math. **1369**, Springer, 1989, pp. 183-221.
- [P] J. Palmkvist, A generalization of the Kantor-Koecher-Tits construction, *J. Gen. Lie Theory Appl.* **2** (2008) 226; arXiv:1308.3761 [math.RA].
- [R08] G. Roos, Exceptional symmetric domains, in: Bruce Guilling et al. (eds.), *Symmetries in Complex Analysis*, AMS, Contemporary Mathematics **468** (2008) 157-189; arXiv:0801.4076 [math.CV].
- [S] R.D. Schafer, Structure and representations of nonassociative algebras, *Bull. Amer. Math. Soc.* **61**:6 (1955) 469-484; -, *An Introduction to Nonassociative Algebras*, Dover, N.Y. 1995.
- [S18] O.C. Stoica, The standard model algebra - leptons, quarks, and gauge from the complex Clifford algebra Cl_6 , *Adv. Appl. Clifford Algebras* **28** (2018) 52; arXiv:1702.04336v3 [hep-th].
- [T] I. Todorov, Clifford algebras and spinors, *Bulg. J. Phys.* **58**:1 (2011) 3-28; arXiv:1106.3197 [math-ph].

- [TD] I. Todorov, M. Dubois-Violette, Deducing the symmetry of the standard model from the automorphism and structure groups of the exceptional Jordan algebra, *Int. J. Mod. Phys. A* **33** (2018) 1850118; arXiv:1806.09450 [hep-th].
- [TD17] I. Todorov, S. Drenska, Composition algebras, exceptional Jordan algebra and related groups, *JGSP* **46** (2017) 59-93.
- [T85/16] P.K. Townsend, Jordan formulation of Quantum mechanics: a review, *DAMTP* (1985)-0263; tt arXiv:1612.09228 [quant-ph].
- [W] E. Witten, Quest for unification, tt arXiv:0207124 [quant-ph].
- [WWW] J.C. Wick, A.S. Wightman, E.P. Wigner, The intrinsic parity of elementary particles, *Phys. Rev.* **88** (1952) 101.
- [Y] I. Yokota, Exceptional Lie groups, arXiv:0902.0431 [math.DG] (204 pages).
- [Z] E. Zelmanov, On Isayah Kantor (1936-2006), *Journal of Generalized Lie Theory and Applications* **2:3** (2008) 111.