

ASYMPTOTIC CONES AND BOUNDARIES OF CAT(0) GROUPS

CURTIS KENT AND RUSSELL RICKS

ABSTRACT. It is well known that the Euclidean cone over the Tits boundary of a proper cocompact CAT(0) space isometrically embeds into every asymptotic cone of the space. We explore the relationships between the asymptotic cones of a CAT(0) space and its boundary under both the standard visual (i.e. cone) topology and the Tits metric. We show that the set of asymptotic cones of a proper cocompact CAT(0) space admits canonical connecting maps under which the direct limit is isometric to the Euclidean cone on the Tits boundary. The resulting projection from any asymptotic cone to the Tits boundary is determined by the visual topology; on the other hand, the visual topology can be recovered from the connecting maps between asymptotic cones. We also demonstrate how maps between asymptotic cones induce maps between Tits boundaries.

1. INTRODUCTION

Asymptotic cones and compactifications are two very different approaches to studying the coarse geometry of groups and spaces. In general, asymptotic cones of a metric space X will not be locally compact and not preserve the local geometry of X while compactifications often can be endowed with metrics topologically compatible with the local geometry of X . However, asymptotic cones do reflect well the coarse Euclidean aspects of a space; and for CAT(0) spaces, the asymptotic cones are complete CAT(0) spaces.

Intuitively, an asymptotic cone of a metric space (X, dist) is the limit of the metric spaces $(X, \text{dist}/d_n)$ where dist/d_n is the metric on X scaled by $1/d_n$. Asymptotic cones have the desirable property that quasi-isometric spaces have bi-Lipschitz asymptotic cones. However, an ultrafilter is required to guarantee the existence of the limit. As a consequence, they also have the drawback that a metric space can have uncountably many distinct asymptotic cones depending on the choice of ultrafilter and scaling sequence used in the construction.

There are many connections between the topological structure of the asymptotic cones of a finitely generated group and its combinatorial and algorithmic properties. For example, a group has polynomial growth if and only if it is virtually nilpotent, if and only if every asymptotic cone is locally compact [7, 10]. Also, a finitely generated group is finitely presented and has polynomial Dehn function if all of its asymptotic cones are simply connected [8].

Another approach to studying the coarse geometry of groups and spaces is via boundaries. The visual boundary is one natural compactification for proper CAT(0) and hyperbolic spaces. The visual boundary is the set of large-scale directions and is an isometry invariant of the space. In fact, for hyperbolic groups the visual boundary is a quasi-isometry invariant. However, Croke and Kleiner proved that a group can act geometrically on two CAT(0) spaces with non-homeomorphic visual boundaries [5]. Thus in the CAT(0) setting the visual boundary is not a quasi-isometry invariant.

The boundary of a CAT(0) space can also be endowed with a metric called the Tits metric which reflects the Euclidean structure of the CAT(0) space. With the Tits metric, the set of large-scale directions is no longer a compactification but in some sense better encodes the coarse Euclidean geometry of the space.

The first author was supported by Simons Foundation Collaboration Grant 587001. The second author was partially supported by NSF RTG 1045119.

We will illustrate how the visual boundary, the Tits boundary, and the asymptotic cones of proper CAT(0) spaces relate. It is well known that the Euclidean cone over the Tits boundary admits a canonical isometric embedding into every asymptotic cone. We will show that the set of asymptotic cones of a proper CAT(0) space determine the Tits boundary of the space.

Theorem A (Theorem 4.18). *Let X be a proper cocompact CAT(0) space. The direct limit of asymptotic cones of X induced by the geodesic retraction on X is isometric to the Euclidean cone on the Tits boundary of X . Moreover, the resulting projection maps onto the direct limit are determined by the visual topology on the boundary.*

Thus the set of asymptotic cones of X for a fixed ultrafilter determines the Tits boundary of X . The connecting maps also give rise to an inverse limit, which is related to the countable ultraproduct of the Tits boundary.

Theorem B (Theorem 4.20). *Let X be a proper cocompact CAT(0) space. The inverse limit of asymptotic cones of X induced by the geodesic retraction on X has an inverse limit metric and with this metric there exists a canonical isometric embedding of the ultraproduct of the Euclidean cone on the Tits boundary of X into the inverse limit.*

We leave it as an open question whether or not this embedding is surjective.

In Section 5, we demonstrate how the asymptotic cones determine the visual topology on the boundary.

Theorem C (Theorem 5.2). *The visual topology on the boundary of a proper cocompact CAT(0) space is determined by the geodesic retraction maps between asymptotic cones.*

In Section 6, the direct limit is used to define continuous maps between Tits boundaries of quasi-isometric CAT(0) spaces, which when restricted to Morse geodesics gives a bijection. This gives an alternate proof that a CAT(0) group has a cut-point in some asymptotic cone if and only if it has cut-points in every asymptotic cone if and only if it has a periodic rank one element.

2. VISUAL BOUNDARY

A CAT(0) space is a uniquely geodesic metric space such that every geodesic triangle $\Delta(x, y, z)$ is thinner than the corresponding comparison triangle $\overline{\Delta}(x, y, z)$ in Euclidean \mathbb{R}^2 . This generalizes the property of nonpositive curvature from Riemannian manifolds to the metric setting. We refer the reader to [4] for a more complete account.

The visual boundary of a complete CAT(0) space can be considered as either the set of equivalence classes of geodesic rays (equivalent if they are asymptotic) or the set of based geodesic rays. Here we will use the latter.

For a CAT(0) space (X, dist) and $x, y \in X$, we will use $[x, y]$ to denote the unique unparameterized geodesic from x to y , and $B_\epsilon^{\text{dist}}(x)$ or $B_\epsilon(x)$ to denote the open metric ball of radius ϵ about x .

Definition 2.1 (Visual compactification and boundary). Let X be a CAT(0) metric space. For a fixed $x_0 \in X$, let ∂X be the set of geodesic rays with basepoint x_0 .

For $\alpha \in \partial X$ and $\epsilon, R > 0$, let

$$V(\alpha, R, \epsilon) = \{\beta \in \partial X \mid \text{dist}(\alpha(R), \text{im}(\beta)) < \epsilon\} \cup \{x \in X \mid [x_0, x] \cap B_\epsilon(\alpha(R)) \neq \emptyset\}.$$

If X is a proper CAT(0) metric space (*proper* meaning closed balls are compact), then

$$\{V(\alpha, R, \epsilon) \mid \alpha \in \partial X, \epsilon, R > 0\} \cup \{B_\epsilon(x) \mid x \in X, \epsilon > 0\}$$

is a basis for a compact topology on $\overline{X} = X \cup \partial X$, which we will call the *visual compactification* of X . Notice that ∂X is a closed subspace. The *visual boundary* of X is the set ∂X endowed

with the subspace topology from \overline{X} and will be denoted by $\partial_\infty X$. The visual boundary has basis $\{U(\alpha, R, \epsilon) \mid \alpha \in \partial X, \epsilon, R > 0\}$ where $U(\alpha, R, \epsilon) = V(\alpha, R, \epsilon) \cap \partial X$.

Definition 2.2 (Limit set). Let X be a CAT(0) space and $A \subset X$. The *limit set* of A , denoted $\Lambda(A)$, is the closure of A in the visual compactification intersected with the visual boundary, i.e. $\Lambda(A) = \text{cl}_{\overline{X}}(A) \cap \partial_\infty X$ where $\text{cl}_{\overline{X}}$ is the topological closure operator in \overline{X} .

The visual boundary can be endowed with several natural metrics. Here we will use the following metric since it relates well to the metric on asymptotic cones. Fix $C > 0$. For two geodesics $\alpha, \beta : ([0, \infty), 0) \rightarrow (X, x_0)$ into a CAT(0) space X , let

$$\text{dist}_C(\alpha, \beta) = \frac{1}{\sup\{t \mid \text{dist}(\alpha(t), \beta(t)) \leq C\}}.$$

Lemma 2.3. *The function dist_C is a metric on ∂X .*

Proof. Notice that $\text{dist}_C(\alpha, \beta) \leq \frac{1}{n}$ implies that $\text{dist}(\alpha(t), \beta(t)) \leq \frac{Ct}{n}$ for all $t \leq n$ by the CAT(0) condition. Hence $\text{dist}_C(\alpha, \beta) = 0$ if and only if $\alpha = \beta$. Clearly dist_C is symmetric and we are only left to show the triangle inequality holds for dist_C .

Suppose that $\text{dist}_C(\alpha, \beta) = \frac{1}{n}$ and $\text{dist}_C(\beta, \gamma) = \frac{1}{m}$ for some $m, n \in \mathbb{R}$. For all $t \leq \min\{m, n\}$, we have $\text{dist}(\alpha(t), \gamma(t)) \leq Ct(\frac{1}{n} + \frac{1}{m})$ by the triangle inequality for the metric on X . Hence $\text{dist}\left(\alpha\left(\frac{mn}{m+n}\right), \gamma\left(\frac{mn}{m+n}\right)\right) \leq C$ which implies that

$$\text{dist}_C(\alpha, \gamma) \leq \frac{m+n}{mn} = \frac{1}{m} + \frac{1}{n} = \text{dist}_C(\alpha, \beta) + \text{dist}_C(\beta, \gamma). \quad \square$$

When the metric is understood, we will simply write $B_\epsilon(\alpha)$ for $B_\epsilon^{\text{dist}_C}(\alpha)$. Notice that $B_{\epsilon/(CR)}^{\text{dist}_C}(\alpha) \subset U(\alpha, R, \epsilon)$ as long as $0 < \epsilon \leq C$, and $U(\alpha, 1/\epsilon + C/2, C/2) \subset B_\epsilon^{\text{dist}_C}(\alpha)$, which proves the following observation.

Lemma 2.4. *The metric dist_C induces the visual topology on ∂X .*

Definition 2.5 (Tits/Angle boundaries). For points x, y, z in a CAT(0) space X , we will let $\overline{\angle}_x(y, z)$ denote the comparison angle at x between y and z . If $p_y : [0, a] \rightarrow X$ and $p_z : [0, b] \rightarrow X$ are the unique geodesics in X from x to y and from x to z respectively, then the *angle between y and z at x* is $\angle_x(y, z) = \lim_{t \rightarrow 0} \overline{\angle}_x(p_y(t), p_z(t))$.

For $\alpha, \beta \in \partial X$, let

$$\begin{aligned} \angle(\alpha, \beta) &= \lim_{t \rightarrow \infty} \overline{\angle}_{x_0}(\alpha(t), \beta(t)) = \sup\{\overline{\angle}_{x_0}(\alpha(t), \beta(t')) \mid t, t' > 0\} \text{ and} \\ \angle_{x_0}(\alpha, \beta) &= \lim_{t \rightarrow 0} \overline{\angle}_{x_0}(\alpha(t), \beta(t)) = \inf\{\overline{\angle}_{x_0}(\alpha(t), \beta(t')) \mid t, t' > 0\}. \end{aligned}$$

It is an exercise to show that $\angle(\cdot, \cdot)$ defines a locally geodesic metric on ∂X , which is called the *angle metric*. We will denote ∂X with this metric by $\partial_\angle X$.

The path metric induced by $\angle(\cdot, \cdot)$ is called the Tits metric and is denoted by $\text{dist}_T(\cdot, \cdot)$. The *Tits boundary of X* is ∂X with this metric and will be denoted by $\partial_T X$. Note that dist_T is an extended metric in the sense that it maps into $[0, \infty]$. The Tits distance between any two points in distinct path components of the angle boundary is infinity. We refer the interested reader to [4, Chapter I. 1, II.9] for complete details.

Proofs of the following standard lemmas can be found in [4, Proposition II.9.8] and [4, Proposition II.9.9].

Lemma 2.6. *Let X be a CAT(0) space. For $\alpha, \beta \in \partial X$,*

$$2 \sin\left(\frac{\angle(\alpha, \beta)}{2}\right) = \lim_{t \rightarrow \infty} \frac{\text{dist}(\alpha(t), \beta(t))}{t}.$$

Lemma 2.7 (Flat Sector Theorem). *Let X be a CAT(0) space. If $\alpha, \beta \in \partial X$ such that $\angle(\alpha, \beta) < \pi$ and $\angle(\alpha, \beta) = \angle_{x_0}(\alpha, \beta)$, then the convex hull of the geodesic rays α and β is isometric to a sector in the Euclidean plane bounded by two rays which meet at an angle $\angle(\alpha, \beta)$.*

3. METRIC LIMITS

As we will only be concerned about limits of spaces indexed by a totally ordered sets, we will present the definitions for inverse and direct limits in terms of spaces indexed by a totally ordered set.

Let (J, \geq) be a totally ordered set. A *system of metric spaces* is a collection of metric spaces $\{(X_j, \text{dist}_j) \mid j \in J\}$ together with a collection of non-expansive maps $\{f_{i,j} : X_i \rightarrow X_j \mid \text{for all } i \geq j\}$, called the *connecting maps*, with the property that $f_{i,k} = f_{i,j} \circ f_{j,k}$ for all $i \geq j \geq k$.

Definition 3.1 (Direct limits of metric spaces). Let (X_j, dist_j) be a metric space for each $j \in J$. Write $\bigsqcup_{j \in J} X_j$ for the disjoint union of X_j . Define a pseudometric on $\bigsqcup_{j \in J} X_j$ by $\underline{\text{dist}}(x, y) = \inf\{\text{dist}_k(f_{i,k}(x), f_{j,k}(y)) \mid i, j \geq k\}$ for $x \in X_i$ and $y \in X_j$. We will say that $x, y \in \bigsqcup_{j \in J} X_j$ are equivalent, denoted $x \sim y$, if $\underline{\text{dist}}(x, y) = 0$. The *direct limit* of the system of metric spaces $\{X_i, f_{i,j}\}$ is the set $\bigsqcup_{j \in J} X_j / \sim$ of equivalence classes, together with the metric $\underline{\text{dist}}$. We will denote the direct limit of $\{X_i, f_{i,j}\}$ by $\varinjlim \{X_i, f_{i,j}\}$ or when convenient $\varinjlim X_i$.

It is immediate that the canonical induced maps $f_i : X_i \rightarrow \varinjlim \{X_i, f_{i,j}\}$ are all non-expansive.

Definition 3.2 (Inverse limits of metric spaces). Recall that an extended metric space is a set with a distance function that potentially takes on infinite values but satisfies the rest of the properties of a metric.

If (X_j, dist_j) is a metric space for each $j \in J$, then the metric product of X_j , written $\prod_{j \in J} X_j$, is the extended metric space consisting of functions $x : J \rightarrow \bigcup_{j \in J} X_j$ such that $x(j) \in X_j$ for all $j \in J$, with metric $\overline{\text{dist}}(x, y) = \sup_{j \in J} \text{dist}_j(x(j), y(j))$. The *inverse limit* of a system of metric spaces is the metric subspace of $\prod_{j \in J} X_j$ which consists of those functions x for which $x(j) = f_{i,j}(x(i))$ for every i and j such that $i \geq j$. We will denote the inverse limit by $\varprojlim \{X_i, f_{i,j}\}$ or when convenient by $\varprojlim X_i$.

The inverse limit of complete metric spaces is complete; the direct limit need not be.

Remark 3.3. We will show that the inverse limit of asymptotic cones is a metric space, with all points finite distance because the connecting maps all preserve distance to the basepoint.

Remark 3.4. While the topological inverse/direct limit of a system of metric spaces is also a topological space, the metric on the metric inverse/direct limit need not always induce the topology of the topological inverse/direct limit.

Example 3.5. Fix a basepoint p in a complete CAT(0) space X . The metric inverse limit of spheres $S_r(p)$ (with connecting maps coming from geodesic projection, and metric $\overline{\text{Z}}_p$) is the Tits boundary $\partial_T X$. On the other hand, the topological inverse limit of this system is the visual boundary $\partial_\infty X$. Meanwhile, metric direct limit of this system of spheres is the space of

directions¹, $\Sigma_p X$, at p . Its completion (if X is proper and geodesically complete it is already complete) is called the link at p .

Now consider the system of closed metric balls $\overline{B}_r(p)$ (with connecting maps all geodesic projection). The metric inverse limit consists an isometric copy of X , along with the points of ∂X , all infinite distance from everything else in the space. The topological inverse limit, on the other hand, is the visual compactification $\overline{X} = X \cup \partial_\infty X$ of X [4, p. 263]. Meanwhile, the metric direct limit is a single point.

If we use rescaled copies $\frac{1}{r}X$ of X (using geodesic projection to rescale distances to p), the metric inverse limit is $\text{Cone}(\partial_T X)$, the Euclidean cone over the Tits boundary; the metric direct limit is the tangent cone $\text{Cone}(\Sigma_p X)$ at p , the Euclidean cone over the space of directions. The system of asymptotic cones we will consider resembles this last system, but the details are more involved.

4. DIRECT LIMITS AND INVERSE LIMITS OF ASYMPTOTIC CONES

Ultrafilters and Asymptotic cones.

Definition 4.1. A (*non-principal*) *ultrafilter* ω on a set S is a finitely additive probability measure on the power set of S with values in $\{0, 1\}$ such that $\omega(A) = 0$ for all finite subsets $A \subset S$. We will say that $A \subset S$ is ω -*large* if $\omega(A) = 1$. A property P holds ω -*almost surely* if it holds for some ω -large subset of S .

For $\{a_s \mid s \in S\} \subset X$, a subset of a topological space X indexed by S , we will say that the *ultralimit* of a_s is x , written $\lim^\omega a_s = x$, if for every open neighborhood U of x the set $\{s \mid a_s \in U\}$ is ω -large. It is an exercise to show that every S -indexed subset of a compact space has a unique ultralimit and that ultralimits satisfy the standard properties of limits. If $\{a_s\} \subset X$ is S -indexed but has no ultralimit in X , we will say a_s is ω -*divergent*.

Definition 4.2 (Ultraproducts). Let (X_n, dist_n) be a sequence of metric spaces, ω an ultrafilter on \mathbb{N} , and $e = (e_n) \in \prod_{n=1}^{\infty} X_n$. The *ultraproduct* of (X_n, dist_n) is

$$\prod_e^\omega X_n = \left\{ (x_n) \in \prod_{n=1}^{\infty} X_n \mid \text{for each } (x_n), \text{dist}_n(x_n, e_n) \text{ is uniformly bounded} \right\} / \sim$$

where $(x_n) \sim (y_n)$ if $\lim^\omega \text{dist}_n(x_n, y_n) = 0$. The ultraproduct has metric

$$\text{dist}((x_n), (y_n)) = \lim^\omega \text{dist}_n(x_n, y_n).$$

In general, the ultraproduct depends on both e and ω . The sequence e will be called the *observation sequence* for the ultraproduct.

We will use the simplified notation $\partial_\angle^\omega X = \prod_e^\omega \partial_\angle X$ and $\partial_T^\omega X = \prod_e^\omega \partial_T X$. Since $\partial_\angle X$ is bounded, $\partial_\angle^\omega X$ is independent of the chosen basepoint but in the case of $\partial_T^\omega X$, there is an implied but unspecified choice of basepoint.

Let ω be an ultrafilter on \mathbb{N} and $d = (d_n)$, an ω -divergent sequence of positive real numbers (called a *scaling sequence*). An *asymptotic cone* of X is $\prod_e^\omega (X, \text{dist}/d_n)$ and will be denoted by $\text{Con}^\omega(X, e, d)$.

Definition 4.3 (Euclidean cones). If X is a metric space, let $\text{Cone}(X) = (\mathbb{R}^+ \times X) / \sim$ where $(0, x) \sim (0, x')$ for all $x, x' \in X$. When convenient, we will denote the equivalence class of (t, x) in $\text{Cone}(X)$ by tx .

¹Indeed, $\Sigma_p X$ is formed by taking the set of germs of geodesic segments based at p , with pseudometric \angle_p . Then $\Sigma_p X$ is the metric space of equivalence classes, where geodesic germs are considered equivalent if $\angle_p = 0$.

We can endow $\text{Cone}(X)$ with a metric by

$$\text{dist}^2(tx, t'x') = t^2 + (t')^2 - 2tt' \cos\left(\max\{\pi, \text{dist}(x, x')\}\right).$$

Write $(\text{Cone}(X))^\omega$ for $\prod_e^\omega \text{Cone}(X)$ with $e = (0x)$.

Notice that if X is a complete CAT(0) space, then $\text{Con}^\omega(X, e, d)$, $\text{Cone}(\partial_\perp X)$, $\text{Cone}(\partial_T X)$, $\text{Cone}(\partial_\perp^\omega X)$, $\text{Cone}(\partial_T^\omega X)$, $(\text{Cone}(\partial_\perp X))^\omega$, and $(\text{Cone}(\partial_T X))^\omega$ are all complete CAT(0) spaces.

Proposition 4.4. *Let X be a complete CAT(0) space. The identity map from $\partial_\perp X$ to $\partial_T X$ induces an isometry from $\text{Cone}(\partial_\perp X)$ to $\text{Cone}(\partial_T X)$. The natural map from $\text{Cone}(\partial_\perp^\omega X)$ to $(\text{Cone}(\partial_\perp X))^\omega$ is an isometry. As well, $\partial_T^\omega X$ is homeomorphic to a path component of $\partial_\perp^\omega X$.*

Hence we will often identify $\text{Cone}(\partial_\perp X)$ with $\text{Cone}(\partial_T X)$.

Proof. The angle boundary, $\partial_\perp X$, is a CAT(1) space, see [4, Theorem II.9.13]. Hence the identity map from $\partial_\perp X$ to $\partial_T X$ is an isometry when restricted to an open ball of radius π . It is then immediate that $\text{Cone}(\partial_\perp X)$ is canonically isometric to $\text{Cone}(\partial_T X)$.

Let $F : \text{Cone}(\partial_\perp^\omega X) \rightarrow (\text{Cone}(\partial_\perp X))^\omega$ by $F(t, (\alpha_n)) = ((t, \alpha_n))$. It is a straightforward exercise to verify that F is an isometry.

Recall that the ultraproduct $\partial_T^\omega X$ depends on a fixed observation sequence $e = (e_n)$ in $\partial_T X$ and is the set of sequences of geodesic rays (α_n) such that $\text{dist}_T(\alpha_n, e_n)$ is uniformly bounded ω -almost surely. Thus for $(\alpha_n), (\beta_n) \in \partial_T^\omega X$, there is some M such that $\text{dist}_T(\alpha_n, \beta_n) < M$ ω -almost surely and the ω -limit of the geodesics in $\partial_T X$ from α_n to β_n gives a path from (α_n) to (β_n) in $\partial_\perp^\omega X$. Therefore the identity map from $\partial_T^\omega X$ to $\partial_\perp^\omega X$, takes $\partial_T^\omega X$ into a path component of $\partial_\perp^\omega X$. Suppose that (γ_n) is any element of $\partial_\perp^\omega X$ contained in the same path component as e . Since $\partial_\perp^\omega X$ is locally geodesic, there exists a rectifiable path from (γ_n) to e . Thus $\text{dist}_T(\gamma_n, e_n)$ is uniformly bounded ω -almost surely by the length of this rectifiable path, which implies that $(\gamma_n) \in \partial_T^\omega X$. Therefore $\partial_T^\omega X$ is a path component of $\partial_\perp^\omega X$. \square

Corollary 4.5. *For a complete CAT(0) space X the following are equivalent.*

- (i) *The natural embedding of $\text{Cone}(\partial_T^\omega X)$ into $\text{Cone}(\partial_\perp^\omega X)$ is surjective.*
- (ii) *The diameter of $\partial_T X$ is finite.*

Definition 4.6. Fix $x_0 \in X$. For a fixed scaling sequence $d = (d_n)$ and a non-principal ultrafilter ω , we will define $\Psi_d^\omega : \text{Cone}(\partial_\perp^\omega X) \rightarrow \text{Con}^\omega(X, (x_0), d)$ by

$$\Psi_d^\omega(t, (\alpha_n)) = (\alpha_n(td_n)).$$

Similarly we define $\Psi_d : \text{Cone}(\partial_\perp X) \rightarrow \text{Con}^\omega(X, (x_0), d)$ by

$$\Psi_d(t, \alpha) = (\alpha(td_n)).$$

Notice that for a constant sequence of geodesic, we have $\Psi_d^\omega(t, (\alpha)) = \Psi_d(t, \alpha)$. Thus when convenient, we will identify $\text{Cone}(\partial_\perp X)$ with its canonical diagonal embedding in $\text{Cone}(\partial_\perp^\omega X)$ and consider Ψ_d as a restriction of Ψ_d^ω .

For fixed $(\alpha_n) \in \partial_\perp^\omega X$ and $\alpha \in \partial_\perp X$, the maps $t \mapsto \Psi_d^\omega(t, (\alpha_n))$ and $t \mapsto \Psi_d(t, \alpha)$ are geodesics rays in $\text{Con}^\omega(X, (x_0), d)$ based at (x_0) . Thus we have the following induced maps of boundaries: $\overline{\Psi}_d^\omega : \partial_\perp^\omega X \rightarrow \partial_\perp(\text{Con}^\omega(X, (x_0), d))$ by

$$\overline{\Psi}_d^\omega(\alpha_n)(t) = \Psi_d^\omega(t, (\alpha_n))$$

and $\overline{\Psi}_d : \partial_T X \rightarrow \partial_T(\text{Con}^\omega(X, (x_0), d))$ by

$$\overline{\Psi}_d(\alpha)(t) = \Psi_d(t, \alpha).$$

Recall that a metric space X is *cocompact* if there exists a compact set C such that the translates of C by the full isometry group of X cover X .

Proposition 4.7. *The maps Ψ_d^ω and $\overline{\Psi}_d^\omega$ are non-expansive. If X is proper and cocompact, then Ψ_d^ω and $\overline{\Psi}_d^\omega$ are surjective.*

Proof. By construction of the Euclidean cone, the natural map $\psi : \text{Cone}(\partial_\perp X) \rightarrow X$ given by $\psi(t, \alpha) = \alpha(t)$ is 1-Lipschitz. It follows that Ψ_d^ω is 1-Lipschitz.

Let $(\alpha_n), (\beta_n) \in \partial_\perp^\omega X$. For any $s > 0$,

$$\begin{aligned} \text{dist}((\alpha_n), (\beta_n)) &= \lim_n^\omega \angle(\alpha_n, \beta_n) = \lim_n^\omega \lim_{t \rightarrow \infty} \overline{Z}_{x_0}(\alpha_n(t), \beta_n(t)) \\ &\geq \lim_n^\omega \overline{Z}_{x_0}(\alpha_n(sd_n), \beta_n(sd_n)) \\ &= \overline{Z}_{(x_0)}(\overline{\Psi}_d^\omega(\alpha_n)(s), \overline{\Psi}_d^\omega(\beta_n)(s)). \end{aligned}$$

Since this holds for any $s > 0$,

$$\text{dist}((\alpha_n), (\beta_n)) \geq \lim_{s \rightarrow \infty} \overline{Z}_{(x_0)}(\overline{\Psi}_d^\omega(\alpha_n)(s), \overline{\Psi}_d^\omega(\beta_n)(s)) = \text{dist}(\overline{\Psi}_d^\omega(\alpha_n), \overline{\Psi}_d^\omega(\beta_n)).$$

Thus $\overline{\Psi}_d^\omega$ is 1-Lipschitz.

If X is proper and admits a cocompact action, by [6] there is some constant K (depending only on X and the action) such that for every $(x_n) \in \text{Con}^\omega(X, (x_0), d)$ there exists a sequence of geodesic rays α_n such that $\alpha_n(0) = x_0$ and $\text{dist}(\alpha_n(t_n d_n), x_n) \leq K$ in X , where t_n converges ω -almost surely to $t = \text{dist}((x_0), (x_n))$ in $\text{Con}^\omega(X, (x_0), d)$. Thus $\Psi_d^\omega(t, (\alpha_n)) = (x_n)$, i.e. Ψ_d^ω is surjective.

Showing that $\overline{\Psi}_d^\omega$ is surjective requires a bit more work. Fix $\tilde{\alpha}$ a geodesic in $\text{Con}^\omega(X, (x_0), d)$ based at (x_0) . Since Ψ_d^ω is surjective, there exist $(\alpha_n^k) \in \partial_\perp^\omega X$ such that $\Psi_d^\omega(k, (\alpha_n^k)) = \tilde{\alpha}(k)$. For $k \in \mathbb{N}$, fix a representative (x_n^k) of $\tilde{\alpha}(k)$. Let

$$A_i = \left\{ n \mid \text{dist}(\alpha_n^j(jd_n), x_n^j) < \frac{d_n}{i} \text{ for all } j \leq i \right\}.$$

Notice that A_i is ω -large for each i and forms a nested sequence. Let $D_n = \{i \leq n \mid n \in A_i\}$ and $m_n = \max D_n$ if $D_n \neq \emptyset$ and 1 otherwise. Then m_n diverges ω -almost surely and

$$\text{dist}(\alpha_n^{m_n}(m_n d_n), x_n^{m_n}) < \frac{d_n}{m_n}.$$

Since m_n diverges ω -almost surely, this together with the CAT(0) condition shows that for ω -almost all n

$$\text{dist}(\alpha_n^{m_n}(kd_n), x_n^k) < \frac{d_n}{l},$$

for any fixed $k, l \in \mathbb{N}$.

Thus $\overline{\Psi}_d^\omega(\alpha_n^{m_n}) = \tilde{\alpha}$ and $\overline{\Psi}_d^\omega$ is surjective. \square

Lemma 4.8. *For fixed $\alpha, \beta \in \partial X$,*

$$\angle(\overline{\Psi}_d^\omega(\alpha), \overline{\Psi}_d^\omega(\beta)) = \angle(\alpha, \beta) = \angle_{(x_0)}(\overline{\Psi}_d^\omega(\alpha), \overline{\Psi}_d^\omega(\beta)).$$

In particular, if $\angle(\alpha, \beta) < \pi$, then the geodesics $\overline{\Psi}_d^\omega(\alpha)$ and $\overline{\Psi}_d^\omega(\beta)$ bound a Euclidean sector in $\text{Con}^\omega(X, (x_0), d)$ for every scaling sequence d and every ultrafilter ω .

Proof. By the Flat Sector Theorem, the second conclusion of the lemma will follow from the first. The first comes from the pair of calculations

$$\begin{aligned} \angle(\overline{\Psi}_d(\alpha), \overline{\Psi}_d(\beta)) &= \lim_{t \rightarrow \infty} \overline{Z}_{(x_0)}(\Psi_d(t, \alpha), \Psi_d(t, \beta)) \\ &= \lim_{t \rightarrow \infty} \lim_n^\omega \overline{Z}_{x_0}(\alpha(td_n), \beta(td_n)) \\ &= \lim_{t \rightarrow \infty} \angle(\alpha, \beta) = \angle(\alpha, \beta) \end{aligned}$$

and similarly

$$\begin{aligned} \angle_{(x_0)}(\overline{\Psi}_d(\alpha), \overline{\Psi}_d(\beta)) &= \lim_{t \rightarrow 0} \overline{Z}_{(x_0)}(\Psi_d(t, \alpha), \Psi_d(t, \beta)) \\ &= \lim_{t \rightarrow 0} \lim_n^\omega \overline{Z}_{x_0}(\alpha(td_n), \beta(td_n)) \\ &= \lim_{t \rightarrow 0} \angle(\alpha, \beta) = \angle(\alpha, \beta). \quad \square \end{aligned}$$

Proposition 4.9. *The map Ψ_d is an isometric embedding. Thus Ψ_d^ω restricted to the diagonal is an isometric embedding.*

Proof. If $\angle(\alpha, \beta) < \pi$, the geodesics $\overline{\Psi}_d(\alpha)$ and $\overline{\Psi}_d(\beta)$ bound a Euclidean sector by Lemma 4.8 which implies that the metric on the Euclidean cone and the metric in the asymptotic cone agree. If $\angle(\alpha, \beta) = \pi$ then $\text{dist}((s, \alpha), (t, \beta)) = s + t = \text{dist}(\Psi_d(s, \alpha), \Psi_d(t, \beta))$. Thus Ψ_d is an isometric embedding. \square

Definition 4.10. The ultrafilter ω induces a total order on the set of scaling sequences by $(d'_n) = d' \preceq d = (d_n)$ if $d'_n \leq d_n$ ω -almost surely.

Lemma 4.11. *Every countable set of scaling sequences is bounded with this ultrafilter ordering on the set of scaling sequences.*

Proof. Let $\{d^i = (d_n^i)\}$ be a countable set of scaling sequences. Let $\bar{d} = (\bar{d}_n)$ where $\bar{d}_n = \max\{d_n^i \mid i \leq n\}$. It is immediate that $d^i \preceq \bar{d}$ for all i .

It takes more care to find a lower bound. For each $i \in \mathbb{N}$, let

$$A_i = \{n \in \mathbb{N} \mid d_n^j > i - 1 \text{ for all } j \leq i\}.$$

Since each scaling sequence is ω -divergent, A_i is the finite intersection of finitely many ω -large set. Hence, A_i is ω -large. The sets A_i form a nested sequence, i.e. $\mathbb{N} = A_1 \supset A_2 \supset \dots$, that we will use to defined \underline{d} . Notice that $\bigcap_{i=1}^\infty A_i = \emptyset$, so $\mathbb{N} = \bigcup_{i=1}^\infty (A_i \setminus A_{i+1})$.

For $n \in A_i \setminus A_{i+1}$, let $\underline{d}_n = \min\{d_n^j \mid j \leq i\}$ and let $\underline{d} = (\underline{d}_n)$. Thus $\underline{d}_n \leq d_n^i$ on A_i , which implies that $\underline{d} \preceq d^i$ for each i . The sequence \underline{d} is ω -divergent since $\underline{d}_n > i - 1$ for all $n \in A_i$. \square

Lemma 4.12. *For every countable subset $C \subset \partial_Z^\omega X$, there exists a scaling sequence $d = (d_n)$ such that for all $(\tilde{d}_n) = \tilde{d} \succeq d$, $\Psi_{\tilde{d}}^\omega$ restricted to $\text{Cone}(C)$ is an isometric embedding. In particular, there exists a scaling sequence d such that if $(\alpha_n), (\beta_n) \in C$, then $\angle(\overline{\Psi}_{\tilde{d}}^\omega(\alpha_n), \overline{\Psi}_{\tilde{d}}^\omega(\beta_n)) = \angle_{(x_0)}(\overline{\Psi}_d^\omega(\alpha_n), \overline{\Psi}_d^\omega(\beta_n))$ for all $\tilde{d} \succeq d$.*

Notice that this implies that if $\lim_n^\omega \angle(\alpha_n, \beta_n) < \pi$, then the geodesics $\Psi_d^\omega(t, (\alpha_n))$ and $\Psi_d^\omega(t, (\beta_n))$ bound a Euclidean sector for all $\tilde{d} \succeq d$.

Proof. Let C be a countable subset of $\partial_Z^\omega X$ and fix $(\alpha_n), (\beta_n) \in C$. Then there exists d_n such that $\overline{Z}_{x_0}(\alpha_n(\sqrt{d_n}), \beta_n(\sqrt{d_n})) \geq \angle(\alpha_n, \beta_n) - 1/n$. Thus for any $(\tilde{d}_n) = \tilde{d} \succeq d$ and $t > 0$, we have $t\tilde{d}_n \geq td_n \geq t\sqrt{d_n}$ ω -almost surely, and therefore $\overline{Z}_{x_0}(\alpha_n(t\tilde{d}_n), \beta_n(t\tilde{d}_n)) \geq \overline{Z}_{x_0}(\alpha_n(\sqrt{d_n}), \beta_n(\sqrt{d_n})) \geq \angle(\alpha_n, \beta_n) - 1/n$ ω -almost surely. Then

$$\begin{aligned}
\angle(\overline{\Psi}_d^\omega(\alpha_n), \overline{\Psi}_d^\omega(\beta_n)) &= \lim_{t \rightarrow \infty} \overline{Z}_{(x_0)}(\Psi_d^\omega(t, (\alpha_n)), \Psi_d^\omega(t, (\beta_n))) \\
&= \lim_{t \rightarrow \infty} \lim_n \overline{Z}_{x_0}^\omega(\alpha_n(t\tilde{d}_n), \beta_n(t\tilde{d}_n)) \\
&\leq \lim_{t \rightarrow \infty} \lim_n \omega \angle(\alpha_n, \beta_n) = \lim_n \omega \angle(\alpha_n, \beta_n)
\end{aligned}$$

and similarly

$$\begin{aligned}
\angle_{(x_0)}(\overline{\Psi}_d^\omega(\alpha_n), \overline{\Psi}_d^\omega(\beta_n)) &= \lim_{t \rightarrow 0} \overline{Z}_{(x_0)}(\Psi_d^\omega(t, (\alpha_n)), \Psi_d^\omega(t, (\beta_n))) \\
&= \lim_{t \rightarrow 0} \lim_n \overline{Z}_{x_0}^\omega(\alpha_n(t\tilde{d}_n), \beta_n(t\tilde{d}_n)) \\
&\geq \lim_{t \rightarrow 0} \lim_n \omega (\angle(\alpha_n, \beta_n) - 1/n) = \lim_n \omega \angle(\alpha_n, \beta_n).
\end{aligned}$$

Thus $\angle(\overline{\Psi}_d^\omega(\alpha_n), \overline{\Psi}_d^\omega(\beta_n)) = \angle_{(x_0)}(\overline{\Psi}_d^\omega(\alpha_n), \overline{\Psi}_d^\omega(\beta_n))$.

Case 1: $\lim_n \omega \angle(\alpha_n, \beta_n) < \pi$. Then the geodesics $\Psi_d^\omega(t, (\alpha_n))$ and $\Psi_d^\omega(t, (\beta_n))$ bound a Euclidean sector which implies that $\text{dist}(\Psi_d^\omega(t, (\alpha_n)), \Psi_d^\omega(s, (\beta_n))) = \text{dist}((t, (\alpha_n)), (s, (\beta_n)))$.

Case 2: $\lim_n \omega \angle(\alpha_n, \beta_n) = \pi$. As before, we have $\text{dist}((s, (\alpha_n)), (t, (\beta_n))) = s + t = \text{dist}(\Psi_d^\omega(s, (\alpha_n)), \Psi_d^\omega(t, (\beta_n)))$.

Thus for every pair of elements of C there exists a scaling sequence satisfying the conclusion of the lemma. Since the set of pairs from C is also countable, Lemma 4.11 implies that there exists a sequence d which will satisfy the conclusion of the lemma. \square

Definition 4.13. Fix $x_0 \in X$. Define $\Xi : X \times \mathbb{R}^+ \rightarrow X$ by

$$\Xi(x, t) = \begin{cases} x & \text{for } t \geq \text{dist}(x_0, x) \\ y \in [x_0, x] \text{ such that } \text{dist}(x_0, y) = t & \text{for } t \leq \text{dist}(x_0, x) \end{cases}$$

Notice that Ξ is the canonical geodesic retraction of X to x_0 via projection onto the closed ball of radius t about x_0 .

Fix scaling sequences $(d'_n) = d' \preceq d = (d_n)$. Then Ξ defines a map $\Theta_{d'}^d : \text{Con}^\omega(X, (x_0), d) \rightarrow \text{Con}^\omega(X, (x_0), d')$ by $\Theta_{d'}^d((x_n)) = (\Xi(x_n, \frac{d'_n}{d_n} \text{dist}(x_n, x_0)))$.

Lemma 4.14. *Let X be a CAT(0) metric space. Then $\Theta_{d'}^d$ is a well-defined 1-Lipschitz map which preserves distance to the observation point of the asymptotic cone.*

If, in addition, X is proper and cocompact; then $\Theta_{d'}^d$ is surjective and $\Theta_{d'}^d \circ \Psi_d^\omega = \Psi_{d'}^\omega$ for all $(d'_n) = d' \preceq d = (d_n)$.

Proof. The maps $\Theta_{d'}^d$ are well-defined 1-Lipschitz maps, since $\Xi|_{X \times \{t\}}$ is 1-Lipschitz for any fixed t . By construction, $\Theta_{d'}^d((x_n))$ has a representative (x'_n) where $\text{dist}(x'_n, x_0) = \frac{d'_n}{d_n} \text{dist}(x_n, x_0)$ ω -almost surely. Thus

$$\text{dist}(\Theta_{d'}^d((x_n)), (x_0)) = \lim \omega \frac{\text{dist}(x'_n, x_0)}{d'_n} = \lim \omega \frac{\text{dist}(x_n, x_0)}{d_n} = \text{dist}((x_n), (x_0)).$$

Notice that $\Theta_{d'}^d \circ \Psi_d^\omega(t, (\alpha_n)) = \Theta_{d'}^d((\alpha_n(td_n))) = (\alpha_n(td'_n)) = \Psi_{d'}^\omega(t, (\alpha_n))$. If X is also proper and cocompact, then $\Psi_{d'}^\omega$ is surjective, which implies that $\Theta_{d'}^d$ must also be surjective. \square

Thus $\Theta_{d'}^d$, while not an isometry, does send geodesics in $\text{Con}^\omega(X, (x_0), d)$ based at (x_0) to geodesics in $\text{Con}^\omega(X, (x_0), d')$ based at (x_0) . When convenient, we can consider the function $\Theta_{d'}^d$ induces on the boundaries of $\text{Con}^\omega(X, (x_0), d)$ and $\text{Con}^\omega(X, (x_0), d')$.

Corollary 4.15. *Let X be a proper cocompact $CAT(0)$ space. Then the map*

$$\overline{\Theta}_{d'}^d : \partial_{\angle} \text{Con}^{\omega}(X, (x_0), d) \rightarrow \partial_{\angle} \text{Con}^{\omega}(X, (x_0), d')$$

defined by $\overline{\Theta}_{d'}^d \circ \overline{\Psi}_d^{\omega}(\alpha_n) = \overline{\Psi}_{d'}^{\omega}(\alpha_n)$ is a non-expansive surjection for all $d' \preceq d$.

Proof. Let $(d'_n) = d' \preceq d = (d_n)$. If $\overline{\Psi}_d^{\omega}(\alpha_n) = \overline{\Psi}_d^{\omega}(\beta_n)$, then $(\alpha_n(td_n)) = (\beta_n(td_n))$ for all t which implies that $(\alpha_n(td'_n)) = (\beta_n(td'_n))$ for all t . Hence, $\overline{\Psi}_{d'}^{\omega}(\alpha_n) = \overline{\Psi}_{d'}^{\omega}(\beta_n)$. Therefore $\overline{\Theta}_{d'}^d$ is well-defined.

Then

$$\begin{aligned} \angle(\overline{\Theta}_{d'}^d \circ \overline{\Psi}_d^{\omega}(\alpha_n), \overline{\Theta}_{d'}^d \circ \overline{\Psi}_d^{\omega}(\beta_n)) &= \angle(\overline{\Psi}_{d'}^{\omega}(\alpha_n), \overline{\Psi}_{d'}^{\omega}(\beta_n)) \\ &= \lim_{t \rightarrow \infty} \overline{Z}_{(x_0)}(\Psi_{d'}^{\omega}(t, (\alpha_n)), \Psi_{d'}^{\omega}(t, (\beta_n))) \\ &= \lim_{t \rightarrow \infty} \overline{Z}_{(x_0)}(\Theta_{d'}^d \circ \Psi_d^{\omega}(t, (\alpha_n)), \Theta_{d'}^d \circ \Psi_d^{\omega}(t, (\beta_n))) \\ &\leq \lim_{t \rightarrow \infty} \overline{Z}_{(x_0)}(\Psi_d^{\omega}(t, (\alpha_n)), \Psi_d^{\omega}(t, (\beta_n))) \\ &= \angle(\overline{\Psi}_d^{\omega}(\alpha_n), \overline{\Psi}_d^{\omega}(\beta_n)) \end{aligned}$$

implies that $\overline{\Theta}_{d'}^d$ is a non-expansive map. Since $\overline{\Psi}_{d'}^{\omega}$ is surjective, so is $\overline{\Theta}_{d'}^d$. \square

Thus $(\text{Con}^{\omega}(X, (x_0), d), \Theta_{d'}^d)$ forms a directed system of metric spaces and one can consider the direct limit $\varinjlim (\text{Con}^{\omega}(X, (x_0), d), \Theta_{d'}^d)$ and the inverse limit $\varprojlim (\text{Con}^{\omega}(X, (x_0), d), \Theta_{d'}^d)$. Before computing the direct limit, we require the following lemma.

Lemma 4.16. *Let X be a proper $CAT(0)$ space. For every $(\alpha_n) \in \partial_{\angle}^{\omega} X$, there exists a scaling sequence d such that $\overline{\Psi}_{d'}^{\omega}(\alpha_n) = \overline{\Psi}_{d'}^{\omega}(\alpha)$ for all $d' \preceq d$ where $\alpha = \lim^{\omega} \alpha_n$ (the limit is taken in $\partial_{\infty} X$). Thus $\Psi_{d'}^{\omega}(t, (\alpha_n)) = \Psi_{d'}^{\omega}(t, \alpha)$ for all t and all $d' \preceq d$.*

Proof. Let $\alpha = \lim^{\omega} \alpha_n$ where the limit is taken in the compact space $\partial_{\infty} X$. Let $\tilde{d}_n = \max\{t \mid \text{dist}(\alpha(t), \alpha_n(t)) \leq C\}$, for some fixed $C > 0$. In other words $\tilde{d}_n = \frac{1}{\text{dist}_C(\alpha, \alpha_n)}$. Since α_n converges to α , ω -almost surely, the sequence $d_n = \sqrt{\tilde{d}_n}$ diverges ω -almost surely. Let $(d'_n) = d' \preceq d = (d_n)$. Then $t \leq d'_n$ ω -almost surely, so

$$\begin{aligned} \text{dist}(\overline{\Psi}_{d'}^{\omega}(\alpha_n)(t), \overline{\Psi}_{d'}^{\omega}(\alpha)(t)) &= \text{dist}((\alpha_n(td'_n), (\alpha(td'_n))) \\ &= \lim_n^{\omega} \frac{\text{dist}(\alpha_n(td'_n), \alpha(td'_n))}{d'_n} \\ &\leq \lim_n^{\omega} \frac{\text{dist}(\alpha_n(d_n'^2), \alpha(d_n'^2))}{d'_n} \\ &\leq \lim_n^{\omega} \frac{C}{d'_n} = 0. \end{aligned} \quad \square$$

Corollary 4.17. *Let X be a proper cocompact $CAT(0)$ space. Then for $x, y \in \text{Con}^{\omega}(X, (x_0), d)$, there exists a scaling sequence $\tilde{d} \preceq d$ such that $\text{dist}(\Theta_{\tilde{d}}^d(x), \Theta_{\tilde{d}}^d(y)) = \text{dist}(\Theta_{\tilde{d}}^d(x), \Theta_{\tilde{d}}^d(y))$ for all $d' \preceq \tilde{d}$.*

Proof. Fix $x, y \in \text{Con}^{\omega}(X, (x_0), d)$. Since X is proper cocompact we can find $s, t \in \mathbb{R}^+$ and geodesics α_n, β_n such that $\Psi_d^{\omega}(s, (\alpha_n)) = x$ and $y = \Psi_d^{\omega}(t, (\beta_n))$. Let $\alpha = \lim^{\omega} \alpha_n$ and $\beta = \lim^{\omega} \beta_n$. We can then find \tilde{d} such that $\Theta_{\tilde{d}}^d(x) = \Psi_{\tilde{d}}^{\omega}(s, (\alpha_n)) = \Psi_{\tilde{d}}^{\omega}(s, \lim^{\omega} \alpha_n)$ and $\Theta_{\tilde{d}}^d(y) =$

$\Psi_d^\omega(t, (\beta_n)) = \Psi_{\tilde{d}}(t, \lim^\omega \beta_n)$. Thus for all $d' \preceq \tilde{d}$

$$\begin{aligned} \text{dist}(\Theta_d^d(x), \Theta_d^d(y)) &= \text{dist}(\Psi_{\tilde{d}}(s, \alpha), \Psi_{\tilde{d}}(t, \beta)) \\ &= \text{dist}(\Psi_{d'}(s, \alpha), \Psi_{d'}(t, \beta)) = \text{dist}(\Theta_{d'}^d(x), \Theta_{d'}^d(y)). \end{aligned} \quad \square$$

Theorem 4.18. *Let X be a proper cocompact CAT(0) space. Then $\lim_{\rightarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$ is isometric to $\text{Cone}(\partial_T X)$. Moreover, for each scaling sequence d the projection map*

$$\Theta_d : \text{Con}^\omega(X, (x_0), d) \rightarrow \lim_{\rightarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$$

is determined by the equation $\Theta_d \circ \Psi_d^\omega(t, (\alpha_n)) = \Theta_d \circ \Psi_d(t, \lim^\omega \alpha_n)$ and $\Theta_d \circ \Psi_d$ is an isometry.

Proof. Recall that by Lemma 4.14, for every $d' \preceq d$, we have that $\Theta_{d'}^d \circ \Psi_d = \Psi_{d'}$. Since $\Theta_{d'}^d \circ \Psi_d^\omega(t, (\alpha_n)) = \Psi_{d'}(t, \lim^\omega \alpha_n)$ for sufficiently small d' (Lemma 4.16), we see that $\Theta_d \circ \Psi_d^\omega(t, (\alpha_n)) = \Theta_d \circ \Psi_d(t, \lim^\omega \alpha_n)$. Thus $\Theta_d \circ \Psi_d$ is surjective.

By Proposition 4.9, Ψ_d is an isometric embedding of $\text{Cone}(\partial_T X)$ into $\text{Con}^\omega(X, (x_0), d)$, which together with the equality $\Theta_{d'}^d \circ \Psi_d = \Psi_{d'}$ shows $\Theta_{d'}^d$ is an isometry when restricted to the image of Ψ_d . Thus $\Theta_d \circ \Psi_d$ is an isometry from $\text{Cone}(\partial_T X)$ to $\lim_{\rightarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$. \square

Remark 4.19. We can now identify $\text{Cone}(\partial_T X)$ with $\lim_{\rightarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$.

Theorem 4.20. *Let X be a proper cocompact CAT(0) space. Then $\lim_{\leftarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$ is a complete metric space with the inverse limit metric and $\text{Cone}(\partial_Z^\omega X)$ isometrically embeds into $\lim_{\leftarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$.*

Proof. Let $x, y \in \lim_{\leftarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$. This means for each scaling sequence d we have $x^d, y^d \in \text{Con}^\omega(X, (x_0), d)$ such that $\Theta_{d'}^d(x^d) = x^{d'}$ and $\Theta_{d'}^d(y^d) = y^{d'}$ for all $d' \preceq d$. The function assigning to each scaling sequence d the real number $\text{dist}(x^d, y^d)$ is an increasing function of d that is bounded by the constant $\text{dist}(x^d, (x_0)) + \text{dist}(y^d, (x_0))$, which is independent of d by Lemma 4.14. Thus $\sup_d \text{dist}(x^d, y^d)$ is finite and we can define a metric (not just an extended metric) on $\lim_{\leftarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d)$ by $\rho((x^d), (y^d)) = \sup_d \text{dist}(x^d, y^d)$. It is easy to verify that ρ does define a complete metric. Lemma 4.14 implies that the functions Ψ_d^ω induce a well-defined map into the inverse limit that is an isometric embedding by Lemma 4.12. \square

5. VISUAL BOUNDARY

We have seen that $\Psi_d^\omega : \partial_Z^\omega X \rightarrow \text{Con}^\omega(X, (x_0), d)$ gives a parametrization of $\text{Con}^\omega(X, (x_0), d)$ which converges to $\text{Cone}(\partial_T X)$ as we allow d to decrease. Thus $\{\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d\}_{d, d'}$ completely determines the Tits boundary $\partial_T(X)$. We now wish to understand the visual boundary $\partial_\infty X$ in terms of $\{\text{Con}^\omega(X, (x_0), d), \Theta_{d'}^d\}_{d, d'}$.

Lemma 5.1. *Let X be a proper CAT(0) space and (α_n) a sequence in ∂X . If $\Psi_d^\omega(t_0, (\alpha_n)) = \Psi_d(t_0, \alpha_0)$ for some scaling sequence d , some $t_0 > 0$ and some $\alpha_0 \in \partial X$, then $\lim^\omega \alpha_n = \alpha_0$.*

Proof. Suppose that $\Psi_d^\omega(t_0, (\alpha_n)) = \Psi_d(t_0, \alpha_0)$ for some $t_0 > 0$. Then

$$\lim_n^\omega \frac{\text{dist}(\alpha_n(t_0 d_n), \alpha_0(t_0 d_n))}{d_n} = 0.$$

Fix $C, \epsilon > 0$. Let A be the ω -large set such that $\text{dist}(\alpha_n(t_0 d_n), \alpha_0(t_0 d_n)) \leq C \epsilon t_0 d_n$ for all $n \in A$. Then the CAT(0) inequality implies that $\text{dist}(\alpha_n(\frac{1}{\epsilon}), \alpha_0(\frac{1}{\epsilon})) \leq C$, and therefore $\text{dist}_C(\alpha_n, \alpha_0) \leq \epsilon$, for all $n \in A \cap B$ where B is the ω -large set such that $d_n \geq \frac{1}{\epsilon t_0}$ for all $n \in B$. Thus $\lim^\omega \alpha_n = \alpha_0$. \square

Theorem 5.2. *Let X be a proper CAT(0) space. For a sequence (α_n) in ∂X , α_n converges to α_0 in the visual boundary of X if and only if for every bijection $\sigma : \mathbb{N} \rightarrow \mathbb{N}$, there exists a scaling sequence d such that $\Psi_d^\omega(t, (\alpha_{\sigma(n)})) = \Psi_d(t, \alpha_0)$ for all t .*

Notice that $\Psi_d^\omega(t, (\alpha_{\sigma(n)})) = \Psi_d(t, \alpha_0)$ for all t is equivalent to $\Theta_d \circ \Psi_d^\omega(t, (\alpha_{\sigma(n)})) = (t, \alpha_0)$ for all t .

Proof. If α_n converges to α_0 in the visual boundary, then $\lim^\omega \alpha_{\sigma(n)} = \alpha_0$ for all bijections σ and the forward implication then follows from Lemma 4.16. Thus we need only show that if for every bijection $\sigma : \mathbb{N} \rightarrow \mathbb{N}$, there exists a scaling sequence d such that $\Psi_d^\omega(t, (\alpha_{\sigma(n)})) = \Psi_d(t, \alpha_0)$ for all t , then α_n converges to α_0 in the visual boundary.

Suppose that there exists a subsequence n_i such that α_{n_i} converges to β . We may assume that $B = \{n_i\}$ has infinite complement in \mathbb{N} . Let A be an ω -large subset of \mathbb{N} with infinite complement. Let $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ be a bijection sending B to A .

Then $\lim^\omega \alpha_{\sigma(n)} = \beta$. By hypothesis, there also exists a scaling sequence d such that $\Psi_d^\omega(t, (\alpha_{\sigma(n)})) = \Psi_d(t, \alpha_0)$ for all t which by Lemma 5.1 implies that $\lim^\omega \alpha_{\sigma(n)} = \alpha_0$. Thus $\alpha_0 = \beta$.

It is then an exercise to use the compactness of $\partial_\infty X$ to show that α_n converges to α_0 , if every convergent subsequence of α_n converges to α_0 . \square

6. QUASI-ISOMETRIES

A map $f : X \rightarrow Y$ of metric spaces is a (λ, C) -quasi-isometric embedding if

$$\frac{1}{\lambda} \text{dist}(x, y) - C \leq \text{dist}(f(x), f(y)) \leq \lambda \text{dist}(x, y) + C \quad \text{for every } x, y \in X$$

and is C -quasi-surjective if $Y \subset \mathcal{N}_C(\text{im}(f))$, i.e., the open C -neighborhood of $\text{im}(f)$ is all of Y . A (λ, C) -quasi-isometry is a C -quasi-surjective (λ, C) -quasi-isometric embedding. A C -quasi-inverse of a map $f : X \rightarrow Y$ is a map $g : Y \rightarrow X$ such that $f \circ g$ is C -close to the identity on Y and $g \circ f$ is C -close to the identity on X . It is a standard exercise to show that every quasi-isometry $f : X \rightarrow Y$ admits a quasi-inverse $g : Y \rightarrow X$, which is itself a quasi-isometry. A quasi-geodesic ray in Y is a quasi-isometric embedding of \mathbb{R}^+ into Y .

Definition 6.1. Let ω be a nonprincipal ultrafilter on \mathbb{N} and $d = (d_n)$ be an ω -divergent sequence of positive real numbers. Suppose that $f : X \rightarrow Y$ is a (λ, C) -quasi-isometric embedding. Then f naturally induces a λ -Lipschitz map $f^\omega : \text{Con}^\omega(X, (x_0), d) \rightarrow \text{Con}^\omega(Y, (f(x_0)), d)$, defined by $f^\omega((x_n)) = (f(x_n))$. If f is a quasi-isometry, f^ω is bi-Lipschitz.

On the other hand, a quasi-isometric embedding (or even a quasi-isometry) $f : X \rightarrow Y$ of proper CAT(0) spaces does not induce a canonical map from ∂X to ∂Y . The reason is that if $\alpha : \mathbb{R}^+ \rightarrow X$ is a geodesic ray, then $f \circ \alpha$ is a quasi-geodesic ray, which in general will not have a unique limit point in $\partial_\infty Y$ (see Proposition 6.8). However, since Y is proper, the pointwise limit $f^d(\alpha) = \lim^\omega [f \circ \alpha(0), f \circ \alpha(d_n)]$ is a geodesic ray in Y . This defines a map $f^d : \partial X \rightarrow \partial Y$. In general, f^d will depend on both ω and the scaling sequence d .

Remark 6.2. The functions $\Theta_d, \Theta_d^d, \Psi_d, \Psi_d^\omega$ are defined for all CAT(0) spaces and have domains and ranges which depend on the chosen CAT(0) space. In most cases we will allow the chosen CAT(0) space to change without changing our notation for the functions $\Theta_d, \Theta_d^d, \Psi_d, \Psi_d^\omega$. When considering a quasi-isometric embedding $f : X \rightarrow Y$, we will always assume that the fixed basepoint in Y used to define the maps $\Theta_d, \Theta_d^d, \Psi_d, \Psi_d^\omega$ is the image of the fixed basepoint in X .

We will now relate the functions Ψ_d, Θ_d , and f^d .

Lemma 6.3. *Let $f : X \rightarrow Y$ be a quasi-isometric embedding of a proper CAT(0) spaces, ω a nonprincipal ultrafilter on \mathbb{N} , and $d = (d_n)$ an ω -divergent sequence of positive real numbers. For every $\alpha \in \partial X$ there exists a scaling sequence $\tilde{d} = (\tilde{d}_n)$ such that for all $d' \preceq \tilde{d}$,*

$$\Theta_{d'}^d \circ f^\omega \circ \Psi_d(1, \alpha) = \Psi_{d'}(1, f^d(\alpha)). \quad (6.4)$$

In particular,

$$\Theta_d \circ f^\omega \circ \Psi_d(1, \alpha) = (1, f^d(\alpha)). \quad (6.5)$$

Proof. Let $f : X \rightarrow Y$ be a quasi-isometric embedding of proper CAT(0) spaces, ω a nonprincipal ultrafilter on \mathbb{N} , and $d = (d_n)$ an ω -divergent sequence of positive real numbers. Fix $\alpha \in \partial X$ and let γ_n be the geodesic from $f \circ \alpha(0)$ to $f \circ \alpha(d_n)$. Let $\tilde{d}_n = \max\{t \mid \text{dist}(f^d(\alpha)(t^2), \gamma_n(t^2)) \leq 1\}$. Notice that \tilde{d}_n is ω -divergent since γ_n converges to $f^d(\alpha)$ by definition.

For $d' = (d'_n) \preceq (\tilde{d}_n)$, we have

$$\begin{aligned} \Theta_{d'}^d \circ f^\omega \circ \Psi_d(1, \alpha) &= \Theta_{d'}^d(f \circ \alpha(d_n)) = \Theta_{d'}^d(\gamma_n(d_n)) \\ &= (\gamma_n(d'_n)) = (f^d(\alpha)(d'_n)) = \Psi_{d'}(1, f^d(\alpha)). \quad \square \end{aligned}$$

Proposition 6.6. *Let $f : X \rightarrow Y$ be a (L, C) -quasi-isometric embedding of a proper CAT(0) spaces, ω a nonprincipal ultrafilter on \mathbb{N} , and $d = (d_n)$ an ω -divergent sequence of positive real numbers. The function f^d is L -Lipschitz continuous when the boundaries are endowed with the Tits or angle metric.*

Proof. First observe that for all $\alpha, \beta \in \partial X$ and any ω -divergent sequence $d' = (d'_n)$,

$$\begin{aligned} 2 \sin\left(\frac{\angle(\alpha, \beta)}{2}\right) &= \lim_{t \rightarrow \infty} \frac{\text{dist}(\alpha(t), \beta(t))}{t} \\ &= \lim_n \omega \frac{\text{dist}(\alpha(d'_n), \beta(d'_n))}{d'_n} = \text{dist}(\Psi_{d'}(1, \alpha), \Psi_{d'}(1, \beta)). \end{aligned}$$

Fix $\alpha, \beta \in \partial X$ and any ω -divergent sequence \tilde{d} satisfying Lemma 6.3. We have

$$\begin{aligned} 2 \sin\left(\frac{\angle(f^d(\alpha), f^d(\beta))}{2}\right) &= \text{dist}(\Psi_{\tilde{d}}(1, f^d(\alpha)), \Psi_{\tilde{d}}(1, f^d(\beta))) \\ &= \text{dist}(\Theta_{\tilde{d}}^d \circ f^\omega \circ \Psi_d(1, \alpha), \Theta_{\tilde{d}}^d \circ f^\omega \circ \Psi_d(1, \beta)) \\ &\leq L \text{dist}(\Psi_d(1, \alpha), \Psi_d(1, \beta)) \\ &= 2L \sin\left(\frac{\angle(\alpha, \beta)}{2}\right). \end{aligned}$$

Thus for every $\epsilon > 0$, we find $\angle(f^d(\alpha), f^d(\beta)) \leq L(1 + \epsilon)\angle(\alpha, \beta)$ for all sufficiently close $\alpha, \beta \in \partial_\angle X$. Hence f^d is $L(1 + \epsilon)$ -Lipschitz by the triangle inequality. It follows that f^d is L -Lipschitz on $\partial_\angle X$. Since the angle and Tits metrics are equal wherever either has value $< \pi$, f^d is L -Lipschitz on $\partial_T X$, too. \square

In general f^d will not define a continuous function on the visual boundary, as illustrated by the following example.

Example 6.7. Let G be the group introduced by Croke and Kleiner in [5] that acts properly cocompactly on CAT(0) spaces X and Y , which have non-homeomorphic visual boundaries. Then there exist G -equivariant quasi-isometries $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that both $f \circ g$ and $g \circ f$ are uniformly close to the identity map. Since G has rank one, the set of Morse geodesics of both $\partial_\infty X$ and $\partial_\infty Y$ are dense.

If for some d , both f^d and g^d induced continuous maps on the visual boundary, then Lemma 6.15 would imply that $f^d \circ g^d$ is the identity on a dense set of $\partial_\infty Y$ and hence the identity function on all of $\partial_\infty Y$. Similarly, $g^d \circ f^d$ would be the identity function on $\partial_\infty X$. Thus f^d would give a homeomorphism from $\partial_\infty X$ to $\partial_\infty Y$, a contradiction. Thus, for every d , at least one of $f^d : \partial_\infty X \rightarrow \partial_\infty Y$ and $g^d : \partial_\infty Y \rightarrow \partial_\infty X$ is not continuous.

Alternatively, Ruane and Bowers give an example of two actions of a group G on the product of a tree with \mathbb{R} such that the G -equivariant induced quasi-isometry does not induce a continuous map of the boundary [3].

In [13], it is shown that the limit set of a geodesic under a quasi-isometry can be any connected, compact subset of Euclidean space. One might hope that for the sufficiently small scaling sequences the maps f^d would tend to choose a favorite limit point, i.e. f^d stabilize for a fixed α and sufficiently small scaling sequence d . However the following proposition illustrates that this is not the case.

Proposition 6.8. *Let $f : X \rightarrow Y$ be a quasi-isometric embedding of proper CAT(0) spaces and fix $\alpha \in \partial X$. Then $\Lambda(\text{im}(f \circ \alpha)) = \{f^d(\alpha) \mid d \text{ is a scaling sequence}\}$. For every $\gamma \in \Lambda(\text{im}(f \circ \alpha))$ and scaling sequence d , there exist scaling sequences d' , \tilde{d} such that $d' \preceq d \preceq \tilde{d}$ and $f^{\tilde{d}}(\alpha) = \gamma = f^{d'}(\alpha)$.*

Proof. Let $f : X \rightarrow Y$ be a quasi-isometric embedding of CAT(0) spaces and fix a scaling sequence $d = (d_n)$. Let $\gamma \in \Lambda(\text{im}(f \circ \alpha))$, i.e. γ is in the limit set of $f \circ \alpha$. Then there exist \bar{d}_n such that $f \circ \alpha(\bar{d}_n)$ converges (not just ω -almost surely) to γ in \bar{Y} which implies that $\lim^\omega [f \circ \alpha(0), f \circ \alpha(\bar{d}_n)] = \gamma$.

Let $d'_n = \max\{\bar{d}_i \mid \bar{d}_i \leq d_n\}$ and $\tilde{d}_n = \min\{\bar{d}_i \mid \bar{d}_i \geq d_n\}$. Then for $d' = (d'_n)$ and $\tilde{d} = (\tilde{d}_n)$, we have $d' \preceq d \preceq \tilde{d}$ and it is elementary to check that $f^{d'}(\alpha) = \gamma = f^{\tilde{d}}(\alpha)$. \square

Theorem 6.9. *Let X be a proper cocompact CAT(0) space such that $\partial_T(X)$ is compact. Then $\partial_T(X)$ is homeomorphic to $\partial_\infty X$ and if there exists a quasi-isometry $f : X \rightarrow Y$, then f^d is a bi-Lipschitz equivalence from $\partial_T X$ to $\partial_T Y$.*

Proof. There is always a continuous bijection from $\partial_T X$ to $\partial_\infty X$. If $\partial_T X$ is compact, then this is a homeomorphism.

Suppose there exists a quasi-isometry $f : X \rightarrow Y$. Fix a scaling sequence $d = (d_n)$. If $\partial_T X$ is compact, then $\partial_\infty X$ is isometric to $\partial_T X$ and $\text{Cone}(\partial_\infty X)$ is locally compact. Thus both $\text{Con}^\omega(X, (x_0), d)$ and $\text{Con}^\omega(Y, (f(x_0)), d)$ are proper metric spaces, since Ψ_d^ω is a 1-Lipschitz surjection that preserves distances to the basepoint. Then $\Psi_d(1, \cdot)$ embeds $\partial_T Y$ as a closed subset of a closed ball of the proper metric space, $\text{Con}^\omega(Y, (f(x_0)), d)$. Thus $\partial_T Y$ is also compact and $\partial_\infty Y = \partial_T Y$.

Thus Ψ_d is a surjection when viewed as a map from $\text{Cone}(\partial_T X)$ to $\text{Con}^\omega(X, (x_0), d)$ or as a map from $\text{Cone}(\partial_T Y)$ to $\text{Con}^\omega(Y, (f(x_0)), d)$. By Lemma 4.9, Θ_d is an isometry when restricted to the image of Ψ_d . Thus $f^d = \Theta_d \circ f^\omega \circ \Psi_d|_{\{1\} \times \partial_T X}$ is bi-Lipschitz. \square

It follows that compactness of the Tits boundary is a quasi-isometry invariant. However, this restriction is so severe it essentially only holds for Euclidean flats. More precisely, we give the following alternate proof of an unpublished result of Bosché [2, Proposition 3].

Theorem 6.10. *Let X be a proper CAT(0) space admitting a properly discontinuous, cocompact group action by isometries. If $\partial_T X$ is compact then X has a quasi-dense Euclidean flat.*

Proof. As noted in the proof of Theorem 6.9, this implies that $\text{Con}^\omega(X, (x_0), d)$ is a proper metric space, which implies that $\text{Con}^\omega(X, (x_0), d)$ is a Euclidean flat by [11, 12]. Thus the Tits

boundary of X is a sphere by Lemma 4.7 and Proposition 4.9. Thus by [9, Proposition 2.1], X contains a flat F with $\partial_\infty F = \partial_\infty X$.

We now only need show that F is quasi-dense. By [6] there is some constant K (depending only on X and the action) such that for every $x, y \in X$ there exists a geodesic ray α based at x such that $\text{dist}(\text{im}(\alpha), y) \leq K$. Suppose that there exists a point $y \in X$ such that $\text{dist}(y, F) \geq 2K$. Let $x \in F$ be the closest point projection of y in F and α a geodesic ray based at x such that $\text{dist}(\text{im}(\alpha), y) \leq K$. Then $\angle_x(y, \alpha) \leq \pi/6$ by comparison with Euclidean geometry (law of sines), and $\angle_x(y, z) \geq \pi/2$ for any $z \in F \setminus \{x\}$. Thus for any geodesic ray β in F based at x , we have that $\angle_x(\alpha, \beta) \geq \pi/4$. Thus $\alpha \notin \partial F$, which contradicts the fact that $\partial_\infty F = \partial_\infty X$. Thus $X \subset \mathcal{N}_{2K}(F)$ and F is quasi-dense. \square

Thus by Bieberbach's Theorem [1], we have the following corollary.

Corollary 6.11. *Every finitely generated group which acts properly discontinuously, cocompactly, and by isometries on a CAT(0) space with compact Tits boundary is virtually free abelian.*

The Hausdorff distance between two subsets A, B of a metric space X is

$$\text{dist}_H(A, B) = \max \left\{ \sup_{a \in A} \left\{ \inf_{b \in B} \text{dist}(a, b) \right\}, \sup_{b \in B} \left\{ \inf_{a \in A} \text{dist}(a, b) \right\} \right\}.$$

Lemma 6.12. *Let α and β be quasi-geodesic rays in an arbitrary metric space. If $\text{im}(\alpha) \subset \mathcal{N}_M(\text{im}(\beta))$ for some M , then the Hausdorff distance $\text{dist}_H(\text{im}(\alpha), \text{im}(\beta))$ between $\text{im}(\alpha)$ and $\text{im}(\beta)$ is finite.*

Proof. Let X be a metric space, and let α and β be (L', C') -quasi-geodesic and (L, C) -quasi-geodesic rays, respectively, in X such that $\text{im}(\alpha) \subset \mathcal{N}_M(\text{im}(\beta))$ for some M .

We need to show there exists M' such that $\text{im}(\beta) \subset \mathcal{N}_{M'}(\text{im}(\alpha))$. Let $t_0 = 0$ and, for $n \in \mathbb{N}$, fix t_n such that $\text{dist}(\beta(t_n), \alpha(n)) \leq M$. Notice that t_n is unbounded which implies that the union of the closed intervals between t_n and t_{n+1} is $\mathbb{R}^{\geq 0}$. By the triangle inequality, $\text{dist}(\beta(t_n), \beta(t_{n+1})) \leq 2M + (L' + C')$ which implies that $|t_n - t_{n+1}| \leq L(2M + L' + C' + C)$.

Thus for every t in the interval between t_n and t_{n+1} , we have $\text{dist}(\beta(t), \beta(t_n)) \leq L^2(2M + L' + C' + C) + C$ and hence $\text{dist}(\beta(t), \alpha(n)) \leq L^2(2M + L' + C' + C) + C + M$.

Therefore $\text{im}(\beta) \subset \mathcal{N}_{L^2(2M+L'+C'+C)+C+M}(\text{im}(\alpha))$. This proves the lemma. \square

Definition 6.13 (Morse). A (quasi-)geodesic γ is called a *Morse (quasi-)geodesic*, if for every $L \geq 1$ and $C \geq 0$ there exists an $M = M(L, C)$ such that every (L, C) -quasi-geodesic with endpoints on γ remains M -close to γ .

Lemma 6.14. *Let $f : X \rightarrow Y$ be a (L, C) -quasi-isometry of proper CAT(0) spaces. If α is a Morse ray in ∂X , then $f^d(\alpha)$ is a Morse ray in ∂Y and there exists an M such that $\text{dist}_H(f \circ \alpha, f^d(\alpha)) < M$.*

Proof. It is immediate that $f \circ \alpha$ is Morse. So for every n , $[f \circ \alpha(0), f \circ \alpha(d_n)] \subset \mathcal{N}_{M'}(\text{im}(f \circ \alpha))$ for some M' depending only on α, L, C , which implies that $\text{im}(f^d(\alpha)) \subset \mathcal{N}_{M'+1}(\text{im}(f \circ \alpha))$. By Lemma 6.12, $\text{im}(f \circ \alpha)$ and $\text{im}(f^d(\alpha))$ are finite Hausdorff distance apart. Since any quasi-geodesic ray that is finite Hausdorff distance from a Morse quasi-geodesic ray is also Morse, $f^d(\alpha)$ is a Morse geodesic ray at $f(x_0)$. \square

Lemma 6.15. *Let $f : X \rightarrow Y$ be a quasi-isometry of proper CAT(0) metric spaces with quasi-inverse $g : Y \rightarrow X$. If $\alpha \in \partial X$ is a Morse ray, then $f^d(\alpha) = f^d(\alpha)$ and $g^d \circ f^d(\alpha) = \alpha$ for all scaling sequences d, d' .*

Proof. Since $\text{dist}_H(f \circ \alpha, f^d(\alpha)), \text{dist}_H(f \circ \alpha, f^{d'}(\alpha))$ are both finite then $f^d(\alpha)$ and $f^{d'}(\alpha)$ are asymptotic geodesics based at $f(x_0)$. Hence $f^d(\alpha) = f^{d'}(\alpha)$.

Since g is a quasi-isometry and $f^d(\alpha)$ is a Morse geodesic, $\text{dist}_H(g \circ f^d(\alpha), g^{d'} \circ f^d(\alpha))$ and $\text{dist}_H(g \circ f(\alpha), g \circ f^d(\alpha))$ are both finite. Since g is a quasi-inverse of f , $\text{dist}_H(\alpha, g \circ f(\alpha))$ is also finite. Thus $g^{d'} \circ f^d(\alpha)$ and α are asymptotic, and therefore equal. \square

Although Morse rays are well-behaved under quasi-isometries, other rays are generally not.

Example 6.16. Consider the family of maps $f: X \rightarrow X$ on $X = \mathbb{R}^2$ given in polar coordinates by $f(r, \theta) = (r, \theta + h(r))$, where h is differentiable on $r > 0$. (The case $h(r) = \log(r)$ is the classic “logarithmic spiral” map.) A short calculation reveals the Jacobian of f at $(x, y) = (r \cos \theta, r \sin \theta)$ is, up to multiplying on the left and right by orthogonal matrices,

$$\begin{pmatrix} 1 & 0 \\ 0 & r \end{pmatrix} \begin{pmatrix} 1 & 0 \\ h'(r) & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{r} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ rh'(r) & 1 \end{pmatrix}$$

for $(x, y) \neq (0, 0)$. Thus we see that f is Lipschitz if and only if $rh'(r)$ is bounded. If so, f^{-1} is also clearly well-defined and Lipschitz, so f is bi-Lipschitz and therefore a quasi-isometry.

Again, the case $h(r) = \log(r)$ is the classic “logarithmic spiral” map, and it is easy to see that f^d can take any given point in ∂X to any other point of ∂X by adjusting d .

The case $h(r) = \log(\log(r))$ also has the property that f^d can map any point in ∂X to any other by choosing the appropriate d . However, in this case, for every d the induced bi-Lipschitz map $f^\omega: \text{Con}^\omega(X, (x_0), d) \rightarrow \text{Con}^\omega(X, (f(x_0)), d)$ is an isometry, since $rh'(r) \rightarrow 0$ as $r \rightarrow \infty$.

More examples can be constructed from the above by using the following general lemma.

Lemma 6.17. *Let Y be a geodesic metric space and $\pi: Y \rightarrow X$ be a locally isometric covering map. If $\tilde{f}: Y \rightarrow Y$ is a continuous lift of an L -Lipschitz map $f: X \rightarrow X$, then \tilde{f} is L -Lipschitz.*

Proof. Assume $L > 0$ (the case $L = 0$ is trivial). By the triangle inequality, it suffices to prove \tilde{f} is locally L -Lipschitz—that is, for every $y \in Y$ there is some $\delta > 0$ such that $\tilde{f}|_{B_\delta(y)}$ is L -Lipschitz. So let $y \in Y$ and $x = \pi(y)$. By hypothesis on π , there exists $\epsilon > 0$ such that $\pi|_{B_\epsilon(\tilde{f}(y))}: B_\epsilon(\tilde{f}(y)) \rightarrow B_\epsilon(f(x))$ is an isometry, and $\delta > 0$ such that $\pi|_{B_\delta(y)}: B_\delta(y) \rightarrow B_\delta(x)$ is an isometry. By continuity of \tilde{f} , we may assume $\delta > 0$ is small enough that $\tilde{f}(B_\delta(y)) \subset B_\epsilon(\tilde{f}(y))$.

So let $p, q \in B_\delta(y)$. Since $\pi|_{B_\epsilon(\tilde{f}(y))}$ and $\pi|_{B_\delta(y)}$ are isometries, we find

$$\begin{aligned} \text{dist}(\tilde{f}(p), \tilde{f}(q)) &= \text{dist}(\pi(\tilde{f}(p)), \pi(\tilde{f}(q))) = \text{dist}(f(\pi(p)), f(\pi(q))) \\ &\leq L \cdot \text{dist}(\pi(p), \pi(q)) = L \cdot \text{dist}(p, q). \end{aligned}$$

Thus we have shown that \tilde{f} is locally L -Lipschitz, which proves the lemma. \square

Corollary 6.18. *Let X be a locally $\text{CAT}(0)$ metric space with universal cover \tilde{X} . If $f: X \rightarrow X$ is bi-Lipschitz then its lift $\tilde{f}: \tilde{X} \rightarrow \tilde{X}$ is bi-Lipschitz.*

Example 6.19. Let $X = \mathbb{R}^2 \setminus B_1(x)$ and let \tilde{X} be its universal cover with the induced $\text{CAT}(0)$ metric, which is proper. Consider the family of maps $f: X \rightarrow X$ on X given in polar coordinates by $f(r, \theta) = (r, \theta + h(r))$, where h is differentiable on $r \geq 1$. From Example 6.16, we see that f is bi-Lipschitz if and only if $rh'(r)$ is bounded, so by Corollary 6.18, $\tilde{f}: \tilde{X} \rightarrow \tilde{X}$ is bi-Lipschitz under the same conditions.

Now notice that $\partial\tilde{X}$ is isometric (under the Tits metric) to the disjoint union of a line and two points, corresponding to the directions $\theta \in \mathbb{R}$ and $\theta = \pm\infty$, respectively. The isolated points $\theta = \pm\infty$ are Morse, so they are fixed by every f^d . The other points, however, are not.

The case $h(r) = \log(r)$ has $f^d(\theta) = +\infty$ for all $\theta \in \mathbb{R} \subset \partial\tilde{X}$. So does $h(r) = \log(\log(r))$, and in this case \tilde{f}^ω is always an isometry. Thus while Morse rays map to Morse rays, they may not be the only rays that do.

The case $h(r) = \log(r) \sin(\log(\log(r)))$ has, for any given $\theta_1 \in \mathbb{R}$ and $\theta_2 \in \mathbb{R} \cup \{\pm\infty\}$, some d such that $f^d(\theta_1) = \theta_2$. So does $h(r) = \log(\log(r)) \sin(\log(\log(\log(r))))$, and in this case \tilde{f}^ω is always an isometry. Thus even the Tits component of $f^d(\alpha)$ is, in general, not solely determined by f and α , but can also depend on d .

Proposition 6.20. *Let $\alpha, \beta \in \partial X$ for X a proper CAT(0) space. Then the following are equivalent.*

- (i) *The Tits distance from α to β is finite.*
- (ii) *For every (or some) d , $\text{dist}_T(\overline{\Psi}_d(\alpha), \overline{\Psi}_d(\beta))$ is finite.*
- (iii) *For every (or some) d and any $t > 0$, $\Psi_d(t, \alpha), \Psi_d(t, \beta)$ are contained in the same component of $\text{Con}^\omega(X, (x_0), d) \setminus \{(x_0)\}$.*

Proof. The implication (i) \Rightarrow (ii) follows immediately from Lemma 4.8, and (ii) \Rightarrow (iii) is trivial. Thus we need only show (iii) \Rightarrow (i).

Let $\gamma' : [0, 1] \rightarrow \text{Con}^\omega(X, (x_0), d) \setminus \{(x_0)\}$ be a path from $\Psi_d(1, \alpha)$ to $\Psi_d(1, \beta)$. Since $\text{dist}(\text{im}(\gamma'), (x_0)) > 0$, we can use the geodesic retraction of $\text{Con}^\omega(X, (x_0), d) \setminus \{(x_0)\}$ to find a path $\gamma : [0, 1] \rightarrow \text{Con}^\omega(X, (x_0), d) \setminus \{(x_0)\}$ from $\Psi_d(t_0, \alpha)$ to $\Psi_d(t_0, \beta)$ such that $\text{dist}(\gamma(s), (x_0)) = t_0$ for all s .

Then we can fix k sufficiently large such that $\text{dist}(\gamma(\frac{i}{k}), \gamma(\frac{i+1}{k})) < t_0$ for all $0 \leq i \leq k-1$. Fix α_n^i a geodesic from x_0 to x_n^i such that $(x_n^i) = \gamma(\frac{i}{k})$. Let $\alpha_i = \lim_n^\omega \alpha_n^i$. Notice that $\alpha = \alpha_0$ and $\beta = \alpha_k$. Fix $l \in \mathbb{N}$. Then there exists an ω -large set, A , such that $\text{dist}(\alpha_i(l), \alpha_n^i(l)) < 1$ and $\text{dist}(\alpha_n^i(t_0 d_n), \alpha_n^{i+1}(t_0 d_n)) \leq \frac{3}{2} t_0 d_n$ for all $n \in A$ and for $0 \leq i \leq k$.

Thus for any $n \in A$ such that $t_0 d_n > l$, we have

$$\frac{\frac{3}{2} t_0 d_n + 2}{t_0 d_n} \geq \frac{\text{dist}(\alpha_i(l), \alpha_{i+1}(l))}{l}$$

by the CAT(0) inequality. Thus $\angle(\alpha_i, \alpha_{i+1}) < \pi$ and $\text{dist}_T(\alpha, \beta) < k\pi$. \square

Corollary 6.21. *Let X be a proper CAT(0) space. Then $\text{Con}^\omega(X, (x_0), d)$ has a cut-point if and only if the Tits diameter of X is infinite.*

Since, under a cocompact action, having infinite Tits diameter is equivalent to having a periodic Morse geodesic, we have the following.

Corollary 6.22. *Let X be a proper cocompact CAT(0) space. If $\text{Con}^\omega(X, (x_0), d)$ has a cut-point, then all asymptotic cones of X have cut-points and X contains a periodic Morse geodesic.*

7. QUESTIONS

Theorem 4.20 shows that the limit of asymptotic cones contains a canonical copy of $\text{Cone}(\partial_Z^\omega X) = (\text{Cone}(\partial_T X))^\omega$ for proper cocompact CAT(0) spaces. However, it is not clear even in simple cases, for example where X is a tree, if this embedding is surjective.

Question 7.1. *Is the inverse limit $\lim_{\leftarrow} (\text{Con}^\omega(X, (x_0), d), \Theta_d^d)$ isometric to $\text{Cone}(\partial_Z^\omega X)$?*

Example 6.7 shows that in general f^d cannot be a homeomorphism. However, it would be interesting to consider if f^d is a homotopy equivalence or if f^d preserves homotopy groups.

Question 7.2. *Let $f : X \rightarrow Y$ be a quasi-isometry of proper cocompact CAT(0) spaces. Does the continuous map $f^d : \partial_T X \rightarrow \partial_T Y$ induce an injective homomorphism of homotopy groups?*

REFERENCES

- [1] Ludwig Bieberbach. über die Bewegungsgruppen der Euklidischen Räume. *Math. Ann.*, 70(3):297–336, 1911.
- [2] Aurélien Bosché. Tits compact $\text{cat}(0)$ spaces. <http://arxiv.org/abs/1106.0149v2>.
- [3] Philip L. Bowers and Kim Ruane. Boundaries of nonpositively curved groups of the form $G \times \mathbf{Z}^n$. *Glasgow Math. J.*, 38(2):177–189, 1996.
- [4] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [5] Christopher B. Croke and Bruce Kleiner. Spaces with nonpositive curvature and their ideal boundaries. *Topology*, 39(3):549–556, 2000.
- [6] Ross Geoghegan and Pedro Ontaneda. Boundaries of cocompact proper $\text{CAT}(0)$ spaces. *Topology*, 46(2):129–137, 2007.
- [7] M. Gromov. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.*, (53):53–73, 1981.
- [8] M. Gromov. Asymptotic invariants of infinite groups. In *Geometric group theory, Vol. 2 (Sussex, 1991)*, volume 182 of *London Math. Soc. Lecture Note Ser.*, pages 1–295. Cambridge Univ. Press, Cambridge, 1993.
- [9] Bernhard Leeb. *A characterization of irreducible symmetric spaces and Euclidean buildings of higher rank by their asymptotic geometry*, volume 326 of *Bonner Mathematische Schriften [Bonn Mathematical Publications]*. Universität Bonn, Mathematisches Institut, Bonn, 2000.
- [10] P. Pansu. Croissance des boules et des géodésiques fermées dans les nilvariétés. *Ergodic Theory Dynam. Systems*, 3(3):415–445, 1983.
- [11] F. Point. Groups of polynomial growth and their associated metric spaces. *J. Algebra*, 175(1):105–121, 1995.
- [12] Mark Sapir. On groups with locally compact asymptotic cones. *Internat. J. Algebra Comput.*, 25(1-2):37–40, 2015.
- [13] Dan Staley. Erratic behavior of $\text{CAT}(0)$ geodesics under G -equivariant quasi-isometries. *Geom. Dedicata*, 159:169–184, 2012.