## COBOUNDARIES OF NONCONVENTIONAL ERGODIC AVERAGES

#### I. ASSANI

ABSTRACT. Let  $(X, \mathcal{A}, \mu)$  be a probability measure space and let  $T_i$ ,  $1 \le i \le H$ , be invertible bi measurable measure preserving transformations on this measure space. We give a sufficient condition for the product of H bounded functions  $f_1, f_2, ..., f_H$  to be a coboundary. This condition turns out to be also necessary when one seeks bounded coboundaries.

### 1. INTRODUCTION

The purpose of this short article is to answer a question brought to our attention by S. Donoso (<sup>1</sup>) during the 2017 ETDS workshop held at Chapel Hill.

To this end we refine the setting in [2].

**Definition 1.1.** A probability measure preserving system  $(X, \mathcal{F}, \mu, T_1, T_2, ..., T_H)$  is a combination of a probability measure space  $(X, \mathcal{F}, \mu)$  and  $T_i, 1 \leq i \leq H$  bi-measurable invertible measure preserving maps acting on this probability space.

Given a probability measure preserving system  $(X, \mathcal{F}, \mu, T_1, T_2, ..., T_H)$ ,  $\mu_{\Delta}$  is the diagonal measure on  $X^H$ ,  $\Phi = T_1 \times T_2 \times \cdots \times T_H$ , and  $\nu$  is the diagonal-orbit measure of  $\Phi$ , i.e.

$$u(A) = \frac{1}{3} \sum_{n \in \mathbb{Z}} \frac{1}{2^{|n|}} \mu_{\Delta}(\Phi^{-n}A).$$

We note that  $\nu$  is nonsingular, since  $\frac{1}{3}\nu(A) \leq \nu(\Phi^{-1}A) \leq 2\nu(A)$ ..

**Definition 1.2.** The diagonal orbit system of the probability measure preserving system  $(X, \mathcal{F}, \mu, T_1, T_2, ..., T_H)$  is the system  $(X^H, \mathcal{F}^H, \nu, \Phi)$ .

#### Remarks

- (1) The maps  $T_i$  do not necessarily commute.
- (2) The nonsingularity of  $\Phi$  with respect to  $\nu$  implies the following simple but key lemma (This lemma does not seem to hold when one replaces  $\nu$  with the diagonal measure  $\mu_{\Delta}$  on  $(X^H, \mathcal{F}^H)$ , defined by the equation  $\int F(x_1, x_2, ..., x_H) d\mu_{\Delta} = \int F(x, x, ..., x) d\mu$ ).

**Lemma 1.3.** Let  $F_n$  be a sequence of measurable functions defined on  $X^H$ . If  $F_n$  converges v a.e. then the sequence  $G_n = F_n \circ \Phi$  converges v a.e. as well.

Department of Mathematics, UNC Chapel Hill, NC 27599, assani@email.unc.edu.

<sup>&</sup>lt;sup>1</sup>He indicated that this question was mentioned to him by J.P. Conze and Y. Kifer

*Proof.* Let  $A = \{z \in X^H; F_n(z) \text{ converges}\}$  and  $B = \{z \in X^H; G_n(z) \text{ converges}\}$ . We have  $B = \Phi^{-1}(A)$ . Therefore if  $\nu(A^c) = 0$  we have  $\nu(\Phi^{-1}(A^c)) = 0$  by the non singularity of  $\nu$ .

We wish to prove the following proposition.

**Proposition 1.4.** Let  $(X, \mathcal{F}, \mu, T_1, ..., T_H)$  be a measure preserving system, and  $f_1, f_2, ..., f_H \in L^{\infty}(\mu)$  and  $1 \le p < \infty$ .

(1) Let us assume that the supremum of the nonconventional ergodic sums is  $L^p$ -bounded, i.e.

$$\sup_{N} \left\| \sum_{n=1}^{N} \prod_{i=1}^{H} f_{i} \circ T_{i}^{n} \right\|_{L^{p}(\nu)} < \infty.$$

(2) Then the product of the functions is a coboundary in  $L^p(X^H, \nu)$ , i.e. if  $\Phi = T_1 \times T_2 \times \cdots \times T_H$ , there exists  $V \in L^p(X^H, \nu)$  such that

$$\bigotimes_{i=1}^{H} f_i = V - V \circ \Phi, \nu\text{-a.e.}$$

Therefore, for  $\mu_{\Delta}$ -a.e.  $(x_1, x_2, ..., x_H) \in X^H$ , we have

$$f_1(x_1)f_2(x_2)\cdots f_H(x_H) = V(x_1, x_2, \dots, x_H) - V(T_1x_1, T_2x_2, \dots, T_Hx_H).$$

We give only the proof for the case p = 1. We use the following a.e.-convergence result obtained by Komlós in 1967. When  $1 the reflexivity of <math>L^p(v)$  allows to bypass this lemma. For  $p = \infty$  the assumptions (1) and (2) in the statement of Theorem 1.4 are equivalent. We state it separately as a corollary.

**Lemma 1.5** ([3]). Let  $(X, \mathcal{F}, \mu)$  be a probability measure space, and  $(g_n)$  be a sequence in  $L^1(\mu)$ . Assume that  $\liminf_n \|g_n\|_{L^1(\mu)} < \infty$ . Then there exists a subsequence  $(g_{n_k})_k$  and a function  $g \in L^1(\mu)$  such that for  $\mu$ -a.e.  $x \in X$ ,

$$\lim_{K\to\infty}\frac{1}{K}\sum_{k=1}^K g_{n_k}(x) = g(x).$$

*Proof of Proposition* 1.4. We show that the techniques used for an invariant measure can be applied to our current nonsingular setting. The assumption made tells us that, if  $F = f_1 \otimes f_2 \otimes \cdots \otimes f_d$ , we have

$$\lim_{N\to\infty} \left\| \frac{1}{N} \sum_{n=1}^{N} F \circ \Phi^n \right\|_{L^1(\nu)} = 0.$$

Therefore, there exists a subsequence  $N_k$  of natural numbers such that

$$\lim_{N_k\to\infty}\frac{1}{N_k}\sum_{n=1}^{N_k}F\circ\Phi^n(z)=0$$

for  $\nu$ -a.e.  $z \in X^H$ . Because  $\Phi$  is nonsingular with respect to  $\nu$ , we also know that

$$\lim_{N_k \to \infty} \frac{1}{N_k} \sum_{n=1}^{N_k} F \circ \Phi^{n+1}(z) = 0$$

for  $\nu$ -a.e.  $z \in X^H$  for the same subsequence  $(N_k)$ . Set

$$D_{N_k} = \frac{1}{N_k} \sum_{n=1}^{N_k} \sum_{j=0}^{n-1} F \circ \phi^j.$$

Since  $\sup_n \left\|\sum_{j=0}^{n-1} F \circ \phi^j\right\|_{L^1(\nu)} < \infty$ ,  $\liminf_k \left\|D_{N_k}\right\|_{L^1(\nu)} < \infty$ . Thus, we may apply Lemma 1.5 to show that there exists a subsequence of  $(D_{N_k})$  (which remain denoted as  $(D_{N_k})$ ) such that the averages  $\frac{1}{K} \sum_{k=1}^{K} D_{N_k}$  converge  $\nu$ -a.e to a function  $V \in L^1(\nu)$ . Similarly, by Lemma 1.3 we have

$$V \circ \Phi(z) = \lim_{K} \frac{1}{K} \sum_{k=1}^{K} D_{N_k} \circ \Phi(z)$$

for  $\nu$ -a.e.  $z \in X^H$ . Therefore,

$$V - V \circ \Phi = \lim_{K} \left( \frac{1}{K} \sum_{k=1}^{K} F - \frac{1}{K} \sum_{k=1}^{K} \left( \frac{1}{N_k} \sum_{n=1}^{N_k} F \circ \Phi^n \right) \right) = F$$

Since  $V - V \circ \Phi = F$  for  $\nu$ -a.e., the equality certainly holds for  $\mu_{\Delta}$ -a.e. (the construction of  $\nu$  guarantees that  $V \in L^1(\mu_{\Delta})$ ).

**Corollary 1.6.** Let  $(X, \mathcal{F}, \mu, T_1, ..., T_H)$  be a measure preserving system, and  $f_1, f_2, ..., f_H \in L^{\infty}(\mu)$  The following statements are equivalent.

(1) The supremum of the nonconventional ergodic sums is  $L^{\infty}$ -bounded, i.e.

$$\sup_{N} \left\| \sum_{n=1}^{N} \prod_{i=1}^{H} f_{i} \circ T_{i}^{n} \right\|_{L^{\infty}(\nu)} < \infty.$$

(2) The product of the functions is a coboundary in  $L^{\infty}(X^{H}, \nu)$ , i.e. if  $\Phi = T_1 \times T_2 \times \cdots \times T_H$ , there exists  $V \in L^{\infty}(X^{H}, \nu)$  such that

$$\bigotimes_{i=1}^{H} f_i = V - V \circ \Phi, \nu\text{-a.e.}$$

*Therefore, for*  $\mu$ *-a.e.*  $x \in X$ *, we have* 

$$f_1(x)f_2(x)\cdots f_H(x) = V(x, x, \dots, x) - V(T_1x, T_2x, \dots, T_Hx).$$

*Proof.* The implication 1) implies 2) can be obtained by following the same path as in the proof of the previous proposition. The only thing to check is that  $V \in L^{\infty}(\nu)$ . But this follows from the fact that if

$$\sup_{N} \left\| \sum_{n=1}^{N} \prod_{i=1}^{H} f_{i} \circ T_{i}^{n} \right\|_{L^{\infty}(\nu)} < C < \infty$$

#### I. ASSANI

then  $||D_{N_k}||_{L^{\infty}} \leq C$ . From this observation one can conclude that the limit function V is also in  $L^{\infty}(\nu)$ . For the reverse implication, if  $F = f_1 \times f_2 \dots \times f_H$  is a coboundary in  $L^{\infty}(\nu)$  (i.e.  $F = V - V \circ \Phi$  where  $V \in L^{\infty}(\nu)$ ) then

$$\sum_{n=1}^{N} F \circ \Phi^n = V - V \circ \Phi^{N+1}$$

and

$$\|\sum_{n=1}^{N} F \circ \Phi^{n}\|_{L^{\infty}(\nu)} = \|V - V \circ \Phi^{N+1}\|_{L^{\infty}(\nu)} \le 2\|V\|_{L^{\infty}(\nu)}$$

# Remark

The assumption

$$\sup_{N} \left\| \sum_{n=1}^{N} \prod_{i=1}^{H} f_{i} \circ T_{i}^{n} \right\|_{L^{p}(\nu)} < \infty.$$

is satisfied when

$$\sup_{N,m\in\mathbb{Z}}\left\|\sum_{n=1}^{N}\prod_{i=1}^{H}f_{i}\circ T_{i}^{n+m}\right\|_{L^{p}(\mu)}<\infty.$$

This last condition is equivalent to

$$\sup_{N\in\mathbb{Z}}\left\|\sum_{n=0}^{N}\prod_{i=1}^{H}f_{i}\circ T_{i}^{n}\right\|_{L^{p}(\mu)}<\infty.$$

which may be easier to check in the applications.

## References

[1] **I.Assani**: "A note on the equation y = (I - T)x in  $L^1$ " Illinois J. of Math. vol 43, 3, (1999) p.540-541.

[2] I. Assani:"Pointwise recurrence for commuting measure preserving transformations" Preprint , arXiv:1312.5270v2, 2015

[3] J. Komlos:"A generalization of a problem of Steinhaus" Acta Math. Acad. Sci. Hungar. 18:217-229, 1967.