

Squares of Tribonacci numbers*

Kunle Adegoke †

Department of Physics and Engineering Physics,
Obafemi Awolowo University, 220005 Ile-Ife, Nigeria

Abstract

We prove some identities for the squares of generalized Tribonacci numbers. Various summation identities involving these numbers are derived.

Contents

1	Introduction	1
2	Squares of Tribonacci numbers	3
3	Partial sums	4
4	Generating function	7
5	Cubes of Tribonacci numbers	7

1 Introduction

For $r \geq 3$, the generalized Tribonacci numbers are defined by

$$\mathcal{T}_r = \mathcal{T}_{r-1} + \mathcal{T}_{r-2} + \mathcal{T}_{r-3}, \quad (1.1)$$

with arbitrary initial values \mathcal{T}_0 , \mathcal{T}_1 and \mathcal{T}_2 .

By substituting $\mathcal{T}_{r-2} = \mathcal{T}_{r-1} - \mathcal{T}_{r-3} - \mathcal{T}_{r-4}$ into the recurrence relation (1.1), a useful alternative recurrence relation is obtained for $r \geq 4$:

$$\mathcal{T}_r = 2\mathcal{T}_{r-1} - \mathcal{T}_{r-4}, \quad (1.2)$$

with arbitrary initial values \mathcal{T}_0 , \mathcal{T}_1 , \mathcal{T}_2 and \mathcal{T}_3 .

Extension of the definition of \mathcal{T}_r to negative subscripts is provided by writing the recurrence relation (1.2) as

$$\mathcal{T}_{-r} = 2\mathcal{T}_{-r+3} - \mathcal{T}_{-r+4}. \quad (1.3)$$

*AMS Classification: 11B37, 11B39, 65B10

†adegoke00@gmail.com

When $\mathcal{T}_0 = 0$, $\mathcal{T}_1 = 1$ and $\mathcal{T}_2 = 1$, we have the Tribonacci sequence, denoted $\{\mathcal{T}_r\}$, $r \in \mathbb{Z}$.

Our purpose in writing this note is to establish the following identities:

$$\begin{aligned}\mathcal{T}_r^2 - 2\mathcal{T}_{r-1}^2 - 3\mathcal{T}_{r-2}^2 - 6\mathcal{T}_{r-3}^2 + \mathcal{T}_{r-4}^2 + \mathcal{T}_{r-6}^2 &= 0, \\ \mathcal{T}_{r+2}^2 - 4\mathcal{T}_{r+1}^2 + \mathcal{T}_r^2 + 14\mathcal{T}_{r-2}^2 - 4\mathcal{T}_{r-3}^2 - 2\mathcal{T}_{r-4}^2 - 8\mathcal{T}_{r-5}^2 + \mathcal{T}_{r-6}^2 + \mathcal{T}_{r-8}^2 &= 0, \\ \mathcal{T}_{r+3}^2 - 3\mathcal{T}_{r+2}^2 - 4\mathcal{T}_r^2 + 2\mathcal{T}_{r-1}^2 - 10\mathcal{T}_{r-2}^2 - 4\mathcal{T}_{r-3}^2 + \mathcal{T}_{r-5}^2 + \mathcal{T}_{r-6}^2 &= 0, \\ \mathcal{T}_{r+1}^2 - 4\mathcal{T}_r^2 + 2\mathcal{T}_{r-2}^2 + 16\mathcal{T}_{r-3}^2 + 4\mathcal{T}_{r-4}^2 - 2\mathcal{T}_{r-6}^2 - \mathcal{T}_{r-7}^2 &= 0\end{aligned}$$

and

$$\mathcal{T}_{r+2}^2 - 2\mathcal{T}_{r+1}^2 - 2\mathcal{T}_r^2 - 8\mathcal{T}_{r-1}^2 - 2\mathcal{T}_{r-2}^2 - 6\mathcal{T}_{r-3}^2 + 2\mathcal{T}_{r-4}^2 + \mathcal{T}_{r-6}^2 = 0.$$

We will also develop the associated summation identities. Specifically, we shall evaluate

$$\begin{aligned}\sum_{j=0}^k x^j \mathcal{T}_j^2, \quad \sum_{j=0}^k \mathcal{T}_{2j}^2, \quad \sum_{j=1}^k \mathcal{T}_{2j-1}^2, \quad \sum_{j=0}^k (-1)^j \mathcal{T}_{2j}^2, \quad \sum_{j=1}^k (-1)^{j-1} \mathcal{T}_{2j-1}^2, \\ \sum_{j=0}^k \mathcal{T}_{4j}^2, \quad \sum_{j=1}^k \mathcal{T}_{4j-3}^2, \quad \sum_{j=0}^k \mathcal{T}_{4j-1}^2, \quad \sum_{j=0}^k j \mathcal{T}_j^2, \quad \sum_{j=0}^k j^2 \mathcal{T}_j^2 \quad \text{and} \quad \sum_{j=0}^{\infty} x^j \mathcal{T}_j^2.\end{aligned}$$

Presently we derive some identities that we shall need.

Rearranging the identity (1.2) and squaring, we have

$$4\mathcal{T}_{r-1}\mathcal{T}_r = 4\mathcal{T}_{r-1}^2 - \mathcal{T}_{r-4}^2 + \mathcal{T}_r^2, \quad (1.4)$$

$$4\mathcal{T}_{r-1}\mathcal{T}_{r-4} = 4\mathcal{T}_{r-1}^2 + \mathcal{T}_{r-4}^2 - \mathcal{T}_r^2 \quad (1.5)$$

and

$$2\mathcal{T}_r\mathcal{T}_{r-4} = 4\mathcal{T}_{r-1}^2 - \mathcal{T}_{r-4}^2 - \mathcal{T}_r^2. \quad (1.6)$$

Rearranging identity (1.2) and multiplying through by $4\mathcal{T}_{r-3}$ to obtain

$$8\mathcal{T}_{r-1}\mathcal{T}_{r-3} = 4\mathcal{T}_r\mathcal{T}_{r-3} + 4\mathcal{T}_{r-4}\mathcal{T}_{r-3}, \quad (1.7)$$

and using identities (1.4) and (1.5) to resolve the right hand side gives

$$8\mathcal{T}_{r-1}\mathcal{T}_{r-3} = 4\mathcal{T}_r^2 + 2\mathcal{T}_{r-3}^2 - \mathcal{T}_{r+1}^2 + 4\mathcal{T}_{r-4}^2 - \mathcal{T}_{r-7}^2. \quad (1.8)$$

We also require the following results from a previous paper [1].

Lemma 1 (Partial sum of an n^{th} order sequence). *Let $\{X_j\}$ be any arbitrary sequence, where X_j , $j \in \mathbb{Z}$, satisfies a n^{th} order recurrence relation $X_j = f_1 X_{j-c_1} + f_2 X_{j-c_2} + \dots + f_n X_{j-c_n} = \sum_{m=1}^n f_m X_{j-c_m}$, where f_1, f_2, \dots, f_n are arbitrary non-vanishing complex functions, not dependent on j , and c_1, c_2, \dots, c_n are fixed integers. Then, the following summation identity holds for arbitrary x and non-negative integer k :*

$$\sum_{j=0}^k x^j X_j = \frac{\sum_{m=1}^n \left\{ x^{c_m} f_m \left(\sum_{j=1}^{c_m} x^{-j} X_{-j} - \sum_{j=k-c_m+1}^k x^j X_j \right) \right\}}{1 - \sum_{m=1}^n x^{c_m} f_m}.$$

Lemma 2 (Generating function). *Under the conditions of Lemma 1, if additionally $x^k X_k$ vanishes in the limit as k approaches infinity, then*

$$S_\infty(x) = \sum_{j=0}^{\infty} x^j X_j = \frac{\sum_{m=1}^n \left(x^{c_m} f_m \sum_{j=1}^{c_m} x^{-j} X_{-j} \right)}{1 - \sum_{m=1}^n x^{c_m} f_m},$$

so that $S_\infty(x)$ is a generating function for the sequence $\{X_j\}$.

We note that a special case of Lemma 1 was proved in [5].

2 Squares of Tribonacci numbers

Theorem 1. *The following identities hold for any integer r :*

$$\mathcal{T}_r^2 - 2\mathcal{T}_{r-1}^2 - 3\mathcal{T}_{r-2}^2 - 6\mathcal{T}_{r-3}^2 + \mathcal{T}_{r-4}^2 + \mathcal{T}_{r-6}^2 = 0, \quad (2.1)$$

$$\mathcal{T}_{r+2}^2 - 4\mathcal{T}_{r+1}^2 + \mathcal{T}_r^2 + 14\mathcal{T}_{r-2}^2 - 4\mathcal{T}_{r-3}^2 - 2\mathcal{T}_{r-4}^2 - 8\mathcal{T}_{r-5}^2 + \mathcal{T}_{r-6}^2 + \mathcal{T}_{r-8}^2 = 0, \quad (2.2)$$

$$\mathcal{T}_{r+3}^2 - 3\mathcal{T}_{r+2}^2 - 4\mathcal{T}_r^2 + 2\mathcal{T}_{r-1}^2 - 10\mathcal{T}_{r-2}^2 - 4\mathcal{T}_{r-3}^2 + \mathcal{T}_{r-5}^2 + \mathcal{T}_{r-6}^2 = 0, \quad (2.3)$$

$$\mathcal{T}_{r+1}^2 - 4\mathcal{T}_r^2 + 2\mathcal{T}_{r-2}^2 + 16\mathcal{T}_{r-3}^2 + 4\mathcal{T}_{r-4}^2 - 2\mathcal{T}_{r-6}^2 - \mathcal{T}_{r-7}^2 = 0 \quad (2.4)$$

and

$$\mathcal{T}_{r+2}^2 - 2\mathcal{T}_{r+1}^2 - 2\mathcal{T}_r^2 - 8\mathcal{T}_{r-1}^2 - 2\mathcal{T}_{r-2}^2 - 6\mathcal{T}_{r-3}^2 + 2\mathcal{T}_{r-4}^2 + \mathcal{T}_{r-6}^2 = 0. \quad (2.5)$$

Proof. To prove identity (2.1), write

$$\mathcal{T}_r - \mathcal{T}_{r-1} = \mathcal{T}_{r-2} + \mathcal{T}_{r-3}, \quad (2.6)$$

square both sides and use the identity (1.4) to resolve the cross-products $\mathcal{T}_r \mathcal{T}_{r-1}$ and $\mathcal{T}_{r-2} \mathcal{T}_{r-3}$.

Identity (2.2) is proved by multiplying through identity (1.2) by $8\mathcal{T}_{r-2}$, obtaining $8\mathcal{T}_r \mathcal{T}_{r-2} = 16\mathcal{T}_{r-1} \mathcal{T}_{r-2} - 8\mathcal{T}_{r-4} \mathcal{T}_{r-2}$ and using identities (1.4) and (1.8) to resolve the cross-products.

To prove identity (2.3), rearrange relation (2.6) as $\mathcal{T}_r - \mathcal{T}_{r-2} = \mathcal{T}_{r-1} + \mathcal{T}_{r-3}$, square both sides and use the identity (1.8) to resolve the cross-products.

Identity (2.4) is proved by rearranging relation (2.6) as $\mathcal{T}_r - \mathcal{T}_{r-3} = \mathcal{T}_{r-1} + \mathcal{T}_{r-2}$, multiplying through by \mathcal{T}_{r-3} to obtain $\mathcal{T}_r \mathcal{T}_{r-3} - \mathcal{T}_{r-3}^2 = \mathcal{T}_{r-1} \mathcal{T}_{r-3} + \mathcal{T}_{r-2} \mathcal{T}_{r-3}$ and using identities (1.4), (1.5) and (1.8) to resolve the cross-products.

To prove identity (2.5) multiply through the recurrence relation (1.1) by \mathcal{T}_r to obtain $\mathcal{T}_r^2 - \mathcal{T}_r \mathcal{T}_{r-1} - \mathcal{T}_r \mathcal{T}_{r-2} - \mathcal{T}_r \mathcal{T}_{r-3} = 0$. Now use the identities (1.4), (1.5) and (1.8) to resolve the cross-products. □

3 Partial sums

From the identity (2.1) of Theorem 1 and using $\{X_r\} \equiv \{\mathcal{T}_r^2\}$, $r \in \mathbb{Z}$, in Lemma 1, the next result follows.

Theorem 2. *The following identity holds for integer k and arbitrary x :*

$$\begin{aligned}
& (1 - 3x - x^2 - x^3)(1 + x + x^2 - x^3) \sum_{j=0}^k x^j \mathcal{T}_j^2 \\
&= (2 + 3x + 6x^2 - x^3 - x^5)(\mathcal{T}_{-1}^2 - x^{k+1} \mathcal{T}_k^2) \\
&\quad + (3 + 6x - x^2 - x^4)(\mathcal{T}_{-2}^2 - x^{k+1} \mathcal{T}_{k-1}^2) \\
&\quad + (6 - x - x^3)(\mathcal{T}_{-3}^2 - x^{k+1} \mathcal{T}_{k-2}^2) \\
&\quad - (1 + x^2)(\mathcal{T}_{-4}^2 - x^{k+1} \mathcal{T}_{k-3}^2) \\
&\quad - x(\mathcal{T}_{-5}^2 - x^{k+1} \mathcal{T}_{k-4}^2) - (\mathcal{T}_{-6}^2 - x^{k+1} \mathcal{T}_{k-5}^2).
\end{aligned}$$

In particular,

$$\begin{aligned}
8 \sum_{j=0}^k \mathcal{T}_j^2 &= 9(\mathcal{T}_k^2 - \mathcal{T}_{-1}^2) + 7(\mathcal{T}_{k-1}^2 - \mathcal{T}_{-2}^2) + 4(\mathcal{T}_{k-2}^2 - \mathcal{T}_{-3}^2) \\
&\quad - 2(\mathcal{T}_{k-3}^2 - \mathcal{T}_{-4}^2) - (\mathcal{T}_{k-4}^2 - \mathcal{T}_{-5}^2) - (\mathcal{T}_{k-5}^2 - \mathcal{T}_{-6}^2)
\end{aligned} \tag{3.1}$$

and

$$\begin{aligned}
8 \sum_{j=0}^k (-1)^j \mathcal{T}_j^2 &= 7(\mathcal{T}_{-1}^2 + (-1)^k \mathcal{T}_k^2) - 5(\mathcal{T}_{-2}^2 + (-1)^k \mathcal{T}_{k-1}^2) \\
&\quad + 8(\mathcal{T}_{-3}^2 + (-1)^k \mathcal{T}_{k-2}^2) - 2(\mathcal{T}_{-4}^2 + (-1)^k \mathcal{T}_{k-3}^2) \\
&\quad + (\mathcal{T}_{-5}^2 + (-1)^k \mathcal{T}_{k-4}^2) - (\mathcal{T}_{-6}^2 + (-1)^k \mathcal{T}_{k-5}^2).
\end{aligned} \tag{3.2}$$

A variant of (3.1) was given by Maiorano [2].

For the Tribonacci numbers, these results translate to:

$$\begin{aligned}
& (1 - 3x - x^2 - x^3)(1 + x + x^2 - x^3) \sum_{j=0}^k x^j T_j^2 \\
&= -(2 + 3x + 6x^2 - x^3 - x^5)x^{k+1} T_k^2 \\
&\quad + (3 + 6x - x^2 - x^4)(1 - x^{k+1} T_{k-1}^2) \\
&\quad + (6 - x - x^3)(1 - x^{k+1} T_{k-2}^2) \\
&\quad + (1 + x^2)x^{k+1} T_{k-3}^2 - x(4 - x^{k+1} T_{k-4}^2) \\
&\quad - (9 - x^{k+1} T_{k-5}^2),
\end{aligned} \tag{3.3}$$

$$8 \sum_{j=0}^k T_j^2 = 9T_k^2 + 7T_{k-1}^2 + 4T_{k-2}^2 - 2T_{k-3}^2 - T_{k-4}^2 - T_{k-5}^2 + 2 \tag{3.4}$$

and

$$8 \sum_{j=0}^k (-1)^j T_j^2 = (-1)^k (7T_k^2 - 5T_{k-1}^2 + 8T_{k-2}^2 - 2T_{k-3}^2 + T_{k-4}^2 - T_{k-5}^2) - 2. \quad (3.5)$$

Shah [3] proved a shorter version of (3.4) and reported a variant of same, attributed to Zeitlin [4].

Combining identity (3.4) and identity (3.5), and using the summation identity

$$\sum_{j=0}^{2k} f_j = \sum_{j=0}^k f_{2j} + \sum_{j=1}^k f_{2j-1}, \quad (3.6)$$

we obtain

$$8 \sum_{j=0}^k T_{2j}^2 = 8T_{2k}^2 + T_{2k-1}^2 + 6T_{2k-2}^2 - 2T_{2k-3}^2 - T_{2k-5}^2 \quad (3.7)$$

and

$$8 \sum_{j=1}^k T_{2j-1}^2 = T_{2k}^2 + 6T_{2k-1}^2 - 2T_{2k-2}^2 - T_{2k-4}^2 + 2. \quad (3.8)$$

Using $f_j = i^j T_j^2$ in identity (3.6), where $i = \sqrt{-1}$ is the imaginary unit, allows us to write

$$\sum_{j=0}^k (-1)^j T_{2j}^2 = \operatorname{Re} \sum_{j=0}^{2k} i^j T_j^2, \quad \sum_{j=1}^k (-1)^{j-1} T_{2j-1}^2 = \operatorname{Im} \sum_{j=0}^{2k} i^j T_j^2. \quad (3.9)$$

Using $x = i$ in identity (3.3) and taking note of identity (3.9), we obtain the following alternating versions of identities (3.7) and (3.8):

$$8 \sum_{j=0}^k (-1)^j T_{2j}^2 = (-1)^k (7T_{2k}^2 + 3T_{2k-1}^2 - 6T_{2k-2}^2 - T_{2k-4}^2 + T_{2k-5}^2) + 2 \quad (3.10)$$

and

$$8 \sum_{j=1}^k (-1)^{j-1} T_{2j-1}^2 = (-1)^k (T_{2k}^2 - 9T_{2k-1}^2 - 6T_{2k-2}^2 + T_{2k-4}^2 + T_{2k-5}^2) + 2. \quad (3.11)$$

Replacing k with $2k$ in identities (3.7) and (3.10) we find

$$16 \sum_{j=0}^k T_{4j}^2 = 15T_{4k}^2 + 4T_{4k-1}^2 - 2T_{4k-3}^2 - T_{4k-4}^2 + 2, \quad (3.12)$$

by their addition, and

$$16 \sum_{j=1}^k T_{4j-2}^2 = T_{4k}^2 - 2T_{4k-1}^2 + 12T_{4k-2}^2 - 2T_{4k-3}^2 + T_{4k-4}^2 - 2T_{4k-5}^2 - 2, \quad (3.13)$$

by their subtraction.

Doing similar analysis with identities (3.8) and (3.11), we find

$$16 \sum_{j=1}^k T_{4j-3}^2 = 2T_{4k}^2 - 3T_{4k-1}^2 - 8T_{4k-2}^2 + T_{4k-5}^2 + 4 \quad (3.14)$$

and

$$16 \sum_{j=0}^k T_{4j-1}^2 = 15T_{4k-1}^2 + 4T_{4k-2}^2 - 2T_{4k-4}^2 - T_{4k-5}^2. \quad (3.15)$$

Weighted sums of the form $\sum_{j=0}^k j^p T_j^2$, p a non-negative integer, can be evaluated by making $\sum_{j=0}^k x^j T_j^2$ subject in the identity of Theorem 2, setting $x = e^y$, differentiating p times with respect to y and then setting $y = 0$. The simplest of such evaluations are the following:

$$\begin{aligned} 8 \sum_{j=0}^k j T_j^2 &= (9k - 2)\mathcal{T}_k^2 + 7(k - 1)\mathcal{T}_{k-1}^2 + 4(k - 2)\mathcal{T}_{k-2}^2 \\ &\quad - 2k\mathcal{T}_{k-3}^2 - k\mathcal{T}_{k-4}^2 - (k - 1)\mathcal{T}_{k-5}^2 \\ &\quad + 11\mathcal{T}_{-1}^2 + 14\mathcal{T}_{-2}^2 + 12\mathcal{T}_{-3}^2 \\ &\quad - 2\mathcal{T}_{-4}^2 - \mathcal{T}_{-5}^2 - 2\mathcal{T}_{-6}^2 \end{aligned} \quad (3.16)$$

and

$$\begin{aligned} 8 \sum_{j=0}^k j^2 T_j^2 &= (9k^2 - 4k + 6)\mathcal{T}_k^2 + (7k^2 - 14k + 7)\mathcal{T}_{k-1}^2 \\ &\quad + (4k^2 - 16k + 10)\mathcal{T}_{k-2}^2 - (2k^2 + 6)\mathcal{T}_{k-3}^2 \\ &\quad - (k^2 + 2)\mathcal{T}_{k-4}^2 - (k^2 - 2k + 3)\mathcal{T}_{k-5}^2 \\ &\quad - 19\mathcal{T}_{-1}^2 - 28\mathcal{T}_{-2}^2 - 30\mathcal{T}_{-3}^2 \\ &\quad + 8\mathcal{T}_{-4}^2 + 3\mathcal{T}_{-5}^2 + 6\mathcal{T}_{-6}^2. \end{aligned} \quad (3.17)$$

In particular,

$$\begin{aligned} 8 \sum_{j=0}^k j T_j^2 &= (9k - 2)T_k^2 + 7(k - 1)T_{k-1}^2 + 4(k - 2)T_{k-2}^2 \\ &\quad - 2kT_{k-3}^2 - kT_{k-4}^2 - (k - 1)T_{k-5}^2 + 4 \end{aligned} \quad (3.18)$$

and

$$\begin{aligned} 8 \sum_{j=0}^k j^2 T_j^2 &= (9k^2 - 4k + 6)T_k^2 + (7k^2 - 14k + 7)T_{k-1}^2 \\ &\quad + (4k^2 - 16k + 10)T_{k-2}^2 - (2k^2 + 6)T_{k-3}^2 \\ &\quad - (k^2 + 2)T_{k-4}^2 - (k^2 - 2k + 3)T_{k-5}^2 + 8. \end{aligned}$$

4 Generating function

Theorem 3. Let $\mathcal{G}(x) = \sum_{j=0}^{\infty} x^j \mathcal{T}_j^2$ be the generating function of the squares of generalized Tribonacci numbers. $\mathcal{G}(x)$ is given by

$$\begin{aligned} & (1 - 3x - x^2 - x^3)(1 + x + x^2 - x^3)\mathcal{G}(x) \\ &= (2 + 3x + 6x^2 - x^3 - x^5)\mathcal{T}_{-1}^2 \\ &+ (3 + 6x - x^2 - x^4)\mathcal{T}_{-2}^2 + (6 - x - x^3)\mathcal{T}_{-3}^2 \\ &- (1 + x^2)\mathcal{T}_{-4}^2 - x\mathcal{T}_{-5}^2 - \mathcal{T}_{-6}^2, \end{aligned}$$

provided $x^k \mathcal{T}_k^2$ vanishes as k approaches infinity.

In particular, we have that $G(x)$, the generating function of the squares of Tribonacci numbers, is given by

$$G(x) = \sum_{j=0}^{\infty} x^j T_j^2 = \frac{x(1 - x - x^2 - x^3)}{(1 - 3x - x^2 - x^3)(1 + x + x^2 - x^3)}, \quad (4.1)$$

valid for x for which $x^k T_k^2$ vanishes as k approaches infinity.

5 Cubes of Tribonacci numbers

We end this note by giving an identity for the cubes of generalized Tribonacci numbers. The identity can be proved by expressing each \mathcal{T}_{r-j} , $j \in \{1, 2, 3, 4, 5, 6, 7, 9\}$ as a linear combination of \mathcal{T}_r , \mathcal{T}_{r-8} and \mathcal{T}_{r-10} and substituting into the identity of Theorem 4.

Theorem 4. The following identity holds for integer r :

$$\begin{aligned} & \mathcal{T}_r^3 - 4\mathcal{T}_{r-1}^3 - 9\mathcal{T}_{r-2}^3 - 34\mathcal{T}_{r-3}^3 + 24\mathcal{T}_{r-4}^3 - 2\mathcal{T}_{r-5}^3 \\ &+ 40\mathcal{T}_{r-6}^3 - 14\mathcal{T}_{r-7}^3 - \mathcal{T}_{r-8}^3 - 2\mathcal{T}_{r-9}^3 + \mathcal{T}_{r-10}^3 = 0. \end{aligned}$$

References

- [1] K. Adegoke, Weighted Tribonacci sums, *arXiv:1804.06449[math.CA]* (2018).
- [2] P. J. Maiorano, Sum of squares of tribonacci numbers, *The Mathematics Teacher* **89:7** (1996), 591.
- [3] D. V. Shah, Some Tribonacci identities, *Mathematics Today* **27** (2011), 1–9.
- [4] D. Zeitlin, On summation formulas and identities for Fibonacci numbers, *The Fibonacci Quarterly* **5:1** (1967), 1–43.
- [5] D. Zeitlin, On the sums $\sum_{k=0}^n k^p$ and $\sum_{k=0}^n (-1)^k k^p$, *Proceedings of the American Mathematical Society* **2** (1964), 105–107.