

SOBOLEV-LIKE HILBERT SPACES INDUCED BY ELLIPTIC OPERATORS

TETIANA KASIRENKO, VLADIMIR MIKHAILETS, AND ALEKSANDR MURACH

ABSTRACT. We investigate properties of function spaces induced by the inner product Sobolev spaces $H^s(\Omega)$ over a bounded Euclidean domain Ω and by an elliptic differential operator A on $\overline{\Omega}$. The domain and the coefficients of A are of the class C^∞ . These spaces consist of all distributions $u \in H^s(\Omega)$ such that $Au \in H^\lambda(\Omega)$ and are endowed with the corresponding graph norm, with $s, \lambda \in \mathbb{R}$. We prove an interpolation formula for these spaces and discuss their application to elliptic boundary-value problems.

1. INTRODUCTION

This paper is devoted to the function spaces induced by the scale $\{H^s(\Omega) : s \in \mathbb{R}\}$ of inner product Sobolev spaces over a bounded Euclidean domain Ω and by an elliptic partial differential operator A on $\overline{\Omega}$. The boundary of Ω and the coefficients of A are supposed to be infinitely smooth. We investigate the inner product spaces $H_{A,\lambda}^s(\Omega)$, where $s, \lambda \in \mathbb{R}$, that consist of all distributions $u \in H^s(\Omega)$ subject to the condition $Au \in H^\lambda(\Omega)$ and that are endowed with the corresponding graph norm. We call $H_{A,\lambda}^s(\Omega)$ the Sobolev-like space induced by A . Such spaces are used in the theory of elliptic differential operators and elliptic boundary-value problems.

Specifically, the space $H_{A,0}^0(\Omega)$ is the domain of the maximal operator that corresponds to the unbounded operator A defined on $C^\infty(\overline{\Omega})$ and acting in $L_2(\Omega)$ [1, 2, 3]. Lions and Magenes [4, 5] systematically use the space $H_{A,0}^s(\Omega)$ for $s < \text{ord } A =: 2q$ in the theory of solvability of regular elliptic boundary-value problems in distribution spaces. In the case of the Dirichlet problem, the space $H_{A,-q}^s(\Omega)$ for $s < q$ is also useful [6]. These spaces serve as domains of Fredholm bounded operators that correspond to elliptic problems. A more general class of the spaces $H_{A,\lambda}^s(\Omega)$ is considered in [7, 8]. Some of their versions are introduced for L_p -Sobolev spaces [4, 5, 9, 10], weighted Sobolev spaces [10, 11, 12], and generalized Sobolev spaces [8, 13, 14] and are also used in the theory of elliptic differential equations.

The purpose of this paper is to investigate some properties of the spaces $H_{A,\lambda}^s(\Omega)$ concerning their completeness, separability, and dense subsets. Besides, we will prove an interpolation formula for these spaces. We also give their application to elliptic boundary-value problems.

2010 *Mathematics Subject Classification.* 46E35, 35J30, 35J40.

Key words and phrases. Sobolev space, elliptic operator, complex interpolation, elliptic problem, Fredholm operator.

2. MAIN RESULTS

Let $1 \leq n \in \mathbb{Z}$ and $s \in \mathbb{R}$. We let $H^s(\mathbb{R}^n)$ denote the complex inner product Sobolev space of order s over \mathbb{R}^n . By definition, this space consists of all tempered distributions w in \mathbb{R}^n that their Fourier transform \widehat{w} is locally Lebesgue integrable over \mathbb{R}^n and satisfies the condition

$$\|w\|_{s,\mathbb{R}^n}^2 := \int_{\mathbb{R}^n} (1 + |\xi|^2)^s |\widehat{w}(\xi)|^2 d\xi < \infty.$$

The space $H^s(\mathbb{R}^n)$ is Hilbert and separable with respect to the norm $\|\cdot\|_{s,\mathbb{R}^n}$. If Ω is an open subset of \mathbb{R}^n , then the Sobolev space $H^s(\Omega)$ of order s over Ω is defined to consist of the restrictions $u := w \upharpoonright \Omega$ of all distributions $w \in H^s(\mathbb{R}^n)$ to Ω . This space is Hilbert and separable with respect to the norm

$$\|u\|_{s,\Omega} := \inf \{ \|w\|_{s,\mathbb{R}^n} : w \in H^s(\mathbb{R}^n), u = w \upharpoonright \Omega \}$$

and is continuously embedded in the linear topological space $\mathcal{D}'(\Omega)$ of all distributions in Ω .

Henceforth we suppose that $n \geq 2$ and that Ω is a bounded domain whose boundary $\Gamma := \partial\Omega$ is an infinitely smooth closed manifold of dimension $n - 1$, the C^∞ -structure on Γ being induced by \mathbb{R}^n . In the closed domain $\overline{\Omega} := \Omega \cup \Gamma$, we consider an arbitrary elliptic linear partial differential expression

$$A := \sum_{|\mu| \leq 2q} a_\mu(x) \frac{\partial^{|\mu|}}{\partial x_1^{\mu_1} \dots \partial x_n^{\mu_n}}$$

of an even order $2q \geq 2$ with complex-valued coefficients $a_\mu \in C^\infty(\overline{\Omega})$. Here, $\mu := (\mu_1, \dots, \mu_n)$ is a multi-index with nonnegative integer-valued components, $|\mu| := \mu_1 + \dots + \mu_n$, and $x = (x_1, \dots, x_n)$ is an arbitrary point in \mathbb{R}^n . The ellipticity condition for A means that

$$\sum_{|\mu|=2q} a_\mu(x) \xi_1^{\mu_1} \dots \xi_n^{\mu_n} \neq 0$$

for every point $x \in \overline{\Omega}$ and each vector $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n \setminus \{0\}$. In the $n = 2$ case, we suppose in addition that A is properly elliptic on $\overline{\Omega}$ (see, e.g., [15, Section 5.2.1]).

We associate some spaces with the elliptic formal differential operator A . Choosing $s, \lambda \in \mathbb{R}$ arbitrarily, we let $H_{A,\lambda}^s(\Omega)$ denote the linear space

$$\{u \in H^s(\Omega) : Au \in H^\lambda(\Omega)\}$$

endowed with the graph norm

$$\|u\|_{s,A,\lambda} := (\|u\|_{s,\Omega}^2 + \|Au\|_{\lambda,\Omega}^2)^{1/2}.$$

Here and below, Au is understood in the sense of the theory of distributions in Ω .

Theorem 1. *Let $s, \lambda \in \mathbb{R}$. The following assertions are true:*

- (i) *The space $H_{A,\lambda}^s(\Omega)$ is Hilbert and separable.*

- (ii) *The equality of linear spaces $H_{A,\lambda}^s(\Omega)$ and $H^s(\Omega)$ holds if and only if $\lambda \leq s - 2q$. If $\lambda \leq s - 2q$, the norms in these spaces are equivalent.*
- (iii) *The set $C^\infty(\overline{\Omega})$ is dense in $H_{A,\lambda}^s(\Omega)$.*
- (iv) *If $s \leq 1/2$ and $\lambda \leq 1/2 - 2q$, then the set $C_0^\infty(\Omega)$ is dense in $H_{A,\lambda}^s(\Omega)$.*

Here and below, $C_0^\infty(\Omega)$ denotes the set of all functions from $C^\infty(\overline{\Omega})$ that vanish near the boundary of Ω .

Note that, if $\lambda > s - 2q$, the space $H_{A,\lambda}^s(\Omega)$ depends on the coefficients of A , even when all of them are constant. For example, if A_1 and A_2 are elliptic constant-coefficient linear differential expressions, the embedding $H_{A_1,0}^0(\Omega) \subseteq H_{A_2,0}^0(\Omega)$ implies that $A_2 = \alpha A_1 + \beta$ for certain $\alpha, \beta \in \mathbb{C}$. This follows from Hörmander's result [2, Theorem 3.1].

It is well known that the Sobolev scale $\{H^s(\Omega) : s \in \mathbb{R}\}$ has the following interpolation property: if $s_0, s_1 \in \mathbb{R}$ and $0 < \theta < 1$, then

$$(1) \quad [H^{s_0}(\Omega), H^{s_1}(\Omega)]_\theta = H^{(1-\theta)s_0 + \theta s_1}(\Omega)$$

up to equivalence of norms (see, e.g., [15, Section 4.3.1, Theorem 1]). Here and below, $[H_0, H_1]_\theta$ denotes the result of the complex interpolation with the parameter θ of a compatible pair of Hilbert (or, more generally, Banach) spaces H_0 and H_1 (see, e.g., [15, Section 1.9]). The pair $[H_0, H_1]$ of these spaces is called compatible if they are continuously embedded in a certain linear Hausdorff space.

Let us formulate a version of the interpolation property (1) for the spaces $H_{A,\lambda_j}^{s_j}(\Omega)$, where $j \in \{0, 1\}$. In view of assertion (ii) of Theorem 1, it is worthwhile to restrict ourselves to the case when $\lambda_j \geq s_j - 2q$.

Theorem 2. *Suppose that $s_0, s_1, \lambda_0, \lambda_1$, and θ are real numbers satisfying the inequalities $\lambda_0 \geq s_0 - 2q$, $\lambda_1 \geq s_1 - 2q$, and $0 < \theta < 1$. Put $s := (1 - \theta)s_0 + \theta s_1$ and $\lambda := (1 - \theta)\lambda_0 + \theta\lambda_1$. Then*

$$(2) \quad [H_{A,\lambda_0}^{s_0}(\Omega), H_{A,\lambda_1}^{s_1}(\Omega)]_\theta = H_{A,\lambda}^s(\Omega)$$

up to equivalence of norms.

3. PROOF OF THE MAIN RESULTS

We will give a joint proof of Theorems 1 and 2.

Let us first prove that the space $H_{A,\lambda}^s(\Omega)$ is Hilbert for arbitrary $s, \lambda \in \mathbb{R}$. Evidently, the norm in it is induced by a graph inner product. Besides, the space $H_{A,\lambda}^s(\Omega)$ is complete. Indeed, if (u_k) is a Cauchy sequence in this space, then there exist the limits $u := \lim u_k$ in $H^s(\Omega)$ and $f := \lim Au_k$ in $H^\lambda(\Omega)$ because the last two spaces are complete. The differential operator A is continuous on $\mathcal{D}'(\Omega)$; therefore, $Au = \lim Au_k = f$ in $\mathcal{D}'(\Omega)$. Here, $u \in H^s(\Omega)$ and $f \in H^\lambda(\Omega)$. Hence, $u \in H_{A,\lambda}^s(\Omega)$, and $\lim u_k = u$ in the space $H_{A,\lambda}^s(\Omega)$. Thus, this space is complete.

Let us now prove assertion (ii) of Theorem 1. Since $\text{ord } A = 2q$, the differential operator A acts continuously from $H^s(\Omega)$ to $H^{s-2q}(\Omega)$ for every $s \in \mathbb{R}$. Hence, if

$\lambda \leq s - 2q$, then $H^s(\Omega) = H_{A,\lambda}^s(\Omega)$ up to equivalence of norms due to the continuous embedding $H^{s-2q}(\Omega) \subseteq H^\lambda(\Omega)$.

Conversely, assume that $H^s(\Omega) = H_{A,\lambda}^s(\Omega)$ for certain $s, \lambda \in \mathbb{R}$. Choose an arbitrary distribution $w \in H^s(\mathbb{R}^n)$ such that $\text{supp } w \subset \Omega$, and put $u := w \upharpoonright \Omega$. Since $u \in H^s(\Omega)$, we have the inclusion $Au \in H^\lambda(\Omega)$ by our assumption. Hence, $\chi u \in H^{\lambda+2q}(\Omega)$ for every function $\chi \in C_0^\infty(\Omega)$ by virtue of the ellipticity of A (see, e.g., [16, Theorem 7.4.1]). We can choose the function χ so that $\chi = 1$ on $\text{supp } w$. Therefore $u \in H^{\lambda+2q}(\Omega)$, which implies the inclusion $w \in H^{\lambda+2q}(\mathbb{R}^n)$. Thus,

$$\{w \in H^s(\mathbb{R}^n) : \text{supp } w \subset \Omega\} \subseteq H^{\lambda+2q}(\mathbb{R}^n).$$

According to [16, Theorem 2.2.2], this embedding implies that

$$(1 + |\xi|)^{\lambda+2q} \leq c(1 + |\xi|)^s$$

for every $\xi \in \mathbb{R}^n$ with some number $c > 0$ not depending on ξ . Hence, $\lambda + 2q \leq s$. We have proved assertion (ii) of Theorem 1.

To prove Theorem 2 and the remaining part of Theorem 1, we use a theorem on interpolation of some Hilbert spaces induced by linear continuous operators. Before we formulate this result, let us make some notation. If H , Φ and Ψ are Hilbert spaces subject to the continuous embedding $\Phi \subseteq \Psi$ and if $T : H \rightarrow \Psi$ is a continuous linear operator, we put

$$(H)_{T,\Phi} := \{u \in H : Tu \in \Phi\}$$

and endow the linear space $(H)_{T,\Phi}$ with the graph norm

$$\|u\|_{(H)_{T,\Phi}} := (\|u\|_H^2 + \|Tu\|_\Phi^2)^{1/2}.$$

This norm does not depend on Ψ , and the space $(H)_{T,\Phi}$ is Hilbert. The latter is proved in a quite similar way as the proof of the completeness of $H_{A,\lambda}^s(\Omega)$.

Proposition 1. *Assume that six separable Hilbert spaces X_0, Y_0, Z_0, X_1, Y_1 , and Z_1 and three linear mappings T, R , and S are given and satisfy the following conditions:*

- (i) *The pairs $[X_0, X_1]$ and $[Y_0, Y_1]$ are compatible.*
- (ii) *The spaces Z_0 and Z_1 are linear manifolds in a certain locally convex topological space E .*
- (iii) *The continuous embeddings $Y_0 \subseteq Z_0$ and $Y_1 \subseteq Z_1$ hold.*
- (iv) *The mapping T is given on $X_0 + X_1$ and defines the bounded operators $T : X_0 \rightarrow Z_0$ and $T : X_1 \rightarrow Z_1$.*
- (v) *The mapping R is given on E and defines the bounded operators $R : Z_0 \rightarrow X_0$ and $R : Z_1 \rightarrow X_1$.*
- (vi) *The mapping S is given on E and defines the bounded operators $S : Z_0 \rightarrow Y_0$ and $S : Z_1 \rightarrow Y_1$.*
- (vii) *The equality $TRu = u + Su$ holds for every $u \in E$.*

Then

$$(3) \quad [(X_0)_{T,Y_0}, (X_1)_{T,Y_1}]_\theta = ([X_0, X_1]_\theta)_{T,[Y_0,Y_1]_\theta}$$

up to equivalence of norms for every $\theta \in (0, 1)$.

This result is proved by Lions and Magenes [12, Chapter 1, Theorem 14.3] in a more general case of Banach spaces.

Let us now prove Theorem 2. Our reasoning is motivated by [12, Chapter 2, Proof of Theorem 4.2] and [8, Section 3.3.4]. We choose an integer $r \geq 1$ arbitrarily and consider the linear differential expression $A^r A^{r+} + 1$ of order $4qr$. Here, as usual, A^{r+} denotes the differential expression which is formally adjoint to the r -th iteration A^r of A . Given an integer $\sigma \geq 2qr$, we let $H_D^\sigma(\Omega)$ denote the linear manifold of all distributions $u \in H^\sigma(\Omega)$ such that $\partial_\nu^j u = 0$ on Γ for each $j \in \{0, \dots, 2qr - 1\}$. Here, ∂_ν is the operator of the differentiation with respect to the inward normal to the boundary Γ of Ω . This definition is reasonable in view of the theorem on traces of distributions from $H^\sigma(\Omega)$ (see, e.g., [15, Section 4.7.1]). According to this theorem, $H_D^\sigma(\Omega)$ is a (closed) subspace of $H^\sigma(\Omega)$. We have the isomorphism

$$A^r A^{r+} + 1 : H_D^\sigma(\Omega) \leftrightarrow H^{\sigma-4qr}(\Omega)$$

for every integer $\sigma \geq 2qr$ (see, e.g., [8, Lemma 3.1]). Let $(A^r A^{r+} + 1)^{-1}$ denote the inverse of this isomorphism. This inverse defines the continuous linear operator

$$(4) \quad (A^r A^{r+} + 1)^{-1} : H^l(\Omega) \rightarrow H^{l+4qr}(\Omega)$$

for every integer $l \geq -2qr$. It follows from the interpolation formula (1) that this operator is well defined and continuous for every real $l \geq -2qr$.

In Proposition 1, we put $X_j := H^{s_j}(\Omega)$, $Y_j := H^{\lambda_j}(\Omega)$, $Z_j := H^{s_j-2q}(\Omega)$, $E := H^{\min\{s_0, s_1\}-2q}(\Omega)$, and $T := A$. Evidently, conditions (i)–(iv) of this proposition are fulfilled. Besides, subjecting r to the restrictions

$$s_j - 2q \geq -2qr \quad \text{and} \quad s_j - 2q - \lambda_j \geq -4qr$$

for each $j \in \{0, 1\}$, we put $R := A^{r-1} A^{r+} (A^r A^{r+} + 1)^{-1}$ and $S := -(A^r A^{r+} + 1)^{-1}$. Owing to (4), we have the continuous linear operators

$$R : Z_j = H^{s_j-2q}(\Omega) \rightarrow H^{s_j}(\Omega) = X_j$$

and

$$S : Z_j = H^{s_j-2q}(\Omega) \rightarrow H^{s_j-2q+4qr}(\Omega) \subseteq H^{\lambda_j}(\Omega) = Y_j$$

for each $j \in \{0, 1\}$ (the last embedding is continuous); i.e., conditions (v) and (vi) are fulfilled as well. The last condition (vii) is satisfied because

$$TRu = (A^r A^{r+} + 1 - 1)(A^r A^{r+} + 1)^{-1}u = u + Su$$

for every $u \in E$. Thus, owing to Proposition 1 and the interpolation formula (1), we conclude that

$$\begin{aligned} [H_{A, \lambda_0}^{s_0}(\Omega), H_{A, \lambda_1}^{s_1}(\Omega)]_\theta &= [(X_0)_{T, Y_0}, (X_1)_{T, Y_1}]_\theta \\ &= ([X_0, X_1]_\theta)_{T, [Y_0, Y_1]_\theta} = H_{A, \lambda}^s(\Omega) \end{aligned}$$

up to equivalence of norms, which proves Theorem 2.

Let us prove the remaining part of Theorem 1. We make use of the following supplement to Proposition 1:

Proposition 2. *Suppose that the hypothesis of Proposition 1 is fulfilled. If the Hilbert spaces X_0 , Y_0 , X_1 , and Y_1 are separable and satisfy the dense continuous embeddings $X_1 \subseteq X_0$ and $Y_1 \subseteq Y_0$, then the Hilbert spaces $(X_0)_{T,Y_0}$ and $(X_1)_{T,Y_1}$ are also separable and satisfy the dense continuous embedding $(X_1)_{T,Y_1} \subseteq (X_0)_{T,Y_0}$.*

This result is proved in [17, Theorem 4.1] for a more general method of interpolation between Hilbert spaces than that used in the paper (see also [8, Section 3.3.2]).

According to Proposition 2, the spaces $H_{A,\lambda_0}^{s_0}(\Omega)$ and $H_{A,\lambda_1}^{s_1}(\Omega)$ from Theorem 2 are separable, and the second of them is continuously and densely embedded in the first provided that $s_0 \leq s_1$ and $\lambda_0 \leq \lambda_1$. This specifically implies assertion (i) of Theorem 1 concerning the separability of $H_{A,\lambda}^s(\Omega)$ for all $s \in \mathbb{R}$ and $\lambda > s - 2q$. If $\lambda \leq s - 2q$, this separability follows from assertion (ii).

It remains to prove assertions (iii) and (iv) of Theorem 1. Let $s, \lambda \in \mathbb{R}$; if $\lambda > s - 2q$, we have the dense continuous embedding

$$(5) \quad H^{s_1}(\Omega) = H_{A,s_1-2q}^{s_1}(\Omega) \subseteq H_{A,\lambda}^s(\Omega)$$

for every number $s_1 \in \mathbb{R}$ such that $s_1 \geq s$ and $s_1 - 2q \geq \lambda$. This has been noted in the previous paragraph. Since the set $C^\infty(\overline{\Omega})$ is dense in $H^{s_1}(\Omega)$, this embedding implies assertion (iii) in the case of $\lambda > s - 2q$. In the opposite case, this assertion is a consequence of assertion (ii).

Finally, assume that $s \leq 1/2$ and $\lambda \leq 1/2 - 2q$. If $\lambda > s - 2q$, assertion (iv) follows from (5) and the density of $C_0^\infty(\Omega)$ in $H^{s_1}(\Omega)$ for $s_1 \leq 1/2$ (see, e.g., [15, Section 4.3.2, Theorem 1 (a)]). If $\lambda \leq s - 2q$, this assertion follows from this density in view of assertion (ii). Our proof is complete.

4. APPLICATION

Let us discuss an application of the space $H_{A,\lambda}^s(\Omega)$ to elliptic boundary-value problems. We consider a boundary-value problem that consists of the elliptic equation

$$(6) \quad Au = f \quad \text{in } \Omega$$

and the boundary conditions

$$(7) \quad B_j u = g_j \quad \text{on } \Gamma, \quad j = 1, \dots, q.$$

Here, each

$$B_j := \sum_{|\mu| \leq m_j} b_{j,\mu}(x) \frac{\partial^{|\mu|}}{\partial x_1^{\mu_1} \dots \partial x_n^{\mu_n}}$$

is a linear boundary partial differential expression on Γ of order $m_j \leq 2q - 1$ with complex-valued coefficients $b_{j,\mu} \in C^\infty(\Gamma)$. We suppose that the boundary-value problem (6), (7) is elliptic in Ω (see the definition in, e.g., [19, Section 1.2]). Put $m := \max\{m_1, \dots, m_q\}$; the case of $m \geq 2q$ is possible.

It is known [18, Theorem 7] that the mapping

$$(8) \quad u \mapsto (Au, Bu) = (Au, B_1 u, \dots, B_q u),$$

where $u \in C^\infty(\overline{\Omega})$, extends uniquely (by continuity) to a Fredholm continuous linear operator

$$(A, B) : H^s(\Omega) \rightarrow H^{s-2q}(\Omega) \oplus \bigoplus_{j=1}^q H^{s-m_j-1/2}(\Gamma)$$

for every real number $s > m + 1/2$. Moreover, the kernel of this operator lies in $C^\infty(\overline{\Omega})$ and together with the index of the operator does not depend on s . Here and below, $H^\sigma(\Gamma)$ denotes the inner product Sobolev space over Γ of order $\sigma \in \mathbb{R}$. Let N and \varkappa respectively denote these kernel and index.

This fundamental property of problem (6), (7) does not remain for $s \leq m + 1/2$. Specifically, if $s \leq m_j + 1/2$, then the mapping $u \mapsto B_j u$, where $u \in C^\infty(\overline{\Omega})$, cannot be extended to a continuous linear operator from $H^s(\Omega)$ to the linear topological space $\mathcal{D}'(\Gamma)$ of all distributions on Γ . But it is possible to prove a version of this property for every $s \leq m + 1/2$ provided that we use some spaces $H_{A,\lambda}^s(\Omega)$ instead of $H^s(\Omega)$.

Theorem 3. *Let numbers $s, \lambda \in \mathbb{R}$ satisfy the conditions $s \leq m + 1/2$, $\lambda > -1/2$, and $\lambda > m + 1/2 - 2q$. Then the mapping (8), where $u \in C^\infty(\overline{\Omega})$, extends uniquely (by continuity) to a Fredholm continuous linear operator*

$$(9) \quad (A, B) : H_{A,\lambda}^s(\Omega) \rightarrow H^\lambda(\Omega) \oplus \bigoplus_{j=1}^q H^{s-m_j-1/2}(\Gamma).$$

The kernel of this operator equals N , and the index equals \varkappa .

If $s \in \mathbb{Z}$ and $s \leq 0$, Theorem 3 is proved in a somewhat similar way as that used in [7] and [8, Sections 4.4.2 and 4.4.3] for regular elliptic boundary-value problems. (The corresponding reasoning is given in [20, Section 5] for $m \geq 2q$). In the general situation we prove Theorem 3 with the help of the interpolation formula (2). Choose $l \in \mathbb{Z}$ such that $l < s$ and $l \leq 0$. We consider the Fredholm continuous operators

$$(10) \quad (A, B) : H_{A,\lambda}^l(\Omega) \rightarrow H^\lambda(\Omega) \oplus \bigoplus_{j=1}^q H^{l-m_j-1/2}(\Gamma)$$

and

$$(11) \quad (A, B) : H^{\lambda+2q}(\Omega) \rightarrow H^\lambda(\Omega) \oplus \bigoplus_{j=1}^q H^{\lambda+2q-m_j-1/2}(\Gamma)$$

(note that $\lambda + 2q > 1/2$ by the hypothesis of the Theorem). They have the common kernel N and the common index \varkappa . Since $l < s < \lambda + 2q$, there exists a number $\theta \in (0, 1)$ such that $s = (1 - \theta)l + \theta(\lambda + 2q)$. Applying the interpolation with the parameter θ to these operators, we conclude by [8, Theorem 1.5] that a restriction of the first operator is a Fredholm continuous operator between the spaces

$$[H_{A,\lambda}^l(\Omega), H^{\lambda+2q}(\Omega)]_\theta$$

and

$$\left[H^\lambda(\Omega) \oplus \bigoplus_{j=1}^q H^{l-m_j-1/2}(\Gamma), H^\lambda(\Omega) \oplus \bigoplus_{j=1}^q H^{\lambda+2q-m_j-1/2}(\Gamma) \right]_\theta.$$

Moreover, this operator has the same kernel and index as operators (10) and (11). Owing to Theorem 2, the first space coincides with $H_{A,\lambda}^s(\Omega)$ up to equivalence of norms (remark that $H^{\lambda+2q}(\Omega) = H_{A,\lambda}^{\lambda+2q}(\Omega)$). Besides, the second space coincides up to equivalence of norms with the target space of operator (9) due to an analog of the interpolation formula (1) for Sobolev spaces over Γ (see, e.g. [12, Chapter 1, Theorem 7.7]). Thus, the latter Fredholm operator acts between the spaces indicated in (9). It is an extension by continuity of the mapping (8), where $u \in C^\infty(\overline{\Omega})$, in view of assertion (iii) of Theorem 1. The proof of Theorem 3 is complete.

Remark that, in the important case of regular elliptic boundary-value problems, Theorem 3 is proved by Lions and Magenes [4, 5] in the framework of L_p -Sobolev spaces provided that $\lambda = 0$ and $s \notin \{1/p - k : 1 \leq k \in \mathbb{Z}\}$. In the case where $s \geq 0$ and $m \leq 2q - 1$, this theorem is contained in the result formulated in [19, p. 86].

REFERENCES

- [1] O. A. Ladyzenskaya, *On the closure of the elliptic operator* (Russian), Doklady Akad. Nauk SSSR (N.S.) **79**, (1951). 723–725.
- [2] L. Hörmander, *On the theory of general partial differential equations*, Acta Math. **94** (1955), no. 1, 161–248.
- [3] M. Š. Birman, *Characterization of elliptic differential operators with the maximal domain of definition* (Russian), Vestnik Leningrad. Univ. Ser. Math. Mech. Astr. **19** (1957), no. 4, 177–183.
- [4] J.-L. Lions, E. Magenes, *Problèmes aux limites non homogènes, V*, Ann. Sci. Norm. Sup. Pisa **16** (1962), 1–44.
- [5] J.-L. Lions, E. Magenes, *Problèmes aux limites non homogènes, VI*, J. d’Analyse Math. **11** (1963), 165–188.
- [6] J.-L. Lions, E. Magenes, *Problèmes aux limites non homogènes, II*, Ann. Inst. Fourier (Grenoble) **11** (1961), 137–178.
- [7] Murach A. A., *Extension of some Lions–Magenes theorems*, Methods Funct. Anal. Topology, **15** (2009), no. 2, 152–167.
- [8] V. A. Mikhailets, A. A. Murach, *Hörmander spaces, interpolation, and elliptic problems*, de Gruyter Stud. Math., vol. 60, De Gruyter, Berlin, 2014.
- [9] G. Geymonat, *Sul problema di Dirichlet per le equazioni lineari ellittiche*, Ann. Sci. Norm. Sup. Pisa **16** (1962), 225–284.
- [10] E. Magenes, *Spazi di interpolazione ed equazioni a derivate parziali*, Atti VII Congr. Un. Mat. Italiana (Genoa, 1963), Edizioni Cremonese, Rome, 1965, 134–197.
- [11] J.-L. Lions, E. Magenes, *Problèmes aux limites non homogènes, VII*, Ann. Mat. Pura Appl. (4) **63** (1963), 201–224.
- [12] J.-L. Lions and E. Magenes, *Non-Homogeneous Boundary-Value Problems and Applications, vol. I*, Grundlehren Math. Wiss., vol. 181, Springer-Verlag, New York–Heidelberg, 1972.
- [13] V. A. Mikhailets and A. A. Murach, *Elliptic problems and Hörmander spaces*, Modern analysis and applications. The Mark Krein Centenary Conference. Vol. 2: Differential operators and mechanics, 447–470, Oper. Theory Adv. Appl. **191**, Birkhäuser, Basel, 2009.

- [14] V. A. Mikhailets, A. A. Murach, *The refined Sobolev scale, interpolation, and elliptic problems*, Banach J. Math. Anal. **6** (2012), no. 2, 211–281.
- [15] H. Triebel, *Interpolation Theory, Function Spaces, Differential Operators* [2-nd edn], Johann Ambrosius Barth, Heidelberg, 1995.
- [16] L. Hörmander, *Linear Partial Differential Operators*, Grundlehren Math. Wiss., vol. 116, Springer, Berlin, 1963.
- [17] V. A. Mikhailets, A. A. Murach, *A regular elliptic boundary-value problem for a homogeneous equation in a two-sided refined scale of spaces*, Ukrainian Math. J. **58** (2006), no. 11, 1748–1767.
- [18] J. Peetre, *Another approach to elliptic boundary-value problems*, Comm. Pure Appl. Math. **14** (1961), no. 4, 711–731.
- [19] M. S. Agranovich, *Elliptic boundary problems*, Partial differential equations, IX, 1–144, Encyclopaedia Math. Sci., **79**, Springer, Berlin, 1997.
- [20] A. V. Anop, T. M. Kasirenko, A. A. Murach, *Nonregular elliptic boundary-value problems and Hörmander spaces*, Ukraïn. Mat. Zh. **70** (2018), no. 3, 299–317.

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3
TERESHCHENKIVS'KA, KYIV, 01601, UKRAINE
E-mail address: kasirenko@imath.kiev.ua

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3
TERESHCHENKIVS'KA, KYIV, 01601, UKRAINE
E-mail address: mikhailets@imath.kiev.ua

INSTITUTE OF MATHEMATICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 3
TERESHCHENKIVS'KA, KYIV, 01601, UKRAINE
E-mail address: murach@imath.kiev.ua