

# Finite groups determined by an inequality of the orders of their elements

Marius Tărnăuceanu

October 1, 2012

## Abstract

In this note we introduce and characterize a class of finite groups for which the element orders satisfy a certain inequality. This is contained in some well-known classes of finite groups.

**MSC (2010):** Primary 20D10, 20D20; Secondary 20D15, 20D25, 20E34.

**Key words:** finite groups, element orders, CP-groups, Frobenius groups.

## 1 Introduction

Let  $CP_1$ ,  $CP$  and  $CN$  be the classes of finite groups in which the centralizers of all nontrivial elements contain only elements of prime order, of prime power order and are nilpotent, respectively. Clearly, we have  $CP_1 \subset CP \subset CN$ . Moreover, the classes  $CP_1$  and  $CP$  consist of exactly those finite groups all of whose elements have prime order and prime power order, respectively. They have been studied in many papers, as [1]-[3], [5]-[7] and [13].

In the following we consider the finite groups  $G$  such that

$$(*) \quad o(xy) \leq \max\{o(x), o(y)\}, \text{ for all } x, y \in G.$$

These form another interesting subclass of  $CP$ , that will be denoted by  $CP_2$ . Its exhaustive description is the main goal of this note.

Most of our notation is standard and will not be repeated here. Basic notions and results on group theory can be found in [4, 8, 9, 11].

First of all, we observe that if a finite group  $G$  belongs to  $\text{CP}_2$ , then for every  $x, y \in G$  satisfying  $o(x) \neq o(y)$  we have

$$o(xy) = \max\{o(x), o(y)\},$$

that is the order map is very close to a monoid homomorphism from  $(G, \cdot)$  to  $(\mathbb{N}^*, \max)$ .

An immediate characterization of finite groups contained in  $\text{CP}_2$  is indicated in the following theorem.

**Theorem A.** *Let  $G$  be a finite group and set  $\pi_e(G) = \{o(x) \mid x \in G\}$ . Then the following conditions are equivalent:*

- a)  $G$  belongs to  $\text{CP}_2$ .
- b) For every  $\alpha \in \pi_e(G)$ , the set  $G_\alpha = \{x \in G \mid o(x) \leq \alpha\}$  is a normal subgroup of  $G$ .

Next, we will focus on establishing some connections between  $\text{CP}_2$  and the previous classes  $\text{CP}$  and  $\text{CP}_1$ .

**Proposition B.** *The class  $\text{CP}_2$  is properly contained in the class  $\text{CP}$ .*

On the other hand, by taking  $\sigma = (12)(34), \tau = (235) \in A_5$ , one obtains

$$5 = o(\sigma\tau) > 3 = \max\{o(\sigma), o(\tau)\},$$

and therefore  $\text{CP}_2$  does not contain the alternating group  $A_5$ . Since  $A_5$  belongs to  $\text{CP}_1$ , we conclude that  $\text{CP}_1$  is not contained in  $\text{CP}_2$ . It is obvious that the converse inclusion also fails (for example, any abelian  $p$ -group belongs to  $\text{CP}_2$ , but not to  $\text{CP}_1$ ).

### Remarks.

1. Other two remarkable classes of finite  $p$ -groups, more large as the class of abelian  $p$ -groups, are contained in  $\text{CP}_2$ : regular  $p$ -groups (see Theorem 3.14 of [11], II, page 47) and  $p$ -groups whose subgroup lattices are modular (see Lemma 2.3.5 of [10]). Moreover, by the main theorem of [12], we infer that the powerful  $p$ -groups for  $p$  odd also belong to  $\text{CP}_2$ .

2. The smallest nonabelian  $p$ -group contained in  $\text{CP}_2$  is the quaternion group  $Q_8$ , while the smallest  $p$ -group not contained in  $\text{CP}_2$  is the dihedral group  $D_8$ . Notice that all quaternion groups  $Q_{2^n}$ , for  $n \geq 4$ , as well as all dihedral groups  $D_n$ , for  $n \neq 1, 2, 4$ , are not contained in  $\text{CP}_2$ .
3. The class  $\text{CP}_2$  contains finite groups which are not  $p$ -groups, too. The smallest example of such a group is  $A_4$ . Remark that the groups  $A_n$ ,  $n \geq 5$ , does not belong to  $\text{CP}_2$ , and this is also valid for the symmetric groups  $S_n$ ,  $n \geq 3$ .

Clearly,  $\text{CP}_2$  is closed under subgroups. On the other hand, the above results imply that  $\text{CP}_2$  is not closed under direct products or extensions. The same thing can be said with respect to homomorphic images, as shows the following example.

**Example.** Let  $p$  be a prime and  $G$  be the semidirect product of an elementary abelian  $p$ -group  $A$  of order  $p^p$  by a cyclic group of order  $p^2$ , generated by an element  $x$  which permutes the elements of a basis of  $A$  cyclically. Then it is easy to see that  $G$  belongs to  $\text{CP}_2$ ,  $x^p \in Z(G)$  and the quotient  $Q = \frac{G}{\langle x^p \rangle}$  is isomorphic to a Sylow  $p$ -subgroup of  $S_{p^2}$ . Obviously, in  $Q$  a product of two elements of order  $p$  can have order  $p^2$ , and hence it does not belong to  $\text{CP}_2$ .

The next result collects other basic properties of the finite groups contained in  $\text{CP}_2$ .

**Proposition C.** *Let  $G$  be a finite group contained in  $\text{CP}_2$ . Then:*

- a) *There is a prime  $p$  dividing the order of  $G$  such that  $F(G) = O_p(G)$ .*
- b) *Both  $Z(G)$  and  $\Phi(G)$  are  $p$ -groups.*
- c)  *$Z(G)$  is trivial if  $G$  is not a  $p$ -group.*

We are now able to present our main result, that gives a complete description of the class  $\text{CP}_2$ .

**Theorem D.** *A finite group  $G$  is contained in  $\text{CP}_2$  if and only if one of the following statements holds:*

- a)  *$G$  is a  $p$ -group and  $\Omega_n(G) = \{x \in G \mid x^{p^n} = 1\}$ , for all  $n \in \mathbb{N}$ .*

- b)  $G$  is a Frobenius group of order  $p^\alpha q^\beta$ ,  $p < q$ , with kernel  $F(G)$  of order  $p^\alpha$  and cyclic complement.

Since all  $p$ -group and all groups of order  $p^\alpha q^\beta$  are solvable, Theorem D leads to the following corollary.

**Corollary E.** *The class  $CP_2$  is properly contained in the class of finite solvable groups.*

**Remark.** The finite supersolvable groups and the CLT-groups constitute two important subclasses of the finite solvable groups. Since  $A_4$  belongs to  $CP_2$ , we infer that  $CP_2$  is not included in these classes. Conversely, a finite supersolvable group or a CLT-group does not necessarily possess the structure described above, and thus they are not necessarily contained in the class  $CP_2$ .

As we already have seen, both  $CP_1$  and  $CP_2$  are subclasses of  $CP$ , and each of them is not contained in the other. Consequently, an interesting problem is to find the intersection of these subclasses. This can be made by using again Theorem D.

**Corollary F.** *A finite group  $G$  is contained in the intersection of  $CP_1$  and  $CP_2$  if and only if one of the following statements holds:*

- a)  $G$  is a  $p$ -group of exponent  $p$ .
- b)  $G$  is a Frobenius group of order  $p^\alpha q$ ,  $p < q$ , with kernel  $F(G)$  of order  $p^\alpha$  and exponent  $p$ , and cyclic complement. Moreover, in this case we have  $G' = F(G)$ .

**Remark.**  $A_4$  is an example of a group of type b) in the above corollary. Mention that for such a group  $G$  the number of Sylow  $q$ -subgroups is  $p^\alpha$ . It is also clear that  $G$  possesses a nontrivial partition consisting of Sylow subgroups:  $F(G)$  and all conjugates of a Frobenius complement.

Finally, we indicate a natural problem concerning the class of finite groups introduced in our paper.

**Open problem.** Give a precise description of the structure of finite  $p$ -groups contained in  $CP_2$ .

## 2 Proofs of the main results

**Proof of Theorem A.** Assume first that  $G$  belongs to  $\text{CP}_2$ . Let  $\alpha \in \pi_e(G)$  and  $x, y \in G_\alpha$ . Then, by  $(*)$ , we have

$$o(xy) \leq \max\{o(x), o(y)\} \leq \alpha,$$

which shows that  $xy \in G_\alpha$ . This proves that  $G_\alpha$  is a subgroup of  $G$ . Moreover,  $G_\alpha$  is normal in  $G$  because the order map is constant on each conjugacy class.

Conversely, let  $x, y \in G$  and put  $\alpha = o(x), \beta = o(y)$ . By supposing that  $\alpha \leq \beta$ , one obtains  $x, y \in G_\beta$ . Since  $G_\beta$  is a subgroup of  $G$ , it follows that  $xy \in G_\beta$ . Therefore

$$o(xy) \leq \beta = \max\{o(x), o(y)\},$$

completing the proof. ■

**Proof of Proposition B.** Let  $G$  be finite group in  $\text{CP}_2$  and take  $x \in G$ . It is well-known that  $x$  can be written as a product of (commuting) elements of prime power orders, say  $x = x_1 x_2 \cdots x_k$ . Then the condition  $(*)$  implies that

$$\prod_{i=1}^k o(x_i) = o(x) \leq \max\{o(x_i) \mid i = \overline{1, k}\},$$

and so  $k = 1$ . Hence  $x$  is of prime power order, i.e.  $G$  is contained in  $\text{CP}$ .

Obviously, the inclusion of  $\text{CP}_2$  in  $\text{CP}$  is strict (we already have seen that  $A_5$  belongs to  $\text{CP}$ , but not to  $\text{CP}_2$ ). ■

**Proof of Proposition C.**

- a) We know that  $F(G)$  is the product of the subgroups  $O_p(G)$ , where  $p$  runs over the prime divisors of  $|G|$ . Suppose that there are two distinct primes  $p$  and  $q$  dividing the order of  $F(G)$ . This leads to the existence of two elements  $x$  and  $y$  of  $F(G)$  such that  $o(x) = p$  and  $o(y) = q$ . Since  $F(G)$  is nilpotent, we obtain  $xy = yx$  and so  $o(xy) = pq$ , a contradiction. Thus  $F(G) = O_p(G)$ , for a prime divisor  $p$  of  $|G|$ .
- b) It is well-known that both  $Z(G)$  and  $\Phi(G)$  are normal nilpotent subgroup of  $G$ . By the maximality of  $F(G)$ , it follows that  $Z(G)$  and  $\Phi(G)$  are contained in  $F(G)$ , and therefore they are also  $p$ -groups.

- c) Assume that  $Z(G)$  is not trivial and take  $x \in Z(G)$  with  $o(x) = p$ . If  $G$  is not a  $p$ -group, it contains an element  $y$  of prime order  $q \neq p$ . Then  $o(xy) = pq$ , contradicting Proposition B. ■

**Proof of Theorem D.** If  $G$  is a  $p$ -group, then the conclusion is obvious.

Assume now that  $G$  is not a  $p$ -group. We will proceed by induction on  $|G|$ . Since  $G$  belongs to  $\text{CP}_2$ , all the numbers in  $\pi_e(G)$  are prime powers. Let  $q^n$  be the largest number of  $\pi_e(G)$ , where  $q$  is a prime, and let  $N = \{g \in G \mid o(g) < q^n\}$ . Then  $N \trianglelefteq G$  and  $\exp(G/N) = q$ . Since  $|N| < |G|$ , by the inductive hypothesis it follows that either  $N$  is a  $p$ -group or  $N$  is a Frobenius group with kernel  $K$  of order  $p^\alpha$  and cyclic complement  $H$  of order  $r^\beta$ , where  $p, r$  are distinct primes. We will prove that in both cases  $G$  is a Frobenius group whose kernel and complement are  $p$ -groups.

**Case 1.**  $N$  is a  $p$ -group.

Since  $G$  is not a  $p$ -group, we can take  $Q \in \text{Syl}_q(G)$ , where  $p \neq q$ . So  $G = N \rtimes Q$ . Since every element of  $N$  is of prime power order, we have  $C_N(h) = 1$  for all  $1 \neq h \in Q$ . Thus,  $G$  is a Frobenius group with kernel  $N$  and complement  $Q$ .

**Case 2.**  $N$  is a Frobenius group.

**Subcase 2.1.**  $q \neq p$  and  $q \neq r$ .

By a similar argument as that of Case 1, we know that  $G$  is a Frobenius group with kernel  $N$ . But  $N$  is not nilpotent, a contradiction.

**Subcase 2.2.**  $q = r$ .

Let  $Q \in \text{Syl}_q(G)$ . Then  $G = K \rtimes Q$ . By a similar argument as that of Case 1, we know that  $G$  is a Frobenius group with kernel  $K$  and complement  $Q$ .

**Subcase 2.3.**  $q = p$ .

We observe that all elements of  $G \setminus N$  are of order  $q^n$ , and  $g^q \in K$ , where  $g \in G \setminus N$  and  $K \in \text{Syl}_q(N)$ . So if  $N_G(H) \cap (G \setminus N) \neq 1$ , then  $N_G(H) \cap K \neq 1$ . But  $N$  is a Frobenius group and  $N_N(H) = H$ . It follows that  $N_G(H) = H$ . Since  $H$  is cyclic,  $N_G(H) = C_G(H)$ . One obtains that  $G$  is  $r$ -nilpotent and thus  $G = P \rtimes H$ , where  $P \in \text{Syl}_p(G)$ . A similar argument as that of Case 1 shows that  $G$  is again a Frobenius group with kernel  $P$  and complement  $H$ .

Finally, we prove that  $H$  is cyclic. By Burnside's Theorem we only need to prove that  $H$  is not a 2-group. If not, let  $L = \{g \in G \mid o(g) = 2\}$ . Then

$L \trianglelefteq G$ . It follows that  $K \times L \leq G$ , where  $K$  is the Frobenius kernel. This contradicts the fact that all elements of  $G$  are of prime power order. ■

**Proof of Corollary F.** The equivalence follows directly by Theorem D. In this way, we have to prove only that  $G' = F(G)$  in the case b).

Obviously,  $G' \subseteq F(G)$ . For the converse inclusion, let  $x \in F(G)$  be a nontrivial element. Then  $o(x) = p$ . If  $y$  is an arbitrary element of order  $q$  in  $G$ , then we have

$$o(xy) \leq \max\{o(x), o(y)\} = q,$$

and therefore  $o(xy) \in \{p, q\}$ . If we assume that  $o(xy) = p$ , it results

$$q = o(y) = o(x^{-1}xy) \leq \max\{o(x^{-1}), o(xy)\} = p,$$

a contradiction. This shows that  $o(xy) = q$ . Then there is  $z \in G$  such that  $xy \in \langle y \rangle^z$ , say  $xy = z^{-1}y^kz$  with  $k \in \mathbb{Z}$ . Since the element

$$xy^{1-k} = z^{-1}y^kzy^{-k} = [z, y^k]$$

has order  $p$ , we infer that  $k$  must be equal to 1. Hence

$$x = [z, y] \in G',$$

which completes the proof. ■

**Acknowledgements.** The author is grateful to the reviewer for its remarks which improve the previous version of the paper.

## References

- [1] Bannuscher, W., Tiedt, G., *On a theorem of Deaconescu*, Rostock. Math. Kolloq. **47** (1994), 23-27.
- [2] Brandl, R., *Finite groups all of whose elements are of prime power order*, Boll. Un. Mat. Ital. A **18** (1981), 491-493.
- [3] Deaconescu, M., *Classification of finite groups with all elements of prime order*, Proc. Amer. Math. Soc. **106** (1989), 625-629.

- [4] Gorenstein, D., *Finite simple groups*, Plenum Press, New York-London, 1982.
- [5] Heineken, H., *On groups all of whose elements have prime power order*, Math. Proc. Royal Irish Acad. **106** (2006), 191-198.
- [6] Higman, G., *Groups and rings having automorphisms without nontrivial fixed elements*, J. London Math. Soc. **32** (1957), 321-334.
- [7] Higman, G., *Finite groups in which every element has prime power order*, J. London Math. Soc. **32** (1957), 335-342.
- [8] Huppert, B., *Endliche Gruppen*, I, Springer Verlag, Berlin-Heidelberg-New York, 1967.
- [9] Isaacs, I.M., *Finite group theory*, Amer. Math. Soc., Providence, R.I., 2008.
- [10] Schmidt, R., *Subgroup lattices of groups*, de Gruyter Expositions in Mathematics **14**, de Gruyter, Berlin, 1994.
- [11] Suzuki, M., *Group theory*, I, II, Springer Verlag, Berlin, 1982, 1986.
- [12] Wilson, L., *On the power structure of powerful  $p$ -groups*, J. Group Theory **5** (2002), 129-144.
- [13] Yang, W., Zhang, Z., *Locally soluble infinite groups in which every element has prime power order*, Southeast Asian Bull. Math. **26** (2003), 857-864.

Marius Tărnăuceanu  
Faculty of Mathematics  
“Al.I. Cuza” University  
Iași, Romania  
e-mail: [tarnauc@uaic.ro](mailto:tarnauc@uaic.ro)