

VOLTERRA TYPE OPERATORS ON WEIGHTED BANACH SPACES OF ANALYTIC FUNCTIONS

QINGZE LIN

ABSTRACT. Smith et al. recently gave the sufficient and necessary conditions for the boundedness of Volterra type operators on Banach spaces of bounded analytic functions when the symbol functions are univalent. In this paper, we give the complete characterizations of the conditions for the boundedness and compactness of Volterra type operators T_g and their companion operators S_g between weighted Banach spaces of analytic functions, which essentially generalize their works.

1. Introduction

Let \mathbb{D} be the unit disk of the complex plane \mathbb{C} and $H(\mathbb{D})$ be the space consisting of all analytic functions on the unit disk. For any analytic function $g \in H(\mathbb{D})$, the Volterra type operator T_g is defined by

$$(T_g f)(z) = \int_0^z f(\omega)g'(\omega)d\omega, \quad z \in \mathbb{D}, f \in H(\mathbb{D}).$$

Its companion operator S_g is defined by

$$(S_g f)(z) = \int_0^z f'(\omega)g(\omega)d\omega, \quad z \in \mathbb{D}, f \in H(\mathbb{D}).$$

The boundedness and compactness of these two operators on some spaces of analytic functions were extensively studied. Pommerenke [19] firstly studied the boundedness of T_g on Hardy-Hilbert space H^2 . After his work, Aleman, Siskakis and Cima [1, 2] studied the boundedness and compactness of T_g on Hardy space H^p , in which they showed that T_g is bounded (compact) on H^p , $0 < p < \infty$, if and only if $g \in BMOA$ ($g \in VMOA$). For the related works, see [11]. Here, $BMOA$ and $VMOA$ denote the spaces of $f \in H(\mathbb{D})$ with boundary values of bounded mean oscillation and vanishing mean oscillation, respectively. What's more, Aleman and Siskakis [3] studied the boundedness and compactness of T_g on Bergman spaces and Xiao [21] studied T_g and S_g on Q_p spaces.

Recently, Li et al. [12] studied T_g and S_g on analytic Morrey spaces. Lin et al. [13] characterized the boundedness of T_g and S_g acting on the derivative Hardy spaces S^p ($1 \leq p < \infty$). Miihkinen [16] investigated the strict singularity of T_g on Hardy space H^p and showed that the strict singularity of T_g coincides with its compactness on H^p . Mengestie [15] obtained a complete description of the boundedness and compactness of the product of Volterra type operators and composition operators on weighted Fock spaces. Furthermore, by applying the Carleson embedding theorem and the Littlewood-Paley formula, Constantin and

2010 *Mathematics Subject Classification.* 47G10, 30H05.

Key words and phrases. Volterra type operator, boundedness, compactness, Banach space.

Peláez [7] obtained the boundedness and compactness of T_g on weighted Fock spaces and investigated the invariant subspaces of the Volterra operator T_z on such spaces.

Now, for any $\alpha \geq 0$, we define the weighted Banach spaces H_α^∞ of analytic functions given by

$$H_\alpha^\infty = \{f \in H(\mathbb{D}) : \|f\|_{H_\alpha^\infty} := \sup_{z \in \mathbb{D}} \{(1 - |z|^2)^\alpha |f(z)|\} < \infty\}.$$

When $\alpha = 0$, H_0^∞ is the usual Banach space of bounded analytic functions. The norms and essential norms of several bounded linear operators on such spaces had been extensively studied. For instance, Montes-Rodríguez [17] obtained the essential norms of weighted composition operators on H_α^∞ .

Although the necessary and sufficient conditions for $T_g : H_0^\infty \rightarrow H_0^\infty$ to be bounded are characterized recently by Contreras, Peláez, Pommerenke and Rättyä [8], the conditions characterizing the symbol g use the Cauchy transforms which are very difficult to verified. However, Anderson, Jovovic and Smith [4] conjectured that $T_g : H_0^\infty \rightarrow H_0^\infty$ is bounded if and only if

$$\sup_{0 \leq \theta < 2\pi} \int_0^1 |g'(re^{i\theta})| dr < +\infty.$$

While this condition of symbol g is sufficient for the boundedness of $T_g : H_0^\infty \rightarrow H_0^\infty$, it was recently proved to be not necessary by Smith, Stolyarov and Volberg in [20], where a counterexample was given. However, when the symbol g is univalent, this conjecture was proved to be affirmative in the same paper. Following their works, Eklund et. al. [9] studied the boundedness and compactness of T_g between H_α^∞ and H_0^∞ , where $0 \leq \alpha < 1$ and g is univalent. It should be noticed that Basalote et al. [6] characterized the boundedness, compactness and weak compactness of T_g acting between different weighted Banach spaces of analytic functions with the quasi-normal weight, and what's more, they applied the characterization of compactness to analyze the behavior of semigroups of composition operators on such spaces.

In this paper, using the ideas from [9, 20], we give the sufficient and necessary conditions for the boundedness and compactness of $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ and $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ when $0 \leq \alpha, \beta$, which generalize their works.

2. The boundedness of Volterra type operators

For any $0 < \eta < \pi, 0 < r < 1$, and $0 \leq \theta < 2\pi$, denote by $\mathcal{B}(\Omega_{\eta,\theta}^r)$ the set of all analytic functions F defined in the open sector

$$\Omega_{\eta,\theta}^r := \left\{ z \in \mathbb{D} : 0 < |z| < r, \theta - \frac{\eta}{2} < \arg z < \theta + \frac{\eta}{2} \right\},$$

such that

$$|F'(z)| \leq \frac{C_F}{|z|}, \quad \text{for } z \in \Omega_{\eta,\theta}^r.$$

Here, C_F is a constant depending only on function f .

By the similar arguments from [9, 20], we obtain the following two lemmas.

Lemma 1. *Let $0 < \gamma < \eta < \pi, 0 \leq \theta < 2\pi$ and $\varepsilon > 0$. Then there exists a positive number $\delta(\varepsilon)$ such that for any $F \in \mathcal{B}(\Omega_{\gamma,\theta}^{1/2})$, there is a real harmonic function u defined in $\Omega_{\eta,\theta}^1$ such that:*

$$(1) \quad |Re(F(xe^{i\theta})) - u(xe^{i\theta})| \leq \varepsilon, \quad \text{for } x \in (0, \delta(\varepsilon)];$$

(2) $|\tilde{u}(z)| \leq C(\varepsilon, \gamma, \eta, C_F) < \infty$, for $z \in \Omega_{\eta, \theta}^1$, where \tilde{u} is the conjugate of u such that $\tilde{u}(\frac{1}{2}e^{i\theta}) = 0$.

Lemma 2. Let $0 < \gamma < \eta < \pi$, $0 \leq \theta < 2\pi$. If $\psi_{\eta, \theta} : \Omega_{\eta, \theta}^1 \rightarrow \mathbb{D}$ is a conformal map such that $\psi_{\eta, \theta}(\frac{1}{2}e^{i\theta}) = 0$ and $\psi_{\eta, \theta}(0) = e^{i\theta}$, then there exists a constant $C_1(\gamma, \eta)$ such that

$$\frac{|\psi'_{\eta, \theta}(z)|}{1 - |\psi_{\eta, \theta}(z)|^2} \leq \frac{C_1(\gamma, \eta)}{|z|}$$

for all $z \in \Omega_{\gamma, \theta}^{1/2}$.

Recall that the Bloch space \mathcal{B} is defined by

$$\mathcal{B} = \{f \in H(\mathbb{D}) : \|f\|_{\mathcal{B}} := |f(0)| + \sup_{z \in \mathbb{D}} \{(1 - |z|^2)|f'(z)|\} < \infty\}.$$

Now we are able to give the sufficient and necessary condition for the boundedness of Volterra type operators $T_g : H_{\alpha}^{\infty} \rightarrow H_{\beta}^{\infty}$.

Theorem 1. If $g \in H(\mathbb{D})$ such that $\log(g') \in \mathcal{B}$ and $0 \leq \alpha, \beta$, then $T_g : H_{\alpha}^{\infty} \rightarrow H_{\beta}^{\infty}$ is bounded if and only if

$$\limsup_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t^2)^{\beta} \int_0^t \frac{|g'(re^{i\theta})|}{(1 - r^2)^{\alpha}} dr < +\infty.$$

Proof. Assume first that

$$\limsup_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t^2)^{\beta} \int_0^t \frac{|g'(re^{i\theta})|}{(1 - r^2)^{\alpha}} dr = +\infty.$$

Then choose a nondecreasing positive sequence $\{t_{0, n}\}_{n=1}^{\infty}$, with $t_{0, n} \rightarrow 1^-$, such that for each $t_{0, n}$, there is an angle θ_n , with $0 \leq \theta_n < 2\pi$, such that

$$(1 - t_{0, n}^2)^{\beta} \int_0^{t_{0, n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^{\alpha}} dr \geq n.$$

Define the function $F_n : \Omega_{\eta, \theta_n}^1 \rightarrow \mathbb{D}$ by

$$F_n(z) = -i \log(g' \circ \psi_{\eta, \theta_n}(z)).$$

Since $\log(g') \in \mathcal{B}$, there is a constant $K > 0$ such that

$$|g''(z)| \leq \frac{K|g'(z)|}{(1 - |z|^2)} \quad \text{for all } z \in \mathbb{D}.$$

Then by Lemma 2, we have

$$|F'_n(z)| = \frac{|g''(\psi_{\eta, \theta_n}(z))|}{|g'(\psi_{\eta, \theta_n}(z))|} \cdot |\psi'_{\eta, \theta_n}(z)| \leq \frac{K |\psi'_{\eta, \theta_n}(z)|}{1 - |\psi_{\eta, \theta_n}(z)|^2} \leq \frac{K \cdot C_1(\gamma, \eta)}{|z|},$$

for all $z \in \Omega_{\gamma, \theta_n}^{1/2}$. Hence, the restriction of F_n to $\Omega_{\gamma, \theta_n}^{1/2}$ belongs to $\mathcal{B}(\Omega_{\gamma, \theta_n}^{1/2})$.

By Lemma 1 with $\varepsilon = \frac{\pi}{3}$, we have a positive number $\delta(\frac{\pi}{3})$ such that for any $F_n \in \mathcal{B}(\Omega_{\gamma, \theta_n}^{1/2})$, there is a real harmonic function u_n defined in $\Omega_{\eta, \theta_n}^1$ such that:

$$(1) |\operatorname{Re}(F_n(xe^{i\theta_n})) - u_n(xe^{i\theta_n})| \leq \frac{\pi}{3}, \quad \text{for } x \in (0, \delta(\frac{\pi}{3}));$$

(2) $|\tilde{u}_n(z)| \leq C(\frac{\pi}{3}, \gamma, \eta, K \cdot C_1(\gamma, \eta)) =: C_2(\gamma, \eta) < \infty$, for $z \in \Omega_{\eta, \theta_n}^1$, where \tilde{u}_n is the conjugate of u_n such that $\tilde{u}_n(\frac{1}{2}e^{i\theta_n}) = 0$.

From (1) and the definition of conformal function ψ_{η, θ_n} , we see that there exists a r_η with $0 < r_\eta < 1$ such that

$$\left| \arg g'(re^{i\theta_n}) - u_n(\psi_{\eta, \theta_n}^{-1}(re^{i\theta_n})) \right| \leq \frac{\pi}{3}, \quad \text{for } r \in [r_\eta, 1].$$

Now define

$$h_n(z) := e^{-i(u_n(z) + i\tilde{u}_n(z))},$$

and consider the sequence $\{f_n\}_{n=1}^\infty$ of functions defined by

$$f_n(z) := \frac{h_n \circ \psi_{\eta, \theta_n}^{-1}(z)}{(1 - e^{-2\theta_n i} z^2)^\alpha}.$$

Then by (2), the sequence $\{f_n\}_{n=1}^\infty$ is bounded in H_α^∞ with

$$\begin{aligned} \|f_n\|_{H_\alpha^\infty} &= \sup_{z \in \mathbb{D}} \left\{ (1 - |z|^2)^\alpha \left| \frac{h_n \circ \psi_{\eta, \theta_n}^{-1}(z)}{(1 - e^{-2\theta_n i} z^2)^\alpha} \right| \right\} \\ &\leq \|h_n \circ \psi_{\eta, \theta_n}^{-1}(z)\|_{H_0^\infty} \\ &\leq e^{C_2(\gamma, \eta)}. \end{aligned}$$

Now choose n large enough such that $r_\eta \leq t_{0,n} < 1$, we observe that

$$\begin{aligned} \|(T_g f_n)\|_{H_\beta^\infty} &= \sup_{z \in \mathbb{D}} \left\{ (1 - |z|^2)^\beta \left| \int_0^z f_n(\zeta) g'(\zeta) d\zeta \right| \right\} \\ &\geq (1 - t_{0,n}^2)^\beta \left| \int_0^{t_{0,n}} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) e^{i\theta_n} dr \right| \\ &\geq \left| \operatorname{Re} \left((1 - t_{0,n}^2)^\beta \int_{r_\eta}^{t_{0,n}} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) dr \right) \right| \\ &\quad - \left| \operatorname{Re} \left((1 - t_{0,n}^2)^\beta \int_0^{r_\eta} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) dr \right) \right| \end{aligned}$$

First, we see that

$$\begin{aligned} \left| \operatorname{Re} \left((1 - t_{0,n}^2)^\beta \int_0^{r_\eta} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) dr \right) \right| &\leq \int_0^{r_\eta} |f_n(re^{i\theta_n})| |g'(re^{i\theta_n})| dr \\ &\leq \|f_n\|_{H_\alpha^\infty} \int_0^{r_\eta} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \\ &\leq \frac{e^{C_2(\gamma, \eta)} C_3(\eta)}{(1 - r_\eta^2)^\alpha}, \end{aligned}$$

where $C_3(\eta) := \sup_{|z| \leq r_\eta} \{|g'(z)|\}$.

Next, to estimate another integral, we see that

$$\begin{aligned}
& \left| \operatorname{Re} \left((1 - t_{0,n}^2)^\beta \int_{r_\eta}^{t_{0,n}} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) dr \right) \right| \\
&= \left| (1 - t_{0,n}^2)^\beta \operatorname{Re} \left(\int_{r_\eta}^{t_{0,n}} \frac{g'(re^{i\theta_n})}{(1 - r^2)^\alpha} e^{-i(u_n(\psi_{\eta,\theta_n}^{-1}(re^{i\theta_n})) + i\tilde{u}_n(\psi_{\eta,\theta_n}^{-1}(re^{i\theta_n})))} dr \right) \right| \\
&= \left| (1 - t_{0,n}^2)^\beta \int_{r_\eta}^{t_{0,n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} e^{\tilde{u}_n(\psi_{\eta,\theta_n}^{-1}(re^{i\theta_n}))} \times \right. \\
&\quad \left. \operatorname{Re} \left(e^{i(\arg(g'(re^{i\theta_n})) - u_n(\psi_{\eta,\theta_n}^{-1}(re^{i\theta_n})))} \right) dr \right| \\
&\geq \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma,\eta)} \left| (1 - t_{0,n}^2)^\beta \int_{r_\eta}^{t_{0,n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \right| \\
&\geq \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma,\eta)} (1 - t_{0,n}^2)^\beta \left(\left| \int_0^{t_{0,n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \right| - \left| \int_0^{r_\eta} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \right| \right) \\
&\geq \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma,\eta)} \left(\left| (1 - t_{0,n}^2)^\beta \int_0^{t_{0,n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \right| - \left| \int_0^{r_\eta} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \right| \right) \\
&\geq \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma,\eta)} \left(n - \left| \int_0^{r_\eta} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \right| \right) \\
&\geq \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma,\eta)} \left(n - \frac{C_3(\eta)}{(1 - r_\eta^2)^\alpha} \right).
\end{aligned}$$

Hence, for n large enough such that $r_\eta \leq t_{0,n} < 1$, we have

$$\|(T_g f_n)\|_{H_\beta^\infty} \geq \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma,\eta)} n - e^{-C_2(\gamma,\eta)} \left(e^{2C_2(\gamma,\eta)} + \cos\left(\frac{\pi}{3}\right) \right) \frac{C_3(\eta)}{(1 - r_\eta^2)^\alpha}.$$

Letting $n \rightarrow \infty$, it follows that $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is not bounded.

Conversely, suppose that

$$\limsup_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t^2)^\beta \int_0^t \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr < +\infty,$$

then there exists $N > 0$, $0 < t_0 < 1$ such that

$$\sup_{0 \leq \theta < 2\pi} (1 - t^2)^\beta \int_0^t \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr \leq N \quad \text{whenever } t_0 < t < 1.$$

For any $f \in H_\alpha^\infty$, we have

$$\begin{aligned}
\|T_g(f)\|_{H_\beta^\infty} &= \sup_{z \in \mathbb{D}} \left\{ (1 - |z|^2)^\beta \left| \int_0^z f(\zeta) g'(\zeta) d\zeta \right| \right\} \\
&= \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R < 1} \left\{ (1 - R^2)^\beta \left| \int_0^{Re^{i\theta}} f(\zeta) g'(\zeta) d\zeta \right| \right\} \\
&= \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R < 1} \left\{ (1 - R^2)^\beta \left| \int_0^R f(re^{i\theta}) g'(re^{i\theta}) e^{i\theta} dr \right| \right\} \\
&\leq \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R \leq t_0} \left\{ (1 - R^2)^\beta \int_0^R |f(re^{i\theta})| |g'(re^{i\theta})| dr \right\} \\
&\quad + \sup_{0 \leq \theta < 2\pi} \sup_{t_0 < R < 1} \left\{ (1 - R^2)^\beta \int_0^R |f(re^{i\theta})| |g'(re^{i\theta})| dr \right\} \\
&\leq \|f\|_{H_\alpha^\infty} \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R \leq t_0} \left\{ (1 - R^2)^\beta \int_0^R \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr \right\} \\
&\quad + \|f\|_{H_\alpha^\infty} \sup_{0 \leq \theta < 2\pi} \sup_{t_0 < R < 1} \left\{ (1 - R^2)^\beta \int_0^R \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr \right\} \\
&\leq \|f\|_{H_\alpha^\infty} (M_{t_0} + N),
\end{aligned}$$

where $M_{t_0} := \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R \leq t_0} \left\{ (1 - R^2)^\beta \int_0^R \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr \right\}$. Accordingly, $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is bounded. \square

Remark 1. From the proof of Theorem 1, we see that the condition

$$\limsup_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t^2)^\beta \int_0^t \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr < +\infty$$

is sufficient for the boundedness of $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ without the requirement that $\log(g') \in \mathcal{B}$.

Remark 2. If $g \in H(\mathbb{D})$ is univalent, then by [18], it holds that $\log(g') \in \mathcal{B}$, thus Theorem 1 holds for the univalent case.

Remark 3. It should be noticed that Basallote et. al. [6] had given the following complete characterization of the boundedness of $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ for the general symbol $g \in H(\mathbb{D})$ when $\beta > 0$: $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is bounded if and only if

$$\sup_{z \in \mathbb{D}} (1 - |z|^2)^{\beta+1-\alpha} |g'(z)| < \infty.$$

As a corollary, we give the following result obtained by Smith, Stolyarov and Volberg in [20].

Corollary 1. *Let $g \in H(\mathbb{D})$ such that $\log(g') \in \mathcal{B}$. Then $T_g : H_0^\infty \rightarrow H_0^\infty$ is bounded if and only if*

$$\sup_{0 \leq \theta < 2\pi} \int_0^1 |g'(re^{i\theta})| dr < +\infty.$$

Proof. By Theorem 1, $T_g : H_0^\infty \rightarrow H_0^\infty$ is bounded if and only if

$$\limsup_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} \int_0^t |g'(re^{i\theta})| dr < +\infty.$$

Since $\int_0^t |g'(re^{i\theta})| dr$ is increasing as $t \rightarrow 1^-$, it follows that

$$\begin{aligned} \limsup_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} \int_0^t |g'(re^{i\theta})| dr &= \sup_{0 \leq \theta < 2\pi} \limsup_{t \rightarrow 1^-} \int_0^t |g'(re^{i\theta})| dr \\ &= \sup_{0 \leq \theta < 2\pi} \int_0^1 |g'(re^{i\theta})| dr. \end{aligned}$$

□

Recall that the disk algebra \mathcal{A} is the space consisting of all analytic functions f defined on \mathbb{D} which can be continuously extended to $\overline{\mathbb{D}}$ and endowed with the norm $\|f\| = \sup_{z \in \mathbb{D}} |f(z)|$.

Corollary 2. *Let $g \in H(\mathbb{D})$ such that $\log(g') \in \mathcal{B}$. Then T_g acting on the disk algebra \mathcal{A} is bounded if and only if $g \in \mathcal{A}$ and*

$$\sup_{0 \leq \theta < 2\pi} \int_0^1 |g'(re^{i\theta})| dr < +\infty.$$

Proof. From [4, Proposition 2.17] we see that for $g \in H(\mathbb{D})$, $T_g : \mathcal{A} \rightarrow \mathcal{A}$ is bounded if and only if $g \in \mathcal{A}$ and $T_g : H_0^\infty \rightarrow H_0^\infty$ is bounded. Thus, the corollary follows from Corollary 1. □

Proposition 1. *If $g \in H(\mathbb{D})$, then the following statements are equivalent:*

- (1) $S_g : \mathcal{A} \rightarrow \mathcal{A}$ is bounded;
- (2) $S_g : H_0^\infty \rightarrow H_0^\infty$ is bounded;
- (3) $T_g : H_0^\infty \rightarrow H_0^\infty$ is bounded.

Proof. From [4] we see that for $g \in H(\mathbb{D})$, $S_g : H_0^\infty \rightarrow H_0^\infty$ is bounded if and only if $T_g : H_0^\infty \rightarrow H_0^\infty$ is bounded.

Assume first that $S_g : H_0^\infty \rightarrow H_0^\infty$ is bounded, then for any polynomial p , since $gp' \in H_0^\infty$, we have

$$(S_g p)(z) = \int_0^z g(\zeta) p'(\zeta) d\zeta \in \mathcal{A}.$$

It is well known that polynomials are dense in \mathcal{A} , it follows that $S_g : \mathcal{A} \rightarrow \mathcal{A}$ is bounded.

Conversely, assume that $S_g : \mathcal{A} \rightarrow \mathcal{A}$ is bounded. Let $f \in H_0^\infty$ and $f_r(z) = f(rz)$ for $0 < r < 1$, then we have $f_r \in \mathcal{A}$. For any $z \in \mathbb{D}$,

$$|(S_g f)(z)| = \left| \lim_{r \rightarrow 1^-} (S_g f_r)(z) \right| \leq \lim_{r \rightarrow 1^-} \|S_g\| \|f_r\|_{H_0^\infty} \leq \|S_g\| \|f\|_{H_0^\infty},$$

which implies that $S_g : H_0^\infty \rightarrow H_0^\infty$ is bounded. □

Corollary 3. *Let $g \in H(\mathbb{D})$ such that $\log(g') \in \mathcal{B}$. Then S_g acting on the disk algebra \mathcal{A} (or H_0^∞) is bounded if and only if*

$$\sup_{0 \leq \theta < 2\pi} \int_0^1 |g'(re^{i\theta})| dr < +\infty.$$

Now for the boundedness of the companion operators $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$, due to Corollary 3, we just need to consider the case when $\alpha > 0$. In this case, since the weight $(1 - |z|^2)^\alpha$ is normal, we have $H_\alpha^\infty = \mathcal{B}_{\alpha+1}^\infty$, where $\mathcal{B}_\alpha^\infty$ is a weighted Bloch space defined as follows:

$$\mathcal{B}_\alpha^\infty = \{f \in H(\mathbb{D}) : \|f\|_{\mathcal{B}_\alpha^\infty} = \sup_{z \in \mathbb{D}} (1 - |z|^2)^\alpha |f'(z)| < \infty\}.$$

Keep in mind that $H_\alpha^\infty = \mathcal{B}_{\alpha+1}^\infty$ and consider the sequence $\{g_n\}_{n=1}^\infty$ of functions defined by

$$g_n(z) := \int_0^z \frac{f_n(\zeta) d\zeta}{(1 - e^{-2\theta n i \zeta^2})},$$

with f_n defined in the proof of Theorem 1, we have

Theorem 2. *Let $g \in H(\mathbb{D})$ such that $\log(g) \in \mathcal{B}$ and $0 < \alpha, 0 \leq \beta$. Then $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is bounded if and only if*

$$\limsup_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t^2)^\beta \int_0^t \frac{|g(re^{i\theta})|}{(1 - r^2)^{\alpha+1}} dr < +\infty.$$

The following corollary is the most interesting part of Theorem 2.

Corollary 4. *Let $g \in H(\mathbb{D})$ such that $\log(g) \in \mathcal{B}$ and $0 < \alpha$. Then $S_g : H_\alpha^\infty \rightarrow H_0^\infty$ is bounded if and only if*

$$\sup_{0 \leq \theta < 2\pi} \int_0^1 \frac{|g(re^{i\theta})|}{(1 - r^2)^{\alpha+1}} dr < +\infty.$$

If $\beta > 0$, then we can give the following complete characterization of the boundedness of $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ for the general symbol $g \in H(\mathbb{D})$.

Theorem 3. *If $g \in H(\mathbb{D})$ and $0 \leq \alpha, 0 < \beta$, then $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is bounded if and only if*

$$\sup_{z \in \mathbb{D}} (1 - |z|^2)^{\beta-\alpha} |g(z)| < \infty.$$

Proof. If $0 < \beta$, then $H_\beta^\infty = \mathcal{B}_{\beta+1}^\infty$. Thus the boundedness of $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is equivalent to the boundedness of $M_g D : H_\alpha^\infty \rightarrow H_{\beta+1}^\infty$, where M_g is the multiplication operator and D is the differentiation operator, respectively. Now by choosing $\varphi(z) \equiv z$ in [14, Theorem 7], the boundedness of $M_g D : H_\alpha^\infty \rightarrow H_{\beta+1}^\infty$ is equivalent to the following condition

$$\sup_{z \in \mathbb{D}} (1 - |z|^2)^{\beta-\alpha} |g(z)| < \infty.$$

□

3. The compactness of Volterra type operators

In this section, we firstly characterize the compactness of $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ and then consider the corresponding questions of its companion operator $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$. For $0 \leq \alpha$, we introduce the notation:

$$\overline{B} := \{f \in H_\alpha^\infty : \|f\|_{H_\alpha^\infty} \leq 1\}$$

By standard arguments, we prove the following lemma.

Lemma 3. *Let $0 \leq \alpha, \beta$. If $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is bounded, then the following two statements are equivalent:*

- (1) $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact;
- (2) For any bounded sequence $\{f_n\}_{n=1}^\infty \subset H_\alpha^\infty$ such that $f_n(z) \rightarrow 0$ uniformly on any compact subset of \mathbb{D} as $n \rightarrow \infty$, it holds that $\|T_g(f_n)\|_{H_\beta^\infty} \rightarrow 0$ as $n \rightarrow \infty$.

Proof. Assume that (2) holds. Let $\{k_n\}_{n=1}^\infty \subset \overline{B}$, then $\{k_n\}$ is a normal family and thus, there exists $k \in H_\alpha^\infty$ and a subsequence $\{k_{n_j}\}_{j=1}^\infty$ such that $k_{n_j} \rightarrow k$ uniformly on any compact subset of \mathbb{D} as $j \rightarrow \infty$. Let $\{f_j\}_{j=1}^\infty = \{k_{n_j} - k\}_{j=1}^\infty$. By (2), $\|T_g(f_j)\|_{H_\beta^\infty} = \|T_g(k_{n_j}) - T_g(k)\|_{H_\beta^\infty} \rightarrow 0$ as $j \rightarrow \infty$. Since by assumption $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is bounded, it follows that $T_g(k) \in H_\beta^\infty$. Accordingly, $T_g(k_{n_j}) \rightarrow T_g(k)$ in H_β^∞ as $j \rightarrow \infty$, which implies that $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact.

Conversely, assume that $\{f_n\}_{n=1}^\infty \subset \overline{B}$, that $f_n \rightarrow 0$ uniformly on any compact subset of \mathbb{D} as $n \rightarrow \infty$, and that $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact. To prove (2), it suffices to show that there exists a subsequence $\{f_{n_j}\}_{j=1}^\infty$ such that $f_{n_j} \rightarrow 0$ as $j \rightarrow \infty$. It is easy to show that $T_g(f_n) \rightarrow 0$ uniformly on any compact subset of \mathbb{D} as $n \rightarrow \infty$. Since $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact, there is a subsequence $\{f_{n_j}\}_{j=1}^\infty$ and $f \in H_\beta^\infty$ such that $\|T_g(f_{n_j}) - f\|_{H_\beta^\infty} \rightarrow 0$ as $j \rightarrow \infty$. Since $T_g(f_{n_j}) \rightarrow 0$ uniformly on any compact subset of \mathbb{D} as $j \rightarrow \infty$, it follows that $f = 0$ and hence $\|T_g(f_{n_j})\|_{H_\beta^\infty} \rightarrow 0$ as $j \rightarrow \infty$. This completes the proof. \square

Theorem 4. *Let $g \in H(\mathbb{D})$ such that $\log(g') \in \mathcal{B}$ and $0 \leq \alpha, \beta$. Then $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact if and only if*

$$\lim_{t_2 \rightarrow 1^-} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t_1^2)^\beta \int_{t_2}^{t_1} \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr = 0.$$

Proof. Assume first that

$$\lim_{t_2 \rightarrow 1^-} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t_1^2)^\beta \int_{t_2}^{t_1} \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr = 0.$$

To prove that $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact, choose any sequence $\{f_n\}_{n=1}^\infty \subset H_\alpha^\infty$ such that there exists a positive number W such that $\sup_{n \in \mathbb{N}} \|f_n\|_{H_\alpha^\infty} \leq W$ and $f_n \rightarrow 0$ uniformly on any compact subset of \mathbb{D} as $n \rightarrow \infty$. Let $\epsilon > 0$, then there exists $t_{2,\epsilon}$ with $0 < t_{2,\epsilon} < 1$ such that

$$\limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t_1^2)^\beta \int_{t_{2,\epsilon}}^{t_1} \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr < \epsilon.$$

Moreover, there exists $t_{1,\epsilon}$ with $t_{2,\epsilon} < t_{1,\epsilon} < 1$ such that

$$\sup_{0 \leq \theta < 2\pi} (1 - t_1^2)^\beta \int_{t_{2,\epsilon}}^{t_1} \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr < 2\epsilon, \quad \text{whenever } t_{1,\epsilon} < t_1 < 1.$$

Now choose $N_\epsilon > 0$ such that

$$\sup_{|z| \leq t_{1,\epsilon}} |f_n(z)| < \frac{\epsilon}{M_{1,\epsilon} \cdot t_{1,\epsilon}}, \quad \text{whenever } n > N_\epsilon,$$

where $M_{1,\epsilon} := \sup_{0 \leq |z| \leq t_{1,\epsilon}} \{|g'(z)|\}$.

Then for $n > N_\epsilon$, we have

$$\begin{aligned}
\|T_g(f_n)\|_{H_\beta^\infty} &= \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R < 1} \left\{ (1-R^2)^\beta \left| \int_0^R f_n(re^{i\theta}) g'(re^{i\theta}) e^{i\theta} dr \right| \right\} \\
&\leq \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R < 1} \left\{ (1-R^2)^\beta \int_0^R |f_n(re^{i\theta})| |g'(re^{i\theta})| dr \right\} \\
&\leq \sup_{0 \leq \theta < 2\pi} \sup_{0 \leq R \leq t_{1,\epsilon}} \left\{ (1-R^2)^\beta \int_0^R |f_n(re^{i\theta})| |g'(re^{i\theta})| dr \right\} \\
&\quad + \sup_{0 \leq \theta < 2\pi} \sup_{t_{1,\epsilon} < R} \left\{ (1-R^2)^\beta \int_0^{t_{2,\epsilon}} |f_n(re^{i\theta})| |g'(re^{i\theta})| dr \right\} \\
&\quad + \sup_{0 \leq \theta < 2\pi} \sup_{t_{1,\epsilon} < R} \left\{ (1-R^2)^\beta \int_{t_{2,\epsilon}}^R |f_n(re^{i\theta})| |g'(re^{i\theta})| dr \right\} \\
&\leq \epsilon + \epsilon + 2W\epsilon = 2(1+W)\epsilon,
\end{aligned}$$

which implies that $\lim_{n \rightarrow \infty} \|T_g(f_n)\|_{H_\beta^\infty} = 0$ and hence, by Lemma 3, $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact.

Conversely, suppose that $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact and so it is bounded. If

$$c := \lim_{t_2 \rightarrow 1^-} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1-t_1^2)^\beta \int_{t_2}^{t_1} \frac{|g'(re^{i\theta})|}{(1-r^2)^\alpha} dr > 0,$$

then by Theorem 1, we see that $c < +\infty$.

Let $\{t_{2,n}\}_{n=1}^\infty$ be a sequence such that $0 < t_{2,n} < t_{2,n+1} < 1$, $\lim_{n \rightarrow \infty} t_{2,n} = 1$ and

$$\begin{aligned}
c &= \inf_{n \in \mathbb{N}} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1-t_1^2)^\beta \int_{t_{2,n}}^{t_1} \frac{|g'(re^{i\theta})|}{(1-r^2)^\alpha} dr \\
&= \lim_{n \rightarrow \infty} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1-t_1^2)^\beta \int_{t_{2,n}}^{t_1} \frac{|g'(re^{i\theta})|}{(1-r^2)^\alpha} dr.
\end{aligned}$$

Then for any $n \in \mathbb{N}$, we can choose $t_{1,n}$ with $t_{2,n} < t_{1,n}$ such that

$$\sup_{0 \leq \theta < 2\pi} (1-t_{1,n}^2)^\beta \int_{t_{2,n}}^{t_{1,n}} \frac{|g'(re^{i\theta})|}{(1-r^2)^\alpha} dr > c - \frac{1}{n^2}.$$

Also, we can choose an angle θ_n with $0 \leq \theta_n < 2\pi$ such that

$$(1-t_{1,n}^2)^\beta \int_{t_{2,n}}^{t_{1,n}} \frac{|g'(re^{i\theta_n})|}{(1-r^2)^\alpha} dr > c - \frac{1}{n}.$$

Define the sequence $\{k_n\}_{n=1}^\infty$ by choosing k_n as the largest positive integer such that $t_{2,n}^{k_n} \geq \frac{1}{3}$ for any $n \in \mathbb{N}$, then since $\lim_{n \rightarrow \infty} \frac{\log \frac{1}{3}}{\log t_{2,n}} = +\infty$, it follows that $\lim_{n \rightarrow \infty} k_n = +\infty$.

Now define the sequence $\{s_n(z)\}_{n=1}^\infty$ by

$$s_n(z) := f_n(z) z^{k_n}, \quad z \in \mathbb{D},$$

where $f_n(z)$ is defined in the proof of Theorem 1, that is,

$$f_n(z) := \frac{h_n \circ \psi_{\eta, \theta_n}^{-1}(z)}{(1 - e^{-2\theta_n i} z^2)^\alpha}.$$

Thus,

$$\|s_n\|_{H_\infty^\alpha} \leq e^{C_2(\gamma, \eta)},$$

and $s_n \rightarrow 0$ uniformly on any compact subset of \mathbb{D} .

Choosing n large enough such that $t_{2,n} > r_\eta$, then

$$\begin{aligned} \|T_g(s_n)\|_{H_\beta^\infty} &\geq \left| (1 - t_{1,n}^2)^\beta \int_0^{t_{1,n}} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) r^{k_n} dr \right| \\ &\geq \left| \operatorname{Re} \left((1 - t_{1,n}^2)^\beta \int_0^{t_{1,n}} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) r^{k_n} dr \right) \right| \\ &\geq \left| \operatorname{Re} \left((1 - t_{1,n}^2)^\beta \int_{r_\eta}^{t_{1,n}} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) r^{k_n} dr \right) \right| \\ &\quad - \left| \operatorname{Re} \left((1 - t_{1,n}^2)^\beta \int_0^{r_\eta} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) r^{k_n} dr \right) \right| \end{aligned}$$

First, we have

$$\begin{aligned} \left| \operatorname{Re} \left((1 - t_{1,n}^2)^\beta \int_0^{r_\eta} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) r^{k_n} dr \right) \right| &\leq \int_0^{r_\eta} |f_n(re^{i\theta_n})| |g'(re^{i\theta_n})| r^{k_n} dr \\ &\leq \|f_n\|_{H_\infty^\alpha} \int_0^{r_\eta} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} r^{k_n} dr \\ &\leq \frac{e^{C_2(\gamma, \eta)} C_3(\eta)}{(1 - r_\eta^2)^\alpha} r_\eta^{k_n}, \end{aligned}$$

where $C_3(\eta) := \sup_{|z| \leq r_\eta} \{|g'(z)|\}$.

Next, to estimate another integral, we obtain

$$\begin{aligned} &\left| \operatorname{Re} \left((1 - t_{1,n}^2)^\beta \int_{r_\eta}^{t_{1,n}} f_n(re^{i\theta_n}) g'(re^{i\theta_n}) r^{k_n} dr \right) \right| \\ &= \left| (1 - t_{1,n}^2)^\beta \operatorname{Re} \left(\int_{r_\eta}^{t_{1,n}} \frac{g'(re^{i\theta_n})}{(1 - r^2)^\alpha} e^{-i(u_n(\psi_{\eta, \theta_n}^{-1}(re^{i\theta_n})) + i\tilde{u}_n(\psi_{\eta, \theta_n}^{-1}(re^{i\theta_n})))} r^{k_n} dr \right) \right| \\ &= \left| (1 - t_{1,n}^2)^\beta \int_{r_\eta}^{t_{1,n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} e^{\tilde{u}_n(\psi_{\eta, \theta_n}^{-1}(re^{i\theta_n}))} \times \right. \\ &\quad \left. \operatorname{Re} \left(e^{i(\arg(g'(re^{i\theta_n})) - u_n(\psi_{\eta, \theta_n}^{-1}(re^{i\theta_n})))} \right) r^{k_n} dr \right| \\ &\geq \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma, \eta)} \left| (1 - t_{1,n}^2)^\beta \int_{r_\eta}^{t_{1,n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} r^{k_n} dr \right| \\ &\geq \cos\left(\frac{\pi}{3}\right) r_{2,n}^{k_n} e^{-C_2(\gamma, \eta)} \left| (1 - t_{1,n}^2)^\beta \int_{t_{2,n}}^{t_{1,n}} \frac{|g'(re^{i\theta_n})|}{(1 - r^2)^\alpha} dr \right| \\ &\geq \frac{1}{3} \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma, \eta)} \left(c - \frac{1}{n}\right). \end{aligned}$$

Therefore, it follows that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \|T_g(s_n)\|_{H_\beta^\infty} &\geq \lim_{n \rightarrow \infty} \left(\frac{1}{3} \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma, \eta)} \left(c - \frac{1}{n}\right) - \frac{e^{C_2(\gamma, \eta)} C_3(\eta)}{(1 - r_\eta^2)^\alpha} r_\eta^{k_n} \right) \\ &= \frac{1}{3} \cos\left(\frac{\pi}{3}\right) e^{-C_2(\gamma, \eta)} c > 0, \end{aligned}$$

which, according to Lemma 3, is in contradiction with the assumption that $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact. Thus the proof is complete. \square

Remark 4. From the proof of Theorem 4, we also see that the condition

$$\lim_{t_2 \rightarrow 1^-} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} (1 - t_1^2)^\beta \int_{t_2}^{t_1} \frac{|g'(re^{i\theta})|}{(1 - r^2)^\alpha} dr = 0$$

is sufficient for the compactness of $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ without the requirement that $\log(g') \in \mathcal{B}$.

Remark 5. It should be noticed that Basallote et. al. [6] had given the following complete characterization of the compactness of $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ for the general symbol $g \in H(\mathbb{D})$ when $\beta > 0$: $T_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact if and only if

$$\lim_{|z| \rightarrow 1^-} (1 - |z|^2)^{\beta+1-\alpha} |g'(z)| = 0.$$

Corollary 5. Let $g \in H(\mathbb{D})$ such that $\log(g') \in \mathcal{B}$. Then the following statements are equivalent:

- (1) $T_g : H_0^\infty \rightarrow H_0^\infty$ is compact;
- (2) $T_g : \mathcal{A} \rightarrow \mathcal{A}$ is compact;
- (3) $\lim_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} \int_t^1 |g'(re^{i\theta})| dr = 0$.

Proof. By Theorem 4, $T_g : H_0^\infty \rightarrow H_0^\infty$ is compact if and only if

$$\lim_{t_2 \rightarrow 1^-} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} \int_{t_2}^{t_1} |g'(re^{i\theta})| dr = 0.$$

Since $\int_{t_2}^{t_1} |g'(re^{i\theta})| dr$ is increasing as $t_1 \rightarrow 1^-$, it follows that

$$\begin{aligned} \lim_{t_2 \rightarrow 1^-} \limsup_{t_1 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} \int_{t_2}^{t_1} |g'(re^{i\theta})| dr &= \lim_{t_2 \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} \limsup_{t_1 \rightarrow 1^-} \int_{t_2}^{t_1} |g'(re^{i\theta})| dr \\ &= \lim_{t \rightarrow 1^-} \sup_{0 \leq \theta < 2\pi} \int_t^1 |g'(re^{i\theta})| dr. \end{aligned}$$

The equivalence of (1) and (2) follows from [8, Theorem 1.7]. \square

Now for the compactness of the companion operators $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$, we have the following complete characterization for the general symbol $g \in H(\mathbb{D})$.

Theorem 5. If $g \in H(\mathbb{D})$ and $0 \leq \alpha, \beta$, then the following statements hold:

- (1) $S_g : H_\alpha^\infty \rightarrow H_0^\infty$ is compact if and only if $g = 0$;
- (2) If $0 < \beta$, then $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is compact if and only if

$$\lim_{|z| \rightarrow 1^-} (1 - |z|^2)^{\beta-\alpha} |g(z)| = 0.$$

Proof. (1) Anderson et. al. [4] had proved that $S_g : H_0^\infty \rightarrow H_0^\infty$ is compact if and only if $g = 0$. Since $H_0^\infty \subset H_\alpha^\infty$ for any $0 \leq \alpha$, it follows that $S_g : H_\alpha^\infty \rightarrow H_0^\infty$ is compact if and only if $g = 0$.

(2) If $0 < \beta$, then $H_\beta^\infty = \mathcal{B}_{\beta+1}^\infty$. Thus the compactness of $S_g : H_\alpha^\infty \rightarrow H_\beta^\infty$ is equivalent to the compactness of $M_g D : H_\alpha^\infty \rightarrow H_{\beta+1}^\infty$, where M_g is the multiplication operator and D is the differentiation operator, respectively. Now by choosing $\varphi(z) \equiv z$ in [14, Theorem 8], the compactness of $M_g D : H_\alpha^\infty \rightarrow H_{\beta+1}^\infty$ is equivalent to the following condition

$$\lim_{|z| \rightarrow 1^-} (1 - |z|^2)^{\beta-\alpha} |g(z)| = 0.$$

□

REFERENCES

- [1] A. Aleman, J. Cima, *An integral operator on H^p and Hardy's inequality*, J. Anal. Math. 85 (2001), 157-176.
- [2] A. Aleman, A. Siskakis, *An integral operator on H^p* , Complex Variables Theory Appl. 28 (1995), no.2, 149-158.
- [3] A. Aleman, A. Siskakis, *Integration operators on Bergman spaces*, Indiana Univ. Math. J. 46 (1997), no.2, 337-356.
- [4] A. Anderson, M. Jovovic, W. Smith, *Some integral operators acting on H^∞* , Integral Equations Operator Theory 80 (2014), no. 2, 275-291.
- [5] A. Anderson, *Some closed range integral operators on spaces of analytic functions*, Integral Equations Operator Theory 69 (2011), 87-99.
- [6] M. Basallote, M. Contreras, C. Hernández-Mancera, M. Martín, P. Paúl, *Volterra operators and semigroups in weighted Banach spaces of analytic functions*, Collect. Math. 65 (2014), no. 2, 233-249.
- [7] O. Constantin, J. Peláez, *Integral operators, embedding theorems and a Littlewood-Paley formula on weighted Fock spaces*, J. Geom. Anal. 26 (2016), no. 2, 1109-1154.
- [8] M. Contreras, J. Peláez, C. Pommerenke, J. Rättyä, *Integral operators mapping into the space of bounded analytic functions*, J. Funct. Anal. 271 (2016), no. 10, 2899-2943.
- [9] T. Eklund, M. Lindström, M. Pirasteh, A. Sanatpour, N. Wikman, *Generalized Volterra operators mapping between Banach spaces of analytic functions*, Monatsh. Math. (2018). <https://doi.org/10.1007/s00605-018-1216-5>.
- [10] G. Hardy, J. Littlewood, *Some properties of fractional integrals. II*, Math. Z. 34 (1932), no. 1, 403-439.
- [11] J. Laitila, S. Miihkinen, P. Nieminen, *Essential norms and weak compactness of integration operators*, Arch. Math. 97 (2011), no. 1, 39-48.
- [12] P. Li, J. Liu, Z. Lou, *Integral operators on analytic Morrey spaces*, Sci. China Math. 57 (2014), no. 9, 1961-1974.
- [13] Q. Lin, J. Liu, Y. Wu, *Volterra type operators on $S^p(\mathbb{D})$ spaces*, J. Math. Anal. Appl. 461 (2018), 1100-1114.
- [14] J. Manhas, R. Zhao, *Products of weighted composition operators and differentiation operators between Banach spaces of analytic functions*, Acta Sci. Math. (Szeged) 80 (2014), 665-679.
- [15] T. Mengestie, *Product of Volterra type integral and composition operators on weighted Fock spaces*, J. Geom. Anal. 24 (2014), no. 2, 740-755.
- [16] S. Miihkinen, *Strict singularity of a Volterra-type integral operator on H^p* , Proc. Amer. Math. Soc. 145 (2017), no. 1, 165-175.
- [17] A. Montes-Rodríguez, *Weighted composition operators on weighted Banach spaces of analytic functions*, J. London Math. Soc. 61 (2000), no. 2, 872-884.
- [18] Ch. Pommerenke, *Boundary behaviour of conformal maps*, Grundlehren der Mathematischen Wissenschaften 299, Springer-Verlag, Berlin (1992).
- [19] Ch. Pommerenke, *Schlichte Funktionen und analytische Funktionen von beschränkter mittlerer Oszillation*, (German) Comment. Math. Helv. 52 (1977), no. 4, 591-602.
- [20] W. Smith, D. Stolyarov, A. Volberg, *Uniform approximation of Bloch functions and the boundedness of the integration operator on H^∞* , Adv. Math. 314 (2017), 185-202.

- [21] J. Xiao, *The Q_p Carleson measure problem*, Adv. Math. 217 (2008), no.5, 2075-2088.

SCHOOL OF APPLIED MATHEMATICS, GUANGDONG UNIVERSITY OF TECHNOLOGY, GUANGZHOU,
GUANGDONG, 510520, P. R. CHINA
E-mail address: `gdlqz@e.gzhu.edu.cn`