

Two types of permutation polynomials with special forms

Dabin Zheng^a, Mu Yuan^a, Long Yu^b

^a*Hubei Province Key Laboratory of Applied Mathematics,
Faculty of Mathematics and Statistics, Hubei University, Wuhan 430062, China*
^b*School of Mathematics and Physics, Hubei Polytechnic University,
Huangshi 435003, China*

Abstract

Let q be a power of a prime and \mathbb{F}_q be a finite field with q elements. In this paper, we propose four families of infinite classes of permutation trinomials having the form $cx - x^s + x^{qs}$ over \mathbb{F}_{q^2} , and investigate the relationship between this type of permutation polynomials with that of the form $(x^q - x + \delta)^s + cx$. Based on this relation, many classes of permutation trinomials having the form $(x^q - x + \delta)^s + cx$ without restriction on δ over \mathbb{F}_{q^2} are derived from known permutation trinomials having the form $cx - x^s + x^{qs}$.

Keywords: Finite field, Permutation polynomial, Symbolic computation

1. Introduction

Let q be a power of a prime. Let \mathbb{F}_q be a finite field with q elements and \mathbb{F}_q^* denote its multiplicative group. A polynomial $f(x) \in \mathbb{F}_q[x]$ is called a *permutation polynomial* (PP) if its associated polynomial mapping $f : c \mapsto f(c)$ from \mathbb{F}_q to itself is a bijection. Permutation polynomials over finite fields have important applications in cryptography, coding theory and combinatorial design theory. So, finding new permutation polynomials is of great interest in both theoretical and applied aspects. The study of permutation polynomials has a long history [6, 12] and many recent results are surveyed in [14].

Permutation polynomials with few terms attracts many researchers' attention due to their simple algebraic representation and wide applications in

Email addresses: dzheng@hubu.edu.cn (Dabin Zheng), yuanmu847566@outlook.com (Mu Yuan), longyu@mails.ccn.edu.cn (Long Yu)

coding theory, combinatorial designs and cryptography. The recent progress on construction of permutation binomials and trinomials can be seen in [2, 7, 11, 19, 18, 20, 21, 22, 23, 25, 38] and the references therein. Permutation polynomials of the form $x + \gamma \text{Tr}_q^{q^n}(x^k)$ were investigated in [3, 4] for some special $\gamma \in \mathbb{F}_{q^n}$. Very recently, G.Kyureghyan and M.E. Zieve [15] found almost all permutation polynomials over \mathbb{F}_{q^n} of this form for $\gamma \in \mathbb{F}_{q^n}^*$ and $q^n \leq 5000$, where q is odd. And Li et al. [18] studied permutation behavior of this type of polynomials over the finite fields with even characteristic. In both [15] and [18], most permutation polynomials having the form $x + \gamma \text{Tr}_q^{q^n}(x^k)$ occur in case of $n = 2$, i.e., they are trinomial permutations. Inspired by this, we construct four families of infinite classes of permutation trinomials over \mathbb{F}_{q^2} with the following form

$$cx - x^s + x^{qs}, \quad (1)$$

where s is a positive integer and $c \in \mathbb{F}_{q^2}$. We use the well-known result (Lemma 1) provided by Park, Lee, Wang and Zieve to prove the first three classes of permutation trinomials. For the proof of permutation behavior of the last class of trinomials, symbolic computation method related to Gröbner bases and resultants is used.

On the other hand, in order to derive new Kloosterman sums identities over \mathbb{F}_{q^n} , Helleseth and Zinoviev [13] found a class of permutation polynomials related to the form

$$(x^{q^i} - x + \delta)^s + cx \quad (2)$$

over \mathbb{F}_{q^n} , and Ding and Yuan [9] first studied this type of permutation polynomials. Following line of work in [9], many permutation polynomials with this form have been constructed [5, 21, 26, 27, 29, 31, 32, 33, 34, 35, 36, 37]. Most known permutation polynomials having this form are related to δ . In this paper, we find a close relationship between the two types of permutation polynomials with the forms (1) and (2) respectively. Based on this relationship, many classes of permutation polynomials having the form (2) without restriction on δ are derived from known permutation polynomials of the form (1).

The remainder of this paper is organized as follows. In section 2, we introduce some preliminary results and investigate the relationship between the two types of permutation polynomials with the forms (1) and (2) respectively. Section 3 presents four families of infinite classes of permutation trinomials with the form (1). In section 4, many new classes of permutation polynomials with the form (2) are derived from known permutation polynomials with the form (1). Finally, concluding remark is given in section 5.

2. Preliminaries

Let q be a power of a prime, and \mathbb{F}_q be a finite field with q elements. Let k be a divisor of n . The trace function from \mathbb{F}_{q^n} to \mathbb{F}_{q^k} is defined by

$$\mathrm{Tr}_{q^k}^{q^n}(x) = \sum_{i=0}^{n/k-1} x^{q^{ik}},$$

where $x \in \mathbb{F}_{q^n}$. Let d be a positive integer with $d \mid (q-1)$, and μ_d denote the set of d th roots of unity in \mathbb{F}_q^* , i.e.,

$$\mu_d = \left\{ x \in \mathbb{F}_q^* \mid x^d = 1 \right\}.$$

Many people investigate the permutation behavior of the polynomials with the form $x^r h(x^s)$ for $s \mid (q-1)$ over \mathbb{F}_q . The following lemma is well-known which transforms the problem of proving $x^r h(x^s)$ permutes \mathbb{F}_q into the problem of investigating whether it permutes $\mu_{(q-1)/d}$.

Lemma 1. [24][28][39] *Let d, r be positive integers with $d \mid (q-1)$, and let $h(x) \in \mathbb{F}_q[x]$. Then $x^r h(x^{(q-1)/d})$ permutes \mathbb{F}_q if and only if both*

- (1) $\gcd(r, (q-1)/d) = 1$ and
- (2) $x^r h(x)^{(q-1)/d}$ permutes $\mu_d := \{x \in \mathbb{F}_q^* \mid x^d = 1\}$.

The following proposition shows a close relationship between the two types of permutation polynomials with the forms (1) and (2).

Proposition 2. *Let m, k be integers with $0 < k < m$ and $\ell = \gcd(k, m)$. Let \mathbb{F}_{q^m} be a finite field and $g(x) \in \mathbb{F}_{q^m}[x]$. For $\delta \in \mathbb{F}_{q^m}$ and $c \in \mathbb{F}_{q^\ell}^*$, if $h(x) = g(x)^{q^k} - g(x) + cx$ permutes \mathbb{F}_{q^m} then $f(x) = g(x^{q^k} - x + \delta) + cx$ permutes \mathbb{F}_{q^m} .*

PROOF. For $\alpha \in \mathbb{F}_{q^m}$, it is sufficient to show that the equation

$$f(x) = g(x^{q^k} - x + \delta) + cx = \alpha \tag{3}$$

has only one solution in \mathbb{F}_{q^m} . Denote by $y = x^{q^k} - x + \delta$. Taking q^k -th power on both sides of (3) and subtracting (3) we have

$$h(y) = g(y)^{q^k} - g(y) + cy = \alpha^{q^k} - \alpha + c\delta.$$

Since h permutes \mathbb{F}_{q^m} we obtain $y = h^{-1}(\alpha^{q^k} - \alpha + c\delta)$. So, the only one solution of (3) in \mathbb{F}_{q^m} is $x = c^{-1}(\alpha - g \circ h^{-1}(\alpha^{q^k} - \alpha + c\delta))$.

Remark 3. In Proposition 2, let $m = 2, k = 1$ and $g(x) = x^s$ and $c \in \mathbb{F}_q^*$. By Proposition 2, if $cx - x^s + x^{qs}$ permutes \mathbb{F}_{q^2} then $(x^q - x + \delta)^s + cx$ permutes \mathbb{F}_{q^2} . Moreover, if there are no restrictions on δ and $c = 1$ then the inverse of Proposition 2 also holds.

Proposition 4. Let \mathbb{F}_{q^m} be the degree m extension of the finite field \mathbb{F}_q . Let $g(x) \in \mathbb{F}_q[x]$ be a polynomial over \mathbb{F}_q . For any $\delta \in \mathbb{F}_{q^m}$, $f(x) = g(x^q - x + \delta) + x$ permutes \mathbb{F}_{q^m} if and only if $h(x) = g(x)^q - g(x) + x$ permutes \mathbb{F}_{q^m} .

PROOF. By Proposition 2 we only prove the necessity. Denote by $\varphi(x) = x^q - x + \delta$, $\alpha = \text{Tr}_q^{q^m}(\delta)$, and $S_\alpha = \{x \in \mathbb{F}_{q^m} \mid \text{Tr}_q^{q^m}(x) = \alpha\}$. By Theorem 2.25 in [16] we have that $S_\alpha = \{x^q - x + \delta \mid x \in \mathbb{F}_{q^m}\}$. For $\alpha, \alpha' \in \mathbb{F}_q$ with $\alpha \neq \alpha'$ we have that $S_\alpha \cap S_{\alpha'} = \emptyset$ and $\cup_{\alpha \in \mathbb{F}_q} S_\alpha = \mathbb{F}_{q^m}$.

It is easy to verify that $\varphi \circ f(x) = h \circ \varphi(x)$ for any $x \in \mathbb{F}_{q^m}$, i.e., the following diagram is commutative,

$$\begin{array}{ccc} \mathbb{F}_{q^m} & \xrightarrow{f} & \mathbb{F}_{q^m} \\ \downarrow \varphi & & \downarrow \varphi \\ S_\alpha & \xrightarrow{h} & S_\alpha \end{array}$$

If $f(x)$ is a bijection on \mathbb{F}_{q^m} , then from above diagram we know that $h(x)$ is a surjective mapping from S_α to itself. So, $h(x)$ is a bijection on S_α for any $\alpha \in \mathbb{F}_q$, moreover $\mathbb{F}_{q^m} = \cup_{\alpha \in \mathbb{F}_q} S_\alpha$. Thus, $h(x)$ permutes \mathbb{F}_{q^m} . \square

3. Four classes of permutation polynomials with the form $cx - x^s + x^{sq}$ over \mathbb{F}_{q^2}

In this section we present four classes of permutation polynomials with the form $cx - x^s + x^{sq}$ for some proper index s and parameter $c \in \mathbb{F}_{q^2}^*$.

Theorem 5. Let $s = \frac{3q^2+2q-1}{4}$ and $c \in \mathbb{F}_{q^2}^*$. The polynomial $f(x) = cx - x^s + x^{qs}$ permutes \mathbb{F}_{q^2} in each of the following cases:

- (1) $q \equiv 1 \pmod{8}$ and $(-\frac{2}{c})^{\frac{q+1}{2}} = 1$;
- (2) $q \equiv 5 \pmod{8}$ and $(\frac{2}{c})^{\frac{q+1}{2}} = 1$.

PROOF. We only prove the case (1). The proof of the case (2) is similar and the details are omitted.

Since $q \equiv 1 \pmod{8}$ we know that $4 \mid (3q + 5)$. The polynomial $f(x)$ is rewritten as

$$f(x) = cx - x^{\frac{3q+5}{4}(q-1)+1} + x^{(\frac{3q+5}{4}q+1)(q-1)+1}.$$

Let $u = \frac{3q+5}{4}$. By Lemma 1, $f(x)$ permutes \mathbb{F}_{q^2} if and only if

$$g(x) = x(c + x^{1-u} - x^u)^{q-1}$$

permutes μ_{q+1} .

Let $\mu_{\frac{q+1}{2}}$ be the cyclic subgroup of μ_{q+1} with order $\frac{q+1}{2}$. It is easy to verify that $\mu_{\frac{q+1}{2}}$ is exactly all square elements of μ_{q+1} . Since $q \equiv 1 \pmod{8}$ we have that -1 is non-square element of μ_{q+1} . Then $-\mu_{\frac{q+1}{2}}$ is the set of all non-square elements of μ_{q+1} . Next we show that $g(x)$ permutes $\mu_{\frac{q+1}{2}}$ and $-\mu_{\frac{q+1}{2}}$ respectively.

Since $q \equiv 1 \pmod{8}$ we have that squaring is a bijection on $\mu_{\frac{q+1}{2}}$. For $x \in \mu_{\frac{q+1}{2}}$, there exists a unique $y \in \mu_{\frac{q+1}{2}}$ such that $x = y^2$. Then

$$x^u = (y^2)^u = y^{\frac{3q+5}{2}} = y, \quad x^{1-u} = x \cdot x^{-u} = y^2 \cdot y^{-1} = y.$$

In this case, $g(x) = c^{q-1}x$, and it permutes $\mu_{\frac{q+1}{2}}$.

For $x \in -\mu_{\frac{q+1}{2}}$, there exists a unique $y \in \mu_{\frac{q+1}{2}}$ such that $x = -y^2$ with the same reason above. Due to $(-\frac{2}{c})^{\frac{q+1}{2}} = 1$, i.e., $\frac{c}{2} \in -\mu_{\frac{q+1}{2}}$ and $y \in \mu_{\frac{q+1}{2}}$ we have that $c - 2y \neq 0$. So,

$$\begin{aligned} g(x) &= g(-y^2) = -y^2(c + (-1)^{1-u}y - (-1)^uy)^{q-1} \\ &= -y^2 \frac{(c - 2y)^q}{c - 2y} = -y^2 \frac{c^q - 2y^{-1}}{c - 2y} \\ &= \frac{2}{c}y. \end{aligned}$$

Since $\frac{c}{2} \in -\mu_{\frac{q+1}{2}}$ and $y \in \mu_{\frac{q+1}{2}}$ we have $\frac{2}{c}y \in -\mu_{\frac{q+1}{2}}$. So, $g(x)$ permutes $-\mu_{\frac{q+1}{2}}$. Combining these two cases, we have that $g(x)$ permutes μ_{q+1} . \square

Similarly, we obtain another family of infinite class of permutation polynomials over \mathbb{F}_{q^2} as follows.

Theorem 6. *Let $s = \frac{(q+1)^2}{4}$ and $c \in \mathbb{F}_{q^2}^*$. The polynomial $f(x) = cx - x^s + x^{qs}$ permutes \mathbb{F}_{q^2} in each of the following cases:*

- (1) $q \equiv 5 \pmod{8}$ and $\left(-\frac{2}{c}\right)^{\frac{q+1}{2}} = 1$;
(2) $q \equiv 1 \pmod{8}$ and $\left(\frac{2}{c}\right)^{\frac{q+1}{2}} = 1$.

By direct verification and simplification we get a family of infinite class of permutation polynomials over \mathbb{F}_{q^2} as follows.

Theorem 7. *Let q be a power of a prime with $q \equiv 1 \pmod{3}$, and $s = \frac{q^2+q+1}{3}$. Then $f(x) = x - x^s + x^{qs}$ is a permutation polynomial over \mathbb{F}_{q^2} .*

PROOF. Since $q \equiv 1 \pmod{3}$ we know that $3 \mid (q+2)$. The polynomial $f(x)$ is rewritten as

$$f(x) = x - x^{\frac{q+2}{3}(q-1)+1} + x^{\left(\frac{q(q+2)}{3}+1\right)(q-1)+1}.$$

Let $u = \frac{q+2}{3}$. By Lemma 1, $f(x)$ permutes \mathbb{F}_{q^2} if and only if

$$g(x) = x(1 + x^{1-u} - x^u)^{q-1}$$

permutes μ_{q+1} .

First, we show that $1 + x^{1-u} - x^u = 0$ has no roots in μ_{q+1} . Otherwise, assume that there is $x \in \mu_{q+1}$ satisfying this equation, i.e.,

$$1 + x^{1-u} - x^u = 0, \quad x^{q+1} = 1. \quad (4)$$

These imply $x^{2u-1} - x^{u-1} = x^{q+1}$, i.e., $x^{2u-2} - x^{u-2} = x^q$. This leads to $x^{2-2u} - x^{2-u} = x$, then $x^{1-2u} - x^{1-u} = 1$, i.e., $x^{2u-1} + x^u = 1$. This equality together with (4) implies $x^{u-1} + x^u = 0$, i.e., $x = -1$. On the other hand, it is easy to verify that $x = -1$ does not satisfy the first equation in (4) since $3 \nmid q$. So, we get a contradiction.

For any $x \in \mu_{q+1}$, we have $x^{3u} = x$ since $u = \frac{q+2}{3}$, and

$$g(x) = x \frac{1 + x^{u-1} - x^{-u}}{1 + x^{1-u} - x^u} = \frac{x + x^u - x^{1-u}}{1 + x^{1-u} - x^u} = \frac{x^u + x^{3u} - x^{2u}}{1 + x^{2u} - x^u} = x^u.$$

So, $g(x)$ permutes μ_{q+1} since $\gcd(u, q+1) = 1$. □

The main technique in the proof of above theorems is provided in Lemma 1. This method doesn't work for the following theorem. Using the method provided by Dobbertin [8] and Kyureghyan and Zieve [15] we present a family of infinite class of permutation polynomials over \mathbb{F}_{q^4} . To this end, we first give two preliminary lemmas.

Lemma 8. *There exists $x \in \mathbb{F}_{q^4}^*$ such that $x^{2q^2} + x^{q^2+1} + x^2 = 0$ if and only if $3 \mid q$ and $x^{q^2-1} = 1$.*

PROOF. Assume that $x \in \mathbb{F}_{q^4}^*$ satisfies

$$x^{2q^2} + x^{q^2+1} + x^2 = 0. \quad (5)$$

Let $y = x^{q^2-1}$, then $y \in \mu_{q^2+1}$ and $y^2 + y + 1 = 0$. It follows that $y^3 = 1$. Since $\gcd(3, q^2 + 1) = 1$, we must have $y = 1$ and $3 \mid q$. Conversely, it is obvious. \square

Similarly, we have the following lemma.

Lemma 9. *There exists $x \in \mathbb{F}_{q^4}^*$ such that $x^{2q^2} - x^{q^2+1} + x^2 = 0$ if and only if $3 \mid q$ and $x^{q^2-1} = -1$.*

Theorem 10. *Let q be a power of an odd prime and $s = q^3 + q^2 - q$. Then the polynomial $f(x) = x - x^s + x^{q^2s}$ permutes \mathbb{F}_{q^4} .*

PROOF. We prove that $f(x) = \alpha$ has only one root in \mathbb{F}_{q^4} for any $\alpha \in \mathbb{F}_{q^4}$. To this end, we discuss the proof according to the following four situations.

Case I: $\alpha = 0$ and $f(x) = 0$. It is clear that this equation has a solution $x = 0$. Next we show that there is no $x \in \mathbb{F}_{q^4}^*$ satisfying $f(x) = 0$. Otherwise, we have

$$1 - x^{q^3+q^2-q-1} + x^{-q^3+q} = 0.$$

Denote by $y = x^{q^2-1}$, then $y \in \mu_{q^2+1}$. From above equation we get

$$y^{q+1} - y^{-q} = 1. \quad (6)$$

Calculation of $(6)^q - (6)^{q^3}$ together with $y^{q^2} = y^{-1}$, we obtain

$$\begin{aligned} 0 &= y^{q^2+q} - y^{-q^2} - y^{q^3+1} + y^{-1} \\ &= y^{q-1} - y - y^{-q+1} + y^{-1} \\ &= (y^{q-1} - y)(y^{-q} + 1). \end{aligned}$$

It follows that $y^{q-2} = 1$ since $y \neq -1$. This fact together with $y^{q^2+1} = 1$ implies that

$$y^{\gcd(q-2, q^2+1)} = y^{\gcd(q-2, 5)} = 1.$$

So, we have $y^5 = 1$. On the other hand, substituting $y^{q-2} = 1$ into (6) we get $y^5 = y^2 + 1$. This together with $y^5 = 1$ implies that $y = 0$. This is a contradiction. So, $f(x) = 0$ has only one solution $x = 0$ in \mathbb{F}_{q^4} .

To discuss case of $\alpha \neq 0$, we define two sets as follows,

$$S_{\pm} = \left\{ x \in \mathbb{F}_{q^4}^* \mid x^{2q^2} \pm x^{q^2+1} + x^2 = 0 \right\}.$$

Case II: $\alpha \in S_+$ and $f(x) = \alpha$. By Lemma 8, we have $3 \mid q$ and $\alpha^{q^2} - \alpha = 0$. In this case we have

$$\begin{aligned} 0 &= \alpha^{q^2} - \alpha = f(x)^{q^2} - f(x) \\ &= x^{q^2} - x^{-q^3+q+1} + x^{q^3+q^2-q} - (x - x^{q^3+q^2-q} + x^{-q^3+q+1}) \\ &= x^{q^2} - x - x^{q^3+q^2-q} + x^{-q^3+q+1} \\ &= x^{-q^3-q}(x - x^{q^2})^q(x^{q^2+q^3} + x^{1+q}). \end{aligned}$$

If $x - x^{q^2} = 0$ then $f(x) = x = \alpha$. If $x^{q^2+q^3} = -x^{1+q}$ then $x^{(q+1)(q^2-1)} = -1$. Since $x \in \mathbb{F}_{q^4}^*$, we have $x^{\gcd(2(q+1)(q^2-1), q^4-1)} = x^{2(q+1)} = 1$. It implies that $x^{(1+q)(q^2-1)} = 1$. This leads to a contradiction. Thus $f(x) = \alpha$ has only one root $x = \alpha$.

Case III: $\alpha \in S_-$ and $f(x) = \alpha$. By similar proof to Case II, we know that $f(x) = \alpha$ has only one solution $x = \alpha$.

Denote by $T = \mathbb{F}_{q^4}^* \setminus (S_+ \cup S_-)$. From the proving process of Cases II and III, we know that $f(x)$ is a permutation over S_+ and S_- respectively. Moreover, $f(T) \subseteq T$. It remains to show that $f(x)$ is also a permutation over the set T .

Case IV: $\alpha \in T$. We will show that $f(x) = \alpha$ has only one $x \in T$ satisfying this equation. Denote by

$$y = x^q, \quad z = x^{q^2}, \quad w = x^{q^3}, \quad \beta = \alpha^q, \quad \gamma = \alpha^{q^2}, \quad \delta = \alpha^{q^3}.$$

The equation $f(x) = \alpha$ is reduced to

$$x + \frac{xy}{w} - \frac{zw}{y} = \alpha. \quad (7)$$

Taking q th, q^2 th and q^3 th power on both sides of (7) respectively, we obtain

$$y + \frac{yz}{x} - \frac{wx}{z} = \beta, \quad (8)$$

$$z + \frac{zw}{y} - \frac{yx}{w} = \gamma, \quad (9)$$

$$w + \frac{wx}{z} - \frac{yz}{x} = \delta, \quad (10)$$

respectively. Adding (7) and (9) we have $z = \alpha + \gamma - x$. Similarly, we get $w = \beta + \delta - y$ from (8) and (10). Substituting the two equalities into (7) and (8) respectively, we obtain

$$\begin{cases} (-\gamma + x)y^2 + ((\alpha + 2\gamma - x)(\beta + \delta))y + (-\alpha - \gamma + x)(\beta + \delta)^2 = 0 \\ (x^2 - (\alpha + \gamma)x + (\alpha + \gamma)^2)y - \delta x^2 - \beta(\alpha + \gamma)x = 0. \end{cases} \quad (11)$$

Since $x \in T$, from Lemma 8 we have

$$x^2 - (\alpha + \gamma)x + (\alpha + \gamma)^2 = x^2 + xz + z^2 \neq 0.$$

From the second equation in (11) we get

$$y = \frac{x(\beta\alpha + \beta\gamma + \delta x)}{\alpha^2 + 2\alpha\gamma + \gamma^2 - (\alpha + \gamma)x + x^2} \triangleq H(x). \quad (12)$$

Substituting this y into the first equation in (11), we obtain a equality $A(x) = 0$, where $A(x) \in \mathbb{F}_{q^4}^*[\alpha, \beta, \gamma, \delta][x]$ and its degree on x is 5.

Suppose that $f(x) = \alpha$ have another root $X \in T$ different from x , then $A(X) = 0$. Replacing $\alpha, \beta, \gamma, \delta$ in coefficients of $A(X)$ with x, y, z, w respectively, and multiplying wx^2yz^2 on both sides of $A(X) = 0$, we get

$$(X - x)B(X)C(X) = 0, \quad (13)$$

where

$$\begin{aligned} B(X) &= yw(x^2 + xz + z^2)X^2 + wxz(z + x)^2(w + y) \\ &\quad - (z + x)(w^2xz + wx^2y + wxyz + wyz^2 - xy^2z)X, \\ C(X) &= (w^2x^2 - wxyz + y^2z^2)X^2 + yz(z + x)^3(w + y) \\ &\quad + (z + x)(w^2xz + wx^2y + wxyz - wyz^2 - xy^2z - 2y^2z^2)X. \end{aligned}$$

Hence $B(X) = 0$ or $C(X) = 0$. It has been verified that $yw(x^2 + xz + z^2) \neq 0$ above. Assume that $w^2x^2 - wxyz + y^2z^2 = 0$. Then

$$0 = w^2x^2 - wxyz + y^2z^2 = (yz)^{2q^2} - (yz)^{q^2+1} + (yz)^2. \quad (14)$$

By Lemma 9 we have $(yz)^{q^2-1} = -1$, i.e., $y^{(q^2-1)(q+1)} = -1$. Since $\gcd(2(q^2-1)(q+1), q^4-1) = 2(q^2-1)$, we have $y^{2(q^2-1)} = 1$. This is a contradiction with $y^{(q^2-1)(q+1)} = -1$. So, the degree of $B(X)$ and $C(X)$ on X is 2, respectively.

(i) When $B(X) = 0$ in (13). Taking q th power on both sides of this equality we get $B'(x^q) = 0$, where the coefficients of $B'(X)$ are q th power of

corresponding coefficients of $B(X)$. Replacing $Y = X^q$ with $H(X)$ in (12) we obtain

$$0 = B'(H(X)) = C(X)D(X), \quad (15)$$

where

$$\begin{aligned} D(X) = & (w^2x^2 + w^2xz + zxwy + xy^2z + y^2z^2)X^2 + wx^2(z+x)^2(w+y) \\ & - (z+x)(2w^2x^2 + w^2xz + wx^2y + zxwy - wyz^2 + xy^2z)X. \end{aligned}$$

If $C(X) = 0$ from (15) then X is a common root of $B(X)$ and $C(X)$. Then the resultant of these two polynomials should be 0. So,

$$\begin{aligned} 0 = Res(B, C, X) = & zx(z+x)^4(w+y)^2(w^2x^2 - zxwy + y^2z^2) \\ & \times (w^2xz + w^2z^2 + wxyz + x^2y^2 + xy^2z)^{q+1}. \end{aligned} \quad (16)$$

It has been verified that $w^2x^2 - zxwy + y^2z^2 \neq 0$ above. Moreover, we have that $(z+x)(w+y) \neq 0$. In fact, if $x+z=0$ or $w+y=0$ we can derive that $B(X) = D(X) = x^2y^2X^2 = 0$. From (16) we have

$$w^2xz + w^2z^2 + wxyz + x^2y^2 + xy^2z = 0. \quad (17)$$

Combining (17) and $B(X) = 0$ we obtain that

$$yw(x^2 + xz + z^2)(X - x)^2 = 0.$$

This is a contradiction with that $x^2 + xz + z^2 \neq 0$ and $X \neq x$.

So, from (15) we must have $D(X) = 0$. If the leading coefficient of $D(X)$ is nonzero, i.e., $w^2x^2 + w^2xz + zxwy + xy^2z + y^2z^2 \neq 0$ then

$$0 = Res(B, D, X) = xwyz(z+x)^6(w+y)^2(w^2x^2 - zxwy + y^2z^2)^{q+1}.$$

This is impossible from the discussion in previous paragraph. Thus, the leading coefficient of $D(X)$ is zero, i.e.,

$$L(x) = w^2x^2 + w^2xz + zxwy + xy^2z + y^2z^2 = 0. \quad (18)$$

The fact that $wx^2(z+x)^2(w+y) \neq 0$ implies that the degree of $D(X)$ is 1. The resultant of $B(X)$ and $D(X)$ is zero. Then

$$0 = Res(B(X), D(X), X) = (z+x)^4xw(w+y)U(x), \quad (19)$$

where

$$\begin{aligned} U(x) = & -yw^3x^5 + (2zw^4 + yzw^3 + 2y^2zw^2x^4)x^4 + (3z^2w^4 + 3yz^2w^3 \\ & + 6y^2z^2w^2 + 2y^3z^2w + y^4z^2)x^3 + (z^3w^4 - 2yz^3w^3 + y^2z^3w^2 \\ & + y^4z^3)x^2 - (2yz^4w^3 + y^2z^4w^2 + 2y^3z^4w)x + y^2z^5w^2. \end{aligned}$$

From (19) we have $U(x) = 0$. By calculation of the resultant of $L(x)$ and $U(x)$ we get

$$0 = \text{Res}(L(x), U(x), x) = (w^2 + y^2)^2(w + y)^6 w^4 z^{10} y^4. \quad (20)$$

In this case, it is verified that $y + w \neq 0$ above. From (20) we have $w^2 = -y^2$, i.e., $y^{2(q^2-1)} = -1$ and $y^{4(q^2-1)} = 1$. Then $y^{\gcd(4(q^2-1), q^4-1)} = y^{2(q^2-1)} = 1$. This is a contradiction. Therefore, we prove that $B(X)$ can not be zero.

(ii) Similarly, we can prove that $C(x)$ in (13) is not zero. Thus, $f(x) = \alpha$ has only one root in T , and $f(x)$ permutes T . \square

4. Permutation polynomials with the form $(x^q - x + \delta)^s + cx$ over \mathbb{F}_{q^2}

In this section we give many classes of permutation polynomials having the form $(x^q - x + \delta)^s + cx$ without restriction on δ over \mathbb{F}_{q^2} . These permutation polynomials are derived from known permutation trinomials with the form (1) by using Proposition 2.

4.1. Permutation polynomials over finite fields with odd characteristic

First, four families of infinite classes of permutation polynomials of the form $(x^q - x + \delta)^s + cx$ over \mathbb{F}_{q^2} with odd characteristic are derived from permutation trinomials introduced in Section 3.

Theorem 11. *Let $\delta \in \mathbb{F}_{q^2}$ and $c \in \mathbb{F}_q^*$. The polynomial*

$$f(x) = (x^q - x + \delta)^{\frac{3q^2+2q-1}{4}} + cx$$

is a permutation over \mathbb{F}_{q^2} in each of the following cases:

- (1) $q \equiv 1 \pmod{8}$ and $c = -2$;
- (2) $q \equiv 5 \pmod{8}$ and $c = 2$.

PROOF. Let $s = \frac{3q^2+2q-1}{4}$ and $g(x) = x^s$. From Theorem 5 we know that $h(x) = cx + g(x)^s - g(x)^{qs}$ permutes \mathbb{F}_{q^2} under the case of $q \equiv 1 \pmod{8}$ and $(-2/c)^{\frac{q+1}{2}} = 1$ or the case of $q \equiv 5 \pmod{8}$ and $(2/c)^{\frac{q+1}{2}} = 1$ respectively. These are exact cases listed in (1) and (2) respectively since $c \in \mathbb{F}_q$. By Proposition 2 we obtain that $f(x)$ permutes \mathbb{F}_{q^2} under one of conditions listed above. \square

Similarly, one easily verifies the following three theorems.

Theorem 12. Let $\delta \in \mathbb{F}_{q^2}$ and $c \in \mathbb{F}_q^*$. The polynomial

$$f(x) = (x^q - x + \delta)^{\frac{(q+1)^2}{4}} + cx$$

is a permutation over \mathbb{F}_{q^2} in each of the following cases:

- (1) $q \equiv 1 \pmod{8}$ and $c = 2$;
- (2) $q \equiv 5 \pmod{8}$ and $c = -2$.

Theorem 13. Let q be a power of a prime with $q \equiv 1 \pmod{3}$. For $\delta \in \mathbb{F}_{q^2}$, the polynomial

$$f(x) = (x^q - x + \delta)^{\frac{q^2+q+1}{3}} + x$$

is a permutation over \mathbb{F}_{q^2} .

Theorem 14. Let q be a power of an odd prime. For any $\delta \in \mathbb{F}_{q^2}$, the polynomial

$$f(x) = (x^q - x + \delta)^{q^3+q^2-q} + x$$

is a permutation over \mathbb{F}_{q^4} .

4.2. Permutation polynomials over finite fields with even characteristic

By Proposition 2 many classes of permutation polynomials having the form $(x^q + x + \delta)^s + cx$ without restriction on δ are derived from known permutation trinomials having the form $cx + x^s + x^{qs}$ over \mathbb{F}_{q^2} with even characteristic. To this end, we first recall known permutation trinomials with the form $cx + x^s + x^{qs}$ over finite fields with even characteristic in the following lemmas.

Lemma 15. [17] Let k be a positive integer and $q = 2^k$. The trinomials $f(x) = cx + x^s + x^{qs}$ are permutation polynomials over \mathbb{F}_{q^2} in each of the following cases.

- (1) $s = 2q - 1$. The positive integer k and $c \in \mathbb{F}_{q^2}$ satisfy one of the following conditions:
 - i) k is even and $c = 1$;
 - ii) k is odd and $c^3 = 1$.
- (2) $s = \frac{(3q-2)(q^2+q+1)}{3}$, and k is even and $c \in \mathbb{F}_{q^2}$ satisfies $c^3 = 1$.
- (3) $s = \frac{q+4}{5}$, and k is odd and $c \in \mathbb{F}_{q^2}$ satisfies $c^3 = 1$.
- (4) $s = \frac{3q+1}{4}$, and $c \in \mathbb{F}_q$ satisfies that $x^3 + x + c = 0$ having no solution in \mathbb{F}_q .

- (5) $s = \frac{q+6}{7}$ and $c = 1$.
- (6) $s = \frac{q^2+3q+2}{6}$, and k is odd and $c \in \mathbb{F}_{q^2}$ satisfies $c^{\frac{q+1}{3}} = 1$.
- (7) $s = \frac{q^2-2q+4}{3}$, and k is even and $c = 1$.
- (8) $s = \frac{Q^3+Q^2-Q+1}{2}$ and $Q = 2^{\frac{k}{2}}$ for an even k , and $c \in \mathbb{F}_Q^*$.

Lemma 16. [19][1] Let k be a positive integer and $q = 2^k$. The trinomials $f(x) = cx + x^s + x^{qs}$ are permutation polynomials over \mathbb{F}_{q^2} in each of the following cases.

- (1) $s = \frac{-1}{2^{k'}-1}(2^k - 1) + 1$, where k' is a positive integer with $\gcd(2^{k'} - 1, 2^k + 1) = 1$ and $c \in \mathbb{F}_{2^{k'}}^* \cap \mathbb{F}_q$.
- (2) $s = \frac{1}{2^{k'}+1}(2^k - 1) + 1$, where k' is a positive integer with $\gcd(2^{k'} + 1, 2^k + 1) = 1$ and $c \in \mathbb{F}_{2^{k'}} \cap \mathbb{F}_q$.

Remark 17. It is easy to see that each s in Lemmas 15 and 16 can be rewritten as the form $s = i(q - 1) + 1$, here i is interpreted as modulo $q + 1$ when it is negative or proper fraction.

By Proposition 2, the coefficient c of x in polynomials of the form (2) should be in \mathbb{F}_q^* , and the permutation polynomials in the following theorem are derived directly from permutation trinomials in Lemmas 15 and 16.

Theorem 18. Let k be a positive integer and $q = 2^k$. For $\delta \in \mathbb{F}_{q^2}$, $f(x) = (x^q + x + \delta)^s + cx$ permutes \mathbb{F}_{q^2} in each of the following cases:

- (1) $s = 2q - 1$ and $c = 1$.
- (2) $s = \frac{(3q-2)(q^2+q+1)}{3}$ and k is even, and $c \in \mathbb{F}_q$ satisfies $c^3 = 1$.
- (3) $s = \frac{q+4}{5}$, $c = 1$ and k is odd.
- (4) $s = \frac{3q+1}{4}$, and $c \in \mathbb{F}_q$ satisfies that $x^3 + x + c = 0$ having no solution in \mathbb{F}_q .
- (5) $s = \frac{q+6}{7}$ and $c = 1$.
- (6) $s = \frac{q^2+3q+2}{6}$, $c = 1$ and k is odd.
- (7) $s = \frac{q^2-2q+4}{3}$, $c = 1$ and k is even.
- (8) $s = \frac{Q^3+Q^2-Q+1}{2}$ and $Q = 2^{\frac{k}{2}}$ for an even k , and $c \in \mathbb{F}_Q^*$.
- (9) $s = \frac{-1}{2^{k'}-1}(2^k - 1) + 1$, and k' is a positive integer with $\gcd(2^{k'} - 1, 2^k + 1) = 1$, and $c \in \mathbb{F}_{2^{k'}} \cap \mathbb{F}_q$.
- (10) $s = \frac{1}{2^{k'}+1}(2^k - 1) + 1$, and k' is a positive integer with $\gcd(2^{k'} + 1, 2^k + 1) = 1$, and $c \in \mathbb{F}_{2^{k'}} \cap \mathbb{F}_q$.

Remark 19. *The permutation polynomial in case (1) has been proposed in Theorem 1 of [26] and Theorem 3.2 of [29]. The permutation polynomial in case (3) has been constructed in [10]. The permutation polynomial in case (4) has been constructed in Theorem 2.1 of [37] and Theorem 3.1 of [29]. Even though the representation of permutation polynomials having the form (2) in those papers may be different, they have the same corresponding trinomials of the form (1) if s is written as $i(q-1)+1$ and i is interpreted as modulo $q+1$ when it is negative or proper fraction. As far as we know, Table 1 presents all permutation polynomials having the form $(x^q+x+\delta)^s+cx$ without restriction on δ over \mathbb{F}_{q^2} with even characteristic, where $\Omega = \{c \in \mathbb{F}_q \mid x^3+x+c=0 \text{ has no roots in } \mathbb{F}_q\}$.*

Table 1: PPs of the form $(x^{2^k}+x+\delta)^s+cx$ without restriction on δ over $\mathbb{F}_{2^{2k}}$

No.	k	$s = i(2^k - 1) + 1$	c	Reference
1	k is even	$i = 2$ or -1	$c = 1$	[26][29]
2	all k	$i = 0$ or 1	$c = 1$	[29][31]
3	all k	$i = \frac{1}{2}$	$c = 1$	[29]
4	k is even	$i = \frac{1}{3}$ or $\frac{2}{3}$	$c = 1$	[30]
5	k is odd	$i = \frac{1}{5}$ or $\frac{4}{5}$	$c = 1$	[10]
6	a positive integer	$i = \frac{1}{4}$ or $\frac{3}{4}$	$c \in \Omega$	[37][29]
7	k is even	$i = \frac{1}{2^k-2}$ or $\frac{-4}{2^k-2}$	$c^3 = 1$	Theorem 18
8	a positive integer	$i = \frac{1}{7}$ or $\frac{6}{7}$	$c = 1$	Theorem 18
9	k is odd	$i = \frac{2^k+4}{6}$ or $\frac{2-2^k}{6}$	$c^{\frac{q+1}{3}} = 1$	Theorem 18
10	k is even	$i = \frac{2^k-1}{3}$ or $\frac{4-2^k}{3}$	$c = 1$	Theorem 18
11	k is even	$i = \frac{2^{k/2}+1}{2}$ or $\frac{1-2^{k/2}}{2}$	$c \in \mathbb{F}_{2^{k/2}}^*$	Theorem 18
12	$\gcd(2^{k'}-1, 2^k+1) = 1$	$i = \frac{-1}{2^{k'}-1}$ or $\frac{2^{k'}}{2^{k'}-1}$	$c \in \mathbb{F}_{2^k}^* \cap \mathbb{F}_{2^{k'}}$	Theorem 18
13	$\gcd(2^{k'}+1, 2^k+1) = 1$	$i = \frac{1}{2^{k'}+1}$ or $\frac{2^{k'}}{2^{k'}+1}$	$c \in \mathbb{F}_{2^k}^* \cap \mathbb{F}_{2^{k'}}$	Theorem 18

5. Concluding remark

In this paper, we proposed four families of infinite classes of permutation trinomials with the form (1). From known permutation trinomials with the form (1), many classes of permutation polynomials with the form (2) are derived. As part of the future work, observe that the relationship between the two types of polynomials can be generalized to the case of multiple

monomials. It would be interesting to find more permutation polynomials with the form (1) or (2) by using this relation.

References

- [1] D. Bartoli, L. Quoos, Permutation polynomials of the type $x^r g(x^s)$ over $\mathbb{F}_{q^{2n}}$, *Designs Codes Cryptography*, Doi:10.1007/s10623-017-0415-8.
- [2] L.A. Bassalygo, V.A. Zinoviev, Permutation and complete permutation polynomials, *Finite Field Appl.* 33(2015) 198-211.
- [3] P. Charpin, G. Kyureghyan, On a Class of Permutation Polynomials over \mathbb{F}_{2^n} , *Sequence and their Application-SETA 2008*, *Lecture Notes in Comput. Sci.* 5203, Springer,(2008) 368-376.
- [4] P. Charpin, G. Kyureghyan, Monomial functions with linear structure and permutation polynomials, *Finite Fields: Theory and Applications*, *Contemp. Math.* 518, Amer. Math. Soc. 3(16)(2010) 99-111.
- [5] N. Cepak, P. Charpin, E. Pasalic, Permutations via linear translators, *Finite Fields Appl.* 45(2017) 19-42.
- [6] L.E. Dickson, The analytic representation of substitutions on a power of a prime number of letters with a discussion of the linear group, *Ann. Math.* 11(1896) 65-120.
- [7] C. Ding, L. Qu, Q. Wang, J. Yuan, P. Yuan, Permutation trinomials over finite fields with even characteristic, *SIAM J. Discrete Math.* 29(2015) 79-92.
- [8] H. Dobbertin, Uniformly representable permutation polynomials. *Sequence and their Applications-SETA 2001*, Springer, 2(9)(2002) 1-22.
- [9] J. Yuan, C. Ding, Four classes of permutation polynomials of \mathbb{F}_{2^m} , *Finite Fields Appl.* 13(4)(2007) 869-876.
- [10] R. Gupta, R.K. Sharma, Further results on permutation polynomials of the form $(x^{p^m} - x + \delta)^s + x$ over $\mathbb{F}_{p^{2m}}$, *Finite Fields Appl.* 50(2018) 196-208.
- [11] R. Gupta, R.K. Sharma, Some new classes of permutation trinomials over finite fields with even characteristic, *Finite Fields Appl.* 41(2016) 89-96.

- [12] Ch. Hermite, Sur les fonctions de sept letters, C. R. Acad. Sci. Paris, 15(1863) 750-757.
- [13] T. Helleseth, V. Zinoviev, New Kloosterman sums identities over \mathbb{F}_{2^m} for all m , Finite Fields Appl. 9(2)(2003) 187-193.
- [14] X. Hou, Permutation polynomials over finite fields—A survey of recent advances, Finite Fields Appl. 32(2015) 82-119.
- [15] G. Kyureghyan, M.E. Zieve, Permutation polynomials of the form $X + \gamma\text{Tr}(X^k)$, Contemporary Developments in Finite Fields and Appl. doi:10.1142/9789814719261-0011 (2016)
- [16] R. Lidl, H. Niederreiter, Finite Fields, Encycl. Math. Appl., Cambridge University Press, 1997.
- [17] K. Li, L. Qu, X. Chen, C. Li, Permutation polynomials of the form $cx + \text{Tr}_q^{q^l}(x^a)$ and permutation trinomials over finite fields with even characteristic, Cryptogr. Commun. (2017) 1-24.
- [18] K. Li, L. Qu, Q. Wang, New Constructions of Permutation Polynomials of the Form $x^r h(x^{q-1})$ over \mathbb{F}_{q^2} , arXiv:1708.01165v1 [cs.IT], (2017).
- [19] N. Li, T. Helleseth, New Permutation Trinomials From Niho Exponents over Finite Fields with Even Characteristic, arXiv:1606.03768, 2016.
- [20] N. Li, T. Helleseth, Several classes of permutation trinomials from Niho exponents, Cryptogr. Commun. (2016) 1-13.
- [21] N. Li, T. Helleseth, X. Tang, Further results on a class of permutation polynomials over finite fields, Finite Fields Appl. 22(2013) 16-23.
- [22] K. Li, L. Qu, X. Chen, New classes of permutation bionmials and permutation trinomials over finite fields, Finite Fields Appl. 43(2017) 69-85.
- [23] J. Ma, G. Ge, A note on permutation polynomials over finite fields, Finite Fields Appl. 48(2017) 261-270.
- [24] Y.H. Park, J.B. Lee, Permutation polynomials and group permutation polynomials, Bull. Aust. Math. Soc, 63(2001) 67-74.

- [25] Z. Tu, X. Zeng, L. Hu, Several classes of complete permutation polynomials, *Finite Fields Appl.* 25(2014) 182-193.
- [26] Z. Tu, X. Zeng, X. Jiang, Two classes of permutation polynomials having the form $(x^{2^m} + x + \delta)^s + x$. *Finite Fields Appl.* 31(2015) 12-24.
- [27] Z. Tu, X. Zeng, C. Li, T. Helleseth, Permutation polynomials of the form $(x^{p^m} - x + \delta)^s + L(x)$ over the finite field $\mathbb{F}_{p^{2m}}$ of odd characteristic, *Finite Fields Appl.* 34(2015) 2-35.
- [28] Q. Wang, Cyclotomic mapping permutation polynomials over finite fields, *Proceedings of the 2007 international conference on Sequences, subsequences, and consequences (SSC 2007)*, Pages 119-128, Los Angeles, CA, USA, 2007.
- [29] L. Wang, B. Wu, Z. Liu, Further results on permutation polynomials of the form $(x^p - x + \delta)^s + x$ over $\mathbb{F}_{p^{2m}}$, *Finite Fields Appl.* 44(2017) 92-112.
- [30] G. Xu, X. Cao, S. Xu, Further results on permutation polynomials of the form $(x^{p^m} - x + \delta)^s + L(x)$ over $\mathbb{F}_{p^{2m}}$, *J. Algebra Appl.* 15(2016) 1650098.
- [31] P. Yuan, C. Ding, Permutation polynomials over finite fields from a powerful lemma, *Finite Fields Appl.* 17(6)(2011) 560-574.
- [32] P. Yuan, C. Ding, Further results on permutation polynomials over finite fields, *Finite Fields Appl.* 27(2014) 88-103.
- [33] P. Yuan, Y. Zheng, Permutation polynomials from piecewise functions, *Finite Fields Appl.* 35(2015) 215-230.
- [34] D. Zheng, Z. Chen, More classes of permutation polynomials of the form $(x^{2^m} + x + \delta)^s + L(x)$, *App. Algebra Engrg. Comm. Comput.* 28(3)(2017) 215-223.
- [35] X. Zeng, X. Zhu, L. Hu, Two new permutation polynomials with the form $(x^{2^m} + x + \delta)^s + x$ over \mathbb{F}_{2^n} , *App. Algebra Engrg. Comm. Comput.* 21(2)(2010) 145-150.
- [36] Z. Zha, L. Hu, Two classes of permutation polynomials over finite fields, *Finite Fields Appl.* 18(4)(2012) 781-790.

- [37] Z. Zha, L. Hu, Some classes of permutation polynomials of the form $(x^{p^m} - x + \delta)^s + x$ over $\mathbb{F}_{p^{2m}}$, *Finite Fields Appl.* 40(2016) 150-162.
- [38] Z. Zha, L. Hu, S. Fan, Further results on permutation trinomials over finite fields with even characteristic, *Finite Fields Appl.* 45(2017) 43-52.
- [39] M.E. Zieve, On some permutation polynomials over \mathbb{F}_q of the form $x^r h(x^{(q-1)/d})$, *Proc. Am. Math. Soc.* 137(2009) 209-216.