

Upper bound for the first non-zero eigenvalue of the p -Laplacian

Sheela Verma
 Department of Mathematics & Statistics
 Indian Institute of Technology Kanpur
 Kanpur, India
 sheela@iitk.ac.in

Abstract

Let M be a closed hypersurface in \mathbb{R}^n and Ω be a bounded domain such that $M = \partial\Omega$. In this article, we obtain an upper bound for the first non-zero eigenvalue of the following problems.

- Closed eigenvalue problem:

$$\Delta_p u = \lambda_p |u|^{p-2} u \quad \text{on } M.$$

- Steklov eigenvalue problem:

$$\begin{aligned} \Delta_p u &= 0 & \text{in } \Omega, \\ |\nabla u|^{p-2} \frac{\partial u}{\partial \nu} &= \mu_p |u|^{p-2} u & \text{on } M. \end{aligned}$$

Keywords: p -Laplacian, Closed eigenvalue problem, Steklov eigenvalue problem, Center-of-mass.
Mathematics subject classification: 35P15, 58J50.

1 Introduction

The p -Laplace operator, defined as $\Delta_p u := -\operatorname{div}(|\nabla u|^{p-2} \nabla u)$, is the nonlinear generalization of the usual Laplace operator.

Many interesting results, providing the sharp upper bounds for the first non zero eigenvalue of the usual Laplacian ($p = 2$) have been obtained. In [4], Bleeker and Weiner obtained a sharp upper bound of the first non-zero eigenvalue of Laplacian in terms of the second fundamental form on a hypersurface M in \mathbb{R}^n . In [5], Reilly gave an upper bound for the first non-zero eigenvalue in terms of higher order mean curvatures for a compact n -dimensional manifold isometrically immersed in \mathbb{R}^{n+p} , which improves the earlier estimate. This result was later extended to submanifolds of simply connected space forms in various ways (see [6, 10]). These upper bounds are extrinsic in the sense that they depend either on the length of the second fundamental form or the higher order mean curvatures of M .

Let M be a hypersurface in a rank-1 symmetric space. In [11], an upper bound for the first non-zero eigenvalue of M was obtained in terms of the integral of the first non-zero eigenvalue of the geodesic spheres centered at the centre of gravity of M .

For a closed hypersurface M contained in a ball of radius less than $\frac{i(\mathbb{M}(k))}{4}$ and bounding a convex domain such that $\partial\Omega = M$ in the simply connected space form $\mathbb{M}(k)$, $k = 0$ or 1 , Santhanam [12] proved that

$$\frac{\lambda_1(M)}{\lambda_1(S(R))} \leq \frac{\operatorname{Vol}(M)}{\operatorname{Vol}(S(R))},$$

where $S(R) (= \partial B(R))$ is the geodesic sphere of radius $R > 0$ such that $\text{Vol}(B(R)) = \text{Vol}(\Omega)$. A similar result was also obtained for $k = -1$.

In this article, we extend the results in [12] to p -Laplacian for a closed hypersurface $M \subset \mathbb{R}^n$. In particular, we consider the closed eigenvalue problem

$$\Delta_p u = \lambda_p |u|^{p-2} u \quad \text{on } M, \quad (1)$$

where M is a closed hypersurface in \mathbb{R}^n and find an upper bound for the first non-zero eigenvalue of this problem.

Let M be a closed hypersurface in \mathbb{R}^n and Ω be the bounded domain such that $M = \partial\Omega$. Consider the following problem

$$\begin{aligned} \Delta f &= 0 & \text{in } \Omega, \\ \frac{\partial f}{\partial \nu} &= \mu f & \text{on } \partial\Omega, \end{aligned}$$

where ν is the outward unit normal on the boundary $\partial\Omega$ and μ is a real number. This problem is known as Steklov eigenvalue problem and was introduced by Steklov [1] for bounded domains in the plane in 1902. This problem is important as the set of eigenvalues of the Steklov problem is same as the set of eigenvalues of the well known Dirichlet-Neumann map.

There are several results which estimate the first non-zero eigenvalue μ_1 of the Steklov eigenvalue problem [3, 7, 8, 9, 13]. The first isoperimetric upper bound for μ_1 was given by Weinstock [2] in 1954. He proved that among all simply connected planar domains with analytic boundary of fixed perimeter, the circle maximizes μ_1 . In [3], Payne obtained a two sided bound for the first non-zero Steklov eigenvalue on a convex plain domain in terms of minimum and maximum curvature. The lower bound in [3] has been generalized by Escobar [7] to 2-dimensional compact manifold with non-negative Gaussian curvature. Using the Weinstock inequality, Escobar [8] proved that for a fix volume, among all bounded simply connected domain in 2-dimensional simply connected space forms, geodesic balls maximize the first non-zero Steklov eigenvalue. This result has been extended to non-compact rank-1 symmetric spaces in [13]. We prove the similar result for the first non-zero eigenvalue of the eigenvalue problem

$$\begin{aligned} \Delta_p u &= 0 & \text{in } \Omega, \\ |\nabla u|^{p-2} \frac{\partial u}{\partial \nu} &= \mu_p |u|^{p-2} u & \text{on } M, \end{aligned} \quad (2)$$

where Ω is a bounded domain in \mathbb{R}^n such that $M = \partial\Omega$ and ν is outward unit normal on M .

In Section 2, we state our main results. In section 3, we state some basic facts about the first non-zero eigenvalues of problem (1) and (2), and prove some results which will be required in the later sections. Followed by this, in section 4, 5 and 6, we provide the proof of results stated in section 2.

2 Statement of the results

We state a variation of centre of mass theorem. This is crucial for our proof of main results.

Theorem 1. *Let Ω be a bounded domain in \mathbb{R}^n and $M = \partial\Omega$. Then for every real number $1 < p < \infty$, there exists a point $t \in \overline{\Omega}$ depending on p and normal coordinate system (X_1, X_2, \dots, X_n) centered at t such that for $1 \leq i \leq n$,*

$$\int_M |X_i|^{p-2} X_i = 0.$$

Now we state our main results.

The following theorem provides an upper bound for the first non-zero eigenvalue $\lambda_{1,p}$ of the closed eigenvalue problem (1).

Theorem 2. Let M be a closed hypersurface in \mathbb{R}^n bounding a bounded domain Ω . Let $R > 0$ be such that $\text{Vol}(\Omega) = \text{Vol}(B(R))$, where $B(R)$ is a ball of radius R . Then the first non-zero eigenvalue $\lambda_{1,p}$ of the closed eigenvalue problem (1) satisfies

$$\lambda_{1,p} \leq n^{\frac{p-2}{2}} \lambda_1(S(R))^{\frac{p}{2}} \left(\frac{\text{Vol}(M)}{\text{Vol}(S(R))} \right). \quad (3)$$

Furthermore, for $p = 2$, the upper bound (3) is sharp and the equality holds if and only if M is a geodesic sphere of radius R (see [12]).

If equality holds in (3) then M is a geodesic sphere and $p = 2$.

In case of Steklov eigenvalue problem, we have the following upper bound for the first non-zero eigenvalue $\mu_{1,p}$.

Theorem 3. Let Ω be a bounded domain in \mathbb{R}^n with smooth boundary M and $R > 0$ be such that $\text{Vol}(\Omega) = \text{Vol}(B(R))$, where $B(R)$ is a ball of radius R . Then the first non-zero eigenvalue $\mu_{1,p}$ of problem (2) satisfies the following inequality.

- For $1 < p < 2$,

$$\mu_{1,p} \leq \frac{1}{R^{p-1}}. \quad (4)$$

- For $p \geq 2$,

$$\mu_{1,p} \leq \frac{n^{p-2}}{R^{p-1}}. \quad (5)$$

Furthermore, for $p = 2$, equality holds in (4) and (5) iff M is a geodesic sphere of radius R (see [13]).

If equality holds in (4) and (5) then M is a geodesic sphere of radius R and $p = 2$.

3 Preliminaries

In this section, we state some basic facts about the first non-zero eigenvalue of the eigenvalue problems (1) and (2). We will also prove some results that are needed in subsequent sections.

Let u_1 be an eigenfunction corresponding to the eigenvalue λ_p of closed eigenvalue problem (1) and u_2 be an eigenfunction corresponding to the eigenvalue μ_p of the Steklov eigenvalue problem (2). Then λ_p and μ_p satisfy

$$\lambda_p \int_M |u_1|^p = \int_M \|\nabla^M u_1\|^p,$$

$$\mu_p \int_M |u_1|^p = \int_\Omega \|\nabla u_1\|^p.$$

This shows that all eigenvalues of problems (1) and (2) are non-negative.

Let $\lambda_{1,p}$ and $\mu_{1,p}$ be the first non-zero eigenvalues of the closed and steklov eigenvalue problems, respectively. Then the variational characterization for $\lambda_{1,p}$ and $\mu_{1,p}$ is given by

$$\lambda_{1,p} = \inf \left\{ \frac{\int_M \|\nabla^M u\|^p}{\int_M |u|^p} : \int_M |u|^{p-2} u = 0, u(\neq 0) \in C^1(M) \right\},$$

$$\mu_{1,p} = \inf \left\{ \frac{\int_\Omega \|\nabla u\|^p}{\int_M |u|^p} : \int_M |u|^{p-2} u = 0, u(\neq 0) \in C^1(\Omega) \right\}.$$

Remark 4. If $p = 2$, then the condition $\int_M |u|^{p-2} u = \int_M u = 0$ is equivalent to say that the test function must be orthogonal to the constant function in L^2 -norm.

Let M be a closed hypersurface in \mathbb{R}^n and Ω be a bounded domain in \mathbb{R}^n such that $M = \partial\Omega$. Fix a point $q \in \Omega$. Then for every point $s \in M$, the line joining q and s may intersect M at some points other than s . For every point $s \in M$, let $r(s) := d(q, s)$ and for every $u \in \mathbb{S}^{n-1}$, let $\beta(u) := \max \{ \beta > 0 \mid q + \beta u \in M, \beta \in \mathbb{R} \}$. Let $A := \{ q + \beta(u)u \mid u \in \mathbb{S}^{n-1} \}$. Then $A \subseteq M$.

Lemma 1. Let Ω be a bounded domain in \mathbb{R}^n with smooth boundary M and $R > 0$ be such that $\text{Vol}(\Omega) = \text{Vol}(B(R))$, where $B(R)$ is a ball of radius R . Fix a point $q \in \overline{\Omega}$, then

$$\int_M r^p(s) ds \geq R^p \text{Vol}(S(R)). \quad (6)$$

Further, equality holds in (6) iff M is a geodesic sphere of radius R centered at q .

Proof. For a point $s \in A$, let γ_s be the unique unit speed geodesic joining q and s with $\gamma_s(0) = q$. Let $u = \gamma'_s(0)$ and $t_s(u) = d(q, s)$. Let $\theta(s)$ be the angle between the outward unit normal $\nu(s)$ to M and the radial vector $\partial r(s)$. Let du be the spherical volume density of the unit sphere \mathbb{S}^{n-1} . Then

$$\begin{aligned} \int_M r^p(s) ds &\geq \int_A r^p(s) ds \\ &= \int_{\mathbb{S}^{n-1}} (t_s(u))^p \sec \theta(s) (t_s(u))^{n-1} du \\ &\geq \int_{\mathbb{S}^{n-1}} (t_s(u))^{n+p-1} du \\ &= (n+p-1) \int_{\mathbb{S}^{n-1}} \int_0^{t_s(u)} r^{n+p-2} dr du \\ &\geq (n+p-1) \int_{\Omega} r^{p-1} dV \end{aligned}$$

and

$$\begin{aligned} \int_{\Omega} r^{p-1} dV &= \int_{\Omega \cap B(R)} r^{p-1} dV + \int_{\Omega \setminus \Omega \cap B(R)} r^{p-1} dV \\ &= \int_{B(R)} r^{p-1} dV - \int_{B(R) \setminus \Omega \cap B(R)} r^{p-1} dV + \int_{\Omega \setminus \Omega \cap B(R)} r^{p-1} dV \\ &\geq \int_{B(R)} r^{p-1} dV - \int_{B(R) \setminus \Omega \cap B(R)} r^{p-1} dV + \int_{\Omega \setminus \Omega \cap B(R)} R^{p-1} dV \\ &= \int_{B(R)} r^{p-1} dV + \int_{B(R) \setminus \Omega \cap B(R)} (R^{p-1} - r^{p-1}) dV \\ &\geq \int_{B(R)} r^{p-1} dV \\ &= \int_{\mathbb{S}^{n-1}} \int_0^R r^{n+p-2} dr du \\ &= \int_{\mathbb{S}^{n-1}} \frac{R^{n+p-1}}{n+p-1} du \\ &= \frac{R^p}{n+p-1} \text{Vol}(S(R)). \end{aligned} \quad (7)$$

We have used the fact that $R \leq r$ in $(\Omega \setminus \Omega \cap B(R))$ in (7).

Further, equality holds in (6) iff $\sec \theta(s) = 1$ for all points $s \in M$ and $\text{Vol}(B(R) \setminus \Omega \cap B(R)) = 0$. Note that $\sec \theta(s) = 1$ iff $\theta(s) = 0$ for all points $s \in M$. Therefore outward unit normal

$\nu(s) = \partial r(s)$ for all points $s \in M$. This shows that $\Omega = B(q, R)$ and M is a geodesic sphere of radius R . \square

Above lemma is the generalization of the Lemma 1 in [12].

Lemma 2. *Let $n \in \mathbb{N}$ and y_1, y_2, \dots, y_n be non-negative real numbers. Then for every real number $\gamma \geq 1$, the following inequality holds.*

$$(y_1 + y_2 + \dots + y_n)^\gamma \geq y_1^\gamma + y_2^\gamma + \dots + y_n^\gamma. \quad (8)$$

Proof. Let $n \in \mathbb{N}$ and y_1, y_2, \dots, y_n be non-negative real numbers. Let $\gamma \geq 1$. Then inequality (8) can be written as

$$\left(\frac{y_1}{y_1 + y_2 + \dots + y_n}\right)^\gamma + \left(\frac{y_2}{y_1 + y_2 + \dots + y_n}\right)^\gamma + \dots + \left(\frac{y_n}{y_1 + y_2 + \dots + y_n}\right)^\gamma \leq 1.$$

Therefore, it is enough to show that $a_1^\gamma + a_2^\gamma + \dots + a_n^\gamma \leq 1$ for non-negative real numbers a_i such that $a_1 + a_2 + \dots + a_n = 1$.

Since $0 \leq a_i \leq 1$ and $\gamma \geq 1$, then $a_i^\gamma \leq a_i$. Therefore, $a_1^\gamma + a_2^\gamma + \dots + a_n^\gamma \leq a_1 + a_2 + \dots + a_n = 1$. This proves the Lemma. \square

Next we estimate $\sum_{i=1}^n \|\nabla^M x_i\|^2$.

Lemma 3. *Let M be a closed hypersurface in \mathbb{R}^n and Ω be a bounded domain such that $M = \partial\Omega$. For a fixed point $t \in \bar{\Omega}$, let (x_1, x_2, \dots, x_n) be the normal coordinate system centered at t . Then*

$$\sum_{i=1}^n \|\nabla^M x_i\|^2 = (n-1).$$

Proof. Observe that $\|\nabla x_i(p)\| = 1$ for $1 \leq i \leq n$ and a point $p \in \mathbb{R}^n$. Let ν be the outward unit normal on M . Then

$$\begin{aligned} \sum_{i=1}^n \|\nabla^M x_i\|^2 &= \sum_{i=1}^n \left(\|\nabla x_i\|^2 - \langle \nabla x_i, \nu \rangle^2 \right) \\ &= \sum_{i=1}^n \|\nabla x_i\|^2 - \|\nu\|^2 \\ &= n-1. \end{aligned} \quad \square$$

For a Riemannian geometric proof of above lemma, see [6].

4 Proof of Theorem 1

Proof. Given a point $x \in \mathbb{R}^n$, we write (x_1, \dots, x_n) , the standard Euclidean coordinate system centered at origin. For $1 < p < \infty$, define a function $f : \bar{\Omega} \rightarrow \mathbb{R}$ by

$$f(t_1, \dots, t_n) = \frac{1}{p} \int_M \sum_{i=1}^n |x_i - t_i|^p dx_1 \cdots dx_n.$$

The function f is non-negative on $\bar{\Omega}$. Let α be its infimum. Then there exists a sequence (t_1^j, \dots, t_n^j) in $\bar{\Omega}$ such that

$$\frac{1}{p} \int_M \sum_{i=1}^n |x_i - t_i^j|^p \longrightarrow \alpha \quad \text{in } \mathbb{R} \quad \text{as } j \longrightarrow \infty. \quad (9)$$

Observe that the sequence (t_1^j, \dots, t_n^j) is bounded. Therefore it has a convergent subsequence, without loss of generality, we denote it by (t_1^j, \dots, t_n^j) itself, which converges to $t = (t_1, \dots, t_n) \in \mathbb{R}^n$. Thus $t \in \overline{\Omega}$. Then

$$\sum_{i=1}^n |x_i - t_i^j|^p \longrightarrow \sum_{i=1}^n |x_i - t_i|^p \quad \text{as } j \longrightarrow \infty$$

and

$$\frac{1}{p} \int_M \sum_{i=1}^n |x_i - t_i^j|^p \longrightarrow \frac{1}{p} \int_M \sum_{i=1}^n |x_i - t_i|^p \quad \text{as } j \longrightarrow \infty.$$

Therefore,

$$\frac{1}{p} \int_M \sum_{i=1}^n |x_i - t_i|^p = \alpha \quad \text{and} \quad f(t_1, \dots, t_n) = \alpha.$$

Since f attains its minimum at $t = (t_1, \dots, t_n)$, we have $(\nabla f)_t = 0$. Therefore for each $1 \leq i \leq n$,

$$\langle \nabla f, e_i \rangle_{(t_1, \dots, t_n)} = \int_M |X_i|^{p-2} X_i = 0,$$

where $\{e_i, 1 \leq i \leq n\}$ is the standard orthonormal basis of \mathbb{R}^n and $X_i := (x_i - t_i), 1 \leq i \leq n$. This proves the theorem. \square

We will use the above theorem to show the existence of a point $t = (t_1, \dots, t_n) \in \overline{\Omega}$, such that the coordinate functions with respect to t are test functions for the eigenvalue problems (1) and (2).

5 Proof of Theorem 2

Proof. Let M be a closed hypersurface in \mathbb{R}^n and Ω be the bounded domain such that $M = \partial\Omega$. Let $R > 0$ be such that $\text{Vol}(\Omega) = \text{Vol}(B(R))$. The variational characterization for $\lambda_{1,p}$ is given by

$$\lambda_{1,p} = \inf \left\{ \frac{\int_M \|\nabla^M u\|^p}{\int_M |u|^p} : \int_M |u|^{p-2} u = 0, u(\neq 0) \in C^1(M) \right\}.$$

By Theorem 1, there exists a point $t \in \overline{\Omega}$ such that

$$\int_M |x_i|^{p-2} x_i = 0 \quad \text{for } 1 \leq i \leq n,$$

where (x_1, \dots, x_n) denotes the normal coordinate system centered at t . Therefore, for all $p > 1$,

$$\lambda_{1,p} \int_M \sum_{i=1}^n |x_i|^p \leq \int_M \sum_{i=1}^n \|\nabla^M x_i\|^p \quad \text{for } 1 \leq i \leq n. \quad (10)$$

Now, we divide the proof of the theorem into the following two cases.

Case 1. $1 < p \leq 2$.

Since $|\frac{x_i}{r}| \leq 1$, it follows that

$$|x_i|^p = r^p \left| \frac{x_i}{r} \right|^p \geq r^p \left| \frac{x_i}{r} \right|^2 \quad \text{for } 1 \leq i \leq n. \quad (11)$$

Therefore,

$$r^p = r^p \sum_{i=1}^n \left| \frac{x_i}{r} \right|^2 \leq r^p \sum_{i=1}^n \left| \frac{x_i}{r} \right|^p = \sum_{i=1}^n |x_i|^p.$$

For $1 < p < 2$, using Hölder's inequality, we obtain

$$\sum_{i=1}^n \|\nabla^M x_i\|^p \leq \left(\sum_{i=1}^n \|\nabla^M x_i\|^2 \right)^{\frac{p}{2}} n^{\frac{2-p}{2}}.$$

This combining with Lemma 3 gives

$$\sum_{i=1}^n \|\nabla^M x_i\|^p \leq (n-1)^{\frac{p}{2}} n^{\frac{2-p}{2}}.$$

Observe that the above inequality is also true for $p = 2$. By substituting above values in inequality (10), we get

$$\lambda_{1,p} \int_M r^p \leq (n-1)^{\frac{p}{2}} n^{\frac{2-p}{2}} \text{Vol}(M). \quad (12)$$

By substituting $\int_M r^p \geq R^p \text{Vol}(S(R))$ in above equation, we get

$$\lambda_{1,p} R^p \text{Vol}(S(R)) \leq (n-1)^{\frac{p}{2}} n^{\frac{2-p}{2}} \text{Vol}(M).$$

As a consequence, we have

$$\lambda_{1,p} \leq n^{\frac{2-p}{2}} \lambda_1(S(R))^{\frac{p}{2}} \left(\frac{\text{Vol}(M)}{\text{Vol}(S(R))} \right).$$

This proves Theorem 2 for $1 < p \leq 2$.

Equality in (3) implies equality in Lemma 1 and equality in (11), which implies that M is a geodesic sphere of radius R and $p = 2$.

Case 2. $p \geq 2$.

For $p > 2$, by Hölder's inequality, we have

$$\sum_{i=1}^n |x_i|^2 \leq \left(\sum_{i=1}^n (|x_i|^2)^{\frac{p}{2}} \right)^{\frac{2}{p}} n^{\frac{p-2}{p}}.$$

Therefore,

$$n^{\frac{2-p}{2}} r^p \leq \sum_{i=1}^n |x_i|^p. \quad (13)$$

Observe that equality holds in the above inequality for $p = 2$, so (13) holds for $p \geq 2$. Now we estimate $\sum_{i=1}^n \|\nabla^M x_i\|^p$. Since $\frac{p}{2} \geq 1$ and $\|\nabla^M x_i\|^2 \geq 0$, for each $1 \leq i \leq n$, it follows from Lemma 2 that

$$\begin{aligned} \sum_{i=1}^n \|\nabla^M x_i\|^p &= \sum_{i=1}^n (\|\nabla^M x_i\|^2)^{\frac{p}{2}} \\ &\leq \left(\sum_{i=1}^n \|\nabla^M x_i\|^2 \right)^{\frac{p}{2}} \\ &= (n-1)^{\frac{p}{2}}. \end{aligned} \quad (14)$$

The last inequality follows from Lemma 3. By substituting values from (13) and (14) in (10), we get

$$\lambda_{1,p} n^{\frac{2-p}{2}} \int_M r^p \leq (n-1)^{\frac{p}{2}} \text{Vol}(M). \quad (15)$$

By substituting $\int_M r^p \geq R^p \text{Vol}(S(R))$ from Lemma 1 in above inequality, we have

$$\lambda_{1,p} n^{\frac{2-p}{2}} R^p \text{Vol}(S(R)) \leq (n-1)^{\frac{p}{2}} \text{Vol}(M).$$

Therefore,

$$\lambda_{1,p} \leq n^{\frac{p-2}{2}} \lambda_1(S(R))^{\frac{p}{2}} \left(\frac{\text{Vol}(M)}{\text{Vol}(S(R))} \right).$$

This proves Theorem 2 for $p \geq 2$.

If equality holds in (3), then equality holds in (6) and also in (13). Equality in (6) implies that M is a geodesic sphere of radius R and equality in (13) holds iff $p = 2$. Otherwise, $p > 2$ and equality in (13) implies that $|x_i| = c$, for some constant c and $1 \leq i \leq n$. Therefore, each point of M is of the form $(\pm c, \pm c, \pm c, \dots, \pm c)$, for some constant c . This contradicts our assumption that M is the boundary of a bounded domain Ω . \square

6 Proof of Theorem 3

Proof. Let Ω be a bounded domain in \mathbb{R}^n with smooth boundary $\partial\Omega = M$ and $R > 0$ be such that $\text{Vol}(\Omega) = \text{Vol}(B(R))$, where $B(R)$ is a ball of radius R . The variational characterization for $\mu_{1,p}$ is given by

$$\mu_{1,p} = \inf \left\{ \frac{\int_{\Omega} \|\nabla u\|^p}{\int_M |u|^p} : \int_M |u|^{p-2} u = 0, u(\neq 0) \in C^1(\Omega) \right\}.$$

By Theorem 1, there exists a point $t \in \bar{\Omega}$ such that

$$\int_M |x_i|^{p-2} x_i = 0, \quad \text{for all } 1 \leq i \leq n,$$

where $(x_1, x_2, x_3, \dots, x_n)$ denotes the normal coordinate system centered at t . By considering each x_i as test function, we have

$$\mu_{1,p} \int_M \sum_{i=1}^n |x_i|^p \leq \int_{\Omega} \sum_{i=1}^n \|\nabla x_i\|^p. \quad (16)$$

Now we consider the following two cases to prove the theorem.

Case 1. $1 < p \leq 2$.

By similar argument as in (11), we get

$$r^p \leq \sum_{i=1}^n |x_i|^p.$$

By Hölder's inequality,

$$\sum_{i=1}^n \|\nabla x_i\|^p \leq \left(\sum_{i=1}^n \|\nabla x_i\|^2 \right)^{\frac{p}{2}} n^{\frac{2-p}{2}} = n.$$

By substituting above values in (16), we get

$$\mu_{1,p} \int_M r^p \leq n \text{Vol}(\Omega).$$

By substituting $\int_M r^p \geq R^P \text{Vol}(S(R))$ from Lemma 1, we have

$$\mu_{1,p} R^P \text{Vol}(S(R)) \leq n \text{Vol}(\Omega).$$

Since $\text{Vol}(\Omega) = \text{Vol}(B(R))$ and $\frac{\text{Vol}(B(R))}{\text{Vol}(S(R))} = \frac{R}{n}$, we get

$$\mu_{1,p} \leq \frac{1}{R^{p-1}}.$$

Case 2. $p \geq 2$.

From (13), we have

$$n^{\frac{2-p}{2}} r^p \leq \sum_{i=1}^n |x_i|^p \quad \text{for all } p \geq 2.$$

By Lemma 2, we have

$$\begin{aligned} \sum_{i=1}^n \|\nabla x_i\|^p &\leq \sum_{i=1}^n (\|\nabla x_i\|^2)^{\frac{p}{2}} \\ &\leq \left(\sum_{i=1}^n \|\nabla x_i\|^2 \right)^{\frac{p}{2}} \\ &\leq n^{\frac{p}{2}}. \end{aligned}$$

By substituting above values in (16), we get

$$\mu_{1,p} n^{\frac{2-p}{2}} \int_M r^p \leq n^{\frac{p}{2}} \text{Vol}(\Omega).$$

We use Lemma 1 again to get

$$\mu_{1,p} n^{\frac{2-p}{2}} R^P \text{Vol}(S(R)) \leq n^{\frac{p}{2}} \text{Vol}(\Omega).$$

Since $\text{Vol}(\Omega) = \text{Vol}(B(R))$ and $\frac{\text{Vol}(B(R))}{\text{Vol}(S(R))} = \frac{R}{n}$, above equation becomes

$$\mu_{1,p} \leq \frac{n^{p-2}}{R^{p-1}}.$$

Equality case will follow same as in Theorem 2. This completes the proof. \square

Acknowledgment

I am very grateful to Prof. G. Santhanam for his support and constructive suggestions which led to improvements in the article.

References

- [1] M. W. Stekloff, les problemes fondamentaux de la physique mathematique, *Ann. Sci. Ecole Norm* **19** 445-490 (1902).
- [2] R. Weinstock, Inequalities for a classical Eigenvalue problem, *Rational Mech. Anal* **3** 745-753 (1954).
- [3] L. E. Payne, Some isoperimetric inequalities for Harmonic functions, *SIAM J. Math. Anal.* **1** 354-359 (1970).

- [4] D. Bleecker, J. Weiner, Extrinsic bounds on λ_1 of Δ on a compact manifold, *Comment. Math. Helv.* **51** 601-609 (1976).
- [5] R. Reilly, On the first eigenvalue of the Laplacian for compact submanifold of Euclidean space, *Comment. Math. Helv.* **52** 525-533 (1977).
- [6] E. Heintze, Extrinsic upperbounds for λ_1 , *Math. Ann.* **280** 389-402 (1988).
- [7] J. F. Escobar, The geometry of the first non-zero Stekloff eigenvalue, *Journal of Functional Analysis* **150**(2) 544-556 (1997).
- [8] J. F. Escobar, An Isoperimetric Inequality and the First Steklov Eigenvalue, *Journal of Functional Analysis* **165**(1) 101-116 (1999).
- [9] J. F. Escobar, A Comparison Theorem for the First Non-zero Steklov Eigenvalue, *Journal of Functional Analysis* **178**(1) 143-155 (2000).
- [10] J. F. Grosjean, Upper bounds for the first eigenvalue of the Laplacian on compact submanifolds, *Pacific. J. Math.* **206** 93-112 (2002).
- [11] Santhanam G, A sharp upper bound for the first eigenvalue of the Laplacian of compact hypersurfaces in rank-1 symmetric spaces, *Proc. Indian Acad. Sci. (Math. Sci.)* **117**(3) 307-315 (2007).
- [12] G. Santhanam, Isoperimetric upper bounds for the first eigenvalue, *Proc. Indian Acad. Sci. (Math. Sci.)* **122**(3) 375-384 (2012).
- [13] Binoy, G. Santhanam, Sharp upperbound and a comparison theorem for the first nonzero Steklov eigenvalue, *J. Ramanujan Math. Soc.* **29**(2) 133-154 (2014).