

# High dimensional robust sparse regression

Liu Liu  
liuliu@utexas.edu

Yanyao Shen  
shenyanyao@utexas.edu

Tianyang Li  
lty@cs.utexas.edu

Constantine Caramanis  
constantine@utexas.edu

The University of Texas at Austin

## Abstract

We provide a novel – and to the best of our knowledge, the first – algorithm for high dimensional sparse regression with corruptions in explanatory and/or response variables. Our algorithm recovers the true sparse parameters in the presence of a constant fraction of arbitrary corruptions. Our main contribution is a robust variant of Iterative Hard Thresholding. Using this, we provide accurate estimators with sub-linear sample complexity. Our algorithm consists of a novel randomized outlier removal technique for robust sparse mean estimation that may be of interest in its own right: it is orderwise more efficient computationally than existing algorithms, and succeeds with high probability, thus making it suitable for general use in iterative algorithms. We demonstrate the effectiveness on large-scale sparse regression problems with arbitrary corruptions.

## 1 Introduction

Learning in the presence of arbitrarily (even adversarially) corrupted outliers in the training data has a long history in Robust Statistics [19, 16, 27], and has recently received much renewed attention. The high dimensional setting poses particular challenges as outlier removal via preprocessing is essentially impossible when the number of variables scales with the number of samples. We propose a computationally efficient estimator for outlier-robust sparse regression that has near-optimal sample complexity, and is the first algorithm resilient to a constant fraction of arbitrary outliers with corrupted covariates and/or response variables. Unless we specifically mention otherwise, all future mentions of outliers mean corruptions in covariates and/or response variables.

Specifically, the problem we solve is as follows. We are given  $n$  samples  $\mathbf{z}_i = (y_i, \mathbf{x}_i)$ , with  $\mathbf{x}_i \in \mathbb{R}^d$ . Following Huber’s  $\epsilon$ -contamination model [18], these samples are drawn from the mixture

$$\mathbf{z}_i \sim (1 - \epsilon)P + \epsilon Q, \quad (1)$$

where  $P$  represents the uncorrupted distribution, with  $y_i = \mathbf{x}_i^\top \boldsymbol{\beta}^* + \xi_i$ , and  $Q$  is an arbitrary distribution, representing the outliers (see Section 1.3 for complete details and definitions). Outlier-robust regression is a classical problem within robust statistics (e.g., [25]), yet even in the low-dimensional setting, efficient algorithms robust to corruption in the covariates have proved elusive, until recent breakthroughs in [24, 11] and [20], which built on important results in Robust Mean Estimation [9, 21] and Sums of Squares [2], respectively.

In the sparse setting, the parameter  $\boldsymbol{\beta}^*$  we seek to recover is also  $k$ -sparse, and a key goal is to provide recovery guarantees with sample complexity scaling with  $k$ , and sublinearly with  $d$ . Without outliers, by now classical results (e.g., [13])  $n = O(k \log d)$  samples are enough to give

recovery guarantees on  $\beta^*$  with and without additive noise. In the additive noise setting, exact recovery of  $\beta^*$  with a constant  $\epsilon$ -fraction of arbitrary outliers does not seem to be possible. Instead, we seek to give an efficient, sample-complexity optimal algorithm that recovers  $\beta^*$  to within accuracy depending on  $\epsilon$ , the fraction of outliers. In the case of no additive noise, we seek algorithms that can guarantee exact recovery, independent of  $\epsilon$ .

## 1.1 Related work

The last 10 years have seen a resurgence in interest in robust statistics, including the problem of resilience to outliers in the data. Important problems attacked have included PCA [30, 29, 21, 9], and more recently robust regression (as in this paper) [24, 11, 20] and robust mean estimation [21, 1], among others. We focus now on the recent work most related to the present paper.

**Robust regression.** Earlier work in robust regression considers corruption only in the output, and shows that algorithms nearly as efficient as for regression without outliers succeeds in parameter recovery, even with a constant fraction of outliers [22, 23, 4, 5]. Yet these algorithms (and their analysis) focus on corruption in  $y$ , and do not seem to extend to the setting of corrupted covariates – the setting of this work.

In the low dimensional setting, there has been remarkable recent progress. The work in [20] shows that the Sum of Squares (SOS) based semidefinite hierarchy can be used for solving robust regression. In fact, their results apply to a corruption model that allows the adversary to select an  $\epsilon$ -fraction of points to corrupt, and hence is stronger than Huber’s  $\epsilon$ -contamination model (where the corrupted fraction is chosen at random). Essentially concurrent to the SOS work, [24, 11] develop a framework for robust gradient descent for empirical risk minimization, by using robust mean estimation as a subroutine to compute robust gradients at each iteration. Though this latter work applies to Huber’s  $\epsilon$ -contamination model, on the other hand, computationally it scales better than the algorithms in [20], as although the Sum of Squares SDP framework gives polynomial time algorithms, they are often not practical [17].

Much less appears to be known in the high-dimensional regime. Work in [8] first analyzed high dimensional sparse regression with arbitrary corruptions in covariates. They show that replacing the standard inner product in Matching Pursuit with a trimmed version, one can recover from an  $\epsilon$ -fraction of outliers, with  $\epsilon = O(1/\sqrt{d})$ . The more recent work in low-dimension regression discussed above does not shed light on the sparse (high dimensional) setting. In particular, it is not clear how to extend the techniques in [24] and [11] to the sparse case, while maintaining correctly scaling sample complexity.

Another approach follows as a byproduct of a recent algorithm for robust sparse mean estimation, in [1]. However, their error guarantee scales with  $\|\beta^*\|_2$ , and moreover, does not provide *exact recovery* in the adversarial corruption case without stochastic noise (i.e., noise variance  $\sigma^2 = 0$ ). We note that this is an inevitable consequence of their approach, as they directly use sparse mean estimation on  $\{y_i x_i\}$ , rather than considering Maximum Likelihood Estimation.

**Robust mean estimation.** A key idea in [24, 11] is to leverage recent breakthroughs in robust mean estimation. Very recently, [9, 21] provided the first robust mean estimation algorithms that can handle a constant fraction of outliers. The key idea there relies on an ellipsoid algorithm. Following their work, [1] extended the ellipsoid algorithm from [9] to also study the high dimensional extension to robust sparse mean estimation. They show that  $d$ -dimensional  $k$ -sparse mean estimation with a constant fraction of outliers can be done with  $n = \Omega(k^2 \log(d))$  samples. The  $k^2$  term appears to be necessary, as  $n = \Omega(k^2)$  follows from an oracle-based lower bound [12].

## 1.2 Main contributions

The main results of our work, and our innovations in the context of the above body of work, are as follows.

- We provide an algorithm that is resilient to a small but constant fraction,  $\epsilon$ , of arbitrary outliers (corruptions in the covariates, and/or labels). Our algorithm requires  $n = \Omega(k^2 \log d)$  samples. The extra factor of  $k$  in the  $k^2$  term is not information-theoretically optimal, and is due to our use of sparse PCA as a subroutine. The  $k^2$  term is not likely improvable for sparse PCA (see, e.g., [3]), and we conjecture it is not improvable for this problem either.<sup>1</sup> Our result is based on Iterative Hard Thresholding (IHT) [6]; specifically, we provide a stability result showing that IHT works with any robust sparse mean estimation algorithm. This is the content of Theorem 2.1.
- Our result can be viewed as a meta-theorem, as it allows the use of any sparse mean estimation subroutine that can provide guarantees with high probability. Then, these guarantees provided carry over to the full IHT-based algorithm. Corollary 2.1 uses a robust sparse mean estimation subroutine based on a version of the ellipsoid algorithm, given and analyzed in [1]. The proof of its performance in [1] hinges on obtaining an upper bound on the sparse operator norm (their Lemmas A.2 and A.3). As we show via counterexample (see Appendix A), the statement of Lemma A.3 seems to be incorrect, and the general approach of upper bounding the sparse operator norm may not work. Nevertheless, the algorithm performance they claim is correct, as we show through a quite different avenue (see Lemma D.3 in Appendix D.3).

Using this ellipsoid algorithm, our results show that with an  $\epsilon$ -fraction of arbitrary corruptions, we recover  $\beta^*$  within additive error  $O(\sigma\epsilon)$ . (We note that this is a stronger guarantee than the low-dimensional robust regression result in [24], which has a  $O(\sigma\sqrt{\epsilon})$  guarantee.) In particular, we obtain exact recovery if either the fraction of outliers goes to zero (this is just ordinary sparse regression), or in the presence of a constant fraction of outliers but with the additive noise term going to zero (this is the case of robust sparse linear equations). To the best of our knowledge, this is the first result that shows exact recovery for robust sparse linear equations with a constant fraction of outliers.

- We then prove a result that may be of interest in its own right: we provide a novel robust sparse mean estimation algorithm that is based on a filtering algorithm for sequentially screening and removing potential outliers, rather than on the ellipsoid algorithm. The filtering algorithm is faster by at least a quadratic factor in the dimension ( $d^2$ ), and also provides exponential probability guarantees, unlike, e.g., the filtering-style algorithm of [10]. This sharper concentration requires a more careful Martingale-bound analysis. This is the content of Theorem 3.1. Combining this with Theorem 2.1 gives us an algorithm for robust sparse regression that is at least a factor of  $d^2$  faster than the ellipsoid-based algorithm. This speedup comes at a cost, as our final result guarantees recovering  $\beta^*$  within an additive error of  $O(\sigma\sqrt{\epsilon})$ , similarly to [24]. Nevertheless, the result is strong enough to guarantee exact recovery when either  $\sigma$  or  $\epsilon$  goes to zero.

## 1.3 Setup, Notation and Outline

We consider Huber’s  $\epsilon$ -contamination model described above: samples  $\{\mathbf{z}_1, \dots, \mathbf{z}_n\}$  are drawn from  $(1 - \epsilon)P + \epsilon Q$ .  $P$  represents the authentic samples, and  $Q$  the adversarially corrupted samples.  $P$  follows the standard Gaussian-design AWGN linear model:  $\mathbf{z}_i = (\mathbf{x}_i, y_i)$ , with

$$y_i = \mathbf{x}_i^\top \beta^* + \xi_i, \tag{2}$$

---

<sup>1</sup>We note that results in [12] show a statistical query-based lower bound of  $\Omega(k^2)$  on sparse mean estimation.

where  $\beta^* \in \mathbb{R}^d$  is the model parameter, and assumed to be  $k$ -sparse. We assume that  $\mathbf{x}_i \sim \mathcal{N}(0, I_d)$  and  $\xi_i \sim \mathcal{N}(0, \sigma^2)$ . The parameter  $\epsilon$  represents the fraction of outliers. Throughout, we assume that it is a (small) constant, *independent of dimension or other problem parameters*. As in [1, 9], we pre-process via a naive pruning step, removing ‘‘obvious’’ outliers; we henceforth assume that all authentic and corrupted points are within a radius bounded by a polynomial in  $n$ ,  $d$  and  $1/\epsilon$ .

Given samples from the  $\epsilon$ -contamination model eq. (1), we define two subsets of  $[n]$ :  $\mathcal{G}, \mathcal{B} \subseteq \{1, \dots, n\}$ , where  $i \in \mathcal{G}$  if the corresponding sample is drawn from  $P$ , and  $i \in \mathcal{B}$  otherwise. Of course, the membership of each point is unknown.

**Notation.** We denote the Hard Thresholding operator of sparsity  $k$  by  $P_k$ . We define the  $k$ -sparse operator norm as  $\|M\|_{k, \text{op}} = \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq k} |\mathbf{v}^\top M \mathbf{v}|$ , where  $M$  is not required to be positive-semidefinite (p.s.d.). We define the infinity norm for a matrix  $M$  as  $\|M\|_\infty = \max_{i,j} |M_{ij}|$ . Given index set  $\mathcal{S}$ ,  $\mathbf{v}^{\mathcal{S}}$  is the vector restricted to indices  $\mathcal{S}$ . Similarly,  $M^{\mathcal{S}\mathcal{S}}$  is the sub-matrix on indices  $\mathcal{S} \times \mathcal{S}$ . We let  $\otimes$  denote the Kronecker product, and for a vector  $\mathbf{u}$ , we denote the outer product by  $\mathbf{u}^{\otimes 2} = \mathbf{u}\mathbf{u}^\top$ . We use  $\mathbb{E}_{\mathbf{g} \in \mathcal{S}}$  to denote the expectation operator obtained by the uniform distribution over all samples  $\mathbf{g}$  in a set  $\mathcal{S}$ . In proofs, we use  $C_1, C_2$  to denote constants that are independent of dimension, but whose value can change from line to line. Finally, we use the notation  $\tilde{O}(\cdot)$  to hide the dependency on  $\text{poly} \log(1/\epsilon)$  in our bounds.

**Outline.** The rest of this paper is organized as follows. In Section 2, we introduce the main algorithm Algorithm 1, and provide theoretical guarantees. In Section 3, we introduce a new and computationally more efficient algorithm (Algorithm 2) for robust sparse mean estimation. We provide the experimental results in Section 4. All proofs are collected in the Appendix.

## 2 Robust sparse regression with corrupted gradients

The overall flow of the algorithm is straightforward: we use IHT with robust sparse gradient updates, and show that the overall algorithm inherits results from robust sparse mean estimation.

### 2.1 Methodology

We define notation for the stochastic gradient corresponding to the  $i^{\text{th}}$  point,  $\mathbf{x}_i$ , and the gradient corresponding to the collection of authentic samples:

$$\mathbf{g}_i^t = \mathbf{x}_i (\mathbf{x}_i^\top \beta^t - y_i), \quad \mathbf{G}^t = \mathbb{E}_P (\mathbf{g}_i^t),$$

where recall that  $P$  is the distribution of the authentic points, as defined in eq. (1). Now, in the *uncorrupted case* where all samples follow the statistical model eq. (2), a single iteration of IHT updates  $\beta^t$  via

$$\beta^{t+1} = P_k \left( \beta^t - \frac{1}{n} \sum_{i=1}^n \mathbf{g}_i^t \right). \quad (3)$$

Since  $\mathbb{E}_P (\mathbf{x}_i \mathbf{x}_i^\top) = I_d$ , the mean of all authentic gradients  $\mathbf{G}^t$  can be calculated as

$$\mathbf{G}^t = \mathbb{E}_P (\mathbf{g}_i^t) = \mathbb{E}_P (\mathbf{x}_i \mathbf{x}_i^\top (\beta^t - \beta^*)) = \beta^t - \beta^*.$$

Therefore,  $\mathbf{G}^t = \beta^t - \beta^*$  is guaranteed to be  $2k$ -sparse.

Thus in the  $\epsilon$ -contamination model eq. (1), we use a Robust Sparse Gradient Estimator (RSGE) to estimate the unknown true  $\mathbf{G}^t$  from  $\{\mathbf{g}_i^t\}_{i=1}^n$ .

For a fixed  $k$ -sparse parameter  $\beta$ , we drop the superscript  $t$ , and use  $\mathbf{g}_i$  in place of  $\mathbf{g}_i^t$ , and  $\mathbf{G}$  in place of  $\mathbf{G}^t$ ; hence,  $\mathbf{G} = \mathbf{G}(\beta)$  denotes the gradient over the authentic samples, at the point  $\beta$ .

---

**Algorithm 1** Robust sparse regression with corrupted gradients

---

- 1: **Input:** Data samples  $\{y_i, \mathbf{x}_i\}_{i=1}^N$ , RSGE subroutine.
  - 2: **Output:** The estimation  $\hat{\beta}$
  - 3: Split samples into  $T$  subsets of size  $n$ .
  - 4: Initialize with  $\beta^0 = \mathbf{0}$ .
  - 5: **for**  $t = 0$  to  $T - 1$ , **do**
  - 6:     At current  $\beta^t$ , calculate all gradients for current  $n$  samples:  $\mathbf{g}_i^t = \mathbf{x}_i (\mathbf{x}_i^\top \beta^t - y_i)$ ,  $i \in [n]$ .
  - 7:     The initial input set is  $\{\mathbf{g}_i^t\}_{i=1}^n$ . We use a RSGE to get  $\hat{\mathbf{G}}^t$ .
  - 8:     Update the parameter:  $\beta^{t+1} = P_k(\beta^t - \hat{\mathbf{G}}^t)$ .
  - 9: Output the estimation  $\hat{\beta} = \beta^T$ .
- 

**Definition 2.1** (RSGE). Given  $n(k, d, \epsilon, \nu)$   $\epsilon$ -corrupted samples  $\{\mathbf{g}_i\}_{i=1}^n$  from eq. (1), we call  $\hat{\mathbf{G}}(\beta)$  a  $\psi(\epsilon)$ -RSGE, if given  $\{\mathbf{g}_i\}_{i=1}^n$ ,  $\hat{\mathbf{G}}(\beta)$  guarantees

$$\left\| \hat{\mathbf{G}}(\beta) - \mathbf{G}(\beta) \right\|_2^2 \leq \alpha \|\mathbf{G}(\beta)\|_2^2 + \psi(\epsilon),$$

with probability at least  $1 - \nu$ , where  $\alpha \in (0, 0.1)$  is a constant.

Algorithm 1 takes any RSGE as a subroutine, and runs the modified version of IHT with the estimated sparse gradient at each iteration. Our results require sample splitting to maintain independence between subsequent iterations, though we believe this is an artifact of our analysis.

## 2.2 Main results for robust sparse regression

RSGE introduces a controlled amount of error in every single IHT update step. Theorem 2.1 gives a convergence guarantee of Algorithm 1 by showing that IHT does not accumulate too much error.

**Theorem 2.1** (Main theorem). *Suppose we observe  $N(k, d, \epsilon, \nu)$   $\epsilon$ -corrupted samples from eq. (1). Algorithm 1, with  $\psi(\epsilon)$ -RSGE defined in Definition 2.1, outputs  $\hat{\beta}$ , such that*

$$\left\| \hat{\beta} - \beta^* \right\|_2 = O\left(\sqrt{\psi(\epsilon)}\right), \quad (4)$$

with probability at least  $1 - \nu$ , by setting  $T = \Theta\left(\log\left(\frac{\|\beta^*\|_2}{\sqrt{\psi(\epsilon)}}\right)\right)$ , and  $N(k, d, \epsilon, \nu) = n(k, d, \epsilon, \nu/T)T$ .

We give the proof of Theorem 2.1 in Appendix C.

In [1], the authors propose a robust sparse regression estimator based on robust sparse mean estimation on  $\{y_i \mathbf{x}_i\}$ , leveraging the fact that  $\mathbb{E}_P(y_i \mathbf{x}_i) = \beta^*$ . With sample complexity  $N = \Omega\left(\frac{k^2 \log(d/\nu)}{\epsilon^2}\right)$ , this algorithm produces a  $\tilde{\beta}$  such that, with probability at least  $1 - \nu$ ,

$$\left\| \tilde{\beta} - \beta^* \right\|_2^2 = \tilde{O}\left(\epsilon^2 \left(\|\beta^*\|_2^2 + \sigma^2\right)\right). \quad (5)$$

Using Theorem 2.1, we show that we can obtain significantly stronger statistical guarantees; in particular, guarantees that yield exact recovery when  $\sigma = 0$ .

Indeed, the ellipsoid-algorithm-based robust sparse mean estimator from [1] gives a RSGE, which we can combine with Theorem 2.1 to obtain stronger results. We state these as Corollary 2.1. We note again that the analysis in [1] has a flaw. Their Lemma A.3 is incorrect, as our counterexample in Appendix A demonstrates. We provide a correct route of analysis in Lemma D.3 of Appendix D.

**Corollary 2.1.** *Under the same setting as in Theorem 2.1, if we use the ellipsoid algorithm for robust sparse gradient estimation, it requires sample complexity  $N(k, d, \epsilon, \nu) = \Omega\left(\frac{k^2 \log(dT/\nu)}{\epsilon^2}\right)T$ , and guarantees  $\psi(\epsilon) = \tilde{O}(\epsilon^2 \sigma^2)$ . Hence, Algorithm 1 outputs  $\hat{\beta}$ , such that*

$$\left\| \hat{\beta} - \beta^* \right\|_2 = \tilde{O}(\sigma \epsilon), \quad (6)$$

with probability at least  $1 - \nu$ , by setting  $T = \Theta\left(\log\left(\frac{\|\beta^*\|_2}{\epsilon \sigma}\right)\right)$ .

*Proof sketch.* The ellipsoid-based algorithm deals with outliers by trying to optimize the set of weights on each of the samples – ideally outliers would receive lower weight and hence their impact would be minimized. Since the set of weights is convex, this can be approached using a separation oracle. The separation oracle in this case depends on a convex relaxation of Sparse PCA. With each call to the relaxation of sparse PCA, we obtain an optimal value,  $\lambda^*$ , and optimal solution,  $H^*$ , to the problem:

$$\lambda^* = \max_H \text{Tr}\left(\left(\hat{\Sigma} - F\left(\hat{\mathbf{G}}\right)\right) \cdot H\right), \quad \text{subject to } H \succcurlyeq 0, \|H\|_{1,1} \leq 2k, \text{Tr}(H) = 1. \quad (7)$$

Here,  $\hat{\mathbf{G}}$ ,  $\hat{\Sigma}$  are weighted first and second order moment estimates from  $\epsilon$ -corrupted samples, and  $F: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$  is a function with closed-form

$$F(\mathbf{G}) \equiv \mathbb{E}_P\left(\left(\mathbf{g}_i - \mathbf{G}\right)\left(\mathbf{g}_i - \mathbf{G}\right)^\top\right), \quad (8)$$

which calculates the underlying true covariance matrix. We provide more details about the calculation of  $F(\cdot)$ , as well as some smoothness properties, in Appendix B.

The key to the proof relies on showing that  $\lambda^*$  controls the quality of the weights of the current iteration, i.e., small  $\lambda^*$  means good weights and thus a good current solution. Showing this relies on using  $\lambda^*$  to control  $\left(\hat{\Sigma} - F\left(\hat{\mathbf{G}}\right)\right)$ . Lemma A.3 in [1] claims that  $\lambda^* \geq \left\|\left(\hat{\Sigma} - F\left(\hat{\mathbf{G}}\right)\right)\right\|_{2k, \text{op}}$ . As we show in Appendix A, however, this need not hold. This is because the trace norm maximization eq. (7) is not a valid convex relaxation for the  $2k$ -sparse operator norm when the term  $\hat{\Sigma} - F\left(\hat{\mathbf{G}}\right)$  is not p.s.d. (which indeed it need not be). We provide a different line of analysis in Lemma D.3, essentially showing that even without the claimed (incorrect) bound,  $\lambda^*$  can still provide the control we need. With the corrected analysis for  $\lambda^*$ , the ellipsoid algorithm guarantees  $\left\|\hat{\mathbf{G}} - \mathbf{G}\right\|_2^2 = \tilde{O}\left(\epsilon^2\left(\|\mathbf{G}\|_2^2 + \sigma^2\right)\right)$  with probability at least  $1 - \nu$ . Therefore, the algorithm provides an  $\tilde{O}(\epsilon^2 \sigma^2)$ -RSGE.  $\square$

From a computational viewpoint, the time complexity of Algorithm 1 depends on the RSGE in each iterate. The time complexity of the ellipsoid algorithm is indeed polynomial in the dimension, but it requires  $O(d^2)$  calls to a relaxation of Sparse PCA ([7]). In the next section, we introduce a faster algorithm as a RSGE, which only requires  $O(n)$  calls of Sparse PCA (recall that  $n = O(k^2 \log d)$ ).

### 3 Computationally efficient robust sparse mean estimation

In this section, we propose a different robust sparse mean estimator which only requires  $O(n)$  Sparse PCA calls. Similarly to the ellipsoid algorithm, the core ingredient is (a relaxation of) Sparse PCA. We use calls to Sparse PCA to discard samples, and hence the number of calls we need is linear in the sample size; in contrast, the ellipsoid algorithm in [9, 1] updates each sample’s weight via an

---

**Algorithm 2** Computationally efficient Robust Sparse Gradient Estimator
 

---

- 1: **Input:** A set  $S_{\text{in}}$ , parameter  $\rho_{\text{sep}}$ .
- 2: **Output:** A set  $S_{\text{out}}$  or sparse mean vector  $\widehat{\mathbf{G}}$ .
- 3: Compute the sample mean  $\widetilde{\mathbf{G}} = \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}}(\mathbf{g})$ , and  $\widehat{\mathbf{G}} = P_{4k}(\widetilde{\mathbf{G}})$ .
- 4: Compute the sample covariance matrix  $\widehat{\Sigma} = \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}}(\mathbf{g} - \widehat{\mathbf{G}})(\mathbf{g} - \widehat{\mathbf{G}})^\top$ .
- 5: Solve the following convex program:

$$\max_H \text{Tr} \left( (\widehat{\Sigma} - F(\widehat{\mathbf{G}})) \cdot H \right), \quad \text{subject to } H \succcurlyeq 0, \|H\|_{1,1} \leq 2k, \text{Tr}(H) = 1. \quad (9)$$

Let  $\lambda^*$  be the optimal value, and  $H^*$  be the corresponding solution.

- 6: **if**  $\lambda^* \leq \rho_{\text{sep}}$  **then return** with  $\widehat{\mathbf{G}}$ .
- 7: Calculate projection score for each  $\mathbf{g}_i \in S_{\text{good}}$ :

$$\tau_i = \text{Tr} \left( H^* \cdot (\mathbf{g}_i - \widehat{\mathbf{G}})(\mathbf{g}_i - \widehat{\mathbf{G}})^\top \right). \quad (10)$$

- 8: Randomly remove a sample  $r$  from  $S_{\text{in}}$  according to

$$\Pr(\mathbf{g}_i \text{ is removed}) = \frac{\tau_i}{\sum_{i \in S_{\text{in}}} \tau_i}. \quad (11)$$

- 9: **return** the set  $S_{\text{out}} = S_{\text{in}} \setminus \{r\}$ .
- 

optimization approach that uses Sparse PCA as a separation oracle, and hence it requires at least  $O(d^2)$  calls to the separation oracle [7] (in this case Sparse PCA).

Our proposed RSGE (Algorithm 2) attempts to remove one outlier at each iteration, as long as a good solution has not already been identified. It first estimates the gradient  $\widehat{\mathbf{G}}$  by hard thresholding (line 3) and then estimates the corresponding sample covariance matrix  $\widehat{\Sigma}$  (line 4). By solving (a relaxation of) Sparse PCA, we obtain a scalar  $\lambda^*$  as well as a matrix  $H^*$ . If  $\lambda^*$  is smaller than the predetermined threshold  $\rho_{\text{sep}}$ , we have a certificate that the effect of the outliers is well-controlled. Otherwise, we compute a score for each sample based on  $H^*$ , and discard one of the samples according to a probability distribution where each sample's probability of being discarded is proportional to the score we have computed.

Algorithm 2 can be easily transplanted to other robust sparse functional estimation problems, whenever we have a covariance function  $F(\mathbf{G})$ . For example, taking  $F(\cdot)$  to be the fixed matrix  $I_d$ , we have an algorithm for robust sparse mean estimation for  $\mathcal{N}(\mu, I_d)$ , where  $\mu$  is  $k$ -sparse.

### 3.1 Guarantees for Algorithm 2

To use Algorithm 2 as a RSGE given  $n$  gradient samples as  $S_{\text{in}}$ , we call Algorithm 2 repeatedly on  $S_{\text{in}}$  and then on its output,  $S_{\text{out}}$ , until it returns a robust estimator  $\widehat{\mathbf{G}}$ . The next theorem provides guarantees on this iterative application of Algorithm 2.

**Theorem 3.1.** *Suppose we observe  $n = \Omega\left(\frac{k^2 \log(d/\nu)}{\epsilon}\right)$   $\epsilon$ -corrupted samples from eq. (1). Let  $S_{\text{in}}$  be an  $\epsilon$ -corrupted set of gradient samples  $\{\mathbf{g}_i^t\}_{i=1}^n$ . If we run Algorithm 2 iteratively with initial set  $S_{\text{in}}$ , and subsequently on  $S_{\text{out}}$ , and use  $\rho_{\text{sep}} = C_\gamma(U + \sigma^2)$ , then this repeated use of Algorithm 2 will stop after at most  $\frac{1.1\gamma}{\gamma-1}\epsilon n$  iterations, and output  $\widehat{\mathbf{G}}^t$ , such that*

$$\left\| \widehat{\mathbf{G}}^t - \mathbf{G}^t \right\|_2^2 = \widetilde{O}\left(\epsilon \left( \left\| \mathbf{G}^t \right\|_2^2 + \sigma^2 \right)\right),$$

with probability at least  $1 - \nu - \exp(-\Theta(\epsilon n))$ . Here,  $C_\gamma$  is a constant depending on  $\gamma$ , where  $\gamma \geq 2$  is a constant, and  $U$  is any constant factor bound on  $\|\mathbf{G}^t\|_2^2$ .

Thus, Theorem 3.1 shows that with high probability, Algorithm 2 provides a Robust Sparse Gradient Estimator where  $\psi(\epsilon) = \tilde{O}(\epsilon\sigma^2)$ . For example, we can take  $\nu = d^{-\Theta(1)}$ . We note that this significantly improves the probability bound in [11]. Combining now with Theorem 2.1, we obtain a strong guarantee for robust sparse regression.

**Corollary 3.1.** *Under the same setting as Theorem 2.1, using Algorithm 2 for robust sparse gradient estimation with  $N(k, d, \epsilon, \nu/T) = \Omega\left(\frac{k^2 \log(dT/\nu)}{\epsilon}\right)$  samples, and  $T = \Theta\left(\log\left(\frac{\|\beta^*\|_2}{\sqrt{\epsilon}\sigma}\right)\right)$ , then*

$$\left\|\hat{\beta} - \beta^*\right\|_2 = \tilde{O}(\sqrt{\epsilon}\sigma) \quad (12)$$

with probability at least  $1 - \nu - T \exp(-\Theta(\epsilon n))$ .

**Remark** (Recovery error dependencies on  $\epsilon$ ). The scaling of  $\epsilon$  in eq. (12) is  $\tilde{O}(\sqrt{\epsilon})$ , which matches  $\tilde{O}(\sqrt{\epsilon})$  in [11] and [24] for low dimensional linear regression. These guarantees are worse than  $\tilde{O}(\epsilon)$  achieved by ellipsoid methods; this seems to be the price to pay for the faster algorithm.

The proof of Theorem 3.1 is given in Appendix F. The key step in Algorithm 2 is outlier removal eq. (11) based on the solution of Sparse PCA's convex relaxation eq. (9).

### 3.2 Outlier removal in Algorithm 2

In Algorithm 2, we denote samples in the input set  $S_{\text{in}}$  as  $\mathbf{g}_i$ . This input set  $S_{\text{in}}$  can be partitioned into two parts:  $S_{\text{good}} = \{i : i \in \mathcal{G} \text{ and } i \in S_{\text{in}}\}$ , and  $S_{\text{bad}} = \{i : i \in \mathcal{B} \text{ and } i \in S_{\text{in}}\}$ . Lemma 3.1 shows that Algorithm 2 can detect when the contribution of  $\sum_{i \in S_{\text{good}}} \tau_i$  is small, and hence the outlier removal step is likely to discard an outlier.

**Lemma 3.1.** *Suppose we observe  $n = \Omega\left(\frac{k^2 \log(d/\nu)}{\epsilon}\right)$   $\epsilon$ -corrupted samples from eq. (1). Let  $S_{\text{in}}$  be an  $\epsilon$ -corrupted set of gradient samples  $\{\mathbf{g}_i^t\}_{i=1}^n$ . Algorithm 2 computes  $\lambda^*$  that satisfies*

$$\lambda^* \geq \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\hat{\mathbf{G}}) \right) \mathbf{v}. \quad (13)$$

Furthermore, if  $\lambda^* \geq \rho_{\text{sep}} = C_\gamma \left( \|\mathbf{G}^t\|_2^2 + \sigma^2 \right)$ , then with probability at least  $1 - \nu$ , the contribution of  $\sum_{i \in S_{\text{good}}} \tau_i$  is small:

$$\sum_{i \in S_{\text{good}}} \tau_i \leq \frac{1}{\gamma} \sum_{i \in S_{\text{in}}} \tau_i, \quad (14)$$

where  $\tau_i$  is defined in eq. (10),  $C_\gamma$  is a constant depending on  $\gamma$ , and  $\gamma \geq 2$  is a constant.

The proofs are collected in Appendix E. In a nutshell, eq. (13) is a natural convex relaxation for the sparsity constraint  $\{\mathbf{v} : \|\mathbf{v}\|_2 = 1, \|\mathbf{v}\|_0 \leq 2k\}$ . The upper bound eq. (14) is obtained through concentration inequalities for the samples in  $S_{\text{good}}$  and the condition that  $\lambda^*$  is large.

Based on Lemma 3.1, if  $\lambda^* \leq \rho_{\text{sep}}$ , eq. (13) can guarantee that  $\left\|\hat{\mathbf{G}}^t - \mathbf{G}^t\right\|_2^2$  is small. We can show that eq. (11) in line 8 is more likely to throw out samples of  $S_{\text{bad}}$  rather than  $S_{\text{good}}$ . Iteratively applying Algorithm 2 on the remaining samples, we can remove those outliers with large effect, and keep the remaining outliers' effect well-controlled. This leads to the final bounds in Theorem 3.1.

<sup>2</sup>Similarly to [9, 10, 1, 11], our results seem to require this side information.






Figure 1: Simulations for Algorithm 2 showing the dependence of relative MSE on different quantities. For each parameter, we choose corresponding sample complexity  $n \propto k^2 \log(d)/\epsilon$ . Different curves for  $\epsilon \in \{0.1, 0.15, 0.2\}$  are the average of 15 trials. Consistent with the theory, the rescaled relative MSE's are nearly independent of sparsity and dimension. Furthermore, by rescaling for different  $\epsilon$ , three curves have the same magnitude.

## 4 Numerical Results

### 4.1 Robust sparse mean estimation

We first demonstrate the performance of Algorithm 2 for robust sparse mean estimation, and then move to Algorithm 1 for robust sparse regression. For the robust sparse gradient estimation, we generate samples through  $\mathbf{g}_i = \mathbf{x}_i \mathbf{x}_i^\top \mathbf{G} - \mathbf{x}_i \xi_i$ , where the unknown true mean  $\mathbf{G}$  is  $k$ -sparse. The authentic  $\mathbf{x}_i$ 's are generated from  $\mathcal{N}(0, I_d)$ . We set  $\sigma = 0$ , since the main part of the error in robust sparse mean estimation is  $\mathbf{G}$ . Each entry of  $\mathbf{G}$  is either  $+1$  or  $-1$ , hence  $\|\mathbf{G}\|_2^2 = k$ .

The outliers are specially designed: the norm of the outliers is  $\|\mathbf{G}\|_2$ , and the directions are orthogonal to  $\mathbf{G}$ . Through this construction, outliers cannot be easily removed by simple pruning, and the directions of outliers can cause large effects on the estimation of  $\mathbf{G}$ . We plot the relative MSE of parameter recovery, defined as  $\|\hat{\mathbf{G}} - \mathbf{G}\|_2^2 / \|\mathbf{G}\|_2^2$ , with respect to different sparsities and dimensions.

**Parameter error vs. sparsity  $k$ .** We fix the dimension to be  $d = 50$ . We solve the trace norm maximization in Algorithm 2 using CVX [15]. We solve robust sparse gradient estimation under different levels of outlier fraction  $\epsilon$  and different sparsity values  $k$ .

**Parameter error vs. dimension  $d$ .** We fix  $k = 5$ . We use a Sparse PCA solver from [14] which is much more efficient for higher dimensions. We run robust sparse gradient estimation Algorithm 2 under different levels of outlier fraction  $\epsilon$  and different dimensions  $d$ .

For each parameter, the corresponding number of samples required for the authentic data is  $n \propto k^2 \log(d)/\epsilon$  according to Theorem 3.1. Therefore, we add  $\epsilon n / (1 - \epsilon)$  outliers (so that the outliers are an  $\epsilon$ -fraction of the total samples), and then run Algorithm 2. According to Theorem 3.1, the rescaled relative MSE:  $\|\hat{\mathbf{G}} - \mathbf{G}\|_2^2 / (\epsilon \|\mathbf{G}\|_2^2)$  should be independent of the parameters  $\{\epsilon, k, d\}$ . We plot this in fig. 1, and these plots validate our theorem on the sample complexity in robust sparse mean estimation problems.

### 4.2 Robust sparse regression

We use Algorithm 1 for robust sparse regression. Similarly as in Section 4.1, we use Algorithm 2 as our Robust Sparse Gradient Estimator, and leverage the Sparse PCA solver from [14]. In the simulation, we fix  $d = 500$ , and  $k = 5$ , hence the corresponding sample complexity is  $n \propto 1/\epsilon$ . However, we do not use the sample splitting technique in the simulations.

The entries of the true parameter  $\boldsymbol{\beta}^*$  are set to be either  $+1$  or  $-1$ , hence  $\|\boldsymbol{\beta}^*\|_2^2 = k$  is fixed. The authentic  $\mathbf{x}_i$ 's are generated from  $\mathcal{N}(0, I_d)$ , and the authentic  $y_i = \mathbf{x}_i^\top \boldsymbol{\beta}^* + \xi_i$  as in eq. (2). We




Figure 2: Empirical illustration of the linear convergence of the Algorithm 1. In all cases, we fix  $k = 5$ ,  $d = 500$ , and choose the sample complexity  $n \propto 1/\epsilon$ . (2a) considers different  $\epsilon$  with fixed  $\sigma^2 = 0.1$ . (2b) considers different  $\sigma^2$  with fixed  $\epsilon = 0.1$ . As expected, the convergence is linear, and flatten out at the level of the final error.

set the covariates of the outliers as  $A$ , where  $A$  is a random  $\pm 1$  matrix of dimension  $\epsilon n / (1 - \epsilon) \times d$ , and set the responses of outliers to  $-A\beta^*$ .

To show the performance of Algorithm 1 under different settings, we use different levels of  $\epsilon$  and  $\sigma$  in fig. 2, and track the parameter error  $\|\beta^t - \beta^*\|_2^2$  of Algorithm 1 in each iteration. Consistent with the theory, the algorithm displays linear convergence, and the error curves flatten out at the level of the final error. Furthermore, Algorithm 1 achieves machine precision when  $\sigma^2 = 0$  in fig. 2b.

## 5 Acknowledgments

The authors would like to thank Simon S. Du for helpful discussions.

## References

- [1] Sivaraman Balakrishnan, Simon S. Du, Jerry Li, and Aarti Singh. Computationally efficient robust sparse estimation in high dimensions. In *Proceedings of the 2017 Conference on Learning Theory*, 2017.
- [2] Boaz Barak and David Steurer. Proofs, beliefs, and algorithms through the lens of sum-of-squares.
- [3] Quentin Berthet and Philippe Rigollet. Complexity theoretic lower bounds for sparse principal component detection. In *Conference on Learning Theory*, pages 1046–1066, 2013.
- [4] Kush Bhatia, Prateek Jain, and Purushottam Kar. Robust regression via hard thresholding. In *Advances in Neural Information Processing Systems*, pages 721–729, 2015.
- [5] Kush Bhatia, Prateek Jain, and Purushottam Kar. Consistent robust regression. In *Advances in Neural Information Processing Systems*, pages 2107–2116, 2017.
- [6] Thomas Blumensath and Mike E Davies. Iterative hard thresholding for compressed sensing. *Applied and computational harmonic analysis*, 27(3):265–274, 2009.
- [7] Sébastien Bubeck. Convex optimization: Algorithms and complexity. *Foundations and Trends® in Machine Learning*, 8(3-4):231–357, 2015.
- [8] Yudong Chen, Constantine Caramanis, and Shie Mannor. Robust sparse regression under adversarial corruption. In *International Conference on Machine Learning*, pages 774–782, 2013.
- [9] Ilias Diakonikolas, Gautam Kamath, Daniel M Kane, Jerry Li, Ankur Moitra, and Alistair Stewart. Robust estimators in high dimensions without the computational intractability. In *Foundations of Computer Science (FOCS), 2016 IEEE 57th Annual Symposium on*, pages 655–664. IEEE, 2016.
- [10] Ilias Diakonikolas, Gautam Kamath, Daniel M. Kane, Jerry Li, Ankur Moitra, and Alistair Stewart. Being robust (in high dimensions) can be practical. In *International Conference on Machine Learning*, pages 999–1008, 2017.
- [11] Ilias Diakonikolas, Gautam Kamath, Daniel M Kane, Jerry Li, Jacob Steinhardt, and Alistair Stewart. Sever: A robust meta-algorithm for stochastic optimization. *arXiv preprint arXiv:1803.02815*, 2018.
- [12] Ilias Diakonikolas, Daniel M Kane, and Alistair Stewart. Statistical query lower bounds for robust estimation of high-dimensional gaussians and gaussian mixtures. In *Foundations of Computer Science (FOCS), 2017 IEEE 58th Annual Symposium on*, pages 73–84. IEEE, 2017.
- [13] David L Donoho. Compressed sensing. *IEEE Transactions on information theory*, 52(4):1289–1306, 2006.
- [14] Alexandre d’Aspremont, Francis Bach, and Laurent El Ghaoui. Optimal solutions for sparse principal component analysis. *Journal of Machine Learning Research*, 9(Jul):1269–1294, 2008.
- [15] Michael Grant, Stephen Boyd, and Yinyu Ye. Cvx: Matlab software for disciplined convex programming, 2008.
- [16] Frank R Hampel, Elvezio M Ronchetti, Peter J Rousseeuw, and Werner A Stahel. *Robust statistics: the approach based on influence functions*, volume 196. John Wiley & Sons, 2011.
- [17] Samuel B Hopkins, Tselil Schramm, Jonathan Shi, and David Steurer. Fast spectral algorithms from sum-of-squares proofs: tensor decomposition and planted sparse vectors. In *Proceedings of the forty-eighth annual ACM symposium on Theory of Computing*, pages 178–191. ACM, 2016.
- [18] Peter J Huber. Robust estimation of a location parameter. *The annals of mathematical statistics*, pages 73–101, 1964.
- [19] Peter J Huber. Robust statistics. In *International Encyclopedia of Statistical Science*, pages 1248–1251. Springer, 2011.
- [20] Adam Klivans, Pravesh K. Kothari, and Raghu Meka. Efficient Algorithms for Outlier-Robust Regression. *arXiv preprint arXiv:1803.03241*, 2018.
- [21] Kevin A Lai, Anup B Rao, and Santosh Vempala. Agnostic estimation of mean and covariance. In *Foundations of Computer Science (FOCS), 2016 IEEE 57th Annual Symposium on*, pages 665–674. IEEE, 2016.

- [22] Xiaodong Li. Compressed sensing and matrix completion with constant proportion of corruptions. *Constructive Approximation*, 37(1):73–99, 2013.
- [23] Nam H Nguyen and Trac D Tran. Exact recoverability from dense corrupted observations via  $\ell_1$ -minimization. *IEEE transactions on information theory*, 59(4):2017–2035, 2013.
- [24] Adarsh Prasad, Arun Sai Suggala, Sivaraman Balakrishnan, and Pradeep Ravikumar. Robust estimation via robust gradient estimation. *arXiv preprint arXiv:1802.06485*, 2018.
- [25] Peter J Rousseeuw and Annick M Leroy. *Robust regression and outlier detection*, volume 589. John wiley & sons, 2005.
- [26] Charles M Stein. Estimation of the mean of a multivariate normal distribution. *The annals of Statistics*, pages 1135–1151, 1981.
- [27] John W Tukey. Mathematics and the picturing of data. In *Proceedings of the International Congress of Mathematicians, Vancouver, 1975*, volume 2, pages 523–531, 1975.
- [28] Vincent Q Vu, Juhee Cho, Jing Lei, and Karl Rohe. Fantope projection and selection: A near-optimal convex relaxation of sparse PCA. In *Advances in neural information processing systems*, pages 2670–2678, 2013.
- [29] Huan Xu, Constantine Caramanis, and Shie Mannor. Outlier-robust PCA: the high-dimensional case. *IEEE transactions on information theory*, 59(1):546–572, 2013.
- [30] Huan Xu, Constantine Caramanis, and Sujay Sanghavi. Robust PCA via outlier pursuit. *IEEE Transactions on Information Theory*, 58(5):3047–3064, 2012.

## A Correcting Lemma A.3 in [1]’s proof

A key part of the proof of the main theorem in [1] is to obtain an upper bound on the  $k$ -sparse operator norm. Specifically, Lemmas A.2 and A.3 in [1] aim to show:

$$\lambda^* \geq \left\| \sum_{i=1}^{|S_{\text{in}}|} w_i \left( \mathbf{g}_i - \widehat{\mathbf{G}}(w) \right)^{\otimes 2} - F \left( \widehat{\mathbf{G}}(w) \right) \right\|_{2k, \text{op}} \geq \frac{\|P_{2k}(\widetilde{\Delta}(w))\|_2^2}{5\epsilon}, \quad (15)$$

where  $\widehat{\mathbf{G}}(w) = P_{4k} \left( \sum_{i=1}^{|S_{\text{in}}|} w_i \mathbf{g}_i \right)$ ,  $\widetilde{\Delta}(w) = \sum_{i=1}^{|S_{\text{in}}|} w_i \mathbf{g}_i - \mathbf{G}^3$ , and recall  $\lambda^*$  is the solution to the SDP as given in Algorithm 2.

Lemma A.3 asserts the first inequality above, and Lemma A.2 the second. As we show below, Lemma A.3 cannot be correct. Specifically, the issue is that the quantity inside the second term in eq. (15) may not be positive semidefinite. In this case, the convex optimization problem whose solution is  $\lambda^*$  is not a valid relaxation, and hence the  $\lambda^*$  they obtain need not a valid upper bound. Indeed, we give a simple example below that illustrates precisely this potential issue.

Fortunately, not all is lost – indeed, as our results imply, the main results in [1] are correct. The key is to show that while  $\lambda^*$  does not upper bound the sparse operator norm, it does, however, upper bound the quantity

$$\max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \sum_{i=1}^{|S_{\text{in}}|} w_i \left( \mathbf{g}_i - \widehat{\mathbf{G}}(w) \right)^{\otimes 2} - F \left( \widehat{\mathbf{G}}(w) \right) \right) \mathbf{v}. \quad (16)$$

We show this in Appendix D. More specifically, in Lemma D.3, we replace the  $2k$ -sparse operator norm in the second term of eq. (15) by the term in eq. (16). We show this can be used to complete the proof in Appendix D.4.

We now provide a counterexample that shows the first inequality in (15) cannot hold. The main argument is that the convex relaxation for sparse PCA is a valid upper bound of the sparse operator norm only for positive semidefinite matrices. Specifically, denoting  $E = \widehat{\Sigma}(w) - F \left( \widehat{\mathbf{G}}(w) \right)$  as the matrix in eq. (16), [1] solves the following convex program:

$$\max_H \text{Tr}(E \cdot H), \quad \text{subject to } H \succeq 0, \|H\|_{1,1} \leq k, \text{Tr}(H) = 1.$$

Since  $\widehat{\Sigma}(w) - F \left( \widehat{\mathbf{G}}(w) \right)$  is no longer a p.s.d. matrix, the trace maximization above may not be a valid convex relaxation, and thus not an upper bound. Let us consider a specific example, in robust sparse mean estimation for  $\mathcal{N}(\mu, I_d)$ , where function  $F(\cdot)$  is a fixed identity matrix  $I_d$ . We choose  $k = 1$ ,  $\mu = [1, 0]^\top$ , and  $d = 2$ . Suppose we observe data to be  $x_1 = [2.5, 0]^\top$ ,  $x_2 = [0, 0]^\top$ , and the weights for  $x_1$  and  $x_2$  are the same. Then, we can compute the following matrices as:

$$\widehat{\Sigma} = \begin{bmatrix} 1.5625 & 0 \\ 0 & 0 \end{bmatrix}, F = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, E = \widehat{\Sigma} - F = \begin{bmatrix} 0.5625 & 0 \\ 0 & -1 \end{bmatrix}.$$

It is clear that  $\left\| \widehat{\Sigma} - F \right\|_{k, \text{op}} = 1$ . Solving the convex relaxation  $\max_H \text{Tr}(E \cdot H)$  or  $\max_H \text{Tr}(\widehat{\Sigma} \cdot H)$  gives answer  $H^* = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  and the corresponding  $\lambda^* = 0.5625$ , which is clearly not an upper bound of  $\left\| \widehat{\Sigma} - F \right\|_{k, \text{op}}$ . Hence  $\lambda^* \geq \left\| \widehat{\Sigma} - F \right\|_{k, \text{op}}$  cannot hold in general.

<sup>3</sup>The  $\{w_i\}$  are weights, and these are defined precisely in Section D, but are not required for the present discussion or counterexample.

## B Covariance structure in robust sparse mean estimation

Both the ellipsoid algorithm and Algorithm 2 require a closed form expression of the true covariance function  $F(\mathbf{G})$ , in order to handle different probabilistic distributions of uncorrupted samples. Indeed, the robust sparse mean estimation algorithm in [1] uses the covariance structure given by  $F$  to detect outliers. The accuracy of robust sparse mean estimation explicitly depends on the properties of  $F(\mathbf{G})$ . Two important properties of  $F(\mathbf{G})$ , relate to its smoothness, as controlled by two parameters  $L_{\text{cov}}$  and  $L_{\text{F}}$ . We first provide a closed-form expression for  $F$ , and then define precisely the smoothness parameters  $L_{\text{cov}}$  and  $L_{\text{F}}$ , and show how these can be controlled.

**The closed form expression of  $F(\mathbf{G})$ .** Recall from eq. (8) that  $F(\mathbf{G})$  is defined as,

$$\begin{aligned} F(\mathbf{G}) &= \mathbb{E} \left( (\mathbf{g}_i - \mathbf{G})(\mathbf{g}_i - \mathbf{G})^\top \right) \\ &= \mathbb{E} \left( (\mathbf{x}\mathbf{x}^\top - I) \mathbf{G}\mathbf{G}^\top (\mathbf{x}\mathbf{x}^\top - I) \right) + \sigma^2 I \\ &= \mathbb{E} \left( \mathbf{x}\mathbf{x}^\top \mathbf{G}\mathbf{G}^\top \mathbf{x}\mathbf{x}^\top \right) - 2 \mathbb{E} \left( \mathbf{x}\mathbf{x}^\top \mathbf{G}\mathbf{G}^\top \right) + \mathbf{G}\mathbf{G}^\top + \sigma^2 I. \end{aligned}$$

Next, we apply the Stein-type Lemma [26] for  $\mathbf{x} \sim \mathcal{N}(0, I_d)$ , and a function  $f(x)$  whose second derivative exists:

$$\mathbb{E} (f(x) \mathbf{x}\mathbf{x}^\top) = \mathbb{E} (f(x)) I + \mathbb{E} (\nabla^2 f(x)). \quad (17)$$

By eq. (17), we have

$$F(\mathbf{G}) = \|\mathbf{G}\|_2^2 I + \mathbf{G}\mathbf{G}^\top + \sigma^2 I.$$

**The smoothness properties of  $\|F\|_{\text{op}}$ .**  $L_{\text{cov}}$  is defined as

$$L_{\text{cov}} = \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} |\mathbf{v}^\top F(\mathbf{G}) \mathbf{v}|.$$

For  $\forall \mathbf{G}, \widehat{\mathbf{G}}$ , there exist a  $L_{\text{F}}$  and a universal constant  $C$  such that,

$$\left\| F(\mathbf{G}) - F(\widehat{\mathbf{G}}) \right\|_{\text{op}} \leq L_{\text{F}} \left\| \mathbf{G} - \widehat{\mathbf{G}} \right\|_2 + C \left\| \mathbf{G} - \widehat{\mathbf{G}} \right\|_2^2.$$

$L_{\text{cov}}$  is upper bounded by the top eigenvalue of  $F(\mathbf{G})$ ,

$$L_{\text{cov}} \leq \|F(\mathbf{G})\|_2 \leq 2 \|\mathbf{G}\|_2^2 + \sigma^2. \quad (18)$$

For the  $L_{\text{F}}$  term,

$$\left\| F(\mathbf{G}) - F(\widehat{\mathbf{G}}) \right\|_{\text{op}} \quad (19)$$

$$= \left\| 2\mathbf{G}^\top (\mathbf{G} - \widehat{\mathbf{G}}) I - \left\| \mathbf{G} - \widehat{\mathbf{G}} \right\|_2^2 I + \mathbf{G} (\mathbf{G} - \widehat{\mathbf{G}})^\top + (\mathbf{G} - \widehat{\mathbf{G}}) \mathbf{G}^\top - (\mathbf{G} - \widehat{\mathbf{G}}) (\mathbf{G} - \widehat{\mathbf{G}})^\top \right\|_{\text{op}} \quad (20)$$

$$\leq 4 \|\mathbf{G}\|_2 \left\| \mathbf{G} - \widehat{\mathbf{G}} \right\|_2 + 2 \left\| \mathbf{G} - \widehat{\mathbf{G}} \right\|_2^2. \quad (21)$$

Therefore, we can choose  $L_{\text{F}} = 4 \|\mathbf{G}\|_2$  and  $C = 2$ .

## C Robust sparse regression with RSGE

**Theorem C.1** (Theorem 2.1). *Suppose we observe  $N(k, d, \epsilon, \nu)$   $\epsilon$ -corrupted samples from eq. (1). Algorithm 1, with  $\psi(\epsilon)$ -RSGE defined in Definition 2.1, outputs  $\hat{\beta}$ , such that*

$$\left\| \hat{\beta} - \beta^* \right\|_2 = O\left(\sqrt{\psi(\epsilon)}\right), \quad (22)$$

with probability at least  $1 - \nu$ , by setting  $T = \Theta\left(\log\left(\frac{\|\beta^*\|_2}{\sqrt{\psi(\epsilon)}}\right)\right)$ , and  $N(k, d, \epsilon, \nu) = n(k, d, \epsilon, \nu/T)T$ .

*Proof.* By splitting  $N$  samples into  $T$  sets (each set has sample size  $n$ ), Algorithm 1 collects a fresh batch of samples with size  $n(k, d, \epsilon, \nu/T)$  at each iteration  $t \in [T]$ . Definition 2.1 shows that for the fixed gradient expectation  $\mathbf{G}^t$ , the estimate for the gradient  $\hat{\mathbf{G}}^t$  satisfies:

$$\left\| \hat{\mathbf{G}}^t - \mathbf{G}^t \right\|_2^2 \leq \alpha \|\beta^t - \beta^*\|_2^2 + \psi \quad (23)$$

with probability at least  $1 - \nu/T$ , where  $\alpha < 0.1$  is a constant. Here, we use the abbreviation  $\psi$  for the  $\psi(\epsilon)$  in Definition 2.1.

Next, let  $z^t = \beta^t - \hat{\mathbf{G}}^t$ . By the triangle inequality and the definition of Hard Thresholding operator, we have

$$\|\beta^{t+1} - \beta^*\|_2 \leq \|\beta^{t+1} - z^t\|_2 + \|\beta^* - z^t\|_2 \leq 2\|\beta^* - z^t\|_2. \quad (24)$$

Next, we study the  $t$ -th iteration of Algorithm 1. Notice that since by definition,  $\mathbf{G}^t = \beta^t - \beta^*$ , by combining eq. (23) and eq. (24), we have

$$\begin{aligned} \|\beta^{t+1} - \beta^*\|_2^2 &\leq 4\|\beta^* - z^t\|_2^2 \\ &= 4\left\| \beta^* - \beta^t + \hat{\mathbf{G}}^t \right\|_2^2 \\ &\leq 4\alpha \|\beta^t - \beta^*\|_2^2 + 4\psi. \end{aligned}$$

By solving this recursion and using a union bound, we have

$$\|\beta^t - \beta^*\|_2^2 \leq (4\alpha)^t \|\beta^0 - \beta^*\|_2^2 + \frac{4\psi}{1 - 4\alpha} = (4\alpha)^t \|\beta^*\|_2^2 + \frac{4\psi}{1 - 4\alpha},$$

with probability at least  $1 - \nu$ . Hence, by the assumption  $\alpha < 0.1$ , we can set  $T = \Theta\left(\log\left(\frac{\|\beta^*\|_2}{\sqrt{\psi(\epsilon)}}\right)\right)$ .

Finally, we have  $\left\| \hat{\beta} - \beta^* \right\|_2 = O\left(\sqrt{\psi(\epsilon)}\right)$  with probability at least  $1 - \nu$ .  $\square$

## D Proofs for the ellipsoid algorithm in robust sparse regression

In this section, we prove guarantees for the ellipsoid algorithm in robust sparse regression. We first give preliminary definitions of error terms defined on  $S_{\text{good}}$  and  $S_{\text{in}}$ , and then prove Lemma D.1. Next, we prove concentration results for gradients of uncorrupted sparse linear regression in Lemma D.2. In Lemma D.3, we provide lower bounds for the  $2k$ -sparse largest eigenvalue defined in eq. (16). Finally, we prove Corollary 2.1 based on previous Lemmas in Appendix D.4.

## D.1 Preliminary definitions and properties related to $S_{\text{good}}, S_{\text{bad}}$

Here, we state again the definitions of  $S_{\text{good}}, S_{\text{bad}}$  and  $S_{\text{in}}$  defined in Section 3.2. In Algorithm 2, we denote the input set as  $S_{\text{in}}$ , which can be partitioned into two parts:  $S_{\text{good}} = \{i : i \in \mathcal{G} \text{ and } i \in S_{\text{in}}\}$ , and  $S_{\text{bad}} = \{i : i \in \mathcal{B} \text{ and } i \in S_{\text{in}}\}$ . Note that  $S_{\text{in}} = S_{\text{good}} \cup S_{\text{bad}}$ , and  $n = |S_{\text{in}}|$ . For the convenience of our analysis, we define the following error terms:

$$\begin{aligned}\tilde{\Delta}_{S_{\text{good}}} &= \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}}(\mathbf{g}) - \mathbf{G}, \\ \widehat{\Delta}_{S_{\text{good}}} &= P_{4k}(\mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}}(\mathbf{g})) - \mathbf{G}, \\ \tilde{\Delta} &= \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}}(\mathbf{g}) - \mathbf{G}, \\ \widehat{\Delta} &= P_{4k}(\mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}}(\mathbf{g})) - \mathbf{G}.\end{aligned}$$

These error terms are defined under a uniform distribution over samples, whereas previous papers using ellipsoid algorithms [9, 1] consider a set of balanced weighted distribution. More specifically, the weights in our setting are defined as:

$$\tilde{w}_i = \frac{1}{n}, \quad \forall i \in S_{\text{good}} \cup S_{\text{bad}}.$$

The balanced weighted distribution is defined to satisfy:

$$0 \leq w_i \leq \frac{1}{(1-2\epsilon)n}, \quad \forall i \in S_{\text{good}} \cup S_{\text{bad}}, \quad \sum_{i \in S_{\text{in}}} w_i = 1.$$

Notice that  $\sum_{i \in S_{\text{bad}}} \tilde{w}_i = O(\epsilon)$ , and  $\sum_{i \in S_{\text{bad}}} w_i = O\left(\frac{\epsilon}{1-2\epsilon}\right)$  with high probability, which intuitively says that both types of distributions have  $O(\epsilon)$  weights over all bad samples. We are interested in considering uniform weighted samples since this formulation helps us analyze the filtering algorithm more conveniently, as we show in the following sections.

Work in [1] shows the connection of these different error terms, which is restated in the following Lemma:

**Lemma D.1** (Lemma A.1 in [1]). *Suppose  $G$  is  $2k$ -sparse. Then we have the following result:*

$$\frac{1}{5} \|\widehat{\Delta}\|_2 \leq \|P_{2k}(\tilde{\Delta})\|_2 \leq 4 \|\widehat{\Delta}\|_2.$$

## D.2 Concentration bounds for gradients in $S_{\text{good}}$

We first prove concentration bounds for gradients for sparse linear regression in the uncorrupted case. The following is essentially Lemma D.1 in [1], and the proof is an immediate corollary of their result.

**Lemma D.2.** *Suppose we observe gradient samples  $\{\mathbf{g}\}_{i=1}^n$  from eq. (2) with  $n = \Omega\left(\frac{k \log(d/\nu)}{\epsilon^2}\right)$ . Denote this set as  $\mathcal{G}$ . Then, there is a  $\delta = \tilde{O}(\epsilon)$ , such that with probability at least  $1 - \nu$ , we have for any subset  $\mathcal{S} \subset [d]$ ,  $|\mathcal{S}| \leq 2k$ ,*

$$\|\mathbb{E}_{\mathbf{g} \in_u \mathcal{G}}(\mathbf{g}^{\mathcal{S}}) - \mathbf{G}^{\mathcal{S}}\|_2 \leq \delta (\|\mathbf{G}\|_2 + \sigma), \quad (25)$$

$$\left\| \mathbb{E}_{\mathbf{g} \in_u \mathcal{G}}(\mathbf{g}^{\mathcal{S}} - \mathbf{G}^{\mathcal{S}})^{\otimes 2} - F(\mathbf{G})^{\mathcal{S}\mathcal{S}} \right\|_{\text{op}} \leq \delta (\|\mathbf{G}\|_2^2 + \sigma^2). \quad (26)$$

*Proof of Lemma D.2.* The main difference from Lemma D.1 of [1] is that we consider a uniform distribution over all samples instead of a balanced weighted distribution. Furthermore, eqs. (25)



and (26) are the concentration inequalities for the mean and covariance of the collected gradient samples  $\{\mathbf{g}_i\}$  in the good set with the form:

$$\mathbf{g}_i = \mathbf{x}_i \mathbf{x}_i^\top \mathbf{G} - \mathbf{x}_i \xi_i,$$

which is equivalent to Lemma D.1 in [1], where they consider  $y_i \mathbf{x}_i = \mathbf{x}_i \mathbf{x}_i^\top \boldsymbol{\beta} + \mathbf{x}_i \xi_i$ . Therefore, by setting all weights to  $\frac{1}{n}$  in Lemma D.1 of [1] we obtain the desired concentration properties.  $\square$

### D.3 Relationship between the first and second moment of samples in $S_{\text{in}}$

Similar to [9, 1], in this part, we show an important connection between the covariance deviation (the empirical covariance of  $S_{\text{in}}$  minus the true covariance of authentic data) and the mean deviation (the empirical mean of  $S_{\text{in}}$  minus the true mean of authentic data). When the mean deviation (in  $\ell_2$  sense) is large, the following Lemma implies that the covariance deviation must also be large. As a result, when the magnitude of the covariance deviation is large, the current set of samples (or the current weights of all samples) needs to be adjusted; when the magnitude of the covariance deviation is small, the average of current sample set (or the weighted sum of samples using current weights) provides a good enough estimate of the model parameter. Moreover, the same principle holds when we use an approximation of the true covariance, which can be efficiently estimated.

Unlike Lemma A.2 in [1], in eq. (29), eq. (30), we provide lower bounds for the  $2k$ -sparse largest eigenvalue, instead of the  $2k$ -sparse operator norm. As we discussed in Appendix A,  $\lambda^*$  is the convex relaxation of finding the  $2k$ -sparse largest eigenvalue (instead of the  $2k$ -sparse operator norm). In the statement of the following Lemma, for the purpose of consistency, we consider the uniform distribution of weights. However, the proof and results can be easily extended to the setting with the balanced distribution of weights. This is due to the similarity between the two types of weight representation, as discussed in Appendix D.1.

**Lemma D.3.** *Suppose  $|S_{\text{bad}}| \leq 2\epsilon|S_{\text{in}}|$ , and the gradient samples in  $S_{\text{good}}$  satisfy*

$$\left\| P_{2k} \left( \tilde{\Delta}_{S_{\text{good}}} \right) \right\|_2 \leq c (\|\mathbf{G}\|_2 + \sigma) \delta, \quad (27)$$

$$\left\| \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} (\mathbf{g} - \mathbf{G})^{\otimes 2} - F(\mathbf{G}) \right\|_{2k, \text{op}} \leq c \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) \delta, \quad (28)$$

where  $c$  is a constant. Suppose  $\delta = \Omega(\epsilon)$ , and  $\left\| P_{2k} \left( \tilde{\Delta} \right) \right\|_2 \geq C_1 (\|\mathbf{G}\|_2 + \sigma) \delta$ , where  $C_1$  is a large constant. Then we have,

$$\max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\mathbf{G}) \right) \mathbf{v} \geq \frac{\left\| P_{2k} \left( \tilde{\Delta} \right) \right\|_2^2}{4\epsilon}, \quad (29)$$

$$\max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\hat{\mathbf{G}}) \right) \mathbf{v} \geq \frac{\left\| P_{2k} \left( \tilde{\Delta} \right) \right\|_2^2}{5\epsilon}. \quad (30)$$

*Proof.* The proof outline is similar to [9] and [1], but we focus on the  $2k$ -sparse largest eigenvalue.

Let  $\mathcal{S} = \arg \max_{S' \subset [d], |S'| \leq 2k} \left\| \tilde{\Delta}^{S'} \right\|_2$ . Then  $\tilde{\Delta}^{\mathcal{S}} = \left\| P_{2k} \left( \tilde{\Delta} \right) \right\|_2 \geq C_1 (\|\mathbf{G}\|_2 + \sigma) \delta$  according to the assumption. Using  $n$  to denote the size of  $S_{\text{in}}$ , we have a lower bound for the sum over bad samples:

$$\left\| \frac{1}{n} \sum_{i \in S_{\text{bad}}} (\mathbf{g}_i^{\mathcal{S}} - \mathbf{G}^{\mathcal{S}}) \right\|_2 = \left\| \tilde{\Delta}^{\mathcal{S}} - \frac{1}{n} \sum_{i \in S_{\text{good}}} (\mathbf{g}_i^{\mathcal{S}} - \mathbf{G}^{\mathcal{S}}) \right\|_2$$

$$\begin{aligned}
&\geq \left\| \tilde{\Delta}^S \right\|_2 - \left\| \frac{1}{n} \sum_{i \in S_{\text{good}}} (\mathbf{g}_i^S - \mathbf{G}^S) \right\|_2 \\
&\stackrel{(i)}{\geq} \left\| \tilde{\Delta}^S \right\|_2 - c(\|\mathbf{G}\|_2 + \sigma) \delta \\
&\stackrel{(ii)}{\geq} \frac{\left\| \tilde{\Delta}^S \right\|_2}{1.1},
\end{aligned}$$

where (i) follows from eq. (27) and the assumptions; (ii) follows from that we choose  $C_1$  large enough.

By PSD-ness of covariance matrices in [9], we have

$$\frac{1}{|S_{\text{bad}}|} \sum_{i \in S_{\text{bad}}} (\mathbf{g}_i^S - \mathbf{G}^S) (\mathbf{g}_i^S - \mathbf{G}^S)^\top \succcurlyeq \left( \frac{1}{|S_{\text{bad}}|} \sum_{i \in S_{\text{bad}}} (\mathbf{g}_i^S - \mathbf{G}^S) \right) \left( \frac{1}{|S_{\text{bad}}|} \sum_{i \in S_{\text{bad}}} (\mathbf{g}_i^S - \mathbf{G}^S) \right)^\top.$$

Therefore, because  $|S_{\text{bad}}| \leq 2\epsilon |S_{\text{in}}|$ , we have

$$\left\| \frac{1}{n} \sum_{i \in S_{\text{bad}}} (\mathbf{g}_i^S - \mathbf{G}^S)^{\otimes 2} \right\|_{\text{op}} \geq \frac{\left\| \frac{1}{n} \sum_{i \in S_{\text{bad}}} (\mathbf{g}_i^S - \mathbf{G}^S) \right\|_2^2}{2\epsilon} \geq \frac{\left\| \tilde{\Delta}^S \right\|_2^2}{2.5\epsilon}. \quad (31)$$

Hence, we can define a vector  $\mathbf{v}_0$  ( $2k$ -sparse) as follows:

$$\mathbf{v}_0 = \arg \max_{\|\mathbf{v}\|_2=1} \mathbf{v}^\top \left( \sum_{i \in S_{\text{bad}}} \frac{1}{n} (\mathbf{g}_i^S - \mathbf{G}^S)^{\otimes 2} \right) \mathbf{v}.$$

For this  $\mathbf{v}_0$ , we have

$$\begin{aligned}
&\mathbf{v}_0^\top \left( \frac{1}{n} \sum_{i=1}^n (\mathbf{g}_i^S - \mathbf{G}^S)^{\otimes 2} - F(\mathbf{G})^{SS} \right) \mathbf{v}_0 \\
&\geq \mathbf{v}_0^\top \left( \frac{1}{n} \sum_{i \in S_{\text{bad}}} (\mathbf{g}_i^S - \mathbf{G}^S)^{\otimes 2} \right) \mathbf{v}_0 \\
&\quad - \left\| \frac{1}{n} \sum_{i \in S_{\text{good}}} (\mathbf{g}_i^S - \mathbf{G}^S)^{\otimes 2} - \frac{|S_{\text{good}}|}{n} F(\mathbf{G})^{SS} \right\|_{\text{op}} - \left\| \frac{|S_{\text{bad}}|}{n} F(\mathbf{G})^{SS} \right\|_{\text{op}} \\
&\stackrel{(i)}{\geq} \frac{\left\| \tilde{\Delta}^S \right\|_2^2}{2.5\epsilon} - c(\|\mathbf{G}\|_2^2 + \sigma^2) \delta - 2\epsilon(\|\mathbf{G}\|_2^2 + \sigma^2) \\
&\stackrel{(ii)}{\geq} \frac{\left\| \tilde{\Delta}^S \right\|_2^2}{3\epsilon}, \quad (32)
\end{aligned}$$

where (i) follows from eq. (28) and eq. (31); (ii) follows from the assumption that  $\epsilon$  is sufficiently small.

Applying eq. (32) on our target  $\mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\mathbf{G})$ , we have

$$\mathbf{v}_0^\top \left( \frac{1}{n} \sum_{i=1}^n (\mathbf{g}_i^S - \hat{\mathbf{G}}^S)^{\otimes 2} - F(\mathbf{G})^{SS} \right) \mathbf{v}_0$$

$$\begin{aligned}
&= \mathbf{v}_0^\top \left( \frac{1}{n} \sum_{i=1}^n (\mathbf{g}_i^S - \mathbf{G}^S)^{\otimes 2} - F(\mathbf{G})^{SS} - \widehat{\Delta}^S (\widetilde{\Delta}^S)^\top - \widetilde{\Delta}^S (\widehat{\Delta}^S)^\top + (\widehat{\Delta}^S)^{\otimes 2} \right) \mathbf{v}_0 \\
&\stackrel{(i)}{\geq} \mathbf{v}_0^\top \left( \frac{1}{n} \sum_{i=1}^n (\mathbf{g}_i^S - \mathbf{G}^S)^{\otimes 2} - F(\mathbf{G})^{SS} \right) \mathbf{v}_0 - 24 \left( \left\| \widetilde{\Delta}^S \right\|_2^2 \right) \\
&\stackrel{(ii)}{\geq} \frac{\left\| \widetilde{\Delta}^S \right\|_2^2}{4\epsilon}, \tag{33}
\end{aligned}$$

where (i) follows from Lemma D.1; (ii) follows from eq. (32) and  $\epsilon$  is sufficiently small. It is easy to see that  $\mathbf{v}_0$  provides a lower bound for the maximum of  $\{\mathbf{v} : \|\mathbf{v}\|_2 = 1, \|\mathbf{v}\|_0 \leq 2k\}$  in eq. (29).

By eq. (33), we already know that

$$\mathbf{v}_0^\top \left( \frac{1}{n} \sum_{i=1}^n (\mathbf{g}_i^S - \widehat{\mathbf{G}}^S)^{\otimes 2} - F(\mathbf{G})^{SS} \right) \mathbf{v}_0 \geq \frac{\left\| \widetilde{\Delta}^S \right\|_2^2}{4\epsilon}.$$

By our assumptions on  $F$ , we have

$$\begin{aligned}
\left\| F(\mathbf{G}) - F(\widehat{\mathbf{G}}) \right\|_{2k, \text{op}} &\leq L_F \left\| \widehat{\Delta} \right\|_2 + C \left\| \widehat{\Delta} \right\|_2^2 \\
&\stackrel{(i)}{\leq} 5L_F \left\| \widetilde{\Delta}^S \right\|_2 + 5C \left\| \widetilde{\Delta}^S \right\|_2^2,
\end{aligned}$$

where (i) follows from Lemma D.1. Since  $\delta = \Omega(\epsilon)$ , we obtain eq. (30) by using the triangle inequality.  $\square$

## D.4 Proof of Corollary 2.1

Equipped with Lemma D.1, Lemma D.2 and Lemma D.3, we can now prove Corollary 2.1.

**Corollary D.1** (Corollary 2.1). *Under the same setting as Theorem 2.1, if we use the ellipsoid algorithm for robust sparse gradient estimation, it requires sample complexity  $N(k, d, \epsilon, \nu) = \Omega\left(\frac{k^2 \log(dT/\nu)}{\epsilon^2}\right) T$ , and guarantees  $\psi(\epsilon) = \widetilde{O}(\epsilon^2 \sigma^2)$ . Hence, Algorithm 1 outputs  $\widehat{\beta}$ , such that*

$$\left\| \widehat{\beta} - \beta^* \right\|_2 = \widetilde{O}(\sigma \epsilon), \tag{34}$$

with probability at least  $1 - \nu$ , by setting  $T = \Theta\left(\log\left(\frac{\|\beta^*\|_2}{\epsilon \sigma}\right)\right)$ .

*Proof.* Consider the  $t$ -th iteration, we omit  $t$  in  $\mathbf{g}_i^t$  and  $\mathbf{G}^t$ . Function  $F(\mathbf{G})$  is calculated as

$$F(\mathbf{G}) = \|\mathbf{G}\|_2^2 I + \mathbf{G} \mathbf{G}^\top + \sigma^2 I,$$

by Appendix B. The accuracy in robust sparse estimation on gradients depends on two parameters for  $F(\mathbf{G})$ :  $L_{\text{cov}} = 2 \|\mathbf{G}\|_2^2 + \sigma^2$ , and  $L_F = 4 \|\mathbf{G}\|_2$ , which are calculated in Appendix B.

Under the statistical model and the contamination model described in Theorem 2.1, we can set the parameters  $L_{\text{cov}}$  and  $L_F$  in the ellipsoid algorithm.

The ellipsoid algorithm considers all possible sample weights in a convex set and finds the optimal weight for each sample. The algorithm iteratively uses a separation oracle, which solves the convex relaxation of Sparse PCA at each iteration:

$$\lambda^* = \max_H \text{Tr} \left( \left( \widehat{\Sigma} - F(\widehat{\mathbf{G}}) \right) \cdot H \right), \quad \text{subject to } H \succcurlyeq 0, \|H\|_{1,1} \leq 2k, \text{Tr}(H) = 1, \tag{35}$$

To prove the Main Theorem (Theorem 3.1) in [1], the only modification is to replace the lower bound of  $\lambda^*$  in Lemma A.3 of [1].

A weighted version of Lemma D.3 implies

$$\max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} v^\top \left( \sum_{i=1}^{|S_{\text{in}}|} w_i (\mathbf{g}_i - \widehat{\mathbf{G}}(w))^{\otimes 2} - F(\widehat{\mathbf{G}}(w)) \right) v \geq \frac{\|P_{2k}(\widetilde{\Delta}(w))\|_2^2}{5\epsilon}, \quad (36)$$

where  $\widehat{\mathbf{G}}(w) = P_{4k} \left( \sum_{i=1}^{|S_{\text{in}}|} w_i \mathbf{g}_i \right)$ , and  $\widetilde{\Delta}(w) = \sum_{i=1}^{|S_{\text{in}}|} w_i \mathbf{g}_i - \mathbf{G}$ . Then,  $\lambda^*$  in the ellipsoid algorithm satisfies

$$\lambda^* \geq \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} v^\top \left( \sum_{i=1}^{|S_{\text{in}}|} w_i (\mathbf{g}_i - \widehat{\mathbf{G}}(w))^{\otimes 2} - F(\widehat{\mathbf{G}}(w)) \right) v, \quad (37)$$

since trace norm maximization is the convex relaxation of finding the  $2k$ -sparse largest eigenvalue.

Combining eq. (36) and eq. (37), we have

$$\lambda^* \geq \frac{\|P_{2k}(\widetilde{\Delta}(w))\|_2^2}{5\epsilon}, \quad (38)$$

which recovers the correctness of the separation oracle in the ellipsoid algorithm, and the Main Theorem (Theorem 3.1) in [1].

Finally, the ellipsoid algorithm guarantees that, with sample complexity  $\Omega\left(\frac{k^2 \log(d/\nu)}{\epsilon^2}\right)$ , the estimate  $\widehat{\mathbf{G}}$  satisfies

$$\|\widehat{\mathbf{G}} - \mathbf{G}\|_2^2 = \widetilde{O}(\epsilon^2 (L_{\text{F}}^2 + L_{\text{cov}})) = \widetilde{O}(\epsilon^2 (\|\mathbf{G}\|_2^2 + \sigma^2)), \quad (39)$$

with probability at least  $1 - \nu$ . This exactly gives us a  $\widetilde{O}(\epsilon^2 \sigma^2)$ -RSGE. Hence, we can apply eq. (39) as the RSGE in Theorem 2.1 to prove Corollary 2.1.  $\square$

## E Proofs for Lemma 3.1

In this section, we consider a single iteration of Algorithm 1 at the  $t$ -th step. For clarity, we omit the superscript  $t$  in both  $\mathbf{g}_i^t$  and  $\mathbf{G}^t$ .

In order to show guarantees for Lemma 3.1, we leverage previous results Lemma D.2 and Lemma D.3. We state Lemma E.1 as a modification of Lemma D.2 by replacing  $\epsilon$  by  $\sqrt{\epsilon}$ , using concentration results in Lemma D.2, and replacing  $\epsilon$  by  $\sqrt{\epsilon}$ . We state Lemma E.2 as a modification of Lemma D.3 by replacing  $\delta = \Omega(\epsilon)$  with  $\delta = \Omega(\sqrt{\epsilon})$ , since the results for  $\delta = \Omega(\epsilon)$  implies the results for  $\delta = \Omega(\sqrt{\epsilon})$ .

The reason we modify the above is to prove guarantees for our computationally efficient RSGE described in Algorithm 2. Our motivation for calculating the score for each sample according to eq. (10) is to make sure that all the scores  $\tau_i$  are positive (notice that the scores calculated based on the original non-p.s.d matrix may be negative). Based on this, we show that the sum of scores over all bad samples is a large constant ( $> 1$ ) times larger than the sum of scores over all good samples. When finding an upper bound for  $\sum_{i \in S_{\text{good}}} \tau_i$ , we compromise an  $\epsilon$  factor in the value of  $\lambda^*$ , which results in an  $\sqrt{\epsilon}$  factor in the recovery guarantee.

**Lemma E.1.** *Suppose we observe gradient samples  $\{\mathbf{g}\}_{i=1}^n$  from eq. (2) with  $n = \Omega\left(\frac{k \log(d/\nu)}{\epsilon}\right)$ . Denote this set as  $\mathcal{G}$ . Then there is a  $\delta = \tilde{O}(\sqrt{\epsilon})$  that with probability at least  $1 - \nu$ , we have for any subset  $\mathcal{S} \subset [d]$ ,  $|\mathcal{S}| \leq 2k$ , the followings hold*

$$\left\| \mathbb{E}_{\mathbf{g} \in_u \mathcal{G}} (\mathbf{g}^{\mathcal{S}}) - \mathbf{G}^{\mathcal{S}} \right\|_2 \leq \delta (\|\mathbf{G}\|_2 + \sigma), \quad (40)$$

$$\left\| \mathbb{E}_{\mathbf{g} \in_u \mathcal{G}} (\mathbf{g}^{\mathcal{S}} - \mathbf{G}^{\mathcal{S}})^{\otimes 2} - F(\mathbf{G})^{\mathcal{S}\mathcal{S}} \right\|_{\text{op}} \leq \delta \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right). \quad (41)$$

**Lemma E.2.** *Suppose  $|S_{\text{bad}}| \leq 2\epsilon|S_{\text{in}}|$ , and the gradient samples in  $S_{\text{good}}$  satisfy*

$$\left\| P_{2k} \left( \tilde{\Delta}_{S_{\text{good}}} \right) \right\|_2 \leq c (\|\mathbf{G}\|_2 + \sigma) \delta, \quad (42)$$

$$\left\| \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} (\mathbf{g} - \mathbf{G})^{\otimes 2} - F(\mathbf{G}) \right\|_{2k, \text{op}} \leq c \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) \delta, \quad (43)$$

where  $c$  is a constant. Suppose  $\delta = \Omega(\sqrt{\epsilon})$ , and  $\left\| P_{2k} \left( \tilde{\Delta} \right) \right\|_2 \geq C_1 (\|\mathbf{G}\|_2 + \sigma) \delta$ , where  $C_1$  is a constant. Then we have,

$$\max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\mathbf{G}) \right) \mathbf{v} \geq \frac{\left\| P_{2k} \left( \tilde{\Delta} \right) \right\|_2^2}{4\epsilon}, \quad (44)$$

$$\max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\hat{\mathbf{G}}) \right) \mathbf{v} \geq \frac{\left\| P_{2k} \left( \tilde{\Delta} \right) \right\|_2^2}{5\epsilon}. \quad (45)$$

By Lemma E.1, eq. (42) and eq. (43) in Lemma E.2 are satisfied, provided that we have  $n = \Omega\left(\frac{k \log(d/\tau)}{\epsilon}\right)$ . Now, equipped with Lemma E.1 and Lemma E.2, the effect of good samples can be controlled by concentration inequalities. Based on these, we are ready to prove Lemma 3.1.

**Lemma E.3** (Lemma 3.1). *Suppose we observe  $n = \Omega\left(\frac{k^2 \log(d/\nu)}{\epsilon}\right)$   $\epsilon$ -corrupted samples from eq. (1). Let  $S_{\text{in}}$  be an  $\epsilon$ -corrupted set of gradient samples  $\{\mathbf{g}_i^t\}_{i=1}^n$ . Algorithm 2 computes  $\lambda^*$  that satisfies*

$$\lambda^* \geq \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\hat{\mathbf{G}}) \right) \mathbf{v}. \quad (46)$$

Furthermore, if  $\lambda^* \geq \rho_{\text{sep}} = C_\gamma \left( \|\mathbf{G}^t\|_2^2 + \sigma^2 \right)$ , then with probability at least  $1 - \nu$ , the contribution of  $\sum_{i \in S_{\text{good}}} \tau_i$  is small:

$$\sum_{i \in S_{\text{good}}} \tau_i \leq \frac{1}{\gamma} \sum_{i \in S_{\text{in}}} \tau_i, \quad (47)$$

where  $\tau_i$  is defined in eq. (10),  $C_\gamma$  is a constant depending on  $\gamma$ , and  $\gamma \geq 2$  is a constant.

*Proof.* Since  $\lambda^*$  is the natural convex relaxation of Sparse PCA, we have

$$\begin{aligned} \lambda^* &= \text{Tr} \left( H^* \cdot \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\hat{\mathbf{G}}) \right) \right) \\ &\geq \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\hat{\mathbf{G}}) \right) \mathbf{v}. \end{aligned}$$

Then, we start to prove eq. (47). By Theorem A.1 in [1], we have

$$\text{Tr} \left( H^* \cdot \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} (\mathbf{g} - \hat{\mathbf{G}})^{\otimes 2} - F(\hat{\mathbf{G}}) \right) \right)$$

$$\leq C \left( \left\| \widehat{\Delta} \right\|_2^2 + \left( L_F + k \left\| \widetilde{\Delta}_{S_{\text{good}}} \right\|_\infty \right) \left\| \widehat{\Delta} \right\|_2 + k \left\| \mathbb{E}_{S_{\text{good}}} (g_i - \mathbf{G})^{\otimes 2} - F(\mathbf{G}) \right\|_\infty \right), \quad (48)$$

where  $C$  is a constant. Noticing that  $\left\| \widetilde{\Delta}_{S_{\text{good}}} \right\|_\infty$  and  $\left\| \mathbb{E}_{S_{\text{good}}} (g_i - \mathbf{G})^{\otimes 2} - F(\mathbf{G}) \right\|_\infty$  are unrelated to  $\widehat{\mathbf{G}}$  and only defined on  $S_{\text{good}}$ , [1] showed concentration bounds for these two terms, when  $n = \Omega\left(\frac{k^2 \log(d/\nu)}{\epsilon}\right)$ . The following holds with probability at least  $1 - \nu$ ,

$$\left\| \widetilde{\Delta}_{S_{\text{good}}} \right\|_\infty \leq C_1 \left( L_F + \sqrt{L_{\text{cov}}} \right) \sqrt{\epsilon}/k \quad (49)$$

$$\left\| \mathbb{E}_{S_{\text{good}}} (g_i - \mathbf{G})^{\otimes 2} - F(\mathbf{G}) \right\|_\infty \leq C_1 (L_F^2 + L_{\text{cov}}) \sqrt{\epsilon}/k \quad (50)$$

Now, we focus on the LHS of eq. (47):

$$\begin{aligned} \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} \tau_i &= \text{Tr} \left( H^* \cdot \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} \left( \mathbf{g} - \widehat{\mathbf{G}} \right)^{\otimes 2} \right) \right) \\ &= \text{Tr} \left( H^* \cdot \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} \left( \mathbf{g} - \widehat{\mathbf{G}} \right)^{\otimes 2} - F(\widehat{\mathbf{G}}) \right) \right) + \text{Tr} \left( H^* \cdot F(\widehat{\mathbf{G}}) \right) \\ &\stackrel{(i)}{\leq} C \left( \left\| \widehat{\Delta} \right\|_2^2 + \left( L_F + k \left\| \widetilde{\Delta}_{S_{\text{good}}} \right\|_\infty \right) \left\| \widehat{\Delta} \right\|_2 + k \left\| \mathbb{E}_{S_{\text{good}}} (g_i - \mathbf{G})^{\otimes 2} - F(\mathbf{G}) \right\|_\infty \right) + \text{Tr} \left( H^* \cdot F(\widehat{\mathbf{G}}) \right) \end{aligned}$$

where (i) follows from eq. (48).

To bound the RHS of above, we first bound  $\text{Tr} \left( H^* \cdot F(\widehat{\mathbf{G}}) \right)$ . Because of the constraint of convex programming,  $H^*$  belongs to Fantope  $\mathcal{F}^1$  [28], for any matrix  $A$ , we have  $\text{Tr} (A \cdot H^*) \leq \|A\|_{\text{op}}$ .

Then, we have

$$\begin{aligned} \text{Tr} \left( H^* \cdot F(\widehat{\mathbf{G}}) \right) &= \text{Tr} \left( H^* \cdot F(\mathbf{G}) \right) + \text{Tr} \left( H^* * \left( F(\widehat{\mathbf{G}}) - F(\mathbf{G}) \right) \right) \\ &\leq \|F(\mathbf{G})\|_{\text{op}} + \left\| F(\widehat{\mathbf{G}}) - F(\mathbf{G}) \right\|_{\text{op}} \\ &\stackrel{(i)}{\leq} C_1 \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) + \left\| F(\widehat{\mathbf{G}}) - F(\mathbf{G}) \right\|_{\text{op}} \\ &\stackrel{(ii)}{\leq} C_1 \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) + L_F \left\| \widehat{\Delta} \right\|_2 + C_2 \left\| \widehat{\Delta} \right\|_2^2, \end{aligned} \quad (51)$$

where (i) follows from the expression of  $F(G)$ ; (ii) follows from the smoothness definition of  $F(G)$ .

By plugging in the concentration guarantees eq. (49) and combining eq. (51), we have

$$\begin{aligned} \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} \tau_i &\leq C_2 \left( (L_F^2 + L_{\text{cov}}) \sqrt{\epsilon} + \left( (L_F + \sqrt{L_{\text{cov}}}) \sqrt{\epsilon} + L_F \right) \left\| \widehat{\Delta} \right\|_2 + \left\| \widehat{\Delta} \right\|_2^2 \right) + C_1 \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) \\ &\stackrel{(i)}{\leq} C_2 \left( \|\mathbf{G}\|_2 \left\| \widehat{\Delta} \right\|_2 + \left\| \widehat{\Delta} \right\|_2^2 \right) + C_1 \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) \\ &\leq C_1 \left( \|\mathbf{G}\|_2 \left\| \widehat{\Delta} \right\|_2 + \left\| \widehat{\Delta} \right\|_2^2 + \|\mathbf{G}\|_2^2 + \sigma^2 \right), \end{aligned} \quad (52)$$

where (i) follows from the fact  $\epsilon$  is sufficiently small.

On the other hand, we have a lower bound for  $\mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} \tau_i$ :

$$\mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} \tau_i = \lambda^* + \text{Tr} \left( H^* \cdot F(\widehat{\mathbf{G}}) \right) \geq \lambda^*.$$

Now, under the condition of eq. (14), i.e.,  $\lambda^* \geq \rho_{\text{sep}} = \Theta\left(\|\mathbf{G}\|_2^2 + \sigma^2\right)$ , we analyze eq. (52) in two scenarios. By separating two cases, we can always show  $\lambda^*$  is very large, and the contribution from good samples is limited.

First, if  $\|\widehat{\Delta}\|_2^2 \geq \Theta\left(\|\mathbf{G}\|_2^2 + \sigma^2\right)$ , we have

$$\|\widehat{\Delta}\|_2^2 \gtrsim \|\mathbf{G}\|_2, \quad \|\widehat{\Delta}\|_2 \gtrsim \|\mathbf{G}\|_2^2, \quad \text{and} \quad \|\widehat{\Delta}\|_2^2 \gtrsim \sigma^2.$$

That said, we only need to compare  $\lambda^*$  and  $\|\widehat{\Delta}\|_2^2$ . By Lemma E.2, we have

$$\mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} \tau_i \geq \lambda^* \geq \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq 2k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} (\mathbf{g} - \widehat{\mathbf{G}})^{\otimes 2} - F(\widehat{\mathbf{G}}) \right) \mathbf{v} \geq \frac{\|\widehat{\Delta}\|_2^2}{\epsilon}.$$

Hence, by eq. (52), we have  $\mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} \tau_i \geq \gamma \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} \tau_i$ , where  $\gamma \geq 2$  is a constant.

Second, if  $\|\widehat{\Delta}\|_2^2 \leq \Theta\left(\|\mathbf{G}\|_2^2 + \sigma^2\right)$ , we have

$$\|\mathbf{G}\|_2^2 \gtrsim \|\mathbf{G}\|_2, \quad \|\widehat{\Delta}\|_2 \gtrsim \|\widehat{\Delta}\|_2^2, \quad \text{or} \quad \sigma^2 \gtrsim \|\widehat{\Delta}\|_2^2.$$

That said, we only need to compare  $\lambda^*$  and  $\max\left(\|\mathbf{G}\|_2^2, \sigma^2\right)$ . Since  $\lambda^* \geq \Theta\left(\|\mathbf{G}\|_2^2 + \sigma^2\right)$  by the condition of Lemma 3.1, we still have  $\mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} \tau_i \geq \gamma \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} \tau_i$ , where  $\gamma \geq 2$  is a constant.

Combing all of above, and setting  $\rho_{\text{sep}} = C_\gamma \left(\|\mathbf{G}\|_2^2 + \sigma^2\right)$ , we have

$$\sum_{i \in S_{\text{in}}} \tau_i = |S_{\text{in}}| \mathbb{E}_{\mathbf{g} \in_u S_{\text{in}}} \tau_i \geq \gamma |S_{\text{good}}| \mathbb{E}_{\mathbf{g} \in_u S_{\text{good}}} \tau_i = \gamma \sum_{i \in S_{\text{good}}} \tau_i.$$

□

## F Proofs for Theorem 3.1

In this section, we still consider the  $t$ -th iteration of Algorithm 1 on  $t$ , and omit  $t$  in  $\mathbf{g}^t$  and  $\mathbf{G}^t$ .

In the case of  $\lambda^* \geq C_\gamma \left(\|\mathbf{G}\|_2^2 + \sigma^2\right)$ , Algorithm 2 iteratively removes one sample according to the probability distribution eq. (11). We denote the steps of this outlier removal procedure as  $l = 1, 2, \dots, n$ . The first step of proving Theorem 3.1 is to show we can remove a corrupted samples with high probability at each step, which is a result by Lemma 3.1.

Intuitively, if all subsequent steps are i.i.d., we can expect Algorithm 2 to remove outliers within around  $\epsilon n$  iterations, with exponentially high probability. However, the subsequent steps in Algorithm 2 are not independent. To achieve this, we use a martingale argument.

### F.1 Supermartingale construction

Let  $\mathcal{F}^l$  be the filtration generated by the set of events until step  $l$ . We define the corresponding set  $S_{\text{in}}^l$ ,  $S_{\text{good}}^l$  and  $S_{\text{bad}}^l$  at the step  $l$ . We have that  $S_{\text{in}}^l, S_{\text{good}}^l, S_{\text{bad}}^l \in \mathcal{F}^l$ , and  $|S_{\text{in}}^l| = n - l$ .

We denote a good event  $\mathcal{E}^l$  at step  $l$  as

$$\sum_{i \in S_{\text{bad}}^l} \tau_i \leq (\gamma - 1) \sum_{i \in S_{\text{good}}^l} \tau_i.$$

Then, by the definition of Algorithm 2 and Lemma 3.1, if  $\lambda^* \geq C_\gamma (\|\mathbf{G}\|_2^2 + \sigma^2)$ ,  $\mathcal{E}^l$  is not true; if  $\mathcal{E}^l$  is true, then Algorithm 2 will return a  $\widehat{\mathbf{G}}$ .

In Lemma F.1, we show that at any step  $l$  when  $\mathcal{E}^l$  is not true, the random outlier removal procedure removes a corrupted sample with probability at least  $(\gamma - 1)/\gamma$ .

**Lemma F.1.** *In each subsequent step  $l$ , if  $\mathcal{E}^l$  is not true, then we can remove one remaining outlier from  $S_{\text{in}}^l$  with probability at least  $(\gamma - 1)/\gamma$ :*

$$\Pr(\text{one sample from } S_{\text{bad}}^l \text{ is removed} \mid \mathcal{F}_l) \geq \frac{\gamma - 1}{\gamma}.$$

*Proof of Lemma F.1.* When  $\lambda^* \geq C_\gamma (\|\mathbf{G}\|_2^2 + \sigma^2)$ , Lemma 3.1 implies

$$\sum_{i \in S_{\text{bad}}^l} \tau_i \geq (\gamma - 1) \sum_{i \in S_{\text{good}}^l} \tau_i.$$

Then we randomly remove a sample  $r$  from  $S_{\text{in}}$  according to

$$\Pr(\mathbf{g}_i \text{ is removed} \mid \mathcal{F}_l) = \frac{\tau_i}{\sum_{i \in S_{\text{in}}^l} \tau_i}.$$

Finally,

$$\Pr(\text{one sample from } S_{\text{bad}}^l \text{ is removed} \mid \mathcal{F}_l) = \sum_{i \in S_{\text{bad}}^l} \frac{\tau_i}{\sum_{i \in S_{\text{in}}^l} \tau_i} \geq \frac{\gamma - 1}{\gamma}.$$

□

Since subsequent steps for applying Algorithm 2 on  $S_{\text{in}}$  are not independent, we need martingale arguments to show the total iterations of applying Algorithm 2 is limited.

We use the martingale technique in [29], by defining  $T$ :  $T = \min\{l : \mathcal{E}^l \text{ is true}\}$ . Based on  $T$ , we define a random variable:

$$Y^l = \begin{cases} |S_{\text{bad}}^{T-1}| + \frac{\gamma-1}{\gamma} (T-1), & \text{if } l \geq T \\ |S_{\text{bad}}^l| + \frac{\gamma-1}{\gamma} l, & \text{if } l < T \end{cases}$$

**Lemma F.2** (Lemma 1 in [29]).  *$\{Y^l, \mathcal{F}^l\}$  is a supermartingale.*

Now, equipped with Lemma F.1 and Lemma F.2, we are ready to prove Theorem 3.1.

## F.2 Proof of Theorem 3.1

**Theorem F.1** (Theorem 3.1). *Consider the statistical model in eq. (2), and we observe  $\epsilon$ -corrupted samples from eq. (1) of size  $n = \Omega\left(\frac{k^2 \log(d/\nu)}{\epsilon}\right)$ . Let  $S_{\text{in}}$  be an  $\epsilon$ -corrupted set of gradient samples  $\{\mathbf{g}_i^t\}_{i=1}^n$ . If we run Algorithm 2 iteratively with initial set  $S_{\text{in}}$ , and  $\rho_{\text{sep}} = C_\gamma (\|\mathbf{G}^t\|_2^2 + \sigma^2)$ , it will stop after at most  $\frac{1.1\gamma}{\gamma-1} \epsilon n$  iterations, and output  $\widehat{\mathbf{G}}^t$ , such that*

$$\|\widehat{\mathbf{G}}^t - \mathbf{G}^t\|_2^2 = \tilde{O}\left(\epsilon (\|\mathbf{G}^t\|_2^2 + \sigma^2)\right),$$

*with probability at least  $1 - \nu - \exp(-\Theta(\epsilon n))$ . Here,  $C_\gamma$  is a constant depending on  $\gamma$ , and  $\gamma \geq 2$  is a constant.*



*Proof.* We analyze Algorithm 2 by discussing a series of  $\{\mathcal{E}^l\}$ .

If  $\mathcal{E}^l$  is true, then  $\lambda^* \leq \rho_{\text{sep}} = C_\gamma \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right)$ . By Lemma E.2, we have

$$\lambda^* \geq \max_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_0 \leq k} \mathbf{v}^\top \left( \mathbb{E}_{\mathbf{g} \in \mathcal{u}_{S_{\text{in}}}} \left( \mathbf{g} - \widehat{\mathbf{G}} \right)^{\otimes 2} - F \left( \widehat{\mathbf{G}} \right) \right) \mathbf{v} \geq \frac{\|P_{2k} \left( \widetilde{\Delta}_S \right)\|_2^2}{5\epsilon}.$$

Plugging in  $\lambda^* \leq C_\gamma \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right)$ , we have

$$\frac{1}{5} \|\widehat{\Delta}_S\|_2^2 \stackrel{(i)}{\leq} \|P_{2k} \left( \widetilde{\Delta}_S \right)\|_2^2 \leq 5\epsilon\lambda^* \leq O \left( \epsilon \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) \right),$$

where (i) follows from Lemma D.1. Hence, when  $\mathcal{E}^l$  is true, Algorithm 2 can return a  $\widehat{\mathbf{G}}$ , such that  $\|\widehat{\mathbf{G}} - \mathbf{G}\|_2^2 \leq O \left( \epsilon \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) \right)$

Then, we only need to show  $\bigcup_{l=1}^L \mathcal{E}^l$  is true, where  $L = \frac{1.1\gamma}{\gamma-1} \epsilon n$ , with high probability. That said, we need to upper bound the probability

$$\Pr \left( \bigcap_{l=1}^L \overline{\mathcal{E}^l} \right) = \Pr (T \geq L) \leq \Pr \left( Y^L \geq \frac{\gamma-1}{\gamma} L \right) = \Pr (Y^L \geq 1.1\epsilon n). \quad (53)$$

Then, we can construct the martingale difference according to [29]. Let  $D^l = Y^l - Y^{l-1}$ , where  $Y^0 = \epsilon n$ , and

$$\bar{D}^l = D^l - \mathbb{E} (D^l | D^1, \dots, D^{l-1}).$$

$\{\bar{D}^l\}$  is a martingale difference process, and  $\mathbb{E} (D^l | D^1, \dots, D^{l-1}) \leq 0$ , since  $\{Y^l\}$  is a supermartingale. Now, eq. (53) can be viewed as a bound for the sum of the associated martingale difference sequence.

$$Y^L - Y^0 = \sum_{j=1}^L D^j = \sum_{j=1}^L \bar{D}^j + \sum_{j=1}^L \mathbb{E} (D^j | D^1, \dots, D^{j-1}) \leq \sum_{j=1}^L \bar{D}^j.$$

Since we only remove one example from the  $S_{\text{in}}^l$ , we can guarantee  $|D^l| \leq 1$  and  $|\bar{D}^l| \leq 2$ . For these bounded random variables, by applying Azuma-Hoeffding inequality, we have

$$\begin{aligned} \Pr (Y^L \geq 1.1\epsilon n) &\leq \Pr \left( \sum_{l=1}^L \bar{D}^l \geq 0.1\epsilon n \right) \\ &\leq \exp \left( \frac{-(0.1\epsilon n)^2}{8L} \right). \end{aligned}$$

Plugging in  $L = \frac{1.1\gamma}{\gamma-1} \epsilon n$ , this probability is upper bounded by  $\exp(-\Theta(\epsilon n))$ .

Notice that  $L = \frac{1.1\gamma}{\gamma-1} \epsilon n \leq n/2$ , for a small  $\epsilon$ . Even in the case we do not remove one outlier from  $l = 1$  to  $L$ , we always have  $|S_{\text{bad}}^l| \leq 2\epsilon |S_{\text{in}}^l|$ . Then Lemma E.2 hold and Lemma 3.1 is still valid.

Combining all of above, we have proven that, with exponentially high probability, Algorithm 2 will return a  $\widehat{\mathbf{G}}$  satisfying  $\|\widehat{\mathbf{G}} - \mathbf{G}\|_2^2 \leq O \left( \epsilon \left( \|\mathbf{G}\|_2^2 + \sigma^2 \right) \right)$ , within  $\frac{1.1\gamma}{\gamma-1} \epsilon n$  iterations.  $\square$