

**NUMERICAL SIMULATION OF 2.5-SET OF ITERATED ITO STOCHASTIC
INTEGRALS OF MULTIPLICITIES 1 TO 5 FROM THE TAYLOR–ITO
EXPANSION**

DMITRIY F. KUZNETSOV

ABSTRACT. The article is devoted to the construction of effective procedures of the mean-square approximation of iterated Ito stochastic integrals of multiplicities 1 to 5 from the Taylor–Ito expansion based on multiple Fourier–Legendre series. The results of the article can be applied to the implementation of numerical methods with orders 1.5, 2.0, and 2.5 of strong convergence for Ito stochastic differential equations with multidimensional non-commutative noise.

CONTENTS

1.	Introduction	2
2.	Explicit One-Step Strong Numerical Schemes With Convergence Orders 1.5, 2.0, and 2.5 Based on the Unified Taylor–Ito Expansion	3
3.	Method of Numerical Modeling for Iterated Ito Stochastic Integrals Based on Generalized Multiple Fourier Series	6
4.	Calculation of the Mean-Square Approximation Error in the Method of Generalized Multiple Fourier Series	11
5.	Expansions of Iterated Stratonovich Stochastic Integrals Based on Multiple Fourier– Legendre Series and Multiple Trigonometric Fourier Series	13
6.	Approximation of Iterated Ito Stochastic Integrals Based on Multiple Fourier–Legendre Series	18
7.	Numerical Algorithm With the Order 2.5 of Strong Convergence	25
8.	Conclusions	26
	References	26

MATHEMATICS SUBJECT CLASSIFICATION: 60H05, 60H10, 42B05, 42C10.

KEYWORDS: EXPLICIT ONE-STEP STRONG NUMERICAL METHOD, ITERATED ITO STOCHASTIC INTEGRAL, ITERATED STRATONOVICH STOCHASTIC INTEGRAL, TAYLOR–ITO EXPANSION, GENERALIZED MULTIPLE FOURIER SERIES, MULTIPLE FOURIER–LEGENDRE SERIES, MULTIPLE TRIGONOMETRIC FOURIER SERIES, ITO STOCHASTIC DIFFERENTIAL EQUATION, NUMERICAL INTEGRATION, MEAN-SQUARE CONVERGENCE, EXPANSION, APPROXIMATION.

1. INTRODUCTION

This paper is a continuation of the author's research [1], [2] on numerical methods with rather high orders of strong convergence for Ito stochastic differential equations (SDEs). The definition of strong convergence will be given below. The development of such numerical methods is topical due to a wide range of applications for Ito SDEs [3]-[9]. In particular, these equations arise in optimal stochastic control, signal filtering against the background of random noises, parameter estimation for stochastic systems as well as in stochastic stability and bifurcations analysis [3]-[9]. Also the Ito SDEs represent adequate mathematical models for dynamic systems of different physical origin that are affected by random perturbations. They are used as mathematical models in stochastic mathematical finance, hydrology and seismology, geophysics, chemical kinetics and population dynamics, electrodynamics, medicine and other fields (see [3]-[9]). On the other hand, new numerical methods with rather high orders of strong convergence are needed for the Ito SDEs because one of the elementary numerical methods — the Euler scheme has insufficient accuracy for a series of practical problems under standard assumptions; the details can be found in [3].

This paper follows a promising approach to the numerical integration of Ito SDEs [3], [7]-[9] that is based on the stochastic analogs of the Taylor formula (the so-called Taylor–Ito and Taylor–Stratonovich expansions [3], [7]-[14]) for the solutions of Ito SDEs. This approach includes the finite partitioning of the time variable and also the numerical solution of an Ito SDE at discrete moments of time using the stochastic analogs of the Taylor formula. The numerical methods with orders 1.5, 2.0, and 2.5 of strong convergence will be considered in the article.

This paper employs the so-called unified Taylor–Ito expansion [12]-[21] with a minimum set of iterated Ito stochastic integrals, which is a simplifying factor at the implementation stage of numerical methods. The iterated Ito stochastic integrals figuring in the numerical schemes with a strong convergence of orders 1.5, 2.0, and 2.5 are approximated using the method of generalized multiple Fourier series, which was considered in a series of papers of the author [14]-[57]. As was noted in [1], [2], this method does not require the splitting of an integration interval $[t, T]$ of the iterated Ito stochastic integrals; recall that its length $T - t$ gives the integration step of the numerical methods for Ito SDEs and hence is a sufficiently small value. In accordance with experimental results [14], the splitting of the interval $[t, T]$ leads to an inadmissibly high computational cost. As a rule, this splitting is used in the approximation methods of iterated Ito stochastic integrals based on integral sums [7], [8], [58].

As mentioned in [1], [2], in a number of publications [3], [7], [8], numerical schemes with strong convergence of high orders (1.5, 2.0, and 2.5) for the Ito SDEs have been proposed. However, these methods do not contain efficient mean-square approximation procedures for the iterated Ito stochastic integrals in the case of multidimensional non-commutative noises. Generally, the authors [3], [7], [8] introduced some simplifying assumptions on the additivity, commutativity or smallness of the noises, which results in a considerable simplification of the numerical modeling problem of iterated Ito stochastic integrals. Like [1], [2], this paper will partially eliminate this drawback.

Note that the properties of numerical schemes for the Ito SDEs (including the ones with a strong convergence of orders 1.5, 2.0, and 2.5) were well studied in [3], [7], [8]. In particular, their stability was thoroughly analyzed. The goal of this paper is to develop efficient numerical modeling procedures for the iterated Ito stochastic integrals of multiplicities 1 to 5 including the exact calculation and efficient estimation of the mean-square approximation errors of these stochastic integrals.

Let $(\Omega, \mathbf{F}, \mathbf{P})$ be a complete probability space, let $\{\mathbf{F}_t, t \in [0, T]\}$ be a nondecreasing right-continuous family of σ -subfields of \mathbf{F} , and let \mathbf{f}_t be a standard m -dimensional Wiener stochastic process, which is \mathbf{F}_t -measurable for any $t \in [0, T]$. We assume that the components $\mathbf{f}_t^{(i)}$ ($i = 1, \dots, m$) of this process are independent. Consider an Ito SDE in the integral form

$$(1) \quad \mathbf{x}_t = \mathbf{x}_0 + \int_0^t \mathbf{a}(\mathbf{x}_\tau, \tau) d\tau + \int_0^t B(\mathbf{x}_\tau, \tau) d\mathbf{f}_\tau, \quad \mathbf{x}_0 = \mathbf{x}(0, \omega).$$

Here \mathbf{x}_t is some n -dimensional stochastic process satisfying the equation (1). The nonrandom functions $\mathbf{a} : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^n$, $B : \mathbb{R}^n \times [0, T] \rightarrow \mathbb{R}^{n \times m}$ guarantee the existence and uniqueness up to stochastic equivalence of a solution of the equation (1) [59]. The second integral on the right-hand side of (1) is interpreted as an Ito stochastic integral. Let \mathbf{x}_0 be an n -dimensional random variable, which is F_0 -measurable and $M\{|\mathbf{x}_0|^2\} < \infty$ (M denotes a mathematical expectation). We assume that \mathbf{x}_0 and $\mathbf{f}_t - \mathbf{f}_0$ are independent when $t > 0$.

Consider the following iterated Ito stochastic integrals

$$(2) \quad J[\psi^{(k)}]_{T,t} = \int_t^T \psi_k(t_k) \dots \int_t^{t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)},$$

where every $\psi_l(\tau)$ ($l = 1, \dots, k$) is a non-random function on $[t, T]$, $\mathbf{w}_\tau^{(i)} = \mathbf{f}_\tau^{(i)}$ for $i = 1, \dots, m$ and $\mathbf{w}_\tau^{(0)} = \tau$, $i_1, \dots, i_k = 0, 1, \dots, m$.

Note that $\psi_l(\tau) \equiv 1$ ($l = 1, \dots, k$) and $i_1, \dots, i_k = 0, 1, \dots, m$ in [3], [7], [8], [10], [11] and $\psi_l(\tau) \equiv (t - \tau)^{q_l}$ ($l = 1, \dots, k$; $q_1, \dots, q_k = 0, 1, 2, \dots$) and $i_1, \dots, i_k = 1, \dots, m$ in [12]-[28].

Effective solution of the problem of combined mean-square approximation of the iterated Ito stochastic integrals (2) of multiplicities 1 to 5 composes the subject of this article.

2. EXPLICIT ONE-STEP STRONG NUMERICAL SCHEMES WITH CONVERGENCE ORDERS 1.5, 2.0, AND 2.5 BASED ON THE UNIFIED TAYLOR-ITO EXPANSION

Introduce the definition of a strong convergence of a numerical method for the Ito SDEs.

Consider the partition $\{\tau_p\}_{p=0}^N$ of the interval $[0, T]$ with the maximum step of discretization Δ_N such that

$$0 = \tau_0 < \tau_1 < \dots < \tau_N = T.$$

Denote by $\mathbf{y}_{\tau_p} \stackrel{\text{def}}{=} \mathbf{y}_p$, $p = 0, 1, \dots, N$ the discrete approximation of the process \mathbf{x}_t , $t \in [0, T]$ (the solution of the Ito SDE (1)) that corresponds to the maximum discretization step Δ_N .

Definition 1 [3]. *The discrete approximation (numerical method) \mathbf{y}_j , $j = 0, 1, \dots, N$ that corresponds to the maximum discretization step Δ_N is said to be strongly converging with an order $\gamma > 0$ to the process \mathbf{x}_t , $t \in [0, T]$ if there exist a constant $C > 0$ that is independent of Δ_N and j ($j = 0, 1, \dots, N$) and also a value $\delta > 0$ such that*

$$(3) \quad M\{|\mathbf{x}_j - \mathbf{y}_j|\} \leq C(\Delta_N)^\gamma \quad (j = 0, 1, \dots, N)$$

for all $\Delta_N \in (0, \delta)$.

In a series of publications [7], [8], the authors considered the mean-square convergence instead of the strong convergence, which corresponds to the replacement of the condition (3) with

$$(4) \quad (M\{|\mathbf{x}_j - \mathbf{y}_j|^2\})^{1/2} \leq C(\Delta_N)^\gamma \quad (j = 0, 1, \dots, N).$$

In (3) and (4): $\mathbf{x}_{\tau_j} \stackrel{\text{def}}{=} \mathbf{x}_j$, $j = 0, 1, \dots, N$.

Obviously, by virtue of the Lyapunov inequality [3] the mean-square convergence implies the strong convergence. As it has appeared, a rather nontrivial question is which iterated stochastic integrals (Ito or Stratonovich) are preferable for the numerical integration of the Ito SDEs with a correct estimation of the mean-square approximation error. By their external view the approximations of the iterated Stratonovich stochastic integrals are simpler than the corresponding approximations of the iterated Ito stochastic integrals; see the details in Sections 3, 5 below. However, the estimation procedure of the mean-square approximation error turns out to be much easier for the iterated Ito stochastic integrals, which motivates the use of these integrals.

Consider the explicit one-step numerical scheme for the Ito SDEs that is based on the unified Taylor–Ito expansion [14]–[18]

$$\begin{aligned}
\mathbf{y}_{p+1} = & \mathbf{y}_p + \sum_{i_1=1}^m B_{i_1} \hat{I}_{(0)\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \mathbf{a} + \sum_{i_1, i_2=1}^m G_{i_2} B_{i_1} \hat{I}_{(00)\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \\
& + \sum_{i_1=1}^m \left(G_{i_1} \mathbf{a} \left(\Delta \hat{I}_{(0)\tau_{p+1}, \tau_p}^{(i_1)} + \hat{I}_{(1)\tau_{p+1}, \tau_p}^{(i_1)} \right) - L B_{i_1} \hat{I}_{(1)\tau_{p+1}, \tau_p}^{(i_1)} \right) + \\
& + \sum_{i_1, i_2, i_3=1}^m G_{i_3} G_{i_2} B_{i_1} \hat{I}_{(000)\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} + \frac{\Delta^2}{2} L \mathbf{a} + \\
(5) \quad & + \mathbf{v}_{p+1, p} + \mathbf{r}_{p+1, p},
\end{aligned}$$

where

$$\begin{aligned}
\mathbf{v}_{p+1, p} = & \sum_{i_1, i_2=1}^m \left(G_{i_2} L B_{i_1} \left(\hat{I}_{(10)\tau_{p+1}, \tau_p}^{(i_2 i_1)} - \hat{I}_{(01)\tau_{p+1}, \tau_p}^{(i_2 i_1)} \right) - L G_{i_2} B_{i_1} \hat{I}_{(10)\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \right. \\
& \left. + G_{i_2} G_{i_1} \mathbf{a} \left(\hat{I}_{(01)\tau_{p+1}, \tau_p}^{(i_2 i_1)} + \Delta \hat{I}_{(00)\tau_{p+1}, \tau_p}^{(i_2 i_1)} \right) \right) + \\
& + \sum_{i_1, i_2, i_3, i_4=1}^m G_{i_4} G_{i_3} G_{i_2} B_{i_1} \hat{I}_{(0000)\tau_{p+1}, \tau_p}^{(i_4 i_3 i_2 i_1)}, \\
\mathbf{r}_{p+1, p} = & \sum_{i_1=1}^m \left(G_{i_1} L \mathbf{a} \left(\frac{1}{2} \hat{I}_{(2)\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \hat{I}_{(1)\tau_{p+1}, \tau_p}^{(i_1)} + \frac{\Delta^2}{2} \hat{I}_{(0)\tau_{p+1}, \tau_p}^{(i_1)} \right) + \right. \\
& \left. + \frac{1}{2} L L B_{i_1} \hat{I}_{(2)\tau_{p+1}, \tau_p}^{(i_1)} - L G_{i_1} \mathbf{a} \left(\hat{I}_{(2)\tau_{p+1}, \tau_p}^{(i_1)} + \Delta \hat{I}_{(1)\tau_{p+1}, \tau_p}^{(i_1)} \right) \right) + \\
& + \sum_{i_1, i_2, i_3=1}^m \left(G_{i_3} L G_{i_2} B_{i_1} \left(\hat{I}_{(100)\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} - \hat{I}_{(010)\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} \right) + \right. \\
& \left. + G_{i_3} G_{i_2} L B_{i_1} \left(\hat{I}_{(010)\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} - \hat{I}_{(001)\tau_{p+1}, \tau_p}^{(i_3 i_2 i_1)} \right) + \right.
\end{aligned}$$

$$\begin{aligned}
 & +G_{i_3}G_{i_2}G_{i_1}\left(\Delta\hat{I}_{(000)\tau_{p+1},\tau_p}^{(i_3i_2i_1)} + \hat{I}_{(001)\tau_{p+1},\tau_p}^{(i_3i_2i_1)}\right) - \\
 & \quad -LG_{i_3}G_{i_2}B_{i_1}\hat{I}_{(100)\tau_{p+1},\tau_p}^{(i_3i_2i_1)}\Big) + \\
 & + \sum_{i_1,i_2,i_3,i_4,i_5=1}^m G_{i_5}G_{i_4}G_{i_3}G_{i_2}B_{i_1}\hat{I}_{(00000)\tau_{p+1},\tau_p}^{(i_5i_4i_3i_2i_1)} + \\
 & \quad + \frac{\Delta^3}{6}LL\mathbf{a},
 \end{aligned}$$

where $\Delta = T/N$ ($N > 1$) is a constant (for simplicity) integration step, $\tau_p = p\Delta$ ($p = 0, 1, \dots, N$), $\hat{I}_{(l_1\dots l_k)s,t}^{(i_1\dots i_k)}$ denotes an approximation of the iterated Ito stochastic integral of multiplicity k

$$(6) \quad I_{(l_1\dots l_k)s,t}^{(i_1\dots i_k)} = \int_t^s (t - \tau_k)^{l_k} \dots \int_t^{\tau_2} (t - \tau_1)^{l_1} d\mathbf{f}_{\tau_1}^{(i_1)} \dots d\mathbf{f}_{\tau_k}^{(i_k)},$$

$$L = \frac{\partial}{\partial t} + \sum_{i=1}^n \mathbf{a}_i(\mathbf{x}, t) \frac{\partial}{\partial \mathbf{x}_i} + \frac{1}{2} \sum_{j=1}^m \sum_{l,i=1}^n B_{lj}(\mathbf{x}, t) B_{ij}(\mathbf{x}, t) \frac{\partial^2}{\partial \mathbf{x}_l \partial \mathbf{x}_i},$$

$$G_i = \sum_{j=1}^n B_{ji}(\mathbf{x}, t) \frac{\partial}{\partial \mathbf{x}_j}, \quad i = 1, \dots, m,$$

$l_1, \dots, l_k = 0, 1, 2$, $i_1, \dots, i_k = 1, \dots, m$, $k = 1, 2, \dots, 5$, B_i and B_{ij} are the i th column and the ij th element of the matrix function B , \mathbf{a}_i is the i th element of the vector function \mathbf{a} , \mathbf{x}_i is the i th element of the column \mathbf{x} , the functions

$$\begin{aligned}
 & B_{i_1}, \mathbf{a}, G_{i_2}B_{i_1}, G_{i_1}\mathbf{a}, LB_{i_1}, G_{i_3}G_{i_2}B_{i_1}, \mathbf{La}, LL\mathbf{a}, G_{i_2}LB_{i_1}, \\
 & LG_{i_2}B_{i_1}, G_{i_2}G_{i_1}\mathbf{a}, G_{i_4}G_{i_3}G_{i_2}B_{i_1}, G_{i_1}\mathbf{La}, LLB_{i_1}, LG_{i_1}\mathbf{a}, G_{i_3}LG_{i_2}B_{i_1}, G_{i_3}G_{i_2}LB_{i_1}, \\
 & G_{i_3}G_{i_2}G_{i_1}\mathbf{a}, LG_{i_3}G_{i_2}B_{i_1}, G_{i_5}G_{i_4}G_{i_3}G_{i_2}B_{i_1}
 \end{aligned}$$

are calculated at the point (\mathbf{y}_p, p) .

Under the standard conditions [3], [14] the numerical scheme (5) has order 2.5 of strong convergence. The major emphasis below will be placed on the approximation of the iterated Ito stochastic integrals appearing in (5). Therefore, among the standard conditions, note the approximation condition of these integrals [3], [14], which has the form

$$(7) \quad \mathbb{M} \left\{ \left(I_{(l_1\dots l_k)\tau_{p+1},\tau_p}^{(i_1\dots i_k)} - \hat{I}_{(l_1\dots l_k)\tau_{p+1},\tau_p}^{(i_1\dots i_k)} \right)^2 \right\} \leq C\Delta^6,$$

where constant C is independent of Δ .

Note that if we exclude $\mathbf{v}_{p+1,p} + \mathbf{r}_{p+1,p}$ from the right-hand side of (5), then we have an explicit one-step strong numerical scheme of order 1.5 [3], [14]-[18]. The right-hand side of (5) but without the value $\mathbf{r}_{p+1,p}$ define an explicit one-step strong numerical scheme of order 2.0 [3], [14]-[18].

Using the numerical scheme (5) or its modification based on the Taylor–Ito expansion [11], the implicit or multistep analogs of (5) can be constructed; see [3]. The set of the iterated Ito stochastic integrals to be approximated for implementing these modifications is the same as for the numerical scheme (5) itself. Interestingly, the truncated unified Taylor–Ito expansion — the foundation of the numerical scheme (5) — contains 12 different iterated Ito stochastic integrals of the form (6), which cannot be interconnected by linear relations [14]–[18]. The analogous Taylor–Ito expansion [3] contains 17 different iterated Ito stochastic integrals, part of which are interconnected by linear relations and part of which have a higher multiplicity than the iterated Ito stochastic integrals (6). This fact well explains the use of the numerical scheme (5).

One of the main problems arising in the implementation of the numerical scheme (5) is the joint numerical modeling of the iterated Ito stochastic integrals figuring in (5). In the next section, we will consider an efficient numerical modeling method for the iterated Ito stochastic integrals and also demonstrate which stochastic integrals (Ito or Stratonovich) are preferable for numerical modeling with a correct estimation of the mean-square approximation error.

3. METHOD OF NUMERICAL MODELING FOR ITERATED ITO STOCHASTIC INTEGRALS BASED ON GENERALIZED MULTIPLE FOURIER SERIES

An efficient numerical modeling method for the iterated Ito stochastic integrals based on generalized multiple Fourier series was considered in [14] (2006); also see [15]–[57]. This method rests on an important result presented below (Theorems 1, 2).

Suppose that $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$. Define the following function on the hypercube $[t, T]^k$

$$(8) \quad K(t_1, \dots, t_k) = \begin{cases} \psi_1(t_1) \dots \psi_k(t_k) & \text{for } t_1 < \dots < t_k \\ 0 & \text{otherwise} \end{cases}, \quad t_1, \dots, t_k \in [t, T], \quad k \geq 2,$$

and $K(t_1) \equiv \psi_1(t_1)$ for $t_1 \in [t, T]$.

Suppose that $\{\phi_j(x)\}_{j=0}^{\infty}$ is a complete orthonormal system of functions in the space $L_2([t, T])$. The function $K(t_1, \dots, t_k)$ belongs to the space $L_2([t, T]^k)$. At this situation it is well known that the generalized multiple Fourier series of $K(t_1, \dots, t_k) \in L_2([t, T]^k)$ is converging to $K(t_1, \dots, t_k)$ in the hypercube $[t, T]^k$ in the mean-square sense, i.e.

$$\lim_{p_1, \dots, p_k \rightarrow \infty} \left\| K(t_1, \dots, t_k) - \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \prod_{l=1}^k \phi_{j_l}(t_l) \right\|_{L_2([t, T]^k)} = 0,$$

where

$$(9) \quad C_{j_k \dots j_1} = \int_{[t, T]^k} K(t_1, \dots, t_k) \prod_{l=1}^k \phi_{j_l}(t_l) dt_1 \dots dt_k,$$

$$\|f\|_{L_2([t, T]^k)} = \left(\int_{[t, T]^k} f^2(t_1, \dots, t_k) dt_1 \dots dt_k \right)^{1/2}.$$

Consider the partition $\{\tau_j\}_{j=0}^N$ of $[t, T]$ such that

$$(10) \quad t = \tau_0 < \dots < \tau_N = T, \quad \Delta_N = \max_{0 \leq j \leq N-1} \Delta\tau_j \rightarrow 0 \text{ if } N \rightarrow \infty, \quad \Delta\tau_j = \tau_{j+1} - \tau_j.$$

Theorem 1 [14] (2006), [15]-[34], [39]-[49], [51]-[57]. *Suppose that every $\psi_l(\tau)$ ($l = 1, \dots, k$) is a continuous non-random function on $[t, T]$ and $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of continuous functions in the space $L_2([t, T])$. Then*

$$(11) \quad J[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_k \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} - \text{l.i.m.}_{N \rightarrow \infty} \sum_{(l_1, \dots, l_k) \in G_k} \phi_{j_1}(\tau_{l_1}) \Delta \mathbf{w}_{\tau_{l_1}}^{(i_1)} \dots \phi_{j_k}(\tau_{l_k}) \Delta \mathbf{w}_{\tau_{l_k}}^{(i_k)} \right),$$

where $J[\psi^{(k)}]_{T,t}$ is defined by (2),

$$G_k = H_k \setminus L_k, \quad H_k = \{(l_1, \dots, l_k) : l_1, \dots, l_k = 0, 1, \dots, N-1\},$$

$$L_k = \{(l_1, \dots, l_k) : l_1, \dots, l_k = 0, 1, \dots, N-1; l_g \neq l_r \ (g \neq r); g, r = 1, \dots, k\},$$

l.i.m. is a limit in the mean-square sense, $i_1, \dots, i_k = 0, 1, \dots, m$,

$$(12) \quad \zeta_j^{(i)} = \int_t^T \phi_j(s) d\mathbf{w}_s^{(i)}$$

are independent standard Gaussian random variables for various i or j (if $i \neq 0$), $C_{j_k \dots j_1}$ is the Fourier coefficient (9), $\Delta \mathbf{w}_{\tau_j}^{(i)} = \mathbf{w}_{\tau_{j+1}}^{(i)} - \mathbf{w}_{\tau_j}^{(i)}$ ($i = 0, 1, \dots, m$), $\{\tau_j\}_{j=0}^N$ is a partition of the interval $[t, T]$, which satisfies the condition (10).

The convergence in the mean of degree $2n$ ($n \in \mathbb{N}$) [19]-[21], [23]-[28] as well as the convergence with probability 1 [19]-[21], [41], [43] of approximations from Theorem 1 (also see Theorem 2 below) are proved. Moreover, the complete orthonormal systems of Haar and Rademacher–Walsh functions in $L_2([t, T])$ can also be applied in Theorems 1 [14]-[28]. The modification of Theorem 1 for complete orthonormal with weight $r(x) \geq 0$ systems of functions in the space $L_2([t, T])$ can be found in [18]-[21], [52]

In order to evaluate the significance of Theorem 1 for practice we will demonstrate its transformed particular cases for $k = 1, \dots, 5$ [14]-[34], [39]-[49], [51]-[57]

$$(13) \quad J[\psi^{(1)}]_{T,t} = \text{l.i.m.}_{p_1 \rightarrow \infty} \sum_{j_1=0}^{p_1} C_{j_1} \zeta_{j_1}^{(i_1)},$$

$$(14) \quad J[\psi^{(2)}]_{T,t} = \text{l.i.m.}_{p_1, p_2 \rightarrow \infty} \sum_{j_1=0}^{p_1} \sum_{j_2=0}^{p_2} C_{j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \right),$$

$$(15) \quad J[\psi^{(3)}]_{T,t} = \text{l.i.m.}_{p_1, p_2, p_3 \rightarrow \infty} \sum_{j_1=0}^{p_1} \sum_{j_2=0}^{p_2} \sum_{j_3=0}^{p_3} C_{j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \right. \\ \left. - \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

$$(16) \quad J[\psi^{(4)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_4 \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_4=0}^{p_4} C_{j_4 \dots j_1} \left(\prod_{l=1}^4 \zeta_{j_l}^{(i_l)} - \right. \\ - \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \\ - \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} - \\ - \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} + \\ + \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} + \\ + \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} + \\ \left. + \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \right),$$

$$(17) \quad J[\psi^{(5)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_5 \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_5=0}^{p_5} C_{j_5 \dots j_1} \left(\prod_{l=1}^5 \zeta_{j_l}^{(i_l)} - \right. \\ - \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \\ - \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{i_1=i_5 \neq 0\}} \mathbf{1}_{\{j_1=j_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \\ - \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} - \\ - \mathbf{1}_{\{i_2=i_5 \neq 0\}} \mathbf{1}_{\{j_2=j_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_5}^{(i_5)} - \\ - \mathbf{1}_{\{i_3=i_5 \neq 0\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_4=i_5 \neq 0\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} + \\ + \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_3=i_5 \neq 0\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_4}^{(i_4)} + \\ + \mathbf{1}_{\{i_1=i_2 \neq 0\}} \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_4=i_5 \neq 0\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_5}^{(i_5)} + \\ + \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_2=i_5 \neq 0\}} \mathbf{1}_{\{j_2=j_5\}} \zeta_{j_4}^{(i_4)} + \mathbf{1}_{\{i_1=i_3 \neq 0\}} \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_4=i_5 \neq 0\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_2}^{(i_2)} + \\ + \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_2=i_5 \neq 0\}} \mathbf{1}_{\{j_2=j_5\}} \zeta_{j_3}^{(i_3)} + \\ + \mathbf{1}_{\{i_1=i_4 \neq 0\}} \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_3=i_5 \neq 0\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_2}^{(i_2)} + \mathbf{1}_{\{i_1=i_5 \neq 0\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_4}^{(i_4)} + \\ + \mathbf{1}_{\{i_1=i_5 \neq 0\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{i_1=i_5 \neq 0\}} \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_2}^{(i_2)} + \\ + \mathbf{1}_{\{i_2=i_3 \neq 0\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_4=i_5 \neq 0\}} \mathbf{1}_{\{j_4=j_5\}} \zeta_{j_1}^{(i_1)} + \mathbf{1}_{\{i_2=i_4 \neq 0\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_3=i_5 \neq 0\}} \mathbf{1}_{\{j_3=j_5\}} \zeta_{j_1}^{(i_1)} + \\ \left. + \mathbf{1}_{\{i_2=i_5 \neq 0\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_3=i_4 \neq 0\}} \mathbf{1}_{\{j_3=j_4\}} \zeta_{j_1}^{(i_1)} \right),$$

where $\mathbf{1}_A$ is the indicator of the set A .

We will consider the case $i_1, \dots, i_5 = 1, \dots, m$. This case corresponds to the numerical method (5).

For further consideration, let us consider the generalization of formulas (13)–(17) for the case of an arbitrary multiplicity k ($k \in \mathbb{N}$) of the iterated Ito stochastic integral $J[\psi^{(k)}]_{T,t}$ defined by (2). In order to do this, let us introduce some notations. Consider the unordered set $\{1, 2, \dots, k\}$ and separate it into two parts: the first part consists of r unordered pairs (sequence order of these pairs is also unimportant) and the second one consists of the remaining $k - 2r$ numbers. So, we have

$$(18) \quad \left(\underbrace{\{\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}\}}_{\text{part 1}}, \underbrace{\{q_1, \dots, q_{k-2r}\}}_{\text{part 2}} \right),$$

where

$$\{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\},$$

braces mean an unordered set, and parentheses mean an ordered set.

We will say that (18) is a partition and consider the sum with respect to all possible partitions

$$(19) \quad \sum_{\substack{(\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}, \{q_1, \dots, q_{k-2r}\}) \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} a_{g_1 g_2, \dots, g_{2r-1} g_{2r}, q_1 \dots q_{k-2r}}.$$

Below there are several examples of sums in the form (19)

$$\begin{aligned} & \sum_{\substack{(\{g_1, g_2\}) \\ \{g_1, g_2\} = \{1, 2\}}} a_{g_1 g_2} = a_{12}, \\ & \sum_{\substack{(\{g_1, g_2\}, \{g_3, g_4\}) \\ \{g_1, g_2, g_3, g_4\} = \{1, 2, 3, 4\}}} a_{g_1 g_2 g_3 g_4} = a_{1234} + a_{1324} + a_{2314}, \\ & \sum_{\substack{(\{g_1, g_2\}, \{q_1, q_2\}) \\ \{g_1, g_2, q_1, q_2\} = \{1, 2, 3, 4\}}} a_{g_1 g_2, q_1 q_2} = \\ & = a_{12,34} + a_{13,24} + a_{14,23} + a_{23,14} + a_{24,13} + a_{34,12}, \\ & \sum_{\substack{(\{g_1, g_2\}, \{q_1, q_2, q_3\}) \\ \{g_1, g_2, q_1, q_2, q_3\} = \{1, 2, 3, 4, 5\}}} a_{g_1 g_2, q_1 q_2 q_3} = \\ & = a_{12,345} + a_{13,245} + a_{14,235} + a_{15,234} + a_{23,145} + a_{24,135} + \\ & + a_{25,134} + a_{34,125} + a_{35,124} + a_{45,123}, \\ & \sum_{\substack{(\{g_1, g_2\}, \{g_3, g_4\}, \{q_1\}) \\ \{g_1, g_2, g_3, g_4, q_1\} = \{1, 2, 3, 4, 5\}}} a_{g_1 g_2, g_3 g_4, q_1} = \\ & = a_{12,34,5} + a_{13,24,5} + a_{14,23,5} + a_{12,35,4} + a_{13,25,4} + a_{15,23,4} + \\ & + a_{12,54,3} + a_{15,24,3} + a_{14,25,3} + a_{15,34,2} + a_{13,54,2} + a_{14,53,2} \end{aligned}$$

$$+a_{52,34,1} + a_{53,24,1} + a_{54,23,1}.$$

Now we can write (11) as

$$(20) \quad J[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_k \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} + \sum_{r=1}^{[k/2]} (-1)^r \times \right. \\ \left. \times \sum_{\substack{(\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}), \{q_1, \dots, q_{k-2r}\} \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} \prod_{s=1}^r \mathbf{1}_{\{i_{g_{2s-1}} = i_{g_{2s}} \neq 0\}} \mathbf{1}_{\{j_{g_{2s-1}} = j_{g_{2s}}\}} \prod_{l=1}^{k-2r} \zeta_{j_{q_l}}^{(i_{q_l})} \right),$$

where $[x]$ is an integer part of a real number x ; another notations are the same as in Theorem 1.

In particular, from (20) for $k = 5$ we obtain

$$J[\psi^{(5)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_5 \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_5=0}^{p_5} C_{j_5 \dots j_1} \left(\prod_{l=1}^5 \zeta_{j_l}^{(i_l)} - \right. \\ \left. - \sum_{\substack{(\{g_1, g_2\}, \{q_1, q_2, q_3\}) \\ \{g_1, g_2, q_1, q_2, q_3\} = \{1, 2, 3, 4, 5\}}} \mathbf{1}_{\{i_{g_1} = i_{g_2} \neq 0\}} \mathbf{1}_{\{j_{g_1} = j_{g_2}\}} \prod_{l=1}^3 \zeta_{j_{q_l}}^{(i_{q_l})} + \right. \\ \left. + \sum_{\substack{(\{g_1, g_2\}, \{g_3, g_4\}, \{q_1\}) \\ \{g_1, g_2, g_3, g_4, q_1\} = \{1, 2, 3, 4, 5\}}} \mathbf{1}_{\{i_{g_1} = i_{g_2} \neq 0\}} \mathbf{1}_{\{j_{g_1} = j_{g_2}\}} \mathbf{1}_{\{i_{g_3} = i_{g_4} \neq 0\}} \mathbf{1}_{\{j_{g_3} = j_{g_4}\}} \zeta_{j_{q_1}}^{(i_{q_1})} \right).$$

The last equality obviously agrees with (17).

Let us consider the generalization of Theorem 1 for the case of an arbitrary complete orthonormal systems of functions in the space $L_2([t, T])$ and $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$.

Theorem 2 [19] (Sect. 1.11), [41] (Sect. 15). *Suppose that $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$ and $\{\phi_j(x)\}_{j=0}^\infty$ is an arbitrary complete orthonormal system of functions in the space $L_2([t, T])$. Then the following expansion*

$$(21) \quad J[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{p_1, \dots, p_k \rightarrow \infty} \sum_{j_1=0}^{p_1} \dots \sum_{j_k=0}^{p_k} C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} + \sum_{r=1}^{[k/2]} (-1)^r \times \right. \\ \left. \times \sum_{\substack{(\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}), \{q_1, \dots, q_{k-2r}\} \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} \prod_{s=1}^r \mathbf{1}_{\{i_{g_{2s-1}} = i_{g_{2s}} \neq 0\}} \mathbf{1}_{\{j_{g_{2s-1}} = j_{g_{2s}}\}} \prod_{l=1}^{k-2r} \zeta_{j_{q_l}}^{(i_{q_l})} \right)$$

converging in the mean-square sense is valid, where $[x]$ is an integer part of a real number x ; another notations are the same as in Theorem 1.

It should be noted that an analogue of Theorem 2 was considered in [62]. Note that we use another notations [19] (Sect. 1.11), [41] (Sect. 15) in comparison with [62]. Moreover, the proof of an analogue of Theorem 2 from [62] is somewhat different from the proof given in [19] (Sect. 1.11), [41] (Sect. 15).

4. CALCULATION OF THE MEAN-SQUARE APPROXIMATION ERROR IN THE METHOD OF GENERALIZED MULTIPLE FOURIER SEIRES

Note that for the integrals $J[\psi^{(k)}]_{T,t}$ defined by (2) the mean-square approximation error can be exactly calculated and efficiently estimated.

Let $J[\psi^{(k)}]_{T,t}^q$ be the expression on the right-hand side of (21) before passing to the limit $\text{l.i.m.}_{p_1, \dots, p_k \rightarrow \infty}$ for the case $p_1 = \dots = p_k = q$, i.e.

$$(22) \quad J[\psi^{(k)}]_{T,t}^q = \sum_{j_1, \dots, j_k=0}^q C_{j_k \dots j_1} \left(\prod_{l=1}^k \zeta_{j_l}^{(i_l)} + \sum_{r=1}^{\lfloor k/2 \rfloor} (-1)^r \times \right. \\ \left. \sum_{\substack{\{\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}\}, \{q_1, \dots, q_{k-2r}\} \\ \{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}}} \prod_{s=1}^r \mathbf{1}_{\{i_{g_{2s-1}} = i_{g_{2s}} \neq 0\}} \mathbf{1}_{\{j_{g_{2s-1}} = j_{g_{2s}}\}} \prod_{l=1}^{k-2r} \zeta_{j_{q_l}}^{(i_{q_l})} \right)$$

Let us denote

$$\mathbf{M} \left\{ \left(J[\psi^{(k)}]_{T,t} - J[\psi^{(k)}]_{T,t}^q \right)^2 \right\} \stackrel{\text{def}}{=} E_k^q, \\ \int_{[t, T]^k} K^2(t_1, \dots, t_k) dt_1 \dots dt_k \stackrel{\text{def}}{=} I_k.$$

In [17]-[21], [41], [42] it was shown that

$$(23) \quad E_k^q \leq k! \left(I_k - \sum_{j_1, \dots, j_k=0}^q C_{j_k \dots j_1}^2 \right)$$

for the following two cases:

1. $i_1, \dots, i_k = 1, \dots, m$ and $T - t \in (0, +\infty)$,
2. $i_1, \dots, i_k = 0, 1, \dots, m$ and $T - t \in (0, 1)$.

The value E_k^q can be calculated exactly.

Theorem 3 [19] (Sect. 1.12), [42] (Sect. 6). *Suppose that $\{\phi_j(x)\}_{j=0}^\infty$ is an arbitrary complete orthonormal system of functions in the space $L_2([t, T])$ and $\psi_1(\tau), \dots, \psi_k(\tau) \in L_2([t, T])$. Then*

$$(24) \quad E_k^q = I_k - \sum_{j_1, \dots, j_k=0}^q C_{j_k \dots j_1} \mathbf{M} \left\{ J[\psi^{(k)}]_{T,t} \sum_{(j_1, \dots, j_k)} \int_t^T \phi_{j_k}(t_k) \dots \int_t^{t_2} \phi_{j_1}(t_1) d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)} \right\},$$

where $i_1, \dots, i_k = 1, \dots, m$; the expression

$$\sum_{(j_1, \dots, j_k)}$$

means the sum with respect to all possible permutations (j_1, \dots, j_k) . At the same time if j_r swapped with j_q in the permutation (j_1, \dots, j_k) , then i_r swapped with i_q in the permutation (i_1, \dots, i_k) ; another notations are the same as in Theorems 1, 2.

Note that

$$\mathbb{M} \left\{ J[\psi^{(k)}]_{T,t} \int_t^T \phi_{j_k}(t_k) \dots \int_t^{t_2} \phi_{j_1}(t_1) d\mathbf{f}_{t_1}^{(i_1)} \dots d\mathbf{f}_{t_k}^{(i_k)} \right\} = C_{j_k \dots j_1}.$$

Therefore, for the special case of pairwise different numbers i_1, \dots, i_k as well as for the case $i_1 = \dots = i_k$ from Theorem 3 it follows that [18]-[21], [29], [42]

$$(25) \quad E_k^q = I_k - \sum_{j_1, \dots, j_k=0}^q C_{j_k \dots j_1}^2,$$

$$E_k^q = I_k - \sum_{j_1, \dots, j_k=0}^q C_{j_k \dots j_1} \left(\sum_{(j_1, \dots, j_k)} C_{j_k \dots j_1} \right),$$

where

$$\sum_{(j_1, \dots, j_k)}$$

is a sum with respect to all possible permutations (j_1, \dots, j_k) .

Consider some examples [18]-[21], [29], [42] of application of Theorem 3 ($i_1, i_2, i_3 = 1, \dots, m$)

$$(26) \quad E_2^q = I_2 - \sum_{j_1, j_2=0}^q C_{j_2 j_1}^2 - \sum_{j_1, j_2=0}^q C_{j_2 j_1} C_{j_1 j_2} \quad (i_1 = i_2),$$

$$(27) \quad E_3^q = I_3 - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_1 j_2} C_{j_3 j_2 j_1} \quad (i_1 = i_2 \neq i_3),$$

$$(28) \quad E_3^q = I_3 - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^q C_{j_2 j_3 j_1} C_{j_3 j_2 j_1} \quad (i_1 \neq i_2 = i_3),$$

$$(29) \quad E_3^q = I_3 - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1} C_{j_1 j_2 j_3} \quad (i_1 = i_3 \neq i_2).$$

The values E_4^q and E_5^q were calculated exactly for all possible combinations of $i_1, \dots, i_5 = 1, \dots, m$ in [18]-[21], [42].

5. EXPANSIONS OF ITERATED STRATONOVICH STOCHASTIC INTEGRALS BASED ON MULTIPLE FOURIER–LEGENDRE SERIES AND MULTIPLE TRIGONOMETRIC FOURIER SEIRES

In contrast to the iterated Ito stochastic integrals, the iterated Stratonovich stochastic integrals have simpler expansions than (11) and (21), but the calculation (or estimation) of the mean-square approximation errors for the latter is a much more difficult problem than for the former. Study this issue in detail.

Introduce the following iterated Stratonovich stochastic integrals

$$(30) \quad J^*[\psi^{(k)}]_{T,t} = \int_t^{*T} \psi_k(t_k) \dots \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_k}^{(i_k)},$$

where the notations are the same as in the formula (2) (in this paper, we use the definition of the Stratonovich stochastic integral from [3]).

Consider a slightly modified and extended theoretical result that adapts Theorems 1, 2 for the iterated Stratonovich stochastic integrals (30) of multiplicities 2 to 4 (some old results).

Theorem 4 [15]-[21], [26]-[28], [32], [40], [43]-[47], [53]. *Assume that $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in $L_2([t, T])$. In addition, assume that $\psi_2(s)$ is a continuously differentiable function on the interval $[t, T]$ and $\psi_1(s)$, $\psi_3(s)$ are twice continuously differentiable functions on the interval $[t, T]$. Then*

$$(31) \quad J^*[\psi^{(k)}]_{T,t} = \text{l.i.m.}_{q \rightarrow \infty} \sum_{j_1, \dots, j_k=0}^q C_{j_k \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_k}^{(i_k)},$$

where $k = 2, 3, 4$. At that $\psi_1(s), \dots, \psi_k(s) \equiv 1$ and $i_1, \dots, i_k = 0, 1, \dots, m$ in (31) for $k = 4$, while $i_1, \dots, i_k = 1, \dots, m$ in (31) for $k = 2, 3$; the other notations are the same as in Theorems 1, 2.

Recently, a new approach to the expansion and mean-square approximation of iterated Stratonovich stochastic integrals has been obtained [19] (Sect. 2.10–2.16), [44] (Sect. 7–13), [45] (Sect. 13–19), [53] (Sect. 5–11), [57] (Sect. 4–9). Let us formulate four theorems that were obtained using this approach.

Theorem 5 [19], [44], [45], [53], [57]. *Suppose that $\{\phi_j(x)\}_{j=0}^\infty$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. Furthermore, let $\psi_1(\tau)$, $\psi_2(\tau)$, $\psi_3(\tau)$ are continuously differentiable nonrandom functions on $[t, T]$. Then, for the iterated Stratonovich stochastic integral of third multiplicity*

$$J^*[\psi^{(3)}]_{T,t} = \int_t^{*T} \psi_3(t_3) \int_t^{*t_3} \psi_2(t_2) \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} d\mathbf{w}_{t_2}^{(i_2)} d\mathbf{w}_{t_3}^{(i_3)} \quad (i_1, i_2, i_3 = 0, 1, \dots, m)$$

the following relations

$$(32) \quad J^*[\psi^{(3)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)},$$

$$(33) \quad \mathbb{M} \left\{ \left(J^*[\psi^{(3)}]_{T,t} - \sum_{j_1, j_2, j_3=0}^p C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \right)^2 \right\} \leq \frac{C}{p}$$

are fulfilled, where $i_1, i_2, i_3 = 0, 1, \dots, m$ in (32) and $i_1, i_2, i_3 = 1, \dots, m$ in (33), constant C is independent of p ,

$$C_{j_3 j_2 j_1} = \int_t^T \psi_3(t_3) \phi_{j_3}(t_3) \int_t^{t_3} \psi_2(t_2) \phi_{j_2}(t_2) \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 dt_2 dt_3$$

and

$$\zeta_j^{(i)} = \int_t^T \phi_j(\tau) d\mathbf{f}_\tau^{(i)}$$

are independent standard Gaussian random variables for various i or j (in the case when $i \neq 0$); another notations are the same as in Theorems 1, 2.

Theorem 6 [19], [44], [45], [53], [57]. Let $\{\phi_j(x)\}_{j=0}^\infty$ be a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. Furthermore, let $\psi_1(\tau), \dots, \psi_4(\tau)$ be continuously differentiable nonrandom functions on $[t, T]$. Then, for the iterated Stratonovich stochastic integral of fourth multiplicity

$$(34) \quad J^*[\psi^{(4)}]_{T,t} = \int_t^{*T} \psi_4(t_4) \int_t^{*t_4} \psi_3(t_3) \int_t^{*t_3} \psi_2(t_2) \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} d\mathbf{w}_{t_2}^{(i_2)} d\mathbf{w}_{t_3}^{(i_3)} d\mathbf{w}_{t_4}^{(i_4)}$$

the following relations

$$(35) \quad J^*[\psi^{(4)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)},$$

$$(36) \quad \mathbb{M} \left\{ \left(J^*[\psi^{(4)}]_{T,t} - \sum_{j_1, j_2, j_3, j_4=0}^p C_{j_4 j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \right)^2 \right\} \leq \frac{C}{p^{1-\varepsilon}}$$

are fulfilled, where $i_1, \dots, i_4 = 0, 1, \dots, m$ in (34), (35) and $i_1, \dots, i_4 = 1, \dots, m$ in (36), constant C does not depend on p , ε is an arbitrary small positive real number for the case of complete orthonormal system of Legendre polynomials in the space $L_2([t, T])$ and $\varepsilon = 0$ for the case of complete orthonormal system of trigonometric functions in the space $L_2([t, T])$,

$$C_{j_4 j_3 j_2 j_1} = \int_t^T \psi_4(t_4) \phi_{j_4}(t_4) \int_t^{t_4} \psi_3(t_3) \phi_{j_3}(t_3) \int_t^{t_3} \psi_2(t_2) \phi_{j_2}(t_2) \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 dt_2 dt_3 dt_4;$$

another notations are the same as in Theorem 5.

Theorem 7 [19], [44], [45], [53], [57]. Assume that $\{\phi_j(x)\}_{j=0}^{\infty}$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$ and $\psi_1(\tau), \dots, \psi_5(\tau)$ are continuously differentiable nonrandom functions on $[t, T]$. Then, for the iterated Stratonovich stochastic integral of fifth multiplicity

$$(37) \quad J^*[\psi^{(5)}]_{T,t} = \int_t^{*T} \psi_5(t_5) \dots \int_t^{*t_2} \psi_1(t_1) d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_5}^{(i_5)}$$

the following relations

$$(38) \quad J^*[\psi^{(5)}]_{T,t} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_5=0}^p C_{j_5 \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_5}^{(i_5)},$$

$$(39) \quad \mathbb{M} \left\{ \left(J^*[\psi^{(5)}]_{T,t} - \sum_{j_1, \dots, j_5=0}^p C_{j_5 \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_5}^{(i_5)} \right)^2 \right\} \leq \frac{C}{p^{1-\varepsilon}}$$

are fulfilled, where $i_1, \dots, i_5 = 0, 1, \dots, m$ in (37), (38) and $i_1, \dots, i_5 = 1, \dots, m$ in (39), constant C is independent of p , ε is an arbitrary small positive real number for the case of complete orthonormal system of Legendre polynomials in the space $L_2([t, T])$ and $\varepsilon = 0$ for the case of complete orthonormal system of trigonometric functions in the space $L_2([t, T])$,

$$C_{j_5 \dots j_1} = \int_t^T \psi_5(t_5) \phi_{j_5}(t_5) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_5;$$

another notations are the same as in Theorems 5, 6.

Theorem 8 [19], [44], [45], [53]. Suppose that $\{\phi_j(x)\}_{j=0}^{\infty}$ is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space $L_2([t, T])$. Then, for the iterated Stratonovich stochastic integral of sixth multiplicity

$$(40) \quad J_{T,t}^{*(i_1 \dots i_6)} = \int_t^{*T} \dots \int_t^{*t_2} d\mathbf{w}_{t_1}^{(i_1)} \dots d\mathbf{w}_{t_6}^{(i_6)}$$

the following expansion

$$J_{T,t}^{*(i_1 \dots i_6)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_6=0}^p C_{j_6 \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_6}^{(i_6)}$$

that converges in the mean-square sense is valid, where $i_1, \dots, i_6 = 0, 1, \dots, m$,

$$C_{j_6 \dots j_1} = \int_t^T \phi_{j_6}(t_6) \dots \int_t^{t_2} \phi_{j_1}(t_1) dt_1 \dots dt_6;$$

another notations are the same as in Theorems 5–7.

Clearly, the expansion (31) is simpler than the expansions (11), (21). However, the calculation of the mean-square approximation error for the expansion (31) turns out to be much more difficult than for the expansions (11), (21). We will demonstrate this fact below.

The cases $k = 1, 2$ are actually not interesting: for $k = 1$, the Ito and Stratonovich stochastic integrals of a smooth nonrandom function equal each other with probability 1 (w. p. 1); for $k = 2$, the Ito stochastic integrals appearing in the numerical scheme (5) differ w. p. 1 from the corresponding Stratonovich stochastic integrals by constant values by virtue of the standard relations between the Ito and Stratonovich stochastic integrals [3]. Consider the triple Stratonovich stochastic integral defined by

$$I_{(000)T,t}^{*(i_1 i_2 i_3)} = \int_t^{*T} \int_t^{*t_3} \int_t^{*t_2} d\mathbf{f}_{t_1}^{(i_1)} d\mathbf{f}_{t_2}^{(i_2)} d\mathbf{f}_{t_3}^{(i_3)} \quad (i_1, i_2, i_3 = 1, \dots, m).$$

In view of the standard relations between the Ito and Stratonovich stochastic integrals [3] and also Theorems 1, 2, and 4 ($k = 3$), we obtain

$$\begin{aligned} & \mathbb{M} \left\{ \left(I_{(000)T,t}^{*(i_1 i_2 i_3)} - I_{(000)T,t}^{*(i_1 i_2 i_3)q} \right)^2 \right\} = \\ & = \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} + \mathbf{1}_{\{i_1=i_2\}} \frac{1}{2} \int_t^T \int_t^\tau ds d\mathbf{f}_\tau^{(i_3)} + \mathbf{1}_{\{i_2=i_3\}} \frac{1}{2} \int_t^T \int_t^\tau d\mathbf{f}_s^{(i_1)} d\tau - I_{(000)T,t}^{*(i_1 i_2 i_3)q} \right)^2 \right\} = \\ & = \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q} + I_{(000)T,t}^{(i_1 i_2 i_3)q} + \right. \right. \\ & \left. \left. + \mathbf{1}_{\{i_1=i_2\}} \frac{1}{2} \int_t^T \int_t^\tau ds d\mathbf{f}_\tau^{(i_3)} + \mathbf{1}_{\{i_2=i_3\}} \frac{1}{2} \int_t^T \int_t^\tau d\mathbf{f}_s^{(i_1)} d\tau - I_{(000)T,t}^{*(i_1 i_2 i_3)q} \right)^2 \right\}, \end{aligned} \quad (41)$$

$$\begin{aligned} I_{(000)T,t}^{(i_1 i_2 i_3)q} & = \sum_{j_1, j_2, j_3=0}^q C_{j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \right. \\ & \left. - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right), \end{aligned} \quad (42)$$

$$I_{(000)T,t}^{*(i_1 i_2 i_3)q} = \sum_{j_1, j_2, j_3=0}^q C_{j_3 j_2 j_1} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)}, \quad (43)$$

where $I_{(000)T,t}^{(i_1 i_2 i_3)q}$ is the approximation defined by (22) (also see (15)) for $k = 3$ and $I_{(000)T,t}^{*(i_1 i_2 i_3)q}$ is the approximation based on Theorem 4 for $k = 3$.

Substituting (42) and (43) into (41) yields

$$\begin{aligned}
 & \mathbb{M} \left\{ \left(I_{(000)T,t}^{*(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q} \right)^2 \right\} = \\
 & = \mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q} + \mathbf{1}_{\{i_1=i_2\}} \left(\frac{1}{2} \int_t^T \int_t^\tau ds d\mathbf{f}_\tau^{(i_3)} - \sum_{j_1, j_3=0}^q C_{j_3 j_1 j_1} \zeta_{j_3}^{(i_3)} \right) + \right. \right. \\
 & \left. \left. + \mathbf{1}_{\{i_2=i_3\}} \left(\frac{1}{2} \int_t^T \int_t^\tau d\mathbf{f}_s^{(i_1)} d\tau - \sum_{j_1, j_3=0}^q C_{j_3 j_3 j_1} \zeta_{j_1}^{(i_1)} \right) - \mathbf{1}_{\{i_1=i_3\}} \sum_{j_1, j_2=0}^q C_{j_1 j_2 j_1} \zeta_{j_2}^{(i_2)} \right)^2 \right\} \leq \\
 & \leq 4 \left(\mathbb{M} \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q} \right)^2 \right\} + \mathbf{1}_{\{i_1=i_2\}} F_q^{(i_3)} + \right. \\
 & \left. + \mathbf{1}_{\{i_2=i_3\}} G_q^{(i_1)} + \mathbf{1}_{\{i_1=i_3\}} H_q^{(i_2)} \right), \tag{44}
 \end{aligned}$$

where

$$\begin{aligned}
 F_q^{(i_3)} &= \mathbb{M} \left\{ \left(\frac{1}{2} \int_t^T \int_t^\tau ds d\mathbf{f}_\tau^{(i_3)} - \sum_{j_1, j_3=0}^q C_{j_3 j_1 j_1} \zeta_{j_3}^{(i_3)} \right)^2 \right\}, \\
 G_q^{(i_1)} &= \mathbb{M} \left\{ \left(\frac{1}{2} \int_t^T \int_t^\tau d\mathbf{f}_s^{(i_1)} d\tau - \sum_{j_1, j_3=0}^q C_{j_3 j_3 j_1} \zeta_{j_1}^{(i_1)} \right)^2 \right\}, \\
 H_q^{(i_2)} &= \mathbb{M} \left\{ \left(\sum_{j_1, j_2=0}^q C_{j_1 j_2 j_1} \zeta_{j_2}^{(i_2)} \right)^2 \right\}.
 \end{aligned}$$

For the case of Legendre polynomials or trigonometric functions we have the equalities (for details, see the proof of Theorem 4 for $k = 3$ in [15]-[21], [26]-[28], [45])

$$\lim_{q \rightarrow \infty} F_q^{(i_3)} = 0, \quad \lim_{q \rightarrow \infty} G_q^{(i_1)} = 0, \quad \lim_{q \rightarrow \infty} H_q^{(i_2)} = 0.$$

However, in accordance with (44) the value

$$\mathbb{M} \left\{ \left(I_{(000)T,t}^{*(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q} \right)^2 \right\}$$

with a finite q can be estimated by the sum of

$$(45) \quad 4M \left\{ \left(I_{(000)T,t}^{(i_1 i_2 i_3)} - I_{(000)T,t}^{(i_1 i_2 i_3)q} \right)^2 \right\},$$

and three additional terms of a rather complex structure. The value (45) can be calculated exactly using Theorem 3 or estimated using (23) for the case $k = 3$.

As is easily observed, this peculiarity will also apply to the iterated Stratonovich stochastic integrals of multiplicities 4 and 5, with the only difference that the number of additional terms like $F_q^{(i_3)}$, $G_q^{(i_1)}$, and $H_q^{(i_2)}$ will be considerably higher and their structure will be more complicated. Therefore, the payment for a relatively simple approximation of the iterated Stratonovich stochastic integrals (Theorems 4–8) in comparison with the iterated Ito stochastic integrals (Theorems 1, 2) is a much more difficult calculation or estimation procedure of their mean-square approximation errors (see Chapter 5 in [19] for detail). This well explains why the main emphasis of the paper is on the approximation of the iterated Ito stochastic integrals figuring in the numerical scheme (5). Their approximation involves Theorems 1, 2 for $k = 1, \dots, 5$ and also a complete orthonormal system of Legendre polynomials in the space $L_2([t, T])$. As was established in [19]–[21], [31], [39], the Legendre polynomials have a series of advantages over the trigonometric functions for the approximation of iterated stochastic integrals using Theorems 1, 2.

6. APPROXIMATION OF ITERATED ITO STOCHASTIC INTEGRALS BASED ON MULTIPLE FOURIER–LEGENDRE SERIES

Consider the approximations of the iterated Ito stochastic integrals that appear in the numerical scheme (5) using Theorems 1, 2 and the complete orthonormal system of Legendre polynomials in the space $L_2([\tau_p, \tau_{p+1}])$ ($\tau_p = p\Delta$, $N\Delta = T$, $p = 0, 1, \dots, N$) [14] (also see [15]–[51], [54]–[56])

$$(46) \quad I_{(0)\tau_{p+1}, \tau_p}^{(i_1)} = \sqrt{\Delta} \zeta_0^{(i_1)},$$

$$(47) \quad I_{(00)\tau_{p+1}, \tau_p}^{(i_1 i_2)q} = \frac{\Delta}{2} \left(\zeta_0^{(i_1)} \zeta_0^{(i_2)} + \sum_{i=1}^q \frac{1}{\sqrt{4i^2 - 1}} \left(\zeta_{i-1}^{(i_1)} \zeta_i^{(i_2)} - \zeta_i^{(i_1)} \zeta_{i-1}^{(i_2)} \right) - \mathbf{1}_{\{i_1=i_2\}} \right),$$

$$(48) \quad I_{(1)\tau_{p+1}, \tau_p}^{(i_1)} = -\frac{\Delta^{3/2}}{2} \left(\zeta_0^{(i_1)} + \frac{1}{\sqrt{3}} \zeta_1^{(i_1)} \right),$$

$$(49) \quad I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} = \sum_{j_1, j_2, j_3=0}^q C_{j_3 j_2 j_1} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

$$I_{(0000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4)q} = \sum_{j_1, j_2, j_3, j_4=0}^q C_{j_4 j_3 j_2 j_1} \left(\prod_{l=1}^4 \zeta_{j_l}^{(i_l)} - \right.$$

$$\begin{aligned}
 & -\mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}}\zeta_{j_3}^{(i_3)}\zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{i_1=i_3\}}\mathbf{1}_{\{j_1=j_3\}}\zeta_{j_2}^{(i_2)}\zeta_{j_4}^{(i_4)} - \\
 & -\mathbf{1}_{\{i_1=i_4\}}\mathbf{1}_{\{j_1=j_4\}}\zeta_{j_2}^{(i_2)}\zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_2=i_3\}}\mathbf{1}_{\{j_2=j_3\}}\zeta_{j_1}^{(i_1)}\zeta_{j_4}^{(i_4)} - \\
 & -\mathbf{1}_{\{i_2=i_4\}}\mathbf{1}_{\{j_2=j_4\}}\zeta_{j_1}^{(i_1)}\zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_3=i_4\}}\mathbf{1}_{\{j_3=j_4\}}\zeta_{j_1}^{(i_1)}\zeta_{j_2}^{(i_2)} + \\
 & \quad + \mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}}\mathbf{1}_{\{i_3=i_4\}}\mathbf{1}_{\{j_3=j_4\}} + \\
 & \quad + \mathbf{1}_{\{i_1=i_3\}}\mathbf{1}_{\{j_1=j_3\}}\mathbf{1}_{\{i_2=i_4\}}\mathbf{1}_{\{j_2=j_4\}} + \\
 & \quad + \mathbf{1}_{\{i_1=i_4\}}\mathbf{1}_{\{j_1=j_4\}}\mathbf{1}_{\{i_2=i_3\}}\mathbf{1}_{\{j_2=j_3\}} \Big),
 \end{aligned} \tag{50}$$

$$\begin{aligned}
 I_{(01)\tau_{p+1},\tau_p}^{(i_1 i_2)q} &= -\frac{\Delta}{2}I_{(00)\tau_{p+1},\tau_p}^{(i_1 i_2)q} - \frac{\Delta^2}{4}\left(\frac{1}{\sqrt{3}}\zeta_0^{(i_1)}\zeta_1^{(i_2)} + \right. \\
 & \left. + \sum_{i=0}^q \left(\frac{(i+2)\zeta_i^{(i_1)}\zeta_{i+2}^{(i_2)} - (i+1)\zeta_{i+2}^{(i_1)}\zeta_i^{(i_2)}}{\sqrt{(2i+1)(2i+5)}(2i+3)} - \frac{\zeta_i^{(i_1)}\zeta_i^{(i_2)}}{(2i-1)(2i+3)} \right) \right),
 \end{aligned} \tag{51}$$

$$\begin{aligned}
 I_{(10)\tau_{p+1},\tau_p}^{(i_1 i_2)q} &= -\frac{\Delta}{2}I_{(00)\tau_{p+1},\tau_p}^{(i_1 i_2)q} - \frac{\Delta^2}{4}\left(\frac{1}{\sqrt{3}}\zeta_0^{(i_2)}\zeta_1^{(i_1)} + \right. \\
 & \left. + \sum_{i=0}^q \left(\frac{(i+1)\zeta_{i+2}^{(i_2)}\zeta_i^{(i_1)} - (i+2)\zeta_i^{(i_2)}\zeta_{i+2}^{(i_1)}}{\sqrt{(2i+1)(2i+5)}(2i+3)} + \frac{\zeta_i^{(i_1)}\zeta_i^{(i_2)}}{(2i-1)(2i+3)} \right) \right)
 \end{aligned} \tag{52}$$

or

$$\begin{aligned}
 I_{(01)\tau_{p+1},\tau_p}^{(i_1 i_2)q} &= \sum_{j_1, j_2=0}^q C_{j_2 j_1}^{01} \left(\zeta_{j_1}^{(i_1)}\zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}} \right), \\
 I_{(10)\tau_{p+1},\tau_p}^{(i_1 i_2)q} &= \sum_{j_1, j_2=0}^q C_{j_2 j_1}^{10} \left(\zeta_{j_1}^{(i_1)}\zeta_{j_2}^{(i_2)} - \mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}} \right); \\
 I_{(2)\tau_{p+1},\tau_p}^{(i_1)} &= \frac{\Delta^{5/2}}{3} \left(\zeta_0^{(i_1)} + \frac{\sqrt{3}}{2}\zeta_1^{(i_1)} + \frac{1}{2\sqrt{5}}\zeta_2^{(i_1)} \right),
 \end{aligned} \tag{53}$$

$$\begin{aligned}
 I_{(001)\tau_{p+1},\tau_p}^{(i_1 i_2 i_3)q} &= \sum_{j_1, j_2, j_3=0}^q C_{j_3 j_2 j_1}^{001} \left(\zeta_{j_1}^{(i_1)}\zeta_{j_2}^{(i_2)}\zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}}\mathbf{1}_{\{j_1=j_2\}}\zeta_{j_3}^{(i_3)} - \right. \\
 & \left. - \mathbf{1}_{\{i_2=i_3\}}\mathbf{1}_{\{j_2=j_3\}}\zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}}\mathbf{1}_{\{j_1=j_3\}}\zeta_{j_2}^{(i_2)} \right),
 \end{aligned} \tag{54}$$

$$(55) \quad I_{(010)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} = \sum_{j_1, j_2, j_3=0}^q C_{j_3 j_2 j_1}^{010} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \right. \\ \left. - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

$$(56) \quad I_{(100)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} = \sum_{j_1, j_2, j_3=0}^q C_{j_3 j_2 j_1}^{100} \left(\zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} - \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_1=j_2\}} \zeta_{j_3}^{(i_3)} - \right. \\ \left. - \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_2=j_3\}} \zeta_{j_1}^{(i_1)} - \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_1=j_3\}} \zeta_{j_2}^{(i_2)} \right),$$

$$(57) \quad I_{(00000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4 i_5)q} = \sum_{j_1, j_2, j_3, j_4, j_5=0}^q C_{j_5 j_4 j_3 j_2 j_1} \left(\prod_{l=1}^5 \zeta_{j_l}^{(i_l)} - \right. \\ - \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \\ - \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \\ - \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_1}^{(i_1)} \zeta_{j_4}^{(i_4)} \zeta_{j_5}^{(i_5)} - \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_5}^{(i_5)} - \\ - \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_3}^{(i_3)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_5}^{(i_5)} - \\ - \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_4}^{(i_4)} - \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_1}^{(i_1)} \zeta_{j_2}^{(i_2)} \zeta_{j_3}^{(i_3)} + \\ + \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_4}^{(i_4)} + \\ + \mathbf{1}_{\{j_1=j_2\}} \mathbf{1}_{\{i_1=i_2\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_5}^{(i_5)} + \\ + \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_4}^{(i_4)} + \mathbf{1}_{\{j_1=j_3\}} \mathbf{1}_{\{i_1=i_3\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_2}^{(i_2)} + \\ + \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_5}^{(i_5)} + \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_2=j_5\}} \mathbf{1}_{\{i_2=i_5\}} \zeta_{j_3}^{(i_3)} + \\ + \mathbf{1}_{\{j_1=j_4\}} \mathbf{1}_{\{i_1=i_4\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_2}^{(i_2)} + \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \zeta_{j_4}^{(i_4)} + \\ + \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \zeta_{j_3}^{(i_3)} + \mathbf{1}_{\{j_1=j_5\}} \mathbf{1}_{\{i_1=i_5\}} \mathbf{1}_{\{j_3=j_4\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_2}^{(i_2)} + \\ + \mathbf{1}_{\{j_2=j_3\}} \mathbf{1}_{\{i_2=i_3\}} \mathbf{1}_{\{j_4=j_5\}} \mathbf{1}_{\{i_4=i_5\}} \zeta_{j_1}^{(i_1)} + \mathbf{1}_{\{j_2=j_4\}} \mathbf{1}_{\{i_2=i_4\}} \mathbf{1}_{\{j_3=j_5\}} \mathbf{1}_{\{i_3=i_5\}} \zeta_{j_1}^{(i_1)} + \\ \left. + \mathbf{1}_{\{j_2=j_5 \neq 0\}} \mathbf{1}_{\{i_2=i_5\}} \mathbf{1}_{\{j_3=j_4 \neq 0\}} \mathbf{1}_{\{i_3=i_4\}} \zeta_{j_1}^{(i_1)} \right),$$

where

$$C_{j_3 j_2 j_1} = \int_{\tau_p}^{\tau_{p+1}} \phi_{j_3}(z) \int_{\tau_p}^z \phi_{j_2}(y) \int_{\tau_p}^y \phi_{j_1}(x) dx dy dz = \\ = \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{8} \Delta^{3/2} \bar{C}_{j_3 j_2 j_1},$$

$$\begin{aligned}
 C_{j_4 j_3 j_2 j_1} &= \int_{\tau_p}^{\tau_{p+1}} \phi_{j_4}(u) \int_{\tau_p}^u \phi_{j_3}(z) \int_{\tau_p}^z \phi_{j_2}(y) \int_{\tau_p}^y \phi_{j_1}(x) dx dy dz du = \\
 &= \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)}}{16} \Delta^2 \bar{C}_{j_4 j_3 j_2 j_1},
 \end{aligned}$$

$$\begin{aligned}
 C_{j_2 j_1}^{01} &= \int_{\tau_p}^{\tau_{p+1}} (\tau_p - y) \phi_{j_2}(y) \int_{\tau_p}^y \phi_{j_1}(x) dx dy = \\
 &= \frac{\sqrt{(2j_1+1)(2j_2+1)}}{8} \Delta^2 \bar{C}_{j_2 j_1}^{01},
 \end{aligned}$$

$$\begin{aligned}
 C_{j_2 j_1}^{10} &= \int_{\tau_p}^{\tau_{p+1}} \phi_{j_2}(y) \int_{\tau_p}^y (\tau_p - x) \phi_{j_1}(x) dx dy = \\
 &= \frac{\sqrt{(2j_1+1)(2j_2+1)}}{8} \Delta^2 \bar{C}_{j_2 j_1}^{10},
 \end{aligned}$$

$$\begin{aligned}
 C_{j_3 j_2 j_1}^{001} &= \int_{\tau_p}^{\tau_{p+1}} (\tau_p - z) \phi_{j_3}(z) \int_{\tau_p}^z \phi_{j_2}(y) \int_{\tau_p}^y \phi_{j_1}(x) dx dy dz = \\
 &= \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{16} \Delta^{5/2} \bar{C}_{j_3 j_2 j_1}^{001},
 \end{aligned}$$

$$\begin{aligned}
 C_{j_3 j_2 j_1}^{010} &= \int_{\tau_p}^{\tau_{p+1}} \phi_{j_3}(z) \int_{\tau_p}^z (\tau_p - y) \phi_{j_2}(y) \int_{\tau_p}^y \phi_{j_1}(x) dx dy dz = \\
 &= \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{16} \Delta^{5/2} \bar{C}_{j_3 j_2 j_1}^{010},
 \end{aligned}$$

$$\begin{aligned}
 C_{j_3 j_2 j_1}^{100} &= \int_{\tau_p}^{\tau_{p+1}} \phi_{j_3}(z) \int_{\tau_p}^z \phi_{j_2}(y) \int_{\tau_p}^y (\tau_p - x) \phi_{j_1}(x) dx dy dz = \\
 &= \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)}}{16} \Delta^{5/2} \bar{C}_{j_3 j_2 j_1}^{100},
 \end{aligned}$$

$$\begin{aligned}
 C_{j_5 j_4 j_3 j_2 j_1} &= \int_{\tau_p}^{\tau_{p+1}} \phi_{j_5}(v) \int_{\tau_p}^v \phi_{j_4}(u) \int_{\tau_p}^u \phi_{j_3}(z) \int_{\tau_p}^z \phi_{j_2}(y) \int_{\tau_p}^y \phi_{j_1}(x) dx dy dz du dv = \\
 &= \frac{\sqrt{(2j_1+1)(2j_2+1)(2j_3+1)(2j_4+1)(2j_5+1)}}{32} \Delta^{5/2} \bar{C}_{j_5 j_4 j_3 j_2 j_1},
 \end{aligned}$$

where

$$\begin{aligned}
\bar{C}_{j_3 j_2 j_1} &= \int_{-1}^1 P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz, \\
\bar{C}_{j_4 j_3 j_2 j_1} &= \int_{-1}^1 P_{j_4}(u) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz, \\
\bar{C}_{j_2 j_1}^{01} &= - \int_{-1}^1 (1+y) P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy, \\
\bar{C}_{j_2 j_1}^{10} &= - \int_{-1}^1 P_{j_2}(y) \int_{-1}^y (1+x) P_{j_1}(x) dx dy, \\
\bar{C}_{j_3 j_2 j_1}^{100} &= - \int_{-1}^1 P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) (x+1) dx dy dz, \\
\bar{C}_{j_3 j_2 j_1}^{010} &= - \int_{-1}^1 P_{j_3}(z) \int_{-1}^z P_{j_2}(y) (y+1) \int_{-1}^y P_{j_1}(x) dx dy dz, \\
\bar{C}_{j_3 j_2 j_1}^{001} &= - \int_{-1}^1 P_{j_3}(z) (z+1) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz, \\
\bar{C}_{j_5 j_4 j_3 j_2 j_1} &= \int_{-1}^1 P_{j_5}(v) \int_{-1}^v P_{j_4}(u) \int_{-1}^u P_{j_3}(z) \int_{-1}^z P_{j_2}(y) \int_{-1}^y P_{j_1}(x) dx dy dz dudv,
\end{aligned}$$

where $P_i(x)$ ($i = 0, 1, 2, \dots$) is the Legendre polynomial and

$$\phi_i(x) = \sqrt{\frac{2i+1}{\Delta}} P_i \left(\left(x - \tau_p - \frac{\Delta}{2} \right) \frac{2}{\Delta} \right), \quad i = 0, 1, 2, \dots$$

Let us consider the exact relations and some estimates for the mean-square approximation errors of iterated Ito stochastic integrals.

Using Theorem 3, we get [16]-[28], [43], [51]

$$(58) \quad \mathbb{M} \left\{ \left(I_{(00)\tau_{p+1}, \tau_p}^{(i_1 i_2)} - I_{(00)\tau_{p+1}, \tau_p}^{(i_1 i_2)q} \right)^2 \right\} = \frac{\Delta^2}{2} \left(\frac{1}{2} - \sum_{i=1}^q \frac{1}{4i^2 - 1} \right) \quad (i_1 \neq i_2),$$

$$\mathbb{M} \left\{ \left(I_{(10)\tau_{p+1}, \tau_p}^{(i_1 i_2)} - I_{(10)\tau_{p+1}, \tau_p}^{(i_1 i_2)q} \right)^2 \right\} = \mathbb{M} \left\{ \left(I_{(01)\tau_{p+1}, \tau_p}^{(i_1 i_2)} - I_{(01)\tau_{p+1}, \tau_p}^{(i_1 i_2)q} \right)^2 \right\} =$$

$$(59) = \frac{\Delta^4}{16} \left(\frac{5}{9} - 2 \sum_{i=2}^q \frac{1}{4i^2 - 1} - \sum_{i=1}^q \frac{1}{(2i-1)^2(2i+3)^2} - \sum_{i=0}^q \frac{(i+2)^2 + (i+1)^2}{(2i+1)(2i+5)(2i+3)^2} \right) \quad (i_1 \neq i_2),$$

$$(60) \quad \begin{aligned} & \mathbb{M} \left\{ \left(I_{(10)\tau_{p+1}, \tau_p}^{(i_1 i_1)} - I_{(10)\tau_{p+1}, \tau_p}^{(i_1 i_1)q} \right)^2 \right\} = \mathbb{M} \left\{ \left(I_{(01)\tau_{p+1}, \tau_p}^{(i_1 i_1)} - I_{(01)\tau_{p+1}, \tau_p}^{(i_1 i_1)q} \right)^2 \right\} = \\ & = \frac{\Delta^4}{16} \left(\frac{1}{9} - \sum_{i=0}^q \frac{1}{(2i+1)(2i+5)(2i+3)^2} - 2 \sum_{i=1}^q \frac{1}{(2i-1)^2(2i+3)^2} \right). \end{aligned}$$

Applying (25)–(29), we obtain

$$\begin{aligned} & \mathbb{M} \left\{ \left(I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} = \frac{\Delta^3}{6} - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 \quad (i_1 \neq i_2, i_1 \neq i_3, i_2 \neq i_3), \\ & \mathbb{M} \left\{ \left(I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} = \frac{\Delta^3}{6} - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^q C_{j_2 j_3 j_1} C_{j_3 j_2 j_1} \quad (i_1 \neq i_2 = i_3), \\ & \mathbb{M} \left\{ \left(I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} = \frac{\Delta^3}{6} - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1} C_{j_1 j_2 j_3} \quad (i_1 = i_3 \neq i_2), \\ & \mathbb{M} \left\{ \left(I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} = \frac{\Delta^3}{6} - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_1 j_2} C_{j_3 j_2 j_1} \quad (i_1 = i_2 \neq i_3). \end{aligned}$$

At the same time using the estimate (23) for $i_1, \dots, i_5 = 1, \dots, m$, we have

$$\begin{aligned} & \mathbb{M} \left\{ \left(I_{(01)\tau_{p+1}, \tau_p}^{(i_1 i_2)} - I_{(01)\tau_{p+1}, \tau_p}^{(i_1 i_2)q} \right)^2 \right\} \leq 2 \left(\frac{\Delta^4}{4} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{01})^2 \right), \\ & \mathbb{M} \left\{ \left(I_{(10)\tau_{p+1}, \tau_p}^{(i_1 i_2)} - I_{(10)\tau_{p+1}, \tau_p}^{(i_1 i_2)q} \right)^2 \right\} \leq 2 \left(\frac{\Delta^4}{12} - \sum_{j_1, j_2=0}^q (C_{j_2 j_1}^{10})^2 \right), \\ (61) \quad & \mathbb{M} \left\{ \left(I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} \leq 6 \left(\frac{\Delta^3}{6} - \sum_{j_3, j_2, j_1=0}^q C_{j_3 j_2 j_1}^2 \right), \end{aligned}$$

$$(62) \quad \mathbb{M} \left\{ \left(I_{(0000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4)} - I_{(0000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4)q} \right)^2 \right\} \leq 24 \left(\frac{\Delta^4}{24} - \sum_{j_1, j_2, j_3, j_4=0}^q C_{j_4 j_3 j_2 j_1}^2 \right),$$

$$(63) \quad \mathbb{M} \left\{ \left(I_{(100)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(100)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} \leq 6 \left(\frac{\Delta^5}{60} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{100})^2 \right),$$

$$(64) \quad \mathbb{M} \left\{ \left(I_{(010)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(010)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} \leq 6 \left(\frac{\Delta^5}{20} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{010})^2 \right),$$

$$(65) \quad \mathbb{M} \left\{ \left(I_{(001)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(001)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)q} \right)^2 \right\} \leq 6 \left(\frac{\Delta^5}{10} - \sum_{j_1, j_2, j_3=0}^q (C_{j_3 j_2 j_1}^{001})^2 \right),$$

$$(66) \quad \mathbb{M} \left\{ \left(I_{(00000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4 i_5)} - I_{(00000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4 i_5)q} \right)^2 \right\} \leq 120 \left(\frac{\Delta^5}{120} - \sum_{j_1, j_2, j_3, j_4, j_5=0}^q C_{j_5 i_4 i_3 i_2 j_1}^2 \right).$$

As was emphasized in [1], [2], [19]-[21], the Fourier–Legendre coefficients $\bar{C}_{j_3 j_2 j_1}$ and $\bar{C}_{j_4 j_3 j_2 j_1}$ (as well as the Fourier–Legendre coefficients $\bar{C}_{j_3 j_2 j_1}^{001}$, $\bar{C}_{j_3 j_2 j_1}^{010}$, $\bar{C}_{j_3 j_2 j_1}^{100}$, $\bar{C}_{j_5 j_4 j_3 j_2 j_1}$) can be exactly calculated using symbolic transformation packages like Derive. The exact values of these Fourier–Legendre coefficients calculated in Derive were presented in tabular form in the monographs [14]-[28]. Note that the mentioned Fourier–Legendre coefficients do not depend on the integration step $\tau_{p+1} - \tau_p$ of the numerical method, which can be variable.

Recently, the database with 270,000 exactly calculated Fourier–Legendre coefficients was described [54]. This database was used in the software package, which is written in the Python programming language for the implementation of explicit one-step strong numerical methods with orders 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 of convergence for Ito SDEs with non-commutative noise. The optimization of the mean-square approximation procedures for iterated Ito stochastic integrals from these numerical schemes can be found in [56].

Generally speaking, the minimum values q that guarantee the fulfillment of the condition (7) for each of approximations (see above) are different and abruptly decreasing with the growth of orders of smallness with respect to Δ of approximations of iterated Ito stochastic integrals.

For pairwise different $i_1, \dots, i_5 = 1, \dots, m$ Theorem 3 gives

$$(67) \quad \mathbb{M} \left\{ \left(I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)6} \right)^2 \right\} = \frac{\Delta^3}{6} - \sum_{j_1, j_2, j_3=0}^6 C_{j_3 j_2 j_1}^2 \approx 0.01956000\Delta^3,$$

$$(68) \quad \mathbb{M} \left\{ \left(I_{(100)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(100)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)2} \right)^2 \right\} = \frac{\Delta^5}{60} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{100})^2 \approx 0.00815429\Delta^5,$$

$$(69) \quad \mathbb{M} \left\{ \left(I_{(010)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(010)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)2} \right)^2 \right\} = \frac{\Delta^5}{20} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{010})^2 \approx 0.01739030\Delta^5,$$

$$(70) \quad \mathbb{M} \left\{ \left(I_{(001)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)} - I_{(001)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)2} \right)^2 \right\} = \frac{\Delta^5}{10} - \sum_{j_1, j_2, j_3=0}^2 (C_{j_3 j_2 j_1}^{001})^2 \approx 0.02528010\Delta^5,$$

$$(71) \quad \mathbb{M} \left\{ \left(I_{(0000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4)} - I_{(0000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4)2} \right)^2 \right\} = \frac{\Delta^4}{24} - \sum_{j_1, j_2, j_3, j_4=0}^2 C_{j_4 j_3 j_2 j_1}^2 \approx 0.02360840 \Delta^4,$$

$$(72) \quad \mathbb{M} \left\{ \left(I_{(00000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4 i_5)} - I_{(00000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4 i_5)1} \right)^2 \right\} = \frac{\Delta^5}{120} - \sum_{j_1, j_2, j_3, j_4, j_5=0}^1 C_{j_5 j_4 j_3 j_2 j_1}^2 \approx 0.00759105 \Delta^5.$$

Recall that the value Δ acts as the integration step of the numerical method (5) for the Ito SDE (1), thereby being rather small. Hence, even for $q = 6, 2$, and 1 the mean-square approximation errors (67)–(72) of the iterated Ito stochastic integrals of multiplicities 3 to 5 are sufficiently small. Note that in [3], [7], [8] the iterated stochastic integrals were approximated using the trigonometric Fourier expansion of the multidimensional Brownian bridge process and the mean-square approximation error of the iterated stochastic integrals were estimated by the value

$$\frac{C_1 \Delta^2}{q},$$

where C_1 is a constant and Δ, q have the same meaning as in (47). Clearly, such an approach is rougher than the one involving Theorem 3.

Note that the number q must be the same for all approximations of iterated stochastic integrals from the considered collection in the approach from [3], [7], [8] while the numbers q can be chosen different for different stochastic integrals from the considered collection in the method based on Theorems 1–3.

On the basis of the presented expansions (see above) of iterated Ito stochastic integrals we can see that increasing of multiplicities of these integrals or degree indexes of their weight functions leads to increasing of orders of smallness with respect to Δ in the mean-square sense for iterated stochastic integrals. This leads to a sharp decrease of member quantities (the numbers q) in expansions of iterated Ito stochastic integrals, which are required for achieving the acceptable accuracy of approximation.

7. NUMERICAL ALGORITHM WITH THE ORDER 2.5 OF STRONG CONVERGENCE

In this section, we will write the formulas and recommendations on the numerical method (5) with the order 2.5 of strong convergence as an algorithm.

Let the Fourier–Legendre coefficients

$$\bar{C}_{j_3 j_2 j_1}, \quad \bar{C}_{j_4 j_3 j_2 j_1}, \quad \bar{C}_{j_3 j_2 j_1}^{001}, \quad \bar{C}_{j_3 j_2 j_1}^{010}, \quad \bar{C}_{j_3 j_2 j_1}^{100}, \quad \bar{C}_{j_5 j_4 j_3 j_2 j_1}$$

be precalculated [54].

Algorithm 1.

Step 1. Specify the initial parameters of the problem: the integration interval $[0, T]$, the integration step Δ (e.g., the constant one $\Delta = T/N$, where $N > 1$; a variable step is also admissible), the initial condition \mathbf{y}_0 , and the constant C appearing in the condition (7).

Step 2. Let $p = 0$.

Step 3. Choose the minimum values q under which the right-hand sides of (58)–(66) are not exceeding the right-hand side of the inequality (7).

Step 4. Generate a sequence of independent standard Gaussian random variables $\zeta_l^{(i)}$ ($l = 0, 1, \dots, q + 2$; $i = 1, \dots, m$). Here the number q is a maximum from the numbers q chosen at Step 3.

Step 5. Model the iterated Ito stochastic integrals

$$I_{(0)\tau_{p+1}, \tau_p}^{(i_1)}, I_{(1)\tau_{p+1}, \tau_p}^{(i_1)}, I_{(2)\tau_{p+1}, \tau_p}^{(i_1)}, I_{(00)\tau_{p+1}, \tau_p}^{(i_1 i_2)}, I_{(10)\tau_{p+1}, \tau_p}^{(i_1 i_2)}, I_{(01)\tau_{p+1}, \tau_p}^{(i_1 i_2)}, I_{(000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)},$$

$$I_{(100)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)}, I_{(010)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)}, I_{(001)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3)}, I_{(0000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4)}, I_{(00000)\tau_{p+1}, \tau_p}^{(i_1 i_2 i_3 i_4 i_5)}$$

using the formulas (46)–(57) with the values q chosen at Step 3.

Step 6. Find \mathbf{y}_{p+1} by the formula (5).

Step 7. If $p < N - 1$, then assign $p = p + 1$ and go back to Step 4. Otherwise proceed to Step 8.

Step 8. End of the Algorithm 1.

8. CONCLUSIONS

In this paper, the efficient mean-square approximation procedures for the iterated Ito stochastic integrals of multiplicities 1 to 5 that are based on the multiple Fourier–Legendre series have been developed. These results can be used for the implementation of the numerical method (5) with the strong order 2.5 of convergence for the Ito SDEs with multidimensional non-commutative noises.

REFERENCES

- [1] Kuznetsov D.F. On numerical modeling of the multidimensional dynamic systems under random perturbations with the 1.5 and 2.0 orders of strong convergence [In English]. Automation and Remote Control, 79, 7 (2018), 1240-1254. DOI: <http://doi.org/10.1134/S0005117918070056>
- [2] Kuznetsov D.F. To numerical modeling with strong orders 1.0, 1.5, and 2.0 of convergence for multidimensional dynamical systems with random disturbances. arXiv:1802.00888 [math.PR]. 2018, 28 pp. [In English].
- [3] Kloeden P.E., Platen E. Numerical solution of stochastic differential equations. Berlin: Springer, 1992, 632 pp.
- [4] Arato M. Linear stochastic systems with constant coefficients. A statistical approach. Berlin–Heidelberg–N.Y.: Springer, 1982, 289 pp.
- [5] Shiriaev A.N. Foundations of Financial Mathematics. Vol. 2, Moscow: Fazis, 1998, 544 pp. [In Russian]
- [6] Liptser R.Sh., Shiriaev A.N. Statistics of Stochastic Processes: Nonlinear Filtering and Related Problems. Moscow: Nauka, 1974, 696 pp. [In Russian]
- [7] Milstein G.N. Numerical Integration of Stochastic Differential Equations. Sverdlovsk: Ural University Press, 1988, 225 pp. [In Russian]
- [8] Milstein G.N., Tretyakov M.V. Stochastic numerics for mathematical physics. Berlin: Springer, 2004, 616 pp.
- [9] Kloeden P.E., Platen E., Schurz H. Numerical solution of SDE through computer experiments. Berlin: Springer, 1994, 292 pp.
- [10] Platen E., Wagner W. On a Taylor formula for a class of Ito processes. Probab. Math. Statist. 3 (1982), 37-51.
- [11] Kloeden P.E., Platen E. The Stratonovich and Ito-Taylor Expansions. Math. Nachr. 151 (1991), 33-50.
- [12] Kulchitskiy O.Yu., Kuznetsov D.F. The unified Taylor-Ito expansion. Journal of Mathematical Sciences (N. Y.). 99, 2 (2000), 1130-1140. DOI: <http://doi.org/10.1007/BF02673635>
- [13] Kuznetsov D.F. Four new forms of the Taylor-Ito and Taylor-Stratonovich expansions and its application to the high-order strong numerical methods for Ito stochastic differential equations. arXiv:2001.10192 [math.PR], 2020, 90 pp. [In English].

- [14] Kuznetsov D.F. Numerical Integration of Stochastic Differential Equations. 2. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2006, 764 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-227> Available at: <http://www.sde-kuznetsov.spb.ru/06.pdf> (ISBN 5-7422-1191-0)
- [15] Kuznetsov D.F. Strong Approximation of Multiple Ito and Stratonovich Stochastic Integrals: Multiple Fourier Series Approach. 2nd Edition. [In English]. Polytechnical University Publishing House, Saint-Petersburg, 2011, 284 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-233> Available at: <http://www.sde-kuznetsov.spb.ru/11a.pdf> (ISBN 978-5-7422-3162-2)
- [16] Kuznetsov D.F. Multiple Ito and Stratonovich Stochastic Integrals: Fourier-Legendre and Trigonometric Expansions, Approximations, Formulas. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 2017, no. 1, 385 pp. (A.1–A.385). DOI: <http://doi.org/10.18720/SPBPU/2/z17-3> Available at: <http://diffjournal.spbu.ru/EN/numbers/2017.1/article.2.1.html>
- [17] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With Programs on MATLAB, 5th Edition. [In Russian]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 2 (2017), A.1-A.1000. DOI: <http://doi.org/10.18720/SPBPU/2/z17-4> Available at: <http://diffjournal.spbu.ru/EN/numbers/2017.2/article.2.1.html>
- [18] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With MATLAB Programs, 6th Edition. [In Russian]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 4 (2018), A.1-A.1073. Available at: <http://diffjournal.spbu.ru/EN/numbers/2018.4/article.2.1.html>
- [19] Kuznetsov D.F. Strong approximation of iterated Ito and Stratonovich stochastic integrals based on generalized multiple Fourier series. Application to numerical solution of Ito SDEs and semilinear SPDEs. arXiv:2003.14184 [math.PR], 2022, 923 pp. [In English].
- [20] Kuznetsov D.F. Strong Approximation of Iterated Ito and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series. Application to Numerical Solution of Ito SDEs and Semilinear SPDEs. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 4 (2020), A.1-A.606. Available at: <http://diffjournal.spbu.ru/EN/collection.html>
- [21] Kuznetsov D.F. Mean-Square Approximation of Iterated Itô and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series. Application to Numerical Integration of Itô SDEs and Semilinear SPDEs. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 4 (2021), A.1-A.788. Available at: <http://diffjournal.spbu.ru/EN/numbers/2021.4/article.1.9.html>
- [22] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With MatLab programs, 1st Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2007, 778 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-228> Available at: <http://www.sde-kuznetsov.spb.ru/07b.pdf> (ISBN 5-7422-1394-8)
- [23] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With MatLab programs, 2nd Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2007, XXXII+770 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-229> Available at: <http://www.sde-kuznetsov.spb.ru/07a.pdf> (ISBN 5-7422-1439-1)
- [24] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With MatLab programs, 3rd Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2009, XXXIV+768 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-230> Available at: <http://www.sde-kuznetsov.spb.ru/09.pdf> (ISBN 978-5-7422-2132-6)
- [25] Kuznetsov D.F. Stochastic Differential Equations: Theory and Practice of Numerical Solution. With MatLab programs. 4th Edition. [In Russian]. Polytechnical University Publishing House, Saint-Petersburg, 2010, XXX+786 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-231> Available at: <http://www.sde-kuznetsov.spb.ru/10.pdf> (ISBN 978-5-7422-2448-8)
- [26] Kuznetsov D.F. Multiple Stochastic Ito and Stratonovich Integrals and Multiple Fourier Series. [In Russian]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 3 (2010), A.1-A.257. DOI: <http://doi.org/10.18720/SPBPU/2/z17-7> Available at: <http://diffjournal.spbu.ru/EN/numbers/2010.3/article.2.1.html>
- [27] Kuznetsov D.F. Strong Approximation of Multiple Ito and Stratonovich Stochastic Integrals: Multiple Fourier Series Approach. 1st Edition. [In English]. Polytechnical University Publishing House, Saint-Petersburg, 2011, 250 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-232> Available at: <http://www.sde-kuznetsov.spb.ru/11b.pdf> (ISBN 978-5-7422-2988-9)
- [28] Kuznetsov D.F. Multiple Ito and Stratonovich Stochastic Integrals: Approximations, Properties, Formulas. [In English]. Polytechnical University Publishing House, Saint-Petersburg, 2013, 382 pp. DOI: <http://doi.org/10.18720/SPBPU/2/s17-234> Available at: <http://www.sde-kuznetsov.spb.ru/13.pdf> (ISBN 978-5-7422-3973-4)

- [29] Kuznetsov D.F. Development and application of the Fourier method for the numerical solution of Ito stochastic differential equations. [In English]. *Computational Mathematics and Mathematical Physics*, 58, 7 (2018), 1058-1070. DOI: <http://doi.org/10.1134/S0965542518070096>
- [30] Kuznetsov D.F. On numerical modeling of the multidimensional dynamic systems under random perturbations with the 2.5 order of strong convergence. [In English]. *Automation and Remote Control*, 80, 5 (2019), 867-881. DOI: <http://doi.org/10.1134/S0005117919050060>
- [31] Kuznetsov D.F. Comparative analysis of the efficiency of application of Legendre polynomials and trigonometric functions to the numerical integration of Itô stochastic differential equations. [In English]. *Computational Mathematics and Mathematical Physics*, 59, 8 (2019), 1236-1250. DOI: <http://doi.org/10.1134/S0965542519080116>
- [32] Kuznetsov D.F. Expansion of multiple Stratonovich stochastic integrals of second multiplicity based on double Fourier-Legendre series summarized by Prinsheim method [In Russian]. *Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online)*, 1 (2018), 1-34. Available at: <http://diffjournal.spbu.ru/EN/numbers/2018.1/article.1.1.html>
- [33] Kuznetsov D.F. Application of the method of approximation of iterated Ito stochastic integrals based on generalized multiple Fourier series to the high-order strong numerical methods for non-commutative semilinear stochastic partial differential equations. arXiv:1905.03724 [math.GM], 2019, 41 pp. [In English].
- [34] Kuznetsov D.F. Application of the method of approximation of iterated stochastic Ito integrals based on generalized multiple Fourier series to the high-order strong numerical methods for non-commutative semilinear stochastic partial differential equations. *Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online)*, 3 (2019), 18-62. Available at: <http://diffjournal.spbu.ru/EN/numbers/2019.3/article.1.2.html>
- [35] Kuznetsov D.F. A method of expansion and approximation of repeated stochastic Stratonovich integrals based on multiple Fourier series on full orthonormal systems. [In Russian]. *Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online)*, 1 (1997), 18-77. Available at: <http://diffjournal.spbu.ru/EN/numbers/1997.1/article.1.2.html>
- [36] Kuznetsov D.F. Problems of the numerical analysis of Ito stochastic differential equations. [In Russian]. *Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online)*, 1 (1998), 66-367. Available at: <http://diffjournal.spbu.ru/EN/numbers/1998.1/article.1.3.html> Hard Cover Edition: 1998, SPbGTU Publishing House, 204 pp. (ISBN 5-7422-0045-5)
- [37] Kuznetsov D.F. Mean square approximation of solutions of stochastic differential equations using Legendre polynomials. [In English]. *Journal of Automation and Information Sciences (Begell House)*, 2000, 32 (Issue 12), 69-86. DOI: <http://doi.org/10.1615/JAutomatInfScien.v32.i12.80>
- [38] Kuznetsov D.F. New representations of explicit one-step numerical methods for jump-diffusion stochastic differential equations. [In English]. *Computational Mathematics and Mathematical Physics*, 41, 6 (2001), 874-888. Available at: <http://www.sde-kuznetsov.spb.ru/01b.pdf>
- [39] Kuznetsov D.F. Comparative analysis of the efficiency of application of Legendre polynomials and trigonometric functions to the numerical integration of Ito stochastic differential equations. arXiv:1901.02345 [math.GM], 2019, 40 pp. [In English]
- [40] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals based on generalized multiple Fourier series. [In English]. *Ufa Mathematical Journal*, 11, 4 (2019), 49-77. DOI: <http://doi.org/10.13108/2019-11-4-49> Available at: http://matem.anrb.ru/en/article?art_id=604.
- [41] Kuznetsov D.F. Expansion of iterated Ito stochastic integrals of arbitrary multiplicity based on generalized multiple Fourier series converging in the mean. arXiv:1712.09746 [math.PR]. 2022, 111 pp. [In English].
- [42] Kuznetsov D.F. Exact calculation of the mean-square error in the method of approximation of iterated Ito stochastic integrals based on the generalized multiple Fourier series. arXiv:1801.01079 [math.PR]. 2018, 68 pp. [In English].
- [43] Kuznetsov D.F. Mean-square approximation of iterated Ito and Stratonovich stochastic integrals of multiplicities 1 to 6 from the Taylor-Ito and Taylor-Stratonovich expansions using Legendre polynomials. arXiv:1801.00231 [math.PR]. 2017, 106 pp. [In English].
- [44] Kuznetsov D.F. The hypotheses on expansion of iterated Stratonovich stochastic integrals of arbitrary multiplicity and their partial proof. arXiv:1801.03195 [math.PR]. 2022, 138 pp. [In English].
- [45] Kuznetsov D.F. Expansions of iterated Stratonovich stochastic integrals based on generalized multiple Fourier series: multiplicities 1 to 6 and beyond. arXiv:1712.09516 [math.PR]. 2022, 203 pp. [In English].
- [46] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of multiplicity 3 based on generalized multiple Fourier series converging in the mean: general case of series summation. arXiv:1801.01564 [math.PR]. 2018, 65 pp. [In English].
- [47] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of multiplicity 2 based on double Fourier-Legendre series summarized by Pringsheim method. arXiv:1801.01962 [math.PR]. 2018, 49 pp. [In English].
- [48] Kuznetsov D.F. Development and application of the Fourier method to the mean-square approximation of iterated Ito and Stratonovich stochastic integrals. arXiv:1712.08991 [math.PR]. 2017, 57 pp. [In English].

- [49] Kuznetsov D.F. Explicit one-step strong numerical methods of orders 2.0 and 2.5 for Ito stochastic differential equations based on the unified Taylor-Ito and Taylor-Stratonovich expansions. arXiv:1802.04844 [math.PR]. 2018, 31 pp. [In English].
- [50] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of arbitrary multiplicity based on generalized iterated Fourier series converging pointwise. arXiv:1801.00784 [math.PR]. 2018, 77 pp. [In English].
- [51] Kuznetsov D.F. Strong numerical methods of orders 2.0, 2.5, and 3.0 for Ito stochastic differential equations based on the unified stochastic Taylor expansions and multiple Fourier-Legendre series. arXiv:1807.02190 [math.PR]. 2018, 42 pp. [In English].
- [52] Kuznetsov D.F. Expansion of iterated stochastic integrals with respect to martingale Poisson measures and with respect to martingales based on generalized multiple Fourier series. arXiv:1801.06501v4 [math.PR]. 2018, 40 pp. [In English].
- [53] Kuznetsov D.F. Expansion of iterated Stratonovich stochastic integrals of fifth and sixth multiplicity based on generalized multiple Fourier series. arXiv:1802.00643 [math.PR]. 2022, 126 pp. [In English].
- [54] Kuznetsov M.D., Kuznetsov D.F. Implementation of strong numerical methods of orders 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 for Ito SDEs with non-commutative noise based on the unified Taylor-Ito and Taylor-Stratonovich Expansions and multiple Fourier-Legendre series. arXiv:2009.14011 [math.PR], 2020, 343 pp. [In English].
- [55] Kuznetsov D.F. Application of multiple Fourier-Legendre series to the implementation of strong exponential Milstein and Wagner-Platen methods for non-commutative semilinear SPDEs. Proceedings of the XIII International Conference on Applied Mathematics and Mechanics in the Aerospace Industry (AMMAI-2020). MAI, Moscow, 2020, pp. 451-453. Available at: <http://www.sde-kuznetsov.spb.ru/20e.pdf>
- [56] Kuznetsov M.D., Kuznetsov D.F. Optimization of the mean-square approximation procedures for iterated Ito stochastic integrals of multiplicities 1 to 5 from the unified Taylor-Ito expansion based on multiple Fourier-Legendre series arXiv:2010.13564 [math.PR], 2020, 63 pp. [In English].
- [57] Kuznetsov D.F. A new approach to the series expansion of iterated Stratonovich stochastic integrals of arbitrary multiplicity with respect to components of the multidimensional Wiener process. [In English]. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 2 (2022), 83-186. Available at: <http://diffjournal.spbu.ru/EN/numbers/2022.2/article.1.6.html>
- [58] Allen, E. Approximation of triple stochastic integrals through region subdivision. Communications in Applied Analysis (Special Tribute Issue to Professor V. Lakshmikantham). 17 (2013), 355-366.
- [59] Gihman I.I., Skorohod A.V. Stochastic Differential Equations and its Applications. Kiev: Naukova Dumka, 1982, 612 pp. [In Russian]
- [60] Kuznetsov, D.F. New representations of the Taylor-Stratonovich expansion. Journal of Mathematical Sciences (N. Y.). 118, 6 (2003), 5586-5596.
DOI: <http://doi.org/10.1023/A:1026138522239>
- [61] Prigarin S.M., Belov S.M. On one application of the Wiener process decomposition into series. Preprint 1107. Novosibirsk, Siberian Branch of the Russian Academy of Sciences, 1998, 16 pp. [In Russian].
- [62] Rybakov, K.A. Orthogonal expansion of multiple Itô stochastic integrals. Electronic Journal "Differential Equations and Control Processes" ISSN 1817-2172 (online), 3 (2021), 109-140. Available at: <http://diffjournal.spbu.ru/EN/numbers/2021.3/article.1.8.html>

DMITRIY FELIKSOVICH KUZNETSOV
PETER THE GREAT SAINT-PETERSBURG POLYTECHNIC UNIVERSITY,
POLYTECHNICHESKAYA UL., 29,
195251, SAINT-PETERSBURG, RUSSIA
Email address: sde_kuznetsov@inbox.ru