

Efficient, Certifiably Optimal High-Dimensional Clustering

Carson Eisenach* Han Liu†

December 11, 2017

Abstract

We consider SDP relaxation methods for data and variable clustering problems, which have been shown in the literature to have good statistical properties in a variety of settings, but remain intractable to solve in practice. In particular, we propose FORCE, a new algorithm to solve the Peng-Wei K -means SDP. Compared to the naive interior point method, our method reduces the computational complexity of solving the SDP from $\tilde{O}(d^7 \log \epsilon^{-1})$ to $\tilde{O}(d^6 K^{-2} \epsilon^{-1})$. Our method combines a primal first-order method with a dual optimality certificate search, which when successful, allows for early termination of the primal method. We show under certain data generating distributions that, with high probability, FORCE is guaranteed to find the optimal solution to the SDP relaxation and provide a certificate of exact optimality. As verified by our numerical experiments, this allows FORCE to solve the Peng-Wei SDP with dimensions in the hundreds in only tens of seconds. We also consider a variation of the Peng-Wei SDP for the case when K is not known a priori and show that a slight modification of FORCE reduces the computational complexity of solving this problem as well: from $\tilde{O}(d^7 \log \epsilon^{-1})$ using a standard SDP solver to $\tilde{O}(d^4 \epsilon^{-1})$.

1 Introduction

Clustering a set of objects according to some similarity measure is a central task of statistics and machine learning. A classical formulation for data clustering is K -means, which seeks an optimal partition of d points in \mathbb{R}^m into K clusters. In particular, if we denote $G = \{G_1, \dots, G_K\}$ a partition of the points x_1, \dots, x_d , the K -means problem is given by

$$\operatorname{argmin}_G \sum_{s=1}^K \sum_{i \in G_s} \|x_i - \mu_s\|_2^2, \text{ subject to } \mu_s = \frac{1}{|G_s|} \sum_{i \in G_s} x_i. \quad (1.1)$$

*Department of Operations Research and Financial Engineering, Princeton University, Princeton NJ 08544, USA; e-mail: eisenach@princeton.edu

†Tencent AI Lab, Bellevue WA 98004; email: han@tencent.com

For variable clustering, we consider n samples of a random variable $\mathbf{X} \in \mathbb{R}^d$ and have the objective of clustering the variables into K groups of size at least m . Informally, to apply K -means to variable clustering, one can treat $\text{Cov}(\mathbf{X})$ as specifying “distances” between d points.

Applications of clustering methods are wide-ranging – clustering problems appear everywhere from the analysis of medical imaging data to search result groupings on Google. In some problems, such as post-selection inference, optimal clusterings are required. Post-selection inference is the problem of first selecting a statistical model, e.g. through variable clustering, and then applying inferential procedures. Applications where variable clustering and statistical inference questions arise include the analysis of stock pricing, fMRI, and gene expression data.

In both data and variable clustering we seek the optimizer to some selection criterion, for example (1.1), but these problems are generally NP-hard (Dasgupta, 2008; Mahajan et al., 2012). To solve (1.1), Lloyd’s algorithm is a fast heuristic, but it cannot guarantee global optimality (Lloyd, 1982). As an attempt to find a global optimizer, one might run Lloyd’s algorithm many times and choose the best solution found or use an initialization scheme like `kmeans++`, but even these improvements lack global optimality guarantees (Arthur and Vassilvitskii, 2007). Moving beyond questions of global optimality, what if we do not even know K a priori? These are significant challenges, which we address by studying (1) *tractable algorithms for finding global optima*, (2) *certifying global optima* and (3) *clustering when K is unknown*.

In this work we study a convex relaxation of (1.1), originally introduced for data clustering by Peng and Wei (2007). Our investigation is motivated by recent work on variable clustering – Bunea et al. (2016) explores variable clustering for G -Latent Models, a statistical model with latent variable clusters, using this semi-definite programming (SDP) relaxation. They show the relaxation is tight with high probability and that using this relaxation gives the optimal statistical rate for exact cluster recovery (see Section 2). However, even though SDPs are solvable in polynomial time, standard SDP solvers have worst-case $\tilde{O}(d^7 \log \epsilon^{-1})$ running time due to the large number of constraints in the Peng-Wei SDP.

Compared to the variable clustering setting, the Peng-Wei SDP is relatively more studied in the data clustering literature. To apply a convex relaxation to a problem, we want to know when a relaxed problem and the original problem have the same minimizer. When they do, it suffices to solve the easier, relaxed problem. One method is to use duality theory to derive optimality certificates and then leverage the certificate to analyze when the relaxation is *tight*. Recent works employ this approach for data clustering under the Stochastic Ball Model (Awasthi and Bandeira, 2015; Iguchi et al., 2015, 2016) – see Section 2 for more details. Again, the difficulty is that even when the SDP relaxation is tight and we can in theory use it to solve (1.1), doing so is intractable in practice.

Our Contributions

In the contexts described above, the Peng-Wei SDP has desirable *statistical* properties, but solving it remains *computationally* out of reach. As mentioned, such methods are critical for the variable clustering setting as they enable previously out of reach statistical analyses.

To fill this gap, we introduce the FORCE (**F**irst-**O**rders **C**ertifiably **O**ptimal Clustering)

algorithm. The difficulty in solving NP-hard problems, such as K -means, derives from the integer structure of their solutions. The insight underlying the FORCE algorithm is that for clustering problems, when we expect the convex relaxation to be tight, the integer structure of the optimal solution can actually be leveraged to *help solve* the clustering problem. The FORCE algorithm consists of two components: a first-order method to solve \mathcal{P} and a dual solution construction used to certify the optimality of a primal solution.

The idea is that if we have an algorithm to quickly construct a dual solution at G^* and an interior point method to solve an SDP relaxation \mathcal{P} , then while solving \mathcal{P} we can periodically “round” the current iterate and search for a matching dual solution. If the primal and dual objective values match, the algorithm can terminate early. In addition, we often do not need to find the exact optimizer of the relaxed problem as once the current iterate is “sufficiently close” to the optimizer, most common clustering algorithms are likely to recover the same clustering as when applied to the optimizer.

To informally describe FORCE, we abstract away the technical details of our first-order method for the primal SDP by denoting $\mathcal{A}_P(\cdot) : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}^{d \times d}$ a *stateful algorithm* for solving the primal SDP. Each call $\mathcal{A}_P(t)$ should be thought of as performing t iterative updates and then returning the current iterate maintained by \mathcal{A}_P . Let $\mathcal{A}_R : \mathbb{R}^{d \times d} \rightarrow \mathcal{G}$, where \mathcal{G} is the space of all possible clusterings, be an algorithm for rounding an iterate \mathbf{U}_t to a candidate clustering \hat{G} . Finally, let $\mathcal{A}_D : \mathcal{G} \rightarrow \{\text{TRUE}, \text{FALSE}\}$ be a deterministic algorithm that attempts to construct a dual optimality certificate for input \hat{G} and returns true if it succeeds, and false otherwise. We also use this abstraction to frame the discussion of how to select the three complementary components \mathcal{A}_P , \mathcal{A}_R and \mathcal{A}_D . Algorithm 1 describes how FORCE produces a solution to the Peng-Wei SDP and the K -means problem.

In Section 3, we show that FORCE has worst-case computational-complexity $\tilde{\mathcal{O}}(d^6 K^{-2} \epsilon^{-1})$ to produce an ϵ -optimal solution to \mathcal{P} when K is known. When K is unknown, there is a variation on the Peng-Wei SDP that in certain settings has been shown to guarantee exact cluster recovery. With slight modification we can apply FORCE to this SDP too, and the gains are even greater than the case when K is known, with computational complexity $\tilde{\mathcal{O}}(d^4 \epsilon^{-1})$. As shown by our numerical results, FORCE often *requires only seconds for d in the hundreds and produces a certificate of exact optimality*. We also show that our dual certificate exists with high probability under a certain class of generating distributions (Section 3).

Notation. Denote either a clustering of data points or a partition of variables by $G = \{G_1, \dots, G_K\}$ where G_i is a single cluster or variable group. Hats, i.e. \hat{G} , always indicate quantities estimated from data and stars, i.e. G^* , always denote ground truths. For a $n \times n$ matrix \mathbf{M} , $\|\mathbf{M}\|_2$ denotes the largest eigenvalue of \mathbf{M} and $\|\mathbf{M}\|_\infty$ is the matrix ℓ_∞ norm. $\|\mathbf{M}\|_{\max} = \max_{i,j} M_{i,j}$ and $\|\mathbf{M}\|_{\min} = \min_{i,j} M_{i,j}$. Let S and S' be subsets of $[n]$. Then $\mathbf{M}_{S,S'}$ refers to the sub-matrix of M with entries whose row index is in S and column index is in S' . The notation $\tilde{\mathcal{O}}$ is used to suppress poly-log factors of the dimension d .

Algorithm 1 FORCE Clustering Algorithm

Input: h, T **Output:** $\mathbf{U}, \hat{G}, \text{optimal}$ $t \leftarrow 0$ $\text{optimal} \leftarrow \text{FALSE}$ **while** $t < T$ and $\text{optimal} == \text{FALSE}$ **do** $\mathbf{U}_t \leftarrow \mathcal{A}_P(1)$ $t \leftarrow t + 1$ **if** $t \bmod h == 0$ **then** $\hat{G}_t \leftarrow \mathcal{A}_R(\mathbf{U}_t)$ $\text{optimal} \leftarrow \mathcal{A}_D(\hat{G}_t)$ **if** $\text{optimal} == \text{TRUE}$ **then** $\text{return } \mathbf{U}_t, \hat{G}_t, \text{optimal}$ **end if****end if****end while** $\hat{G}_T \leftarrow \mathcal{A}_R(\mathbf{U}_T)$ **return** $\mathbf{U}_T, \hat{G}_T, \text{optimal}$

2 Preliminaries and Related Work

2.1 The K-means SDP and Its Dual

SDP relaxations often provide superior theoretical properties to other common types of convex relaxations, so we focus on these. As explained in Section 1, we study the Peng-Wei SDP, which is given as

$$\begin{aligned} & \underset{\mathbf{U}}{\text{maximize}} && \langle -\mathbf{D}, \mathbf{U} \rangle \\ & \text{subject to} && U_{ab} \geq 0 \text{ for } a, b \\ & && \mathbf{U}\mathbf{1} = \mathbf{1} \\ & && \text{tr}(\mathbf{U}) = K \\ & && \mathbf{U} \succeq 0. \end{aligned} \tag{2.1}$$

The matrix \mathbf{D} typically specifies distances between the points to be clustered. Thus $D_{i,j} = \|x_i - x_j\|_2^2$. If the task, instead of clustering data points, is clustering *variables*, \mathbf{D} can be the negated covariance matrix of the sampled data. The intuition is that this is similar to a distance measure. A solution is called “integer” if $U_{ij} = \frac{1}{|G_a|}$ if $i, j \in G_a$ and 0 otherwise, and it is said to correspond to the K -means solution G . This is also called the “partnership matrix” of the clustering solution G and we denote this matrix $B(G)$.

Because FORCE leverages a dual certificate, we need to find the dual SDP of (2.1),

$$\begin{aligned}
& \underset{y_{a,b}, y_a, y_T}{\text{minimize}} && 2 \sum_{a=1}^d y_a + K y_T \\
& \text{subject to} && \sum_{a=1}^d y_a \mathbf{R}_a + y_T \mathbf{I} \succeq -\mathbf{D} + \sum_{a \leq b} y_{a,b} \mathbf{I}_{a,b} \\
& && y_{a,b} \geq 0 \text{ for all } a \leq b.
\end{aligned} \tag{2.2}$$

The matrices $\mathbf{I}_{a,b}$ and \mathbf{R}_a are defined by $\mathbf{I}_{ab} = \frac{1}{2}(\mathbf{e}_a \mathbf{e}_b^T + \mathbf{e}_b \mathbf{e}_a^T)$ for all $a < b$, $\mathbf{I}_{aa} = \frac{1}{2} \mathbf{e}_a \mathbf{e}_a^T$, and $\mathbf{R}_a = \mathbf{1} \mathbf{e}_a^T + \mathbf{e}_a^T \mathbf{1}$.

2.2 Related Work

Second Order Methods. An obvious approach is to simply solve the SDP relaxation using the standard second-order convex optimization methods (see [Boyd and Vandenberghe \(2004\)](#) for some examples). It should be noted that for SDPs, standard black-box methods provide both an optimal solution to both the SDP and its dual, providing a certificate of approximate optimality. The problem is that for SDP relaxations of (1.1), standard methods have running time at least $\tilde{\mathcal{O}}(d^7 \log \epsilon^{-1})$, rendering them impractical.

Optimality Certificates for Data Clustering. In proving the tightness of (2.1) it is standard to derive an empirically testable condition on an instance of the clustering problem ([Awasthi and Bandeira, 2015](#); [Iguchi et al., 2015, 2016](#)). To do this, recent work on convex relaxations of K -means for data clustering takes a *dual optimality certificate* approach ([Awasthi and Bandeira, 2015](#); [Iguchi et al., 2015, 2016](#)).

Specifically, the strategy is: (1) find an appropriate convex relaxation (denoted \mathcal{P}) of (1.1), (2) find the dual problem (denoted \mathcal{D}) of \mathcal{P} , (3) given a proposed solution to \mathcal{P} construct a solution to \mathcal{D} with matching objective value, (4) derive a deterministic condition that can be checked on a given instance of \mathcal{P} and proposed solution to \mathcal{P} that guarantees the construction in (3) exists. The deterministic condition found in step (4) can then be analyzed to find the necessary assumptions on the data generating distribution to give the following guarantee: *with high probability a random instance of \mathcal{P} will satisfy the condition at the optimal solution G^* to \mathcal{P} .* To use the condition from step (4), all that remains is a way to “quickly” find optimal solutions to \mathcal{P} and then test the condition at the proposed optimal solution.

All three of the aforementioned works either explicitly or implicitly depend upon Theorem 4 from [Iguchi et al. \(2015\)](#) which characterizes optimal primal, dual solution pairs to (2.1) and (2.2). Specifically, the authors show that the following are equivalent: (1) \mathbf{B}^* is an optimal solution to (2.1), (2) every solution to (2.2) satisfies $y_{a,b} = 0$ for $a, b \in G_i^*$ and $\mathbf{Q}_{G_i^*, G_i^*} \mathbf{1} = 0$ for all i , and (3) every solution to (2.2) satisfies $\mathbf{y}_{G_i^*} = \mathbf{L}_{G_i^*, G_i^*}^{-1} (-\mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - y_T \mathbf{1})$. \mathbf{L} is a block-diagonal matrix determined by G^* , where the diagonal blocks are defined as $\mathbf{L}_{G_i^*, G_i^*} = |G_i^*| \mathbf{I} + \mathbf{1} \mathbf{1}^T$ and the off-diagonal blocks are zero. From this characterization of all optimal solution pairs, one can see that there are fewer “independent variables” in (2.2) than appear at first glance. The dual solutions used in [Awasthi and](#)

Bandeira (2015); Iguchi et al. (2016) differ from each other mainly in their choice of assignment to $y_{a,b}$. The choice of $y_{a,b}$ in turn determines what testable condition they can derive and leverage to prove tightness results and certify optimal clusterings.

The condition given in Iguchi et al. (2016) can be reduced to an eigenvalue condition on an empirical quantity, and so they introduce a “power iterator hypothesis test” which is used to certify the optimality of a clustering solution. Coupled with a fast method to find the optimal solution, their optimality certificate might be effective. Unfortunately, Iguchi et al. (2016) only offer a fast algorithm for the $K = 2$ case, and their method cannot be directly applied to variable clustering since it operates directly on the data points to be clustered, not merely the matrix \mathbf{D} .

In Section 3.2 we give a different choice of $y_{a,b}$, this one tailored to variable clustering. Further, our dual solution construction has an intuitive interpretation and is very easy to construct algorithmically for use in practice to certify optimal clusterings. Our choice of $y_{a,b}$ also yields a testable property, which we term the *Large Diagonal Blocks Property* (Property 1, see Section 3.2). If an instance satisfies this property, then FORCE is guaranteed to find both the optimal clustering to (1.1) and a certificate of optimality. The FORCE certificate has the nice property that it is actually a solution to the dual problem and thus it is easy to use the certificate to verify optimality.

Convex Relaxation Methods for Variable Clustering. Bunea et al. (2016) studies variable clustering in cluster-based Gaussian graphical models where the observed random vector $\mathbf{X} := (X_1, \dots, X_d)$ has some latent structure. In particular, they consider the G -Latent model, where there exists an unknown partition $G^* := \{G_1^*, \dots, G_K^*\}$ of the indices $\{1, \dots, d\}$. Under this model, there exists a K -dimensional latent random vector \mathbf{Z} such that $\mathbf{X} = \mathbf{A}\mathbf{Z} + \mathbf{E}$, where \mathbf{E} is a mean zero error vector and \mathbf{A} is the assignment matrix corresponding to G^* – that is, $A_{jk} = \mathbb{I}\{j \in G_k\}$. It is assumed \mathbf{Z} is centered and the noise \mathbf{E} is independent of \mathbf{Z} . We denote $\text{Cov}(\mathbf{Z}) = \mathbf{C}^*$ and $\text{Cov}(\mathbf{E}) = \mathbf{\Gamma}^*$, a diagonal matrix.

To recover the true group partition G^* , the authors propose using (2.1) with $\mathbf{D} = \widehat{\mathbf{\Gamma}} - \widehat{\mathbf{\Sigma}}$, a penalized covariance matrix estimator (we refer to this as the PECOK Estimator). Because a priori the group structure is unknown, an estimator $\widehat{\mathbf{\Gamma}}$ of $\mathbf{\Gamma}^*$ is somewhat involved so we omit the details here. For our purposes, we are concerned with its rate of convergence in the max-norm. Bunea et al. (2016) show that if $\mathbf{X}_1, \dots, \mathbf{X}_n$ are generated from a G -Latent Model, then there exist constants c_1, c_2 such that with probability at least $1 - c_2/d^2$,

$$\|\widehat{\mathbf{\Gamma}} - \mathbf{\Gamma}^*\|_\infty \leq c_1 \|\mathbf{\Gamma}^*\|_\infty \sqrt{\log d/n}. \quad (2.3)$$

The authors also show that under the G -Latent model, if $\Delta(\mathbf{C}^*)$ is sufficiently large, then the optimizer to (2.1) is integer with high probability and it corresponds to the true clustering. Specifically, they show that if $\log d \leq c_1 n$ and

$$\Delta(\mathbf{C}^*) \leq c_2 \|\mathbf{\Gamma}\|_\infty \left(\sqrt{\log d/mn} + \sqrt{\log d/m^2n} + \log d/n + d/mn \right),$$

then with probability at least $1 - c_3/d$ the optimizer to (2.1) is $\mathbf{X}^* = B(G^*)$.

Bunea et al. (2016) show that the optimal solution to (2.1) recovers G^* with high probability, but they do not give an algorithm for *solving* (2.1) or a method for *certifying* the optimality of a proposed solution to (2.1). Our work with the FORCE algorithm provides both of those. In Section

A we show that the FORCE certificate exists with high probability under the G -Latent Model, and is “nearly optimal” (in a sense to be made precise later), which incidentally also implies tightness of (2.1) under G -Latent Model (albeit with slightly stronger assumptions than the result shown in Bunea et al. (2016)).

Other Clustering Methods. There are too many clustering methods to discuss in depth here, but some popular methods include spectral clustering (Kumar and Kannan, 2010; Awasthi and Sheffet, 2012) and local linkage based clustering schemes. For our motivating application, these spectral clustering results are not useful as they are tailored towards data clustering and Bunea et al. (2016) show that a variant of spectral clustering is provably suboptimal in terms of its ability to guarantee exact recovery of the variable clusters. Likewise, linkage based schemes do not have strong global optimality guarantees. As a subroutine of FORCE, we make use of Lloyd’s Algorithm (Lloyd, 1982) and CLINK (Defays, 1977), a linkage based clustering method.

Approaches for Speeding Up SDPs. Given the goal of finding an optimizer to (2.1), there are several methods for quickly finding optimal solutions to SDPs. One well known approach is the matrix multiplicative weights algorithm due to Arora et al. (2005). For our problem, when K is known the MMW algorithm requires $\tilde{\mathcal{O}}(K^2 d^2 \alpha^{-2} \epsilon^{-2})$ where α is related to a lower bound on the optimal value of the SDP under a rescaling and ϵ is the multiplicative accuracy. Typically, we have $\alpha \geq c/d$ for some constant c . In this case, the running time bound becomes $\tilde{\mathcal{O}}(K^2 d^4 \epsilon^{-2})$, which is worse than ours in its dependence upon ϵ . We note that ϵ in our result does not have the exact same meaning as in their result; in ours it is related to an additive error and in theirs, a multiplicative error. We also mention that we implemented a MMW algorithm for the K -means SDP, but found it to be unstable. We suspect this is due to the presence of the equality constraints as an equality constraint is not satisfied exactly by an iterate of the MMW algorithm.

Instead we follow the approach of converting the semi-definite program into an equivalent eigenvalue maximization problem using the technique in Renegar (2014), which allows us to achieve better worst case bounds. This is described in more detail in the next section.

2.3 First-Order Methods for SDP

Based on the discussion above, we employ the technique from Renegar (2014) to derive a method for the primal K -means SDP. The author proposes trading the positive semidefinite constraint for a non-smooth objective function. Consider an SDP in standard form,

$$\begin{aligned} & \underset{\mathbf{U}}{\text{minimize}} && \langle \mathbf{D}, \mathbf{U} \rangle \\ & \text{subject to} && \langle \mathbf{A}_i, \mathbf{U} \rangle = b_i \text{ for } i = 1, \dots, p \\ & && \mathbf{U} \succeq 0, \end{aligned} \tag{2.4}$$

where $\mathbf{A}_i \in \mathcal{S}^{n \times n}$, $\mathbf{D} \in \mathcal{S}^{n \times n}$ and $b_i \in \mathbb{R}$. Denote the optimal value of (2.4) by u^* and \mathcal{C} the set of feasible solutions. Given some strictly feasible solution \mathbf{F} for (2.4), a projection can be defined from \mathbf{F} onto the border of the positive semi-definite cone by

$$P_{\mathbf{F}}(\mathbf{U}) = \mathbf{F} + \frac{1}{1 - \lambda_{\min, \mathbf{F}}(\mathbf{U})} (\mathbf{U} - \mathbf{F}),$$

where $\lambda_{\min, F}(\mathbf{U}) = \lambda_{\min}(\mathbf{F}^{-1/2}\mathbf{U}\mathbf{F}^{-1/2})$. $P_{\mathbf{F}}(\mathbf{U})$ lies at the intersection of the line segment between \mathbf{F} and \mathbf{U} and the positive semi-definite cone. Clearly if $\mathbf{U} \in \mathcal{S}_+^{n \times n}$ then $P_{\mathbf{F}}(\mathbf{U}) \in \mathcal{S}_+^{n \times n}$. Now, let $u_0 \in \mathbb{R}$ satisfying $u_0 < \langle \mathbf{D}, \mathbf{F} \rangle$. Renegar (2014) shows in their Theorem 2.2 that if \mathbf{V}^* is a global optimum for

$$\begin{aligned} & \underset{\mathbf{V}}{\text{maximize}} && \lambda_{\min, F}(\mathbf{V}) \\ & \text{subject to} && \langle \mathbf{A}_i, \mathbf{V} \rangle = b_i \text{ for } i = 1, \dots, p \\ & && \langle \mathbf{D}, \mathbf{V} \rangle = u_0, \end{aligned} \tag{2.5}$$

then $P_{\mathbf{F}}(\mathbf{V}^*)$ is optimal for (2.4). In addition, if \mathbf{U}^* is optimal for (2.4), then

$$\mathbf{V}^* = \mathbf{F} + \frac{\langle \mathbf{D}, \mathbf{F} \rangle - u_0}{\langle \mathbf{D}, \mathbf{F} \rangle - u^*} (\mathbf{U}^* - \mathbf{F})$$

is optimal for (2.5).

If $\mathbf{F} = \mathbf{I}$, Renegar (2014) demonstrates convergence using Nesterov’s projected gradient descent with a constant step size. The objective function of (2.5) is not smooth, but it is still convex, and thus at all points in its domain there exists a sub-gradient. The proof of convergence, however, relies on the fact that $\lambda_{\min}(\mathbf{U})$ is 1-Lipschitz. Thus if $\mathbf{F} \neq \mathbf{I}$, the necessary choice of step size and the resulting rate of convergence is not as clear. See Appendix F for more details.

The advantage of solving (2.5) is that though it may be difficult to compute the projection onto the PSD cone in (2.4), it may be cheap to project onto \mathcal{C}_λ , the feasible set of (2.5), in each iteration of gradient ascent. Though typically the projected sub-gradient method is presented as a gradient update followed by a projection onto \mathcal{C}_λ , it is equivalent to first project the gradient onto $\mathcal{C}_\lambda^\perp = \{\mathbf{U} | \langle \mathbf{A}_i, \mathbf{U} \rangle = 0, \langle \mathbf{D}, \mathbf{U} \rangle = 0\}$ and then performing the update. For our purposes, it is easier to work with the Karush-Kuhn-Tucker (KKT) conditions for the projection onto $\mathcal{C}_\lambda^\perp$,

$$\mathcal{P}_{\mathcal{C}_\lambda^\perp}(\mathbf{U}) = \underset{\mathbf{V} \in \mathcal{D}^\perp}{\operatorname{argmin}} \frac{1}{2} \|\mathbf{V} - \mathbf{U}\|_F^2, \tag{2.6}$$

which we show in Section 3.1 can be reduced to solving a relatively small system of linear equations.

Applying Nesterov Smoothing

To speed up convergence, we use a smoothing technique from Nesterov (2005). By smoothing, we can apply Nesterov’s accelerated projected gradient descent and obtain a fast convergence rate. The function $\lambda_{\min}(\mathbf{V})$ is smoothed as $f_\mu(\mathbf{V}) = -\mu \log \left(\sum_j \exp(-\lambda_j(\mathbf{V})/\mu) \right)$. Though there are some necessary technical details to ensure the solution is sufficiently close to optimality for the original problem, Renegar’s method can be thought of as using Nesterov’s accelerated projected gradient descent to solve f_μ and then projecting back to a solution of the original SDP. The authors note that their approach can be applied to problems with general \mathbf{F} , but do not analyze how the convergence rate is effected by a different choice of \mathbf{F} . These are necessary to analyze FORCE and we provide them, for completeness, in Appendix F. The algorithm is called the “Smoothed Scheme” and a version for arbitrary \mathbf{F} is given as Algorithm 7, also in Appendix F.

3 The FORCE Algorithm

Because we consider clustering in the high-dimensional setting, a fast algorithm to solve (2.1) is critical. For the design \mathcal{A}_P , second-order methods have an appealing iteration complexity, but the per iteration cost is prohibitive for (2.1). This is because the cost of each iteration depends not only on the dimension d of the K -means-type problem, *but also on the number of constraints* – in (2.1), this is $\mathcal{O}(d^2)$. As a result, the overall running time to solve the primal problem is at best $\tilde{\mathcal{O}}(d^7 \log \epsilon^{-1})$.

First-order methods, by contrast, may have a higher iteration complexity, but a lower per-iteration cost. As is described below, the FORCE algorithm uses a first-order approach based on the one described in Section 2 as \mathcal{A}_P . By doing so, each iterative update requires only $\mathcal{O}(d^3)$ computations because we avoid the projection onto the PSD cone and the hefty cost induced by the d^2 entry-wise non-negativity constraints.

In this section we consider FORCE when K is known (for K unknown, see Section 4). It proceeds as follows:

1. Section 3.1 derives \mathcal{A}_P by demonstrating the generic approach in Renegar (2014) can be applied to (2.1) by converting it to an eigenvalue maximization, finding the projection onto \mathcal{C}^\perp , and demonstrating the existence of a strictly feasible solution;
2. Section 3.2 derives \mathcal{A}_D by giving an intuitive optimality certificate construction;
3. Section 3.3 provides bounds on the running time of FORCE when K is fixed.

3.1 Primal Step

The generic first-order method to solving SDPs described in Section 2 is only applicable to SDPs in standard form with at least one strictly feasible solution. In addition, the projection onto the set of constraints must be relatively inexpensive to compute or no advantage is offered over standard second-order methods. As it stands, (2.1) is not in standard form, the complexity of computing the projection is unclear and the identity matrix \mathbf{I} is not feasible. In this section, we first give the conversion of (2.1) to an eigenvalue maximization problem, then demonstrate the computational complexity of the projection step, and lastly prove the existence of a strictly feasible solution. Therefore we can choose the iterative method \mathcal{A}_P to be a version of Renegar’s Smoothed-Scheme (Algorithm 7 in Appendix F).

Conversion to an Eigenvalue Maximization Problem

To apply Renegar's method and obtain a first-order algorithm for (2.1), we must first convert to standard form. We first write (2.1) in terms of matrix (in)equality constraints

$$\begin{aligned}
& \underset{\mathbf{U}}{\text{maximize}} && \langle -\mathbf{D}, \mathbf{U} \rangle \\
& \text{subject to} && \langle \mathbf{I}_{a,b}, \mathbf{U} \rangle \geq 0 \text{ for all } a \leq b \\
& && \langle \mathbf{R}_a, \mathbf{U} \rangle = 2 \text{ for all } a \\
& && \langle \mathbf{I}, \mathbf{U} \rangle = K \\
& && \mathbf{U} \succeq 0.
\end{aligned} \tag{3.1}$$

To complete the transformation into standard form, we introduce slack variables for the $\mathbf{I}_{a,b}$ inequality constraints by constructing the variable

$$\mathbf{U}' = \left[\begin{array}{c|c} \mathbf{U} & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{U}_C \end{array} \right],$$

where \mathbf{U}_C is a $d^2 \times d^2$ diagonal matrix of slack variables – hence the subscript C , corresponding to the constraint set. Denote

$$\mathbf{I}'_{a,b} = \left[\begin{array}{c|c} \mathbf{I}_{a,b} & \mathbf{0} \\ \hline \mathbf{0} & \frac{-1}{2} \text{diag}(\mathbf{e}_{a,b}) \end{array} \right], \mathbf{R}'_a = \left[\begin{array}{c|c} \mathbf{R}_a & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} \end{array} \right], \mathbf{I}' = \left[\begin{array}{c|c} \mathbf{I} & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} \end{array} \right] \text{ and } \mathbf{D}' = \left[\begin{array}{c|c} \mathbf{D} & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{0} \end{array} \right]. \tag{3.2}$$

Above, $\mathbf{e}_{a,b}$ denotes the d^2 -dimensional vector of 0s with 1s in only the $((a-1)d+b)^{th}$ and $((b-1)d+a)^{th}$ positions and encodes the slack variables for the non-negativity constraints. The standard form of (2.1) is thus

$$\begin{aligned}
& \underset{\mathbf{U}'}{\text{minimize}} && \langle \mathbf{D}', \mathbf{U}' \rangle \\
& \text{subject to} && \langle \mathbf{I}'_{ab}, \mathbf{U}' \rangle = 0 \text{ for all } a \leq b \\
& && \langle \mathbf{R}'_a, \mathbf{U}' \rangle = 2 \text{ for all } a \\
& && \langle \mathbf{I}'_{d \times d}, \mathbf{U}' \rangle = K \\
& && \mathbf{U}' \succeq 0.
\end{aligned} \tag{3.3}$$

Note that a negation of the optimal value of (3.3) is necessary for equivalence of objective values, but it does not affect the optimal solution so we ignore it. Given a strictly feasible solution \mathbf{F} and \mathbf{U}_0 such that $\langle -\mathbf{D}, \mathbf{F} \rangle < \langle -\mathbf{D}, \mathbf{U}_0 \rangle = -u_0$ to (2.1), it is clear that

$$\mathbf{F}' = \left[\begin{array}{c|c} \mathbf{F} & \mathbf{0} \\ \hline \mathbf{0} & \text{diag}(\text{vec}(\mathbf{F})) \end{array} \right] \text{ and } \mathbf{U}'_0 = \left[\begin{array}{c|c} \mathbf{U}_0 & \mathbf{0} \\ \hline \mathbf{0} & \text{diag}(\text{vec}(\mathbf{U}_0)) \end{array} \right]$$

are the necessary pair \mathbf{F}' , and \mathbf{U}'_0 to apply the method described in Section 2 to (3.3). Turning (3.3) into an equivalent eigenvalue maximization problem gives

$$\begin{aligned}
& \underset{\mathbf{V}'}{\text{maximize}} && \lambda_{\min} \left((\mathbf{F}')^{-1/2} \mathbf{V}' (\mathbf{F}')^{-1/2} \right) \\
& \text{subject to} && \langle \mathbf{I}'_{ab}, \mathbf{V}' \rangle = 0 \text{ for all } a, b \\
& && \langle \mathbf{R}'_a, \mathbf{V}' \rangle = 2 \text{ for all } a \\
& && \langle \mathbf{I}', \mathbf{V}' \rangle = K \\
& && \langle \mathbf{D}', \mathbf{V}' \rangle = u_0.
\end{aligned} \tag{3.4}$$

Note that

$$\lambda_{\min} \left(\mathbf{F}'^{-\frac{1}{2}} \mathbf{V}' \mathbf{F}'^{-\frac{1}{2}} \right) = \min \left\{ \lambda_{\min} \left(\mathbf{F}^{-1/2} \mathbf{V} \mathbf{F}^{-1/2} \right), \min_{i,j} \frac{X_{i,j}}{F_{i,j}^{-1}} \right\}$$

because the eigenvalues of a block diagonal matrix are the combined eigenvalues of the blocks.

Constraint Set Projection

To project onto $\mathcal{C}_\lambda^\perp$, we must find the optimizer for $\mathcal{P}_{\mathcal{C}_\lambda^\perp}(\mathbf{U}')$. Notationally, $(U_C)_{a,b}$ refers to the $((a-1)d+b)^{\text{th}}$ diagonal entry in \mathbf{U}_C as it is a diagonal matrix of the slack variables. Because a projection onto a convex set has a unique minimizer, any point satisfying the KKT conditions for the projection must be the projected point. Solving for the projection gives the following system of $d+2$ equations in $d+2$ unknowns:

$$\begin{aligned}
\sum_{b=1}^d U_{ab} + \sum_{b=1}^d (U_C)_{ab} &= \sum_{b=1}^d y_b^* + dy_a^* + y_T^* + \left[\sum_{b=1}^d D_{ab} \right] \lambda^* && \text{for } a \in [d] \\
\text{tr}(\mathbf{U}) + \text{tr}(\mathbf{U}_C) &= \sum_{a=1}^d y_a^* + dy_T^* + \text{tr}(\mathbf{D})\lambda^* \\
\text{tr}(\mathbf{D}\mathbf{U}) + \text{tr}(\mathbf{D}\mathbf{U}_C) &= 2 \sum_{a=1}^d \left[\sum_{b=1}^d D_{ab} \right] y_a^* + \text{tr}(\mathbf{D})y_T^* + \text{tr}(\mathbf{D}\mathbf{D})\lambda^*.
\end{aligned} \tag{3.5}$$

Solving (3.5), we get the projected matrix

$$\mathcal{P}_{\mathcal{C}_\lambda^\perp}(\mathbf{V}^*) = \left[\begin{array}{c|c} \mathbf{V}^* & \mathbf{0} \\ \hline \mathbf{0} & \text{diag}(\text{vec}(\mathbf{V}^*)) \end{array} \right], \text{ where } \mathbf{V}^* = \frac{1}{2} \left[\mathbf{U} + \mathbf{U}_C - \sum_{a=1}^d \mathbf{R}_a Y_a^* - y_T^* \mathbf{I} - \lambda^* \mathbf{D} \right]. \tag{3.6}$$

Existence of a Strictly Feasible Solution

Unlike for the SDP's considered by Renegar (2014), \mathbf{I} is not feasible for (2.1) as $K < d$, $\text{tr}(\mathbf{I}) = d \neq K$. We also note that the intuitive idea to find a possibly suboptimal clustering \widehat{G} and use $\mathbf{F} = B(\widehat{G})$ is not possible because strict feasibility for (2.1) requires all $\mathbf{F}_{ij} > 0$.

Nonetheless, there are valid choices of \mathbf{F} . Consider matrices of the form $\mathbf{F} = a\mathbf{I} + b\mathbf{1}\mathbf{1}^T$, where $a, b > 0$. Such matrices clearly satisfy $\mathbf{F}_{ij} > 0$ and $\mathbf{F} \succ 0$, so all that remains is to choose a and b

such that $\langle \mathbf{F}, \mathbf{I} \rangle = K$ and $\mathbf{F}\mathbf{1} = \mathbf{1}$. Multiplying these expressions out, simplifying and solving the resulting system of equations gives

$$a = \frac{K-1}{d-1} \text{ and } b = \frac{d-K}{d^2-d}.$$

Lemma 3.1 summarizes the properties of \mathbf{F} .

Lemma 3.1. Given d and K , define

$$\mathbf{F}_{d,K} := \frac{K-1}{d-1}\mathbf{I} + \frac{d-K}{d^2-d}\mathbf{1}\mathbf{1}^T.$$

$\mathbf{F}_{d,K}$ is strictly feasible for (2.1) and $\|\mathbf{F}^{-1}\|_2 = \frac{d-1}{K-1}$.

Proof of Lemma 3.1. The first claim follows by the previous discussion and the second follows immediately from Lemma B.1. \square

3.2 FORCE Algorithm: Dual Step

The key idea underlying the FORCE algorithm is to alternate iterations of the primal method with a dual optimality certificate search. Because all instances of (2.1) are strictly feasible, as shown in Lemma 3.1, then for any primal optimal solution there exists a dual solution such that its objective value is exactly equal to the primal. Finding this solution can provide the desired certificate of optimality. However, unlike the primal problem, the dual does not lend itself easily to mapping a solution for the K -means problem onto a feasible solution of the the dual SDP. The approach of our dual solution construction is to derive properties that when satisfied demonstrate the proposed solution is dual feasible, and design our construction accordingly.

FORCE Dual Solution Construction

Let $\widehat{G} = \{\widehat{G}_1, \dots, \widehat{G}_K\}$ be the candidate clustering for which we want to find a dual solution. Because the goal is to certify optimality, take $\widehat{G} = G^*$. Without loss of generality, also assume that the variables are ordered according to G^* , so that $\mathbf{B}^* = B(G^*)$ is block-diagonal. Denote by $d^* = \langle \mathbf{D}, \mathbf{B}^* \rangle$ and $\mathbf{Q} := \sum_{a=1}^d y_a \mathbf{R}_a + y_T \mathbf{I} + \mathbf{D} - \sum_{a \leq b} y_{a,b} \mathbf{I}_{a,b}$. Complementary slackness gives that for $a \in G_i^*$ and $b \in G_i^*$, $y_{a,b} = 0$. If we can somehow “eliminate” the off-diagonal blocks in \mathbf{Q} , finding a dual solution should be very straightforward. This inspires Property 1.

Property 1 (Large Diagonal Blocks Property). An instance \mathbf{D} of a clustering problem satisfies the Large Diagonal Blocks Property if there exists a feasible dual solution with value d^* such that the variables $y_{a,b}$ can be chosen to make off-diagonal blocks of the matrix \mathbf{Q} equal to $\mathbf{0}$.

Intuitively, we expect that in the variable clustering setting Property 1 will frequently hold. Because $-\mathbf{D}$ is estimate of a covariance matrix for a model with block covariance structure, the diagonal blocks should dominate the off-diagonal blocks. If an instance satisfies Property 1, then there is some assignment to y_a and y_T attaining d^* such that choosing $y_{a,b} = 0$ if $a = b$ and $y_{a,b} = y_a + y_b + D_{a,b}$ otherwise, gives $\mathbf{Q} \succeq 0$ and $y_{a,b} \geq 0$ for all a and b .

What remains is to search over assignments to y_a and y_T . In light of the characterization of all dual optimal solutions in Section 2.2, the FORCE dual solution construction can be viewed as a function of y_T :

$$\mathbf{y}_{G_i^*}(\mathbf{D}, y_T) = \mathbf{L}_{G_i^*, G_i^*}^{-1}(-\mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - y_T \mathbf{1}) \text{ and } y_{a,b}(\mathbf{D}, y_T) = \begin{cases} 0, & \text{if } a = b \\ y_a + y_b + D_{a,b}, & \text{o/w.} \end{cases}$$

Performing binary search on the value of y_T , we can obtain such a dual solution if and only if Property 1 is satisfied. The details are given as Algorithm 3 in Appendix C. For variable clustering in G -Latent Models, this is a safe assumption (see Appendix A), but the procedure can be modified to only require that y_T lies in a bounded interval.

3.3 Theoretical Results

On its own, the FORCE Primal Step (see Section 3.1), offers an improved theoretical guarantee over second-order interior point methods. As described informally by Algorithm 1, the FORCE algorithm alternates the FORCE Primal Step and FORCE dual steps. Taking \mathcal{A}_P to be the smoothed-scheme applied to (2.1), described in Section 3.1, \mathcal{A}_D the certificate search described in Section 3.2, and \mathcal{A}_R to be Lloyd’s Algorithm, Algorithm 1 becomes Algorithm 2. By appropriately choosing the dual certificate search frequency h , the convergence rate properties of the primal step transfer to FORCE. These results are summarized as Theorem 3.2.

Algorithm 2 First-Order Certifiable Clustering (FORCE)

Input: $0 < \epsilon < 1$, \mathbf{D} , h , \mathbf{U}_0 , \mathbf{F}

Output: G^*

Run Algorithm 7 with inputs ϵ , \mathbf{D} , \mathbf{U}_0 , \mathbf{F} .

for Every h^{th} gradient update while running Renegar’s Smoothed Scheme **do**

Denote the current iterate as \mathbf{V}_s

Round $P_{\mathbf{F}}(\mathbf{V}_s)$ using Lloyd’s Algorithm to obtain \widehat{G}_s .

Construct \mathbf{B}_s , the partnership matrix corresponding to \widehat{G}_s

Run Algorithm 3 on \mathbf{B}_s .

if Dual and primal objective values match **then**

return \widehat{G}_s

end if

end for

Denote the output of Algorithm 7 as \mathbf{V}_T

Run Lloyd’s algorithm on $P_{\mathbf{F}}(\mathbf{U}_T)$ to obtain \widehat{G}_T

return \widehat{G}_T

Theorem 3.2 (Convergence Rate of FORCE for Fixed K). Assume that the matrix \mathbf{D} satisfies $\|\mathbf{D}\|_2 \leq C$ for some constant C . Then, FORCE (Algorithm 2) terminates after $\widetilde{\mathcal{O}}(d^6 K^{-2} \epsilon^{-1})$ time, giving an ϵ -optimal solution.

Proof of Theorem 3.2. We start by showing that the claim holds for Renegar’s “smoothed scheme” applied to (2.1). Note that for any \mathbf{U} and $\mathbf{V} \in \mathcal{C}$, $\|\mathbf{U} - \mathbf{V}\|_F \leq \sqrt{2d}$. For $\mathbf{F}_{d,K}$, applying Lemma 3.1 gives $\|\mathbf{F}_{d,K}^{-1}\|_2^2 = \frac{d-1}{K-1}$. The convergence rate of the smoothed scheme (given as Theorem F.6) gives that the number of sub-gradient updates required is at most

$$T = \left(2\sqrt{2\log d}\right) \frac{d(d-1)^2}{(K-1)^2} \left(\frac{1}{\epsilon} + \log_{5/4} \left(\frac{\langle \mathbf{D}, \mathbf{F} \rangle - u^*}{\langle \mathbf{D}, \mathbf{F} \rangle - u_0}\right)\right).$$

Because each gradient update requires $\mathcal{O}(d^3)$ operations, the running time of Renegar’s smoothed scheme is bounded by $\tilde{\mathcal{O}}(d^6 K^{-2} \epsilon^{-1})$, proving the first claim.

To show the main claim, we start by analyzing Algorithm 3 which requires at most $C \log C$ iterations of binary search. Pre-computing the transformations for $\mathbf{y}_{G_i^*}$ requires at most $\mathcal{O}(d^3)$ arithmetic operations. Each iteration therefore requires computing a minimum eigenvalue of a d -dimensional matrix. This gives an overall running time of $\tilde{\mathcal{O}}(Cd^3)$ for the dual certificate construction.

Lastly, the rounding step requires $\mathcal{O}(d^2 KI)$ running time, where I is the number of updates Lloyd’s Algorithm performs. We can choose to impose an iteration limit to obtain the necessary running time bounds, or replace it with another clustering method such as CLINK which has a running time of $\mathcal{O}(d^2)$. Combining the bounds given above, the second claim is proved. \square

Theorem 3.3 shows that for variable clustering under the G -Latent model, the FORCE certificate exists with high probability (with respect to the generating distribution) at the true grouping G^* . The rate is nearly optimal in terms of the required cluster separation $\Delta \mathbf{C}^*$ and also implies the tightness of (2.1) with high probability (albeit at a slightly less than optimal rate, see Bunea et al. (2016)). See Appendix A for a more in depth discussion.

Theorem 3.3 (FORCE Certificate Exists with High Probability). Consider the variable clustering setting under the G -Latent model and assume $\log d \leq c_0 n$, where c_0 is an absolute constant. There exist universal constants c_1, c_2 and c_3 such that if

$$\Delta \mathbf{C}^* \geq c_1 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \sqrt{\frac{\log d}{nm^2}} + \sqrt{\frac{d}{nm^2}} + \frac{d}{nm} \right) + c_2 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right),$$

then with probability at least $1 - c_3/d$ the FORCE Dual Certificate exists at G^* , where $\sigma_1 = \max_i C_{i,i}^*$ and $\sigma_2 = (\|\mathbf{\Gamma}^*\|_\infty \max_i C_{i,i}^*)^{1/2}$.

The proof of Theorem 3.3 is significantly more involved, so it is deferred to Appendix A.

4 Extension of FORCE to Unknown K

The motivation and insight behind the FORCE algorithm remains the same when K is unknown, so we do not repeat the full discussion given in Section 3. When K is not known a priori, it can sometimes be estimated simultaneously by altering the SDP relaxation – we can exchange the trace constraint for an appropriately chosen trace penalty. In the variable clustering setting, Bunea et al. (2016) show that (4.1) recovers the optimal solution to (1.1) without requiring K to be known a priori.

K-means Adaptive SDP

We will refer to (4.1) as the K -means Adaptive SDP due to that its ability to *adaptively* select the number of clusters and the optimal clustering simultaneously. The trace penalty is defined by a data driven tuning parameter $\widehat{\kappa}$. It is beyond the scope of this work to consider the theoretical properties of (4.1) for data clustering and the remainder of this section focuses on the variable clustering setting.

$$\begin{aligned}
& \underset{\mathbf{U}}{\text{maximize}} && \langle -\mathbf{D} - \widehat{\kappa}\mathbf{I}, \mathbf{U} \rangle \\
& \text{subject to} && U_{ab} \geq 0 \text{ for } a, b \\
& && \mathbf{U}\mathbf{1} = \mathbf{1} \\
& && \mathbf{U} \succeq 0.
\end{aligned} \tag{4.1}$$

The dual SDP is given by (4.2)

$$\begin{aligned}
& \underset{y_{a,b}, y_a}{\text{minimize}} && 2 \sum_{a=1}^d y_a \\
& \text{subject to} && \sum_{a=1}^d y_a \mathbf{R}_a + \widehat{\kappa}\mathbf{I} \succeq -\mathbf{D} + \sum_{a \leq b} y_{a,b} \mathbf{I}_{a,b} \\
& && y_{a,b} \geq 0 \text{ for all } a \leq b.
\end{aligned} \tag{4.2}$$

Conversion to Eigenvalue Maximization

The conversion to standard form and an eigenvalue maximization problem are nearly identical to those for the case when K is known, so the derivations are omitted. All notation is carried over from Section 3, thus in standard form (4.1) becomes

$$\begin{aligned}
& \underset{\mathbf{U}'}{\text{minimize}} && \langle \mathbf{D}' + \widehat{\kappa}\mathbf{I}', \mathbf{U}' \rangle \\
& \text{subject to} && \langle \mathbf{I}'_{ab}, \mathbf{U}' \rangle = 0 \text{ for all } a \leq b \\
& && \langle \mathbf{R}'_a, \mathbf{U}' \rangle = 2 \text{ for all } a \\
& && \mathbf{U}' \succeq 0,
\end{aligned} \tag{4.3}$$

and the corresponding eigenvalue maximization problem is

$$\begin{aligned}
& \underset{\mathbf{V}'}{\text{maximize}} && \lambda_{\min}(\mathbf{V}') \\
& \text{subject to} && \langle \mathbf{I}'_{ab}, \mathbf{V}' \rangle = 0 \text{ for all } a, b \\
& && \langle \mathbf{R}'_a, \mathbf{V}' \rangle = 1 \text{ for all } a \\
& && \langle \mathbf{D}' + \widehat{\kappa}\mathbf{I}', \mathbf{V}' \rangle = u_0.
\end{aligned} \tag{4.4}$$

Constraint Projection

As in the case when K is known, we must derive a projection onto $\mathcal{C}_\lambda^\perp$. Solving the KKT conditions, we find the projected matrix as

$$\mathcal{P}_{\mathcal{C}_\lambda^\perp}(\mathbf{V}'_*) = \left[\begin{array}{c|c} \mathbf{V}^* & \mathbf{0} \\ \hline \mathbf{0} & \text{diag}(\text{vec}(\mathbf{V}^*)) \end{array} \right],$$

where

$$\mathbf{V}_* = \frac{1}{2} \left[\mathbf{U} + \mathbf{U}_C - \sum_{a=1}^d \mathbf{R}_a Y_a^* - \lambda^* (\mathbf{D} + \widehat{\kappa} \mathbf{I}) \right].$$

Existence of a Feasible Solution

Clearly for (4.1) \mathbf{I} is feasible, but unfortunately is not strictly feasible. Thus we still must appeal to the extensions of the results in Renegar (2014) given in Appendix F. Unlike in the case when K is known, there is no trace constraint imposed and therefore we can find an \mathbf{F} such that for any d , $c_1^{-1} \leq \lambda_{\min}(\mathbf{F}) \leq \lambda_{\max}(\mathbf{F}) \leq c_1$, for some $c_1 \geq 1$. In particular, we can choose

$$\mathbf{F} := \frac{1}{2} \mathbf{I} + \frac{1}{2d} \mathbf{1}\mathbf{1}^T.$$

Clearly, \mathbf{F} is strictly feasible for (4.1). Using the Sherman-Morrison formula, we obtain that

$$\mathbf{F}^{-1} = 2\mathbf{I} - \frac{1}{d} \mathbf{1}\mathbf{1}^T.$$

Furthermore, it is easy to see that $\frac{1}{2} \leq \lambda_{\min}(\mathbf{F}) \leq \lambda_{\max}(\mathbf{F}) \leq 2$. This shows that in the case where K is unknown, we pay only a factor of 4 penalty for \mathbf{I} not being strictly feasible. This is a sharp contrast to the fixed K case, where the penalty is much higher.

4.1 FORCE Dual Step

In order to find a dual certificate, it is useful to determine the form optimal solutions to (4.2) take. Lemma 4.1 characterizes all primal, dual optimal pairs for (4.1). This is the analogue of Theorem 4 from Iguchi et al. (2015) for the case when K is not known beforehand.

Lemma 4.1. The following are equivalent: (1) \mathbf{B}^* is an optimal solution to (2.1), (2) every solution to (2.2) satisfies $y_{a,b} = 0$ for $a, b \in G_i^*$ and $\mathbf{Q}_{G_i^*, G_i^*} \mathbf{1} = 0$ for all i , and (3) every solution to (2.2) satisfies $\mathbf{y}_{G_i^*} = \mathbf{L}_{G_i^*, G_i^*}^{-1} (-\mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - \widehat{\kappa} \mathbf{1})$.

Proof. The proof of Lemma 4.1 follows from complementary slackness and by re-arranging a system of linear equations. For more details, we direct the reader to Iguchi et al. (2015). \square

Now, observe that in (4.2), $\widehat{\kappa}$ plays the same role as y_T in (2.2). Thus, the results and intuition regarding the dual construction hold, with the only difference being that now we do not perform a binary search over y_T . Instead we simply invert a linear system and check if the construction

succeeds or fails. The computational complexity is thus just the cost of solving one $d + 1 \times d + 1$ linear system. The construction is given as

$$\mathbf{y}_{G_i^*}(\mathbf{D}) = \mathbf{L}_{G_i^*, G_i^*}^{-1}(-\mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - \hat{\kappa} \mathbf{1}) \quad \text{and} \quad y_{a,b}(\mathbf{D}) = \begin{cases} 0, & \text{if } a = b \\ y_a + y_b + D_{a,b}, & \text{o/w.} \end{cases} \quad (4.5)$$

Just as the case when K is known, we can use the explicit dual solution construction (4.5) to certify the optimality of a proposed solution G^* .

4.2 The FORCE Algorithm

We can apply a variant of FORCE – the details are given as Algorithm 4 in Appendix C – to (4.1) by substituting the appropriate update and dual construction operations into Algorithm 1. As before, \mathcal{A}_P is the smoothed scheme applied to (4.1) and now \mathcal{A}_D becomes inverting a linear system as in (4.5) and checking dual feasibility. Choosing \mathcal{A}_R is no longer as clear – to obtain a clustering we must obtain \hat{K} . One choice is to round $\text{tr}(P_{\mathbf{F}}(\mathbf{V}_s))$ to the nearest integer to estimate K . This method is theoretically justified because assuming the relaxation is tight and the certificate exists at the optimal solution, this method correctly estimates the true K and an optimality certificate is found.

FORCE with unknown K has all the same properties as the original FORCE algorithm for the case when K is known, but with an improved running time, as is shown below.

4.3 Theoretical Results

The worst-case bound on the running time of FORCE for (4.1) is given by Theorem 4.2 below.

Theorem 4.2 (Convergence Rate of FORCE for Unknown K). For solving (4.1), FORCE terminates after $\tilde{\mathcal{O}}(d^4 \epsilon^{-1})$ time, giving an ϵ -optimal solution.

Proof of Theorem 4.2. To prove the claim, we first show the rate for Renegar’s “Smoothed Scheme” applied to (4.1). Here, clearly $R \leq \sqrt{2}d$ given the constraints $\mathbf{U} \geq 0$ and $\mathbf{U}\mathbf{1} = \mathbf{1}$. Because each update requires $\mathcal{O}(d^3)$ operations – inverting a linear system and performing an eigen-decomposition – applying the convergence rate of the smoothed scheme (given as Theorem F.6) gives the result. The main claim follows because the dual certificate requires $\mathcal{O}(d^3)$ time to construct. \square

Next we address how to choose $\hat{\kappa}$ in practice. The choice is driven by the following consideration: when does the dual certificate exist and when is the SDP relaxation tight? These questions are intimately connected, and so similar to Bunea et al. (2016) we choose

$$\hat{\kappa} := 4\|\hat{\mathbf{\Gamma}}\|_{\infty} \left(\frac{d}{n} + \sqrt{\frac{d}{n}} \right)$$

for variable clustering in G -Latent models when K is unknown. As is made clear below, the choice of constant in $\hat{\kappa}$ could be altered, but we do not explore whether or not some other choice is preferable. Importantly $\hat{\kappa}$ is data-driven in the sense that its selection requires *no knowledge* of the parameters of the generating distribution.

Theorem 4.3 (FORCE Certificate Exists with High Probability for K Unknown). Consider the variable clustering setting under the G -Latent model and assume $\log d \leq c_0 n$. If $\widehat{\kappa} = 4\|\widehat{\mathbf{\Gamma}}\|_\infty \left(d/n + \sqrt{d/n} \right)$, there exist universal constants c_1, c_2 and c_3 such that if

$$\Delta \mathbf{C}^* \geq c_1 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \sqrt{\frac{\log d}{nm^2}} + \sqrt{\frac{d}{nm^2}} + \frac{d}{nm} \right) + c_2 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right),$$

then with probability at least $1 - c_3/d$, the FORCE Dual Certificate exists at G^* , where $\sigma_1 = \max_i C_{i,i}^*$ and $\sigma_2 = \left(\|\mathbf{\Gamma}^*\|_\infty \max_i C_{i,i}^* \right)^{1/2}$.

4.4 Discussion

The additional cost of constraining K to be fixed is imposed directly by the trace constraint. It is somewhat surprising that we should obtain a significantly better worst-case complexity bound when we have *less information* about the structure of the problem at hand. Because FORCE appears to scale better with d in practice than the worst case analysis suggests, it is possible that the analysis of FORCE when K is known can be improved via a different choice of \mathbf{F} , however this direction is left to future work. (4.1) can be applied to data clustering, and we suspect the FORCE algorithm may have strong theoretical properties in that setting when K is unknown, but that analysis is beyond the scope of this work.

5 Numerical Results and Discussion

5.1 Evaluating the FORCE Primal Step

First, we evaluate the primal step of the FORCE algorithm only, with no dual certificate search. The goal is to demonstrate how much the faster the FORCE algorithm is in solving (2.1) and (4.1) than other existing methods. The experimental setup varies d from 20 to 180 in increments of 10. Three levels of K are chosen $K = \mathcal{O}(\log d)$, $K = \mathcal{O}(\sqrt{d})$, and $K = \mathcal{O}(d)$ and all groups are the same size. $2n$ samples are drawn from a randomly generated covariance structure.

For comparison, we test against a MATLAB implementation using the package CVX and the SDPT3 solver. We ran the experiments on Princeton’s Della cluster, and each experimental run was allotted 8 cores on a 2.4 GHz Broadwell compute node. To obtain the running times shown in the plot, each experiment is repeated 100 times and the average running time is the one reported. For this experiment, the stopping criterion for FORCE is when it shows no further improvement in objective function for at least 100 updates.

Figure 1 shows the relative running times of both implementations with different choices for K . Clearly the asymptotic performance is much better than a standard SDP solver. Interestingly, even though our theory suggests that for $K = \mathcal{O}(d)$, fewer iterations should be needed to obtain convergence, we find in practice that this is not true. Understanding this phenomenon better is an interesting direction of future inquiry.




Figure 1: Comparison of FORCE, for both K known and K unknown, versus a MATLAB implementation using the SDPT3 solver. Solid lines correspond to $K = \mathcal{O}(\log d)$, dashed lines to $K = \mathcal{O}(\sqrt{d})$, and dotted lines to $K = \mathcal{O}(d)$.

5.2 Evaluating FORCE

To distinguish FORCE from other clustering algorithms, several high-level criteria that can be used to evaluate an algorithm for solving (2.1) are (1) does it provide a certificate of *exact* optimality in most cases and (2) in the worst case does it converge “quickly” to an ϵ -optimal solution. FORCE meets both these criteria from a theoretical perspective. To verify in practice, we evaluate (1) how often is a provably optimal solution found and (2) is the exact recovery rate of the clustering from running \mathcal{K} on $P_{\mathbf{F}}(\mathbf{Z}_T)$ higher than on \mathbf{D} ? For notational simplicity, we denote by \mathcal{K} the K -means algorithm, which takes input \mathbf{D} and returns a clustering of the rows. By \mathcal{KB} we denote an algorithm that takes input (\mathbf{D}, N) and runs the K -means algorithm N times and returns the best clustering found in terms of objective value. These two algorithms are used as a baseline against which to measure FORCE.

The simulations described in this section demonstrate the effectiveness of FORCE for variable clustering in G -Block Models. In these simulations the following data generating setup is used $d = 400$, $m = 3$, $K = 15$, and $n = 400$. 30 distinct levels of $\Delta(\mathbf{C}^*)$ were chosen. For each level, we run 100 experiments. For each experiment, we generate a new random covariance structure \mathbf{C}^* and $\mathbf{\Gamma}^*$ from which $n = 400$ samples are drawn. All figures referenced herein can be found in Appendix D.

As Figure 2 shows, the percentage of randomly generated instances for which a dual optimality certificate is found approaches 100% rapidly with increasing $\Delta(\mathbf{C}^*)$. We can see a sharp phase transition occurs at the 5th level of $\Delta(\mathbf{C}^*)$ – as expected based on our theoretical results and those in the literature (Bunea et al., 2016). This is the most important metric as applications of variable clustering may require exact recovery of the true variable clusters, not merely “very close” recovery. The other metric we consider is whether or not the first order method we derive is effective in

practice, not just in theory. As can be seen in Figure 2, the final iterate of FORCE produces much better clusterings than either $\widehat{\Sigma}$ or $\widehat{\Sigma} - \widehat{\Gamma}$.

Figure 3 shows how FORCE outperforms simply running K -means many times on either $\widehat{\Sigma}$ or $\widehat{\Sigma} - \widehat{\Gamma}$. Figures 4 and 5 demonstrate that the interior point method used by FORCE does produce an output matrix $B(\mathbf{Z}_T)$ that is “better” than either $\widehat{\Sigma}$ or $\widehat{\Sigma} - \widehat{\Gamma}$ in terms of quality of the resulting clustering solution.

Figure 6 shows several metrics of the performance of the FORCE dual certificate. On the left is a plot of how long (in seconds) it takes FORCE to find a provably optimal solution (for instances where it could find one). The experimental setup for which this plot is generated is the same as above. The code was run on Princeton’s TIGER cluster, which consists of 644 16-core Intel SandyBridge CPU nodes. The code was run on a single node and given access to 8 of the cores. As expected, as $\Delta(\mathbf{C}^*)$ increases it takes less time before the current iterate is sufficiently close to optimal that the optimal clustering is recovered and certified.

On the right in Figure 6 is a comparison of the FORCE certificate versus that in Iguchi et al. (2016). The experimental setup here is different as they provide an implementation for their certificate that is dependent upon access to the data points being clustered (not only upon \mathbf{D}), so we could not directly apply the implementation they provide for their certificate to variable clustering. Thus, we generate samples from the stochastic ball model at varying levels of cluster center separation Δ . Observe that as the phase transition occurs, moving from uncertifiable to certifiable at G^* , the FORCE certificate outperforms the certificate given by Iguchi et al. (2016). In the plot, the red line represents percentage of instances for which some certificate was found for, and the blue and yellow lines show the proportion of instances for which only one of the two certificates was found. This validates the applicability of the FORCE certificate to data clustering as well. This suggests it may be possible to obtain similar theoretical results for certifying data clusters with at most minor modifications to the FORCE certificate.

5.3 Discussion

Our numerical studies clearly indicate the success of FORCE in the variable clustering setting. A direction of future inquiry is to alter either the FORCE certificate or the proof of Theorem 3.3 to obtain the optimal rate, as given in Bunea et al. (2016). We believe that with at most minor modifications, this is possible. Due to the gap in computational complexity between the cases when K is known and K is unknown, we anticipate with further study and perhaps modification to choice of \mathbf{F} , the worst-case bound on running time when K is known can be improved. It also would be of interest to study the properties of FORCE under other generating distributions for variable and data clustering. The FORCE algorithm will be available for use in an upcoming R package named GFORCE.

References

- ARORA, S., HAZAN, E. and KALE, S. (2005). Fast Algorithms for Approximate Semidefinite Programming using the Multiplicative Weights Update Method. In *FOCS*.
- ARTHUR, D. and VASSILVITSKII, S. (2007). k-means++: The Advantages of Careful Seeding. In *SODA*.
- AWASTHI, P. and BANDEIRA, A. S. (2015). Relax, no need to round: integrality of clustering formulations. In *ITCS*.
- AWASTHI, P. and SHEFFET, O. (2012). Improved Spectral-Norm Bounds for Clustering. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*. 37–49.
- BOYD, S. and VANDENBERGHE, L. (2004). *Convex Optimization*. Cambridge University Press.
- BUBECK, S. (2015). Convex Optimization: Algorithms and Complexity. *Foundations and Trends in Machine Learning* **8** 231–357.
- BUNEA, F., GIRAUD, C., ROYER, M. and VERZELEN, N. (2016). PECOK: a convex optimization approach to variable clustering. [arXiv:1606.05100](https://arxiv.org/abs/1606.05100).
- DASGUPTA, S. (2008). The hardness of k -means clustering. Tech. rep.
- DEFAYS, D. (1977). An efficient algorithm for a complete link method. *The Computer Journal* **20** 364–366.
- IGUCHI, T., MIXON, D. G., PETERSON, J. and VILLAR, S. (2015). On the tightness of an SDP relaxation of k-means. [arXiv:1505.04778](https://arxiv.org/abs/1505.04778).
- IGUCHI, T., MIXON, D. G., PETERSON, J. and VILLAR, S. (2016). Probably certifiably correct k-means clustering. *Mathematical Programming* 1–29.
- KUMAR, A. and KANNAN, R. (2010). Clustering with Spectral Norm and the k-means Algorithm. In *FOCS*.
- LLOYD, S. P. (1982). Least Squares Quantization in PCM. *IEEE Transactions on Information Theory* **28** 129–137.
- MAHAJAN, M., NIMBORKAR, P. and VARADARAJAN, K. (2012). The planar k-means problem is NP-hard. *Theoretical Computer Science* **442** 13–21.
- NESTEROV, Y. (2004). *Introductory Lectures on Convex Optimization: A Basic Course*, vol. 87. Springer US.
- NESTEROV, Y. (2005). Smooth minimization of non-smooth functions. *Math. Program., Ser. A* **103** 127–152.

- PENG, J. and WEI, Y. (2007). Approximating K-means-type Clustering via Semidefinite Programming. *SIAM Journal on Optimization* **18** 186–205.
- RENEGAR, J. (2014). Efficient first-order methods for linear programming and semidefinite programming. [arXiv:1409.5832](#).
- RUDELSON, M. and VERSHYNIN, R. (2013). Hanson-Wright Inequality and Sub-Gaussian Concentration. [arXiv:1306.2872](#).
- VERSHYNIN, R. (2011). Introduction to the non-asymptotic analysis of random matrices. [arXiv:1011.3027](#).

A The FORCE Certificate and Tightness of K-means SDP Relaxations

In this section we explore the theoretical properties of the FORCE Certificate proposed in Section 3.2. Because we focus on applicability to variable clustering where \mathbf{D} is a penalized covariance estimator, we intuitively expect that the Large Diagonal Blocks Property (Property 1, see Section 3.2) will be satisfied with high probability. We show that this is true for data generated via a G -Latent Model, and therefore that (1) with high probability FORCE finds a certificate and (2) that for the PECOK estimator (see Section 2), the SDP relaxation is tight with high probability. In some sense, this shows our FORCE algorithm is nearly optimal for clustering variables in the G -Latent model, as under only slightly stronger conditions than when recovery is statistically possible, we can also produce *proof that the recovered clusters are the optimal solution to (1.1)*.

Thus far, we have followed a similar approach to the literature (see Section 1) for certifying K -means clustering solutions, and in this section we perform the final steps by deriving an empirically testable condition from the FORCE certificate (Section 3.2) and showing that under the G -Latent Model, it is satisfied with high probability. This section is organized as follows:

- Section A.1 gives the derivation of a testable condition for the FORCE certificate, and
- Section A.2 demonstrates that under the G -Latent model, the testable condition is satisfied with high probability.

A.1 General Approach to Proving Existence of the FORCE Certificate

Having given the dual construction and the underlying intuition, it remains to analyze when does the construction succeed. By succeed we mean that the output of Algorithm 3 is dual optimal – and thus a certificate for the corresponding primal solution. Given a particular instance of (2.1), we want to understand when does the FORCE Certificate construction succeed (Algorithm 3 gives the procedure explicitly). The two main difficulties in answering this question lie in determining

- how small can y_T be such that all $y_{a,b}$ are nonnegative,
- and how large must y_T be such that $\mathbf{Q}_{\widehat{G}_i, \widehat{G}_i}$ are positive semidefinite.

To approach this, we consider the case when an optimal solution G^* is known and recall the FORCE dual solution for G^* as a function of \mathbf{D} and y_T , which is given by

$$\mathbf{y}_{G_i^*}(\mathbf{D}, y_T) = \mathbf{L}_{G_i^*, G_i^*}^{-1}(-\mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - y_T \mathbf{1}) \quad \text{and} \quad y_{a,b}(\mathbf{D}, y_T) = \begin{cases} 0 & \text{if } a = b \\ y_a + y_b + D_{a,b} & \text{otherwise} \end{cases}$$

We have Lemma A.1 regarding the form of the matrix \mathbf{L} .

Lemma A.1. Let \mathbf{L} be defined as above as in Section 3.2. Then \mathbf{L} is invertible and its inverse is block-diagonal and given as

$$\mathbf{L}_{G_i^*, G_i^*}^{-1} = \frac{1}{|G_i^*|} \mathbf{I} - \frac{1}{2|G_i^*|^2} \mathbf{1}\mathbf{1}^T$$

Furthermore, $\lambda_{\max} \left(\mathbf{L}_{G_i^*, G_i^*}^{-1} \right) = \frac{1}{|G_i^*|}$.

Given our choice of $y_{a,b}$ in the FORCE certificate, it is clear that we must characterize when Property 1 holds – when, for each i , we can choose y_T such that

- (a) for all a and b , $y_{a,b}(\mathbf{D}, y_T) \geq 0$
- (b) and $\mathbf{Q}_i(\mathbf{D}, y_T) := \mathbf{D}_{G_i^*, G_i^*} + \sum_{a \in G_i^*} y_a \mathbf{R}_a + y_T \mathbf{I}$ is positive semidefinite

Importantly, problem (b) requires studying the behavior of points or variables only within the same group, greatly simplifying the analysis. We give the following lemma, Lemma A.2, to characterize the behavior of the minimal eigenvalue of \mathbf{Q}_i . For comparison, this result fulfills a similar role to Theorem 7 in Iguchi et al. (2016), but applies to the FORCE certificate instead.

Lemma A.2. Using the notation above,

$$\lambda_{\min}(\mathbf{Q}_i(\mathbf{D}, y_T)) = y_T + \min\{-y_T, \lambda_{\min}(\mathbf{Q}_i^\perp(\mathbf{D}))\},$$

where

$$\mathbf{Q}_i^\perp(\mathbf{D}) := \frac{1}{|G_i^*|^2} (\mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{1}) \mathbf{1} \mathbf{1}^T - \frac{1}{|G_i^*|} (\mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} + \mathbf{D}_{G_i^*, G_i^*} \mathbf{1} \mathbf{1}^T) + \mathbf{D}_{G_i^*, G_i^*}.$$

Proof. This result is obtained through a series of linear algebraic manipulations to the form of $\mathbf{Q}_i(\mathbf{D}, y_T)$. See Appendix E for a detailed proof. \square

A.2 FORCE Certification under the G-Latent Latent Variable Model

In this section, \mathbf{D} is to be taken as a penalized covariance estimator, in particular the PECOK estimator (see Section 2). In previous sections \mathbf{D} was an arbitrary instance of some clustering problem, either variable *or* data. We show that with high probability, Algorithm 3 succeeds when the within-between group covariance gap, $\Delta \mathbf{C}^*$, is sufficiently large. In an abuse of notation we will write $\mathbf{y}_{G_i^*}(\mathbf{X}, y_T)$ where $\mathbf{X} \in \mathbb{R}^{n \times d}$ is the matrix containing the random sample from the model.

$$\mathbb{E}[\mathbf{y}_{G_i^*}(\mathbf{X}, y_T)] \approx \left(\frac{1}{2} C_{i,i}^* - \frac{1}{2|G_i^*|} y_T \right) \mathbf{1}$$

It follows then that

$$\mathbb{E}[\mathbf{Q}_i(\mathbf{X}, y_T)] \approx y_T \mathbf{I} - \frac{y_T}{|G_i^*|} \mathbf{1} \mathbf{1}^T$$

where the equality is inexact because we do not know the expectation of $\widehat{\mathbf{\Gamma}}$, just its convergence rate. The FORCE construction fails only when y_T cannot be chosen such that there exists a feasible choice of $y_{a,b}$. The main result in this section is the proof of Theorem 3.3, which says that under nearly the same conditions on $\Delta \mathbf{C}^*$ as Bunea et al. (2016), Algorithm 3 produces an optimal dual solution with high probability on input G^* . Phrased differently: with high probability, a dual optimal solution exists such that it is guaranteed to be found by Algorithm 3.

To prove the result, we need to apply Lemma A.2. We denote by \mathcal{E} the event that $\|\widehat{\Gamma} - \Gamma^*\|_\infty \leq c_2 \|\Gamma^*\|_{\max} \sqrt{\frac{\log d}{n}}$, so that we can separate this event from the analysis of Lemma A.3. First we analyze $\mathbf{Q}_i^\perp(\mathbf{X})$ and then we analyze $y_{a,b}$. The analysis is done group by group, and then the theorem is proved by taking a union bound.

Lemma A.3. Assume that $\log d \leq c_0 n$ and that event \mathcal{E} holds. Then,

$$\|\mathbf{Q}_i^\perp(\mathbf{X})\|_2 \leq c_1 \|\Gamma^*\|_\infty \left(\frac{d}{n} + \frac{\log d}{n} + \sqrt{\frac{d}{n}} + \sqrt{\frac{\log d}{n}} \right),$$

with probability at least $1 - \frac{c_2}{d^2}$ for n sufficiently large, where c_0, c_1 and c_2 are absolute constants.

Proof. The proof of this result is obtained using familiar concentration of measure results for sub-Gaussian and sub-exponential random variables. See Appendix E for a detailed proof. \square

All that remains is to analyze $y_{a,b}$. In particular, as described before, in the construction we set $y_{a,b} = D_{a,b} + y_a + y_b$ for $a \in G_i^*$ and $b \in G_j^*$, $i \neq j$. We require the following lemma to complete the proof of the main result.

Lemma A.4. Let i and j be in $[K]$ and $i \neq j$. Define $y'_{a,b}(\mathbf{X}, y_T) := D_{a,b} + y_a + y_b$ for all $a \in G_i^*$ and $b \in G_j^*$. Under the assumption $\log d \leq c_0 n$,

$$\begin{aligned} y'_{a,b} &\geq \frac{1}{2}(C_{i,i}^* + C_{j,j}^* - 2C_{i,j}^*) - \frac{1}{2|G_i^*|} y_T - \frac{1}{2|G_j^*|} y_T \\ &\quad - c_2 \|\Gamma^*\|_\infty \left(\sqrt{\frac{\log d}{n|G_i^*|}} + \frac{\log d}{n|G_i^*|} \right) - c_3 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right). \end{aligned}$$

with probability at least $1 - c_1/d^3$, where $\sigma_1 = \max_i C_{i,i}^*$ and $\sigma_2 = \max\{\max_i C_{i,i}^*, \|\Gamma^*\|_\infty\}$, and c_1, c_2 and c_3 are absolute constants.

Proof. The proof of this result is obtained using familiar concentration of measure results for sub-Gaussian and sub-exponential random variables. See Appendix E for a detailed proof. \square

Now we have all the necessary lemmas to prove the main result.

Proof of Theorem 3.3. For notation, c_i will be used to denote absolute constants. First, assume that event \mathcal{E} occurs and denote by $m = \min_i |G_i^*|$. Next, we select

$$y'_T := \max_i \|\mathbf{Q}_i^\perp(\mathbf{X})\|_2,$$

ensuring that all $\mathbf{Q}_i(\mathbf{X})$ are positive semidefinite. By Lemma A.3 and taking the union bound over all $i \in [K]$,

$$y'_T \leq c_4 \|\Gamma^*\|_\infty \left(\frac{d}{n} + \frac{\log d}{n} + \sqrt{\frac{d}{n}} + \sqrt{\frac{\log d}{n}} \right)$$

with probability at least $1 - c_5/d$.

Furthermore, by taking the union bound over all a and b not in the same group and using Lemma A.4,

$$\begin{aligned} \min y'_{a,b} &\geq \frac{1}{2}(C_{i,i}^* + C_{j,j}^* - 2C_{i,j}^*) - \frac{1}{2m}y_T - \frac{1}{2m}y_T \\ &\quad - c_6 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \frac{\log d}{nm} \right) - c_7 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right) \end{aligned}$$

with probability at least $1 - c_8/d$. Therefore, there exist some absolute constants c_1 , c_2 and c_9 such that if we take $y_T = y'_T$ and

$$\Delta \mathbf{C}^* \geq c_1 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \sqrt{\frac{\log d}{nm^2}} + \sqrt{\frac{d}{nm^2}} + \frac{d}{nm} \right) + c_2 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right),$$

then with probability at least $1 - c_9/d$, $\min_{a,b} y_{a,b} \geq 0$ demonstrating dual feasibility. Taken together with (2.3) and using $\log d \leq c_0 n$, this shows that there exists some absolute constant c_3 such that if

$$\Delta \mathbf{C}^* \geq c_1 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \sqrt{\frac{\log d}{nm^2}} + \sqrt{\frac{d}{nm^2}} + \frac{d}{nm} \right) + c_2 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right)$$

then with probability at least $1 - c_3/d$, y'_T gives a feasible solution to (2.2). Because $y'_T \geq 0$, this solution is on the search path of Algorithm 3, and thus if no other feasible solution is found, then the algorithm will find the solution corresponding to y'_T . This concludes the proof of the theorem. \square

A.3 Extension to Unknown \mathbf{K}

Because of the close similarity of (4.2) to (2.2), the proof of Theorem 4.3 is nearly identical to the proof of Theorem 3.3. Before giving the proof, we note that the constants in Theorem 4.3 are not necessarily the same as those in Theorem 3.3.

Proof of Theorem 4.3. The proof follows the same exact path as that for Theorem 3.3. As before, define

$$y'_T = \max_i \|\mathbf{Q}_i^\perp(\mathbf{X})\|_2,$$

ensuring that all $\mathbf{Q}_i(\mathbf{X})$ are positive semidefinite. Note that in Lemma A.3, we can move all constants in the inequality into the constant c_2/d^2 . Thus by Lemma A.3, and taking the union bound over all $i \in [K]$, we get

$$y'_T \leq 2 \|\mathbf{\Gamma}^*\|_\infty \left(\frac{d}{n} + \sqrt{\frac{d}{n}} \right)$$

with probability at least $1 - c_4/d$. Conditional on the event \mathcal{E} , we see by the triangle inequality that our choice of $\hat{\kappa}$ satisfies $\hat{\kappa} \geq y'_T$.

Next, observe that Lemma A.4 holds if y_T is replaced with $\widehat{\kappa}$. Thus by using Lemma A.4 and taking the union bound over all a and b not in the same group,

$$\begin{aligned} \min y'_{a,b} &\geq \frac{1}{2}(C_{i,i}^* + C_{j,j}^* - 2C_{i,j}^*) - \frac{1}{2m}\widehat{\kappa} - \frac{1}{2m}\widehat{\kappa} \\ &\quad - c_6 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \frac{\log d}{nm} \right) - c_7 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right) \end{aligned}$$

with probability at least $1 - c_8/d$. Therefore, there exists some absolute constants c_1, c_2 and c_9 such that if

$$\Delta \mathbf{C}^* \geq c_1 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \sqrt{\frac{\log d}{nm^2}} + \sqrt{\frac{d}{nm^2}} + \frac{d}{nm} \right) + c_2 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right),$$

then with probability at least $1 - c_9/d$, $\min_{a,b} y_{a,b} \geq 0$ demonstrating dual feasibility. Taken together with (2.3) and using $\log d \leq c_0 n$, this shows that there exists some absolute constant c_3 such that if

$$\Delta \mathbf{C}^* \geq c_1 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{nm}} + \sqrt{\frac{\log d}{nm^2}} + \sqrt{\frac{d}{nm^2}} + \frac{d}{nm} \right) + c_2 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right)$$

then with probability at least $1 - c_3/d$, (4.5) gives a feasible solution to (4.2). This concludes the proof of the theorem. \square

B Miscellaneous Technical Lemmas

B.1 Results for Linear Algebra

Lemma B.1. Let \mathbf{M} be a $d \times d$ real, symmetric matrix of the form

$$\mathbf{M} = a\mathbf{I} + b\mathbf{1}\mathbf{1}^T.$$

where $a, b \in \mathbb{R}$ then \mathbf{M} has eigenvalues $a + b$ with multiplicity 1 and a with multiplicity $d - 1$. If $a, b > 0$, then \mathbf{M} also has the property that

$$\begin{aligned} \mathbf{M}^{1/2} &= \sqrt{a}\mathbf{I} + \frac{\sqrt{a+db} - \sqrt{a}}{d}\mathbf{1}\mathbf{1}^T, \\ \mathbf{M}^{-1} &= \frac{1}{a}\mathbf{I} - \frac{b}{a^2 + abd}\mathbf{1}\mathbf{1}^T, \\ \mathbf{M}^{-1/2} &= \frac{1}{\sqrt{a}}\mathbf{I} - \frac{\sqrt{a+db} - \sqrt{a}}{d\sqrt{a^2 + dab}}\mathbf{1}\mathbf{1}^T. \end{aligned}$$

Proof of Lemma B.1. Using the Sherman-Morrison formula, a matrix of the form $\mathbf{M} = a\mathbf{I} + b\mathbf{1}\mathbf{1}^T$, where $a, b > 0$ has the inverse

$$\mathbf{M}^{-1} = \frac{1}{a}\mathbf{I} - \frac{b}{a^2 + abd}\mathbf{1}\mathbf{1}^T.$$

Because $\mathbf{M} \succ 0$, all eigenvalues are strictly positive and denote by λ_i and q_i the eigenvalues and corresponding eigenvectors. Without loss of generality, let q_i be orthonormal. Then we can write

$\mathbf{M} = \sum_i \lambda_i \mathbf{q}_i \mathbf{q}_i^T$. By the form of \mathbf{M} , clearly $\frac{1}{\sqrt{d}} \mathbf{1}$ is always an eigenvector of \mathbf{M} with eigenvalue $a + db$, so we can take $q_1 = \frac{1}{\sqrt{d}} \mathbf{1}$ and $\lambda_1 = 1$. The remaining q_i span $(\mathbf{1}\mathbf{1}^T)^\perp$ and have corresponding eigenvalues $\lambda_i = a$. Therefore,

$$\mathbf{M}^{1/2} = \frac{\sqrt{a+db}}{\sqrt{d}} \mathbf{1}\mathbf{1}^T + \sum_{i=2}^d \sqrt{a} \mathbf{q}_i \mathbf{q}_i^T.$$

Because this eigen-decomposition is unique, the above gives

$$\mathbf{M}^{1/2} = \sqrt{a} \mathbf{I} + \frac{\sqrt{a+db} - \sqrt{a}}{d} \mathbf{1}\mathbf{1}^T.$$

Using the expression for \mathbf{M}^{-1} given above, it follows that

$$\mathbf{M}^{-1/2} = \frac{1}{\sqrt{a}} \mathbf{I} - \frac{\sqrt{a+db} - \sqrt{a}}{d\sqrt{a^2 + dab}} \mathbf{1}\mathbf{1}^T.$$

□

C Additional Psuedo-code for FORCE Algorithms

Below, additional psuedo-code is given for the FORCE dual certificate construction and the FORCE algorithm when K is unknown.

Algorithm 3 FORCE Optimality Certificate - Binary Search

Input: $\widehat{G} = \{\widehat{G}_1, \dots, \widehat{G}_K\}$, C , \mathbf{D} , K , ϵ

Output: d^* , feasible

Compute d^* corresponding to the primal solution

$L \leftarrow \epsilon$

$U \leftarrow C$

$y_T \leftarrow \text{Null}$

$D \leftarrow \text{Null}$

feasible \leftarrow FALSE

while $\frac{U}{L} > 1 + \epsilon$ **do**

$y_T \leftarrow \frac{U+L}{2}$

$\mathbf{y}_{G_i^*} \leftarrow \mathbf{L}_{G_i^*, G_i^*}^{-1} (-\mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - y_T \mathbf{1})$

$$y_{a,b} \leftarrow \begin{cases} 0 & \text{if } a = b \\ D_{a,b} + y_a + y_b & \text{otherwise} \end{cases}$$

$\lambda \leftarrow \lambda_{\min} \left(\sum_{a=1}^d y_a \mathbf{R}_a + y_T \mathbf{I}^{d \times d} + \mathbf{D} - \sum_{a \leq b} y_{a,b} \mathbf{I}_{a,b} \right)$

if $\lambda \geq 0$ **then**

$U \leftarrow y_T$

$D \leftarrow (y_{a,b}, y_a, y_T)$

else

$L \leftarrow y_T$

end if

end while

if $y_{a,b} \geq 0$ for all a and b in solution candidate D **then**

 feasible \leftarrow TRUE

else

 feasible \leftarrow FALSE

end if

return d^* , feasible

Algorithm 4 First-Order Certifiable Clustering (FORCE) - Unknown K

Input: $0 < \epsilon < 1$, \mathbf{D} , h , \mathbf{U}_0 , \mathbf{F} **Output:** G^* Run Algorithm Renegar's Smoothed Scheme with inputs ϵ , \mathbf{D} , \mathbf{U}_0 , \mathbf{F} **for** Every h^{th} gradient update while running Algorithm Renegar's Smoothed Scheme **do** Denote the current iterate as \mathbf{V}_s Denote by \hat{K} an estimate of K by $\text{tr}(P_{\mathbf{F}}(\mathbf{V}_s))$ (rounded to the nearest integer) Round $P_{\mathbf{F}}(\mathbf{V}_s)$ using Lloyd's Algorithm with $K = \hat{K}$ giving \hat{G}_s Construct \mathbf{B}_s , the partnership matrix corresponding to \hat{G}_s Run Algorithm 3 on \mathbf{B}_s **if** Dual and primal objective values match **then** **return** \hat{G}_s **end if****end for**Denote the output of the Smoothed Scheme as \mathbf{V}_T Estimate \hat{K} by $\text{tr}(P_{\mathbf{F}}(\mathbf{V}_s))$ (rounded to the nearest integer)Run CLINK with $K = \hat{K}$ on $P_{\mathbf{F}}(\mathbf{U}_T)$ to obtain \hat{G}_T **return** \hat{G}_T

D Figures for Numerical Results




Figure 2: On the left is a comparison of \mathcal{K} with inputs $P_{\mathbf{F}}(\mathbf{Z}_T)$, $\hat{\Sigma} - \hat{\Gamma}$, and $\hat{\Sigma}$. The plot on the right shows the percentage of instances where FORCE finds a dual certificate showing exact optimality.




Figure 3: Comparison of FORCE vs \mathcal{KB} . To produce the plot on the left, the purity measure is averaged across all experiments with the same $\Delta(\mathbf{C})$. To produce the plot on the right, the percentage is with respect to all experiments with the same $\Delta(\mathbf{C})$. In the plot, N is a placeholder for the number of times FORCE calls \mathcal{K} as a subroutine.




Figure 4: Comparison of \mathcal{K} with inputs $P_{\mathbf{F}}(\mathbf{Z}_T)$, $\hat{\Sigma} - \hat{\Gamma}$, and $\hat{\Sigma}$. To produce the plot on the left, \mathcal{K} is run 100 times and the average purity measure is then averaged across all experiments with the same $\Delta(\mathbf{C})$. To produce the plot on the right, the exact recovery rate is estimated for each experiment (\mathcal{K} run 100 times) and then averaged across all experiments with the same $\Delta(\mathbf{C})$.




Figure 5: Comparison of \mathcal{K} with inputs $P_{\mathbf{F}}(\mathbf{Z}_T)$, $\hat{\Sigma} - \hat{\Gamma}$, and $\hat{\Sigma}$. To produce the plot, \mathcal{K} is run 100 times and the difference in average purity measure is computed for each experiment. This is then averaged across all experiments with the same $\Delta(\mathbf{C})$.




Figure 6: On the left is a plot of mean and median times until a dual certificate is found by FORCE versus $\Delta(\mathbf{C}^*)$ of the generating distribution. On the right is a comparison between the FORCE certificate and the certificate from [Iguchi et al. \(2016\)](#). It is plotted against Δ , the distance between centers under the stochastic ball model.

E Proof of Results in Appendix A

First we give a proof of the Lemma A.2.

Proof of Lemma A.2. To demonstrate the result, we first find an expression of the minimal eigenvalue of $\mathbf{Q}_i(\mathbf{D}, y_T)$ in terms of y_T and $\mathbf{D}_{G_i^*, G_i^*}$. Then we can apply Lemma A.3 to obtain the result. One way to express the minimum eigenvalue is

$$\operatorname{argmin}_{\mathbf{v} \in \mathcal{S}^{|G_i^*|-1}} \underbrace{\mathbf{v}^T \mathbf{Q}_i(\mathbf{D}, y_T) \mathbf{v}}_{(i)}.$$

Now, for any $\mathbf{v} \in \mathcal{S}^{|G_i^*|-1}$ we can expand (i) as

$$\begin{aligned} (i) &= \sum_{a=1}^{|G_i^*|} \sum_{b=1}^{|G_i^*|} v_a v_b Q_i(\mathbf{D}, y_T)_{a,b} \\ &= \sum_{a=1}^{|G_i^*|} v_a^2 y_T + \sum_{a=1}^{|G_i^*|} \sum_{b=1}^{|G_i^*|} v_a v_b (y_a + y_b) + \sum_{a=1}^{|G_i^*|} \sum_{b=1}^{|G_i^*|} v_a v_b D_{a,b} \\ &= y_T + \underbrace{\mathbf{v}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{v}}_{(ii.a)} + 2 \underbrace{\sum_{a=1}^{|G_i^*|} \sum_{b=1}^{|G_i^*|} v_a v_b y_a}_{(ii.b)}. \end{aligned} \quad (E.1)$$

Via some algebra we obtain

$$(ii.b) = \sum_{a=1}^{|G_i^*|} v_a y_a \sum_{b=1}^{|G_i^*|} v_b = \sum_{a=1}^{|G_i^*|} v_a y_a \mathbf{v}^T \mathbf{1} = \mathbf{v}^T \mathbf{1} \mathbf{y}_{G_i^*}^T \mathbf{v}.$$

From E above we see that the object of interest is now $\mathbf{1} \mathbf{y}_{G_i^*}^T$, a $|G_i^*| \times |G_i^*|$ matrix. Recall that $\mathbf{y}_{G_i^*}^T$ is ultimately a function of y_T and \mathbf{D} . Fortunately, we already have explicit expressions for these quantities. In particular,

$$\begin{aligned} \mathbf{1} \mathbf{y}_{G_i^*}^T &= \mathbf{1} (-\mathbf{1}^T y_T - \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*}) \mathbf{L}_{G_i^*, G_i^*}^{-1} \\ &= -y_T \mathbf{1} \mathbf{1}^T \mathbf{L}_{G_i^*, G_i^*}^{-1} - \mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{L}_{G_i^*, G_i^*}^{-1} \\ &= -y_T \frac{1}{|G_i^*|} \mathbf{1} \mathbf{1}^T + \frac{1}{2|G_i^*|} \mathbf{1} \mathbf{1}^T - \frac{1}{|G_i^*|} \mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} + \frac{1}{2|G_i^*|^2} \mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{1} \mathbf{1}^T \\ &= -\frac{y_T}{2|G_i^*|} \mathbf{1} \mathbf{1}^T - \frac{1}{|G_i^*|} \mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} + \underbrace{\frac{1}{2|G_i^*|^2} \mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{1} \mathbf{1}^T}_{(iii)}. \end{aligned} \quad (E.2)$$

In E.2, observe that (iii) = $\frac{1}{2|G_i^*|^2} (\mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{1}) \mathbf{1} \mathbf{1}^T$. Plugging this back into E.2 gives that

$$\mathbf{1} \mathbf{y}_{G_i^*}^T = \frac{1}{2|G_i^*|^2} (\mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - |G_i^*| y_T) \mathbf{1} \mathbf{1}^T - \frac{1}{|G_i^*|} \mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*}. \quad (E.3)$$

We can substitute E.3 into E, yielding that

$$\begin{aligned} \text{(ii.b)} &= \mathbf{v}^T \left(\frac{1}{|G_i^*|^2} (\mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - |G_i^*| y_T) \mathbf{1} \mathbf{1}^T - \frac{1}{|G_i^*|} \mathbf{1} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} - \frac{1}{|G_i^*|} \mathbf{D}_{G_i^*, G_i^*} \mathbf{1} \mathbf{1}^T \right) \mathbf{v} \\ &= \mathbf{v}^T \left(\mathbf{Q}_i^\perp(\mathbf{D}) - \frac{y_T}{|G_i^*|} \mathbf{1} \mathbf{1}^T - \mathbf{D}_{G_i^*, G_i^*} \right) \mathbf{v}. \end{aligned}$$

Substituting back into (i), we get that

$$\lambda_{\min}(\mathbf{Q}_i(\mathbf{D}, y_T)) = y_T + \lambda_{\min} \left(-\frac{y_T}{|G_i^*|} \mathbf{1} \mathbf{1}^T + \mathbf{Q}_i^\perp(\mathbf{D}) \right)$$

which is nearly the desired result. To proceed, we can see that $\frac{y_T}{|G_i^*|} \mathbf{1} \mathbf{1}^T$ and $\mathbf{Q}_i^\perp(\mathbf{D})$ lie in orthogonal spaces. This is a deterministic statement and does not depend on any particular clustering instance. Indeed, we can check that

$$\mathbf{1}^T \mathbf{Q}_i^\perp(\mathbf{D}) \mathbf{1} = 0$$

This is good, because then their respective eigenspaces are orthogonal giving

$$\lambda_{\min}(\mathbf{Q}_i(\mathbf{D}, y_T)) = y_T + \min\{-y_T, \lambda_{\min}(\mathbf{Q}_i^\perp(\mathbf{D}))\}.$$

□

Proof of Lemma A.3. Step 1: For notation, c_i will be used to denote absolute constants. The first step is to decompose $\mathbf{Q}_i^\perp(\mathbf{X})$. Recall that under the G-Latent model, $\mathbf{D} = -\widehat{\Sigma} + \widehat{\Gamma}$. Substituting that into the expression for $\mathbf{Q}_i^\perp(\mathbf{X})$ gives

$$\begin{aligned} \mathbf{Q}_i^\perp(\mathbf{X}) &= \underbrace{-\frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \widehat{\Sigma}_{G_i^*, G_i^*} \mathbf{1} \right) \mathbf{1} \mathbf{1}^T + \frac{1}{|G_i^*|} \left(\mathbf{1} \mathbf{1}^T \widehat{\Sigma}_{G_i^*, G_i^*} + \widehat{\Sigma}_{G_i^*, G_i^*} \mathbf{1} \mathbf{1}^T \right) - \widehat{\Sigma}_{G_i^*, G_i^*}}_{\text{(i)}} \\ &\quad + \underbrace{\frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \widehat{\Gamma}_{G_i^*, G_i^*} \mathbf{1} \right) \mathbf{1} \mathbf{1}^T - \frac{1}{|G_i^*|} \left(\mathbf{1} \mathbf{1}^T \widehat{\Gamma}_{G_i^*, G_i^*} + \widehat{\Gamma}_{G_i^*, G_i^*} \mathbf{1} \mathbf{1}^T \right) + \widehat{\Gamma}_{G_i^*, G_i^*}}_{\text{(ii)}}. \end{aligned}$$

For (i), we recall that by the definition of the G-Latent model that

$$\widehat{\Sigma}_{G_i^*, G_i^*} = \frac{1}{n} \sum_{j=1}^n \mathbf{X}_{G_i^*}^j \mathbf{X}_{G_i^*}^{jT} = \sum_{j=1}^n (Z_i^j + \mathbf{E}_{G_i^*}^j)(Z_i^j + \mathbf{E}_{G_i^*}^j)^T.$$

Plugging this into (i) and simplifying gives us that

$$\text{(i)} = \frac{1}{n} \sum_{j=1}^n \left(-\frac{\mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \mathbf{1}}{|G_i^*|^2} \mathbf{1} \mathbf{1}^T + \frac{\mathbf{1}^T \mathbf{E}_{G_i^*}^j}{|G_i^*|} \left(\mathbf{1} \mathbf{E}_{G_i^*}^{jT} + \mathbf{E}_{G_i^*}^j \mathbf{1}^T \right) - \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \right).$$

Now we see that, again, the expression for $\mathbf{Q}_i^\perp(\mathbf{X})$ has eight terms. We first show that each concentrates to its mean at the desired rate, and then use the triangle inequality to obtain the final

result. Fortunately, we can subtract the mean for each of the 8 terms to the expression for $\mathbf{Q}_i^\perp(\mathbf{X})$ as the means for (i) are offset by the means for (ii). To give the new decomposition of $\mathbf{Q}_i^\perp(\mathbf{X})$ explicitly,

$$\begin{aligned} \mathbf{Q}_i^\perp(\mathbf{X}) = & - \underbrace{\sum_{j=1}^n \frac{\mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \mathbf{1}}{n|G_i^*|^2} \mathbf{1}\mathbf{1}^T}_{\text{(i).a}} + \underbrace{\sum_{j=1}^n \frac{\mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{1} \mathbf{E}_{G_i^*}^{jT}}{n|G_i^*|}}_{\text{(i).b}} + \underbrace{\sum_{j=1}^n \frac{\mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^j \mathbf{1}^T}{n|G_i^*|}}_{\text{(i).c}} - \underbrace{\frac{1}{n} \sum_{j=1}^n \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT}}_{\text{(i).d}} \\ & + \underbrace{\frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*} \mathbf{1} \right) \mathbf{1}\mathbf{1}^T}_{\text{(ii).a}} - \underbrace{\frac{1}{|G_i^*|} \mathbf{1}\mathbf{1}^T \widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*}}_{\text{(ii).b}} + \underbrace{\frac{1}{|G_i^*|} \widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*} \mathbf{1}\mathbf{1}^T}_{\text{(ii).c}} + \underbrace{\widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*}}_{\text{(ii).d}}. \end{aligned} \quad (\text{E.4})$$

Step 2: For the term (i).a, we can directly apply Lemma E.1. Doing so, it follows immediately that with probability at least $1 - \frac{c_3}{d^2}$

$$\left\| \sum_{j=1}^n \frac{\mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \mathbf{1}}{n|G_i^*|^2} \mathbf{1}\mathbf{1}^T - \frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1} \right) \mathbf{1}\mathbf{1}^T \right\|_2 \leq \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right).$$

For the term (i).c (and so by symmetry (i).b), we observe that it is a rank-1 matrix and has the form $\mathbf{u}\mathbf{v}^T$. For such matrices, $\|\mathbf{u}\mathbf{v}^T\|_2 = \mathbf{u}^T \mathbf{v}$. Therefore, we can apply Lemma E.1 and obtain that with probability at least $1 - \frac{c_4}{d^2}$,

$$\left\| \sum_{j=1}^n \frac{\mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{1}}{n|G_i^*|} \mathbf{E}_{G_i^*}^j \mathbf{1}^T - \frac{1}{|G_i^*|} \mathbf{1}\mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \right\|_2 \leq \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right).$$

Step 3: Now we control the term (i).d, the sample covariance matrix of the errors. We can directly apply Corollary E.3 because $\mathbf{\Gamma}^*$ is diagonal, so we only need rescale the matrix. Thus we obtain that with probability at least $1 - c_5/d^2$

$$\begin{aligned} \left\| \frac{1}{n} \sum_{j=1}^n \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} - \mathbf{\Gamma}_{G_i^*, G_i^*}^* \right\|_2 & \leq \|\mathbf{\Gamma}^*\|_\infty \left(\frac{d}{n} + \frac{\log d}{n} + \frac{2\sqrt{d}}{\sqrt{n}} + \frac{2\sqrt{d \log d}}{n} + \frac{2\sqrt{\log d}}{\sqrt{n}} \right) \\ & \leq c_6 \|\mathbf{\Gamma}^*\|_\infty \left(\frac{d}{n} + \frac{\log d}{n} + \sqrt{\frac{d}{n}} + \sqrt{\frac{\log d}{n}} \right), \end{aligned}$$

where the second inequality follows from using the assumption that $\log d_0 \leq cn$.

Step 4: For the terms in (ii), consider first (ii).a. We see that

$$\left\| \left(\mathbf{1}^T \widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*} \mathbf{1} \right) \mathbf{1}\mathbf{1}^T - \left(\mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1} \right) \mathbf{1}\mathbf{1}^T \right\|_{\max} \leq |G_i^*| \|\widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*} - \mathbf{\Gamma}_{G_i^*, G_i^*}^*\|_{\max}$$

Therefore because we conditioned on \mathcal{E} ,

$$\left\| \frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*} \mathbf{1} \right) \mathbf{1}\mathbf{1}^T - \frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1} \right) \mathbf{1}\mathbf{1}^T \right\|_{\max} \leq \frac{c_7}{|G_i^*|} \|\mathbf{\Gamma}^*\|_\infty \sqrt{\frac{\log d}{n}}$$

Because the matrices above are a multiple of $\mathbf{1}\mathbf{1}^T$, it follows that

$$\left\| \frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \widehat{\Gamma}_{G_i^*, G_i^*} \mathbf{1} \right) \mathbf{1}\mathbf{1}^T - \frac{1}{|G_i^*|^2} \left(\mathbf{1}^T \Gamma_{G_i^*, G_i^*}^* \mathbf{1} \right) \mathbf{1}\mathbf{1}^T \right\|_2 \leq c_7 |\Gamma^*|_\infty \sqrt{\frac{\log d}{n}}$$

Next for (ii).b (and (ii).c by symmetry), we can see that

$$\left\| \frac{1}{|G_i^*|} \mathbf{1}\mathbf{1}^T \widehat{\Gamma}_{G_i^*, G_i^*} - \frac{1}{|G_i^*|} \mathbf{1}\mathbf{1}^T \Gamma_{G_i^*, G_i^*}^* \right\|_2 = \frac{1}{|G_i^*|} \left\| \mathbf{1}\mathbf{1}^T \left(\widehat{\Gamma}_{G_i^*, G_i^*} - \Gamma_{G_i^*, G_i^*}^* \right) \right\|_2. \quad (\text{E.5})$$

But, it is clear that

$$\begin{aligned} \max_{\|\mathbf{v}\|_2=1} \left| \mathbf{v}^T \mathbf{1}\mathbf{1}^T \left(\widehat{\Gamma}_{G_i^*, G_i^*} - \Gamma_{G_i^*, G_i^*}^* \right) \mathbf{v} \right| &\leq \max_{\|\mathbf{v}\|_2=1} \|\mathbf{v}^T\|_2 \|\mathbf{1}\mathbf{1}^T\|_2 \|\widehat{\Gamma}_{G_i^*, G_i^*} - \Gamma_{G_i^*, G_i^*}^*\|_2 \|\mathbf{v}\|_2 \\ &= |G_i^*| \|\widehat{\Gamma}_{G_i^*, G_i^*} - \Gamma_{G_i^*, G_i^*}^*\|_{\max} \end{aligned}$$

where the first inequality follows from three applications of Cauchy-Schwartz, which we can apply as $\widehat{\Gamma}_{G_i^*, G_i^*} - \Gamma_{G_i^*, G_i^*}^*$ is diagonal and we take the absolute values of its diagonal elements. Then by E.5 and \mathcal{E} ,

$$\left\| \frac{1}{|G_i^*|} \mathbf{1}\mathbf{1}^T \widehat{\Gamma}_{G_i^*, G_i^*} - \frac{1}{|G_i^*|} \mathbf{1}\mathbf{1}^T \Gamma_{G_i^*, G_i^*}^* \right\|_2 \leq c_9 |\Gamma^*|_\infty \sqrt{\frac{\log d}{n}}$$

The same result is immediate for (ii).a by (2.3). Therefore by combining the above and applying the triangle inequality to Equation E.4, we get that conditional on event \mathcal{E} then with probability at least $1 - \frac{c_2}{d^2}$

$$\|\mathbf{Q}_i^\perp(\mathbf{X})\|_2 \leq c_1 \|\Gamma^*\|_\infty \left(\frac{d}{n} + \frac{\log d}{n} + \sqrt{\frac{d}{n}} + \sqrt{\frac{\log d}{n}} \right),$$

concluding the proof. \square

Proof of Lemma A.4. For notation, c_i will be used to denote absolute constants. First, under the G-Latent model,

$$y'_{a,b}(\mathbf{X}, y_T) = - \underbrace{\widehat{\Sigma}_{a,b}}_{(i)} + \underbrace{y_a(\mathbf{X}, y_T)}_{(ii)} + \underbrace{y_b(\mathbf{X}, y_T)}_{(iii)}$$

Above, we saw that

$$y_a(\mathbf{X}, y_T) = \frac{1}{2|G_i^*|^2} \mathbf{1}^T \mathbf{D}_{G_i^*, G_i^*} \mathbf{1} - \frac{1}{|G_i^*|} \mathbf{D}_{a, G_i^*} \mathbf{1} - \frac{1}{2|G_i^*|} y_T,$$

and likewise for y_b . Below we denote by $\sigma_1 = \max_i C_{i,i}^*$ and $\sigma_2 = \max\{\max_i C_{i,i}^*, \|\Gamma^*\|_\infty\}$. Following the same decomposition as in Lemma A.3, we get that

$$\begin{aligned} y_a(\mathbf{X}, y_T) &= - \frac{1}{2|G_i^*|^2} \mathbf{1}^T \widehat{\Sigma}_{G_i^*, G_i^*} \mathbf{1} + \frac{1}{2|G_i^*|^2} \mathbf{1}^T \widehat{\Gamma}_{G_i^*, G_i^*} \mathbf{1} + \frac{1}{|G_i^*|} \widehat{\Sigma}_{a, G_i^*} \mathbf{1} - \widehat{\Gamma}_{a,a} - \frac{1}{2|G_i^*|} y_T \\ &= \underbrace{\frac{1}{n} \sum_{l=1}^n \frac{1}{2} (Z_i^l)^2}_{(ii).a} - \underbrace{\frac{1}{2n|G_i^*|^2} \sum_{l=1}^n (\mathbf{1}^T \mathbf{E}_{G_i^*}^l)^2}_{(ii).b} + \underbrace{\frac{1}{n|G_i^*|} \sum_{l=1}^n E_a^l \mathbf{1}^T \mathbf{E}_{G_i^*}^l}_{(ii).c} + \underbrace{\frac{1}{n} \sum_{l=1}^n E_a Z_i^l}_{(ii).d} \\ &\quad + \underbrace{\frac{1}{2|G_i^*|^2} \mathbf{1}^T \widehat{\Gamma}_{G_i^*, G_i^*} \mathbf{1}}_{(ii).e} - \underbrace{\frac{1}{|G_i^*|} \widehat{\Gamma}_{a,a}}_{(ii).f} - \frac{1}{2|G_i^*|} y_T. \end{aligned}$$

As in the proof of Lemma A.3, the means of (ii).b and (ii).c offset the means of (ii).e and (ii).f. To control terms (ii).b and (ii).c, by Lemma E.1 with probability at least $1 - c_4/d^3$,

$$\frac{1}{2n|G_i^*|^2} \sum_{j=1}^n \left(\mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \mathbf{1} - \mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1} \right) \leq \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{n|G_i^*|^2}} + \frac{\log d}{n|G_i^*|} \right).$$

Likewise, by Lemma E.6,

$$\frac{1}{n|G_i^*|} \sum_{i=1}^n \left(E_a \mathbf{E}_{G_i^*}^{jT} \mathbf{1} - \gamma_a^* \right) \geq -\|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{n|G_i^*|}} + \frac{\log d}{n|G_i^*|} \right),$$

with probability at least $1 - c_5/d^3$. From (2.3), we can see that

$$\frac{1}{2|G_i^*|^2} \left(\mathbf{1}^T \widehat{\mathbf{\Gamma}}_{G_i^*, G_i^*} \mathbf{1} - \mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1} \right) \leq \|\mathbf{\Gamma}^*\|_\infty \sqrt{\frac{\log d}{n|G_i^*|}},$$

with probability at least $1 - c_6/d^3$. Similarly from (2.3),

$$\frac{1}{|G_i^*|} \left(\widehat{\Gamma}_{a,a} - \Gamma_{a,a}^* \right) \geq -\|\mathbf{\Gamma}^*\|_\infty \sqrt{\frac{\log d}{n|G_i^*|}},$$

with probability at least $1 - c_6/d^3$. Lastly, term (ii).d can be bounded by using Lemma E.7, which gives that

$$\frac{1}{n} \sum_{l=1}^n E_a Z_l^l \geq -\sigma_2 \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right), \quad (\text{E.6})$$

with probability at least $1 - c_7/d^3$. The same results can be obtained for y_b . For the terms in (i), we expand as before:

$$\widehat{\Sigma}_{a,b} = \underbrace{\frac{1}{n} \sum_{l=1}^n Z_l^l Z_l^l}_{(\text{i}).\text{a}} + \underbrace{\frac{1}{n} \sum_{l=1}^n E_a^l Z_l^l}_{(\text{i}).\text{b}} + \underbrace{\frac{1}{n} \sum_{l=1}^n E_b^l Z_l^l}_{(\text{i}).\text{c}} + \underbrace{\frac{1}{n} \sum_{l=1}^n E_a^l E_b^l}_{(\text{i}).\text{d}}.$$

Terms (i).b and (i).c can be bounded in the same way as (E.6). Term (i).d can be bounded by Lemma E.7, giving that

$$\frac{1}{n} \sum_{l=1}^n E_a^l E_b^l \geq -\|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right),$$

with probability at least $1 - c_8/d^3$. All that remains is to bound the terms (i).a, (ii).a and (iii).a. Fortunately, these correspond to the population quantity $\Delta \mathbf{C}^*$. Observing that this is just a quadratic form of $2n$ -dimensional Gaussian vector, we can applying Lemma E.1. Doing so gives that

$$\frac{1}{2n} \left(\sum_{l=1}^n (Z_l^l)^2 + \sum_{l=1}^n (Z_j^l)^2 - 2 \sum_{l=1}^n Z_l^l Z_j^l \right) \geq \frac{1}{2} (C_{i,i}^* + C_{j,j}^* - C_{i,j}^*) - \sigma_1 \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right)$$

with probability at least $1 - c_1 0/d^3$. Combining all the bounds for (i)-(iii), we can see that, with probability at least $1 - c_1/d^3$

$$\begin{aligned} y'_{a,b} &\geq \frac{1}{2}(C_{i,i}^* + C_{j,j}^* - 2C_{i,j}^*) - \frac{1}{2|G_i^*|}y^T - \frac{1}{2|G_j^*|}y^T \\ &\quad - c_1 \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{\frac{\log d}{n|G_i^*|}} + \frac{\log d}{n|G_i^*|} \right) - c_2 (\sigma_1 + \sigma_2) \left(\sqrt{\frac{\log d}{n}} + \frac{\log d}{n} \right). \end{aligned}$$

□

Lemma E.1. Using the notation and assumptions from previous sections,

$$\left| \sum_{j=1}^n \mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \mathbf{1} - \mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1} \right| \leq \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{|G_i^*|^2 n \log d} + |G_i^*| \log d \right),$$

with probability at least $1 - \frac{c}{d}$, for some absolute constant c .

Proof. To obtain the result, we observe that

$$\sum_{j=1}^n \mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \mathbf{1} - \mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1}$$

is a quadratic form of a $n|G_i^*|$ -dimensional Gaussian random vector with independent entries. In particular, if we choose $\mathbf{M} := (\mathbf{\Gamma}^*)^{1/2} \mathbf{1} \mathbf{1}^T (\mathbf{\Gamma}^*)^{1/2}$ as the matrix in Lemma E.7, we see that $\|\mathbf{M}\|_2 \leq \|\mathbf{\Gamma}^*\|_\infty |G_i^*|$ and $\|\mathbf{M}\|_F \leq \|\mathbf{\Gamma}^*\|_\infty |G_i^*| \sqrt{n}$. Applying the Lemma gives

$$\left| \sum_{j=1}^n \mathbf{1}^T \mathbf{E}_{G_i^*}^j \mathbf{E}_{G_i^*}^{jT} \mathbf{1} - \mathbf{1}^T \mathbf{\Gamma}_{G_i^*, G_i^*}^* \mathbf{1} \right| \leq \|\mathbf{\Gamma}^*\|_\infty \left(\sqrt{|G_i^*|^2 n \log d} + |G_i^*| \log d \right),$$

with probability at least $1 - \frac{c}{d}$ as desired. □

E.1 Results for Random Variables

This section contains results used in Section A.2. The proofs are either given here or a reference is provided. As made clear in Section 2, to show that under a certain generating model the relaxation is tight and the dual certificate exists requires that the largest eigenvalue of a certain matrix concentrates sufficiently quickly. In particular, we are concerned with the rate of concentration in the spectral norm of a sample covariance matrix to its mean: $\|\widehat{\Sigma} - \Sigma^*\|_2$. If we write $\widehat{\Sigma} = \frac{1}{n} \mathbf{X}^T \mathbf{X}$, where \mathbf{X} refers to the $n \times d$ matrix in which the rows are the observations \mathbf{X}_i , we see how such a result is directly applicable to the problem at hand. We repeat the statement of Gordon's Theorem given in Vershynin (2011) below as Proposition E.2. We use the notation from Vershynin (2011) of s_{\min} and s_{\max} to denote the smallest and largest singular values, respectively.

Proposition E.2. Let \mathbf{X} be an $n \times d$ matrix whose entries are independent standard normal random variables. Then

$$\sqrt{n} - \sqrt{d} \leq \mathbb{E}[s_{\min}(\mathbf{X})] \leq \mathbb{E}[s_{\max}(\mathbf{X})] \leq \sqrt{n} + \sqrt{d}$$

Using the result on sub-Gaussian concentration of a Lipschitz function of independent random variables, we immediately obtain the following corollary (also given in [Vershynin \(2011\)](#)).

Corollary E.3. Let \mathbf{X} be an $n \times d$ matrix whose entries are independent standard normal random variables, then for every $t \geq 0$

$$\sqrt{n} - \sqrt{d} - t \leq s_{\min}(\mathbf{X}) \leq s_{\max}(\mathbf{X}) \leq \sqrt{n} + \sqrt{d} + t$$

with probability at least $1 - 2 \exp(-t^2/2)$.

Proof. Observing that the functions s_{\min} and s_{\max} are 1-Lipschitz and using the sub-Gaussian tail bound, the result is immediate from the above. \square

Lemma E.4 (Tail Bound for Sub-Exponential Random Variables). Let X be a sub-exponential random variable with mean μ and parameters α and ν . Then

$$\mathbb{P}(X - \mu \geq t) \leq \begin{cases} \exp(-\frac{t^2}{2\nu^2}) & \text{for } 0 \leq t \leq \frac{\nu^2}{\alpha} \\ \exp(-\frac{t}{2\alpha}) & \text{for } t > \frac{\nu^2}{\alpha} \end{cases}$$

Corollary E.5. Let X be a sub-exponential random variable with mean μ and parameters α and ν . Then

$$\mathbb{P}(X - \mu \geq \nu\sqrt{2t} + 2\alpha t) \leq \exp(-t).$$

Proof. This result follows immediately from [Lemma E.4](#). \square

The following result for quadratic forms of standard multivariate Gaussian random variables can be found in many forms in the literature (for example, [Rudelson and Vershynin \(2013\)](#)).

Lemma E.6 (Hanson-Wright Inequality for Gaussian Random Variables). Let $\mathbf{X} \sim N(0, \mathbf{I})$ be a d -dimensional random vector and let \mathbf{A} be a $d \times d$ matrix in $\mathbb{R}^{d \times d}$. Then

$$\mathbb{P}(|\mathbf{X}^T \mathbf{A} \mathbf{X} - \mathbb{E}[\mathbf{X}^T \mathbf{A} \mathbf{X}]| \geq t) \leq 2 \exp\left(-c \min\left\{\frac{t^2}{\|\mathbf{A}\|_F^2}, \frac{t}{\|\mathbf{A}\|_2}\right\}\right),$$

for some absolute constant c .

In particular, the following corollary is useful.

Corollary E.7. Let $\mathbf{X} \sim N(0, \mathbf{I})$ be a d -dimensional random vector and let \mathbf{A} be a $d \times d$ matrix in $\mathbb{R}^{d \times d}$. Then

$$\mathbb{P}\left(|\mathbf{X}^T \mathbf{A} \mathbf{X} - \mathbb{E}[\mathbf{X}^T \mathbf{A} \mathbf{X}]| \geq \|\mathbf{A}\|_F \sqrt{t} + \|\mathbf{A}\|_2 t\right) \leq 2 \exp(-ct),$$

for some absolute constant c .

F Extension of First-Order SDP Results

This section continues the discussion in Section 2. Because \mathbf{I} is not feasible for (2.1), we need to adapt the approach in Renegar (2014) before using those methods. One way to avoid the $\mathbf{F} \neq \mathbf{I}$ issue, as shown in Renegar (2014), is to instead solve the rotated problem

$$\begin{aligned} & \underset{\mathbf{V}}{\text{maximize}} && \lambda_{\min}(\mathbf{V}) \\ & \text{subject to} && \langle \mathbf{F}^{1/2} \mathbf{A}_i \mathbf{F}^{1/2}, \mathbf{V} \rangle = b_i \text{ for } i = 1, \dots, p \\ & && \langle \mathbf{F}^{1/2} \mathbf{D} \mathbf{F}^{1/2}, \mathbf{V} \rangle = u_0. \end{aligned} \tag{F.1}$$

Rotating the system of constraints is not a satisfactory solution for (2.1) because the easy projection onto $\mathcal{C}_\lambda^\perp$ is lost. Thus we need to carefully analyze the smoothness of the objective function $f_{\mu, \mathbf{F}}$ and obtain similar results as the case when $\mathbf{F} = \mathbf{I}$. This is done in the section below.

F.1 Extension of the Smoothed Scheme to Arbitrary Initial Solutions

For completeness, we give in this section the extension of the results in Renegar (2014) to arbitrary choice of initial feasible solution \mathbf{F} . Similar to the notation in Renegar (2014), we denote the smoothed approximation of $\lambda_{\min, \mathbf{F}}(\mathbf{V})$ as

$$f_{\mu, \mathbf{F}}(\mathbf{V}) = -\mu \log \left(\sum_j \exp \left(-\lambda_j(\mathbf{F}^{-1/2} \mathbf{V} \mathbf{F}^{-1/2}) / \mu \right) \right), \tag{F.2}$$

where λ_j denotes the j^{th} eigenvalue of \mathbf{V} .

Lemma F.1. The function $f_{\mu, \mathbf{F}}(\mathbf{V})$ is $\frac{\|\mathbf{F}^{-1}\|_2^2}{\mu}$ -smooth.

Proof. From Nesterov (2005) we have that

$$f_\mu(\mathbf{V}) = -\mu \log \left(\sum_j \exp \left(-\lambda_j(\mathbf{V}) / \mu \right) \right)$$

is $1/\mu$ -smooth. Denote by $g : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}^{d \times d}$ the mapping $g(\mathbf{V}) = \mathbf{F}^{-1/2} \mathbf{V} \mathbf{F}^{-1/2}$. Using differential notation, we see can obtain that

$$dg(\mathbf{V}) = \mathbf{F}^{-1/2} d\mathbf{V} \mathbf{F}^{-1/2}.$$

By Cauchy invariance, and vectorizing g , we obtain that the Jacobian is $D \text{vec } g(\mathbf{V}) = \mathbf{F}^{-1/2} \otimes \mathbf{F}^{-1/2}$. To simplify the proof, we now view f_μ and $f_{\mu, \mathbf{F}}$ as functions on \mathbb{R}^{d^2} . By the chain rule for the Jacobian,

$$Df_{\mu, \mathbf{F}}(\mathbf{V}) = Df_\mu(g(\mathbf{V})) D \text{vec } g(\mathbf{V}).$$

For any \mathbf{V} and \mathbf{U} in $\mathbb{R}^{d \times d}$, we obtain

$$\begin{aligned}
\|Df_{\mu, \mathbf{F}}(\mathbf{V}) - Df_{\mu, \mathbf{F}}(\mathbf{U})\| &= \|Df_{\mu}(g(\mathbf{V}))D \text{vec } g - Df_{\mu}(g(\mathbf{U}))D \text{vec } g\| \\
&\leq \|D \text{vec } g\|_2 \|Df_{\mu}(g(\mathbf{V})) - Df_{\mu}(g(\mathbf{U}))\| \\
&\leq \frac{\|D \text{vec } g\|_2}{\mu} \|g(\mathbf{V}) - g(\mathbf{U})\| \\
&= \frac{\|D \text{vec } g\|_2}{\mu} \|\mathbf{F}^{-1/2}(\mathbf{V} - \mathbf{U})\mathbf{F}^{-1/2}\| \\
&= \frac{\|D \text{vec } g\|_2}{\mu} \|\mathbf{F}^{-1/2} \otimes \mathbf{F}^{-1/2} \text{vec}(\mathbf{V} - \mathbf{U})\| \\
&\leq \frac{\|\mathbf{F}^{-1/2}\|_2^4}{\mu} \|\mathbf{V} - \mathbf{U}\|,
\end{aligned}$$

proving the result. \square

The smoothed form of (2.5) is

$$\begin{aligned}
&\underset{\mathbf{V}}{\text{maximize}} && f_{\mu, \mathbf{F}}(\mathbf{V}) \\
&\text{subject to} && \langle \mathbf{A}_i, \mathbf{V} \rangle = b_i \text{ for } i = 1, \dots, p \\
&&& \langle \mathbf{D}, \mathbf{V} \rangle = u_0.
\end{aligned} \tag{F.3}$$

The underlying sub-gradient descent method used in Renegar (2014) is from Chapter 3 in Nesterov (2004), adapted to (2.5). The convergence analysis is presented below. We denote the optimal solution to (2.5) as $\mathbf{V}_{u_0}^*$ because the solution is within the level set corresponding to u_0 in the original problem.

Theorem G.2 gives the rate for the accelerated projected sub-gradient method, applied to a smooth objective function. Using Nesterov's acceleration for constrained optimization (Algorithm 8) we can adapt the results in Sections 6 and 7 of Renegar (2014) to the more general problem with arbitrary \mathbf{F} . For (F.3), Algorithm 5 gives more details of Nesterov's acceleration applied to our problem of interest.

Algorithm 5 Nesterov's Accelerated Projected Gradient Descent for (F.3)

Input: $T, \mathbf{U}_1 \in \mathcal{D}, \beta, \{\lambda_t\}$ and $\{\gamma_t\}$

Output: \mathbf{U}_T

```

 $\mathbf{V}_1 \leftarrow \mathbf{U}_1$ 
for  $t \leftarrow 1, \dots, T-1$  do
     $\mathbf{U}_{t+1} = \mathbf{V}_t + \frac{1}{\beta} \mathcal{P}_{\mathcal{C}_{\lambda}^{\perp}}(\nabla f_{\mu, \mathbf{F}}(\mathbf{V}_t))$ 
     $\mathbf{V}_{t+1} = (1 - \gamma_t)\mathbf{U}_{t+1} + \gamma_t\mathbf{U}_t$ 
end for
return  $\mathbf{U}_T$ 

```

In Algorithm 5, $\beta = \frac{\|\mathbf{F}^{-1}\|_2^2}{\mu}$. Notationally, we denote the optimal solution to (F.3) as $\mathbf{V}_{u_0}^*(\mu)$. Theorem F.2 gives the convergence rate.

Theorem F.2 (Analogue to 6.1 in [Renegar \(2014\)](#)). Let $\epsilon' > 0$ and $\mu = \frac{\epsilon'}{2 \log d}$. Applying Algorithm 5 with initial iterate \mathbf{U}_1 satisfying $u_0 = \langle \mathbf{D}, \mathbf{U}_1 \rangle < \langle \mathbf{D}, \mathbf{F} \rangle$ and with

$$T \geq \frac{2\sqrt{\log d} \|\mathbf{F}^{-1}\|_2^2 \|\mathbf{U}_1 - \mathbf{V}_{u_0}^*(\mu)\|_F}{\epsilon'}$$

gives that

$$\lambda_{\min, F}(\mathbf{V}_{u_0}^*) - \lambda_{\min, F}(\mathbf{U}_T) \leq \epsilon'.$$

Proof of Theorem F.2. This follows mainly from [G.2](#) and that

$$\lambda_{\min, F}(\mathbf{U}) - \mu \log d \leq f_{\mu, \mathbf{F}}(\mathbf{U}) \leq \lambda_{\min, F}(\mathbf{U}).$$

□

Corollary F.3 (Analogue to 6.2 in [Renegar \(2014\)](#)). Let $\epsilon' > 0$ and $\mu = \frac{\epsilon'}{2 \log d}$. Applying Algorithm 5 with initial iterate \mathbf{U}_1 satisfying $u_0 = \langle \mathbf{D}, \mathbf{U}_1 \rangle < \langle \mathbf{D}, \mathbf{F} \rangle$ and with

$$T \geq \frac{2\sqrt{\log d} \|\mathbf{F}^{-1}\|_2^2 R}{\epsilon'}$$

gives that

$$\lambda_{\min, F}(\mathbf{V}_{u_0}^*) - \lambda_{\min, F}(\mathbf{U}_T) \leq \epsilon',$$

where

$$R = \max\{\|\mathbf{U} - \mathbf{V}\|_F : \mathbf{U}, \mathbf{V} \text{ are feasible for (2.4) and } \langle \mathbf{D}, \mathbf{U} \rangle \leq \langle \mathbf{D}, \mathbf{F} \rangle, \langle \mathbf{D}, \mathbf{V} \rangle \leq \langle \mathbf{D}, \mathbf{F} \rangle\}.$$

Proof of Corollary F.3. See proof of 6.2 in [Renegar \(2014\)](#). The proof here is the same. The main idea is $\mathbf{V}_{u_0}^*(\mu)$ is feasible for (2.4). □

The Corollary above gives a bound on the solution to (2.5), but what we want is a bound on the solution to 2.1. Clearly, however, this depends on the inputs to the algorithm. This is summarized in the next Corollary.

Corollary F.4 (Analogous to 6.3 in [Renegar \(2014\)](#)). Let $\epsilon' > 0$ and $\mu = \frac{\epsilon'}{6 \log d}$. Assume that

$$\lambda_{\min, F}(\mathbf{U}_1) \geq \frac{1}{6} \text{ and } \frac{\langle \mathbf{D}, \mathbf{F} \rangle - v^*}{\langle \mathbf{D}, \mathbf{F} \rangle - v_0} \leq 3$$

Applying Algorithm 5 with initial iterate \mathbf{U}_1 satisfying $u_0 = \langle \mathbf{D}, \mathbf{U}_1 \rangle < \langle \mathbf{D}, \mathbf{F} \rangle$ and with

$$T \geq \frac{2\sqrt{\log d} \|\mathbf{F}^{-1}\|_2^2 R}{\epsilon}$$

gives that

$$\frac{\langle \mathbf{D}, P_{\mathbf{F}}(\mathbf{U}_T) \rangle - u^*}{\langle \mathbf{D}, \mathbf{F} \rangle - u^*} \leq \epsilon,$$

where

$$R = \max\{\|\mathbf{U} - \mathbf{V}\|_F : \mathbf{U}, \mathbf{V} \text{ are feasible for (2.4) and } \langle \mathbf{D}, \mathbf{U} \rangle \leq \langle \mathbf{D}, \mathbf{F} \rangle, \langle \mathbf{D}, \mathbf{V} \rangle \leq \langle \mathbf{D}, \mathbf{F} \rangle\}.$$

Proof of Corollary F.4. We can apply Corollary F.3 to get the result. \square

From F.4 it is clear that if we can find an initial iterate satisfying a certain closeness to optimality, then we are closer to an algorithm that does not require knowledge of the optimal value as input. This can be accomplished using Algorithm 6 and Algorithm 7. Lemma F.5 establishes the required conditions and gives the rate for Algorithm 6.

Algorithm 6 Smoothed Subscheme for (F.3) (Renegar, 2014)

Input: $\epsilon, \mathbf{U}_0 \in \mathcal{C}$ such that $\langle \mathbf{D}, \mathbf{U}_0 \rangle < \langle \mathbf{D}, \mathbf{F} \rangle$ and $\lambda_{\min, F}(\mathbf{U}_0) = \frac{1}{6}$

Output: \mathbf{U}_L such that $\lambda_{\min, F}(\mathbf{U}_L) = \frac{1}{6}$ and $\frac{\langle \mathbf{D}, \mathbf{F} \rangle - u^*}{\langle \mathbf{D}, \mathbf{F} \rangle - u_L} \leq 3$

$l \leftarrow 0$ (Outer Iterations Counter)

$\mu \leftarrow \frac{1}{6 \log d}$

$T \leftarrow 2\sqrt{\log d} \|\mathbf{F}^{-1}\|_2^2 R$

$u_0 = \langle \mathbf{D}, \mathbf{U}_0 \rangle$

done \leftarrow FALSE

while !done **do**

 Apply Algorithm 5 to (F.3) on level set corresponding to u_l and inputs T, \mathbf{U}_l . Denote the output by \mathbf{V}_l .

if $\lambda_{\min, F}(\mathbf{U}_{l+1}) \leq \frac{1}{3}$ **then**

 done \leftarrow TRUE

else

$\mathbf{U}_{l+1} \leftarrow \mathbf{F} + \frac{5}{6} \frac{1}{1 - \lambda_{\min, F}(\mathbf{V}_l)} (\mathbf{V}_l - \mathbf{F})$

$u_{l+1} = \langle \mathbf{D}, \mathbf{U}_{l+1} \rangle$

$l \leftarrow l + 1$

end if

end while

$\mathbf{V}_L = \mathbf{V}_l$

return \mathbf{V}_L

Algorithm 7 Smoothed Scheme for (F.3) (Renegar, 2014)

Input: $0 < \epsilon < 1$ and \mathbf{U}_0 such that $\langle \mathbf{D}, \mathbf{U}_0 \rangle < \langle \mathbf{D}, \mathbf{F} \rangle$ and $\lambda_{\min, F}(\mathbf{U}_0) = \frac{1}{6}$ and \mathbf{U}_0 feasible for (2.4).

Output: $P_{\mathbf{F}}(\mathbf{V})$

 Apply Algorithm 6 with input \mathbf{U}_0 . Let \mathbf{U}_1 denote its output.

$T \leftarrow \lceil \frac{2\sqrt{\log d} \|\mathbf{F}^{-1}\|_2^2 R}{\epsilon} \rceil$

$\mu \leftarrow \frac{\epsilon}{6 \log d}$

 Apply Algorithm 5 with inputs T, \mathbf{U}_1, μ on (F.3) with level set u_1 . Denote the output by \mathbf{V} .

return $P_{\mathbf{F}}(\mathbf{V})$

Lemma F.5 (Analogue to Proposition 7.1 [Renegar \(2014\)](#)). Assuming inputs as stated, Algorithm 6 terminates with a matrix \mathbf{U}_L which is feasible for (2.4) and satisfies

$$\lambda_{\min, F}(\mathbf{U}_L) = \frac{1}{6}, \frac{\langle \mathbf{D}, \mathbf{F} \rangle - u^*}{\langle \mathbf{D}, \mathbf{F} \rangle - \langle \mathbf{D}, \mathbf{U}_L \rangle} \leq 3.$$

Furthermore, the number of outer iterations L , is bounded by

$$L \leq \log_{5/4} \left(\frac{\langle \mathbf{D}, \mathbf{F} \rangle - u^*}{\langle \mathbf{D}, \mathbf{F} \rangle - u_0} \right),$$

where $u_0 = \langle \mathbf{D}, \mathbf{U}_0 \rangle$.

Proof of Lemma F.5. See the proof of Proposition 7.1. The rate from [Bubeck \(2015\)](#) can be used in place of that from [Nesterov \(2004\)](#). \square

Theorem F.6 (Analogue to Theorem 7.2 [Renegar \(2014\)](#)). Assuming inputs as stated, Algorithm 7 terminates with a matrix \mathbf{U} which is feasible for (2.4) and satisfies

$$\frac{\langle \mathbf{D}, \mathbf{U} \rangle - u^*}{\langle \mathbf{D}, \mathbf{F} \rangle - u^*} \leq \epsilon.$$

Furthermore, the total number of iterations of Algorithm 5 is bounded by

$$2R \|\mathbf{F}^{-1}\|_2^2 \sqrt{\log d} \left(\frac{1}{\epsilon} + \log_{5/4} \left(\frac{\langle \mathbf{D}, \mathbf{F} \rangle - u^*}{\langle \mathbf{D}, \mathbf{F} \rangle - u_0} \right) \right),$$

where $u_0 = \langle \mathbf{D}, \mathbf{U}_0 \rangle$.

Proof of Theorem F.6. Follows from F.5. \square

G Accelerated Projected Gradient Descent

In this section we give, for completeness, a proof of Nesterov's acceleration for smooth, constrained optimization problems. The algorithm is summarized as Algorithm 8. The problem is phrased as a minimization

$$x \in \operatorname{argmin}_{x \in \mathcal{C}} f(x) \tag{G.1}$$

for some β -smooth, convex $f(x)$, Algorithm 8 gives Nesterov's accelerated projected gradient descent over a convex set \mathcal{C} . Following [Bubeck \(2015\)](#) we can define the auxiliary sequences $\{\lambda_t\}$ and $\{\gamma_t\}$.

$$\lambda_0 = 0 \quad \text{and} \quad \lambda_{t+1} = \frac{1 + \sqrt{1 + 4\lambda_t^2}}{2} \quad \text{and} \quad \gamma_t = \frac{1 - \lambda_t}{\lambda_{t+1}}. \tag{G.2}$$

Before the proof, we require Lemma G.1, characterizing β -smoothness in a way that is helpful.

Lemma G.1. Consider any x_t and y in a convex set \mathcal{C} . Let α be the gradient update step-size and let $z_{t+1} = \Pi_{\mathcal{C}}(x_{t+1} - \alpha \nabla f(x_t))$. Then,

$$f(z_{t+1}) - f(y) \leq g^\perp(x_t)^T(x_t - y) - \frac{\alpha}{2} \|g^\perp(x_t)\|_2^2.$$

Algorithm 8 Nesterov's Accelerated Projected Gradient Descent for β -smooth f

Input: $T, \mathcal{C}, x_1 \in \mathcal{C}, \beta, \{\lambda_t\}$ and $\{\gamma_t\}$

Output: z_T

```

 $y_1 \leftarrow x_1$ 
 $z_1 \leftarrow x_1$ 
for  $t \leftarrow 1, \dots, T-1$  do
   $y_{t+1} \leftarrow x_t - \frac{1}{\beta} \nabla f(x_t)$ 
   $z_{t+1} = \Pi_{\mathcal{C}}(y_{t+1})$ 
   $x_{t+1} = (1 - \gamma_t)z_{t+1} + \gamma_t z_t$ 
end for
return  $z_T$ 

```

Proof. This is a common result, so we omit the proof. \square

Theorem G.2 (Adapted from 3.12 in Bubeck (2015)). Let f be a convex, β -smooth function and T be the number of iterations. Then Algorithm 8 satisfies

$$f(z_T) - f(x^*) \leq \frac{2\beta \|x_1 - x^*\|^2}{T^2}.$$

Proof of Theorem G.2. This proof mirrors that in Bubeck (2015) for the unconstrained case. Denote by α the step-size and $g^\perp(x_t)$ the orthogonal projection of $\nabla f(x_t)$ onto \mathcal{C}

$$g^\perp(x_t) = \frac{1}{\alpha} (x_t - \Pi_{\mathcal{C}}(x_t - \alpha \nabla f(x_t)))$$

From Lemma G.1,

$$\begin{aligned} f(z_{t+1}) - f(z_t) &\leq g^\perp(x_t)^T (x_t - z_t) - \frac{1}{2\beta} \|g^\perp(x_t)\|_2^2 \\ &= \beta(x_t - z_{t+1})^T (x_t - z_t) - \frac{\beta}{2} \|x_t - z_{t+1}\|_2^2, \end{aligned} \quad (\text{G.3})$$

where the equality follows by substituting in the update step for z_{t+1} . Similarly, we can find that

$$f(z_{t+1}) - f(x^*) \leq \beta(x_t - z_{t+1})^T (x_t - x^*) - \frac{\beta}{2} \|x_t - z_{t+1}\|_2^2. \quad (\text{G.4})$$

Next, denote the distance between the value at the t^{th} iterate and the optimal value by $\delta_t := f(z_t) - f(x^*)$. To bound δ_t , we can multiply both sides of (G.3) by $(\lambda_t - 1)$ and add (G.4) to obtain the relation

$$\lambda_t \delta_{t+1} - (\lambda_t - 1) \delta_t \leq \beta(x_t - z_{t+1})^T (\lambda_t x_t - (\lambda_t - 1)z_t - x^*) - \frac{\beta}{2} \lambda_t \|x_t - z_{t+1}\|_2^2. \quad (\text{G.5})$$

From the definition of λ_t given in (G.2), we can see that $\lambda_t^2 - \lambda_t = \lambda_{t-1}^2$. Using this, we multiply (G.5) by λ_t on both sides, giving

$$\begin{aligned} \lambda_t^2 \delta_{t+1} + 1 - \lambda_{t-1}^2 \delta_t &\leq \frac{\beta}{2} (2\lambda_t(x_t - z_{t+1})^T (\lambda_t x_t - (\lambda_t - 2)z_t - x^*) - \|\lambda_t(z_{t+1} - x_t)\|_2^2) \\ &= \frac{\beta}{2} (\|\lambda_t x_t - (\lambda_t - 1)z_t - x^*\|_2^2 - \|\lambda_t z_{t+1} - (\lambda_t - 1)z_t - x^*\|_2^2). \end{aligned} \quad (\text{G.6})$$

Now, if we multiply the update step for x_t in Algorithm 8 by λ_{t+1} on both sides we obtain the relation

$$\lambda_{t+1}x_{t+1} - (\lambda_{t+1} - 1)z_{t+1} = \lambda_t z_{t+1} - (\lambda_t - 1)z_t. \quad (\text{G.7})$$

We can define $u_t = \lambda_t x_t - (\lambda_t - 1)z_t - x^*$ and substitute this into (G.6) which gives

$$\lambda_t^2 \delta_{t+1} - \lambda_{t-1}^2 \delta_t^2 \leq \frac{\beta}{2} (\|u_t\|_2^2 - \|u_{t+1}\|_2^2).$$

Summing these from 1 to $T - 1$, we see that they telescope, giving

$$\delta_T \leq \frac{\beta}{2\lambda_{T-1}^2} \|x_1 - x^*\|.$$

Lastly, for $T = 2$, clearly $\lambda_{T-1} \geq \frac{T}{2}$. By an inductive argument, we easily obtain that for any T , $\lambda_{T-1} \geq \frac{T}{2}$. Plugging this in gives

$$f(z_T) - f(x^*) \leq \frac{2\beta \|x_1 - x^*\|^2}{T^2},$$

as desired. □