

SOME MORE AMPLITUHEDRA ARE CONTRACTIBLE

PAVLE V. M. BLAGOJEVIĆ, PAVEL GALASHIN, NEVENA PALIĆ, AND GÜNTER M. ZIEGLER

ABSTRACT. The amplituhedra arise as images of the totally nonnegative Grassmannians by projections that are induced by linear maps. They were introduced in Physics by Arkani-Hamed & Trnka (Journal of High Energy Physics, 2014) as model spaces that should provide a better understanding of the scattering amplitudes of quantum field theories. The topology of the amplituhedra has been known only in a few special cases, where they turned out to be homeomorphic to balls. The amplituhedra are special cases of Grassmann polytopes introduced by Lam (Current Developments in Mathematics 2014, Int. Press). In this paper we show that some further amplituhedra are homeomorphic to balls, and that some more Grassmann polytopes and amplituhedra are contractible.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

1.1. Introduction. Let n and k be integers such that $n \geq k \geq 1$. If $\text{Mat}_{k,n}$ denotes the space of all real $k \times n$ matrices of rank k , then the real Grassmannian $G_k(\mathbb{R}^n)$ — the space of all k -dimensional linear subspaces of \mathbb{R}^n — can be defined as the orbit space $G_k(\mathbb{R}^n) = \text{GL}_k \backslash \text{Mat}_{k,n}$. The totally nonnegative part of the Grassmannian is defined quite analogously.

Definition 1.1 (Postnikov [14, Sec. 3]). Let $n \geq k \geq 1$ be integers, let $\text{Mat}_{k,n}^{\geq 0}$ be the space of all real $k \times n$ matrices of rank k all whose maximal minors are nonnegative, and let GL_k^+ denote the group of all real $k \times k$ matrices with positive determinant, which acts freely on $\text{Mat}_{k,n}^{\geq 0}$ by matrix multiplication from the left. The *totally nonnegative Grassmannian* $G_k^{\geq 0}(\mathbb{R}^n)$ is the orbit space $G_k^{\geq 0}(\mathbb{R}^n) = \text{GL}_k^+ \backslash \text{Mat}_{k,n}^{\geq 0}$.

The totally nonnegative Grassmannian was introduced and studied by Postnikov in 2006 [14, Sec. 3], building on works by Lusztig [12] and by Fomin & Zelevinsky [6]. Subsequently, the geometric and combinatorial properties of the totally nonnegative Grassmannian were studied intensively. Rietsch & Williams showed that the totally nonnegative Grassmannian is contractible [16, Thm. 1.1]; an earlier argument by Lusztig [13, Sec. 4.4] can also be adapted to prove the same. Galashin, Karp & Lam [7, Thm. 1.1] proved that $G_k^{\geq 0}(\mathbb{R}^n)$ is indeed homeomorphic to a closed $k(n-k)$ -dimensional ball.

In 2014, the physicists Arkani-Hamed & Trnka [2, Sec. 9] introduced the amplituhedra as certain images of the totally nonnegative Grassmannians. They conjectured that their geometry describes scattering amplitudes in some quantum field theories. For a gentle introduction to amplituhedra in physics and mathematics consult [4]. Shortly after, Lam introduced Grassmann polytopes [11], which generalize amplituhedra.

Postnikov [14, Def. 3.2, Thm. 3.5] defined a CW structure on the totally nonnegative Grassmannian $G_k^{\geq 0}(\mathbb{R}^n)$ such that each cell, also called a *positroid cell*, is indexed by the associated matroid — a positroid — of rank k on n elements, see also [15]. Furthermore, Rietsch & Williams [16] showed that the closures of positroid cells are contractible and that their boundaries are homotopy equivalent to spheres.

Definition 1.2. Let $k \geq 1$, $m \geq 0$ and $n \geq k+m$ be integers, and let Z be a real $(k+m) \times n$ matrix such that the assignment

$$\tilde{Z}(\text{span}(V)) = \text{span}(VZ^\top) \tag{1}$$

induces a map

$$\tilde{Z}: G_k^{\geq 0}(\mathbb{R}^n) \longrightarrow G_k(\mathbb{R}^{k+m}).$$

Here $V \in \text{Mat}_{k,n}^{\geq 0}$, span denotes the row span of a matrix, and Z^\top is the transpose of the matrix Z . The image $\tilde{Z}(\bar{e})$ of a closed positroid cell \bar{e} in the CW decomposition of the nonnegative Grassmannian $G_k^{\geq 0}(\mathbb{R}^n)$ is called a *Grassmann polytope*, denoted by $P_Z(e)$. If e is the maximal cell, which for this CW

Date: January 28, 2019.

The research by Pavle V. M. Blagojević has received funding from the grant ON 174024 of the Serbian Ministry of Education and Science.

The research by Nevena Palić has received funding from DFG via the Berlin Mathematical School.

This material is based on work supported by the National Science Foundation under Grant No. DMS-1440140 during the Fall of 2017, while all authors were in residence at the Mathematical Sciences Research Institute in Berkeley CA.

decomposition means $\bar{e} = G_k^{\geq 0}(\mathbb{R}^n)$, and all $(k+m) \times (k+m)$ minors of the matrix Z are positive, then the Grassmann polytope $P_Z(e)$ is called an *amplituhedron* and is denoted by $\mathcal{A}_{n,k,m}(Z)$.

The previous definition in particular means that if $v_1, \dots, v_k \in \mathbb{R}^n$ are linearly independent row vectors, then

$$\tilde{Z}(\text{span}\{v_1, \dots, v_k\}) = \text{span}\{v_1 Z^\top, \dots, v_k Z^\top\}.$$

The map \tilde{Z} is said to be *well defined* if $\text{span}(VZ^\top)$ is a k -dimensional subspace of \mathbb{R}^{k+m} for every $V \in \text{Mat}_{k,n}^{\geq 0}$. The fact that the map \tilde{Z} is well defined when Z is a matrix with positive maximal minors was established by Arkani-Hamed & Trnka in [2] and by Karp in [9, Thm. 4.2]. Lam [11, Prop. 15.2], however, considers a larger class of matrices Z for which the map \tilde{Z} is still well defined.

The structure of the amplituhedron is known only in a few cases. In the case $m = 0$ all amplituhedra $\mathcal{A}_{n,k,0}(Z)$ are the point $G_k(\mathbb{R}^k)$, whereas when $m = 1$ Karp & Williams [10, Cor. 6.18] have shown that the amplituhedron is homeomorphic to a ball. For $k = 1$ the amplituhedron is a cyclic polytope of dimension m on n vertices [18], and for $n = k + m$ the map Z is a linear isomorphism, and consequently the amplituhedron is homeomorphic to the totally nonnegative Grassmannian $G_k^{\geq 0}(\mathbb{R}^n)$, which is a ball by [7, Thm. 1.1]. Finally, Galashin, Karp & Lam [7, Thm. 1.2] proved that the cyclically symmetric amplituhedra, amplituhedra arising from particularly chosen matrices Z , are homeomorphic to balls whenever m is even. The topology of other Grassmann polytopes is unknown.

1.2. Main results. Our first result gives a family of contractible Grassmann polytopes.

Theorem 1.3. *Let $k \geq 1$ and $m \geq 0$ be integers, and let Z be a real $(k+m) \times (k+m+1)$ matrix such that the map $\tilde{Z}: G_k^{\geq 0}(\mathbb{R}^{k+m+1}) \rightarrow G_k(\mathbb{R}^{k+m})$ is well defined. Then the Grassmann polytope $P_Z(e)$ is contractible for every positroid cell e in the CW decomposition of $G_k^{\geq 0}(\mathbb{R}^{k+m+1})$.*

The proof of Theorem 1.3 relies on classical results of Smale [17, Main Thm.] and Whitehead [19, Thm. 1] combined with the fact that every Grassmann polytope admits a triangulation (as a topological space), see Theorem 1.6.

The following is a consequence of Smale's result [17, Main Thm.].

Theorem 1.4 (Smale). *Let X and Y be path connected, locally compact, separable metric spaces, and in addition let X be locally contractible. Let $f: X \rightarrow Y$ be a continuous surjective proper map, that is, any inverse image of a compact set is compact. If for every $y \in Y$ the inverse image $f^{-1}(\{y\})$ is contractible, then the induced homomorphism*

$$f_\#: \pi_i(X) \rightarrow \pi_i(Y)$$

is an isomorphism for all $i \geq 0$.

Recall that a continuous map $f: X \rightarrow Y$ between topological spaces X and Y is a *weak homotopy equivalence* if the induced map on the path connected components $f_\#: \pi_0(X) \rightarrow \pi_0(Y)$ is bijective, and for every point $x_0 \in X$ and for every integer $n \geq 1$ the induced map $f_\#: \pi_n(X, x_0) \rightarrow \pi_n(Y, f(x_0))$ is an isomorphism.

Theorem 1.5 ([19, Thm. 1]). *Let X and Y be topological spaces that are homotopy equivalent to CW complexes. Then a continuous map $f: X \rightarrow Y$ is a weak homotopy equivalence if and only if it is a homotopy equivalence.*

Since Theorem 1.5 requires that spaces have the homotopy type of a CW complex, the following theorem is a necessary ingredient in the proof of Theorem 1.3.

Theorem 1.6. *Every Grassmann polytope is semi-algebraic as a subset of a Grassmannian. In particular, every Grassmann polytope is homeomorphic to a semi-algebraic subset of some real affine space, and admits a triangulation.*

Note that Theorem 1.6 claims that every Grassmann polytope $P_Z(e)$ can be triangulated in a classical sense, thus there exists a simplicial complex T and a homeomorphism $T \rightarrow P_Z(e)$. A very similar argument to ours was also given by Arkani-Hamed, Bai & Lam in [1, Appendix J].

In order to apply Theorem 1.4 to the map \tilde{Z} , we need to understand its fibers. Thus we prove the following theorem.

Theorem 1.7. *Let $k \geq 1$ and $m \geq 0$ be integers, and let Z be a real $(k+m) \times (k+m+1)$ matrix such that the map \tilde{Z} is well defined. Then for every positroid cell e and for every point $y \in P_Z(e)$, the inverse image $(\tilde{Z}|_{\bar{e}})^{-1}(\{y\}) = \tilde{Z}^{-1}(\{y\}) \cap \bar{e}$ under the restriction map $\tilde{Z}|_{\bar{e}}: \bar{e} \rightarrow P_Z(e)$ is contractible.*

The proof of Theorem 1.7 is postponed to the next section. Here we show that Theorem 1.7 in combination with Theorems 1.4–1.6 implies our main result.

Proof of Theorem 1.3. Let e be a positroid cell in the CW decomposition of $G_k^{\geq 0}(\mathbb{R}^{k+m+1})$. We apply Theorem 1.4 to the map $\tilde{Z} : \bar{e} \rightarrow P_Z(e)$. The spaces \bar{e} and $P_Z(e)$, as well as the map \tilde{Z} , satisfy the assumptions of Theorem 1.4. Furthermore, Theorem 1.7 implies that for every $y \in \bar{e}$, the fiber $\tilde{Z}^{-1}(\{y\})$ is contractible. Thus, from Theorem 1.4 we have that the map \tilde{Z} is a weak homotopy equivalence.

The closed positroid cell \bar{e} is a CW complex. Furthermore, the Grassmann polytope $P_Z(e)$ is a CW complex, by Theorem 1.6. Thus, from Theorem 1.5, we conclude that the map \tilde{Z} is a homotopy equivalence. Hence, the Grassmann polytope $P_Z(e)$ is homotopy equivalent to the closed positroid cell \bar{e} , which is contractible, see [16, Thm. 1.1]. \square

Theorem 1.3 in particular implies that all amplituhedra $\mathcal{A}_{k+m+1,k,m}(Z)$ are contractible. Our next result shows that if in addition m is even, they are homeomorphic to balls.

Theorem 1.8. *Let $k \geq 1$ be an integer, let $m \geq 0$ be an even integer, and let $Z \in \text{Mat}_{k+m,k+m+1}$ be a matrix with all $(k+m) \times (k+m)$ minors positive. Then the amplituhedron $\mathcal{A}_{k+m+1,k,m}(Z)$ induced by the matrix Z is homeomorphic to a km -dimensional ball.*

The proof of Theorem 1.8 is presented in Section 4. We remark that the combinatorics of the amplituhedron in the case $n = k + m + 1$ with m even is identical to that of a cyclic polytope, see [8].

Acknowledgement. The authors thank Rainer Sinn for sharing the knowledge about semi-algebraic sets, to Thomas Lam, whose great observations increased the generality of the results in this paper, and to Steven Karp for helpful comments. We are grateful to the referee for careful reading of our manuscript and for useful suggestions that improved the quality of our paper.

2. PROOF OF THEOREM 1.7

Let $k \geq 1, m \geq 0$ and $n \geq k + m$ be integers and let Z be a real $(k + m) \times n$ matrix such that the map \tilde{Z} is well defined. Since the action of the group GL_k^+ on $\text{Mat}_{k,n}^{\geq 0}$ is free, there is a fibration

$$\text{GL}_k^+ \longrightarrow \text{Mat}_{k,n}^{\geq 0} \longrightarrow G_k^{\geq 0}(\mathbb{R}^n). \quad (2)$$

The matrix Z , as in Definition 1.2, induces a map

$$\begin{aligned} \widehat{Z} : \text{Mat}_{k,n}^{\geq 0} &\longrightarrow \text{Mat}_{k,k+m}, \\ V &\longmapsto VZ^\top, \end{aligned}$$

which is again well defined, see for example [11, Prop. 15.2].

Let e be a positroid cell in the CW decomposition of $G_k^{\geq 0}(\mathbb{R}^n)$, and let $I_e \subseteq \binom{[n]}{k}$ be the family of nonbases (dependent sets) of cardinality k of the matroid that defines the cell e . The maximal minors of a $k \times n$ matrix are indexed by the set $\binom{[n]}{k}$. Denote by $\text{Mat}_{k,n}^{\geq 0}(e)$ the set of all matrices $V \in \text{Mat}_{k,n}^{\geq 0}$ whose minors indexed by elements of I_e are equal to zero. Then every point in $\bar{e} \subseteq G_k^{\geq 0}(\mathbb{R}^n)$ is represented by a matrix in $\text{Mat}_{k,n}^{\geq 0}(e)$, and the row span of every such matrix lies in \bar{e} . In other words, $\bar{e} = \text{GL}_k^+ \setminus \text{Mat}_{k,n}^{\geq 0}(e)$. Thus the restriction of the fibration (2) is a fibration

$$\text{GL}_k^+ \longrightarrow \text{Mat}_{k,n}^{\geq 0}(e) \longrightarrow \bar{e}. \quad (3)$$

Note that if e is the maximal positroid cell, the set $\text{Mat}_{k,n}^{\geq 0}(e)$ is the whole set $\text{Mat}_{k,n}^{\geq 0}$.

Denote by $\widehat{P_Z(e)}$ the image of the set $\text{Mat}_{k,n}^{\geq 0}(e)$ under the map \widehat{Z} . With a usual abuse of notation, we consider maps $\widehat{Z} : \text{Mat}_{k,n}^{\geq 0}(e) \rightarrow \widehat{P_Z(e)}$ and $\tilde{Z} : \bar{e} \rightarrow P_Z(e)$. Then there exists a commutative diagram of spaces and continuous maps

$$\begin{array}{ccc} \text{Mat}_{k,n}^{\geq 0}(e) & \xrightarrow{\widehat{Z}} & \widehat{P_Z(e)} \\ \downarrow & & \downarrow \\ \bar{e} & \xrightarrow{\tilde{Z}} & P_Z(e), \end{array}$$

where vertical maps send any matrix to its row span.

The proof of Theorem 1.7 splits into the following two lemmas.

Lemma 2.1. *Let $k \geq 1$ and $m \geq 0$ be integers, $n = k + m + 1$, and let Z be a real $(k + m) \times n$ matrix such that the map \tilde{Z} is well defined. Then for every positroid cell e in the CW decomposition of $G_k^{\geq 0}(\mathbb{R}^n)$ and for every $W \in \widehat{P}_Z(e)$, the inverse image $\widehat{Z}^{-1}(\{W\}) \subseteq \text{Mat}_{k,n}^{\geq 0}(e)$ is nonempty and convex.*

Proof. The matrix Z induces a linear map

$$\begin{aligned} \mathbb{R}^n &\longrightarrow \mathbb{R}^{k+m} \\ v &\longmapsto vZ^\top, \end{aligned} \tag{4}$$

where $v \in \mathbb{R}^n$ is a row vector. Since $n = k + m + 1$, the kernel of the map (4) is 1-dimensional. Fix a generator $a \in \mathbb{R}^n$ of that kernel.

Choose an arbitrary point $W \in \widehat{P}_Z(e)$, and let U and V be any two points in $\widehat{Z}^{-1}(\{W\})$. Our goal is to show that for every $\lambda \in [0, 1]$ the convex combination $(1 - \lambda)U + \lambda V$ also belongs to $\widehat{Z}^{-1}(\{W\})$.

Since $UZ^\top = VZ^\top = W$, the rows of the matrix $V - U$ belong to $\ker(Z)$. Consequently, there exists a row vector $x \in \mathbb{R}^k$ such that $V - U = x^\top a$, where a is also considered as a row vector. Thus we have to show that for every $\lambda \in [0, 1]$ the convex combination

$$(1 - \lambda)U + \lambda V = U + \lambda x^\top a \tag{5}$$

belongs to the space $\text{Mat}_{k,n}^{\geq 0}(e)$, this means that every $k \times k$ minor of the matrix (5) is nonnegative, and in addition that all the minors of the matrix (5) indexed by the nonbases $I_e \subseteq \binom{[n]}{k}$ of the matroid corresponding to e are equal to zero.

A $k \times k$ submatrix of the matrix (5) is of the form

$$\begin{pmatrix} u_{1i_1} + \lambda x_1 a_{i_1} & \dots & u_{1i_k} + \lambda x_1 a_{i_k} \\ \vdots & & \vdots \\ u_{ki_1} + \lambda x_k a_{i_1} & \dots & u_{ki_k} + \lambda x_k a_{i_k} \end{pmatrix}, \tag{6}$$

where

$$U = \begin{pmatrix} u_{11} & \dots & u_{1n} \\ \vdots & & \vdots \\ u_{k1} & \dots & u_{kn} \end{pmatrix}, \quad x = (x_1 \dots x_k), \quad a = (a_1 \dots a_n),$$

and $1 \leq i_1 < \dots < i_k \leq n$. The matrix (6) can be transformed using row operations into a matrix that contains the variable λ only in one row. Therefore, every $k \times k$ minor of the matrix (5) is a polynomial of degree at most 1 in the variable λ . Since it takes nonnegative values for $\lambda = 0$ and $\lambda = 1$, it is also nonnegative for all $\lambda \in [0, 1]$. Thus for every $\lambda \in [0, 1]$, the point $(1 - \lambda)U + \lambda V$ belongs to $\text{Mat}_{k,n}^{\geq 0}$. Similarly, if $\{i_1, \dots, i_k\}$ is a nonbasis of the matroid corresponding to e , then the determinant of the matrix (6) is zero for $\lambda = 0$ and $\lambda = 1$, so it is a constant zero-polynomial, meaning that the matrix (5) belongs to $\text{Mat}_{k,n}^{\geq 0}(e)$ for every $\lambda \in [0, 1]$. Consequently the set $\widehat{Z}^{-1}(\{W\})$ is convex. \square

Lemma 2.2. *Let $k \geq 1, m \geq 0$ and $n \geq k + m$ be integers. For every positroid cell e and for every $W \in \widehat{P}_Z(e)$, the inverse images*

$$\widehat{Z}^{-1}(\{W\}) \subseteq \text{Mat}_{k,n}^{\geq 0}(e) \subseteq \text{Mat}_{k,n}^{\geq 0} \quad \text{and} \quad \tilde{Z}^{-1}(\{\text{span}(W)\}) \subseteq \bar{e} \subseteq G_k^{\geq 0}(\mathbb{R}^n)$$

are homeomorphic.

Proof. Let $\varphi: \widehat{Z}^{-1}(\{W\}) \rightarrow \tilde{Z}^{-1}(\{\text{span}(W)\})$ be defined by $\varphi(U) = \text{span}(U)$, where $U \in \widehat{Z}^{-1}(\{W\})$, and span denotes the row span. We prove that φ is a homeomorphism.

Clearly, φ is continuous, so it suffices to find a continuous map $\psi: \tilde{Z}^{-1}(\{\text{span}(W)\}) \rightarrow \widehat{Z}^{-1}(\{W\})$ such that $\varphi \circ \psi$ is the identity map on $\tilde{Z}^{-1}(\{\text{span}(W)\})$ and $\psi \circ \varphi$ is the identity map on $\widehat{Z}^{-1}(\{W\})$. Let $L \in \tilde{Z}^{-1}(\{\text{span}(W)\})$. Then there exists a matrix $K \in \text{Mat}_{k,n}^{\geq 0}(e)$ whose rows span the subspace L . Since

$$\text{span}(KZ^\top) = \text{span}(W),$$

there exists a unique $C \in \text{GL}_k$ such that $KZ^\top = CW$. Now define ψ as $\psi(L) = C^{-1}K$. It can be seen using Cauchy–Binet formula that $\det(C) > 0$, thus, $C^{-1}K \in \text{Mat}_{k,n}^{\geq 0}(e)$. Even though we have defined the map ψ using an arbitrarily chosen matrix K such that $\text{span}(K) = L$, it can be checked directly that the definition of ψ does not depend on a choice of K .

In order to prove that the map ψ is continuous, we need to show that the choice of a matrix K can be made continuously on $\tilde{Z}^{-1}(\{\text{span}(W)\})$. The choice of a matrix K is equivalent to the choice of a positively oriented basis for the subspace $L \subseteq \mathbb{R}^n$. Therefore, we need a continuous section of the fiber

bundle (3) restricted to the set $\tilde{Z}^{-1}(\{\text{span}(W)\})$. Since the base space \bar{e} is contractible, the fiber bundle (3) is trivial. In particular, its restriction on $\tilde{Z}^{-1}(\{\text{span}(W)\})$ is also trivial, so it admits a continuous section. Therefore, the bases for elements of $\tilde{Z}^{-1}(\{\text{span}(W)\})$ can be chosen continuously. On the other hand, the matrix C is a solution of the linear system $KZ^\top = CW$, which depends continuously on K , thus it also depends continuously on L .

Lastly,

$$\varphi(\psi(L)) = \varphi(C^{-1}K) = \text{span}(C^{-1}K) = \text{span}(K) = L,$$

holds for every $L \in \tilde{Z}^{-1}(\{\text{span}(W)\})$, and

$$\psi(\varphi(U)) = \psi(\text{span}(U)) = C^{-1}U,$$

for every $U \in \hat{Z}^{-1}(\{W\})$, where C is the unique $k \times k$ matrix such that $W = \hat{Z}(U) = UZ^\top = CW$, hence C is the identity matrix. \square

Finally, Lemma 2.1 and Lemma 2.2 complete the proof of Theorem 1.7.

3. PROOF OF THEOREM 1.6

Let us fix an arbitrary positroid cell e in the CW decomposition of the totally nonnegative Grassmannian $G_k^{\geq 0}(\mathbb{R}^n)$.

Furthermore, let $d = \binom{k+m}{k}$, and consider the Veronese embedding $\nu: \mathbb{RP}^{d-1} \rightarrow \mathbb{R}^{d \times d}$ given by

$$x = (x_1 : \dots : x_d) \mapsto \left(\frac{x_i x_j}{x_1^2 + \dots + x_d^2} \right)_{1 \leq i, j \leq d},$$

where $x = (x_1 : \dots : x_d) \in \mathbb{RP}^{d-1}$. The embedding ν maps every line $x \in \mathbb{RP}^{d-1}$ to the matrix of the projection $\mathbb{R}^d \rightarrow x$. For more details on the Veronese embedding see for example [3, Sec. 3.4.2].

Consider next, with obvious abuse of notation, the continuous map $\nu: \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{R}^{d \times d}$ given by

$$(x_1, \dots, x_d) \mapsto \left(\frac{x_i x_j}{x_1^2 + \dots + x_d^2} \right)_{1 \leq i, j \leq d} \in \mathbb{R}^{d \times d}.$$

In this way, we obtain the commutative diagram of spaces and maps

$$\begin{array}{ccccccc} \text{Mat}_{k,n}^{\geq 0} & \xrightarrow{\hat{Z}} & \text{Mat}_{k,k+m} & \xrightarrow{\gamma} & \mathbb{R}^d \setminus \{0\} & \xrightarrow{\nu} & \mathbb{R}^{d \times d} \\ \downarrow & & \downarrow & & \downarrow \pi & & \downarrow \text{id} \\ G_k^{\geq 0}(\mathbb{R}^n) & \xrightarrow{\tilde{Z}} & G_k(\mathbb{R}^{k+m}) & \xrightarrow{\gamma} & \mathbb{RP}^{d-1} & \xrightarrow{\nu} & \mathbb{R}^{d \times d}, \end{array}$$

where $\gamma: G_k(\mathbb{R}^{k+m}) \rightarrow \mathbb{RP}^{d-1}$ is the Plücker embedding, $\gamma: \text{Mat}_{k,k+m} \rightarrow \mathbb{R}^d \setminus \{0\}$ maps every matrix to the tuple of its $k \times k$ minors, and $\pi: \mathbb{R}^d \setminus \{0\} \rightarrow \mathbb{RP}^{d-1}$ is the quotient map.

The Grassmann polytope $P_Z(e) = \tilde{Z}(\bar{e})$ is embedded into \mathbb{RP}^{d-1} via γ , the projective space \mathbb{RP}^{d-1} is embedded into the Euclidean space $\mathbb{R}^{d \times d}$ via ν , and thus the image $\nu(\gamma(P_Z(e)))$ is homeomorphic to $P_Z(e)$.

First, we prove that the homeomorphic image of the Grassmann polytope $\nu(\gamma(P_Z(e)))$ is semi-algebraic. The commutativity of the diagram above implies that

$$\nu(\gamma(P_Z(e))) = \nu(\pi(\widehat{\gamma(P_Z(e))})) = \nu(\widehat{\gamma(P_Z(e))}) = \nu(\widehat{\gamma(\hat{Z}(\text{Mat}_{k,n}^{\geq 0}(e)))}).$$

The set $\text{Mat}_{k,n}^{\geq 0}(e) \subseteq \mathbb{R}^{k \times n}$ is semi-algebraic, even algebraic. Since the map \hat{Z} is multiplication by a matrix, the set $\widehat{\gamma(P_Z(e))}$ is also semi-algebraic [5, Cor. 2.4(2)]. Furthermore, the map $\gamma: \text{Mat}_{k,k+m} \rightarrow \mathbb{R}^d \setminus \{0\}$ is a restriction of a polynomial map $\mathbb{R}^{k \times (k+m)} \rightarrow \mathbb{R}^d$, and thus $\widehat{\gamma(P_Z(e))} \subseteq \mathbb{R}^d \setminus \{0\}$ is semi-algebraic by [5, Cor. 2.4(2)] as well. Finally, the map $\nu: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d}$ is a regular rational map, and consequently it maps semi-algebraic sets to the semi-algebraic sets, see [3, Prop. 2.2.7] [5, Cor. 2.9(1)]. Hence, we have proved that $\nu(\gamma(P_Z(e)))$ is semi-algebraic in $\mathbb{R}^{d \times d}$, and consequently the Grassmann polytope $P_Z(e)$ is homeomorphic to a semi-algebraic set. In particular, since $P_Z(e)$ is compact and homeomorphic to a semi-algebraic set it admits a triangulation according to [3, Thm. 9.2.1] [5, Thm. 3.11].

Second, notice that we obtained a bit more. The π inverse image of the embedded Grassmann polytope $P_Z(e)$ via γ can be presented as follows

$$\pi^{-1}(\gamma(P_Z(e))) = \gamma(\widehat{\hat{Z}(\text{Mat}_{k,n}^{\geq 0}(e))}) \cup (-\gamma(\widehat{\hat{Z}(\text{Mat}_{k,n}^{\geq 0}(e))})).$$

Since we proved that $\gamma(\widehat{Z}(\text{Mat}_{k,n}^{\geq 0}(e)))$ is semi-algebraic we can conclude that $\pi^{-1}(\gamma(P_Z(e)))$ is also a semi-algebraic subset of \mathbb{R}^d .

Having in mind that every real projective variety is affine, we can define that a subset X of the real projective space \mathbb{RP}^{d-1} is semi-algebraic if, for example, its preimage $\pi^{-1}(X) \subseteq \mathbb{R}^d$, via the defining quotient map $\mathbb{R}^d \rightarrow \mathbb{RP}^{d-1}$, is semi-algebraic.

Thus, we proved that the Grassmann polytope $P_Z(e)$, when embedded in \mathbb{RP}^{d-1} via the Plücker embedding, is a semi-algebraic subset of the real projective space. \square

4. PROOF OF THEOREM 1.8

Let $k \geq 1$, $m \geq 0$ and $n \geq k + m$ be integers, and suppose in addition that m is even. Let $S \in \text{GL}_n$ be given by

$$S(x_1, \dots, x_n) = (x_2, \dots, x_n, (-1)^{k-1}x_1).$$

Denote by $Z_0 \in \text{Mat}_{k+m,n}$ the matrix whose rows are the eigenvectors of the matrix $S + S^\top$ that correspond to the largest $k + m$ eigenvalues. It was shown in [7, Lemma 3.1] that all $(k + m) \times (k + m)$ minors of the matrix Z_0 are positive, thus it defines an amplituhedron $\mathcal{A}_{n,k,m}(Z_0)$, called *cyclically symmetric amplituhedron*. Galashin, Karp & Lam [7, Thm. 1.2] showed that $\mathcal{A}_{n,k,m}(Z_0)$ is homeomorphic to a closed km -dimensional ball whenever the parameter m is even.

We conclude the proof of Theorem 1.8 by showing that the amplituhedra $\mathcal{A}_{n,k,m}(Z)$ and $\mathcal{A}_{n,k,m}(Z_0)$ are homeomorphic.

From [9, Cor. 1.12(ii)] we know that entries of every nonzero vector of $\ker(Z_0)$ and of $\ker(Z)$ are nonzero, and they alternate in sign. Since $n = k + m + 1$, the kernels of matrices Z and Z_0 are 1-dimensional. Let $a = (a_1, \dots, a_n) \in \mathbb{R}^n$ be a generator of the kernel of Z and let $b = (b_1, \dots, b_n) \in \mathbb{R}^n$ be a generator of the kernel of Z_0 (it follows from the cyclic symmetry of Z_0 that $b_i = (-1)^{i-1}$ for $1 \leq i \leq n$, see [7]). Choose them in such a way that a_1 and b_1 have the same sign. Consequently, for every $1 \leq i \leq n$, the entries a_i and b_i have the same sign. Let D be an $n \times n$ diagonal matrix $D = \text{diag}(\frac{a_1}{b_1}, \dots, \frac{a_n}{b_n})$. The matrix ZD has the same kernel as the matrix Z_0 , and since the diagonal entries of the matrix D are positive, all maximal minors of the matrix ZD are positive. The fact that the matrices ZD and Z_0 have the same kernel implies that they have the same row spans, as well. In particular, there exists a matrix $C \in \text{GL}_{k+m}^+$ such that $Z_0 = CZD$.

Multiplication by D on the right gives a homeomorphism $\widehat{D} : \text{Mat}_{k,n}^{\geq 0} \rightarrow \text{Mat}_{k,n}^{\geq 0}$, which induces a homeomorphism $\widetilde{D} : \text{G}_k^{\geq 0}(\mathbb{R}^n) \rightarrow \text{G}_k^{\geq 0}(\mathbb{R}^n)$. Furthermore, multiplication by C^\top on the right gives a homeomorphism $\widehat{C} : \text{Mat}_{k,k+m} \rightarrow \text{Mat}_{k,k+m}$, thus the induced map $\widetilde{C} : \text{G}_k(\mathbb{R}^{k+m}) \rightarrow \text{G}_k(\mathbb{R}^{k+m})$ is also a homeomorphism. Hence, we obtain the commutative diagram of spaces and maps

$$\begin{array}{ccccccc} \text{Mat}_{k,n}^{\geq 0} & \xrightarrow{\widehat{D}} & \text{Mat}_{k,n}^{\geq 0} & \xrightarrow{\widehat{Z}} & \text{Mat}_{k,k+m} & \xrightarrow{\widehat{C}} & \text{Mat}_{k,k+m} \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \text{G}_k^{\geq 0}(\mathbb{R}^n) & \xrightarrow{\widetilde{D}} & \text{G}_k^{\geq 0}(\mathbb{R}^n) & \xrightarrow{\widetilde{Z}} & \text{G}_k(\mathbb{R}^{k+m}) & \xrightarrow{\widetilde{C}} & \text{G}_k(\mathbb{R}^{k+m}). \end{array}$$

The image of the composition $\widetilde{C} \circ \widetilde{Z} \circ \widetilde{D}$ of the maps in the lower row of the diagram is the cyclically symmetric amplituhedron $\mathcal{A}_{n,k,m}(Z_0)$ and the image of the map \widetilde{Z} is the amplituhedron $\mathcal{A}_{n,k,m}(Z)$. Since the maps \widetilde{C} and \widetilde{D} are homeomorphisms, these two amplituhedra are homeomorphic. Finally, the fact that the cyclically symmetric amplituhedron $\mathcal{A}_{n,k,m}(Z_0)$ is homeomorphic to a km -dimensional ball [7, Thm. 1.2], when m is even, concludes the argument that every amplituhedron $\mathcal{A}_{n,k,m}(Z)$ is homeomorphic to a km -dimensional ball whenever $n = k + m + 1$ and m is even. \square

The proof of Theorem 1.8 gives even more. Let us say that two Grassmann polytopes $P_Z(e), P_{Z'}(e') \subseteq \text{G}_k(\mathbb{R}^{k+m})$ are *projectively equivalent* if there exists a matrix $M \in \text{GL}_{k+m}$ such that

$$P_{Z'}(e') = \{\widetilde{M}(x) \mid x \in P_Z(e)\}.$$

Here \widetilde{M} denotes a map $\text{G}_k(\mathbb{R}^{k+m}) \rightarrow \text{G}_k(\mathbb{R}^{k+m})$ induced by the natural action of M on \mathbb{R}^{k+m} . For $k = 1$, this coincides with the standard notion of projective equivalence for polytopes in the projective space \mathbb{RP}^{k+m-1} . The proof of Theorem 1.8 actually shows that for $n = k + m + 1$, any two amplituhedra are projectively equivalent.

REFERENCES

- [1] Nima Arkani-Hamed, Yuntao Bai, and Thomas Lam, *Positive geometries and canonical forms*, Preprint, March 2017, 123 pages, [arXiv:1703.04541](https://arxiv.org/abs/1703.04541).
- [2] Nima Arkani-Hamed and Jaroslav Trnka, *The amplituhedron*, Journal of High Energy Physics **10** (2014), 030.
- [3] Jacek Bochnak, Michel Coste, and Marie-Françoise Roy, *Real algebraic geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 36, Springer-Verlag, Berlin, 1998, Translated from the 1987 French original, Revised by the authors.
- [4] Jacob Bourjaily and Hugh Thomas, *What is the amplituhedron?*, Notices of the AMS **65** (2018), 167–169.
- [5] Michel Coste, *An Introduction to Semialgebraic Geometry*, Université de Rennes, 2002, 78 pages, <http://gcomte.perso.math.cnrs.fr/M2/CosteIntroToSemialGeo.pdf>.
- [6] Sergey Fomin and Andrei Zelevinsky, *Double Bruhat cells and total positivity*, J. Amer. Math. Soc. **12** (1999), 335–380.
- [7] Pavel Galashin, Steven N. Karp, and Thomas Lam, *The totally nonnegative Grassmannian is a ball*, Preprint, March 2018, 19 pages, [arXiv:1707.02010](https://arxiv.org/abs/1707.02010).
- [8] Pavel Galashin and Thomas Lam, *Parity duality for the amplituhedron*, Preprint, May 2018, 52 pages, [arXiv:1805.00600](https://arxiv.org/abs/1805.00600).
- [9] Steven N. Karp, *Sign variation, the Grassmannian, and total positivity*, J. Combin. Theory Ser. A **145** (2017), 308–339.
- [10] Steven N. Karp and Lauren K. Williams, *The $m = 1$ amplituhedron and cyclic hyperplane arrangements*, International Mathematics Research Notices (2017), rnx140.
- [11] Thomas Lam, *Totally nonnegative Grassmannian and Grassmann polytopes*, Current Developments in Mathematics 2014, Int. Press, Somerville, MA, 2016, pp. 51–152.
- [12] George Lusztig, *Total positivity in reductive groups*, Lie theory and geometry, Progr. Math., vol. 123, Birkhäuser Boston, Boston, MA, 1994, pp. 531–568.
- [13] ———, *Introduction to total positivity*, Positivity in Lie theory: open problems, De Gruyter Exp. Math., vol. 26, de Gruyter, Berlin, 1998, pp. 133–145.
- [14] Alexander Postnikov, *Total positivity, Grassmannians, and networks*, Preprint, September 2006, 85 pages; [arXiv:0609764](https://arxiv.org/abs/0609764).
- [15] Alexander Postnikov, David Speyer, and Lauren Williams, *Matching polytopes, toric geometry, and the totally non-negative Grassmannian*, J. Algebraic Combin. **30** (2009), 173–191.
- [16] Konstanze Rietsch and Lauren Williams, *Discrete Morse theory for totally non-negative flag varieties*, Adv. Math. **223** (2010), 1855–1884.
- [17] Stephen Smale, *A Vietoris mapping theorem for homotopy*, Proc. Amer. Math. Soc. **8** (1957), 604–610.
- [18] Bernd Sturmfels, *Totally positive matrices and cyclic polytopes*, Proceedings of the Victoria Conference on Combinatorial Matrix Analysis (Victoria, BC, 1987), vol. 107, 1988, pp. 275–281.
- [19] John H. C. Whitehead, *Combinatorial homotopy I*, Bulletin Amer. Math. Soc. **55** (1949), 213–245.

INSTITUTE OF MATHEMATICS, FREIE UNIVERSITÄT BERLIN, ARNIMALLEE 2, 14195 BERLIN, GERMANY
 MAT. INSTITUT SANU, KNEZ MIHAILOVA 36, 11000 BEOGRAD, SERBIA
E-mail address: blagojevic@math.fu-berlin.de

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE,
 CAMBRIDGE, MA 02139, USA
E-mail address: galashin@mit.edu

INSTITUTE OF MATHEMATICS, FREIE UNIVERSITÄT BERLIN, ARNIMALLEE 2, 14195 BERLIN, GERMANY
E-mail address: palic@math.fu-berlin.de

INSTITUTE OF MATHEMATICS, FREIE UNIVERSITÄT BERLIN, ARNIMALLEE 2, 14195 BERLIN, GERMANY
E-mail address: ziegler@math.fu-berlin.de