COMBINATORIAL CALABI FLOW WITH SURGERY ON SURFACES

XIANG ZHU, XU XU

ABSTRACT. Computing uniformization maps for surfaces has been a challenging problem and has many practical applications. In this paper, we provide a theoretically rigorous algorithm to compute such maps via combinatorial Calabi flow for vertex scaling of polyhedral metrics on surfaces, which is an analogue of the combinatorial Yamabe flow introduced by Luo [37]. To handle the singularies along the combinatorial Calabi flow, we do surgery on the flow by flipping. Using the discrete conformal theory established in [31, 32], we prove that for any initial Euclidean or hyperbolic polyhedral metric on a closed surface, the combinatorial Calabi flow with surgery exists for all time and converges exponentially fast after finite number of surgeries. The convergence is independent of the combinatorial structure of the initial triangulation on the surface.

Mathematics Subject Classification (2010). 53C44, 52B70.

Keywords. Polyhedral metrics; Discrete uniformization; Combinatorial Calabi flow; Surgery by flipping.

1. INTRODUCTION

1.1. Backgrounds and main results. One of the central topics in modern geometry concerns with the canonical metrics on a given manifold, which is related to special geometric structures on manifolds. The flow method is an important approach for such problems. To achieve this goal, Hamilton [34] introduced the Ricci flow and Calabi [5, 6] introduced the Calabi flow. The Ricci flow has been used to solve many longstanding problems in differential geometry, including the Poincaré conjecture [41, 42, 43] and the sphere theorem [4]. The Calabi flow is

$$\frac{\partial g}{\partial t} = \Delta_g K \cdot g \tag{1.1}$$

on a Riemannian surface, where g is the Riemannian metric, K is the Gauss curvature and Δ_g is the Laplace-Beltrami operator of g. The long time existence and convergence of the Calabi flow (1.1) on closed Riemannian surfaces were established in [9, 13]. The readers can also refer to [7, 8, 10] and the references therein for more information on Calabi flow on surfaces.

However, computing special geometric structures on manifolds has been a challenging problem, even for the uniformization maps on surfaces. Different from the smooth approach, one works on polyhedral manifolds to compute geometric structures on manifolds. This idea dates at least back to

Thurston [50], who used the circle packing metric (a type of polyhedral metric) to study hyperbolic structures on three dimensional manifolds. Thurston ([50], Section 13.7) obtained the combinatorial obstruction for the existence of constant combinatorial curvature circle packing metrics. Motivated by Ricci flow on surfaces [11, 35], Chow-Luo [12] introduced the combinatorial surface Ricci flow for Thurston's circle packing metrics on triangulated surfaces, which provides an effective way to compute metrics with prescribed combinatorial curvatures (specially the uniformization maps) and has many applications (see for example [56] and the references therein). Inspire by [12], Ge [16, 17] introduced a combinatorial Calabi flow for Thurston's Euclidean circle packing metrics on triangulated surfaces and proved the longtime existence and convergence of the flow. Then Ge and the second author [24] and Ge-Hua [18] studied the convergence of combinatorial Calabi flow for Thurston's hyperbolic circle packing metrics on triangulated surfaces. Similar to combinatorial Ricci flow, the combinatorial Calabi flow can also be used to compute circle packing metrics with prescribed combinatorial curvatures [57]. Combinatorial Ricci flow and Calabi flow are further used to study α -curvatures of circle packings on triangulated surfaces and sphere packings on 3-dimensional manifolds [23, 25, 26, 27].

To study the discrete conformal geometry of polyhedral metrics on maniflods, Rocek-Williams [47] and Luo [37] introduced a notion of discrete conformality for Euclidean polyhedral metrics (piecewise linear metrics or PL metrics for short) on triangulated surfaces independently, which is now called vertex scaling. Luo [37] further defined the combinatorial Yamabe flow for PL metrics. Based on Bobenko-Pinkall-Springborn's work [2] on vertex scaling and Penner's work [40] on decorated Teichimüller spaces, Gu-Luo-Sun-Wu [32] and Gu-Guo-Luo-Sun-Wu [31] recently proved a discrete uniformization theorem for Euclidean and hyperbolic polyhedral metrics on surfaces respectively, which provides a constructive proof of the classical uniformization theorem on closed surfaces. Combinatorial Yamabe flows with surgery were introduced in [31, 32], where the long-time existence and convergence were proved. The finiteness of surgeries along the combinatorial Yamabe flow was proved by Wu [51]. The combinatorial Yamabe flow with surgery provides an effective algorithm to compute the uniformization maps on surfaces [49]. Unlike the circle packing case [12, 16, 17, 18, 24], the convergence of the combinatorial Yamabe flow with surgery in [31, 32] is independent of the combinatorial structure of the initial triangulation on the surface. Following [37], Ge [16] introduced the combinatorial Calabi flow for vertex scaling of PL metrics on triangulated surfaces and proved the short time existence. In this paper, by doing surgery along the combinatorial Calabi flow by flipping, we prove the combinatorial Calabi flow with surgery for vertex scaling exists for all time and converges as time tends to infinity. One of the main results of the paper is the following theorem on Euclidean combinatorial Calabi flow with surgery.

Theorem 1.1. Suppose (S, V) is a closed connected marked surface and d_0 is any initial piecewise linear metric on (S, V). Then the combinatorial Calabi flow with surgery exists for all time and converges exponentially fast to a piecewise linear metric d^* with constant combinatorial curvature after finite number of surgeries.

Note that the convergence is independent of the combinatorial structure of the triangulation on the surface, which improves the convergence of combinatorial curvature flows for circle packing metrics on surfaces [12, 16, 17, 18, 24]. The combinatorial Calabi flow with surgery gives an effective way to compute polyhedral metrics on a surface with given combinatorial curvatures [48]. Especially, the combinatorial Calabi flow provides an algorithm to compute the uniformization maps on surfaces.

We further define the hyperbolic combinatorial Calabi flow and the corresponding hyperbolic combinatorial Calabi flow with surgery. Please refer to Section 4 for the definitions. The main result for hyperbolic combinatorial Calabi flow with surgery is as follows.

Theorem 1.2. Suppose (S, V) is a closed connected marked surface with $\chi(S) < 0$ and d_0 is any initial piecewise hyperbolic metric on (S, V). Then the hyperbolic combinatorial Calabi flow with surgery exists for all time and converges exponentially fast to a hyperbolic metric d^* on S after finite number of surgeries.

Theorem 1.2 is an analogue of the results obtained in [9, 13] with initial metric given by a piecewise hyperbolic metric, which is a hyperbolic cone metric. In Section 4, we will prove a generalization of Theorem 1.2.

The main idea of the paper comes from reading of [16, 17, 31, 32, 37]. As the main tools used in this paper come from [31, 32], many notations are taken from [31, 32] for consistence.

1.2. Notations and definitions. Here we give some notations and definitions used in the main results. Suppose S is a closed surface and V is a finite subset of S, (S, V) is called a marked surface. A piecewise linear metric on (S, V) is a flat cone metric with cone points contained in V. The combinatorial curvature K_i at $v_i \in V$ is 2π less the cone angle at v_i . Suppose $\mathcal{T} = (V, E, F)$ is a triangulation of (S, V), where V, E, F represent the set of vertices, edges and faces respectively. We use (S, V, \mathcal{T}) to denote a marked surface (S, V) with a fixed triangulation \mathcal{T} . If a map $l : E \to (0, +\infty)$ satisfies that $l_{rs} < l_{rt} + l_{st}$ for $\{r, s, t\} = \{i, j, k\}$, where $\{i, j, k\}$ is any triangle in F, then l determines a piecewise linear metric on (S, V). Given (S, V) with a triangulation \mathcal{T} and a map $l : E \to (0, +\infty)$ determined by a piecewise linear metric d on (S, V), the vertex scaling [37, 47] of d by a function $u : V \to \mathbb{R}$ is defined to be the piecewise linear metric u * d on (S, V) determined by $u * l : E \to (0, +\infty)$ with

$$(u*l)_{ij} := e^{u_i + u_j} l_{ij}, \quad \forall \{ij\} \in E.$$

The function $u: V \to \mathbb{R}$ is called a conformal factor. Note that the vertex scaling of a piecewise linear metric on a surface depends on the triangulation of the surface.

The Calabi flow (1.1) on smooth Riemannian surfaces deforms the metrics conformally. If g is conformal to a background metric g_0 with $g = e^u g_0$, where u is the conformal factor, then the Calabi flow (1.1) translates into an evolution equation

$$\frac{du}{dt} = \Delta_g K$$

of the conformal factor. The Euclidean combinatorial Calabi flow for piecewise linear metrics is defined similarly.

Definition 1.1 ([16] Definition 2.11). Suppose d_0 is a piecewise linear metric on a triangulated surface (S, V, \mathcal{T}) . The Euclidean combinatorial Calabi flow on (S, V, \mathcal{T}) is defined as

$$\begin{cases} \frac{du_i}{dt} = \Delta^{\mathbb{E}, \mathcal{T}} K_i \\ u_i(0) = 0 \end{cases}, \qquad (1.2)$$

where $u: V \to \mathbb{R}$ is the conformal factor and $\Delta^{\mathbb{E},\mathcal{T}}$ is the Euclidean discrete Laplace operator of $u * d_0$ on (S, V, \mathcal{T}) defined as

$$(\Delta^{\mathbb{E},\mathcal{T}}f)_i = \sum_{j;j\sim i} \omega_{ij}^{\mathbb{E},\mathcal{T}}(f_j - f_i), \ \forall f \in \mathbb{R}^V$$
(1.3)

with

$$\omega_{ij}^{\mathbb{E},\mathcal{T}} = \cot \theta_k^{ij} + \cot \theta_l^{ij}.$$

Here θ_i^{jk} is the inner angle at the vertex v_i in a triangle $\triangle ijk \in F$.

The Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$ is the well-known finiteelements Laplace operator. The Euclidean combinatorial Calabi flow (1.2) is defined for a fixed triangulation \mathcal{T} of (S, V) and may develop singularities, including the conformal factor tends to infinity and some triangle degenerates along the flow. To handle the possible singularities along the flow, we do surgery on the flow by flipping, the idea of which comes from [31, 32, 37]. Note that the weight $\omega_{ij}^{\mathbb{E},\mathcal{T}}$ may be negative and $\omega_{ij}^{\mathbb{E},\mathcal{T}} \geq 0$ if and only if $\theta_k^{ij} + \theta_l^{ij} \leq \pi$, which is the locally Delaunay condition of \mathcal{T} on the edge $\{ij\}$ [3]. To ensure that the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$ has good properties, we require the triangulations along the Euclidean combinatorial Calabi flow (1.2) to be Delaunay, which is equivalent to every edge satisfies the locally Delaunay condition [3]. Note that each piecewise linear metric on (S, V) has at least one Delaunay triangulation [1, 3], so this additional condition is reasonable. Along the Euclidean combinatorial Calabi flow (1.2)on (S, V) with a triangulation \mathcal{T} , if \mathcal{T} is Delaunay in $u(t) * d_0$ for $t \in [0, T]$ and not Delaunay in $u(t) * d_0$ for $t \in (T, T + \epsilon)$, $\epsilon > 0$, there exists an edge $\{ij\} \in E$ such that $\theta_k^{ij}(t) + \theta_l^{ij}(t) \leq \pi$ for $t \in [0, T]$ and $\theta_k^{ij}(t) + \theta_l^{ij}(t) > \pi$ for $t \in (T, T + \epsilon)$. We replace the triangulation \mathcal{T} by a new triangulation \mathcal{T}'

at time t = T by replacing two triangles $\triangle ijk$ and $\triangle ijl$ adjacent to $\{ij\}$ by two new triangles $\triangle ikl$ and $\triangle jkl$. This is called a **surgery by flipping** on the triangulation \mathcal{T} , which is an isometry of (S, V) in the piecewise linear metric $u(T) * d_0$. After the surgery at time t = T, we run the Euclidean combinatorial Calabi flow (1.2) on (S, V, \mathcal{T}') with initial metric coming from the Euclidean combinatorial Calabi flow (1.2) on (S, V, \mathcal{T}) at time t = T.

Vertex scaling, combinatorial Calabi flow and surgery by flipping can also defined for piecewise hyperbolic metrics on surfaces. Please refer to Section 2 and Section 4 for more details.

1.3. Organization of the paper. The paper is organized as follows. In Section 2, we give some preliminaries on discrete conformal geometry, including the definitions of polyhedral metrics, vertex scaling, discrete curvature and discrete Laplace operators. In Section 3, we study the Euclidean combinatorial Calabi flow on surfaces and prove a generalization of Theorem 1.1. In Section 4, we study the hyperbolic combinatorial Calabi flow on surfaces and prove a generalization 5, we give some remarks and propose an interesting question.

2. Polyhedral metrics, discrete curvatures and discrete Laplace operators

2.1. **Polyhedral metrics on surfaces.** The definition of piecewise linear metric on marked surfaces has been given in Section 1. Here we extend the definition to polyhedral metrics on surfaces.

Definition 2.1 ([31, 32]). Suppose (S, V) is a marked surface. A piecewise linear (hyperbolic and spherical respectively) metric on (S, V) is a flat (hyperbolic and spherical respectively) cone metric on S whose cone points are contained in V. Piecewise linear, piecewise hyperbolic and piecewise spherical metrics on marked surfaces are all called polyhedral metrics.

In this paper, we concern only about piecewise linear and piecewise hyperbolic metrics. For simplicity, piecewise linear and piecewise hyperbolic metrics are denoted by PL and PH metrics respectively in the following.

A marked surface (S, V) with a triangulation $\mathcal{T} = \{V, E, F\}$ is called a triangulated surface and denoted by (S, V, \mathcal{T}) . In this paper, a function defined on vertices V is regarded as a column vector and n = #V is used to denote the number of vertices. Moreover, all vertices, marked by $v_1, ..., v_n$, are supposed to be ordered one by one and we often write *i* instead of v_i . We use $\{ij\}$ to denote the edge between v_i and v_j in E and use $\triangle ijk$ to denote the face determined by v_i, v_j and v_k in F.

Geometrically, PL metrics on (S, V) are obtained by isometrically gluing Euclidean triangles along their edges so that the cone points are contained in V, which gives a geometric triangulation \mathcal{T} of (S, V) whose simplices are quotients of the simplices in the disjoint union. A triangulation \mathcal{T} of (S, V)is geometric in a PL metric d on (S, V) if each triangle in \mathcal{T} is isometric to

a Euclidean triangle in d. PH metrics on (S, V) are obtained similarly with Euclidean triangles replaced by hyperbolic triangles. A geometrical triangulation of (S, V) with a PL metric is said to be a *Delaunay triangulation* if the sum of two angles facing each edge is at most π . A geometrical triangulation of (S, V) with a PH metric is said to be a *Delaunay triangulation* if for each edge e adjacent to two hyperbolic triangles t and t', the interior of the circumball of t does not contain the vertices of t' when the quadrilateral $t \cup t'$ is lifted to \mathbb{H}^2 . See [1, 3, 31, 32, 36, 45] for further discussions on Delaunay triangulations of surfaces. Note that a polyhedral metric on a marked surface is independent of the triangulations.

Suppose $\mathcal{T} = \{V, E, F\}$ is a geometric triangulation of (S, V) with a PL or PH metric d, then the metric d determines a map

$$d: E \longrightarrow (0, +\infty)$$
$$\{ij\} \mapsto d_{ij} \triangleq d(\{ij\})$$

such that for any topological triangle $\Delta ijk \in F$, the triangle inequalities $d_{ij} < d_{ik} + d_{jk}, d_{ik} < d_{ij} + d_{jk}, d_{jk} < d_{ij} + d_{ik}$ are satisfied. Conversely, given a map $d : E \to (0, +\infty)$ satisfying the triangle inequalities $d_{ij} < d_{ik} + d_{jk}, d_{ik} < d_{ij} + d_{jk}, d_{jk} < d_{ij} + d_{ik}$ for each triangle $\Delta ijk \in F$, the map $d : E \to (0, +\infty)$ uniquely determines a PL metric on (S, V) by the construction of the polyhedral metrics. Then for a triangulated surface (S, V, \mathcal{T}) , the space of PL metrics is given by

$$\mathbb{R}^{E(\mathcal{T})}_{\Delta} = \{ d \in \mathbb{R}^{E(\mathcal{T})}_{>0} | d_{ij}, d_{ik}, d_{jk} \text{ satisfy the triangle} \\ \text{ inequalities for any } \Delta i j k \in F \}.$$

And there is an injective map $\Phi_{\mathcal{T}} : \mathbb{R}^{E(\mathcal{T})}_{\Delta} \longrightarrow T_{PL}(S, V)$ between the subspace $\mathbb{R}^{E(\mathcal{T})}_{\Delta}$ and the Teichimüller space $T_{PL}(S, V)$ of PL metrics on (S, V). Note that $\mathbb{R}^{E(\mathcal{T})}_{\Delta}$ is a proper subset of $\mathbb{R}^{E(\mathcal{T})}_{>0}$.

2.2. **Discrete curvature.** On a marked surface, the well-known combinatorial curvature is defined as follows.

Definition 2.2. Suppose (S, V) is a marked surface with a polyhedral metric, the combinatorial curvature K_i at $v_i \in V$ is 2π less the cone angle at v_i .

If \mathcal{T} is a geometric triangulation of (S, V), the combinatorial curvature K_i at v_i is

$$K_i = 2\pi - \sum_{\triangle ijk \in F} \theta_i^{jk},$$

where the summation is taken over triangles with v_i as vertex and θ_i^{jk} is the inner angle of $\Delta i j k$ at v_i . Note that the combinatorial curvature K_i is independent of the geometric triangulations of (S, V) with a given polyhedral metric. Combinatorial curvature K satisfies the following discrete Gauss-Bonnet formula [12]

$$\sum_{i=1}^{n} K_i = 2\pi\chi(S) - \lambda Area(S),$$

where Area(S) is the area of the marked surface (S, V) with polyhedral metric d and $\lambda = -1, 0, +1$ respectively when d is a hyperbolic, Euclidean or spherical polyhedral metric respectively.

2.3. Vertex scaling of polyhedral metrics. Vertex scaling of PL metrics on a triangulated surface was introduced by Luo [37] and Rŏcek-Williams [47] independently as an analogy of the conformal transformation of Riemannian metrics.

Definition 2.3 ([37, 47]). Suppose d is a PL metric on a triangulated surface (S, V, \mathcal{T}) and u is a function defined on the vertices V. The vertex scaling of d by u for (S, V, \mathcal{T}) is defined to be the PL metric u * d such that

$$(u * d)_{ij} := d_{ij}e^{u_i + u_j}$$

determines a PL metric u * d on (S, V, \mathcal{T}) . u is called a conformal factor.

Note that u * d determines a PL metric on a triangulated surface (S, V, \mathcal{T}) is equivalent to $u : V \to \mathbb{R}$ is in the following admissible space of the conformal factors

$$\Omega^{\mathbb{E},\mathcal{T}}(d) \triangleq \{ u \in \mathbb{R}^V | d_{rs} e^{u_t} + d_{rt} e^{u_s} > d_{st} e^{u_r}, \{r,s,t\} = \{i,j,k\}, \forall \triangle ijk \in F \}.$$

Luo-Sun-Wu [39] showed that the vertex scaling of the PL metric is an approximation of the conformal transformation in the smooth case.

Vertex scaling of PH metrics on a triangulated surface was introduced by Bobenko-Pinkall-Springborn [2].

Definition 2.4 ([2]). Suppose d is a PH metric on a triangulated surface (S, V, \mathcal{T}) and u is a function defined on the vertices V. The vertex scaling of d by u is defined to be u * d such that

$$\sinh\frac{(u*d)_{ij}}{2} := e^{u_i + u_j} \sinh\frac{d_{ij}}{2}$$

determines a PH metric u * d on (S, V, \mathcal{T}) . u is called a conformal factor.

Similar to the PL metrics, u * d determines a PH metric if and only if the conformal factor u is in the admissible space $\Omega^{\mathbb{H},\mathcal{T}}(d)$ of conformal factors

$$\Omega^{\mathbb{H},\mathcal{T}}(d) \triangleq \{ u \in \mathbb{R}^V | (u * d)_{rs} + (u * d)_{rt} > (u * d)_{st}, \\ \{r,s,t\} = \{i,j,k\}, \forall \triangle ijk \in F \}.$$

The combinatorial curvature K is rigid with respect to the conformal factor.

Theorem 2.1 ([2]). Suppose (S, V, T) is a triangulated surface with a PL or PH metric, then the conformal factor is uniquely determined by the combinatorial curvature K (up to scaling for the PL metric).

Vertex scaling for spherical polyhedral metrics was defined by Bobenko-Pinkall-Springborn [2]. Gu-Luo-Sun-Wu [32] observed that the notions of vertex scaling are related to the Ptolemy identities for all polyhedral surfaces. Note that the definition of vertex scaling for polyhedral metrics depends on the triangulation of the marked surface (S, V).

2.4. Laplace operators on triangulated surfaces. The discrete Laplace operator of a PL metric on a triangulated surface, known as finite elements Laplacian, has been extensively studied in geometry and computer graphics and is defined as follows.

Definition 2.5. Suppose (S, V, \mathcal{T}) is a triangulated surface with a PL metric d. The Euclidean discrete Laplace operator of d on (S, V, \mathcal{T}) is defined to be the map

$$\Delta^{\mathbb{E},\mathcal{T}} : \mathbb{R}^V \longrightarrow \mathbb{R}^V$$
$$f \mapsto \Delta^{\mathbb{E},\mathcal{T}} f$$

where $f: V \to \mathbb{R}$ is a function defined on the vertices and the value of $\Delta^{\mathbb{E},\mathcal{T}} f$ at v_i is

$$(\Delta^{\mathbb{E},\mathcal{T}}f)_i = \sum_{j;j\sim i} \omega_{ij}^{\mathbb{E},\mathcal{T}}(f_j - f_i)$$
(2.1)

with weight

$$\omega_{ij}^{\mathbb{E},\mathcal{T}} = \cot \theta_k^{ij} + \cot \theta_l^{ij}.$$

Here θ_i^{jk} is the inner angle at the vertex v_i in a triangle $\triangle ijk \in F$.

Remark 2.1. Generally, for a triangulated surface (S, V, \mathcal{T}) with a PL metric d, the Euclidean discrete Lapalce operator of d depends on the triangulation \mathcal{T} . However, if the triangulation \mathcal{T} is Delaunay, even though there exists different Delaunay triangulations for the same PL metric d on a marked surface (S, V), the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$ is independent of the Delaunay triangulations of d on (S, V) [3]. In this case, the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$ is independent of the Delaunay triangulations.

Suppose $u: V \to \mathbb{R}$ is a conformal factor defined on the vertices and the vertex v_i is adjacent to v_i , Luo [37] proved

$$\frac{\partial \theta_i^{jk}}{\partial u_j} = \cot \theta_k^{ij}, \quad \frac{\partial K_i}{\partial u_j} = -(\cot \theta_k^{ij} + \cot \theta_l^{ij}),$$

where $\triangle ijk$ and $\triangle ijl$ are adjacent triangles in F.

$$L = (L_{ij})_{N \times N} = \frac{\partial(K_1, \cdots, K_N)}{\partial(u_1, \cdots, u_N)} = \begin{pmatrix} \frac{\partial K_1}{\partial u_1} & \cdots & \frac{\partial K_1}{\partial u_N} \\ \vdots & \ddots & \vdots \\ \frac{\partial K_N}{\partial u_1} & \cdots & \frac{\partial K_N}{\partial u_N} \end{pmatrix}$$
(2.2)

be the Jacobian of K with respect to u. For Euclidean polyhedral metrics, the matrix L has the following property.

Lemma 2.1 ([37]). For a triangulated surface (S, V, \mathcal{T}) with a PL metric d, the matrix L is symmetric and positive semi-definite on $\Omega^{\mathbb{E},\mathcal{T}}(d)$ with kernel $\{t\mathbf{1}|t \in \mathbb{R}\}, \text{ where } \mathbf{1} = (1, \cdots, 1)^T.$

By Lemma 2.1, we have $\sum_{j=1}^{N} \frac{\partial K_i}{\partial u_j} = 0$. Then the Euclidean discrete Laplace operator (2.1) translates into

$$(\Delta^{\mathbb{E},\mathcal{T}}f)_i = \sum_{j;j\sim i} (-\frac{\partial K_i}{\partial u_j})(f_j - f_i) = -\sum_{j\in V} \frac{\partial K_i}{\partial u_j} f_j,$$

which implies that

$$\Delta^{\mathbb{E},\mathcal{T}} = -L$$

in matrix form. In this way, we can take the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$ as a matrix-valued map defined on the admissible space of conformal factors $\Omega^{\mathbb{E},\mathcal{T}}(d)$. If the triangulation \mathcal{T} is Delaunay in u * d, then the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$, as a matrix-valued map, is defined on

$$\Omega_D^{\mathbb{E},\mathcal{T}}(d) \triangleq \{ u \in \mathbb{R}^V | \mathcal{T} \text{ is Delaunay in } u * d \},\$$

which is a subspace of the admissible conformal factors $\Omega^{\mathbb{E},\mathcal{T}}(d)$ [32, 40]. The idea of writing the discrete Laplace operator in matrix form comes from Ge [16, 17].

Note that L is the Jacobian of the curvature K with respect to the conformal factor u, we define the hyperbolic discrete Laplace operator similarly.

Definition 2.6. Suppose (S, V, \mathcal{T}) is a triangulated surface with a PH metric d. The hyperbolic discrete Laplace operator is defined to be the map

$$\begin{split} \Delta^{\mathbb{H},\mathcal{T}} : \mathbb{R}^V & \longrightarrow \mathbb{R}^V \\ f & \mapsto \Delta^{\mathbb{H},\mathcal{T}} f, \end{split}$$

where $f: V \to \mathbb{R}$ is a function defined on the vertices and the value of $\Delta^{\mathbb{H},\mathcal{T}} f$ at v_i is

$$(\Delta^{\mathbb{H},\mathcal{T}}f)_i = -(Lf)_i \tag{2.3}$$

with L formally given by (2.2).

For hyperbolic polyhedral metrics, the matrix ${\cal L}$ has the following property.

Lemma 2.2 ([2], Proposition 6.1.5). For a triangulated surface (S, V, \mathcal{T}) with a PH metric d, the matrix L is symmetric and strictly positive definite on $\Omega^{\mathbb{H},\mathcal{T}}(d)$.

Similar to the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$, the hyperbolic discrete Laplace operator $\Delta^{\mathbb{H},\mathcal{T}}$ is a matrix-valued map defined on the admissible space of conformal factors $\Omega^{\mathbb{H},\mathcal{T}}(d)$. If the triangulation \mathcal{T} is Delaunay in u * d, then the hyperbolic discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$, as a matrix-valued map, is defined on

$$\Omega_D^{\mathbb{H},\mathcal{T}}(d) \triangleq \{ u \in \mathbb{R}^V | \mathcal{T} \text{ is Delaunay in } u * d \},\$$

which is a subspace of the admissible conformal factors $\Omega^{\mathbb{H},\mathcal{T}}(d)$ [31].

Remark 2.2. Generally, for a triangulated surface (S, V, \mathcal{T}) with a PH metric d, the hyperbolic discrete Laplace operator $\Delta^{\mathbb{H},\mathcal{T}}$ depends on the triangulations. However, if the triangulation \mathcal{T} is restricted to be Delaunay, the hyperbolic discrete Laplace operator $\Delta^{\mathbb{H},\mathcal{T}}$ is independent of the Delaunay triangulations of (S, V) for the PH metric d [31]. In this case, the hyperbolic discrete Laplace operator by $\Delta^{\mathbb{H},\mathcal{T}}$ is intrinsic in the sense that it is independent of the Delaunay triangulations.

3. EUCLIDEAN COMBINATORIAL CALABI FLOW

3.1. Euclidean combinatorial Calabi flow on triangulated surfaces. The definition of Euclidean combinatorial Calabi flow on a triangulated marked surface (S, V, \mathcal{T}) is given in Definition 1.1, which could be written in the following matrix form [16]

$$\frac{du}{dt} = -LK.$$

Note that this is essential an ODE system, therefor the Euclidean combinatorial Calabi flow exists in a short time [16]. By direct calculations, we have the combinatorial curvature K evolves according to

$$\frac{dK}{dt} = -(\Delta^{\mathbb{E},\mathcal{T}})^2 K$$

along the Euclidean combinatorial Calabi flow (1.2). By Proposition 2.1, $\sum_{i=1}^{n} u_i$ is invariant along the Euclidean combinatorial Calabi flow (1.2) [16].

Similar to the fact that combinatorial Calabi flow for Thurston's circle packing metrics is the negative gradient flow of combinatorial Calabi energy [16, 17, 24], the combinatorial Calabi flow (1.2) for PL metrics is also the negative gradient flow of combinatorial Calabi energy.

Definition 3.1 ([16, 17]). Suppose d is a polyhedral metric on a marked surface (S, V), the combinatorial Calabi energy of d on (S, V) is defined to be

$$\mathcal{C} = ||K||^2 = \sum_{i=1}^n K_i^2.$$

If K^* is a function defined on V, the modified combinatorial Calabi energy is defined to be

$$\overline{\mathcal{C}} = ||K - K^*||^2 = \sum_{i=1}^n (K_i - K_i^*)^2.$$

Remark 3.1. Combinatorial Calabi energy C and modified combinatorial Calabi energy \overline{C} are independent of the geometric triangulations of d on (S, V).

Note that

$$\nabla_u \mathcal{C} = 2LK = -2\Delta^{\mathbb{E},\mathcal{T}}K$$

The Euclidean combinatorial Calabi flow (1.2) can be written as

$$\frac{du}{dt} = -\frac{1}{2}\nabla_u \mathcal{C}.$$

Proposition 3.1 ([16]). The Euclidean combinatorial Calabi flow (1.2) on a triangulated surface (S, V, T) is the negative gradient flow of combinatorial Calabi energy C and the combinatorial Calabi energy C is decreasing along the flow.

Similar to the stability results in [23, 24, 26, 27, 28], we have the following result for Euclidean combinatorial Calabi flow (1.2).

Theorem 3.1. If the solution of Euclidean combinatorial Calabi flow (1.2) on a triangulated surface (S, V, \mathcal{T}) converges as time tends to infinity, then the limit metric is a constant combinatorial curvature PL metric. Furthermore, suppose there is a constant combinatorial curvature PL metric $d^* =$ $u^* * d_0$ on (S, V, \mathcal{T}) with $\sum_{i=1}^n u_i^* = 0$, there exists a constant $\delta > 0$ such that if the initial modified combinatorial Calabi energy $||K(u(0)) - K(u^*)||^2 < \delta$, then the Euclidean combinatorial Calabi flow (1.2) on (S, V, \mathcal{T}) exists for all time and converges exponentially fast to u^* .

Proof. If u(t) converges as time tends to infinity, then $u(+\infty) = \lim_{t \to +\infty} u(t) \in \Omega^{\mathbb{E},\mathcal{T}}(d_0)$ exists. As K is a smooth function of $u \in \Omega^{\mathbb{E},\mathcal{T}}(d_0)$, we have $K(+\infty) = \lim_{t \to +\infty} K(u(t))$ and $L(+\infty) = \lim_{t \to +\infty} L(u(t))$ exist. Similarly, $\mathcal{C}(+\infty)$ and $\mathcal{C}'(+\infty)$ exist. Note that

$$\mathcal{C}'(t) = -2K^T L^2 K \le 0$$

and $\mathcal{C}(t)$ is uniformly bounded, we have

$$\mathcal{C}'(+\infty) = -2K^T(+\infty)L^2(+\infty)K(+\infty) = 0.$$

By Lemma 2.1, we have $K(+\infty) = \frac{2\pi\chi(S)}{n}(1, 1, \dots, 1)^T$, which implies that $u(+\infty) * d_0$ is a PL metric with constant combinatorial curvature.

By Theorem 2.1, we know that $u^* = u(+\infty)$ is the unique conformal factor such that $u^* * d_0$ has constant combinatorial curvature. Set $\Gamma(u) = \Delta^{\mathbb{E},\mathcal{T}} K = -LK$, then $\Gamma(u^*) = 0$ and

$$D_u \Gamma(u^*) = -L^T L \le 0. \tag{3.1}$$

Note that $rank (D_u \Gamma(u^*)) = n - 1$ and the kernel of $D_u \Gamma(u^*)$ is exactly $c(1, 1, \dots, 1)^T$, $c \in \mathbb{R}$. Along the Euclidean combinatorial Calabi flow (1.2), $\sum_{i=1}^n u_i$ is a constant. This implies that u^* is a local attractor of the Euclidean combinatorial Calabi flow (1.2) and the conclusion follows from the Lyapunov Stability Theorem ([44], Chapter 5). Q.E.D.

Theorem 3.1 does not give the long time existence and convergence of the Euclidean combinatorial Calabi flow (1.2) for general initial PL metrics. To study the long time behavior of the flow for general initial PL metrics, we need to analysis the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$. Note that, for $j \sim i$, the weight of the Euclidean discrete Laplace operator $\Delta^{\mathbb{E},\mathcal{T}}$ is

$$\omega_{ij}^{\mathbb{E},\mathcal{T}} = -\frac{\partial K_i}{\partial u_j} = \cot \theta_k^{ij} + \cot \theta_l^{ij} = \frac{\sin(\theta_k^{ij} + \theta_l^{ij})}{\sin \theta_k^{ij} \sin \theta_l^{ij}}$$

which may be negative and unbounded along the Euclidean combinatorial Calabi flow (1.2) for general initial PL metric. Even through discrete Laplace operators with negative coefficients has many applications in the study of combinatorial curvatures on two and three dimensional manifolds (see [15, 28, 29, 30, 33, 38, 46, 52, 53, 58] for example), a discrete Laplace operator is generally defined with nonnegative weights [14]. For $\Delta^{\mathbb{E},\mathcal{T}}$, the weight is nonnegative is equivalent to $\theta_k^{ij} + \theta_l^{ij} \leq \pi$, which is the Delaunay condition [3]. So it is natural to require the triangulation to be Delaunay, otherwise the coefficients of the discrete Laplace operator will be negative. Note that the admissible space for a Delaunay triangulation

 $D_{PL}(\mathcal{T}) = \{ [d] \in T_{PL}(S, V) | \mathcal{T} \text{ is isotopic to a Delaunay triangulation of } d \}$ is a subspace of the admissible space

 $\Omega(\mathcal{T}) = \{ [d] | d_{ii}, d_{ik}, d_{jk} \text{ satisfy the triangle inequalities for any } \triangle ijk \in F \}.$

3.2. Gu-Luo-Sun-Wu's work on discrete uniformization theorem. One of the main tools used in the proof of Theorem 1.1 for convergence of combinatorial Calabi flow with surgery is the discrete conformal theory developed by Gu-Luo-Sun-Wu [32]. In this subsection, we will briefly recall the theory. For details of the theory, please refer to [32].

Definition 3.2 ([32] Definition 1.1). Two piecewise linear metrics d, d' on (S, V) are discrete conformal if there exist sequences of PL metrics $d_1 = d$, \cdots , $d_m = d'$ on (S, V) and triangulations $\mathcal{T}_1, \cdots, \mathcal{T}_m$ of (S, V) satisfying

- (a): (Delaunay condition) each \mathcal{T}_i is Delaunay in d_i ,
- (b): (Vertex scaling condition) if $\mathcal{T}_i = \mathcal{T}_{i+1}$, there exists a function $u: V \to \mathbb{R}$ so that if e is an edge in \mathcal{T}_i with end points v and v', then the lengths $l_{d_{i+1}}(e)$ and $l_{d_i}(e)$ of e in d_i and d_{i+1} are related by

$$l_{d_{i+1}}(e) = l_{d_i}(e)e^{u(v) + u(v')},$$

(c): if $\mathcal{T}_i \neq \mathcal{T}_{i+1}$, then (S, d_i) is isometric to (S, d_{i+1}) by an isometry homotopic to identity in (S, V).

The space of PL metrics on (S, V) discrete conformal to d is called the conformal class of d and denoted by $\mathcal{D}(d)$.

Recall the following result for Delaunay triangulations.

Lemma 3.1 ([1, 3]). If \mathcal{T} and \mathcal{T}' are Delaunay triangulations of d, then there exists a sequence of Delaunay triangulations $\mathcal{T}_1 = \mathcal{T}, \mathcal{T}_2, \cdots, \mathcal{T}_k = \mathcal{T}'$ so that \mathcal{T}_{i+1} is obtained from \mathcal{T}_i by a diagonal switch.

The diagonal switch in Lemma 3.1 is the surgery by flipping described in the introduction.

The following discrete uniformization theorem was established in [32].

Theorem 3.2 ([32] Theorem 1.2). Suppose (S, V) is a closed connected marked surface and d is a PL metric on (S, V). Then for any $K^* : V \to$ $(-\infty, 2\pi)$ with $\sum_{v \in V} K^*(v) = 2\pi\chi(S)$, there exists a PL metric d', unique up to scaling and isometry homotopic to the identity on (S, V), such that d' is discrete conformal to d and the discrete curvature of d' is K^* . Furthermore, the metric d' can be found using a finite dimensional (convex) variational principle.

Denote the Teichimüller space of all PL metrics on (S, V) by $T_{PL}(S, V)$ and decorated Teichimüller space of all equivalence class of decorated hyperbolic metrics on S - V by $T_D(S - V)$. In the proof of Theorem 3.2, Gu-Luo-Sun-Wu proved the following result.

Theorem 3.3 ([32]). There is a C^1 -diffeomorphism $\mathbf{A} : T_{PL}(S, V) \rightarrow T_D(S, V)$ between $T_{PL}(S, V)$ and $T_D(S-V)$. Furthermore, the space $\mathcal{D}(d) \subset T_{PL}(S, V)$ of all equivalence classes of PL metrics discrete conformal to d is C^1 -diffeomorphic to $\{p\} \times \mathbb{R}^V_{>0}$ under the diffeomorphism \mathbf{A} , where p is the unique hyperbolic metric on S - V determined by the PL metric d on (S, V).

Set $u_i = \ln w_i$ for $w = (w_1, w_2, \dots, w_n) \in \mathbb{R}^n_{>0}$. Using the map **A**, Gu-Luo-Sun-Wu [32] defined the curvature map

$$\mathbf{F} : \mathbb{R}^n \to (-\infty, 2\pi)^n$$
$$u \mapsto K_{\mathbf{A}^{-1}(p, w(u))}$$

and proved the following property of \mathbf{F} .

Proposition 3.2 ([32]). (1) For any $k \in \mathbb{R}$, $\mathbf{F}(v + k(1, 1, \dots, 1)) = \mathbf{F}(v)$.

(2) There exists a C^2 -smooth convex function $W = \int \sum_{i=1}^n \mathbf{F}_i(u) du_i : \mathbb{R}^n \to \mathbb{R}$ so that its gradient ∇W is \mathbf{F} and the restriction $W : \{u \in \mathbb{R}^n | \sum_{i=1}^n u_i = 0\} \to \mathbb{R}$ is strictly convex.

Theorem 3.3 implies that the union of the admissible spaces $\Omega_D^{\mathbb{E},\mathcal{T}}(d')$ of conformal factors such that \mathcal{T} is Delaunay for $d' \in \mathcal{D}(d)$ is \mathbb{R}^n . It is further proved that $\mathbb{R}^n = \bigcup_{\mathcal{T}} \Omega_D^{\mathbb{E},\mathcal{T}}(d')$ is an analytic cell decomposition of \mathbb{R}^n [32]. Furthermore, **F** defined on \mathbb{R}^n is a C^1 -extension of the curvature K defined on the space of conformal factors $\Omega_D^{\mathbb{E},\mathcal{T}}(d')$ for $d' \in \mathcal{D}(d)$. Then we can extend the Euclidean discrete Laplace operator to be defined on \mathbb{R}^n , which is the space of the conformal factors for the discrete conformal class $\mathcal{D}(d)$.

Definition 3.3. Suppose (S, V) is a marked surface with a PL metric d. For a function $f: V \to \mathbb{R}$ on the vertices, the Euclidean discrete conformal Laplace operator of d on (S, V) is defined to be the map

$$\Delta^{\mathbb{E}} : \mathbb{R}^V \longrightarrow \mathbb{R}^V$$
$$f \mapsto \Delta^{\mathbb{E}} f,$$

where the value of $\Delta^{\mathbb{E}} f$ at v_i is

$$\Delta^{\mathbb{E}} f_i = \sum_{j \sim i} (-\frac{\partial \mathbf{F}_i}{\partial u_j}) (f_j - f_i) = -(\widetilde{L}f)_i,$$

where $\widetilde{L}_{ij} = \frac{\partial \mathbf{F}_i}{\partial u_j}$ is an extension of $L_{ij} = \frac{\partial K_i}{\partial u_j}$ for $u \in \Omega_D^{\mathbb{E}, \mathcal{T}}(d'), d' \in \mathcal{D}(d)$.

Remark 3.2. Note that \mathbf{F} is C^1 -smooth on \mathbb{R}^n and $\Delta^{\mathbb{E},\mathcal{T}}$ is independent of the Delaunay triangulations of a PL metric, the operator $\Delta^{\mathbb{E}}$ is well-defined on \mathbb{R}^n . Furthermore, $\Delta^{\mathbb{E}}$ is continuous and piecewise smooth on \mathbb{R}^n as a matrix-valued function ([32], Lemma 5.1).

3.3. Euclidean combinatorial Calabi flow with surgery on surfaces. We can extend the definition of combinatorial Calabi flow to be defined on a conformal class of d as follows.

Definition 3.4. Suppose (S, V) is a marked surface with a PL metric d_0 . The Euclidean combinatorial Calabi flow with surgery defined on the conformal class $\mathcal{D}(d_0)$ is

$$\begin{cases} \frac{du_i}{dt} = \Delta^{\mathbb{E}} \mathbf{F}_i, \\ u_i(0) = 0, \end{cases}$$
(3.2)

where $\Delta^{\mathbb{E}}$ is the Euclidean discrete conformal Laplace operator in Definition 3.3 defined for the PL metric $d(t) = \mathbf{A}^{-1}((\{p\}, e^{u(t)})) \in \mathcal{D}(d_0).$

By Proposition 3.2, $\sum_{i=1}^{n} u_i$ is invariant along the Euclidean combinatorial Calabi flow with surgery (3.2).

We can modify the definition of combinatorial Calabi flow with surgery to search for PL metrics with prescribed combinatorial curvatures and prove a generalization of Theorem 1.1.

Definition 3.5. Suppose (S, V) is a closed connected marked surface with a PL metric d_0 and K^* is a function defined on V. The Euclidean combinatorial Calabi flow with surgery for K^* defined on the conformal class $\mathcal{D}(d_0)$ is defined to be

$$\begin{cases} \frac{du_i}{dt} = \Delta^{\mathbb{E}} (\mathbf{F} - K^*)_i, \\ u_i(0) = 0. \end{cases}$$
(3.3)

If $K^* = 0$, the flow (3.3) is reduced to the flow (3.2). The Euclidean combinatorial Calabi flow with surgery for K^* (3.3) is also a negative gradient flow. The flow (3.3) can be written as

$$\frac{du_i}{dt} = \Delta^{\mathbb{E}}(\mathbf{F} - K^*)_i = -\frac{1}{2}\nabla_{u_i}(||\mathbf{F} - K^*||^2) = -\frac{1}{2}\nabla_{u_i}\overline{\mathcal{C}}_i$$

where $\overline{\mathcal{C}} = ||\mathbf{F} - K^*||^2$. This implies that (3.3) is the negative gradient flow of $\overline{\mathcal{C}}$ and $\overline{\mathcal{C}}$ is decreasing along the flow (3.3).

We have the following generalization of Theorem 1.1 for the Euclidean combinatorial Calabi flow with surgery for K^* (3.3).

Theorem 3.4. Suppose (S, V) is a closed connected marked surface and d_0 is a PL metric on (S, V). For any $K^* : V \to (-\infty, 2\pi)$ with $\sum_{v \in V} K^*(v) = 2\pi\chi(S)$, the combinatorial Calabi flow with surgery for K^* (3.3) exists for all time and converges exponentially fast to a unique PL metric $d^* \in \mathcal{D}(d_0)$ with combinatorial curvature K^* after finite number of surgeries.

Proof: Note that $\sum_{i=1}^{n} u_i$ is invariant along the combinatorial Calabi flow with surgery for K^* (3.3), then the solution u(t) of (3.3) lies in the hyperplane $\Sigma_0 = \{u \in \mathbb{R}^n | \sum_{i=1}^{n} u_i = 0\}$. As K^* is a given function on (S, V) with $\sum_{i=1}^{n} K_i^* = 2\pi\chi(S)$, there exists a unique $u^* \in \Sigma_0$ such that $\mathbf{F}(u^*) = K^*$ by Gu-Luo-Sun-Wu's Theorem 3.2. Define the Ricci potential function W^* on \mathbb{R}^n as

$$W^*(u) = W(u) - \int_{u^*}^{u} \sum_{i=1}^{n} K_i^* du_i = \int_{u^*}^{u} \sum_{i=1}^{n} (\mathbf{F}_i - K_i^*) du_i.$$

 W^* is well-defined and C^2 on \mathbb{R}^n with $\nabla W^* = \mathbf{F} - K^*$ by Proposition 3.2. Furthermore, W^* is convex on \mathbb{R}^n and strictly convex on Σ_0 . So we have $W^*(u^*) = 0$, $\nabla W^*(u^*) = 0$ and $W^*(u) \ge 0$ for any $u \in \Sigma_0$.

On the other hand, the following lemma is a well-known fact from analysis. The reader could refer to [28] (Lemma 4.6) for a proof.

Lemma 3.2. Suppose f(x) is a C^1 -smooth convex function on \mathbb{R}^{n-1} with $\nabla f(x_0) = 0$ for some $x_0 \in \mathbb{R}^{n-1}$, f(x) is strictly convex in a neighborhood of x_0 , then $\lim_{x\to\infty} f(x) = +\infty$.

By Lemma 3.2, we have $\lim_{u\to\infty} W^*|_{\Sigma_0} = +\infty$ and $W^*|_{\Sigma_0}$ is a nonnegative proper function on Σ_0 .

Set $\psi(t) = W^*(u(t))$. Then

$$\psi'(t) = \nabla_u W^* \cdot \frac{du}{dt} = -(\mathbf{F} - K^*)^T \cdot \widetilde{L} \cdot (\mathbf{F} - K^*) \le 0$$

by Proposition 3.2. As $\sum_{i=1}^{n} u_i$ is invariant along the flow and $\psi(t) \ge 0$, by the properness of $W^*|_{\Sigma_0}$ on Σ_0 , we have the solution u(t) lies in a compact subset of Σ_0 , which implies that the solution of the combinatorial Calabi flow (3.3) exists for all time.

By the C^1 -smoothness of **F** and the boundness of the solution u(t) of the flow (3.3), to prove the convergence of the combinatorial Calabi flow with

surgery (3.3), we just need to prove the convergence of the curvature. As the solution u(t) of the combinatorial Calabi flow (3.3) lies in a compact subset of Σ_0 and \widetilde{L} is strictly positive definite on Σ_0 , we have the first nonzero eigenvalue λ_1 of \widetilde{L} , which is continuous in u(t), has a uniform positive lower bound $\sqrt{\frac{\lambda}{2}} > 0$ along the Euclidean combinatorial Calabi flow with surgery (3.3). So we have

$$\overline{\mathcal{C}}'(t) = -2(\mathbf{F} - K^*)^T \cdot \widetilde{L}^2 \cdot (\mathbf{F} - K^*) \le -2\lambda_1^2(t)\overline{\mathcal{C}}(t) \le -\lambda\overline{\mathcal{C}}(t),$$

which implies

$$\overline{\mathcal{C}}(t) \le \overline{\mathcal{C}}(0)e^{-\lambda t}$$

So the curvature \mathbf{F} converges exponentially fast to K^* , which implies that u(t) converges exponentially fast to u^* by the C^1 -smoothness of \mathbf{F} .

In [51], Wu proved the following result.

Theorem 3.5. Suppose $\mathbb{R}^n = \bigcup D_i$ is an analytic cell decomposition. $f(x) \in C^1(\mathbb{R}^n)$ is analytic on each cell D_i , and has a unique minimum point where f has positive hession. Then its gradient flow $\gamma(t)$, which satisfies $\gamma'(t) = -\nabla f(\gamma(t))$, intersects the cell faces D_i finitely many times.

Combining Theorem 3.5, $\overline{\mathcal{C}}(u^*) = 0$, (3.1), the fact that the combinatorial Calabi flow is a negative gradient flow and the fact that $\mathbb{R}^n = \bigcup_{\mathcal{T}} \Omega_D^{\mathbb{E},\mathcal{T}}(d')$ is an analytic cell decomposition of \mathbb{R}^n [32], we get the finiteness of the surgeries along the combinatorial Calabi flow with surgery (3.3). Q.E.D.

Remark 3.3. Ge-Jiang [19] used another extension method introduced by Bobenko-Pinkall-Springborn [2] to extend the combinatorial curvature to the whole space by extending the inner angle of a triangle by constant. This method ensures the long time existence of the extended combinatorial Yamabe flow and was then used to study the combinatorial Ricci flow of inversive distance circle packing metrics [20, 21, 22, 28]. Comparing to Ge-Jiang's method to extend combinatorial curvature flow, the extension of Gu-Luo-Sun-Wu has the following two advantages. The first is that Ge-Jiang's method is used to study the curvature for a fixed triangulation, where there may exist no constant curvature polyhedral metric on the triangulation, while Gu-Luo-Sun-Wu's extension ensures the existence of constant curvature polyhedral metric. The second is that the extension method used by Ge-Jiang can be used to give a continuous extension of the combinatorial curvature, but the derivative of curvature may blow up along the flow, the bound of which is crucial for the long time existence and convergence of combinatorial Calabi flow. The C^1 -smoothness of **F** is crucial for the proof of Theorem 3.4.

4. Hyperbolic combinatorial Calabi flow

In this section, we introduce the combinatorial Calabi flow for piecewise hyperbolic metrics. To handle the singularities that may develop along the flow, we do surgery by flipping again. Then we prove the long time existence and convergence of the hyperbolic combinatorial Calabi flow with surgery. As the results are paralleling to the Euclidean case and the proofs are almost the same, the results in this section are stated without proof.

4.1. Hyperbolic combinatorial Calabi flow on triangulated surfaces. We define the combinatorial Calabi flow for PH metrics similar to the PL case.

Definition 4.1. Suppose d_0 is a PH metric on a triangulated surface (S, V, \mathcal{T}) . The hyperbolic combinatorial Calabi flow on (S, V, \mathcal{T}) is defined as

$$\begin{cases} \frac{du_i}{dt} = \Delta^{\mathbb{H}, \mathcal{T}} K_i, \\ u_i(0) = 0, \end{cases}$$
(4.1)

where $\Delta^{\mathbb{H},\mathcal{T}}$ is the hyperbolic discrete Laplace operator (2.3) defined by the *PH* metric u * d.

Similar to the PL case, we have the short time existence of the hyperbolic combinatorial Calabi flow and the combinatorial curvature K evolves according to

$$\frac{dK}{dt} = -(\Delta^{\mathbb{H},\mathcal{T}})^2 K$$

along the hyperbolic combinatorial Calabi flow (4.1). The hyperbolic combinatorial Calabi flow is also a negative gradient flow of the combinatorial Calabi energy.

Proposition 4.1. The combinatorial Calabi flow (4.1) for hyperbolic polyhedral metrics on a triangulated surface (S, V, \mathcal{T}) is the negative gradient flow of combinatorial Calabi energy C and the combinatorial Calabi energy C is decreasing along the flow.

Theorem 4.1. If the solution of hyperbolic combinatorial Calabi flow (4.1) on a triangulated surface (S, V, \mathcal{T}) converges as time tends to infinity, then the limit metric is a hyperbolic metric. Furthermore, suppose there exists a hyperbolic metric $d^* = u^* * d_0$ on (S, V, \mathcal{T}) , there exists a constant $\delta > 0$ such that if the initial modified Calabi energy $||K(u(0)) - K(u^*)||^2 < \delta$, then the hyperbolic combinatorial Calabi flow (4.1) on (S, V, \mathcal{T}) exists for all time and converges exponentially fast to u^* .

4.2. Gu-Guo-Luo-Sun-Wu's work on discrete uniformization theorem.

Definition 4.2 ([31], Definition 1). Two PH metrics d, d' on a closed marked surface (S, V) are discrete conformal if there exists sequences of PH metrics $d_1 = d$, $d_2, \dots, d_m = d'$ on (S, V) and triangulations $\mathcal{T}_1, \dots, \mathcal{T}_m$ of (S, V) satisfying

(a): each \mathcal{T}_i is Delaunay in d_i ,

(b): if $\mathcal{T}_i = \mathcal{T}_{i+1}$, there exists a function $u : V \to \mathbb{R}$, called a conformal factor, so that if e is an edge in \mathcal{T}_i with end points v and v', then the lengths $x_{d_i}(e)$ and $x_{d_{i+1}}(e)$ of e in metrics d_i and d_{i+1} are related by

$$\sinh \frac{x_{d_{i+1}}(e)}{2} = e^{u(v) + u(v')} \sinh \frac{x_{d_i}(e)}{2},$$

(c): if $\mathcal{T}_i \neq \mathcal{T}_{i+1}$, then (S, d_i) is isometric to (S, d_{i+1}) by an isometry homotopic to the identity in (S, V).

The space of PH metrics on (S, V) discrete conformal to d is called the conformal class of d and denoted by $\mathcal{D}(d)$.

Recall the following result of Delaunay triangulations for PH metrics.

Lemma 4.1 ([31], Proposition 16). Suppose (S, V) is a marked surface with a PH metric d. If \mathcal{T} and \mathcal{T}' are Delaunay triangulations of d, then there exists a sequence of Delaunay triangulations $\mathcal{T}_1 = \mathcal{T}, \mathcal{T}_2, \cdots, \mathcal{T}_k = \mathcal{T}'$ so that \mathcal{T}_{i+1} is obtained from \mathcal{T}_i by a diagonal switch.

A diagonal switch in Lemma 4.1 is referred to a surgery by flipping in the hyperbolic case.

The following discrete uniformization theorem was established by Gu-Guo-Luo-Sun-Wu [31].

Theorem 4.2 ([31], Theorem 3). Suppose (S, V) is a closed connected surface with marked points and d is a PH metric on (S, V). Then for any $K^*: V \to (-\infty, 2\pi)$ with $\sum_{v \in V} K^*(v) > 2\pi\chi(S)$, there exists a unique PH metric d' on (S, V) so that d' is discrete conformal to d and the discrete curvature of d' is K^* . Furthermore, the discrete Yamabe flow with surgery associated to curvature K^* having initial value d converges to d' linearly fast.

Denote the Teichimüller space of all PH metrics on (S, V) by $T_{hp}(S, V)$ and decorated Teichimüller space of all equivalence class of decorated hyperbolic metrics on S - V by $T_D(S - V)$. In the proof of Theorem 4.2, they prove the following result.

Theorem 4.3 ([31], Theorem 22, Corollary 24). There is a C^1 -diffeomorphism $\mathbf{A}: T_{hp}(S, V) \to T_D(S, V)$ between $T_{hp}(S, V)$ and $T_D(S - V)$. Furthermore, the space $\mathcal{D}(d) \subset T_{hp}(S, V)$ of all equivalence classes of PH metrics discrete conformal to d is C^1 -diffeomorphic to $\{p\} \times \mathbb{R}^V_{>0}$ under the diffeomorphism \mathbf{A} , where p is the unique hyperbolic metric on S - V determined by the PH metric d on (S, V).

Set $u_i = \ln w_i, 1 \le i \le n$, for $w = (w_1, w_2, \cdots, w_n) \in \mathbb{R}^n_{>0}$ and $\mathbf{P} = \{x \in (-\infty, 2\pi)^n | \sum_{i=1}^n x_i > 2\pi\chi(S)\}$. Using the map **A**, Gu-Guo-Luo-Sun-Wu defined the curvature map

$$\begin{aligned} \mathbf{F} &: \mathbb{R}^n \to \mathbf{P} \\ & u \mapsto K_{\mathbf{A}^{-1}(p, w(u))} \end{aligned}$$

and proved the following property of \mathbf{F} .

Proposition 4.2 ([31]). There exists a C^2 -smooth strictly convex function $W : \mathbb{R}^n \to \mathbb{R}$ so that its gradient ∇W is **F**.

Theorem 4.3 implies that the union of the admissible spaces $\Omega_D^{\mathbb{H},\mathcal{T}}(d')$ of conformal factors such that \mathcal{T} is Delaunay for $d' \in \mathcal{D}(d)$ is \mathbb{R}^n . Similar to the Euclidean case, $\mathbb{R}^n = \bigcup_{\mathcal{T}} \Omega_D^{\mathbb{H},\mathcal{T}}(d')$ is an analytic cell decomposition of \mathbb{R}^n [31]. Furthermore, \mathbf{F} defined on \mathbb{R}^n is a C^1 -extension of the curvature K defined on the space of conformal factors $\Omega_D^{\mathbb{H},\mathcal{T}}(d')$ for $d' \in \mathcal{D}(d)$. Then we can extend the hyperbolic discrete Laplace operator to be defined on \mathbb{R}^n , which is the total space of the conformal factors for the conformal class $\mathcal{D}(d)$.

Definition 4.3. Suppose (S, V) is a marked surface with a PH metric d. For a function $f: V \to \mathbb{R}$ on the vertices, the hyperbolic discrete conformal Laplace operator is defined to be the map

$$\Delta^{\mathbb{H}} : \mathbb{R}^V \longrightarrow \mathbb{R}^V$$
$$f \mapsto \Delta^{\mathbb{H}} f$$

where the value of $\Delta^{\mathbb{H}} f$ at v_i is

$$\Delta^{\mathbb{H}} f_i = \sum_{j \sim i} \left(-\frac{\partial \mathbf{F}_i}{\partial u_j}\right) (f_j - f_i) = -(\widetilde{L}f)_i$$

where $\widetilde{L}_{ij} = \frac{\partial \mathbf{F}_i}{\partial u_j}$ is an extension of $\frac{\partial K_i}{\partial u_j}$ for a single Delaunay triangulation \mathcal{T} without surgery.

Remark 4.1. F is C^1 -smooth on \mathbb{R}^n and $\Delta^{\mathbb{H},\mathcal{T}}$ is independent of the Delaunay triangulations of a PH metric [31], the operator $\Delta^{\mathbb{H}}$ is well-defined on \mathbb{R}^n . Furthermore, $\Delta^{\mathbb{H}}$ is continuous and piecewise smooth on \mathbb{R}^n as a matrix-valued function ([31], Lemma 26).

4.3. Hyperbolic combinatorial Calabi flow with surgery on surfaces. We can extend the definition of hyperbolic combinatorial Calabi flow to be defined on the discrete conformal class $\mathcal{D}(d_0)$ as follows.

Definition 4.4. Suppose (S, V) is a marked surface with a PH metric d_0 . The hyperbolic combinatorial Calabi flow with surgery defined on the conformal class $\mathcal{D}(d_0)$ is defined to be

$$\begin{cases} \frac{du_i}{dt} = \Delta^{\mathbb{H}} \mathbf{F}_i, \\ u_i(0) = 0. \end{cases}$$

We can modify the definition of hyperbolic combinatorial Calabi flow with surgery to search for PH metrics with prescribing combinatorial curvatures and prove a generalization of Theorem 1.2.

Definition 4.5. Suppose (S, V) is a marked surface with a PH metric d_0 and K^* is a function defined on V. The hyperbolic combinatorial Calabi

flow with surgery for K^* defined on the discrete conformal class $\mathcal{D}(d_0)$ is defined as

$$\begin{cases} \frac{du_i}{dt} = \Delta^{\mathbb{H}} (\mathbf{F} - K^*)_i, \\ u_i(0) = 0. \end{cases}$$
(4.2)

The hyperbolic combinatorial Calabi flow with surgery for K^* (4.2) is also a negative gradient flow. The flow (4.2) could be written in the following form

$$\frac{du_i}{dt} = \Delta^{\mathbb{E}}(\mathbf{F} - K^*)_i = -\frac{1}{2}\nabla_{u_i}(||\mathbf{F} - K^*||^2) = -\frac{1}{2}\nabla_{u_i}\overline{\mathcal{C}},$$

where $\overline{\mathcal{C}} = ||\mathbf{F} - K^*||^2$. This implies that (4.2) is the negative gradient flow of $\overline{\mathcal{C}}$ and $\overline{\mathcal{C}}$ is decreasing along the flow (4.2).

We have the following generalization of Theorem 1.2 for the hyperbolic combinatorial Calabi flow with surgery for K^* (4.2).

Theorem 4.4. Suppose (S, V) is a closed connected marked surface and d_0 is a PH metric on (S, V). For any $K^* : V \to (-\infty, 2\pi)$ with $\sum_{v \in V} K^*(v) > 2\pi\chi(S)$, the hyperbolic combinatorial Calabi flow with surgery (4.2) exists for all time and converges exponentially fast to a unique PH metric $d^* \in \mathcal{D}(d_0)$ with combinatorial curvature K^* after finite number of surgeries.

5. Remarks and questions

We use the discrete uniformization theorems established in [31, 32] to prove the long time existence and convergence of the combinatorial Calabi flow with surgery in Theorem 1.1 and Theorem 1.2. For the combinatorial Yamabe flow with surgery, the long time existence and convergence were proved similarly [31, 32]. Similar results were proved for smooth surface Ricci flow [11, 35] and smooth surface Calabi flow [9, 13]. However, the smooth Ricci flow and Calabi flow can be used to prove the uniformization theorem on Riemannian surfaces [7, 8, 10]. So it is interesting to ask the following question.

Question: Can we reprove the discrete uniformization theorems via the combinatorial Yamabe flow or the combinatorial Calabi flow with surgery?

In [54, 55], the second author generalizes the definition of α -curvatures in [26, 27] for Thurston's circle packing metrics to polyhedral metrics on surfaces. Similar properties of α -curvatures on polyhedral surfaces to those in [26, 27] are established in [54, 55] using combinatorial Yamabe flow and combinatorial Calabi flow with surgery. Furthermore, based on the discrete uniformization theorems in [31, 32], we prove a parameterized discrete uniformization theorem for α -curvatures in [54, 55].

Acknowledgements

The first author thanks Professor Jian Sun for introduction and guidance on

discrete geometry. Part of this work was done when the first author was visiting the School of Mathematics and Statistics, Wuhan University. He would like to thank Wuhan University for its hospitality. The second author also thanks Professor Kai Zheng for communications on Calabi flow. The authors thank Dr. Tianqi Wu for helpful communications. The research of the second author is supported by Hubei Provincial Natural Science Foundation of China under grant no. 2017CFB681, National Natural Science Foundation of China under grant no. 61772379 and no. 11301402 and Fundamental Research Funds for the Central Universities.

References

- [1] F. Aurenhammer, R. Klein, *Voronoi diagrams*. Handbook of computational geometry, 201-290, North-Holland, Amsterdam, 2000.
- [2] A. Bobenko, U. Pinkall, B. Springborn, *Discrete conformal maps and ideal hyperbolic polyhedra*. Geom. Topol. 19 (2015), no. 4, 2155-2215.
- [3] A. Bobenko, B. Springborn, A discrete Laplace-Beltrami operator for simplicial surfaces. Discrete Comput. Geom. 38 (2007), no. 4, 740-756.
- [4] Š. Brendle, R. Schoen, Manifolds with 1/4-pinched curvature are space forms.
 J. Amer. Math. Soc. 22 (2009), no. 1, 287-307
- [5] E. Calabi, *Extremal Kähler metrics*, Seminar on Differential Geometry, pp. 259-290, Ann. of Math. Stud., 102, Princeton Univ. Press, Princeton, N.J., 1982.
- [6] E. Calabi, *Extremal Kähler metrics. II*, Differential geometry and complex analysis, Springer, Berlin, 1985, 95-114.
- [7] S. C. Chang, Global existence and convergence of solutions of Calabi flow on surfaces of genus $h \ge 2$. J. Math. Kyoto Univ. 40 (2000), no. 2, 363-377.
- [8] S. C. Chang, The 2-dimensional Calabi flow. Nagoya Math. J. 181 (2006), 63-73.
- [9] X. X. Chen, Calabi flow in Riemann surfaces revisited: a new point of view, Internat. Math. Res. Notices 2001, no. 6, 275-297.
- [10] X. X. Chen, P. Lu, G. Tian, A note on uniformization of Riemann surfaces by Ricci flow, Proc. Amer. Math. Soc. 134 (2006), no. 11, 3391-3393.
- [11] B. Chow, *The Ricci flow on the 2-sphere*, J. Differential Geometry, 33 (1991), 325-334.
- [12] B. Chow, F. Luo, *Combinatorial Ricci flows on surfaces*, J. Differential Geometry, 63 (2003), 97-129.
- [13] P. T. Chruściel, Semi-global existence and convergence of solutions of the Robinson-Trautman (2-dimensional Calabi) equation, Comm. Math. Phys. 137 (1991), 289-313.
- [14] Chung, Fan R. K. Spectral graph theory. CBMS Regional Conference Series in Mathematics, 92. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1997. xii+207 pp. ISBN: 0-8218-0315-8
- [15] D. Cooper, I. Rivin, Combinatorial scalar curvature and rigidity of ball packings, Math. Res. Lett. 3 (1996), 51-60.
- [16] H. Ge, *Combinatorial methods and geometric equations*, Thesis (Ph.D.)-Peking University, Beijing. 2012.
- [17] H. Ge, Combinatorial Calabi flows on surfaces, Trans. Amer. Math. Soc. 370 (2018), no. 2, 1377-1391.
- [18] H. Ge, B. Hua, On combinatorial Calabi flow with hyperbolic circle patterns, Adv. Math. 333 (2018), 523-538.

- [19] H. Ge, W. Jiang, On the deformation of discrete conformal factors on surfaces. Calc. Var. Partial Differential Equations 55 (2016), no. 6, Art. 136, 14 pp.
- [20] H. Ge, W. Jiang, On the deformation of inversive distance circle packings.
- arXiv:1604.08317 [math.GT]. To appear in Trans. Amer. Math. Soc.
- [21] H. Ge, W. Jiang, On the deformation of inversive distance circle packings, II.
 J. Funct. Anal. 272 (2017), no. 9, 3573-3595.
- [22] H. Ge, W. Jiang, On the deformation of inversive distance circle packings, III.
 J. Funct. Anal. 272 (2017), no. 9, 3596-3609.
- [23] H. Ge, X. Xu, Discrete quasi-Einstein metrics and combinatorial curvature flows in 3-dimension, Adv. Math. 267 (2014), 470-497.
- [24] H. Ge, X. Xu, 2-dimensional combinatorial Calabi flow in hyperbolic background geometry, Differential Geom. Appl. 47 (2016) 86-98.
- [25] H. Ge, X. Xu, α -curvatures and α -flows on low dimensional triangulated manifolds. Calc. Var. Partial Differential Equations 55 (2016), no. 1, Art. 12, 16 pp
- [26] H. Ge, X. Xu, A Discrete Ricci Flow on Surfaces in Hyperbolic Background Geometry, Int. Math. Res. Not. IMRN 2017, no. 11, 3510-3527.
- [27] H. Ge, X. Xu, A combinatorial Yamabe problem on two and three dimensional manifolds, arXiv:1504.05814v2 [math.DG].
- [28] H. Ge, X. Xu, On a combinatorial curvature for surfaces with inversive distance circle packing metrics. J. Funct. Anal. 275 (2018), no. 3, 523-558.
- [29] D. Glickenstein, A combinatorial Yamabe flow in three dimensions, Topology 44 (2005), No. 4, 791-808.
- [30] D. Glickenstein, A maximum principle for combinatorial Yamabe flow, Topology 44 (2005), No. 4, 809-825.
- [31] X. D. Gu, R. Guo, F. Luo, J. Sun, T. Wu, A discrete uniformization theorem for polyhedral surfaces II, J. Differential Geom. 109 (2018), no. 3, 431-466.
- [32] X. D. Gu, F. Luo, J. Sun, T. Wu, A discrete uniformization theorem for polyhedral surfaces, J. Differential Geom. 109 (2018), no. 2, 223-256.
- [33] R. Guo, *Local rigidity of inversive distance circle packing*, Trans. Amer. Math. Soc. 363 (2011) 4757-4776.
- [34] R.S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Differential Geom., 17 (1982), 255-306.
- [35] R.S. Hamilton, *The Ricci flow on surfaces*. Mathematics and general relativity (Santa Cruz, CA, 1986), 237-262, Contemp. Math., 71, Amer. Math. Soc., Providence, RI, 1988.
- [36] G. Leibon, Characterizing the Delaunay decompositions of compact hyperbolic surface. Geom. Topol. 6 (2002), 361-391.
- [37] F. Luo, *Combinatorial Yamabe flow on surfaces*, Commun. Contemp. Math. 6 (2004), no. 5, 765-780.
- [38] F. Luo, Rigidity of polyhedral surfaces, III, Geom. Topol. 15 (2011), 2299-2319.
- [39] F. Luo, J. Sun, T. Wu, Discrete conformal geometry of polyhedral surfaces and its convergence, preprint, 37 pages, 2016.
- [40] R. C. Penner, The decorated Teichmuller space of punctured surfaces. Comm. Math. Phys. 113 (1987), no. 2, 299-339.
- [41] G. Perelman, The entropy formula for the Ricci flow and its geometric applications, arXiv:math/0211159 [math.DG].
- [42] G. Perelman, *Ricci flow with surgery on three-manifolds*, arXiv:math/0303109 [math.DG].
- [43] G. Perelman, Finite extinction time for the solutions to the Ricci flow on certain three-manifolds, arXiv:math/0307245 [math.DG].
- [44] L.S. Pontryagin, Ordinary differential equations, Addison-Wesley Publishing Company Inc., Reading, 1962.

[45] I. Rivin, Euclidean structures of simplicial surfaces and hyperbolic volume. Ann. of Math. 139 (1994), 553-580.

[46] I. Rivin, An extended correction to "Combinatorial Scalar Curvature and Rigidity of Ball Packings," (by D. Cooper and I. Rivin), arXiv:math/0302069v2 [math.MG].

[47] M. Rŏcek, R. M. Williams, *The quantization of Regge calculus*. Z. Phys. C 21 (1984), no. 4, 371-381.

[48] K. Su, C. Li, Y. Zhou, X. Xu, X. Gu, *Discrete Calabi flow: a unified conformal parameterization method*, To appear in Pacific Graphics 2019.

[49] J. Sun, T. Wu, X. D. Gu, F. Luo, Discrete conformal deformation: algorithm and experiments. SIAM J. Imaging Sci. 8 (2015), no. 3, 1421-1456.

[50] W. Thurston, *Geometry and topology of 3-manifolds*, Princeton lecture notes 1976, http://www.msri.org/publications/books/gt3m.

[51] T. Wu, *Finiteness of switches in discrete Yamabe flow*, Master Thesis, Tsinghua University, Beijing, 2014.

[52] X. Xu, On the global rigidity of sphere packings on 3-dimensional manifolds, arXiv:1611.08835v4 [math.GT]. To appear in J. Differential Geometry.

[53] X. Xu, Rigidity of inversive distance circle packings revisited, Adv. Math. 332 (2018), 476-509.

[54] X. Xu, Combinatorial α -curvatures and α -flows on polyhedral surfaces, I, arXiv:1806.04516 [math.GT].

[55] X. Xu, Combinatorial α -curvatures and α -flows on polyhedral surfaces, II, preprint, 2018

[56] W. Zeng, X. Gu, *Ricci flow for shape analysis and surface registration. Theories, algorithms and applications.* Springer Briefs in Mathematics. Springer, New York, 2013. xii+139 pp. ISBN: 978-1-4614-8780-7; 978-1-4614-8781-4

[57] H. Zhao, X. Li, H. Ge, N. Lei, M. Zhang, X. Wang, X. D. Gu, Conformal mesh parameterization using discrete Calabi flow. Comput. Aided Geom. Design 63 (2018), 96-108.

[58] Z. Zhou, Circle patterns with obtuse exterior intersection angles, arXiv:1703.01768v2 [math.GT].

(Xiang Zhu) Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, P.R. China

E-mail address: zhu-x13@mails.tsinghua.edu.cn

(Xu Xu) School of Mathematics and Statistics, Wuhan University, Wuhan 430072, P.R. China

E-mail address: xuxu2@whu.edu.cn