Nonuniform average sampling in multiply generated shift-invariant subspaces of mixed Lebesgue spaces

Qingyue Zhang

College of Science, Tianjin University of Technology, Tianjin 300384, China e-mails: jczhangqingyue@163.com

June 14, 2018

Abstract. In this paper, we study nonuniform average sampling problem in multiply generated shift-invariant subspaces of mixed Lebesgue spaces. We discuss two types of average sampled values: average sampled values $\{\langle f, \psi_a(\cdot - x_j, \cdot - y_k)\rangle : j, k \in \mathbb{J}\}\$ generated by single averaging function and average sampled values $\{ \langle f, \psi_{x_j, y_k} \rangle : j, k \in \mathbb{J} \}$ generated by multiple averaging functions. Two fast reconstruction algorithms for this two types of average sampled values are provided.

Key words. mixed Lebesgue spaces; nonuniform average sampling; shift-invariant subspaces.

2010 MR Subject Classification 94A20, 94A12, 42C15, 41A58

1 Introduction and motivation

In 1961, Benedek firstly proposed mixed Lebesgue spaces [\[1,](#page-13-0)[2\]](#page-13-1). In the 1980s, Fernandez, and Francia, Ruiz and Torrea developed the theory of mixed Lebesgue spaces in integral operators and Calderon–Zygmund operators, respectively $[3, 4]$ $[3, 4]$. Recently, Torres and Ward, and Li, Liu and Zhang studied sampling problem in shift-invariant subspaces of mixed Lebesgue spaces [\[5–](#page-13-4)[8\]](#page-13-5). Mixed Lebesgue spaces generalize Lebesgue spaces. It was proposed due to considering functions that depend on independent quantities with different properties. For a function in mixed Lebesgue spaces, one can consider the integrability of each variable independently. This is distinct from traditional Lebesgue spaces. The flexibility of mixed Lebesgue spaces makes them have a crucial role to play in the study of time based partial differential equations. In this context, we study nonuniform average sampling problem in shift-invariant subspaces of mixed Lebesgue spaces.

Sampling theorem is the theoretical basis of modern pulse coded modulation communication system, and is also one of the most powerful basic tools in signal processing and image processing. In 1948, Shannon formally proposed sampling theorem [\[9,](#page-14-0)[10\]](#page-14-1). Shannon sampling theorem shows that for any $f \in L^2(\mathbb{R})$ with supp $\hat{f} \subseteq [-T, T]$,

$$
f(x) = \sum_{n \in \mathbb{Z}} f\left(\frac{n}{2T}\right) \frac{\sin \pi (2Tx - n)}{\pi (2Tx - n)},
$$

where the series converges uniformly on compact sets and in $L^2(\mathbb{R})$, and

$$
\hat{f}(\xi) = \int_{\mathbb{R}} f(x)e^{-2\pi ix\xi} dx, \qquad \xi \in \mathbb{R}.
$$

However, in many realistic situations, the sampling set is only a nonuniform sampling set. For example, the transmission through the internet from satellites only can be viewed as a nonuniform sampling problem, because there exists the loss of data packets in the transmission. In recent years, there are many results concerning nonuniform sampling problem [\[11](#page-14-2)[–17\]](#page-14-3). Uniform and nonuniform sampling problems also have been generalized to more general shift-invariant spaces [\[18](#page-14-4)[–23\]](#page-14-5) of the form

$$
V(\phi) = \left\{ \sum_{k \in \mathbb{Z}} c(k) \phi(x - k) : \{c(k) : k \in \mathbb{Z}\} \in \ell^{2}(\mathbb{Z}) \right\}.
$$

In the classical sampling theory, the sampling values are the function values of the signal at the sampling points. In practical application, due to the precision of physical equipment and so on, it is impossible to measure the exact value of a signal at a point. And the actual measured value is the local mean value of the signal near this point. Therefore, average sampling has attracted more and more the attentions of researchers [\[24](#page-14-6)[–32\]](#page-15-0).

For the sampling problem in shift-invariant subspaces of mixed Lebesgue spaces, Torres and Ward studied uniform sampling problem for band-limited functions in mixed Lebesgue spaces [\[5,](#page-13-4) [6\]](#page-13-6). Li, Liu and Zhang discussed the nonuniform sampling problem in principal shift-invariant spaces of mixed Lebesgue spaces [\[7,](#page-13-7) [8\]](#page-13-5). In this paper, we discuss nonuniform average sampling problem in multiply generated shift-invariant subspaces of mixed Lebesgue spaces. We discuss two types of average sampled values: average sampled values $\{\langle f, \psi_a(\cdot - x_i, \cdot - y_k)\rangle : j, k \in \mathbb{J}\}\$ generated by single averaging function and average sampled values $\{\langle f, \psi_{x_j,y_k}\rangle : j, k \in \mathbb{J}\}\$ generated by multiple averaging functions. Two fast reconstruction algorithms for this two types of average sampled values are provided.

The paper is organized as follows. In the next section, we give the definitions and preliminary results needed and define multiply generated shift-invariant subspaces in mixed Lebesgue spaces $L^{p,q}(\mathbb{R}^{d+1})$. In Section 3, we give main results of this paper. Section 4 gives some useful propositions and lemmas. In Section 5, we give proofs of main results. Finally, concluding remarks are presented in Section 6.

2 Definitions and preliminary results

In this section, we give some definitions and preliminary results needed in this paper. First of all, we give the definition of mixed Lebesgue spaces $L^{p,q}(\mathbb{R}^{d+1})$.

Definition 2.1 For $1 \leq p, q < +\infty$. $L^{p,q} = L^{p,q}(\mathbb{R}^{d+1})$ consists of all measurable func $tions f = f(x, y)$ *defined on* $\mathbb{R} \times \mathbb{R}^d$ *satisfying*

$$
||f||_{L^{p,q}} = \left[\int_{\mathbb{R}} \left(\int_{\mathbb{R}^d} |f(x,y)|^q dy \right)^{\frac{p}{q}} dx \right]^{\frac{1}{p}} < +\infty.
$$

The corresponding sequence spaces are defined by

$$
\ell^{p,q} = \ell^{p,q}(\mathbb{Z}^{d+1}) = \left\{ c : ||c||_{\ell^{p,q}}^p = \sum_{k_1 \in \mathbb{Z}} \left(\sum_{k_2 \in \mathbb{Z}^d} |c(k_1, k_2)|^q \right)^{\frac{p}{q}} < +\infty \right\}.
$$

In order to control the local behavior of functions, we introduce mixed Wiener amalgam spaces $W(L^{p,q})(\mathbb{R}^{d+1}).$

Definition 2.2 *For* $1 \leq p, q < \infty$ *, if a measurable function f satisfies*

$$
||f||_{W(L^{p,q})}^p := \sum_{n \in \mathbb{Z}} \sup_{x \in [0,1]} \left[\sum_{l \in \mathbb{Z}^d} \sup_{y \in [0,1]^d} |f(x+n, y+l)|^q \right]^{p/q} < \infty,
$$

then we say that f *belongs to the mixed Wiener amalgam space* $W(L^{p,q}) = W(L^{p,q})(\mathbb{R}^{d+1})$ *.*

For $1 \leq p, q < \infty$, let $W_0(L^{p,q})$ denote the space of all continuous functions in $W(L^{p,q})$. For $1 \leq p < \infty$, if a function f satisfies

$$
||f||_{W(L^{p})}^{p} := \sum_{k \in \mathbb{Z}^{d+1}} \text{ess sup}_{x \in [0,1]^{d+1}} |f(x+k)|^{p} < \infty,
$$

then we say that f belongs to the Wiener amalgam space $W(L^p) = W(L^p)(\mathbb{R}^{d+1})$. Obviously, $W(L^p) \subset W(L^{p,p})$.

For $1 \leq p < \infty$, let $W_0(L^p)$ denote the space of all continuous functions in $W(L^p)$. Let B be a Banach space. $(B)^{(r)}$ denotes r copies $B \times \cdots \times B$ of B.

For any $f, g \in L^2(\mathbb{R}^{d+1})$, define their convolution

$$
(f * g)(x) = \int_{\mathbb{R}^{d+1}} f(y)g(x - y)dy.
$$

The following is a preliminary result.

Lemma 2.3 *If* $f \in L^1(\mathbb{R}^{d+1})$ *and* $g \in W(L^{1,1})$ *, then* $f * g \in W(L^{1,1})$ *and*

$$
||f * g||_{W(L^{1,1})} \le ||g||_{W(L^{1,1})} ||f||_{L^{1}}.
$$

Proof. Since

$$
||f * g||_{W(L^{1,1})} = \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d}
$$

$$
\left| \int_{y_1 \in \mathbb{R}} \int_{y_2 \in \mathbb{R}^d} f(y_1, y_2) g(x_1 + k_1 - y_1, x_2 + k_2 - y_2) dy_2 dy_1 \right|
$$

$$
\leq \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d}
$$

$$
\int_{y_1 \in \mathbb{R}} \int_{y_2 \in \mathbb{R}^d} |f(y_1, y_2)| |g(x_1 + k_1 - y_1, x_2 + k_2 - y_2)| dy_2 dy_1
$$
\n
$$
\leq \int_{y_1 \in \mathbb{R}} \int_{y_2 \in \mathbb{R}^d} |f(y_1, y_2)| \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} |g(x_1 + k_1 - y_1, x_2 + k_2 - y_2)| dy_2 dy_1
$$
\n
$$
\leq \int_{y_1 \in \mathbb{R}} \int_{y_2 \in \mathbb{R}^d} |f(y_1, y_2)| \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} |g(x_1 + k_1, x_2 + k_2)| dy_2 dy_1
$$
\n
$$
= \int_{y_1 \in \mathbb{R}} \int_{y_2 \in \mathbb{R}^d} |f(y_1, y_2)| ||g||_{W(L^{1,1})} dy_2 dy_1
$$
\n
$$
= ||g||_{W(L^{1,1})} \int_{y_1 \in \mathbb{R}} \int_{y_2 \in \mathbb{R}^d} |f(y_1, y_2)| dy_2 dy_1
$$
\n
$$
= ||g||_{W(L^{1,1})} ||f||_{L^1},
$$

the desired result in Lemma [2.3](#page-2-0) is following.

2.1 Shift-invariant spaces

For $\Phi = (\phi_1, \phi_2, \dots, \phi_r)^T \in W(L^{1,1})^{(r)}$, the multiply generated shift-invariant space in the mixed Legesgue spaces $L^{p,q}$ is defined by

$$
V_{p,q}(\Phi) = \left\{ \sum_{i=1}^r \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} c_i(k_1, k_2) \phi_i(\cdot - k_1, \cdot - k_2) : \\ c_i = \left\{ c_i(k_1, k_2) : k_1 \in \mathbb{Z}, k_2 \in \mathbb{Z}^d \right\} \in \ell^{p,q}, 1 \le i \le r \right\}.
$$

It is easy to see that the three sum pointwisely converges almost everywhere. In fact, for any $1 \leq i \leq r$, $c_i = \{c_i(k_1, k_2) : k_1 \in \mathbb{Z}, k_2 \in \mathbb{Z}^d\} \in \ell^{p,q}$ derives $c_i \in \ell^{\infty}$. This combines $\Phi = (\phi_1, \phi_2, \dots, \phi_r)^T \in W(L^{1,1})^{(r)}$ gets

$$
\sum_{i=1}^{r} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} |c_i(k_1, k_2) \phi_i(x - k_1, y - k_2)| \leq \sum_{i=1}^{r} ||c_i||_{\infty} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} |\phi_i(x - k_1, y - k_2)|
$$

$$
\leq \sum_{i=1}^{r} ||c_i||_{\infty} ||\phi_i||_{W(L^{1,1})} < \infty \ (a.e.).
$$

The following proposition gives that multiply generated shift-invariant spaces are welldefined in $L^{p,q}$.

Proposition 2.4 *[\[8,](#page-13-5) Theorem 2.8] Assume that* $1 \leq p, q < \infty$ *and* $\Phi = (\phi_1, \phi_2, \cdots, \phi_r)^T \in$ $W(L^{1,1})^{(r)}$. Then for any $C = (c_1, c_2, \dots, c_r)^T \in (\ell^{p,q})^{(r)}$, the function

$$
f = \sum_{i=1}^{r} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} c_i(k_1, k_2) \phi_i(\cdot - k_1, \cdot - k_2)
$$

belongs to $L^{p,q}$ *and there exist* $D_1, D_2 > 0$ *such that*

$$
D_1||f||_{L^{p,q}} \leq \left(\sum_{i=1}^r ||c_i||_{\ell^{p,q}}^2\right)^{1/2} \leq D_2||f||_{L^{p,q}}.
$$

3 Main results

In this section, we mainly discuss nonuniform average sampling in multiply generated shift-invariant spaces. The main results of this section are two fast reconstruction algorithms which allow to exactly reconstruct the signals f in multiply generated shift-invariant subspaces from the average sampled values of f .

3.1 The case of single averaging function

In this subsection, we will give a fast reconstruction algorithm which allows to exactly reconstruct the signals f in multiply generated shift-invariant subspaces from the average sampled values $\{\langle f, \psi_a(\cdot - x_j, \cdot - y_k)\rangle : j, k \in \mathbb{J}\}\$ of f. Before giving the main result of this subsection, we first give some definitions.

Definition 3.1 *A bounded uniform partition of unity* $\{\beta_{j,k} : j,k \in \mathbb{J}\}\$ associated to ${B_\gamma(x_j, y_k) : j, k \in \mathbb{J}}$ *is a set of functions satisfying*

- (i). $0 \leq \beta_{j,k} \leq 1, \forall j, k \in \mathbb{J}$,
- (ii). $\text{supp}\beta_{i,k} \subset B_{\gamma}(x_i, y_k),$
- (iii). $\sum_{j\in\mathbb{J}}\sum_{k\in\mathbb{J}}\beta_{j,k}=1$.

Here $B_{\gamma}(x_j, y_k)$ *is the open ball with center* (x_j, y_k) *and radius* γ *.*

If $f \in W_0(L^{1,1}),$ we define

$$
A_{X,a}f = \sum_{j\in\mathbb{J}}\sum_{k\in\mathbb{J}}\langle f, \psi_a(\cdot - x_j, \cdot - y_k) \rangle \beta_{j,k} = \sum_{j\in\mathbb{J}}\sum_{k\in\mathbb{J}}(f * \psi_a^*)(x_j, y_k)\beta_{j,k},
$$

and define

$$
Q_X f = \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} f(x_j, y_k) \beta_{j,k}
$$

for the quasi-interpolant of the sequence $c(j, k) = f(x_j, y_k)$. Here $\psi_a(\cdot) = 1/a^{d+1}\psi(\cdot/a)$ and $\psi_a^*(x) = \overline{\psi_a(-x)}$. Obviously, one has $A_{X,a}f = Q_X(f * \psi_a^*)$.

In order to describe the structure of the sampling set X , we give the following definition.

Definition 3.2 *If a set* $X = \{(x_j, y_k) : k, j \in \mathbb{J}, x_k \in \mathbb{R}, y_j \in \mathbb{R}^d\}$ *satisfies*

$$
\mathbb{R}^{d+1} = \cup_{j,k} B_{\gamma}(x_j, y_k) \quad \text{for every } \gamma > \gamma_0,
$$

then we say that the set X *is* γ_0 -dense in \mathbb{R}^{d+1} . Here $B_{\gamma}(x_j, y_k)$ *is the open ball with center* (x_j, y_k) *and radius* γ *, and* \mathbb{J} *is a countable index set.*

The following is main result of this subsection.

Theorem 3.3 *Assume that* $\Phi = (\phi_1, \phi_2, \dots, \phi_r)^T \in W_0(L^{1,1})^{(r)}$ whose support is compact *and P is a bounded projection from* $L^{p,q}$ *onto* $V_{p,q}(\Phi)$ *. Let* $\psi \in W_0(L^{1,1})$ *and* $\int_{\mathbb{R}^{d+1}} \psi = 1$ *. Then there are density* $\gamma_0 = \gamma_0(\Phi, \psi) > 0$ *and* $a_0 = a_0(\Phi, \psi) > 0$ *such that any* $f \in V_{p,q}(\Phi)$ *can be reconstructed from its average samples* $\{ \langle f, \psi_a(\cdot - x_j, \cdot - y_k) \rangle : j, k \in \mathbb{J} \}$ *on any* γ ($\gamma \leq \gamma_0$)-dense set $X = \{(x_j, y_k) : j, k \in \mathbb{J}\}\$ and for any $0 < a \leq a_0$, by the following *iterative algorithm:*

$$
\begin{cases}\nf_1 = P A_{X,a} f \\
f_{n+1} = P A_{X,a} (f - f_n) + f_n.\n\end{cases}
$$
\n(3.1)

The iterate f_n converges to f in the $L^{p,q}$ norm. Furthermore, the convergence is geometric, *namely,*

$$
||f - f_n||_{L^{p,q}} \leq M\alpha^n
$$

for some $\alpha = \alpha(\gamma, a, P, \Phi, \psi) < 1$ *and* $M < \infty$.

3.2 The case of multiple averaging functions

In above subsection, we treat the case of single averaging function. However, in practice, we often encounter the case of multiple averaging functions. Thus, the average sampled values can be described by $\{\langle f, \psi_{x_j,y_k}\rangle : j, k \in \mathbb{J}\}\.$ For this case, we recover the functions f exactly by using the following fast algorithm. Before giving the fast algorithm, we first define

$$
A_X f = \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} \langle f, \psi_{x_j, y_k} \rangle \beta_{j,k}.
$$

Theorem 3.4 *Assume that* $\Phi = (\phi_1, \phi_2, \dots, \phi_r)^T \in W_0(L^{1,1})^{(r)}$ whose support is compact and P is a bounded projection from $L^{p,q}$ onto $V_{p,q}(\Phi)$. Let the averaging sampling functions $\psi_{x_j, y_k} \in W(L^{1,1})$ satisfy $\int_{\mathbb{R}^{d+1}} \psi_{x_j, y_k} = 1$ and $\int_{\mathbb{R}^{d+1}} |\psi_{x_j, y_k}| \leq M$, where $M > 0$ is inde*pendent of* (x_j, y_k) *. Then there exist density* $\gamma_0 = \gamma_0(\Phi, M) > 0$ *and* $a_0 = a_0(\Phi, M) > 0$ such that if $X = \{(x_j, y_k) : j, k \in \mathbb{J}\}\$ is $\gamma (\gamma \leq \gamma_0)$ -dense in \mathbb{R}^{d+1} , and if the average $sampling$ functions ψ_{x_j, y_k} satisfy $\text{supp}\,\psi_{x_j, y_k} \subseteq (x_j, y_k) + [-a, a]^{d+1}$ for some $0 < a \le a_0$, *then any* $f \in V_{p,q}(\Phi)$ *can be recovered from its average samples* $\{ \langle f, \psi_{x_j, y_k} \rangle : j, k \in \mathbb{J} \}$ *by the following iterative algorithm:*

$$
\begin{cases}\nf_1 = P A_X f \\
f_{n+1} = P A_X (f - f_n) + f_n.\n\end{cases}
$$
\n(3.2)

In this case, the iterate f_n converges to f in the $L^{p,q}$ -norm. Moreover, the convergence is *geometric, that is,*

 $||f - f_n||_{L^{p,q}} \leq C\alpha^n$

for some $\alpha = \alpha(\gamma, a, P, \Phi, M) < 1$ *and* $C < \infty$.

4 Useful propositions and lemmas

In this section, we introduce some useful propositions and lemmas.

Let f be a continuous function. We define the oscillation (or modulus of continuity) of f by $\csc_{\delta}(f)(x_1, x_2) = \sup_{|y_1| \leq \delta, |y_2| \leq \delta} |f(x_1 + y_1, x_2 + y_2) - f(x_1, x_2)|.$

The following tow propositions are needed in the proof of two lemmas in this section.

Proposition 4.1 [\[8,](#page-13-5) Lemma 3.4] If $\phi \in W_0(L^{1,1})$ whose support is compact, then there $exists \ \delta_0 > 0 \ \text{such that } osc_{\delta}(\phi) \in W_0(L^{1,1}) \ \text{for any } \delta \leq \delta_0.$

Proposition 4.2 [\[8,](#page-13-5) Lemma 3.5] If $\phi \in W_0(L^{1,1})$ whose support is compact, then

$$
\lim_{\delta \to 0} \| osc_{\delta}(\phi)\|_{W(L^{1,1})} = 0.
$$

To prove our main results, we need the following Lemma.

Lemma 4.3 Let $\psi \in L^1(\mathbb{R}^{d+1})$ satisfying $\int_{\mathbb{R}^{d+1}} \psi(x) dx = 1$ and $\psi_a = (1/a^{d+1})\psi(\cdot/a)$. *Then for every* $\phi \in W_0(L^{1,1})$ *whose support is compact,*

$$
\|\phi - \phi * \psi_a^*\|_{W(L^{1,1})} \to 0 \quad as \quad a \to 0^+.
$$

Here a *is any positive real number.*

Proof. Since $\int_{\mathbb{R}^{d+1}} \psi(x)dx = 1$ and $\psi_a^*(x) = \overline{\psi_a(-x)}$, one has

$$
\phi - \phi * \psi_a^* = \int_{\mathbb{R}^{d+1}} (\phi(x) - \phi(x+t)) \overline{\psi_a(t)} dt.
$$

By Proposition [4.1,](#page-6-0) there exists $\delta_0 > 0$ such that $\csc(\phi) \in W_0(L^{1,1})$ for any $\delta \leq \delta_0$. Thus

$$
\|\phi - \phi * \psi_{a}^{*}\|_{W(L^{1,1})} \n= \sum_{k_{1} \in \mathbb{Z}} \sup_{x_{1} \in [0,1]} \sum_{k_{2} \in \mathbb{Z}^{d}} \sup_{x_{2} \in [0,1]^{d}} \sup_{k_{2} \in \mathbb{Z}^{d}} \sup_{x_{2} \in [0,1]^{d}} \left| \int_{\mathbb{R}} \int_{\mathbb{R}^{d}} (\phi(x_{1} + k_{1}, x_{2} + k_{2}) - \phi(x_{1} + k_{1} + t_{1}, x_{2} + k_{2} + t_{2})) \overline{\psi_{a}(t_{1}, t_{2})} dt_{2} dt_{1} \right| \n\leq \sum_{k_{1} \in \mathbb{Z}} \sup_{x_{1} \in [0,1]} \sum_{k_{2} \in \mathbb{Z}^{d}} \sup_{x_{2} \in [0,1]^{d}} \sup_{k_{2} \in \mathbb{Z}^{d}} \sup_{x_{2} \in [0,1]^{d}} \left| \phi(x_{1} + k_{1}, x_{2} + k_{2}) - \phi(x_{1} + k_{1} + t_{1}, x_{2} + k_{2} + t_{2}) \right| |\psi_{a}(t_{1}, t_{2})| dt_{2} dt_{1} \right| \n\leq \sum_{k_{1} \in \mathbb{Z}} \sup_{x_{1} \in [0,1]} \sum_{k_{2} \in \mathbb{Z}^{d}} \sup_{x_{2} \in [0,1]^{d}} \sup_{k_{2} \in \mathbb{Z}^{d}} \sup_{x_{2} \in [0,1]^{d}} \left| \phi(x_{1} + k_{1}, x_{2} + k_{2}) - \phi(x_{1} + k_{1} + t_{1}, x_{2} + k_{2} + t_{2}) \right| |\psi_{a}(t_{1}, t_{2})| dt_{2} dt_{1} \right| \n+ \sum_{k_{1} \in \mathbb{Z}} \sup_{x_{1} \in [0,1]} \sum_{k_{2} \in \mathbb{Z}^{d}} \sup_{x_{2} \in [0,1]^{d}} \sup_{k_{2} \in \mathbb{Z}^{d}} \left| \phi(x_{1} + k_{1}, x_{2} + k_{2}) - \phi(x_{1} + k_{1} + t_{1}, x_{2} + k_{2} + t_{2}) \right| |\psi_{a}(t_{1
$$

$$
\int_{|t_1|>\delta_0} \int_{|t_2| \le \delta_0} |\phi(x_1 + k_1, x_2 + k_2) - \phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2)| |\psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
+ \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} |\phi(x_1 + k_1, x_2 + k_2) - \phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2)| |\psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
= I_1 + I_2 + I_3,
$$

where

$$
I_1 = \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} \exp \left(\int_{|t_1| \le \delta_0} |\phi(x_1 + k_1, x_2 + k_2) - \phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2)| \right) |\psi_a(t_1, t_2)| dt_1 dt_1,
$$

$$
I_2 = \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} \exp \left(-\frac{1}{2} \int_{|t_1| > \delta_0} |\phi(x_1 + k_1, x_2 + k_2) - \phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2)| \, |\psi_a(t_1, t_2)| \, dt_2 dt_1 + \frac{1}{2} \int_{|t_1| > \delta_0} \left| \phi(x_1 + k_1, x_2 + k_2) - \phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2) \right| \, |\psi_a(t_1, t_2)| \, dt_2 dt_1 + \frac{1}{2} \int_{|t_1| > \delta_0} \left| \phi(x_1 + k_1, x_2 + k_2) - \phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2) \right| \, |\psi_a(t_1, t_2)| \, dt_2 dt_1
$$

and

$$
\begin{split} I_3 = & \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} \\ & \qquad \int\limits_{\mathbb{R}} \int\limits_{|t_2| > \delta_0} |\phi(x_1 + k_1, x_2 + k_2) - \phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2)| \left| \psi_a(t_1, t_2) \right| dt_2 dt_1. \end{split}
$$

First of all, we treat I_1 : let $|t| = \max\{|t_1|, |t_2|\}$. Then

$$
I_1 \leq \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} \int_{|t_1| \leq \delta_0} \int_{|t_2| \leq \delta_0} \csc_{|t|}(\phi)(x_1 + k_1, x_2 + k_2)) |\psi_a(t_1, t_2)| dt_2 dt_1
$$

\n
$$
\leq \int_{|t_1| \leq \delta_0} \int_{|t_2| \leq \delta_0} \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} \csc_{|t|}(\phi)(x_1 + k_1, x_2 + k_2)) |\psi_a(t_1, t_2)| dt_2 dt_1
$$

\n
$$
= \int_{|t_1| \leq \delta_0} \int_{|t_2| \leq \delta_0} |\csc_{|t|}(\phi) ||_{W(L^{1,1})} |\psi_a(t_1, t_2)| dt_2 dt_1.
$$

By proposition [4.2,](#page-6-1) for any $\epsilon>0,$ there exists $\delta_1>0\, (\delta_1<\delta_0)$ such that

$$
\|\text{osc}_{\delta}(\phi)\|_{W(L^{1,1})} < \epsilon, \quad \text{for any } \delta \le \delta_1.
$$

Write

$$
\int_{|t_1| \le \delta_0} \int_{|t_2| \le \delta_0} |\n\cos(\kappa_1(t_1))| \psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
= \int_{|t_1| \le \delta_1} \int_{|t_2| \le \delta_1} |\n\cos(\kappa_1(t_1))| \psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
+ \int_{\delta_1 < |t_1| \le \delta_0} \int_{|t_2| \le \delta_0} |\n\cos(\kappa_1(t_1))| \psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
+ \int_{|t_1| \le \delta_1} \int_{|t_2| \le \delta_0} |\n\cos(\kappa_1(t_1))| \psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
= I_4 + I_5 + I_6.
$$

Then

$$
I_4 \leq \int_{|t_1| \leq \delta_1} \int_{|t_2| \leq \delta_1} |\n\cos \xi_1(\phi)| |_{W(L^{1,1})} |\psi_a(t_1, t_2)| dt_2 dt_1
$$

$$
\leq \epsilon \int_{|t_1| \leq \delta_1} \int_{|t_2| \leq \delta_1} |\psi_a(t_1, t_2)| dt_2 dt_1 \leq \epsilon \|\psi\|_{L^1},
$$

$$
I_5 \leq \int_{\delta_1 < |t_1| \leq \delta_0} \int_{|b\infty \delta_0} \left\| \operatorname{osc}_{\delta_0}(\phi) \right\|_{W(L^{1,1})} \left| \psi_a(t_1, t_2) \right| dt_2 dt_1
$$
\n
$$
\leq \left\| \operatorname{osc}_{\delta_0}(\phi) \right\|_{W(L^{1,1})} \int_{\delta_1/a < |s_1|} \int_{s_2 \in \mathbb{R}^d} \left| \psi(s_1, s_2) \right| ds_2 ds_1
$$
\n
$$
\to 0 \quad \text{as } a \to 0^+
$$

and

$$
I_6 \leq \int_{|t_1| \leq \delta_1} \int_{\delta_1 < |t_2| \leq \delta_0} \|\csc_{|t|}(\phi)\|_{W(L^{1,1})} |\psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
\leq \|\csc_{\delta_0}(\phi)\|_{W(L^{1,1})} \int_{s_1 \in \mathbb{R}} \int_{\delta_1/a < |s_2|} |\psi(s_1, s_2)| ds_2 ds_1
$$
\n
$$
\to 0 \quad \text{as } a \to 0^+.
$$

Next, we treat I_2 :

$$
I_2 \leq \int_{|t_1| > \delta_0} \int_{|t_2| \leq \delta_0} \sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} |\phi(x_1 + k_1, x_2 + k_2)| |\psi_a(t_1, t_2)| dt_2 dt_1
$$

+
$$
\int_{|t_1| > \delta_0} \int_{|t_2| \leq \delta_0} |t_2| \leq \delta_0
$$

$$
\sum_{k_1 \in \mathbb{Z}} \sup_{x_1 \in [0,1]} \sum_{k_2 \in \mathbb{Z}^d} \sup_{x_2 \in [0,1]^d} |\phi(x_1 + k_1 + t_1, x_2 + k_2 + t_2)| |\psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
\leq 2 \int \int \int \|\phi\|_{W(L^{1,1})} |\psi_a(t_1, t_2)| dt_2 dt_1
$$
\n
$$
\leq 2 \|\phi\|_{W(L^{1,1})} \int \int \int \psi_a(t_1, t_2) |dt_2 dt_1
$$
\n
$$
|t_1| > \delta_0 |t_2| \leq \delta_0
$$
\n
$$
\leq 2 \|\phi\|_{W(L^{1,1})} \int \int \int \psi(a(t_1, t_2) | ds_2 ds_1)
$$
\n
$$
|s_1| > \delta_0 / a t_2 \in \mathbb{R}^d
$$
\n
$$
\to 0 \quad \text{as } a \to 0^+.
$$

Similarly, we can prove $I_3 \to 0$ as $a \to 0^+$. This completes the proof of Lemma [4.3.](#page-6-2) \Box The following lemma is a generalization of [\[30,](#page-15-1) Lemma 4.1].

Lemma 4.4 Let X be any sampling set which is γ -dense in \mathbb{R}^{d+1} , let $\{\beta_{j,k} : j,k \in \mathbb{J}\}$ be a *bounded uniform partition of unity associated with* X, and let $\phi \in W_0(L^{1,1})$ whose support *is compact. Then there exist constants* C and γ_0 *such that for any* $f = \sum_{k \in \mathbb{Z}^{d+1}} c_k \phi(\cdot - k)$ *and* $\gamma \leq \gamma_0$ *, one has*

$$
||Q_X f||_{L^{p,q}} \leq C||c||_{\ell^{p,q}}||\phi||_{W(L^{1,1})} \quad \text{for any } c = \{c_k : k \in \mathbb{Z}^{d+1}\} \in \ell^{p,q}.
$$

To prove the Lemma [4.4,](#page-9-0) we introduce the following proposition.

Proposition 4.5 [\[7,](#page-13-7) Theorem 3.1] Assume that $1 \leq p, q < \infty$ and $\phi \in W(L^{1,1})$. Then *for any* $c \in \ell^{p,q}$, the function $f = \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} c(k_1, k_2) \phi(\cdot - k_1, \cdot - k_2)$ *belongs to* $L^{p,q}$ *and*

$$
||f||_{L^{p,q}} \leq ||c||_{\ell^{p,q}} ||\phi||_{W(L^{1,1})}.
$$

Proof of Lemma [4.4](#page-9-0). By Proposition [4.5](#page-9-1) and Proposition [4.1,](#page-6-0) one has that there exists $\gamma_0 > 0$ such that for any $\gamma \leq \gamma_0$

$$
||f - Q_X f||_{L^{p,q}} \le ||\text{osc}_{\gamma}(f)||_{L^{p,q}} \le ||c||_{\ell^{p,q}} ||\text{osc}_{\gamma}(\phi)||_{W(L^{1,1})}.
$$

Using Proposition [4.5](#page-9-1) and the proof of $[8, \text{ Lemma } 3.4]$, one obtains that there exists C' such that

$$
||Q_Xf||_{L^{p,q}} = ||f - Q_Xf + f||_{L^{p,q}} \le ||f - Q_Xf||_{L^{p,q}} + ||f||_{L^{p,q}}
$$

\n
$$
\le ||c||_{\ell^{p,q}}||\csc_{\gamma}(\phi)||_{W(L^{1,1})} + ||c||_{\ell^{p,q}} ||\phi||_{W(L^{1,1})}
$$

\n
$$
\le ||c||_{\ell^{p,q}}C'||\phi||_{W(L^{1,1})} + ||c||_{\ell^{p,q}} ||\phi||_{W(L^{1,1})}
$$

\n
$$
\le C||c||_{\ell^{p,q}} ||\phi||_{W(L^{1,1})},
$$

where $C = 1 + C'$

The following lemma plays an important role in the proof Theorem [3.3.](#page-5-0)

Lemma 4.6 Let $\Phi = (\phi_1, \phi_2, \dots, \phi_r)^T \in W_0(L^{1,1})^{(r)}$ whose support is compact and P be *a* bounded projection from $L^{p,q}(\mathbb{R}^{d+1})$ onto $V_{p,q}(\Phi)$. Then there exist $\gamma_0 > 0$ and $a_0 > 0$ *such that for* γ -dense set X with $\gamma \leq \gamma_0$ and for every positive real number $a \leq a_0$, the *operator* $I - PA_{X,a}$ *is a contraction operator on* $V_{p,q}(\Phi)$ *.*

. В последните последните последните последните последните последните последните последните последните последн
В 1990 година от селото на сел

Proof. Putting $f = \sum_{i=1}^r \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} c_i(k_1, k_2) \phi_i(\cdot - k_1, \cdot - k_2) \in V_{p,q}(\Phi)$. Then one has

$$
||f - PA_{X,a}f||_{L^{p,q}} = ||f - PQ_Xf + PQ_Xf - PA_{X,a}f||_{L^{p,q}}
$$

\n
$$
\leq ||f - PQ_Xf||_{L^{p,q}} + ||PQ_Xf - PA_{X,a}f||_{L^{p,q}}
$$

\n
$$
= ||Pf - PQ_Xf||_{L^{p,q}} + ||PQ_Xf - PA_{X,a}f||_{L^{p,q}}
$$

\n
$$
\leq ||P||_{op} (||f - Q_Xf||_{L^{p,q}} + ||Q_Xf - A_{X,a}f||_{L^{p,q}})
$$

\n
$$
= ||P||_{op} (||f - Q_Xf||_{L^{p,q}} + ||Q_Xf - Q_X(f * \psi_a^*)||_{L^{p,q}}).
$$

Firstly, we estimate the first term of the last inequality. By Proposition [4.5,](#page-9-1) Proposition [2.4](#page-3-0) and Cauchy inequality, one has that there exists γ_1 such that for any $\gamma \leq \gamma_1$

$$
||f - Q_X f||_{L^{p,q}} \leq \left\| \sum_{i=1}^r \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} |c_i(k_1, k_2)| \text{osc}_{\gamma}(\phi_i)(\cdot - k_1, \cdot - k_2) \right\|_{L^{p,q}} \leq \sum_{i=1}^r \left\| \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} |c_i(k_1, k_2)| \text{osc}_{\gamma}(\phi_i)(\cdot - k_1, \cdot - k_2) \right\|_{L^{p,q}} \leq \sum_{i=1}^r \|c_i\|_{\ell^{p,q}} \|\text{osc}_{\gamma}(\phi_i)\|_{W(L^{1,1})} \leq \max_{1 \leq i \leq r} \|\text{osc}_{\gamma}(\phi_i)\|_{W(L^{1,1})} \sum_{i=1}^r \|c_i\|_{\ell^{p,q}} \leq \max_{1 \leq i \leq r} \|\text{osc}_{\gamma}(\phi_i)\|_{W(L^{1,1})} \sqrt{r} \left(\sum_{i=1}^r \|c_i\|_{\ell^{p,q}}^2 \right)^{1/2} \leq D_2 \sqrt{r} \max_{1 \leq i \leq r} \|\text{osc}_{\gamma}(\phi_i)\|_{W(L^{1,1})} \|f\|_{L^{p,q}}.
$$
\n(4.1)

Next, we estimate the second term of the last inequality. Let $\phi_i^a = \phi_i - \phi_i * \psi_a^*$ (i = $(1, \dots, r)$. Using $\phi_i \in W_0(L^{1,1})$ $(i = 1, \dots, r)$, $\psi \in L^1$ and Lemma [2.3,](#page-2-0) it follows that $\phi_i^a \in W_0(L^{1,1})$. Note that $f - f * \psi_a^* = \sum_{i=1}^r \sum_{k \in \mathbb{Z}^{d+1}} c_i(k) \phi_i^a(\cdot - k)$. By Lemma [4.4,](#page-9-0) there exist constants $C, \gamma_2 > 0$ such that for any $\gamma \leq \gamma_2$

$$
\|Q_X f - Q_X (f * \psi_a^*)\|_{L^{p,q}} \leq \left\| Q_X \left(\sum_{i=1}^r \sum_{k \in \mathbb{Z}^{d+1}} c_i(k) \phi_i^a(\cdot - k) \right) \right\|_{L^{p,q}}
$$

$$
= \left\| \sum_{i=1}^r Q_X \left(\sum_{k \in \mathbb{Z}^{d+1}} c_i(k) \phi_i^a(\cdot - k) \right) \right\|_{L^{p,q}}
$$

$$
= \sum_{i=1}^r \left\| Q_X \left(\sum_{k \in \mathbb{Z}^{d+1}} c_i(k) \phi_i^a(\cdot - k) \right) \right\|_{L^{p,q}}
$$

$$
\leq C \sum_{i=1}^r \|c_i\|_{\ell^{p,q}} \|\phi_i^a\|_{W(L^{1,1})}.
$$

Using Proposition [2.4](#page-3-0) and Cauchy inequality

$$
\|Q_X f - Q_X (f * \psi_a^*)\|_{L^{p,q}} \leq C \max_{1 \leq i \leq r} \|\phi_i^a\|_{W(L^{1,1})} \sum_{i=1}^r \|c_i\|_{\ell^{p,q}}
$$

$$
\leq C \max_{1 \leq i \leq r} \|\phi_i^a\|_{W(L^{1,1})} \sqrt{r} \left(\sum_{i=1}^r \|c_i\|_{\ell^{p,q}}^2\right)^{1/2}
$$

$$
\leq D_2 \sqrt{r} C \max_{1 \leq i \leq r} \|\phi_i^a\|_{W(L^{1,1})} \|f\|_{L^{p,q}}.
$$

Assume that $\epsilon > 0$ is any positive real number. By Proposition [4.2,](#page-6-1) there exists γ_3 such that $D_2\sqrt{r}\max_{1\leq i\leq r} \|\csc_\gamma(\phi_i)\|_{W(L^{1,1})} \leq \epsilon/2$ for any $\gamma \leq \gamma_3$. Using Lemma [4.3,](#page-6-2) there exists $a_0 > 0$ such that for any $a \le a_0$

$$
D_2\sqrt{r}C\max_{1\leq i\leq r}\|\phi_i^a\|_{W(L^{1,1})}\leq\epsilon/2.
$$

Hence, we choose $\gamma_0 = \min{\{\gamma_1, \gamma_2, \gamma_3\}}$, one has

$$
||f - PA_{X,a}f||_{L^{p,q}} \le \epsilon ||P||_{op} ||f||_{L^{p,q}}
$$
 for any $f \in V_{p,q}(\Phi)$, $\gamma \le \gamma_0$, $a \le a_0$.

To get a contraction operator, we choose $\epsilon ||P||_{op} < 1$.

5 Proofs of main results

In this section, we give proofs of Theorem [3.3](#page-5-0) and Theorem [3.4.](#page-5-1)

5.1 Proof of Theorem [3.3](#page-5-0)

For convenience, let $e_n = f - f_n$ be the error after *n* iterations. Using [\(3.1\)](#page-5-2),

$$
e_{n+1} = f - f_{n+1}
$$

= $f - f_n - PA_{X,a}(f - f_n)$
= $(I - PA_{X,a})e_n$.

Using Lemma [4.6,](#page-9-2) we may choose right γ_0 and a_0 such that for any $\gamma \leq \gamma_0$ and $a \leq a_0$

$$
||I - PA_{X,a}||_{op} = \alpha < 1.
$$

Then we obtain

$$
||e_{n+1}||_{L^{p,q}} \leq \alpha ||e_n||_{L^{p,q}} \leq \alpha^n ||e_1||_{L^{p,q}}.
$$

Wherewith $||e_n||_{L^{p,q}} \to 0$, when $n \to \infty$. This completes the proof.

5.2 Proof of Theorem [3.4](#page-5-1)

We only need to prove $I - PA_X$ is a contraction operator on $V_{p,q}(\Phi)$.

Let
$$
f = \sum_{i=1}^{r} \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} c_i(k_1, k_2) \phi_i(\cdot - k_1, \cdot - k_2) \in V_{p,q}(\Phi)
$$
. One has
\n
$$
||f - PA_Xf||_{L^{p,q}} = ||f - PQ_Xf + PQ_Xf - PA_Xf||_{L^{p,q}}
$$
\n
$$
\leq ||f - PQ_Xf||_{L^{p,q}} + ||PQ_Xf - PA_Xf||_{L^{p,q}}
$$
\n
$$
= ||Pf - PQ_Xf||_{L^{p,q}} + ||PQ_Xf - PA_Xf||_{L^{p,q}}
$$
\n
$$
\leq ||P||_{op} (||f - Q_Xf||_{L^{p,q}} + ||Q_Xf - A_Xf||_{L^{p,q}}). \tag{5.1}
$$

Putting $f_i = \sum_{k_1 \in \mathbb{Z}} \sum_{k_2 \in \mathbb{Z}^d} c_i(k_1, k_2) \phi_i(\cdot - k_1, \cdot - k_2), (1 \leq i \leq r)$. Then $f_i \in V^{p,q}(\phi_i)$ for $1 \leq i \leq r$ and $f = \sum_{i=1}^{r} f_i$. For each f_i , one has

$$
|Q_X f_i - A_X f_i| = \left| \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} (f_i(x_j, y_k) - \langle f_i, \psi_{x_j, y_k} \rangle) \beta_{j,k} \right|
$$

\n
$$
= \left| \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} \left(\int_{\mathbb{R}^{d+1}} (f_i(x_j, y_k) - f_i(t)) \overline{\psi_{x_j, y_k}(t)} dt \right) \beta_{j,k} \right|
$$

\n
$$
\leq \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} \int_{\mathbb{R}^{d+1}} |f_i(x_j, y_k) - f_i(t)| |\psi_{x_j, y_k}(t)| dt \beta_{j,k}
$$

\n
$$
\leq \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} \text{osc}_{a}(f_i)(x_j, y_k) \int_{\mathbb{R}^{d+1}} |\psi_{x_j, y_k}(t)| dt \beta_{j,k}
$$

\n
$$
\leq M \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} \text{osc}_{a}(f_i)(x_j, y_k) \beta_{j,k}
$$

\n
$$
\leq M \sum_{j \in \mathbb{J}} \sum_{k \in \mathbb{J}} \sum_{k \in \mathbb{Z}} \sum_{k \in \mathbb{Z}^{d}} |c_i(k_1, k_2)| \text{osc}_{a}(\phi_i)(x_j - k_1, y_k - k_2) \beta_{j,k}.
$$

By Proposition [4.5](#page-9-1) and Proposition [4.1,](#page-6-0) there exists a_1 such that for any $a \le a_1$

$$
||Q_X f_i - A_X f_i||_{L^{p,q}} \leq MC||c_i||_{\ell^{p,q}}||\text{osc}_a(\phi_i)||_{W(L^{1,1})}.
$$

Therewith

$$
||Q_X f - A_X f||_{L^{p,q}} \leq MC \sum_{i=1}^r ||c_i||_{\ell^{p,q}} ||\operatorname{osc}_a(\phi_i)||_{W(L^{1,1})}.
$$

Thus by the proof of Lemma [4.6](#page-9-2)

$$
||Q_X f - A_X f||_{L^{p,q}} \leq MCD_2 \sqrt{r} \max_{1 \leq i \leq r} ||\csc_a(\phi_i)||_{W(L^{1,1})} ||f||_{L^{p,q}}.
$$
\n(5.2)

Using [\(4.1\)](#page-10-0), [\(5.1\)](#page-12-0) and [\(5.2\)](#page-12-1), one has that there exists γ_1 such that for any $\gamma \leq \gamma_1$ and $a\leq a_1$

$$
||f - PA_Xf||_{L^{p,q}} \le ||P||_{op}(D_2\sqrt{r} \max_{1 \le i \le r} ||\text{osc}_{\gamma}(\phi_i)||_{W(L^{1,1})}
$$

$$
+ MCD_2\sqrt{r} \max_{1 \le i \le r} ||\text{osc}_{a}(\phi_i)||_{W(L^{1,1})})||f||_{L^{p,q}}.
$$

Assume that $\epsilon > 0$ is any positive real number. By Proposition [4.2,](#page-6-1) there exists γ_2 such that for any $\gamma \leq \gamma_2$

$$
D_2\sqrt{r}\max_{1\leq i\leq r} \|\text{osc}_{\gamma}(\phi_i)\|_{W(L^{1,1})} \leq \epsilon/2.
$$

Using Lemma [4.3,](#page-6-2) there exists $a_2 > 0$ such that for any $a \le a_2$

$$
MCD_2\sqrt{r}\max_{1\leq i\leq r}\|\csc_a(\phi_i)\|_{W(L^{1,1})}\leq \epsilon/2.
$$

Hence, we choose $\gamma_0 = \min\{\gamma_1, \gamma_2\}$ and $a_0 = \min\{a_1, a_2\}$, one has

$$
||f - PA_Xf||_{L^{p,q}} \le \epsilon ||P||_{op} ||f||_{L^{p,q}}
$$
 for any $f \in V_{p,q}(\Phi)$, $\gamma \le \gamma_0$, $a \le a_0$.

To get a contraction operator, we choose $\epsilon ||P||_{op} < 1$.

6 Conclusion

In this paper, we study nonuniform average sampling problem in multiply generated shift-invariant subspaces of mixed Lebesgue spaces. We discuss two types of average sampled values. Two fast reconstruction algorithms for this two types of average sampled values are provided. Studying $L^{p,q}$ -frames in multiply generated shift-invariant subspaces of mixed Lebesgue spaces is the goal of future work.

Acknowledgements

This work was supported partially by the National Natural Science Foundation of China (11326094 and 11401435).

References

- [1] Benedek, A., Panzone, R.: The space L^p with mixed norm. Duke Math. J. 28(3), 301–324 (1961)
- [2] Benedek, A., Calderón, A.P., Panzone, R.: Convolution operators on Banach space valued functions. Proc. Nat. Acad. Sci. USA 48(3), 356–365 (1962)
- [3] Fernandez, D.L.: Vector-valued singular integral operators on L^p -spaces with mixed norms and applications. Pac. J. Math. 129(2), 257–275 (1987)
- [4] Francia, J.L., Ruiz, F.J., Torrea, J.L.: Calder´on–Zygmund theory for operatorvalued kernels. Adv. Math. 62(1), 7–48 (1986)
- [5] Torres, R., Ward. E.: Leibniz's Rule, Sampling and wavelets on mixed Lebesgue spaces. J. Fourier Anal. Appl. 21(5), 1053–1076 (2015)
- [6] Ward, E.: New estimates in harmonic analysis for mixed Lebesgue spaces. Ph.D. Thesis, University of Kansas (2010)
- [7] Li, R., Liu, B., Liu, R., Zhang, Q.: Nonuniform sampling in principal shift-invariant subspaces of mixed Lebesgue spaces $L^{p,q}(\mathbb{R}^{d+1})$. J. Math. Anal. Appl. 453(2), 928– 941 (2017)
- [8] Zhang, Q.: Reconstruction of functions in principal shift-invariant subspaces of mixed Lebesgue spaces. [arXiv:1801.05715](http://arxiv.org/abs/1801.05715)
- [9] Shannon, C.: Communication in the presence of noise. Proc. IRE 37(1), 10–21 (1949)
- [10] Shannon, C.: Communication in the presence of noise. Proc. IEEE 72(9), 447–457 (1998)
- [11] Sun, Q.: Nonuniform average sampling and reconstruction of signals with finite rate of innovation. SIAM J. Math. Anal. 38(5), 1389–1422 (2006)
- [12] Benedetto, J.J., Ferreira, P.J.S.G.: Modern sampling theory, Mathematics and applications. Birkhäuser, Boston (2001)
- [13] Marvasti, F.: Nonuniform sampling, theory and practice. Kluwer Academic/Plenum Publishers, New York (2001)
- [14] Venkataramani, R., Bresler, Y.: Sampling theorems for uniform and periodic nonuniform MIMO sampling of multiband signals. IEEE Trans. Signal Process. 51(12), 3152–3163 (2003)
- [15] Heil, C.: Harmonic analysis and applications. Birkhäuser, Boston (2006)
- [16] Sun, W., Zhou, X.: Sampling theorem for multiwavelet subspaces. Chinese Sci. Bull. 44(14), 1283–1286 (1999)
- [17] Zhou, X., Sun, W.: On the sampling theorem for wavelet subspaces. J. Fourier Anal. Appl. 5(4), 347–354 (1999)
- [18] Aldroubi, A., Feichtinger, H.: Exact iterative reconstruction algorithm for multivariate irregularly sampled functions in spline-like spaces: The L^p theory. Proc. Amer. Math. Soc. 126(9), 2677–2686 (1998)
- [19] Aldroubi, A., Gröchenig, K.: Beurling-Landau-type theorems for nonuniform sampling in shift-invariant spaces. J. Fourier Anal. Appl. 6(1), 91–101 (2000)
- [20] Aldroubi, A., Gröchenig, K.: Nonuniform sampling and reconstruction in shiftinvariant spaces. SIAM Rev. 43(4), 585–562 (2001)
- [21] Aldroubi, A., Unser, M.: Sampling procedure in function spaces and asymptotic equivalence with Shannon's sampling theory. Numer. Funct. Anal. Optim. 15(1), 1–21 (1994)
- [22] Vaidyanathan, P.P., Vrcelj, B.: Biorthogonal partners and applications. IEEE Trans. Signal Process. 49, 1013–1027 (2001)
- [23] Zhang, Q., Sun, W.: Invariance of shift invariant spaces. Sci. China Seri. A 55(7), 1395–1401 (2012)
- [24] Sun, W., Zhou, X.: Average sampling in spline subspaces. Appl. Math. Lett., 15, 233–237 (2002)
- [25] Pérez-Villalón, G., Portal, A.: Reconstruction of splines from local average samples. Appl. Math. Lett., 25, 1315– 1319 (2012)
- [26] Ponnaian, D., Shanmugam, Y.: Existence and uniqueness of spline reconstruction from local weighted average samples. Rend. Circ. Mat. Palermo., 63, 97–108 (2014)
- [27] Kang, S., Kwon, K.H.: Generalized average sampling in shift invariant spaces. J. Math. Anal. Appl., 377, 70–78 (2011)
- [28] Atreas, N.D.: On a class of non-uniform average sampling expansions and partial reconstruction in subspaces of $L^2(\mathbb{R})$. Adv. Comput. Math., 36, 21-38 (2012)
- [29] Aldroubi, A.: Non-uniform weighted average sampling and reconstruction in shiftinvarant and wavelet spaces. Appl. Comput. Harmon. Anal., 13, 152–161 (2002)
- [30] Aldroubi, A., Sun, Q., Tang, W.S.: Nonuniform average sampling and reconstruction in multiply generated shift-invariant spaces. Constr. Approx., 20, 173–189 (2004)
- [31] Xian, J., Li, S.: General A-P iterative algorithm in shift-invariant spaces. Acta Math. Sin. Engl. Ser., 25, 545–552 (2009)
- [32] Sun, Q.: Nonuniform average sampling and reconstruction of signals with finite rate of innovation. SIAM J. Math. Anal., 38, 1389–1422 (2006)
- [33] Jia, R.Q., Micchelli, C.A.: Using the refinement equation for the construction of pre-wavelets II: Powers of two. P.J. Laurent, A. Le Mfhaut6 and L.L. Schumaker (Eds.), Curves and Surfaces, Academic Press, New York, 209–246(1991)