

ON DIRICHLET'S LAMBDA FUNCTION

SU HU AND MIN-SOO KIM

ABSTRACT. Let

$$\lambda(s) = \sum_{n=0}^{\infty} \frac{1}{(2n+1)^s},$$

$$\beta(s) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^s},$$

and

$$\eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s}$$

be the Dirichlet lambda function, its alternating form, and the Dirichlet eta function, respectively. According to a recent historical book by Varadarajan ([25, p. 70]), these three functions were investigated by Euler under the notations $N(s)$, $L(s)$, and $M(s)$, respectively.

In this paper, we shall present some additional properties for them. That is, we obtain a number of infinite families of linear recurrence relations for $\lambda(s)$ at positive even integer arguments $\lambda(2m)$, convolution identities for special values of $\lambda(s)$ at even arguments and special values of $\beta(s)$ at odd arguments, and a power series expansion for the alternating Hurwitz zeta function $J(s, a)$, which involves a known one for $\eta(s)$.

1. INTRODUCTION

1.1. Background. According to Varadarajan [25, p. 59], Pietro Mengoli (1625–1686) posed the problem of finding the sum of the series

$$(1.1) \quad \sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \cdots.$$

This problem was first solved by Euler, who communicated his findings in a letter to Daniel Bernoulli, asserting that

$$(1.2) \quad 1 + \frac{1}{4} + \frac{1}{9} + \cdots = \frac{\pi^2}{6},$$

more generally, let

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

2000 *Mathematics Subject Classification.* 11B68, 11S40.

Key words and phrases. Dirichlet lambda function, Dirichlet eta function, Recurrences, Euler polynomials.

be Riemann's zeta function. Then, Euler proved the following formula:

$$(1.3) \quad \zeta(2m) = 1 + \frac{1}{2^{2m}} + \frac{1}{3^{2m}} + \cdots = \frac{(-1)^{m-1} B_{2m} 2^{2m}}{2(2m)!} \pi^{2m},$$

where the B_{2m} are the Bernoulli numbers, defined by the generating function

$$(1.4) \quad \frac{t}{e^t - 1} = 1 - \frac{t}{2} + \sum_{m=1}^{\infty} B_{2m} \frac{t^{2m}}{(2m)!}.$$

One of Euler's proofs of (1.2) from around 1742 is based on the following infinite product expansion of $\frac{\sin x}{x}$:

$$(1.5) \quad \frac{\sin x}{x} = \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2 \pi^2} \right).$$

Euler also provided another proof of (1.2) by starting from the formula

$$(1.6) \quad \frac{1}{2}(\arcsin x)^2 = \int_0^x \frac{\arcsin t}{\sqrt{1-t^2}} dt.$$

First by taking $x = 1$ in the left-hand side, we get $\frac{\pi^2}{8}$. By expanding $\arcsin t$ as a power series and integrating term-by-term on the right-hand side, we obtain the following summation:

$$1 + \frac{1}{3^2} + \frac{1}{5^2} + \cdots = \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2}.$$

After comparing the results on both sides, we have

$$(1.7) \quad \sum_{n=0}^{\infty} \frac{1}{(2n+1)^2} = \frac{\pi^2}{8}.$$

Then, by noticing that

$$(1.8) \quad 1 + \frac{1}{3^2} + \frac{1}{5^2} + \cdots = \zeta(2) - \frac{1}{2^2} \zeta(2) = \frac{3}{4} \zeta(2),$$

we recover (1.2) (see [25, pp. 62–63]).

According to Abramowitz and Stegun's handbook [1, pp. 807–808], the function

$$(1.9) \quad \lambda(s) = \sum_{n=0}^{\infty} \frac{1}{(2n+1)^s} = (1 - 2^{-s}) \zeta(s), \quad \operatorname{Re}(s) > 1,$$

is usually named the Dirichlet lambda function, and was also studied by Euler under the notation $N(s)$ (see [25, p. 70]). In addition, (1.7) provided the special value of $\lambda(s)$ at 2, and the special values of $\lambda(s)$ at any even positive integer $2m$ are calculated by the following formula:

$$(1.10) \quad \lambda(2m) = (-1)^m \frac{\pi^{2m}}{4(2m-1)!} E_{2m-1}(0), \quad m \geq 1,$$

where the Euler polynomials $E_n(x)$ are defined by the following generating function:

$$(1.11) \quad \frac{2}{e^t + 1} e^{tx} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}$$

(see [6, (4.16)] and [11, (2.15)]). The integers $E_n = 2^n E_n(1/2)$, $n \geq 0$, are called Euler numbers, and can also be defined as the coefficients of $t^n/n!$ in the Taylor expansion of $\operatorname{sech}(t)$, $|t| < \pi/2$. For example, $E_0 = 1$, $E_2 = -1$, $E_4 = 5$, and $E_6 = -61$. Euler numbers and polynomials (so called by Scherk in 1825) were introduced in Euler's famous book, *Insitutiones Calculi Differentialis* (1755, pp. 487–491 and p. 522).

Recently, the Euler polynomials have been employed in several different applications, such as semi-classical approximations of quantum probability distributions (see [3]), and various approximations and expansion formulas in discrete mathematics and number theory (see [1, 19]). Euler polynomials can be defined using various methods, depending on their applications (see [9, 24]). For example, the explicit formula

$$(1.12) \quad E_n(x) = \sum_{k=0}^n \binom{n}{k} x^{n-k} E_k(0)$$

shows that $E_n(x)$ is a polynomial of degree n . There are various well-known approaches to the theory of Euler polynomials, and these can be found in the classical papers by Euler [7], Nörlund [18], and Raabe [20].

It may be interesting to note that there is also a connection between the generalized Euler numbers and the ideal class group of the p^{n+1} -th cyclotomic field when p is a prime number. For details, we refer the reader to the recent paper [10], especially [10, Proposition 3.4].

In addition, Euler also studied the function

$$(1.13) \quad \beta(s) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^s}, \quad \operatorname{Re}(s) > 0,$$

($L(s)$ in his notation: see [25, p. 70]). This is the alternating form of the Dirichlet lambda function $\lambda(s)$ (1.9). Furthermore, the constant $\beta(2) = G$ is usually named Catalan's constant (see [21], [27], [1, p. 807], and [8, p. 53]). It is known that the special values of $\beta(s)$ at odd positive integers $2m+1$ are given by

$$(1.14) \quad \beta(2m+1) = (-1)^m \frac{E_{2m}}{2(2m)!} \left(\frac{\pi}{2}\right)^{2m+1},$$

where E_{2m} are the Euler numbers, and $\beta(s)$ admits the following interesting integral representation using a trigonometric function:

$$\beta(s) = \frac{1}{2\Gamma(s)} \int_0^{\infty} \frac{t^{s-1}}{\cosh(t)}, \quad \operatorname{Re}(s) > 0,$$

where Γ denotes the gamma function (see [8, p. 56]).

1.2. Main results. Dating back to Euler, the study of the special values of the Riemann zeta function at positive integers has a long history; hence there is a vast collection of related literature. Here, we mention some that are directly related to our study.

In 1987, Song [23] found the following linear recurrence relation for Riemann's zeta function at even arguments $\zeta(2m)$, using the Fourier series expansion of periodic functions:

$$(-1)^{m+1} \frac{\pi^{2m} \cdot m}{(2m+1)!} + \sum_{k=1}^{m-1} \frac{\pi^{2k}}{(2k+1)!} \zeta(2m-2k) = 0.$$

Recently, Merca [15] obtained an alternative proof for the above equality using the generating function of Bernoulli numbers, and [16] also obtained a homogeneous linear recurrence relation for $\zeta(2m)$ using several tools from symmetric function theory. Extending the above results, in 2017 Merca [17] also introduced a number of infinite families of linear recurrence relations between $\zeta(2m)$.

In 2013, Lettington ([13], [14, (1.24)]) proved the following linear recurrence relation between the special values $\lambda(2m)$ of the Dirichlet lambda function $\lambda(s)$.

Theorem 1.1 (Lettington). *Let m be a positive integer. Then*

$$\lambda(2m) = (-1)^{m-1} \left(\frac{\pi^{2m}}{4(2m)!} + \sum_{k=1}^{m-1} \frac{(-1)^{m-k} \pi^{2k}}{(2k+1)!} \lambda(2m-2k) \right).$$

In this study, based on the integral representation of

$$(1.15) \quad J(s, a) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+a)^s}, \quad \operatorname{Re}(s) > 0,$$

(see (2.3)) and an analogue of the Hermite formula for $J(s, a)$ (see (2.4)), we provide a new proof of Theorem 1.1.

Analogous to Merca's work [17], we further prove a number of infinite families of linear recurrence relations for $\lambda(2m)$ using the generating function of the Euler polynomials and the Euler-type formula (1.10). (See Theorems 1.2, 1.3, 1.8, 1.9, and 1.10 and Corollaries 1.4, 1.5, and 1.7).

Moreover, we prove convolution identities for the special values of $\lambda(s)$ at even arguments and for special values of $\zeta(s)$ and $\beta(s)$ at odd arguments. (See Theorems 5.1 and 5.2).

Finally, in Proposition 6.1 below, we demonstrate a power series expansion for the function $J(s, a)$ (1.15), which implies a known one from Coffey for the Dirichlet eta functions

$$\eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s}, \quad \operatorname{Re}(s) > 0,$$

(see Corollary 6.3).

In the following, we shall employ the usual convention that an empty sum is taken to be zero. For example, if $m = 1$, then we understand that $\sum_{k=1}^{m-1} = 0$.

1.3. Linear recurrence relations for the lambda function at even integers.

Theorem 1.2. *Let m be a positive integer, and α a complex number such that $\alpha \neq \frac{1}{2}$. Then we have*

$$\begin{aligned} & (-1)^{m+1} \frac{\alpha^{2m} - (\alpha - 1)^{2m}}{4} \cdot \frac{\pi^{2m+2}}{(2m)!} \\ & + \sum_{k=1}^m (-1)^k (\alpha^{2k-1} + (\alpha - 1)^{2k-1}) \frac{\pi^{2k}}{(2k-1)!} \lambda(2m - 2k + 2) = 0. \end{aligned}$$

Theorem 1.3. *Let m be a nonnegative integer, and α a complex number. Then we have*

$$\begin{aligned} & (-1)^{m+1} \frac{\alpha^{2m+1} - (\alpha - 1)^{2m+1}}{4} \cdot \frac{\pi^{2m+2}}{(2m+1)!} \\ & + \sum_{k=0}^m (-1)^k (\alpha^{2k} + (\alpha - 1)^{2k}) \frac{\pi^{2k}}{(2k)!} \lambda(2m - 2k + 2) = 0, \end{aligned}$$

where define $0^0 = 1$.

Let us state some special cases of Theorem 1.3.

Corollary 1.4 ($\alpha = 1$). *Let m be a nonnegative integer. Then we have*

$$(-1)^{m+1} \frac{\pi^{2m+2}}{4(2m+1)!} + \sum_{k=0}^m (-1)^k (1 + \delta_k) \frac{\pi^{2k}}{(2k)!} \lambda(2m - 2k + 2) = 0,$$

where $\delta_k = 0$ if $k \neq 0$ and 1 if $k = 0$.

Corollary 1.5 ($\alpha = \frac{1}{2}$). *Let m be a nonnegative integer. Then we have*

$$(-1)^{m+1} \frac{\pi^{2m+2}}{4(2m+1)!} + \sum_{k=0}^m (-1)^k \frac{2^{2m-2k+1} \pi^{2k}}{(2k)!} \lambda(2m - 2k + 2) = 0.$$

Remark 1.6. *If m is a positive integer, then from Corollary 1.5 we have the following result:*

$$\lambda(2m) = (-1)^{m-1} \left(\frac{\pi^{2m}}{2^{2m+1} (2m-1)!} + \sum_{k=1}^{m-1} (-1)^{m-k} \frac{\pi^{2k}}{2^{2k} (2k)!} \lambda(2m - 2k) \right),$$

which is an analogue of a result by Lettington (see Theorem 1.1). We observe that this linear recurrence relation does not require a priori knowledge on the Euler numbers. If m is small, then it is easy to evaluate $\lambda(2m)$ as follows:

$$\lambda(2) = \frac{\pi^2}{8}, \quad \lambda(4) = \frac{\pi^4}{96}, \quad \lambda(6) = \frac{\pi^6}{960}, \quad \lambda(8) = \frac{17\pi^8}{161280}.$$

Corollary 1.7. *Let n be a positive integer.*

(1) For integers $m > 0$, we have

$$\begin{aligned} & \frac{(-1)^{m+1}}{4} \left(2 \sum_{j=1}^{n-1} (-1)^j j^{2m} + (-1)^n n^{2m} \right) \frac{\pi^{2m+2}}{(2m)!} \\ & + (-1)^n \sum_{k=1}^m (-1)^k n^{2k-1} \frac{\pi^{2k}}{(2k-1)!} \lambda(2m-2k+2) = 0. \end{aligned}$$

(2) For integers $m \geq 0$, we have

$$\begin{aligned} & \frac{(-1)^{m+1}}{4} \left(2 \sum_{j=1}^{n-1} (-1)^j j^{2m+1} + (-1)^n n^{2m+1} \right) \frac{\pi^{2m+2}}{(2m+1)!} \\ & + \sum_{k=0}^m (-1)^k ((-1)^n n^{2k} - \delta_k) \frac{\pi^{2k}}{(2k)!} \lambda(2m-2k+2) = 0. \end{aligned}$$

Theorem 1.8. Let m be a positive integer, and α a complex number. Then we have

$$\begin{aligned} & (-1)^m \frac{\pi^{2m+2}}{4(2m)!} ((\alpha-1)^{2m} + (\alpha+1)^{2m} - 2\alpha^{2m}) \\ & + \sum_{k=1}^m (-1)^k ((\alpha-1)^{2k-1} - (\alpha+1)^{2k-1}) \frac{\pi^{2k}}{(2k-1)!} \lambda(2m-2k+2) = 0. \end{aligned}$$

Theorem 1.9. Let m be a nonnegative integer, and α a complex number. Then we have

$$\begin{aligned} & (-1)^m \frac{\pi^{2m+2}}{4(2m+1)!} ((\alpha-1)^{2m+1} - (\alpha+1)^{2m+1}) \\ & + \sum_{k=0}^m (-1)^k ((\alpha-1)^{2k} + (\alpha+1)^{2k} + 2\alpha^{2k}) \frac{\pi^{2k}}{(2k)!} \lambda(2m-2k+2) = 0, \end{aligned}$$

where define $0^0 = 1$.

Theorem 1.10. Let m be a nonnegative integer. Then we have

$$\begin{aligned} & (-1)^{m+1} 2(1 - 3^{-2m-1}) \lambda(2m+2) \\ & = \frac{\pi^{2m+2}}{3^{2m+1}(2m+1)!} + 4 \sum_{k=0}^m (-1)^{k+1} \frac{\pi^{2m-2k}}{3^{2m-2k}(2m-2k)!} \lambda(2k+2). \end{aligned}$$

2. AN ALTERNATIVE PROOF OF THEOREM 1.1

For our purpose, we require the following lemmas.

Lemma 2.1. Let $i = \sqrt{-1}$. Then

$$(1+ix)^a - (1-ix)^a = 2i(1+x^2)^{\frac{a}{2}} \sin(a \tan^{-1} x).$$

Proof. We have the following identity with $y = a \tan^{-1} x$:

$$\sin y = \frac{1}{2i} \left[\left(\frac{1+ix}{1-ix} \right)^{\frac{a}{2}} - \left(\frac{1+ix}{1-ix} \right)^{-\frac{a}{2}} \right].$$

From this, we obtain

$$\sin(a \tan^{-1} x) = \frac{1}{2i} \left[\frac{(1+ix)^a}{(1+x^2)^{\frac{a}{2}}} - \frac{(1-ix)^a}{(1+x^2)^{\frac{a}{2}}} \right],$$

which is the desired result. \square

Remark 2.2. Here, we note that a similar expression to Lemma 2.1 was given by Adamchik (see, e.g., [2, p. 4, (9)]) without proof.

Lemma 2.3. Let m be a positive integer and $i = \sqrt{-1}$. Then we have

$$\sum_{k=1}^m (-1)^k \binom{2m}{2k-1} \left(\frac{t}{\pi}\right)^{2k-1} = \frac{i}{2\pi^{2m}} [(\pi+it)^{2m} - (\pi-it)^{2m}].$$

Proof. Applying the binomial expansion, we have

$$\begin{aligned} i [(\pi+it)^{2m} - (\pi-it)^{2m}] &= i\pi^{2m} \sum_{k=0}^{2m} (1 - (-1)^k) \binom{2m}{k} \left(\frac{it}{\pi}\right)^k \\ &= i\pi^{2m} \sum_{k=1}^m 2 \binom{2m}{2k-1} \left(\frac{it}{\pi}\right)^{2k-1} \\ &= 2\pi^{2m} \sum_{k=1}^m (-1)^k \binom{2m}{2k-1} \left(\frac{t}{\pi}\right)^{2k-1}, \end{aligned}$$

which completes the proof. \square

Lemma 2.4. Let m be a positive integer. Then we have

$$\sum_{k=1}^m (-1)^k \frac{\pi^{2m-2k}}{(2m-2k+1)!} \frac{t^{2k-1}}{(2k-1)!} = -\frac{(\pi^2+t^2)^m}{\pi(2m)!} \sin\left(2m \tan^{-1}\left(\frac{t}{\pi}\right)\right).$$

Proof. Applying the binomial expansion, we have

$$\begin{aligned} &\sum_{k=1}^m (-1)^k \frac{\pi^{2m-2k}}{(2m-2k+1)!} \frac{t^{2k-1}}{(2k-1)!} \\ &= \sum_{k=1}^m (-1)^k \frac{(2m)! \pi^{-2k+1} t^{2k-1}}{(2m-2k+1)! (2k-1)!} \frac{\pi^{2m-1}}{(2m)!} \\ &= \frac{\pi^{2m-1}}{(2m)!} \sum_{k=1}^m (-1)^k \binom{2m}{2k-1} \left(\frac{t}{\pi}\right)^{2k-1} \\ &= \frac{i}{2\pi(2m)!} [(\pi+it)^{2m} - (\pi-it)^{2m}] \\ &\quad \text{(by Lemma 2.3)} \\ &= -\frac{1}{\pi(2m)!} (\pi^2+t^2)^m \sin\left(2m \tan^{-1}\left(\frac{t}{\pi}\right)\right) \\ &\quad \text{(by Lemma 2.1 with } a = 2m \text{ and } x = t/\pi\text{).} \end{aligned}$$

This completes the proof. \square

Lemma 2.5. *Let m be a positive integer. Then we have*

$$\int_0^\infty (\pi^2 + t^2)^m \sin \left(2m \tan^{-1} \left(\frac{t}{\pi} \right) \right) \frac{e^t dt}{e^{2t} - 1} = \frac{\pi^{2m+1}}{4}.$$

Proof. Recall that the function $J(s, a)$ is defined as follows:

$$(2.1) \quad J(s, a) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+a)^s},$$

where $0 < a \leq 1$ (see [28, (1.1)]). Using the classical Hurwitz zeta functions

$$(2.2) \quad \zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s},$$

we also have the integral representation of $J(s, a)$

$$(2.3) \quad \Gamma(s)J(s, a) = \int_0^\infty \frac{e^{(1-a)t} x^{s-1}}{e^t + 1} dt, \quad \operatorname{Re}(s) > 0,$$

(see [28, (3.1)]) and an analogue of Hermite's formula for $J(s, a)$,

$$(2.4) \quad J(s, a) = \frac{a^{-s}}{2} + 2 \int_0^\infty (a^2 + y^2)^{-s/2} \sin \left(s \tan^{-1} \frac{y}{a} \right) \frac{e^{\pi y} dy}{e^{2\pi y} - 1}.$$

The above formula enables $J(s, a)$ to be analytically continued to the whole complex plane. The special values of $J(s, a)$ at non-positive integers can be represented by Euler polynomials as follows:

$$(2.5) \quad J(-m, a) = \frac{(-1)^m}{2} E_m(1-a) = \frac{1}{2} E_m(a), \quad m \geq 0,$$

(see [28, (3.8)]). Letting $t = \pi y$, $a = 1$, and $s = -2m$ in (2.4), we have

$$(2.6) \quad \begin{aligned} J(-2m, 1) - \frac{1}{2} &= -\frac{2}{\pi} \int_0^\infty \left(1 + \left(\frac{t}{\pi} \right)^2 \right)^m \sin \left(2m \tan^{-1} \left(\frac{t}{\pi} \right) \right) \frac{e^t dt}{e^{2t} - 1} \\ &= -\frac{2}{\pi^{2m+1}} \int_0^\infty (\pi^2 + t^2)^m \sin \left(2m \tan^{-1} \left(\frac{t}{\pi} \right) \right) \frac{e^t dt}{e^{2t} - 1}. \end{aligned}$$

Replacing $-m$ by $-2m$ and setting $a = 1$ in (2.5), we have

$$(2.7) \quad J(-2m, 1) = \frac{(-1)^{-2m}}{2} E_{2m}(0) = \frac{1}{2} E_{2m}(1) = 0, \quad m \geq 1,$$

because $E_{2m}(0) = 0$ for $m > 1$. Combining (2.6) and (2.7), we obtain the desired result. \square

Proof of Theorem 1.1. The analytic continuation of $\lambda(s)$ can be established using Riemann's method. According to Euler, we have the following integral representation of the Gamma function:

$$(2.8) \quad \frac{\Gamma(s)}{(2n+1)^s} = \int_0^\infty e^{-(2n+1)t} t^{s-1} dt, \quad \operatorname{Re}(s) > 1.$$

Summing both sides over n , we have

$$\begin{aligned}
 \Gamma(s) \sum_{n=0}^{\infty} \frac{1}{(2n+1)^s} &= \sum_{n=0}^{\infty} \int_0^{\infty} e^{-(2n+1)t} t^{s-1} dt \\
 (2.9) \qquad \qquad \qquad &= \int_0^{\infty} \sum_{n=0}^{\infty} e^{-(2n+1)t} t^{s-1} dt \\
 &= \int_0^{\infty} \frac{e^{-t} t^{s-1}}{1 - e^{-2t}} dt.
 \end{aligned}$$

That is, for $\operatorname{Re}(s) > 1$,

$$(2.10) \qquad \lambda(s) = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{e^t t^{s-1}}{e^{2t} - 1} dt, \quad \operatorname{Re}(s) > 1.$$

Then, applying the integral representation (2.10), we have

$$\begin{aligned}
 &\sum_{k=1}^{m-1} \frac{(-1)^k \pi^{2m-2k}}{(2m-2k+1)!} \lambda(2k) \\
 &= \sum_{k=1}^{m-1} \frac{(-1)^k \pi^{2m-2k}}{(2m-2k+1)!} \frac{1}{\Gamma(2k)} \int_0^{\infty} \frac{e^t t^{2k-1}}{e^{2t} - 1} dt \\
 (2.11) \qquad &= \int_0^{\infty} \left[\sum_{k=1}^{m-1} \frac{(-1)^k \pi^{2m-2k}}{(2m-2k+1)!} \frac{t^{2k-1}}{(2k-1)!} \right] \frac{e^t}{e^{2t} - 1} dt \\
 &= \int_0^{\infty} \left[\sum_{k=1}^m \frac{(-1)^k \pi^{2m-2k}}{(2m-2k+1)!} \frac{t^{2k-1}}{(2k-1)!} \right] \frac{e^t}{e^{2t} - 1} dt \\
 &\quad + (-1)^{m+1} \frac{1}{(2m-1)!} \int_0^{\infty} \frac{e^t t^{2m-1}}{e^{2t} - 1} dt,
 \end{aligned}$$

and applying Lemma 2.4 and (2.10) to the last line of (2.11) with $s = 2m$, we derive the following identity:

$$\begin{aligned}
 &\sum_{k=1}^{m-1} \frac{(-1)^k \pi^{2m-2k}}{(2m-2k+1)!} \lambda(2k) \\
 (2.12) \qquad &= -\frac{1}{\pi(2m)!} \int_0^{\infty} \left[(\pi^2 + t^2)^m \sin \left(2m \tan^{-1} \left(\frac{t}{\pi} \right) \right) \right] \frac{e^t}{e^{2t} - 1} dt \\
 &\quad + (-1)^{m+1} \lambda(2m).
 \end{aligned}$$

Finally, applying Lemma 2.5 to the right-hand side of the above equality, we obtain

$$\sum_{k=1}^{m-1} \frac{(-1)^k \pi^{2m-2k}}{(2m-2k+1)!} \lambda(2k) = -\frac{\pi^{2m}}{4(2m)!} + (-1)^{m+1} \lambda(2m),$$

and the desired result follows from a direct manipulation. \square

3. PROOFS OF THEOREMS 1.2 AND 1.3 AND COROLLARY 1.7

First, we prove the following lemma:

Lemma 3.1. *Let n be a nonnegative integer, and x and α complex numbers. Then*

$$\sum_{k=0}^n (\alpha^{n-k} - (-1)^k (2x - 1 + \alpha)^{n-k}) \binom{n}{k} E_k(x) = 0.$$

Remark 3.2. *Merca proved a similar result for the Bernoulli polynomials (see [17, Theorem 2.1]).*

Remark 3.3. *Letting $\alpha = 0$ in Lemma 3.1 and recalling that $0^0 = 1$, we have*

$$(1 - (-1)^n) E_n(x) - \sum_{k=0}^{n-1} (-1)^k (2x - 1)^{n-k} \binom{n}{k} E_k(x) = 0$$

for $n \geq 1$, which can be written as

$$E_n(x) - \sum_{k=0}^n (-1)^k (2x - 1)^{n-k} \binom{n}{k} E_k(x) = 0.$$

Then, by (1.10) we have the following recurrence relation for $\lambda(s)$ at positive even integers:

$$\lambda(2m+2) + (-1)^{m+1} \pi^{2m+2} \left(\frac{1}{4(2m+1)!} + \sum_{k=0}^m (-1)^{k+1} \frac{\lambda(2k+2)}{\pi^{2k+2} (2m-2k)!} \right) = 0.$$

Proof of Lemma 3.1. We consider (1.11) and the series $\sum_{n=0}^{\infty} \alpha^n \frac{t^n}{n!} = e^{\alpha t}$. For $|t| < \pi$, it holds that

$$(3.1) \quad \frac{2e^{tx}}{e^t + 1} e^{\alpha t} = \left(\sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} \alpha^n \frac{t^n}{n!} \right),$$

which can be written as

$$(3.2) \quad \begin{aligned} \frac{2e^{tx}}{e^t + 1} e^{\alpha t} &= \frac{2e^{x(-t)}}{1 + e^{-t}} e^{(2x+\alpha-1)t} \\ &= \left(\sum_{n=0}^{\infty} (-1)^n E_n(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} (2x + \alpha - 1)^n \frac{t^n}{n!} \right). \end{aligned}$$

By applying Cauchy's rule for the multiplication of two power series to the right-hand sides of (3.1) and (3.2) and then comparing the coefficients of t^n , we obtain the relation

$$\sum_{k=0}^n \binom{n}{k} E_k(x) \alpha^{n-k} = \sum_{k=0}^n (-1)^k \binom{n}{k} E_k(x) (2x + \alpha - 1)^{n-k}$$

for $n \geq 0$, and the desired result follows from a direct manipulation. \square

Proof of Theorem 1.2. Setting $x = 0$ and $n = 2m$ in Lemma 3.1, we have

$$\sum_{k=0}^{2m} (\alpha^{2m-k} - (-1)^k (\alpha - 1)^{2m-k}) \binom{2m}{k} E_k(0) = 0$$

for $m \geq 0$, which can be written as

$$\sum_{k=1}^m \frac{\alpha^{2k-1} + (\alpha - 1)^{2k-1}}{(2k-1)!(2m-2k+1)!} E_{2m-2k+1}(0) = -\frac{\alpha^{2m} - (\alpha - 1)^{2m}}{(2m)!}$$

for $m > 0$, because $E_0(0) = 1$ and $E_{2m}(0) = 0$ for $m > 1$. Then, using Euler's formula (1.10) for the Dirichlet lambda functions, we obtain the relation

$$\begin{aligned} \sum_{k=1}^m \frac{\alpha^{2k-1} + (\alpha - 1)^{2k-1}}{(2k-1)!} (-1)^{m-k+1} \frac{4}{\pi^{2m-2k+2}} \lambda(2m-2k+2) \\ = -\frac{\alpha^{2m} - (\alpha - 1)^{2m}}{(2m)!}, \end{aligned}$$

where $\alpha \neq \frac{1}{2}$, and a direct manipulation implies the desired result. \square

Proof of Theorem 1.3. Letting $n = 2m + 1$ and $x = 0$ in Lemma 3.1, we have

$$\sum_{k=0}^{2m+1} (\alpha^{2m+1-k} - (-1)^k (\alpha - 1)^{2m+1-k}) \binom{2m+1}{k} E_k(0) = 0$$

for $m \geq 0$, which can be written as

$$\sum_{k=0}^m \frac{\alpha^{2k} + (\alpha - 1)^{2k}}{(2k)!(2m-2k+1)!} E_{2m-2k+1}(0) = -\frac{\alpha^{2m+1} - (\alpha - 1)^{2m+1}}{(2m+1)!}$$

for $m \geq 0$, because $E_0(0) = 1$ and $E_{2m}(0) = 0$ for $m > 1$. Then, using Euler's formula (1.10) for the Dirichlet lambda functions, we obtain the relation

$$\begin{aligned} \sum_{k=0}^m \frac{\alpha^{2k} + (\alpha - 1)^{2k}}{(2k)!} (-1)^{m-k+1} \frac{4}{\pi^{2m-2k+2}} \lambda(2m-2k+2) \\ = -\frac{\alpha^{2m+1} - (\alpha - 1)^{2m+1}}{(2m+1)!}, \end{aligned}$$

where $m \geq 0$ (define $0^0 = 1$), and the desired result follows from a direct manipulation. \square

Proof of Corollary 1.7. For $n > 0$, it is easy to observe that

$$(3.3) \quad \sum_{j=1}^n (-1)^j (j^{2m} - (j-1)^{2m}) = 2 \sum_{j=1}^{n-1} (-1)^j j^{2m} + (-1)^n n^{2m}$$

and

$$(3.4) \quad \sum_{j=1}^n (-1)^j (j^{2k-1} + (j-1)^{2k-1}) = (-1)^n n^{2k-1}.$$

Letting $\alpha = j$ ($j \geq 1$) in Theorem 1.2, we have

$$(3.5) \quad \sum_{j=1}^n (-1)^j \left((-1)^{m+1} \frac{j^{2m} - (j-1)^{2m}}{4} \cdot \frac{\pi^{2m+2}}{(2m)!} \right. \\ \left. + \sum_{k=1}^m (-1)^k (j^{2k-1} + (j-1)^{2k-1}) \frac{\pi^{2k}}{(2k-1)!} \lambda(2m - 2k + 2) \right) = 0.$$

Then, substituting (3.3) and (3.4) into (3.5), we obtain

$$(3.6) \quad \frac{(-1)^{m+1}}{4} \left(2 \sum_{j=1}^{n-1} (-1)^j j^{2m} + (-1)^n n^{2m} \right) \frac{\pi^{2m+2}}{(2m)!} \\ + (-1)^n \sum_{k=1}^m (-1)^k n^{2k-1} \frac{\pi^{2k}}{(2k-1)!} \lambda(2m - 2k + 2) = 0,$$

which implies the first part. The second part follows from the same reasoning. \square

4. PROOFS OF THEOREMS 1.8, 1.9, AND 1.10

In this section, we employ the generating functions of the Euler polynomials (1.11) to derive another three families of infinite recurrence relations for the Dirichlet lambda functions at positive even integer arguments.

Lemma 4.1. *Let n be a nonnegative integer, and x and α complex numbers. Then we have*

$$\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{(2x + \alpha - 1)^k + (2x - \alpha - 1)^k}{2} E_{n-k}(x) = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n}{2k} E_{n-2k}(x) \alpha^{2k},$$

where $\lfloor \cdot \rfloor$ is the floor function.

Remark 4.2. *Merca proved a similar result for the Bernoulli polynomials (see [17, Theorem 3.1]).*

Proof of Lemma 4.1. Using the generating function of Euler polynomials $E_n(x)$ (1.11) and the power series expansion of e^t , we obtain

$$\begin{aligned}
(4.1) \quad & \left(\sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} \alpha^{2n} \frac{t^{2n}}{(2n)!} \right) \\
&= \frac{1}{2} \left(\sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} \alpha^n \frac{t^n}{n!} + \sum_{n=0}^{\infty} (-1)^n \alpha^n \frac{t^n}{n!} \right) \\
&= \frac{1}{2} \frac{2e^{xt}}{e^t + 1} (e^{\alpha t} + e^{-\alpha t}) \\
&= \frac{1}{2} \frac{2e^{x(-t)}}{e^{-t} + 1} (e^{(2x+\alpha-1)t} + e^{(2x-\alpha-1)t}) \\
&= \frac{1}{2} \left(\sum_{n=0}^{\infty} (-1)^n E_n(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} ((2x+\alpha-1)^n + (2x-\alpha-1)^n) \frac{t^n}{n!} \right) \\
&= \frac{1}{2} \sum_{n=0}^{\infty} \left(\sum_{k=0}^n (-1)^{n-k} \frac{(2x+\alpha-1)^k + (2x-\alpha-1)^k}{(n-k)!k!} E_{n-k}(x) \right) t^n,
\end{aligned}$$

where the final equality is obtained by Cauchy's rule for multiplying power series. By again applying Cauchy's rule, we derive the following identity:

$$(4.2) \quad \left(\sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} \alpha^{2n} \frac{t^{2n}}{(2n)!} \right) = \sum_{n=0}^{\infty} \left(\sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{\alpha^{2k}}{(n-2k)!(2k)!} E_{n-2k}(x) \right) t^n,$$

where $\lfloor \cdot \rfloor$ is the floor function. Therefore, comparing the coefficients of t^n in (4.1) and (4.2), we obtain the desired result. \square

Proof of Theorem 1.8. Setting $x = 0$ and $n = 2m$ in Lemma 4.1, we have

$$\alpha^{2m} = \frac{1}{2} \sum_{k=0}^{2m} (-1)^{2m-k} \binom{2m}{k} ((\alpha-1)^k + (-\alpha-1)^k) E_{2m-k}(0)$$

for $m \geq 0$. Because $E_{2m}(0) = 0$ ($m > 1$) and $E_0(0) = 1$, after moving the $2m$ -th term to the left-hand side we obtain

$$\begin{aligned}
& \alpha^{2m} - \frac{1}{2} ((\alpha-1)^{2m} + (\alpha+1)^{2m}) \\
&= -\frac{1}{2} \sum_{k=1}^m \binom{2m}{2k-1} ((\alpha-1)^{2k-1} - (\alpha+1)^{2k-1}) E_{2m-(2k-1)}(0)
\end{aligned}$$

for $m \geq 1$. Finally, by applying Euler's formula (1.10) for the Dirichlet lambda functions to the above equality, we obtain the desired result. \square

Proof of Theorem 1.9. Setting $x = 0$ and $n = 2m + 1$ in Lemma 4.1, we have

$$\begin{aligned} & \sum_{k=0}^{\lfloor \frac{2m+1}{2} \rfloor} \binom{2m+1}{2k} E_{2m-2k+1}(0) \alpha^{2k} \\ &= \frac{1}{2} \sum_{k=0}^{2m+1} (-1)^{2m-k+1} \binom{2m+1}{k} ((\alpha-1)^k + (-\alpha-1)^k) E_{2m-k+1}(0) \end{aligned}$$

for $m \geq 0$. Because $E_{2m}(0) = 0$ ($m > 1$) and $E_0(0) = 1$, after moving the $(2m+1)$ -th term to the left-hand side we obtain

$$\begin{aligned} & \sum_{k=0}^m \binom{2m+1}{2k} E_{2m-2k+1}(0) \alpha^{2k} - \frac{1}{2} ((\alpha-1)^{2m+1} - (\alpha+1)^{2m+1}) \\ &= \frac{1}{2} \sum_{k=0}^m (-1)^{2m-2k+1} \binom{2m+1}{2k} ((\alpha-1)^{2k} + (-\alpha-1)^{2k}) E_{2m-2k+1}(0), \end{aligned}$$

or equivalently

$$\begin{aligned} & \sum_{k=0}^m \binom{2m+1}{2k} \left(\frac{1}{2} ((\alpha-1)^{2k} + (\alpha+1)^{2k}) + \alpha^{2k} \right) E_{2m-2k+1}(0) \\ &= \frac{1}{2} ((\alpha-1)^{2m+1} - (\alpha+1)^{2m+1}) \end{aligned}$$

for $m \geq 0$. Finally, by applying Euler's formula (1.10) to the above equality, we obtain the relation

$$\begin{aligned} & \sum_{k=0}^m (-1)^{k+1} \frac{(2m+1)!}{(2k)!} \left(\frac{1}{2} ((\alpha-1)^{2k} + (\alpha+1)^{2k}) + \alpha^{2k} \right) \pi^{2k} \lambda(2m-2k+2) \\ &= \frac{1}{8} (-1)^m \pi^{2m+2} ((\alpha-1)^{2m+1} - (\alpha+1)^{2m+1}) \end{aligned}$$

for $m \geq 0$, which is the desired result. \square

Proof of Theorem 1.10. From the expression for the Euler polynomials

$$E_n(x) = \sum_{k=0}^n \binom{n}{k} x^{n-k} E_k(0),$$

we have

$$\begin{aligned} (4.3) \quad E_{2m+1} \left(\frac{1}{3} \right) &= \sum_{k=0}^{2m+1} \binom{2m+1}{k} E_k(0) \left(\frac{1}{3} \right)^{2m+1-k} \\ &= \left(\frac{1}{3} \right)^{2m+1} + \sum_{k=0}^m \binom{2m+1}{2k+1} E_{2k+1}(0) \left(\frac{1}{3} \right)^{2m-2k} \end{aligned}$$

because $E_{2m}(0) = 0$ ($m > 1$) and $E_0(0) = 1$. It is known from [24, Theorem 3.3] that

$$(4.4) \quad E_n(x) = m^n \sum_{k=0}^{m-1} (-1)^k E_n \left(\frac{x+k}{m} \right) \quad \text{if } 2 \nmid m.$$

Setting $x = 0$ and $m = 3$ in (4.4), and noticing that $E_{2m+1} \left(\frac{2}{3} \right) = -E_{2m+1} \left(\frac{1}{3} \right)$, we have

$$(4.5) \quad E_{2m+1} \left(\frac{1}{3} \right) = \frac{1}{2} (1 - 3^{-2m-1}) E_{2m+1}(0).$$

Comparing (4.3) and (4.5), we find that

$$(4.6) \quad \frac{1}{2} (1 - 3^{-2m-1}) E_{2m+1}(0) = \left(\frac{1}{3} \right)^{2m+1} + \sum_{k=0}^m \binom{2m+1}{2k+1} E_{2k+1}(0) \left(\frac{1}{3} \right)^{2m-2k}.$$

Then, by applying Euler's formula (1.10) for the Dirichlet lambda functions to the above equality, we obtain the desired result. \square

5. CONVOLUTION IDENTITIES

In this section, we prove convolution identities for special values of $\lambda(s)$ at even arguments and for special values of $\beta(s)$ at odd arguments.

Euler derived the following beautiful convolution identity for the Bernoulli numbers:

$$(5.1) \quad \sum_{k=1}^{m-1} \binom{2m}{2k} B_{2k} B_{2m-2k} = -(2m+1) B_{2m}, \quad m \geq 2,$$

which has been generalized by many authors in different directions (see, for example, [5], [6], and [12]). By Euler's formula for the special values of the Riemann zeta function at even arguments,

$$\zeta(2m) = (-1)^{m-1} \frac{(2\pi)^{2m} B_{2m}}{2(2m)!},$$

(5.1) can be written as

$$\sum_{k=1}^{m-1} \zeta(2k) \zeta(2m-2k) = \left(m + \frac{1}{2} \right) \zeta(2m), \quad m \geq 2.$$

Similarly, the following convolution identity for the special values of $\lambda(s)$ at even arguments has been demonstrated in [6, 22] for $m \geq 1$:

$$(5.2) \quad \sum_{k=1}^m \lambda(2k) \lambda(2m-2k+2) = \left(m + \frac{1}{2} \right) \lambda(2m+2).$$

In addition, the following convolution identity for the special values of $\beta(s)$ at odd arguments was proved by Williams (see [27, p. 22, Theorem II]). Here, we present an alternative proof.

Theorem 5.1 (G.T. Williams). *Let m be a nonnegative integer. Then we have*

$$\sum_{k=0}^m \beta(2k+1)\beta(2m-2k+1) = \left(m + \frac{1}{2}\right) \lambda(2m+2).$$

Proof. Applying the generating function of the Euler polynomials, we have

$$\begin{aligned} \frac{2^2 e^t}{(e^t + 1)^2} &= \left(\sum_{n=0}^{\infty} E_n \left(\frac{1}{2}\right) \frac{t^n}{n!} \right) \left(\sum_{m=0}^{\infty} E_m \left(\frac{1}{2}\right) \frac{t^m}{m!} \right) \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^n E_k \left(\frac{1}{2}\right) E_{n-k} \left(\frac{1}{2}\right) \frac{t^n}{k!(n-k)!} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^n \binom{n}{k} E_k \left(\frac{1}{2}\right) E_{n-k} \left(\frac{1}{2}\right) \frac{t^n}{n!}. \end{aligned}$$

We also have

$$\begin{aligned} \frac{1}{2} \left(\frac{2e^{\frac{1}{2}t}}{e^t + 1} \right)^2 &= \frac{d}{dt} \left(\frac{2e^t}{e^t + 1} \right) = \frac{d}{dt} \left(\sum_{n=0}^{\infty} E_n(1) \frac{t^n}{n!} \right) \\ &= \sum_{n=0}^{\infty} E_{n+1}(1) \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients of t^n in the above two equalities, we have

$$(5.3) \quad \sum_{k=0}^n \frac{E_k \left(\frac{1}{2}\right) E_{n-k} \left(\frac{1}{2}\right)}{k!(n-k)!} = 2 \frac{E_{n+1}(1)}{n!}.$$

Then, setting $n = 2m$ in (5.3) and noticing that $E_{2m+1}(1) = (-1)^{2m+1} E_{2m+1}(0)$ and $E_k = 2^k E_k(1/2)$, we obtain

$$(5.4) \quad \sum_{k=0}^{2m} \frac{E_k E_{2m-k}}{2^{2m+1} k!(2m-k)!} = -\frac{1}{(2m)!} E_{2m+1}(0).$$

That is,

$$(5.5) \quad \sum_{k=0}^m \frac{E_{2k} E_{2m-2k}}{2^{2m+1} (2k)!(2m-2k)!} = -\frac{1}{(2m)!} E_{2m+1}(0)$$

because $E_{2k+1} = 0$ for $k \geq 0$. Applying (1.10) and (1.14) to the above equality, we obtain the desired result. \square

Theorem 5.2. *Let m be a positive integer. Then we have*

$$\sum_{k=0}^m 2^{-2k} \zeta(2k) \lambda(2m-2k+2) = 0.$$

Proof. Taking into account (1.4) and (1.11), we have

$$\begin{aligned}
\sum_{n=0}^{\infty} 2^n B_n \left(\frac{x}{2}\right) \frac{t^n}{n!} &= \frac{2te^{\frac{x}{2}(2t)}}{e^{2t} - 1} \\
&= \frac{t}{e^t - 1} \frac{2e^{xt}}{e^t + 1} \\
&= \left(\sum_{n=0}^{\infty} B_n \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \right) \\
&= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} B_k E_{n-k}(x) \right) \frac{t^n}{n!},
\end{aligned}$$

yielding

$$(5.6) \quad 2^n B_n \left(\frac{x}{2}\right) = \sum_{k=0}^n \binom{n}{k} B_k E_{n-k}(x), \quad n \geq 0.$$

If we set $n = 2m + 1$ and $x = 0$ in (5.6) and use $B_{2m+1} = E_{2m}(0) = 0$ ($m \geq 1$), then we obtain

$$(5.7) \quad \sum_{k=0}^m \binom{2m+1}{2k} B_{2k} E_{2m-2k+1}(0) = 0, \quad m \geq 1.$$

Hence, by the well-known relations (1.3) and (1.10), we obtain

$$\sum_{k=0}^m 2^{-2k} \zeta(2k) \lambda(2m - 2k + 2) = 0, \quad m \geq 1,$$

as required. □

6. POWER SERIES EXPANSION FOR $J(s, a)$

We start with

$$\frac{\Gamma(s)}{(n+a)^s} = \int_0^{\infty} e^{-(n+a)t} t^{s-1} dt, \quad \operatorname{Re}(s) > 0.$$

Applying this to (2.1), we obtain

$$\begin{aligned}
\Gamma(s) J(s, a) &= \sum_{n=0}^{\infty} \int_0^{\infty} (-1)^n e^{-(n+a)t} t^{s-1} dt \\
(6.1) \quad &= \int_0^{\infty} \sum_{n=0}^{\infty} (-1)^n e^{-(n+a)t} t^{s-1} dt \\
&= \int_0^{\infty} \frac{e^{-at} t^{s-1}}{e^{-t} + 1} dt
\end{aligned}$$

for $\operatorname{Re}(s) > 0$. That is,

$$(6.2) \quad \Gamma(s) J(s, a) = \int_0^{\infty} \frac{e^{(1-a)t} t^{s-1}}{e^t + 1} dt, \quad \operatorname{Re}(s) > 0,$$

(see (2.3)). By splitting the above integral at x and employing the generating function of the Euler polynomials

$$(6.3) \quad \frac{2e^{zt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(z) \frac{t^n}{n!}, \quad |t| < \pi,$$

for the integral over $[0, x)$ obtain

$$(6.4) \quad \begin{aligned} \Gamma(s)J(s, a) &= \int_x^{\infty} \frac{e^{(1-a)t} t^{s-1}}{e^t + 1} dt + \int_0^x \frac{e^{(1-a)t} t^{s-1}}{e^t + 1} dt \\ &= \int_x^{\infty} \frac{e^{(1-a)t} t^{s-1}}{e^t + 1} dt + \frac{1}{2} \sum_{n=0}^{\infty} \frac{E_n(1-a)}{n!} \int_0^x t^{n+s-1} dt. \end{aligned}$$

To handle the first integral in the above equality, we employ a standard integral representation for the incomplete Gamma function

$$\Gamma(s, x) = \int_x^{\infty} t^{s-1} e^{-t} dt$$

(see, e.g., [1, p. 260] and [4, (1.1)]), together with a geometric series expansion, to obtain

$$(6.5) \quad \begin{aligned} \int_x^{\infty} \frac{e^{(1-a)t} t^{s-1}}{e^t + 1} dt &= \sum_{n=0}^{\infty} (-1)^n \int_x^{\infty} e^{-(n+a)t} t^{s-1} dt \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{1}{(n+a)^s} \Gamma(s, (n+a)x). \end{aligned}$$

From (6.4) and (6.5), we obtain the following expressions of $\Gamma(s)J(s, a)$.

Proposition 6.1. *Let $|x| < \pi$ and $\operatorname{Re}(s) > 0$. Then we have*

$$\Gamma(s)J(s, a) = \sum_{n=0}^{\infty} (-1)^n \frac{\Gamma(s, (n+a)x}{(n+a)^s} + \frac{1}{2} \sum_{n=0}^{\infty} \frac{E_n(1-a)}{n!} \frac{x^{n+s}}{n+s}.$$

The above representation holds in the whole complex plane \mathbb{C} . Furthermore, it implies the special values of $J(s, a)$ at non-positive integers [28, (3.8)] for $k \geq 0$ given by

$$J(-k, a) = \frac{(-1)^k}{2} E_k(1-a) = \frac{1}{2} E_k(a).$$

Taking $a = \frac{1}{2}$ in (2.1), it follows from (1.13) that

$$(6.6) \quad J\left(s, \frac{1}{2}\right) = 2^s \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^s} = 2^s \beta(s).$$

Thus, the above proposition also implies the following result.

Corollary 6.2. *We have*

$$\Gamma(s)\beta(s) = \frac{1}{2^s} \sum_{n=0}^{\infty} (-1)^n \frac{\Gamma(s, (n+1/2)x}{(n+1/2)^s} + \frac{1}{2^{s+1}} \sum_{n=0}^{\infty} \frac{E_n}{2^n n!} \frac{x^{n+s}}{n+s}$$

with the free parameter $x \in [0, \pi)$. In particular, we obtain the special values $\beta(-k) = E_k/2$ of $\beta(s)$ at non-positive integers for $k \geq 0$.

Let

$$\eta(s) = (1 - 2^{1-s})\zeta(s)$$

be the alternating zeta function (in Euler's notation, $M(s)$; see [25, p. 70]). That is,

$$(6.7) \quad \eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s}, \quad \operatorname{Re}(s) > 0,$$

(see [1, pp. 807–808]). Then, taking $a = 1$ in (2.1), we obtain $J(s, 1) = \eta(s)$. The former is sometimes called Lerch's eta function, and the latter is sometimes named Dirichlet's eta function [26]. Proposition 6.1 may have many interesting applications. For example, letting $a = 1$, we recover the following result by Coffey.

Corollary 6.3 (Coffey, [4, p. 1384, Proposition 1]).

$$\Gamma(s)\eta(s) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{\Gamma(s, nx)}{n^s} + \frac{1}{2} \sum_{n=0}^{\infty} \frac{E_n(0)}{n!} \frac{x^{n+s}}{n+s},$$

with the free parameter $x \in [0, \pi)$. In particular, we obtain the exact evaluations $\eta(-k) = (-1)^k E_k(0)/2$ for $k \geq 0$.

Proposition 6.4. For $0 < a \leq 1$, we have

$$\Gamma(s)J(s, a) = \Gamma(s) \frac{(a+1)^{1-s} - a^{1-s}}{2(s-1)} - \sum_{n=1}^{\infty} \frac{2^n \Gamma(s+n)}{(n+1)!} J(s+n, a).$$

Proof. From the power series expansion of e^t , we have

$$(6.8) \quad \frac{1}{2t} e^{2t} - \frac{1}{2t} - 1 = \sum_{n=1}^{\infty} \frac{(2t)^n}{(n+1)!}.$$

In (6.2), by replacing s by $s+n$, multiplying both sides by $\frac{2^n}{(n+1)!}$, and summing from $n=1$ to ∞ , it then follows from (6.8) and (6.2) that

$$(6.9) \quad \begin{aligned} & \sum_{n=1}^{\infty} \frac{2^n}{(n+1)!} \Gamma(s+n) J(s+n, a) \\ &= \int_0^{\infty} \frac{e^{(1-a)t} t^{s-1}}{e^t + 1} \sum_{n=1}^{\infty} \frac{(2t)^n}{(n+1)!} dt \\ &= \int_0^{\infty} \frac{e^{(1-a)t} t^{s-1}}{e^t + 1} \left(\frac{1}{2t} e^{2t} - \frac{1}{2t} - 1 \right) dt \\ &= \frac{\Gamma(s-1)}{2} (J(s-1, a+2) - J(s-1, a)) - \Gamma(s) J(s, a) \\ &= \frac{\Gamma(s-1)}{2} ((a+1)^{1-s} - a^{1-s}) - \Gamma(s) J(s, a). \end{aligned}$$

The final equality holds because $J(s, a+1) + J(s, a) = a^{-s}$. Therefore, the proposition is proven. \square

ACKNOWLEDGEMENT

The authors would like to thank the referee for valuable comments and suggestions.

REFERENCES

- [1] M. Abramowitz and I. Stegun (eds.), *Handbook of mathematical functions with formulas, graphs and mathematical tables*, Dover, New York, 1972.
- [2] V.S. Adamchik, *On the Hurwitz function for rational arguments*, Appl. Math. Comput. **187** (2007) 3–12.
- [3] L.E. Ballentine and S.M. McRae, *Moment equations for probability distributions in classical and quantum mechanics*, Phys. Rev. A **58** (1998) 1799–1809.
- [4] M.W. Coffey, *Series representation of the Riemann zeta function and other results: complements to a paper of Crandall*, Math. Comp. **83** (2014) 1383–1395.
- [5] M.W. Coffey, *Bernoulli identities, zeta relations, determinant expressions, Mellin transforms, and representation of the Hurwitz numbers*, J. Number Theory **184** (2018) 27–67.
- [6] K. Dilcher, *Sums of products of Bernoulli numbers*, J. Number Theory **60** (1996) 23–41.
- [7] L. Euler, *Remarques sur un beau rapport entre les séries des puissances tant directes que réciproques*, 1768, E352 (Eneström Index), The Euler Archive.
- [8] S.R. Finch, *Mathematical Constants*, Cambridge Univ. Press, Cambridge, 2003.
- [9] I.S. Gradshteyn and I.M. Ryzhik, *Table of integrals, series, and products*, Academic Press, New York, 1994.
- [10] S. Hu and M.-S. Kim, *The $(S, \{2\})$ -Iwasawa theory*, J. Number Theory **158** (2016) 73–89.
- [11] S. Hu, D. Kim and M.-S. Kim, *Special values and integral representations for the Hurwitz-type Euler zeta functions*, J. Korean Math. Soc. **55** (2018) 185–210.
- [12] M.-S. Kim and S. Hu, *Sums of products of Apostol-Bernoulli numbers*, Ramanujan J. **28** (2012) 113–123.
- [13] M.C. Lettington, *Fleck’s congruence, associated magic squares and a zeta identity*, Funct. Approx. Comment. Math. **45** (2011) 165–205.
- [14] M.C. Lettington, *A trio of Bernoulli relations, their implications for the Ramanujan polynomials and the special values of the Riemann zeta function*, Acta Arith. **158** (2013) 1–31.
- [15] M. Merca, *On the Song recurrence relation for the Riemann zeta function*, Miskolc Math. Notes **17** (2016) 941–945.
- [16] M. Merca, *Asymptotics of the Chebyshev-Stirling numbers of the first kind*, Integral Transforms Spec. Funct. **27** (2016) 259–267.
- [17] M. Merca, *On families of linear recurrence relations for the special values of the Riemann zeta function*, J. Number Theory **170** (2017) 55–65.
- [18] N.E. Nörlund, *Vorlesungen ber Differenzenrechnung*, Springer, Berlin, 1924.
- [19] Á. Pintér and C. Rakaczki, *On the decomposability of linear combinations of Euler polynomials*, Miskolc Math. Notes **18** (2017) 407–415.
- [20] J.L. Raabe, *Zurückführung einiger Summen und bestmmtiem Integrale auf die Jacob-Bernoullische Function*, J. Reine Angew. Math. **42** (1851) 348–367.
- [21] T. Rivoal and W. Zudilin, *Diophantine properties of numbers related to Catalan’s constant*, Math. Ann. **326** (2003) 705–721.
- [22] R. Sitaramachandrarao and B. Davis, *Some identities involving the Riemann zeta function, II*, Indian J. Pure Appl. Math. **17** (1986) 1175–1186.
- [23] I. Song, *A recursive formula for even order harmonic series*, J. Comput. Appl. Math. **21** (1988) 251–256.
- [24] Z.-W. Sun, *Introduction to Bernoulli and Euler polynomials*, a lecture given in Taiwan on June 6, 2002, <http://maths.nju.edu.cn/~zwsun/BerE.pdf>.

- [25] V.S. Varadarajan, *Euler through time: a new look at old themes*, American Mathematical Society, Providence, RI, 2006.
- [26] https://en.m.wikipedia.org/wiki/Dirichlet_eta_function.
- [27] G.T. Williams, *A new method of evaluating $\zeta(2n)$* , Amer. Math. Monthly **60** (1953) 19–25.
- [28] K.S. Williams and N.Y. Zhang, *Special values of the Lerch zeta function and the evaluation of certain integrals*, Proc. Amer. Math. Soc. **119** (1993) 35–49.

DEPARTMENT OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY,
GUANGZHOU, GUANGDONG 510640, CHINA
E-mail address: mahusu@scut.edu.cn

DIVISION OF MATHEMATICS, SCIENCE, AND COMPUTERS, KYUNGNAM UNIVERSITY,
7(WORYEONG-DONG) KYUNGNAMDAEHAK-RO, MASANHAPPO-GU, CHANGWON-
SI, GYEONGSANGNAM-DO 51767, REPUBLIC OF KOREA
E-mail address: mskim@kyungnam.ac.kr