

STICK NUMBER OF SPATIAL GRAPHS

MINJUNG LEE, SUNGJONG NO, AND SEUNGSANG OH

ABSTRACT. For a nontrivial knot K , Negami found an upper bound on the stick number $s(K)$ in terms of its crossing number $c(K)$ which is $s(K) \leq 2c(K)$. Later, Huh and Oh utilized the arc index $\alpha(K)$ to present a more precise upper bound $s(K) \leq \frac{3}{2}c(K) + \frac{3}{2}$. Furthermore, Kim, No and Oh found an upper bound on the equilateral stick number $s_=(K)$ as follows; $s_=(K) \leq 2c(K) + 2$. As a sequel to this research program, we similarly define the stick number $s(G)$ and the equilateral stick number $s_=(G)$ of a spatial graph G , and present their upper bounds as follows;

$$\begin{aligned} s(G) &\leq \frac{3}{2}c(G) + 2e + \frac{3b}{2} - \frac{v}{2}, \\ s_=(G) &\leq 2c(G) + 2e + 2b - k, \end{aligned}$$

where e and v are the number of edges and vertices of G , respectively, b is the number of bouquet cut-components, and k is the number of non-splittable components.

1. INTRODUCTION

Throughout this paper we work in the piecewise linear category. A graph is a finite set of vertices connected by edges allowing loops and multiple edges. A *spatial graph* is a graph embedded in \mathbb{R}^3 . We consider two spatial graphs to be the same if they are equivalent under ambient isotopy. A *bouquet* is a spatial graph consisting of only one vertex and loops. Note that a knot is a spatial graph consisting of a vertex and a loop.

A *stick spatial graph* is a spatial graph which consists of finite line segments, called *sticks*, as drawn in Figure 1. This presentation of spatial graphs can be considered to be a reasonable mathematical model of polymers because such physical objects have rigidity. Concerning stick spatial graphs, one natural problem may be to determine the number of sticks. The *stick number* $s(G)$ of a spatial graph G is defined to be the minimal number of sticks required to realize the spatial graph as a stick spatial graph.


The stick number of a nontrivial knot K is related to its crossing number $c(K)$ by the following inequalities [3, 6, 11];

$$\frac{7 + \sqrt{8c(K) + 1}}{2} \leq s(K) \leq \frac{3}{2}c(K) + \frac{3}{2}.$$

Mathematics Subject Classification 2010: 57M25, 57M27.

The second author was supported by the BK21 Plus Project through the National Research Foundation of Korea (NRF) grant funded by the Korean Ministry of Education (22A20130011003).

The corresponding author (Seungsang Oh) was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP) (No. NRF-2017R1A2B2007216).

FIGURE 1. Spatial θ -curve 5_1 and its stick presentation

Particularly, a 2-bridge knot K with $c(K) \geq 6$ has a better upper bound $s(K) \leq c(K) + 2$ shown in [4, 5, 10]. There are few knots whose stick number can be determined exactly. Knots with small crossing numbers were investigated by Randell [12]. Jin [7] determined the precise stick number of a (p, q) -torus knot $T(p, q)$ in case of $2 \leq p \leq q \leq 2p$ as $s(T(p, q)) = 2q$. The same result was found independently by Adams *et al.* [1], but for a smaller range of parameters.

As a sequel to this research program, we consider the stick number of spatial graphs. We consider two types of 2-spheres that separate a spatial graph G into two parts. Such a 2-sphere is called a splitting-sphere if it does not meet G , and a cut-sphere if it intersects G in a single vertex which is called a cut vertex. We maximally decompose G into *cut-components* by cutting G along a maximal set \mathcal{S} of splitting-spheres and cut-spheres where any two spheres are either disjoint or intersect each other in a cut vertex. G is said to be *non-splittable* if there is no splitting-sphere separating G . Each cut-component is non-splittable. In particular, if such a cut-component is a bouquet, it is called a *bouquet cut-component*. The crossing number $c(G)$ is the minimal number of double points in any generic projection of G into the plane \mathbb{R}^2 . Here, these double points are disjoint from the projected vertices of the spatial graph.

Theorem 1. *Let G be any spatial graph with e edges, v vertices and b bouquet cut-components. Then*

$$s(G) \leq \frac{3}{2}c(G) + 2e + \frac{3b}{2} - \frac{v}{2}.$$

For a knot K , this result implies $s(K) \leq \frac{3}{2}c(K) + 3$ which is very close to the previously known upper bound. Furthermore for both the unlink with n components and the trivial θ_n -curve even for any large n , the stick number $s(G)$ is actually equal to the value $2e + \frac{3b}{2} - \frac{v}{2}$ of G (obviously $c(G) = 0$). This implies that this sum of the last three terms is best possible.

We are also interested in another quantity concerning stick spatial graphs. An *equilateral spatial graph* is a stick spatial graph which consists of uniform length sticks. The *equilateral stick number* $s_=(G)$ of a spatial graph G is defined to be the minimal number of sticks required to construct an equilateral spatial graph representation of G .

Even for knots, little is known about the equilateral stick number. Rawdon and Scharein [13] used algorithms in the software KnotPlot to compute upper bounds for the equilateral stick number of all prime knots with up to 10 crossings. They showed that all such knots except seven can be constructed with the same number of equal length sticks as their stick numbers. Recently, Kim, No and Oh [8] found an upper bound for a nontrivial knot K as follows;

$$s_=(K) \leq 2c(K) + 2.$$

Theorem 2. *Let G be any spatial graph with e edges and b bouquet cut-components. Suppose that G has k non-splittable components. Then*

$$s_=(G) \leq 2c(G) + 2e + 2b - k.$$

For a nontrivial knot K , this result implies $s_=(K) \leq 2c(K) + 3$ which is very close to the previously known upper bound. Furthermore for both the unlink with n components and the trivial θ_n -curve even for any large n , the stick number $s_=(G)$ is equal to the value $2e + 2b - k$ of G , implying that this sum of the last three terms is best possible.

2. ARC INDEX OF SPATIAL GRAPHS

In an *arc presentation* of a spatial graph G , G is constructed in an open-book with finitely many half-planes so that it meets each half-plane in exactly one simple arc with two different end-points on the binding axis. Therefore the binding axis contains all vertices of G and finitely many points from the interiors of edges of G , and each edge of G may pass from one page to another across the binding axis. See Figure 2 for an arc presentation of the θ -curve 5_1 with eight pages. The *arc index* $\alpha(G)$ is defined to be the minimal number of pages among all possible arc presentations of G .




FIGURE 2. Arc presentation of 5_1

Bae and Park [2] established an upper bound on the arc index of a non-trivial knot in terms of the crossing number. The authors [9] generalized the argument used in [2] to find an upper bound on the arc index of a spatial graph as follows. This upper bound result is crucial in the proofs of our two main theorems.

Theorem 3. (Lee-No-Oh [9]) *Let G be any spatial graph with e edges and b bouquet cut-components. Then*

$$\alpha(G) \leq c(G) + e + b.$$

3. CIRCULAR ARC PRESENTATIONS

We introduce a variant of an arc presentation of a spatial graph G , called a *circular arc presentation*, which is more useful to construct a stick presentation of G in Section 4. First, take a circular disk D in \mathbb{R}^2 whose boundary actually indicates the binding axis combined with point at infinity. Now draw straight chords instead of arcs of the arc presentation. During this procedure all the page numbers and the under/over crossings are preserved. The chords are denoted by l_1, \dots, l_n in the order of the page numbers, so that if l_i and l_j share a crossing in the interior of D and $i < j$, then l_i passes under l_j . Note that $\alpha(G) = n$. The points of the graph lying on the boundary of D are called the *binding points*. Note that each chord has two different binding points at its ends. See Figure 3.




FIGURE 3. Circular arc presentation of 5_1

For a binding point b , the *initiating page number* $p(b)$ is the smallest page number among all chords incident to b . A chord l_i is said to have an *initiating end* if this end point has the initiating page number i . Then all chords are distinguished into three types as follows; a *bi-initiating chord* if it has two initiating ends on both sides, a *uni-initiating chord* if it has an initiating end only on one side, and a *non-initiating chord* if it has no initiating ends. In Figure 3, the initiating page numbers of the seven binding points in clockwise order starting at point at infinity are 1, 2, 6, 3, 1, 4, and 2. Furthermore, l_1 and l_2 are bi-initiating chords, l_3 , l_4 and l_6 are uni-initiating chords, and the others are non-initiating chords.

4. STICK PRESENTATIONS

In this section we will prove Theorem 1.

Proof of Theorem 1. Let G be a spatial graph with e edges and v vertices, and b is the number of bouquet cut-components of G . Assume that $\alpha(G) = n$ and C is its circular arc presentation with n chords l_1, \dots, l_n . Further assume

that C has n_2 , n_1 and n_0 numbers of bi-initiating, uni-initiating and non-initiating chords where $n_2 + n_1 + n_0 = n$.

The projection $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ is defined by $\pi(x, y, z) = (x, y)$. For each $i = 1, \dots, n$, put a horizontal line segment h_i in $D \times \{i\}$ so that $\pi(h_i) = l_i$ as in Figure 4.




FIGURE 4. Simple stack of n chords

Now, deform these line segments h_i in an inductive manner on i , meanwhile their projections are preserved. The segment h_1 stays in itself and so its highest z -coordinate is $z_1 = 1$. Assume that h_1, h_2, \dots, h_{k-1} are deformed inductively with their highest integer-valued z -coordinates $z_1 < z_2 < \dots < z_{k-1}$, respectively.

Deforming h_k depends on the type of the chord l_k as in Figure 5. If l_k is a bi-initiating chord, raise up h_k into $D \times \{z_{k-1} + 1\}$ with its highest z -coordinate $z_k = z_{k-1} + 1$. If l_k is a uni-initiating chord, then exactly one end b of l_k has the initiating page number $p(b) = k$. Let b' denote the other end with its initiating page number $p(b') = j' < k$. Raise up h_k obliquely so that its one end is $h_{j'} \cap \pi^{-1}(b')$ and the other end is $b \times \{z_k\}$ for a sufficiently large integer $z_k > z_{k-1}$. More precisely, the interior of the triangle between this new h_k and a horizontal line segment H in $D \times \{z_k\}$ with $\pi(H) = l_k$ has no intersection with any other h_i , $i < k$. Lastly, if l_k is a non-initiating chord, then both ends b and b' of l_k have the initiating page numbers $p(b) = j < k$ and $p(b') = j' < k$. Bend h_k at the center $c \times \{k\}$ into two connected line segments and raise up h_k so that its ends are $h_j \cap \pi^{-1}(b)$ and $h_{j'} \cap \pi^{-1}(b')$, and the center point goes to $c \times \{z_k\}$ for a sufficiently large integer $z_k > z_{k-1}$. Eventually the interiors of the two triangles between this new h_k and a horizontal line segment H in $D \times \{z_k\}$ with $\pi(H) = l_k$ have no intersection with any other h_i , $i < k$.

Then the new stick spatial graph $G' = \cup_{i=1}^n h_i$ in $D \times [1, \infty)$ is isotopic to G and consists of $n_2 + n_1 + 2n_0 = n + n_0$ sticks as desired. Therefore,

$$s(G) \leq \alpha(G) + n_0.$$

On the other hand, split all chords in C slightly away from binding points. Then there are total $2n$ ends of the n chords. Note that the number of binding points is $n - e + v$ by an easy combinatorial calculation. Since each binding point is an initiating end of exactly one of chords incident to it, there are exactly $n - e + v$ initiating ends among these $2n$ ends. Thus $n + e - v$ ends are not initiating. Since non-initiating chords have no initiating ends,




FIGURE 5. Bi-initiating chord, uni-initiating chord and non-initiating chord cases

the number n_0 of non-initiating chords is at most $\frac{n+e-v}{2}$. This implies

$$s(G) \leq \frac{3}{2}\alpha(G) + \frac{e}{2} - \frac{v}{2}.$$

Now we complete the proof by applying Theorem 3. \square

5. EQUILATERAL STICK PRESENTATIONS

In this section we will prove Theorem 2.

Proof of Theorem 2. We will follow the main argument in the proof of Theorem 1.1 in [8]. Let G be a spatial graph with e edges and b bouquet cut-components. Suppose that G has k non-splittable components. Let G_1 be a non-splittable component of G .

Assume that $\alpha(G_1) = n$ and A is its arc presentation with n arcs. Let M be a sufficiently large number. Now we stretch A so that each arc is deformed to two connected sticks of length M each on the related page, while all binding points are fixed, as drawn in Figure 6. Already we obtain an equilateral spatial graph of G_1 consisting of $2n$ sticks of length M . Therefore, $s_=(G_1) \leq 2\alpha(G_1)$.

Now we will reduce one more stick from A . Assume that the top binding point has m sticks incident to, say d_1, \dots, d_m . Let e_1, \dots, e_m denote the other m sticks connected to d_1, \dots, d_m , respectively. As in Figure 7, delete all m sticks d_1, \dots, d_m . Rotate the stick e_1 counterclockwise along its endpoint at the binding axis on the related page until e_1 is close enough to the binding axis. Now rotate each e_i , $i = 2, \dots, m$, counterclockwise along its endpoint at the binding axis on the related page so that the distance between the other endpoints of e_1 and e_i which are not at the binding axis is M . Glue a new stick f_i of length M to these endpoints as a substitution for d_i . Clearly this formation does not change the spatial graph type, but we reduce one stick from the original. Hence the resulting equilateral spatial graph G_1 satisfies

$$s_=(G_1) \leq 2\alpha(G_1) - 1.$$



FIGURE 6. Equilateral stick presentation of 5_1 

FIGURE 7. Reducing one more stick

Furthermore, if G can be split into two components G_1 and G_2 , then obviously $s_=(G) = s_=(G_1) + s_=(G_2)$. Theorem 2 follows directly from Theorem 3.

Note that unlike the case of nontrivial knots in [8], it is possible that there is an arc connecting the top and the bottom binding points, for example, the trivial θ_n -curve. We cannot repeat the same argument at the bottom binding point. \square

REFERENCES

- [1] C. Adams, B. Brennan, D. Greilsheimer and A. Woo, *Stick numbers and composition of knots and links*, J. Knot Theory Ramifications **6** (1997) 149–161.
- [2] Y. Bae and C. Park, *An upper bound of arc index of links*, Math. Proc. Cambridge Philos. Soc. **129** (2000) 491–500.
- [3] J. Calvo, *Characterizing polygons in \mathbb{R}^3* , in Physical knots: Knotting, Linking, and Folding Geometric Objects in \mathbb{R}^3 , Contemporary Mathematics, Vol. 304 (American Mathematical Society, 2002) pp. 37–53.
- [4] E. Furstenberg, J. Li and J. Schneider, *Stick knots*, Chaos Solitons Fractals **9** (1998) 561–568.

- [5] Y. Huh, S. No and S. Oh, *Stick numbers of 2-bridge knots and links*, Proc. Amer. Math. Soc. **139** (2011) 4143–4152.
- [6] Y. Huh and S. Oh, *An upper bound on stick number of knots*, J. Knot Theory Ramifications **20** (2011) 741–747.
- [7] G. T. Jin, *Polygonal indices and superbridges indices of torus knots and links*, J. Knot Theory Ramifications **6** (1997) 281–289.
- [8] H. Kim, S. No and S. Oh, *Equilateral stick number of knots*, J. Knot Theory Ramifications **23** (2014) 1460008.
- [9] M. Lee, S. No and S. Oh, *Arc index of spatial graphs*, J. Graph Theory (in press).
- [10] C. McCabe, *An upper bound on edge numbers of 2-bridge knots and links*, J. Knot Theory Ramifications **7** (1998) 797–805.
- [11] S. Negami, *Ramsey theorems for knots, links, and spatial graphs*, Trans. Amer. Math. Soc. **324** (1991) 527–541.
- [12] R. Randell, *An elementary invariant of knots*, J. Knot Theory Ramifications **3** (1994) 279–286.
- [13] E. Rawdon and R. Scharein, *Upper bounds for equilateral stick numbers*, in Physical knots: Knotting, Linking, and Folding Geometric Objects in \mathbb{R}^3 , Contemporary Mathematics, Vol. 304 (American Mathematical Society, 2002) pp. 55–76.

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, SEOUL 02841, KOREA
E-mail address: mjmlj@korea.ac.kr

DEPARTMENT OF STATISTICS, EWHA WOMANS UNIVERSITY, SEOUL 03760, KOREA
E-mail address: sungjongno84@gmail.com

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, SEOUL 02841, KOREA
E-mail address: seungsang@korea.ac.kr