

MODELS OF CURVES OVER DVRS

TIM DOKCHITSER

ABSTRACT. Let C be a smooth projective curve over a discretely valued field K , defined by an affine equation $f(x, y) = 0$. We construct a model of C over the ring of integers of K using a toroidal embedding associated to the Newton polygon of f . We show that under ‘generic’ conditions it is regular with normal crossings, and determine when it is minimal, the global sections of its relative dualising sheaf, and the tame part of the first étale cohomology of C .

1. INTRODUCTION

The purpose of this paper is to construct regular models and study invariants of arithmetic surfaces using a toric resolution derived from the defining equation. Let K be a field with a discrete valuation v_K and residue field k , and C/K a smooth projective curve specified by an affine equation $f(x, y) = 0$. We show that the Newton polytope Δ_v of f with respect to v_K , under a ‘generic’ condition called Δ_v -regularity, determines explicitly

- the minimal regular normal crossings model of C over O_K ;
- whether C and $\text{Jac } C$ have good, semistable and tame reduction;
- the action $\text{Gal}(\bar{K}/K) \curvearrowright H_{\text{ét}}^1(C_{\bar{K}}, \mathbb{Q}_l)$ when C is tamely ramified, and the action on its wild inertia invariants in general, for $l \neq \text{char } k$;
- a basis of global sections of the relative dualising sheaf;
- the reduction map on points from the generic to the special fibre.

A regular model is usually constructed by starting with any model over O_K , and repeatedly blowing up at points and components of the special fibre and taking normalisations. In effect, the toric resolution replaces repeated blow-ups along coordinate axes, and for Δ_v -regular curves one such resolution is enough to get a good model.

1.1. Regular model. Our main objects are the Newton polytopes of the defining equation $f = \sum a_{ij}x^i y^j$ of the curve C over K and over O_K ,

$$\begin{aligned} \Delta &= \text{convex hull}((i, j) \mid a_{ij} \neq 0) \subset \mathbb{R}^2, \\ \Delta_v &= \text{lower convex hull}((i, j, v_K(a_{ij})) \mid a_{ij} \neq 0) \subset \mathbb{R}^2 \times \mathbb{R}. \end{aligned}$$

(The lower convex hull is those points Q in the convex hull such that $Q - (0, 0, \epsilon)$ is not in it for any $\epsilon > 0$.) Thus, Δ is a 2-dimensional convex polygon, and above every point $P \in \Delta$ there is a unique point $(P, v(P)) \in \Delta_v$. The function $v : \Delta \rightarrow \mathbb{R}$ is piecewise affine, and breaks Δ into 2-dimensional v -faces and 1-dimensional v -edges, the images of faces and edges of the polytope Δ_v under the homeomorphic projection to Δ . We will associate

- denominators $\delta_L, \delta_F \in \mathbb{N}$ to every v -edge L and v -face F (see 3.2),
- proper 1-dimensional schemes \bar{X}_F/k to every v -face F , and finite 0-dimensional schemes X_L/k to every v -edge L (see 3.4-3.7),

Date: January 12, 2021.

This research is supported by the EPSRC grant EP/M016838/1.

- slopes $[s_L^1, s_L^2]$ at every v -edge L from adjacent v -faces (see 3.12).
- $\Delta(\mathbb{Z}) = \text{interior}(\Delta) \cap \mathbb{Z}^2$, the set of interior integer points.

We say that f (or C) is Δ_v -regular essentially¹ if all \bar{X}_F, X_L are smooth. Under these conditions, C is the completion of $\{f = 0\} \subset \mathbb{G}_m^2$ in the toric variety with fan Δ ; see §2. In particular, the genus of C is $|\Delta(\mathbb{Z})|$ and it has a basis of regular differentials $x^i y^j \frac{dx}{xyf'_y}$ indexed by $(i, j) \in \Delta(\mathbb{Z})$.

As an example, let $\pi \in K$ be a uniformiser, and take a curve (left)

$$C: y^2 + \pi x^3 y + x^3 + \pi^5 = 0$$

Then Δ is a quadrangle, with values of v on $\Delta \cap \mathbb{Z}^2$ as above (middle). It has v -faces F_1, F_2 with $\delta_{F_1} = 6, \delta_{F_2} = 3$, and five v -edges (4 outer, 1 inner). Here C has genus 2 and is Δ_v -regular, for any ground field K .

In §4 we construct a proper flat model \mathcal{C}_Δ/O_K of C/K , using a toroidal embedding associated to Δ_v . For Δ_v -regular curves the main result is

Theorem 1.1 (3.13+8.12). *If f is Δ_v -regular, then \mathcal{C}_Δ/O_K is a regular model of C/K with strict normal crossings. Its special fibre \mathcal{C}_k/k consists of:*

- (1) A complete smooth curve \bar{X}_F/k of multiplicity δ_F , for every v -face F .
- (2) For every v -edge L with slopes $[s_1^L, s_2^L]$ pick $\frac{m_i}{d_i} \in \mathbb{Q}$ so that

$$s_1^L = \frac{m_0}{d_0} > \frac{m_1}{d_1} > \dots > \frac{m_r}{d_r} > \frac{m_{r+1}}{d_{r+1}} = s_2^L \quad \text{with} \quad \left| \frac{m_i}{d_i} - \frac{m_{i+1}}{d_{i+1}} \right| = 1.$$

Then L gives $|X_L(\bar{k})|$ chains of \mathbb{P}^1 s of length r from² \bar{X}_{F_1} to \bar{X}_{F_2} and multiplicities $\delta_L d_1, \dots, \delta_L d_r$.³ The absolute Galois group $\text{Gal}(k^s/k)$ permutes the chains through its action on $X_L(\bar{k})$.

If f'_y is not identically zero, then the differentials

$$\pi^{\lfloor v(i,j) \rfloor} x^{i-1} y^{j-1} \frac{dx}{f'_y}, \quad (i, j) \in \Delta(\mathbb{Z})$$

form an O_K -basis of global sections of the relative dualising sheaf $\omega_{\mathcal{C}_\Delta/O_K}$.

In the example above, the special fibre is shown on the right. See Theorem 3.14 for a slightly more general statement, for arbitrary curves.

1.2. Reduction. There is a natural description, in terms of Δ_v , of the dual graph of the special fibre (6.1), of the minimal regular model with normal crossings (5.7) and of the reduction map of points to the special fibre (3.19). From these one can also deduce the following criterion for good, semistable, and tame⁴ reduction for Δ_v -regular curves and their Jacobians.

¹The condition for outer v -faces can be made slightly weaker, see 3.9.

²If L is at the boundary of Δ , it has only one adjacent v -face and there is no \bar{X}_{F_2} .

³If $r = 0$, this is interpreted as \bar{X}_{F_1} meeting \bar{X}_{F_2} transversally at $|X_L(\bar{k})|$ points in the inner case, and no contribution from L in the outer case. In the outer case, this happens precisely when $\delta_L = \delta_{F_1}$.

⁴By ‘tame’ we mean semistable over some tamely ramified extension of K ; this is automatically the case if $\text{char } k = 0$ or $\text{char } k > 2 \text{ genus}(C) + 1$

Write $\Delta(\mathbb{Z})^F \subseteq \Delta(\mathbb{Z})$ for points that are in the interiors of v -faces, and $\Delta(\mathbb{Z})^L$ for the others (lying on v -edges). We let subscripts, such as $\Delta(\mathbb{Z})_{\mathbb{Z}}$ or $\Delta(\mathbb{Z})_{\mathbb{Z}_p}^F$, indicate that we further restrict to points with $v(P) \in \mathbb{Z}$ or $v(P) \in \mathbb{Z}_p$, respectively. (When $p = 0$, the latter is interpreted as an empty condition). See §1.6 for other notation.

Theorem 1.2 (=6.7). *Suppose C/K is Δ_v -regular, of genus ≥ 1 . Then*

- (1) C is Δ_v -regular over every finite tame extension K'/K .
 - (2) C has good reduction $\Leftrightarrow \Delta(\mathbb{Z}) = F(\mathbb{Z})$ for some v -face F with $\delta_F = 1$.
 - (3) C is semistable \Leftrightarrow every principal v -face F (see 5.1) has $\delta_F = 1$.
 - (4) C is tame \Leftrightarrow every principal v -face F has $\text{char } k \nmid \delta_F$.
- In this case, C is Δ_v -regular over every finite extension K'/K .*

Suppose k is perfect, and let J be the Jacobian of C . Then

- (5) J has good reduction $\Leftrightarrow \Delta(\mathbb{Z}) = \Delta(\mathbb{Z})_{\mathbb{Z}}^F$.
- (6) J is semistable $\Leftrightarrow \Delta(\mathbb{Z}) = \Delta(\mathbb{Z})_{\mathbb{Z}}$.
- (7) J is tame $\Leftrightarrow \Delta(\mathbb{Z}) = \Delta(\mathbb{Z})_{\mathbb{Z}_p}$, where $p = \text{char } k$. In this case,
 - J has potentially good reduction $\Leftrightarrow \Delta(\mathbb{Z})^L = \emptyset$, and
 - J has potentially totally toric reduction $\Leftrightarrow \Delta(\mathbb{Z})^F = \emptyset$.

1.3. Étale cohomology. One of the fundamental invariants of a curve C over a global field is its global Galois representation and the associated L -function $L(C, s)$. To study it in practice, one needs to understand the corresponding local Galois representations. They can be quite intricate when C has bad reduction, but for tame Δ_v -regular curves we can give a complete description, as follows.

Let K be complete with perfect residue field k of characteristic $p \geq 0$. Assume that C/K is Δ_v -regular and fix $l \neq p$. The representation in question is the étale cohomology of C with the action of the absolute Galois group,

$$G_K = \text{Gal}(K^s/K) \curvearrowright H^1(C) = H_{\text{ét}}^1(C_{\bar{K}}, \mathbb{Q}_l).$$

Let $I_{\text{wild}} \triangleleft I_K \triangleleft G_K$ be the wild inertia and the inertia subgroups. It turns out that the action of I_K on the subspace of wild inertia invariants $H^1(C)^{I_{\text{wild}}=1}$ (which is all of $H^1(C)$ if C is tame) depends only on Δ_v , and has the following elementary description.

Fix a uniformiser π of K . Each point $P \in \Delta(\mathbb{Z})_{\mathbb{Z}_p}$ defines a tame character $\chi_P : I_K \rightarrow \{\text{roots of unity}\}$ by $\sigma \mapsto \sigma(\pi^{v(P)})/\pi^{v(P)}$. Let $V_{\text{tame}}^{\text{ab}}, V_{\text{tame}}^{\text{toric}}$ be the unique continuous representations of I_K over \mathbb{Q}_l that decompose over \mathbb{Q}_l as

$$V_{\text{tame}}^{\text{ab}} \cong_{\mathbb{Q}_l} \bigoplus_{P \in \Delta(\mathbb{Z})_{\mathbb{Z}_p}^F} (\chi_P \oplus \chi_P^{-1}), \quad V_{\text{tame}}^{\text{toric}} \cong_{\mathbb{Q}_l} \bigoplus_{P \in \Delta(\mathbb{Z})_{\mathbb{Z}_p}^L} \chi_P.$$

Then there is an isomorphism of I_K -modules (Theorem 6.4),

$$H^1(C)^{I_{\text{wild}}=1} \cong V_{\text{tame}}^{\text{ab}} \oplus V_{\text{tame}}^{\text{toric}} \otimes \text{Sp}_2.$$

In particular, the dimension of the wild inertia invariants is $2|\Delta(\mathbb{Z})_{\mathbb{Z}_p}^L|$.

When k is finite, we describe $H^1(C)^{I_{\text{wild}}=1}$ as a full G_K -representation. To a v -face F of Δ we associate a scheme $\bar{X}_F^{\text{tame}}/O_K$, that depends only on the coefficients a_P for P along F . It is tame, defined by an equation that has only one v -face, and its H^1 admits an explicit description in terms of point counting (Theorem 7.3). Similarly we get schemes X_L^{tame}/O_K

for v -edges L . Then there is an isomorphism of G_K -representations (see Theorem 7.2)

$$H^1(C)^{I_{\text{wild}}=1} \ominus \mathbf{1} \cong \bigoplus_{\substack{F \\ v\text{-faces}}} (H^1(\bar{X}_F^{\text{tame}}) \ominus \mathbf{1}) \oplus \bigoplus_{\substack{L \text{ inner} \\ v\text{-edges}}} (H^0(X_L^{\text{tame}}) \otimes \text{Sp}_2).$$

1.4. Applications. As an illustration, we

- recover Tate’s algorithm for elliptic curves (§9),
- determine regular models of Fermat curves $x^p + y^p = 1$ over \mathbb{Z}_p and tame extensions of \mathbb{Z}_p , that seem to be unavailable (§10),
- compute regular models and their differentials for some curves from the literature, in an elementary way (3.16, 3.18, 8.15).

Generally, on the computational side our motivation came from determining arithmetic invariants of curves that enter the Birch–Swinnerton-Dyer conjecture, such as the differentials, conductor and Tamagawa numbers. On the theoretical side, for Δ_v -regular curves we also recover explicit versions of various classical results: existence of (minimal) model with normal crossings, Saito’s criterion for wild reduction ([Sai, Thm 3]; see [Hal] as well), and Kisin’s theorem on the continuity of l -adic representations for H^1 of curves in the tame case (see 7.6). It also extends some computations of regular models as tame quotients [Vie, LoD, CES] to the wild case.

1.5. Remarks. (a) Note that Theorem 1.1 applies to curves with wild reduction, because the construction is ‘from the bottom’ (toroidal embedding) rather than ‘from the top’ (taking a quotient of a semistable model over some larger field). In fact, Δ_v -regularity is automatic for the most ‘irregular’ reduction types, the ones where the only points in $\Delta_v \cap \mathbb{Z}^3$ are vertices of Δ_v . (This is the case with the example above.) Then the shape of the special fibre of their regular models does not depend on the residue characteristic at all. This explains why for example elliptic curves $y^2 = x^3 + \pi$, $y^2 = x^3 + \pi^5$, $y^2 = x^3 + \pi x$, $y^2 = x^3 + \pi^3 x$ are always of Kodaira type II, II*, III, III*, irrespectively of whether $\text{char } k$ is 2, 3 or ≥ 5 .

(b) Also, the theorem produces a regular model with normal crossings, which is somewhat more pleasant than the minimal regular model for classification and computational purposes (as it can be encoded combinatorially in the dual graph of the special fibre), and sometimes for theoretical purposes as well; see e.g. [Sai, Hal].

(c) The technique appears to be useful for arithmetic schemes of higher dimension as well, and over more general local rings. However, we only treat the case of curves over DVRs here, as it seems sufficiently interesting and already sufficiently involved.

(d) Although we have chosen to formulate the results for global equations, the construction of the regular model is really a local process (in effect a version of a repeated blow up).

(e) In a way, our results for arithmetic surfaces are a natural extension of two classical lower-dimensional analogues — 1-dimensional schemes over a field, and over a DVR. For a plane curve over a field, Baker’s theory computes the genus, points at infinity and differentials from a Newton polygon of the defining equation; we review this in §2. And for univariate $f(x)$ with coefficients in a complete DVR, the Newton polygon of f reflects how f factors, in effect forcing a decomposition of $H_{\text{ét}}^0$ of the scheme $f = 0$; the results from §1.3 simply extend this to bivariate $f(x, y)$ and their $H_{\text{ét}}^1$.

(f) Finally, there is an obvious question as to what extent the theory extends to arbitrary curves, not just the Δ_v -regular ones. It appears that it does, but this requires additional work. We postpone this to a future paper.


1.6. Notation. Throughout the paper we use the following notation:

K, v_K	field with a discrete valuation; in §2, K is any field
O_K, k, p	ring of integers, residue field, char k
π	a fixed choice of a uniformiser of K
f	$= \sum a_{ij}x^i y^j$, equation in $K[x, y]$, not 0 or a monomial
f'_y	derivative of f with respect to y ; in §2, §8 we assume $f'_y \neq 0$, equivalently $K(C)/K(x)$ is separable
C	smooth projective curve over K birational to $f = 0$
Δ	Newton polygon of f over K in \mathbb{R}^2 , as in §1.1
Δ_v	Newton polytope of f over O_K in \mathbb{R}^3 , as in §1.1
v	piecewise affine function $v : \Delta \rightarrow \mathbb{R}$ induced by v_K , as in §1.1
∂	boundary (of a convex polygon in \mathbb{R}^2)
L, F	v -edges and v -faces of Δ , the images of 1- and 2-dimensional faces of Δ_v under the homeomorphic projection $\Delta_v \rightarrow \Delta$
$\Delta(\mathbb{Z})$	interior(Δ) $\cap \mathbb{Z}^2$; similarly $L(\mathbb{Z}), F(\mathbb{Z})$
$\bar{\Delta}(\mathbb{Z})$	closure(Δ) $\cap \mathbb{Z}^2$; similarly $\bar{L}(\mathbb{Z}), \bar{F}(\mathbb{Z})$
$\delta_P, \delta_L, \delta_F$	the denominator of $v(P)$ for $P \in \bar{\Delta}(\mathbb{Z})$, and the common denominators of $v(P)$ for $P \in \bar{L}(\mathbb{Z})$ or $\bar{F}(\mathbb{Z})$ (3.2)
$f _L, f _F$	restriction of f to a v -edge or a v -face (3.4)
\bar{f}_L, \bar{f}_F	reduction to $k[t], k[x, y]$ of the restrictions (3.5)
X_L, X_F	affine schemes $\{\bar{f} _L = 0\} \subset \mathbb{G}_{m,k}$ and $\{\bar{f} _F = 0\} \subset \mathbb{G}_{m,k}^2$ (3.7)
\bar{X}_F	completion of X_F with respect to its Newton polygon (3.7)
r.n.c.	flat proper regular O_K -model with normal crossings of a curve over K (exists in genus ≥ 1 , see [Liu, 9/1.7])
m.r.n.c.	minimal r.n.c. model (unique, see [Liu, 9/3.36])
\mathcal{C}_Δ/O_K	flat proper model of C/K constructed in §4, r.n.c. if C is Δ_v -regular (cf. 3.13, 3.14)
$\mathcal{C}_\Delta^{\min}/O_K$	m.r.n.c. model of a Δ_v -regular curve (3.13+5.7)
K^s, \bar{K}	separable and algebraic closure of K
G_K	$\text{Gal}(K^s/K)$, the absolute Galois group
$I_K \triangleleft G_K$	the inertia group
$I_{\text{wild}} \triangleleft I_K$	the wild inertia subgroup
Jac C	Jacobian of C
l	prime different from $p = \text{char } k$
Sp_2	standard representation $I_K \rightarrow \text{GL}_2(\mathbb{Q}_l), \sigma \mapsto \begin{pmatrix} 1 & t(\sigma) \\ 0 & 1 \end{pmatrix}$, where $t : I_K \rightarrow \mathbb{Z}_l$ is the l -adic tame character (see [TaN, 4.1.4])
\oplus, \ominus	direct sum and difference of representations
$\mathbf{1}$	trivial representation
$\mathbb{G}_{m,K}$	$\text{Spec } K[t, t^{-1}]$, affine line minus the origin
m_{*j}, m_{i*}	j th column and i th row of a matrix $M = (m_{ij})$
$\tilde{m}_{*j}, \tilde{m}_{i*}$	j th column and i th row of a matrix $M^{-1} = (\tilde{m}_{ij})$


All representations are finite-dimensional, models are flat and proper, lattices are affine, and polygons are convex lattice polygons (vertices in \mathbb{Z}^2).

See Table 1 for examples of curves and the special fibres of their m.r.n.c. models, with a glossary of notation used in the pictures.

Δ_v -regular



not Δ_v -regular



$xy^2 = x^4 + x^2 + \pi^3$ (good elliptic curve meeting III*). On the left is Δ broken into v -edges and v -faces, with the values of v on $\Delta \cap \mathbb{Z}^2$, and the special fibre of the m.r.n.c. model $\mathcal{C}_\Delta^{\min}$ of 3.13, 5.7.

Components of positive genus are marked g1, g2 etc. $y^2 = (x-1)(x-2)(x-3)(x-\pi)(x-2\pi)(x-3\pi)(x-4\pi)$ Dashed blue lines mark empty chains of \mathbb{P}^1 's, that is main components meeting transversally. Here, the special fibre consists of two genus 1 curves meeting at two points.

Removable 2-faces (= not touching interior points $\Delta(\mathbb{Z})$) are grey; they do not affect the m.r.n.c. model; see §5. Here $y^2 = x^5 + x + \pi^6$ has the same model as $y^2 = x^5 + x$, one genus 2 component (good reduction).

$y^2 = x^2(x-1)(x+2)(x+3) + \pi^4$ Contractible 2-faces (= meeting $\Delta(\mathbb{Z})$ in one point P with $v(P) \in \mathbb{Z}$) are grey and marked with an arc; see §5. They give a chain of \mathbb{P}^1 's between 2-faces at the end of the arc, here from the genus 1 component X_F to itself.

$(y-1)^2 = (x-1)(x-2)(x-3)^2(x-4) + \pi^4$: this is (iv) shifted by (3,1). It has a node at (3,1), and its depth cannot be determined from Δ_v alone. (Here and below the conditions of Thm. 3.13 are *not* satisfied and, in fact, \mathcal{C}_Δ is not regular.)

$(y-1)^2 = (x-1)^2(x-2)^2 + \pi^5$: as in (v), except the equation is reducible mod π , and there are two components meeting at two nodes, again of undetermined depth.

$(y-1)^2 = (x-2)^5 + \pi^2$: singular point from a 2-face, which is not a node; here, a cusp $y^2 = x^5$ at (2,1).

$y^2 = (x-1)(x-2)(x-3)^2(x-4) + \pi^4$: (iv) shifted by (3,0); singular point from an outer 1-face.

$x^4y^2 = x(y-x)^2 + \pi^3$: singular point from an inner 1-face.

TABLE 1. Examples and notation in the pictures. For the sample equations (right column) suppose char $k = 0$, say.

2. BAKER'S THEOREM

By a classical result going back to Baker in 1893 [Bak], under generic conditions the genus of a plane curve is the number of interior integral points in its Newton polygon Δ . Modern versions, over a general field, have been proved in [KWZ, Lemma 3.4], [BP, Thm 4.2] and [Bee, Thm 2.4]. There are extensions to higher-dimensional varieties and their invariants, notably over \mathbb{C} [Kou, Kho, BKK],[Bat, §4]; see also [DL, Thm 1.3] over finite fields. The question which curves admit a ‘Baker model’ is addressed in detail in [CV]. Here we give a slightly different (and elementary) proof of Baker’s theorem. It describes regular differentials and points at infinity as well, and we will need the steps later.

Notation 2.1. Let K be any field. Fix a polynomial, which is not zero or a monomial,

$$f = \sum_{i,j} a_{ij} x^i y^j \quad \in K[x, y].$$

Recall that the *Newton polygon* Δ of f is

$$\Delta = \text{convex hull}((i, j) \mid a_{ij} \neq 0) \subset \mathbb{R}^2.$$

It is the smallest convex set containing the exponents of non-zero monomials, and by assumptions it is a convex polygon with at least two vertices. Write

$$\begin{aligned} \Delta(\mathbb{Z}) &= \mathbb{Z}^2 \cap \text{interior of } \Delta, \\ \bar{\Delta}(\mathbb{Z}) &= \mathbb{Z}^2 \cap \Delta. \end{aligned}$$

If Δ is not a line segment, we call its 1-dimensional faces *edges*. We use ‘ $L \subset \partial\Delta$ ’ as a shorthand for ‘for L an edge of Δ ’. For such an edge L , write

$$L^* = L^*_{(\Delta)} = \text{unique affine function } \mathbb{Z}^2 \rightarrow \mathbb{Z} \text{ with } L^*|_L = 0 \text{ and } L^*|_{\Delta} \geq 0.$$

When Δ is a line segment, we view it as having two edges L_1, L_2 and $\Delta(\mathbb{Z}) = \emptyset$, and pick L_i^* as above (not unique in this case) so that $L_1^* = -L_2^*$.

If $L \cap \mathbb{Z}^2 = \{(i_0, j_0), \dots, (i_r, j_r)\}$, ordered⁵ along L , set

$$f_L(t) = \sum_{n=0}^r a_{i_n, j_n} t^n \in K[t].$$

Theorem 2.2. *Let $f \in K[x, y]$, with Newton polygon $\Delta \subset \mathbb{R}^2$. If $C_0 : f = 0$ is a smooth⁶ curve in $\mathbb{G}_{m,K}^2$ and f_L is square-free for every edge L , then*

- (1) *The non-singular complete curve C birational to C_0 has genus $|\Delta(\mathbb{Z})|$.*
- (2) *C has a basis of regular differentials, non-vanishing on C_0 ,*

$$\omega_{ij} = x^{i-1} y^{j-1} \frac{dx}{f_y}, \quad (i, j) \in \Delta(\mathbb{Z}).$$

(If $f'_y = 0$, equivalently $K(C)/K(x)$ is inseparable, then swap $x \leftrightarrow y$.)

- (3) *There is a natural bijection that preserves $\text{Gal}(K^s/K)$ -action,*

$$C(\bar{K}) \setminus C_0(\bar{K}) \xrightarrow{1:1} \coprod_{L \subset \partial\Delta} \{\text{roots of } f_L \text{ in } \bar{K}^\times\}.$$

- (4) *If P corresponds to a root of f_L via (3), then*

$$\begin{aligned} (4a) \quad \text{ord}_P x^i y^j &= L^*(i, j) - L^*(0, 0) && \text{for } (i, j) \in \mathbb{Z}^2; \\ (4b) \quad \text{ord}_P \omega_{ij} &= L^*(i, j) - 1 && \text{for } (i, j) \in \Delta(\mathbb{Z}). \end{aligned}$$

⁵The two possible orders give $f_L(t)$ vs $f_L(1/t)$; this choice does not affect anything

⁶smooth=non-singular over \bar{K} , and square-freeness is also smoothness; recall that Δ =line segment is allowed (non-connected curves) but not Δ =point or \emptyset (not a curve)

Proof. Let C be the non-singular complete curve birational to C_0 , possibly non-connected. The statements are easy when Δ is a line segment (C_0 =union of \mathbb{G}_m s, C =union of \mathbb{P}^1 s), so assume that $\text{vol}(\Delta) > 0$. We will cover C by charts C_L , $C_0 \hookrightarrow C_L \hookrightarrow C$, indexed by the edges L of Δ .

First, let $\phi : (i, j) \mapsto mj - ni + s$ be any affine function $\mathbb{Z}^2 \rightarrow \mathbb{Z}$ which is non-negative on Δ and zero on some vertex; so $l = (m, n)$ is a primitive generator of the line $L = \phi^{-1}(0)$, going counterclockwise around Δ .⁷ Write $mn' - m'n = 1$ with $m', n' \in \mathbb{Z}$. In terms of $X = x^m y^n, Y = x^{m'} y^{n'}$,

$$f(x, y) = X^2 Y^2 (f_L(X) + Yh(X, Y)), \quad h \in K[X^{\pm 1}, Y].$$

The univariate polynomial f_L is a non-zero constant when $L \cap \Delta$ =vertex, and is separable (by assumption) of positive degree when $L \cap \Delta$ =edge. In both cases,

$$C_L : f_L(X) + Yh(X, Y) = 0$$

is a smooth curve in $\mathbb{G}_m \times \mathbb{A}^1$ with $C_L \cap \mathbb{G}_m^2 = C_0$; so $C_0 \hookrightarrow C_L \hookrightarrow C$. The \bar{K} -points in $C_L \setminus C_0$ have $Y = 0$ and correspond to the roots of f_L in $(K^s)^\times$, with the same Galois action. At these points,

$$\begin{aligned} (X =) \quad x^m y^n & \text{ is regular non-vanishing,} \\ (Y =) \quad x^{m'} y^{n'} & \text{ is regular vanishing.} \end{aligned}$$

Conversely, pick $P \in C \setminus C_0$. Either x or y has a zero or a pole at P , so

$$(i, j) \mapsto \text{ord}_P x^i y^j$$

is a non-trivial linear map $\mathbb{Z}^2 \rightarrow \mathbb{Z}$. It has a rank 1 kernel spanned by some primitive vector $(b, -a)$. Changing its sign if necessary, it follows from the above construction that P is 'visible' on C_L obtained from $\phi : (i, j) \mapsto ai + bj + c$, for a unique c . Finally,

$$L^*(i, j) = ai + bj + c = \text{ord}_P x^i y^j + c, \quad L^*(0, 0) = c.$$

This proves (3) and (4a).

(4b) Suppose P corresponds to a root of f_L via (3).

It suffices to prove $\text{ord}_P \omega_{ij} = L^*(i, j) - 1$ for *one* $(i, j) \in \mathbb{Z}^2$ and use (4a) to get all (i, j) . By Lemma A.3, this claim is invariant under $\text{SL}_2(\mathbb{Z}) \subset \text{Aut } \mathbb{G}_m^2$, reducing to the case

$$L \subset y\text{-axis}, \quad C_0 : f_L(y) + xG(x, y) = 0, \quad \omega_{11} = \frac{dx}{f_L + xG'_y}.$$

Since $f_L(y)$ is assumed to be separable and $P \in y$ -axis, the denominator of ω_{11} does not vanish at P . Because dx is regular non-vanishing on $\mathbb{A}^1 \times \mathbb{G}_m$, we get $\text{ord}_P \omega_{11} = 0$, as required.

(1), (2) The total degree of any ω_{ij} is the sum of $\text{ord}_P \omega_{ij} = L^*(i, j) - 1$ over all $L \subset \partial\Delta$, taken with multiplicity $\deg f_L$. An elementary calculation (Lemma A.2) shows that this sum is $2|\Delta(\mathbb{Z})| - 2$. This proves (1), as the degree of a differential form on a complete genus g curve is $2g - 2$. Finally, the ω_{ij} with $(i, j) \in \Delta(\mathbb{Z})$ form a basis of regular differentials, as there are g of them and there cannot be a linear relation between monomials $x^i y^j$ of degree $< \deg f$. (See Remark 2.3 for a slightly stronger statement.) \square


Remark 2.3. If $f = \sum a_{ij} x^i y^j$ satisfies the conditions of the theorem, the differentials ω_{ij} with $(i, j) \in \bar{\Delta}(\mathbb{Z})$ have exactly one linear relation between them up to scalars⁸, namely

⁷Thus $\phi = L^*$ in the case $L \cap \Delta$ =edge

⁸It is not hard to see that this is a basis of differentials that are regular on C_0 and have at most simple poles on $C \setminus C_0$, though we will not need this.

$\sum a_{ij}\omega_{ij} = 0$. This is clear from the minimality (=square-freeness) of the defining equation $f = 0$.


Example 2.4. The following are genus 0 curves, with $\Delta(\mathbb{Z}) = \emptyset$:



Example: Newton polygons of genus 0 curves (say char $K \nmid 2mn$)
 $y^n = 1$, $y = x^n + 1$, $x^2 + y^2 = 1$, $xy + x + y = 1$, $x^n - 2 = y(x^m - 1)$

Conversely, Rabinowitz [Rab, Thm 1], extending Arkinstall [Ark, Lemma 1] shows that a convex lattice polygon $\Delta \subset \mathbb{R}^2$ with $\Delta(\mathbb{Z}) = \emptyset$ is $\text{GL}_2(\mathbb{Z})$ -equivalent to one of the pictures above: a line segment, a right triangle with legs 1, n or 2, 2, or a trapezium of height 1 (right-angled as above, if desired).

Example 2.5. Curves of genus ≤ 4 always have an equation satisfying Baker’s theorem if K is large enough, but not in general in higher genus [CV]. For instance, in genus 3, every smooth projective curve C/K is either hyperelliptic or a plane quartic (but not both). If, say, $K = \bar{K}$, then C has an equation $f = 0$ with Δ either



A basis of regular differentials is $\frac{dx}{f'_y}, x\frac{dx}{f'_y}, y\frac{dx}{f'_y}$ and $\frac{dx}{f'_y}, x\frac{dx}{f'_y}, x^2\frac{dx}{f'_y}$, respectively. Similarly, in genus ≤ 2 all curves have hyperelliptic models when $K = \bar{K}$, and there are seven (disjoint) types in genus 4 [CV, §7].

Remark 2.6. Let $C_0 : f = 0$ be an equation in $\mathbb{G}_{m,K}^2$, with $\text{vol}(\Delta) > 0$.

(a) If C_0 satisfies the conditions of Theorem 2.2, the proof shows that the associated smooth complete curve C is covered by charts

$$(2.7) \quad C = \bigcup_{L \subset \partial\Delta} C_L, \quad C_L \hookrightarrow \mathbb{G}_m \times \mathbb{A}^1 \text{ closed.}$$

These $\mathbb{G}_m \times \mathbb{A}^1$ cover, up to a finite set, a toric variety \mathbb{T}_Δ with polytope Δ , and C is simply the closure of C_0 in \mathbb{T}_Δ . It is often called the toric resolution of C_0 on \mathbb{T}_Δ (see [CDV, §2.1], [Koe], [Cas]) or tropical compactification [Te]. Two standard examples are

Δ	\mathbb{T}_Δ	\supset	C
triangle on $(0, 0), (n, 0), (0, n)$	\mathbb{P}^2	\supset	curve of degree n ,
rectangle $[n, 0] \times [0, m]$	$\mathbb{P}^1 \times \mathbb{P}^1$	\supset	curve of degree (m, n) .

(b) The theorem constructs C_L as in (2.7) and the closure $C \subset \mathbb{T}_\Delta$ of C_0 for any f with Newton polygon Δ , whether C_0 is smooth or not. We call this scheme the *completion* C_0^Δ of C_0 with respect to its Newton polygon. Since \mathbb{T}_Δ is a complete variety, C_0^Δ is a proper scheme over K .

(c) When C_0 is smooth, the squarefreeness condition in Baker’s theorem to ensure smoothness of C is sufficient but not quite necessary. We say that C_0 is *outer regular* at an edge L of Δ if C_L is smooth.

When C_0 is outer regular at all L , in other words when C_0^Δ is smooth, the curve C_0 is often called nondegenerate with respect to Δ [CDV, §2, Def. 1]. For example, $(x-1)^2 + (y-1)^2 = 1$

is smooth in $\mathbb{G}_{m,\mathbb{C}}^2$, and its closure in $\mathbb{T}_\Delta = \mathbb{P}_{\mathbb{C}}^2$ is smooth as well. The restriction of f to the two edges of Δ on the coordinate axes is not squarefree, but C_0 is outer regular there.

(d) Fix Δ with $\text{vol}(\Delta) > 0$ and let $V = \{\text{vertices of } \Delta\}$. All $f \in \bar{K}[x, y]$ with Newton polygon Δ form a family $\mathcal{C} \subset \mathcal{P} \times \mathbb{T}_\Delta$ with a parameter variety $\mathcal{P} \cong \mathbb{A}^{\bar{\Delta}(\mathbb{Z}) \setminus V} \times \mathbb{G}_m^V$, flat over \mathcal{P} . Its generic member $f = 0$ is smooth⁹ [CDV, §2, Prop 1], irreducible (dimension count), and hence connected. Consequently, *any* C_0 with Newton polygon Δ is connected (Zariski connectedness), and

$$C_0^\Delta \text{ has arithmetic genus } |\Delta(\mathbb{Z})|,$$

since arithmetic genus is constant in flat families [Har, III.9.13]. In particular, when C_0 is an integral scheme and so defines a curve, we have

$$\text{geometric genus of } C_0 \leq |\Delta(\mathbb{Z})| \quad (\text{Baker's inequality}).$$

3. SETUP AND MAIN THEOREM

From now on, K is a field with a discrete valuation $v : K^\times \rightarrow \mathbb{Z}$, ring of integers O_K , uniformiser π and residue field k . As in §1.1, fix

$$f = \sum_{i,j} a_{ij} x^i y^j \in K[x, y],$$

and consider its Newton polytopes over K and over O_K ,

$$\begin{aligned} \Delta &= \text{convex hull} \left((i, j) \mid a_{ij} \neq 0 \right) \subset \mathbb{R}^2, \\ \Delta_v &= \text{lower convex hull} \left((i, j, v_K(a_{ij})) \mid a_{ij} \neq 0 \right) \subset \mathbb{R}^2 \times \mathbb{R}. \end{aligned}$$

Assume from now on that $\text{vol}(\Delta) > 0$. Above every point $P \in \Delta$ there is a unique point $(P, v(P)) \in \Delta_v$. This defines a piecewise affine function $v : \Delta \rightarrow \mathbb{R}$, and the pair (Δ, v) determines Δ_v .

Definition 3.1 (Faces). The images of the 0-, 1- and 2-dimensional (open) faces of the polytope Δ_v under the homeomorphic projection $\Delta_v \rightarrow \Delta$ are called *v-vertices*, *v-edges* and *v-faces* of Δ . Thus, *v-vertices* are points in \mathbb{Z}^2 , *v-edges* (denoted L) are homeomorphic to an open interval, and *v-faces* (denoted F) to an open disk.

Notation 3.2 (Integer points, denominators). For *v-edges* and *v-faces* write

$$L(\mathbb{Z}) = L \cap \mathbb{Z}^2, \quad F(\mathbb{Z}) = F \cap \mathbb{Z}^2.$$

As before, $\Delta(\mathbb{Z}) = (\text{interior of } \Delta) \cap \mathbb{Z}^2$, and we write $\bar{L}(\mathbb{Z})$, $\bar{F}(\mathbb{Z})$, $\bar{\Delta}(\mathbb{Z})$ to include points on the boundary. We decompose $\Delta(\mathbb{Z})$ into

$$\begin{aligned} \Delta(\mathbb{Z})^F &= \{P \in \Delta(\mathbb{Z}) \mid P \in F(\mathbb{Z}) \text{ for some } v\text{-face } F\}, \\ \Delta(\mathbb{Z})^L &= \{P \in \Delta(\mathbb{Z}) \mid P \notin F(\mathbb{Z}) \text{ for any } v\text{-face } F\}. \end{aligned}$$

We also use subscripts to restrict to points with $v(P)$ in a given set, such as

$$F(\mathbb{Z})_{\mathbb{Z}_p} = \{P \in F(\mathbb{Z}) \mid p \nmid \text{den}(v(P))\} = \{P \in F(\mathbb{Z}) \mid v(P) \in \mathbb{Z}_p\}.$$

(When $p = 0$, the above is interpreted as an empty condition.) We write $\Delta(\mathbb{Z})_{\mathbb{Z}}^F$ etc. for multiple constraints. The *denominator* δ_λ of a *v-edge* or a *v-face* λ is, equivalently (see Lemma A.1 for (1) \Leftrightarrow (3)),

- (1) common denominator of $v(P)$ for $P \in \bar{\lambda}(\mathbb{Z})$;
- (2) smallest $m \geq 1$ for which $\bar{\lambda}(\mathbb{Z})_{\frac{1}{m}\mathbb{Z}} = \bar{\lambda}(\mathbb{Z})$;

⁹This is subtle when $\text{char } K > 0$ and fails in higher dimensions [CDV, §2, below Cor. 1]

(3) index of the affine lattice Λ_λ spanned by $\bar{\lambda}(\mathbb{Z})$ in its saturation in \mathbb{Z}^2 .

If F is a v -face, and we write


$$v_F = \text{unique extension of } v|_F \text{ to an affine function } \mathbb{Z}^2 \rightarrow \mathbb{Q},$$

then $\Lambda_F = v_F^{-1}(\mathbb{Z})$ and \mathbb{Z}^2/Λ_F is cyclic of order δ_F , again by Lemma A.1.

Example 3.3. Let $C : f = 0$ be the genus 4 curve defined by

$$f = y^3 + \pi^2 x^2 y^2 + y^2 + \pi^3 x^6 + \pi x^3 + \pi^3.$$

Here Δ_v has v -faces F_1, F_2, F_3 with $\delta_F=3,3,6$, v -edges with $\delta_L=1,1,1,2,3,3,3$ and five v -vertices (from the terms of f except for $\pi^2 x^2 y^2$ that has too high valuation). Below (left) is a picture of Δ broken into v -faces, and the values of v on $\bar{\Delta}(\mathbb{Z})$. ‘Capital’ ones indicate that the corresponding coefficient of f has exactly that valuation (and not larger):



In this example, $|\Delta(\mathbb{Z})| = 4$, with $|\Delta(\mathbb{Z})^F| = |\Delta(\mathbb{Z})^L| = 2$.

To produce the regular model and describe the equations for the components of the special fibre (picture on the right), we need to extract monomials from f that come from a given face, restrict them to a sublattice, and reduce the resulting equation:

Definition 3.4 (Restriction). Let $g = \sum_{i \in \mathbb{Z}^n} c_i \mathbf{x}^i \in K[\mathbf{x}]$ be a polynomial in n variables $\mathbf{x} = (x_1, \dots, x_n)$, and $\emptyset \neq S \subset \mathbb{Z}^n$. Take Λ to be the smallest affine lattice with $S \subset \Lambda \subset \mathbb{Z}^n$, say of rank r . Choose an isomorphism $\phi : \mathbb{Z}^r \rightarrow \Lambda$, write $\mathbf{y} = (y_1, \dots, y_r)$ and define the *restriction*


$$g|_S = \sum_{i \in \phi^{-1}(S)} c_{\phi(i)} \mathbf{y}^i.$$

Different choices of ϕ change $g|_S$ by a $\text{GL}_r(\mathbb{Z})$ -transformations of the variables and multiplication by monomials. For a v -vertex/edge/face λ , set

$$f|_\lambda = f|_{\bar{\lambda}(\mathbb{Z})_{\mathbb{Z}}}.$$

Definition 3.5 (Reduction). Suppose $h \in K[x, y]$, and there are $c, m, n \in \mathbb{Z}$ such that $\tilde{h}(x, y) = \pi^c h(\pi^m x, \pi^n y)$ has coefficients in O_K , and $h \in K[x, y]$ and $\tilde{h} \bmod \pi \in k[x, y]$ have the same Newton polygon. Then we say that $\tilde{h} \bmod \pi$ is the *reduction* of h , and denote it \bar{h} .

Example 3.6. Let $f = y^2 - x^3 - \pi$, and F the unique v -face of Δ . In the notation of 3.4 and 3.5, the lattice $\Lambda = \mathbb{Z}(3, 0) + \mathbb{Z}(0, 2)$ has index $\delta_F = 6$ in \mathbb{Z}^2 , and $f|_F = f|_{\bar{F}(\mathbb{Z})_{\mathbb{Z}}} = f|_\Lambda$ and its reduction $\bar{f}|_F$ are as follows:



Definition 3.7 (Components X_F, X_L). In the notation of 3.4, 3.5, define k -schemes

- (1) $X_L: \{\overline{f|_L} = 0\} \subset \mathbb{G}_{m,k}$ for each v -edge L of Δ ,
- (2) $X_F: \{\overline{f|_F} = 0\} \subset \mathbb{G}_{m,k}^2$ for each v -face F of Δ ,
- (3) $\bar{X}_F =$ completion of X_F with respect to its Newton polygon (2.6b); it is a geometrically connected scheme, proper over k (2.6d).

The vertices and edges of the Newton polygon of $\overline{f|_F}$ correspond to those of F (cf. example above), and so there is a bijection from Baker's Theorem,

$$(3.8) \quad \bar{X}_F(\bar{k}) \setminus X_F(\bar{k}) \quad \longleftrightarrow \quad \prod_{L \subset \partial F} X_L(\bar{k}).$$

We can now define Δ_v -regularity. Modulo one allowed exception, it is simply saying that all X_F need to satisfy Baker's Theorem assumptions:

Definition 3.9 (Δ_v -regularity). We say that C (or f) is Δ_v -regular if all X_F, X_L are smooth over k , except that for v -edges $L \subset \partial F$ with $L \subset \partial \Delta$ and $\delta_L = \delta_F$, we drop the assumption that X_L is smooth but require \bar{X}_F to be outer regular at L , i.e. smooth at the points that correspond to L via (3.8).

It is not hard to see that regularity does not depend on the choice of π and those made in 3.4.

Example 3.10. The curve C/K of Example 3.3 is Δ_v -regular iff $\text{char } k \neq 3$. (There is only one v -edge to check, all other restrictions are linear).

Remark 3.11 (Base change). Suppose K'/K is an extension of discretely valued fields of ramification degree e , with residue field extension k'/k .

For a v -face F , write X'_F for the scheme X_F computed for f as an element of $K'[x, y]$. Let Λ_F be the affine lattice spanned by $\bar{F}(\mathbb{Z})_{\mathbb{Z}}$, so that \mathbb{Z}^2/Λ_F is cyclic of order δ_F (see 3.2). Let $\Lambda_F \subseteq \Lambda'_F \subseteq \mathbb{Z}^2$ be the unique intermediate lattice with $(\Lambda'_F : \Lambda_F) = \text{gcd}(\delta_F, e)$. Let $\phi : \mathbb{G}_{m,k'}^2 = \text{Spec } k'[\Lambda'_F] \rightarrow \text{Spec } k[\Lambda_F] = \mathbb{G}_{m,k}^2$ be the corresponding cover. Then

$$X'_F \cong X_F \times_{\mathbb{G}_{m,k}^2, \phi} \mathbb{G}_{m,k'}^2,$$

and a similar statement holds for v -edges. Thus,

- (1) If $\text{gcd}(\delta_F, e) = 1$, then X_F/k smooth $\Leftrightarrow X'_F$ is.
- (2) If $p \nmid \text{gcd}(\delta_F, e)$, then ϕ is étale, so X_F/k smooth $\Rightarrow X'_F$ is.
- (3) If $p \nmid \text{gcd}(\delta_F, e)$ for every v -face F of Δ , e.g. K'/K is tame, then

$$C/K \text{ is } \Delta_v\text{-regular} \quad \Rightarrow \quad C/K' \text{ is } \Delta_v\text{-regular}.$$

Finally, we need slopes of two v -faces F_1, F_2 meeting at a v -edge L . They will control the chains of \mathbb{P}^1 s between \bar{X}_{F_1} and \bar{X}_{F_2} :

Definition 3.12 (Slopes). Every v -edge L is either *inner*, bounding two v -faces, say F_1 and F_2 , or *outer*, bounding F_1 only. Choose $P_0, P_1 \in \mathbb{Z}^2$ with $L_{(F_1)}^*(P_0) = 0$, $L_{(F_1)}^*(P_1) = 1$ (see 2.1). The slopes $[s_1^L, s_2^L]$ at L are

$$s_1^L = \delta_L (v_1(P_1) - v_1(P_0)), \quad s_2^L = \begin{cases} \delta_L (v_2(P_1) - v_2(P_0)) & \text{for } L \text{ inner,} \\ \lfloor s_1^L - 1 \rfloor & \text{for } L \text{ outer,} \end{cases}$$

where v_i is the unique affine function $\mathbb{Z}^2 \rightarrow \mathbb{Q}$ that agrees with v on F_i .


In §4 from Δ_v we construct a scheme \mathcal{C}_Δ/O_K . The reader is invited to skip the construction completely since regularity and the special fibre of \mathcal{C}_Δ can be described solely in terms of the above data. The results are:

Theorem 3.13 (Model in the Δ_v -regular case). *Suppose $C : f = 0$ is Δ_v -regular. Then \mathcal{C}_Δ/O_K is a regular model of C/K with strict normal crossings. Its special fibre \mathcal{C}_k/k is as follows:*

- (1) *Every v -face F of Δ gives a complete smooth curve \bar{X}_F/k of multiplicity δ_F and genus $|F(\mathbb{Z})_{\mathbb{Z}}|$.*
- (2) *For every v -edge L with slopes $[s_1^L, s_2^L]$ pick $\frac{m_i}{d_i} \in \mathbb{Q}$ so that*

$$s_1^L = \frac{m_0}{d_0} > \frac{m_1}{d_1} > \dots > \frac{m_r}{d_r} > \frac{m_{r+1}}{d_{r+1}} = s_2^L \quad \text{with} \quad \left| \frac{m_i}{d_i} - \frac{m_{i+1}}{d_{i+1}} \right| = 1.$$

Then L gives $|X_L(\bar{k})|$ chains of \mathbb{P}^1 s of length r from \bar{X}_{F_1} to \bar{X}_{F_2} (and open-ended in the outer case) and multiplicities $\delta_L d_1, \dots, \delta_L d_r$.¹⁰ The group G_k permutes the chains by its natural action on $X_L(\bar{k})$.



Contributions to \mathcal{C}_k from an inner v -edge L_1 and an outer v -edge L_2

This is a special case of the following more general version:

Theorem 3.14 (General case). *Suppose $f(x, y) = 0$ defines a 1-dimensional scheme $C_0 \subset \mathbb{G}_{m, K}^2$. Then \mathcal{C}_Δ/O_K is a proper flat model of the completion $C = C_0^\Delta$. Its special fibre \mathcal{C}_k is a union of closed subschemes \bar{X}_F indexed by v -faces F and $X_L \times_k \Gamma_L$ indexed by v -edges L , all 1-dimensional over k :*

- (1) *The X_L, X_F, \bar{X}_F are as in 3.7. The \bar{X}_F are geometrically connected, have arithmetic genus $|F(\mathbb{Z})_{\mathbb{Z}}|$, and come with multiplicity δ_F in the special fibre (i.e. defined by $\bar{f}_F^{\delta_F} = 0$ in \mathcal{C}_k).*
- (2) *For every v -edge L with slopes $[s_1^L, s_2^L]$ pick $\frac{n_i}{d_i} \in \mathbb{Q}$ so that*

$$s_1^L = \frac{n_0}{d_0} > \frac{n_1}{d_1} > \dots > \frac{n_r}{d_r} > \frac{n_{r+1}}{d_{r+1}} = s_2^L \quad \text{with} \quad \left| \frac{n_i}{d_i} - \frac{n_{i+1}}{d_{i+1}} \right| = 1.$$

Let $\Gamma_L = \Gamma_L^1 \cup \dots \cup \Gamma_L^r$ be a chain of \mathbb{P}_k^1 s, with multiplicities $\delta_L d_i$, meeting transversally: $\infty \in \Gamma_L^i$ is identified with $0 \in \Gamma_L^{i+1}$;

when $r = 0$, let $\Gamma_L = \text{Spec } k$, viewing it as $0 \in \Gamma_L^1$ and $\infty \in \Gamma_L^r$.

- (3) *The subscheme $X_L \times \{0\} \subset X_L \times \Gamma_L^1$ is identified with the subscheme of $\bar{X}_{F_1} \setminus X_{F_1}$ that corresponds to L via (3.8); similarly for $X_L \times \{\infty\} \subset X_L \times \Gamma_L^r$ and $\bar{X}_{F_2} \setminus X_{F_2}$ when L is inner.*
- (4) *The intersections of \bar{X}_{F_i} ($i = 1, 2$) with $X_L \times \Gamma_L$ and, when $r = 0$, of \bar{X}_{F_1} with \bar{X}_{F_2} at X_L are transversal. In other words, the intersection, as a scheme, is given by $\bar{f}_L^{\delta_L} = 0$.*

The model \mathcal{C}_Δ is given by explicit charts in §4. It is geometrically regular at

- the smooth locus of X_F , for every v -face F ,
- (smooth locus of X_L) $\times \Gamma_L$, for every v -edge L ,

¹⁰If $r = 0$, this is interpreted as \bar{X}_{F_1} meeting \bar{X}_{F_2} transversally at $|X_L(\bar{k})|$ points in the inner case, and no contribution from L in the outer case. In the outer case, this happens precisely when $\delta_L = \delta_{F_1}$.

- the smooth points of $\bar{X}_F \setminus X_F$ that correspond to L via (3.8), when $L \subset \partial F$ is outer with $\delta_L = \delta_F$, and $r = 0$ in (2).

If C_0 is Δ_v -regular, then C is smooth, and C_Δ/O_K is its r.n.c. model.

The theorems are proved in §4. See Table 1 for some examples, both Δ_v -regular and not. We end this section with a few comments:

Remark 3.15 (Hirzebruch-Jung). To see that sequences as in 3.14(2) exist, take all numbers in $[s_2^L, s_1^L] \cap \mathbb{Q}$ of denominator $\leq \max(\text{den } s_1^L, \text{den } s_2^L)$ in decreasing order. This is essentially a Haros (=Farey) series, and so satisfies the determinant condition [HW, Ch. III, Thm. 28]. Now repeatedly remove, in any order, terms of the form (*loc. cit.*, Thm. 29)

$$\dots > \frac{a}{b} > \frac{a+c}{b+d} > \frac{c}{d} > \dots \quad \mapsto \quad \dots > \frac{a}{b} > \frac{c}{d} > \dots,$$

until this is no longer possible; this corresponds to blowing down \mathbb{P}^1 s of self-intersection -1 , see (3.17) below. The resulting minimal sequence is unique¹¹. If $(s_2^L, s_1^L) \cap \mathbb{Z} = \{N, \dots, N+a\}$ is non-empty, it has the form

$$s_1^L = \frac{n_0}{d_0} > \dots > \frac{n_k}{d_k} > N+a > \dots > N+1 > N > \frac{n_l}{d_l} > \dots > \frac{n_{r+1}}{d_{r+1}} = s_2^L,$$

with d_0, \dots, d_k decreasing and d_l, \dots, d_{r+1} increasing. If $s_2^L > 0$, say (else shift by an integer), the numbers $N > \frac{n_l}{d_l} > \dots > \frac{n_{r+1}}{d_{r+1}}$ are the approximants of the Hirzebruch-Jung continued fraction expansion of s_2^L , see [Ful, §2.6]. (Similarly, for the first terms consider the expansion of $1 - s_1^L$.) These are designed to resolve toric singularities, which is essentially what the proof of 3.14 does, so their appearance is expected.

Remark 3.16 (Minimal vs normal crossings). If C/K is Δ_v -regular, the theorem constructs a r.n.c. model, in fact m.r.n.c. after an easy modification (see §5). Recall that this is *not* the same as the minimal regular model: the latter has generally fewer components but more complicated singularities in the special fibre. A r.n.c. model is encoded by the dual graph and the component multiplicities, so it is somewhat easier to work with (and to draw, for the matter). Note that if a component X of multiplicity m meets Y_1, \dots, Y_n of multiplicities m_1, \dots, m_n transversally, then the self-intersection

$$(3.17) \quad X \cdot X = -\frac{m_1 + \dots + m_n}{m} \quad (\in -\mathbb{N}).$$

If X has genus 0 and $X \cdot X = -1$, then X may be blown down. However, when $n \geq 3$, the result is no longer r.n.c. For example, take the genus 12 curve C/\mathbb{Q}_7 from [GRS²],

$$x^7 = \frac{y^3 - 2y^2 - y + 1}{y^3 - y^2 - 2y + 1}.$$

Letting $y \mapsto \frac{2y-3}{3y-2}$ and clearing the denominator, we get a model (tree-like with no positive genus components, in agreement with [GRS², App. A])




¹¹E.g., m.r.n.c. models would not be unique otherwise

The components X_{F_1}, X_{F_2} (multiplicity 21) have self-intersection -1 . If they are blown down, we find components of multiplicity 14 meeting ‘7,6 and 1’, so they get self-intersection -1 and can be blown down again. Ditto for ‘6’ and then ‘3’, resulting in a minimal regular model which has one multiplicity 1 component with two (bad) singularities on the special fibre.

Example 3.18. Here is an example for Theorem 3.14 (and not just 3.13). Consider the curve C_5/\mathbb{Q}_2 from [PSS, §13.3]. Letting $x \mapsto x+1, y \mapsto xy+1$ we find an equation

$$(3x^3+6x^2)y^3+9xy^2-(2x^3-6)y-4x^2-6x-4=0.$$

Here Δ_v and the special fibre of the model \mathcal{C}_Δ are as follows:



At the v -face F_1 (in red), the reduced equation is¹² $\overline{f_{F_1}} = XY + X + Y + 1$ and it gives two irreducible components $X_{F_1^a} : X = 1$ and $X_{F_1^b} : Y = 1$ meeting transversally at $P = (1, 1)$; both have multiplicity $\delta_{F_1} = 4$ on the special fibre. The point P is not regular, and the regular model has a chain of \mathbb{P}^1 s with multiplicity 4 from $X_{F_1^a}$ to $X_{F_1^b}$ (zigzag on the right picture). Its length is not determined by Δ_v , and requires a computation in the local ring at P . (It is actually 5, see [PSS, §13.3].)

Remark 3.19 (Reduction of points). Let \mathcal{C}/O_K be any regular model of a curve C/K . Write N for the set of multiplicity 1 components of the special fibre \mathcal{C}_k of \mathcal{C} , and $X^{ns} \subseteq X$ for the smooth locus of X for $X \in N$. There is a natural reduction map

$$\text{red}: C(K) \rightarrow \mathcal{C}_k(k),$$

landing in $\coprod_{X \in N} X^{ns}(k)$ (see [Liu, 9/1.32]), and onto it if K is Henselian. (In particular, if $N = \emptyset$, then $C(K) = \emptyset$.)

Now suppose C/K is Δ_v -regular and $\mathcal{C} = \mathcal{C}_\Delta$. The components $X \in N$ come from three sources, and are indexed by:

- (1_i) triples (L, r, n) where L is an *inner* v -edge with $\delta_L = 1$, r a root of $\overline{f|_L}$ in k^\times , and $n \in (s_2^L, s_1^L) \cap \mathbb{Z}$, plus
- (1_o) pairs (L, r) where $L \subset \partial F$ is an *outer* v -edge, $\delta_L = 1$, $\delta_F > 1$ and r a root of $\overline{f|_L}$ in k^\times , plus
- (2) v -faces F with $\delta_F = 1$.

Write C_0 for the curve $f = 0$ in \mathbb{G}_m^2 , and let $P \in C(K)$ be a point.

(a) Suppose $P \in C(K) \setminus C_0(K)$. By Baker’s theorem 2.2(3) it corresponds to a root r of f_L for an edge L of Δ . The edge breaks into v -edges that correspond to factors of f_L by its Newton polygon over the completion of K . The factor that has r as a root gives the v -edge indexing the component to which P reduces (Case (1_o)).

(b) Suppose $P \in C_0(K)$, so $x(P), y(P) \in K^\times$. The linear form

$$\mathcal{L}_P(i, j, z) = v(x(P))i + v(y(P))j + z$$

¹²as clear from the picture of Δ_v on the left, thanks to the scarcity of units in \mathbb{F}_2

Write $\tilde{L} = P_0 + \mathbb{R}\nu$ with $\nu = (v_x, v_y, v_z)$ such that $\delta\nu \in \mathbb{Z}^3$ is primitive and (v_x, v_y) goes counterclockwise along ∂F_1 . Write $P_1 - P_0 = (w_x, w_y)$ and $\omega_i = (w_x, w_y, \frac{n_i}{\delta d_i})$; the choices force $v_x w_y - v_y w_x = 1$. We also make one modification:

$$\text{If } L \text{ is outer, redefine } d_{r+1} = 0, n_{r+1} = -1, \omega_{r+1} = (0, 0, -\frac{1}{\delta}). \quad (\dagger)$$

These modified charts will be responsible for the ‘most outer’ points of \mathcal{C}_Δ .

In all cases, define planes in \mathbb{R}^3 (see above picture, when L is inner),

$$\mathcal{P}_{L,i} = \mathbb{R}\nu + \mathbb{R}\omega_i \quad i = 0, \dots, r+1.$$

The planes $P_0 + \mathcal{P}_{L,i}$ rotate around \tilde{L} ; the first one contains \tilde{F}_1 , and the last either contains \tilde{F}_2 (L inner) or is vertical (L outer). Let

$$\mathcal{P}_{L,i}^{\perp+} \subset \mathcal{P}_{L,i}^\perp$$

be the ray ($\cong \mathbb{R}_+$) for which the half-space $P_0 + \mathcal{P}_{L,i} + \mathcal{P}_{L,i}^{\perp+}$ contains Δ_v .

4.2. Toroidal embedding. For toroidal embeddings over a DVR, we follow [K2MS, §IV.3].

For a v -edge L , define polyhedral cones in $\mathbb{R}^2 \times \mathbb{R}_+$,

$$\begin{aligned} \text{0-dimensional cone} & \quad \sigma_0 = \{0\} \\ \text{1-dimensional cones} & \quad \sigma_{L,i} = \mathcal{P}_{L,i}^{\perp+} \quad (0 \leq i \leq r+1) \\ \text{2-dimensional cones} & \quad \sigma_{L,i,i+1} = \mathcal{P}_{L,i}^{\perp+} + \mathcal{P}_{L,i+1}^{\perp+} \quad (0 \leq i \leq r). \end{aligned}$$

The set Σ of all such cones from all L is a fan. The associated toric scheme

$$T_\Sigma = \bigcup_{\sigma \in \Sigma} T_\sigma, \quad T_\sigma = \text{Spec } O_K[\sigma^\vee \cap \mathbb{Z}^3]$$

(denoted $X_\Sigma = \bigcup X_\sigma$ in [K2MS]) is flat and separated over O_K (though not proper, as there are no 3-dimensional cones). Its generic fibre naturally contains $\mathbb{G}_{m,K}^2$, and we define

$$\mathcal{C}_\Delta = \text{closure of } \{f = 0\} \subset \mathbb{G}_{m,K}^2 \text{ in } T_\Sigma.$$

4.3. Charts. Let us describe the charts T_σ explicitly. Fix L as in 4.1, and tweak $\nu, \omega_i, -\omega_{i+1}$ into a basis of \mathbb{Z}^3 : take $\delta\nu$ (which is primitive) for the first basis vector. For the other two, shift $d_i\omega_i$ by an appropriate multiple $k_i\nu$ to get rid of the denominator in the last coordinate. In other words, pick $k_i \in \mathbb{Z}$ with

$$k_i \equiv -n_i \cdot (\delta v_z)^{-1} \pmod{\delta} \quad (0 \leq i \leq r+1).$$

As $\delta\nu$ is primitive, δv_z is invertible mod δ , so this is possible; when $\delta = 1$, set $k_i = 0$, say. For $0 \leq i \leq r$ define

$$M_{L,i} = \begin{pmatrix} \delta v_x & d_i w_x + k_i v_x & -d_{i+1} w_x - k_{i+1} v_x \\ \delta v_y & d_i w_y + k_i v_y & -d_{i+1} w_y - k_{i+1} v_y \\ \delta v_z & \frac{n_i}{\delta} + k_i v_z & -\frac{n_{i+1}}{\delta} - k_{i+1} v_z \end{pmatrix} = \begin{pmatrix} \delta v_x & d_i w_x & -d_{i+1} w_x \\ \delta v_y & d_i w_y & -d_{i+1} w_y \\ \delta v_z & \frac{n_i}{\delta} & -\frac{n_{i+1}}{\delta} \end{pmatrix} \begin{pmatrix} 1 & \frac{k_i}{\delta} & -\frac{k_{i+1}}{\delta} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = ST.$$

Our choices guarantee that $M = M_{L,i}$ has integer entries. Moreover,

$$\begin{aligned} \det M &= \det S = \delta v_x \frac{-d_i w_y n_{i+1} + d_{i+1} w_y n_i}{\delta} - \delta v_y \frac{-d_i w_x n_{i+1} + d_{i+1} w_x n_i}{\delta} + \delta v_z \cdot 0 \\ &= \delta v_x \frac{w_y}{\delta} - \delta v_y \frac{w_x}{\delta} = 1. \end{aligned}$$

Thus $M \in \text{SL}_3(\mathbb{Z})$, with inverse $M^{-1} = (\tilde{m}_{ij}) = T^{-1}S^{-1}$,

$$(4.1) \quad M_{L,i}^{-1} = \begin{pmatrix} 1 & -k_i/\delta & k_{i+1}/\delta \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} w_y/\delta & -w_x/\delta & 0 \\ n_{i+1}v_y - \delta d_{i+1}v_z w_y & -n_{i+1}v_x + \delta d_{i+1}v_z w_x & \delta d_{i+1} \\ n_i v_y - \delta d_i v_z w_y & -n_i v_x + \delta d_i v_z w_x & \delta d_i \end{pmatrix}.$$

Because of the choices of ν, ω_i and their orientation, we have

$$\begin{aligned} \mathcal{P}_{L,i} &= \mathbb{R}\nu + \mathbb{R}\omega_i = \mathbb{R}m_{*1} + \mathbb{R}m_{*2}, & \sigma_{L,i} &= \mathcal{P}_{L,i}^{\perp+} = \mathbb{R}_+ \tilde{m}_{3*}, \\ \mathcal{P}_{L,i+1} &= \mathbb{R}\nu + \mathbb{R}\omega_{i+1} = \mathbb{R}m_{*1} + \mathbb{R}m_{*3}, & \sigma_{L,i+1} &= \mathcal{P}_{L,i+1}^{\perp+} = \mathbb{R}_+ \tilde{m}_{2*}, \\ \mathcal{P}_{L,i} \cap \mathcal{P}_{L,i+1} &= \mathbb{R}\nu = \mathbb{R}m_{*1}, & \sigma_{L,i,i+1} &= \langle \mathcal{P}_{L,i}^{\perp+}, \mathcal{P}_{L,i+1}^{\perp+} \rangle = \mathbb{R}_+ \tilde{m}_{2*} + \mathbb{R}_+ \tilde{m}_{3*}, \end{aligned}$$

and monomial exponents from the dual cones are

$$\begin{aligned} \sigma_{L,i}^{\vee} \cap \mathbb{Z}^3 &= \mathbb{Z}m_{*1} + \mathbb{Z}m_{*2} + \mathbb{Z}_+m_{*3} \\ \sigma_{L,i+1}^{\vee} \cap \mathbb{Z}^3 &= \mathbb{Z}m_{*1} + \mathbb{Z}_+m_{*2} + \mathbb{Z}m_{*3} \\ \sigma_{L,i,i+1}^{\vee} \cap \mathbb{Z}^3 &= \mathbb{Z}m_{*1} + \mathbb{Z}_+m_{*2} + \mathbb{Z}_+m_{*3}. \end{aligned}$$

In other words, the coordinate transformation

$$\begin{aligned} (X, Y, Z) &= (x^{m_{11}}y^{m_{21}}\pi^{m_{31}}, x^{m_{12}}y^{m_{22}}\pi^{m_{32}}, x^{m_{13}}y^{m_{23}}\pi^{m_{33}}) = (x, y, \pi) \bullet M, \\ (x, y, \pi) &= (X, Y, Z) \bullet M^{-1} \end{aligned}$$

gives ring homomorphisms (as $\tilde{m}_{23}, \tilde{m}_{33} \geq 0$, the denominator makes sense)

$$K[x^{\pm 1}, y^{\pm 1}] \stackrel{M}{\cong} \frac{O_K[X^{\pm 1}, Y^{\pm 1}, Z^{\pm 1}]}{(\pi - X^{\tilde{m}_{13}}Y^{\tilde{m}_{23}}Z^{\tilde{m}_{33}})} \longleftrightarrow \frac{O_K[X^{\pm 1}, Y, Z]}{(\pi - X^{\tilde{m}_{13}}Y^{\tilde{m}_{23}}Z^{\tilde{m}_{33}})} = R,$$

and this is the inclusion $O_K[\sigma_0^{\vee}] \hookrightarrow O_K[\sigma_{L,i,i+1}^{\vee}]$ of **[K²MS]**. Thus,

$$\begin{aligned} T_{\sigma_{L,i,i+1}} &= \text{Spec } R, & T_{\sigma_{L,i+1}} &= \text{Spec } R[Z^{-1}], \\ T_{\sigma_{L,i}} &= \text{Spec } R[Y^{-1}], & T_{\sigma_0} &= \text{Spec } R[Y^{-1}, Z^{-1}], \end{aligned}$$

and these are all flat O_K -schemes of relative dimension 2. Glueing the $T_{\sigma_{L,i,i+1}}$ for varying L and i along their common opens gives T_{Σ} .

4.4. The closure of $f = 0$ in T_{Σ} . The model \mathcal{C}_{Δ} is covered by schematic closures of $\{f = 0\} \subset \mathbb{G}_{m,K}^2$ in the various $T_{\sigma_{L,i,i+1}}$. Explicitly, view f as a polynomial in x, y, π whose non-zero coefficients u_{ij} are in O_K^{\times} ,

$$(4.2) \quad f = \sum_{i,j} u_{ij} \pi^{v(a_{ij})} x^i y^j, \quad u_{ij} = \frac{a_{ij}}{\pi^{v(a_{ij})}}.$$

Let $M = (m_{ij}) \in \text{SL}_3(\mathbb{Z})$, for the moment arbitrary. Write $M^{-1} = (\tilde{m}_{ij})$, suppose $\tilde{m}_{23} \geq 0, \tilde{m}_{33} > 0$, and change variables $x = X^{\tilde{m}_{11}}Y^{\tilde{m}_{21}}Z^{\tilde{m}_{31}}, y = \dots, \pi = \dots$ according to M^{-1} . For some unique $n_Y, n_Z \in \mathbb{Z}$, we have

$$f = Y^{n_Y} Z^{n_Z} \mathcal{F}_M(X, Y, Z), \quad \mathcal{F}_M \in O_K[X^{\pm 1}, Y, Z], \quad Y \nmid \mathcal{F}_M, \quad Z \nmid \mathcal{F}_M.$$

Thus \mathcal{F}_M and f have the same coefficients u_{ij} , just different monomials, and $\mathcal{F}_M = 0$ defines a complete intersection

$$\mathcal{F}_M(X, Y, Z) = \pi - X^{\tilde{m}_{13}}Y^{\tilde{m}_{23}}Z^{\tilde{m}_{33}} = 0 \tag{*}$$

in $\text{Spec } O_K[X^{\pm 1}, Y, Z]$. Now let $M = M_{L,i}$ from 4.3. This Spec is $T_{\sigma_{L,i,i+1}}$, and (*) defines the open set $\mathcal{C}_{\Delta} \cap T_{\sigma_{L,i,i+1}} \subseteq \mathcal{C}_{\Delta}$. These cover all of \mathcal{C}_{Δ} .

4.5. Special fibre. The special fibre $\pi=0$ of T_Σ meets $(*)$ in $Y^{\tilde{m}_{23}}Z^{\tilde{m}_{33}}=0$. Its underlying reduced subscheme is $Z=0$ in the case (\dagger) , and $YZ=0$ otherwise. To find $(*) \cap \{Z=0\}$, write $f = Y^{n_Y}Z^{n_Z}\mathcal{F}_M$ as above. Noting that $Z=0 \Rightarrow \pi=0$, we obtain a k -scheme

$$(*) \cap \{Z=0\} = \text{Spec } k[X^{\pm 1}, Y]/(\mathcal{F}_M(X, Y, 0)).$$

The non-zero terms in $\mathcal{F}(X, Y, 0) \in k[X, Y]$ come from monomials $\pi^{v(a_{ij})}x^i y^j$ in (4.2) that, when rewritten in X, Y, Z , have minimal exponent ($=n_Z$) of Z . In other words, they minimise the following linear form $\mathbb{Z}^3 \rightarrow \mathbb{Z}$ on Δ_v ,

$$\phi: (\alpha, \beta, \gamma) \mapsto \text{ord}_Z(x^\alpha y^\beta \pi^\gamma) = \tilde{m}_{31}\alpha + \tilde{m}_{32}\beta + \tilde{m}_{33}\gamma.$$

Geometrically, $\mathcal{P} = \ker \phi$ is the plane spanned by the two columns of \tilde{M} , and ϕ is minimised on some face $\tilde{\lambda} \subset \Delta_v$ parallel to \mathcal{P} . Let λ be the projection of $\tilde{\lambda}$ onto Δ ; it is a v -vertex/edge/face. With an appropriate choice of the basis of the lattice in 3.4, 3.5, we find that

$$\bar{\mathcal{F}}_0(X, Y) = \bar{f}_\lambda.$$

This, and an identical computation for $(*) \cap \{Y=0\}$ to $M = M_{L,i}$ gives

$$\begin{aligned} \mathcal{F}_M(X, 0, 0) &= \bar{f}_L(X) \\ \mathcal{F}_M(X, Y, 0) &= \begin{cases} \bar{f}_{F_1}(X, Y) & i = 0 \\ \bar{f}_L(X) & i > 0 \end{cases} \\ \mathcal{F}_M(X, 0, Z) &= \begin{cases} \bar{f}_{F_2}(X, Y) & i = r, \text{ } L \text{ inner} \\ \bar{f}_L(X) & i < r. \end{cases} \end{aligned} \quad (**)$$

For a fixed v -face F and varying $L \subset \partial F$, the subsets $\bar{f}_F = 0$ of $\mathbb{G}_{m,k} \times \mathbb{A}_k^1$ are precisely the charts that define the completion \bar{X}_F . Glueing the pieces together, we find that \mathcal{C}_k is the union of \bar{X}_F over all v -faces F , and $X_L \times \Gamma_L$ over v -edges L , as claimed in Theorem 3.14 (and correct multiplicities). Since X_L are finite and so proper over k , and the \bar{X}_F and Γ_L are proper as well, \mathcal{C}_k/k is proper. By [Liu, 3/3.28] (or EGA IV.15.7.10), \mathcal{C}_Δ/O_K is proper.

4.6. Regularity. Let $P \in \mathcal{C}_k(\bar{k})$ be a point. It corresponds to an ideal

$$m_P = (\pi, X - x_0, Y - y_0, Z - z_0) \subset O_K[X^{\pm 1}, Y, Z]/(*).$$

To prove the claims in 3.14 when P is regular, we compute $\dim m_P/m_P^2$. The equation $\pi = X^{\tilde{m}_{13}}Y^{\tilde{m}_{23}}Z^{\tilde{m}_{33}}$ of $(*)$ eliminates π (recall that $\tilde{m}_{23}, \tilde{m}_{33} \geq 0$), and the assumptions of 3.14 on \bar{f}_L, \bar{f}_F eliminate either X or Y from m_P/m_P^2 ; so \mathcal{C} is regular at P under those assumptions.

Finally, suppose \mathcal{C} is Δ_v -regular. Then \mathcal{C} is geometrically regular at every point of \mathcal{C}_k , as above. As geometric regularity is an open condition, this implies that the generic fibre, which is \bar{C} by construction, is smooth. So \mathcal{C} is a regular model of \bar{C} in this case.

This proves Theorems 3.14 and 3.13. \square

5. MINIMAL MODEL

From now on, assume that C has genus ≥ 1 .

Here, in Theorem 5.7, we describe the *minimal regular normal crossings* (m.r.n.c.) model, a canonical regular model obtained from any normal crossings model by ‘removing unnecessary stuff’, that is repeatedly blowing down genus 0 components of self-intersection -1 . In our case, ‘unnecessary stuff’ comes from v -faces that avoid $\Delta(\mathbb{Z})$. We will see later that $\Delta(\mathbb{Z})$ also plays a crucial role in the description of étale cohomology and differentials.

Definition 5.1. Let F be a v -face of Δ . We say that

- F is *removable* if $F \cap \Delta(\mathbb{Z}) = \emptyset$.
- F is *contractible* if $F \cap \Delta(\mathbb{Z})$ has a unique point P , all v -edges $L \subset \partial F$ with $P \notin L(\mathbb{Z})$ are outer, and $v(P) \in \mathbb{Z}$.
- F is *principal* in all other cases.

We will refer to chains of \mathbb{P}^1 s from Theorem 3.13(2) as *inner chains* or *outer chains*, depending on whether they come from an inner or outer v -edge.

Proposition 5.2. *Let C/K be a Δ_v -regular curve, and \bar{X}_F a component of C_k from a v -face F , as in 3.13. The following conditions are equivalent:*

- (a₀) \bar{X}_F has genus 0 and meets the rest of the special fibre either at one point or at two points one of which lies on an outer chain.
- (b₀) F is $\text{GL}_2(\mathbb{Z})$ -equivalent to an $n \times 1$ right-angled triangle whose short sides are either outer v -edges (as in Table 2, top left) or one outer v -edge and one inner, of length 1, bounding a removable face.
- (c₀) F is removable.

Similarly, the following conditions are equivalent:

- (a₁) \bar{X}_F has genus 0 and meets the rest of the special fibre at two points, that are not outer chains.
- (b₁) F is $\text{GL}_2(\mathbb{Z})$ -equivalent to a polygon in Table 2, bottom row (an $n \times 1$ or 2×2 right-angled triangle, a chopped version of it, or a right-angled trapezoid with parallel sides $1, n$), with inner v -edges as shaded and of denominator 1.
- (c₁) F is contractible.




TABLE 2. Removable and contractible faces. Shading indicates the direction towards other parts of Δ (inner v -edges)

Proof. ¹³ Denote the four shapes in (b₁) by S_1, S_2, S_3, S_4 (cf. Table 2). The edges of S_3 and S_4 span \mathbb{Z}^2 , so such faces have $\delta_F = 1$; similarly $\delta_F|n$ for S_1 and $\delta_F|2$ for S_2 .

(b_i) \Rightarrow (c_i) Clear.

(b_i) \Rightarrow (a_i) By Theorem 3.13, outer v -edges contribute nothing, and inner ones give one chain (S_1, S_4) or two (S_2, S_3).

(a_i) \Rightarrow (b_i) Depending on the case, we have

$$(5.3) \quad 1 \text{ or } 2 = \# \text{ chains from } X_F = \sum_L |X_L(\bar{k})| = \sum_L |L(\mathbb{Z})_{\mathbb{Z}}|,$$

summing over inner v -edges L , and outer v -edges $L \subset \partial F$ with $\delta_L < \delta_F$ (see Theorem 3.13).

Write \tilde{F} for F restricted to the sublattice Λ spanned by $\bar{F}(\mathbb{Z})_{\mathbb{Z}}$. The genus of X_F is $|F(\mathbb{Z})_{\mathbb{Z}}|$, assumed to be 0. Therefore, by the Arkinstall–Rabinowicz classification of convex

¹³A direct classification-free proof of (a_i) \Leftrightarrow (c_i) would be very much appreciated

polygons with no interior points (Example 2.4), \tilde{F} is one of S_1, \dots, S_4 . As \mathbb{Z}^2/Λ is cyclic of order δ_F (see 3.2),

$$\begin{aligned} \delta_F = 1 &\Rightarrow \text{all } \delta_L = 1 \\ \delta_F > 1 &\Rightarrow \delta_L < \delta_F \text{ for all } L \text{ except possibly those} \\ &\quad \text{parallel to one fixed line in } \mathbb{Z}^2. \end{aligned} \tag{*}$$

Suppose LHS of (5.3) is 1. Then F has one inner face L with $L(\mathbb{Z})_{\mathbb{Z}} = \emptyset$, and all the others are outer with $\delta_L = \delta_F$. This eliminates $\tilde{F} = S_2$. In the S_3 and S_4 -case, $\delta_F = 1$ by (*), but then one inner face with $L(\mathbb{Z})_{\mathbb{Z}} = \emptyset$ is impossible by convexity, as there is no space behind L to contain interior integral points (while $\text{genus}(C) > 0$). So \tilde{F} is S_1 , the inner face gives the unique chain of \mathbb{P}^1 s, and two outer ones have $\delta_L = \delta_F$ which forces $\delta_F = 1$ by (*). Thus, F is S_1 (and, moreover, $n = 1$).

If the LHS of (5.3) is 2, then the same (but more tedious) considerations give the list of possible cases.

(c_i) \Rightarrow (b_i) Similar: as F has no interior integer points, it is one of the S_i by the Arkinstall–Rabinowicz classification; the rest is by inspection. \square

Remark 5.4. Let C/K be Δ_v -regular curve of genus $g > 0$.

- If $g = 1$ with $|\Delta(\mathbb{Z})_{\mathbb{Z}}^L| = 1$, there are no principal v -faces.
- In all other cases, e.g. if $g > 1$, there is always a principal v -face.

Indeed, in the first case, it follows from Theorem 3.13, after contracting the chains, that the special fibre is a reduced n -gon of \mathbb{P}^1 s (Kodaira type I_n). In the second case, if $\Delta(\mathbb{Z})^F \neq \emptyset$ or if $g = 1$, $\Delta(\mathbb{Z})_{\mathbb{Z}}^L = \emptyset$, then there is a principal face by definition. Otherwise for distinct $P, Q \in \Delta(\mathbb{Z})$, every point in the interior of the line segment PQ belongs to principal v -face(s).

Proposition 5.5. *Let C be a Δ_v -regular curve of genus $g > 0$. If $v(P) \in \mathbb{Z}$ for every $P \in \Delta(\mathbb{Z})$, then exactly one of the following holds:*

- $g > 1$ and every principal v -face F has $\delta_F = 1$.
- $g = 1$ and C is a semistable elliptic curve (type I_n with $n \geq 0$).
- $g = 1$, $C(K) = \emptyset$, and up to an isomorphism of the form $x \mapsto x^a y^b \pi^c$, $y \mapsto x^h y^i \pi^j$, Δ_v has one of the three exceptional shapes

$$(5.6) \quad \begin{array}{ccc} \begin{array}{c} 1 \\ | \\ 3/2 \\ | \\ 2 \end{array} \begin{array}{c} \diagdown \\ 1 \\ \diagup \\ 1 \end{array} \begin{array}{c} 1/2 \\ | \\ 1 \end{array} \begin{array}{c} \diagdown \\ 1 \\ \diagup \\ 1 \end{array} \begin{array}{c} 1/2 \\ | \\ 0 \end{array} \\ \begin{array}{c} 2 \end{array} \begin{array}{c} -3/2 \end{array} \begin{array}{c} -1 \end{array} \begin{array}{c} -1/2 \end{array} \begin{array}{c} 0 \end{array} \\ \begin{array}{c} F \end{array} \end{array} \quad \begin{array}{ccc} \begin{array}{c} 1 \end{array} \begin{array}{c} -3/2 \end{array} \begin{array}{c} -2 \end{array} \\ | \quad \quad \quad | \\ 1/2 \quad \quad \quad 3/2 \\ | \quad \quad \quad | \\ 0 \end{array} \begin{array}{c} -1/2 \end{array} \begin{array}{c} -1 \end{array} \\ \begin{array}{c} F \end{array} \end{array} \quad \begin{array}{ccc} \begin{array}{c} 1 \end{array} \begin{array}{c} \diagdown \\ 2/3 \\ \diagup \\ 1/3 \end{array} \begin{array}{c} 4/3 \end{array} \\ | \quad \quad \quad | \\ 1/3 \quad \quad \quad 5/3 \\ | \quad \quad \quad | \\ 0 \end{array} \begin{array}{c} -2/3 \end{array} \begin{array}{c} -4/3 \end{array} \begin{array}{c} -2 \end{array} \\ \begin{array}{c} F \end{array} \end{array}$$

The special fibre C_k consists of a unique genus 1 component \bar{X}_F , of multiplicity 2, 2, 3, respectively.

Proof. Case 2. $g = 1$, $|\Delta(\mathbb{Z})_{\mathbb{Z}}^L| = 1$. See Remark 5.4.

Case 3. $g = 1$, $|\Delta(\mathbb{Z})_{\mathbb{Z}}^F| = 1$. There is a unique principal v -face F . Up to $\text{GL}_2(\mathbb{Z})$ -equivalence, there are 15 possible convex lattice polygons with one interior point [Rab, Thm. 5]. Of these, in 12 cases the coordinates of the vertices and the interior point generate \mathbb{Z}^2 as an affine lattice, so $\delta_F = 1$. The other three shapes (Cases 1, 4, 9 of *loc.cit.*) are as in (5.6), with lattice index 2, 2, 3. Assume we are in one of these, so $\delta_F = 2, 2, 3$, respectively. After rescaling x, y by powers of π , the valuations can be made as in (5.6).

Case 1. $g \geq 2$. Suppose F is a principal v -face with $\delta_F > 1$. If $|F(\mathbb{Z})| < 2$, from 2.4 and the proof of Case 3, the only possibilities for F are

- (a) an $n \times 1$ right triangle; here $|F(\mathbb{Z})| = 0$, $1 < \delta_F \mid n$,
- (b) a 2×2 right triangle; here $|F(\mathbb{Z})| = 0$, $\delta_F = 2$,
- (c) the three cases in (5.6); here $|F(\mathbb{Z})| = 1$, $\delta_F = 2, 2, 3$ respectively.

In (b) and (c), as F is not the whole of Δ , at least one v -edge $L \subset \partial F$ must be inner with $\delta_L > 1$, contradicting $\Delta(\mathbb{Z}) = \Delta(\mathbb{Z})_{\mathbb{Z}}$. Similarly, in (a), as F is not removable or contractible, one of the corners of the v -edge $L \subset \partial F$ of length n must be in $\Delta(\mathbb{Z})$, and this again makes L inner with $\delta_L > 1$.

Finally, if $|F(\mathbb{Z})| \geq 2$, subdivide F into convex lattice polygons one of which has exactly one interior integer point, and apply the same argument to it to show that it is in (c), to get the same contradiction. \square

Theorem 5.7. *Let $C : f = 0$ be a Δ_v -regular curve, with $f = \sum a_{ij}x^i y^j$. Its m.r.n.c. model $\mathcal{C}_{\Delta}^{\min}/O_K$ has the following special fibre \mathcal{C}_k^{\min}/k .*

- (1) *Let $f_{nr} = \sum' a_{ij}x^i y^j$, the sum taken over those (i, j) that lie in the closure of some non-removable v -face of Δ . Then f_{nr} is Δ_v -regular.*
- (2) *Let \mathcal{C} be the model of $f_{nr} = 0$ of §4 and Theorem 3.13, and \mathcal{C}_k its special fibre. The principal v -faces of Δ are in 1-1 correspondence with principal components of \mathcal{C}_k in the sense of Xiao [Xi] (positive genus or meeting the rest of \mathcal{C}_k in ≥ 3 points).*
- (3) *Starting with \mathcal{C}_k , repeatedly contract \mathbb{P}^1 s of self-intersection -1 in the chains between two principal v -faces¹⁴. This gives \mathcal{C}_k^{\min}/k .*

Proof. Follows from Proposition 5.2 and Remark 3.15. \square

6. REDUCTION AND INERTIA

Throughout this section, C is a Δ_v -regular curve, and \mathcal{C}_k the special fibre of the r.n.c. model \mathcal{C}_{Δ} of §4 and Theorem 3.13. Write $\Gamma(\mathcal{C}_k)$ for its *dual graph*: it has geometric components as vertices, with an edge for every double point. Viewing it as a topological space, we compute its homology:

Theorem 6.1 (Homology of the dual graph). *Suppose C is Δ_v -regular. Then $H_1(\Gamma(\mathcal{C}_k), \mathbb{C})$ has dimension $|\Delta(\mathbb{Z})_{\mathbb{Z}}^L|$, and*

$$H_1(\Gamma(\mathcal{C}_k), \mathbb{Z}) \cong \ker \left(\bigoplus_{\substack{\text{inner} \\ v\text{-edges } L}} \mathbb{Z}[X_L(\bar{k})] \xrightarrow{\phi} \mathbb{Z}[v\text{-faces}] \right)$$


as G_k -modules. Here G_k acts naturally on $\mathbb{Z}[X_L(\bar{k})]$, trivially on $\mathbb{Z}[v\text{-faces}]$, and for $L \subset \partial F_1, \partial F_2$ and every $r \in X_L(\bar{k})$, we set $\phi(r) = F_1 - F_2$.

Proof. For every connected directed graph Γ , there is an $\text{Aut } \Gamma$ -equivariant isomorphism $H_1(\Gamma, \mathbb{Z}) \cong \ker \phi$, where

$$\phi: \mathbb{Z}^{\text{edges}} \longrightarrow \mathbb{Z}^{\text{vertices}}, \quad \phi(v_1 \rightarrow v_2) = v_1 - v_2$$

is as above. Now take $\Gamma = \Gamma(\mathcal{C}_k)$, obtained from Theorem 3.13; note that it is connected. The choice of the ordering F_1, F_2 of v -faces whose boundary is a given v -edge L (already implicit in the definition of ϕ in the statement) makes Γ directed. Now modify Γ , by viewing every chain of \mathbb{P}^1 s as one component (i.e. edge), and contracting all outer chains. This produces a new graph Γ' with $H_1(\Gamma, \mathbb{Z}) = H_1(\Gamma', \mathbb{Z})$.

¹⁴if there are no contractible faces, and all chains in 3.13(2) are chosen to be minimal (cf. 3.15), then \mathcal{C}_k is already minimal and there is nothing to contract



We find a planar realisation of Γ' with 2-dimensional faces in 1-1 correspondence with elements of $|\Delta(\mathbb{Z})_{\mathbb{Z}}^L|$; these give a \mathbb{Z} -basis of homology. The statement on the G_k -action follows from Theorem 3.13. \square

Next, recall that $\text{Pic}^0 \mathcal{C}_k$ is an algebraic group over k , an extension of an abelian variety (the *abelian part*) by a linear group; the latter is an extension of a unipotent group by a torus (the *toric part*) [BLR, §9.2].

Corollary 6.2 (Special fibre). *Suppose C/K is Δ_v -regular, and k is perfect.*

- (1) *The abelian part of $\text{Pic}^0 \mathcal{C}_k$ is isomorphic to $\bigoplus_{v\text{-faces } F} \text{Pic}^0 \bar{X}_F$, and has dimension $|\Delta(\mathbb{Z})_{\mathbb{Z}}^F|$.*
- (2) *The toric part of $\text{Pic}^0 \mathcal{C}_k$ has dimension $|\Delta(\mathbb{Z})_{\mathbb{Z}}^L|$ and character group $H_1(\Gamma(\mathcal{C}_k), \mathbb{Z})$ given by Theorem 6.1.*
- (3) *Let $l \neq \text{char } k$, and denote $H^i(-) = H^i(-_{\bar{k}}, \mathbb{Q}_l)$. We have*

$$H^1(\mathcal{C}_k) \ominus \mathbf{1} \cong \bigoplus_{v\text{-faces } F} (H^1(\bar{X}_F) \ominus \mathbf{1}) \oplus \bigoplus_{\substack{\text{inner} \\ v\text{-edges } L}} H^0(\bar{X}_L).$$

Proof. (1), (2) Combine Theorems 3.13, 6.1 with [BLR, 9.2.5/8]. (3) The image of ϕ of Theorem 6.1 is the degree 0 part of $\mathbb{Z}[v\text{-faces}]$, as $\Gamma = \Gamma(\mathcal{C}_k)$ is connected. Tensoring with \mathbb{Q}_l , we find that the toric part is

$$H^1(\Gamma, \mathbb{Z}) \otimes \mathbb{Q}_l \cong H_1(\Gamma, \mathbb{Z}) \otimes \mathbb{Q}_l \cong \bigoplus_L H^0(\bar{X}_L) \ominus \mathbb{Z}[v\text{-faces}] \oplus \mathbf{1}.$$

Moving $\mathbb{Z}[v\text{-faces}]$ into the abelian part gives the claim. \square

Remark 6.3. When K is local and C/K is semistable, this also determines

- the *conductor exponent* $\mathfrak{f}_{C/K} = |\Delta(\mathbb{Z})_{\mathbb{Z}}^L|$ (the toric dimension), and
- the *local root number* $w_{C/K} = \text{multiplicity of } \mathbf{1} \text{ in } H_1(\Gamma, \mathbb{C})$, see e.g. [D2r, Prop. 3.23].

Theorem 6.4 (Inertia action). *Suppose k is perfect, C/K is a Δ_v -regular curve, and $l \neq p$. For $P \in \Delta(\mathbb{Z})_{\mathbb{Z}_p}$ define a tame character*

$$\chi_P : I_K \longrightarrow \{\text{roots of unity}\}, \quad \sigma \longmapsto \sigma(\pi^{v(P)}) / \pi^{v(P)}.$$

Let $V_{\text{tame}}^{\text{ab}}$, $V_{\text{tame}}^{\text{toric}}$ be the unique continuous l -adic representations of I_K that decompose over $\bar{\mathbb{Q}}_l$ as

$$V_{\text{tame}}^{\text{ab}} \cong_{\bar{\mathbb{Q}}_l} \bigoplus_{P \in \Delta(\mathbb{Z})_{\mathbb{Z}_p}^F} (\chi_P \oplus \chi_P^{-1}), \quad V_{\text{tame}}^{\text{toric}} \cong_{\bar{\mathbb{Q}}_l} \bigoplus_{P \in \Delta(\mathbb{Z})_{\mathbb{Z}_p}^L} \chi_P.$$

Then there are isomorphisms of I_K -modules,

$$H_{\text{ét}}^1(C_{\bar{K}}, \mathbb{Q}_l)^{I_{\text{wild}}} \cong (V_l \text{Jac } C)^{I_{\text{wild}}} \cong V_{\text{tame}}^{\text{ab}} \oplus V_{\text{tame}}^{\text{toric}} \otimes \text{Sp}_2.$$

In particular, $\text{Jac } C$ is wildly ramified $\Leftrightarrow \Delta(\mathbb{Z})_{\mathbb{Z}_p} \subsetneq \Delta(\mathbb{Z})$.

Proof. Write $J = \text{Jac } C$. The statement only concerns inertia, so we may assume $k = \bar{k}$ and that K is complete. The first isomorphism is standard.

Now $V_{\text{tame}}^{\text{ab}}, V_{\text{tame}}^{\text{toric}}$ are I_K/I_{wild} -representations that factor through some cyclic group C_d , $p \nmid d$. They are rational (realisable over \mathbb{Q}) by Lemma A.1 (2), and uniquely determined by the dimensions of their invariants under subgroups: for $H < C_d$ of index e ,

$$(6.5) \quad \dim(V_{\text{tame}}^{\text{ab}})^H = 2|\Delta(\mathbb{Z})_{\frac{1}{e}\mathbb{Z}}^F|, \quad \dim(V_{\text{tame}}^{\text{toric}})^H = |\Delta(\mathbb{Z})_{\frac{1}{e}\mathbb{Z}}^L|.$$

On the other side, there is a canonical filtration of I_K -modules [SGA7_I, IX]

$$(V_l J)^{\text{toric}} \subseteq (V_l J)^{I_K} \subseteq V_l J$$

that gives a decomposition

$$V_l J \cong (V_l J)^{\text{ab}} \oplus (V_l J)^{\text{toric}} \otimes \text{Sp}_2.$$

Decompose the wild inertia invariants accordingly,

$$(V_l J)^{I_{\text{wild}}} \cong (V_l J)_{\text{tame}}^{\text{ab}} \oplus (V_l J)_{\text{tame}}^{\text{toric}} \otimes \text{Sp}_2.$$

The I_K/I_{wild} -representations $(V_l J)_{\text{tame}}^{\text{ab}}, (V_l J)_{\text{tame}}^{\text{toric}}$ also factor through finite (cyclic) groups by the Grothendieck monodromy theorem. They are rational, as their characters are independent of l , and a representation of cyclic group with a rational character is rational. We need to show that they are $\cong V_{\text{tame}}^{\text{ab}}$ and $\cong V_{\text{tame}}^{\text{toric}}$. First note that the inertia invariants match:

$$\dim(V_l J)^{I_K} = \dim V_l \text{Pic}^0 \mathcal{C}_k = 2|\Delta(\mathbb{Z})_{\mathbb{Z}}^F| + |\Delta(\mathbb{Z})_{\mathbb{Z}}^L|,$$

with the first equality by Theorem B.1, and the second by Corollary 6.2 plus the fact that the unipotent part of $\text{Pic}^0 \mathcal{C}_k$ has no l -power torsion. By Deligne's argument on Frobenius weights [SGA7_I, I, §6], 'the abelian and toric parts do not mix', so $(V_l J)_{\text{tame}}^{\text{ab}}$ reduces to the abelian part, and $(V_l J)_{\text{tame}}^{\text{toric}}$ to the toric part of $\text{Pic}^0 \mathcal{C}_k$. In other words,

$$\dim(V_l J)_{\text{tame}}^{\text{ab}} = 2|\Delta(\mathbb{Z})_{\mathbb{Z}}^F|, \quad \dim(V_l J)_{\text{tame}}^{\text{toric}} = |\Delta(\mathbb{Z})_{\mathbb{Z}}^L|.$$

The same argument applies over the (unique) tame extension of K of degree e for every $e \geq 1$, because C stays Δ_v -regular in finite tame extensions by Remark 3.11, and the function v on Δ gets multiplied by e . This proves the claimed isomorphism. Finally,

$$\dim V_l J = 2 \dim J = 2g(C) = 2|\Delta(\mathbb{Z})| = 2|\Delta(\mathbb{Z})_{\mathbb{Z}_p}^F| + 2|\Delta(\mathbb{Z})_{\mathbb{Z}_p}^L| + 2|\Delta(\mathbb{Z})_{\mathbb{Z}_p}^L|.$$

Comparing the dimensions with (6.5) for $H = \{1\}$, we see that the last term is non-zero if and only if $V_l J \neq (V_l J)^{I_{\text{wild}}}$. \square

Example 6.6. The curve from Example 3.3 has wild reduction when $\text{char } k$ is 2 or 3, and tame otherwise. In the tame case, C becomes semistable over any extension K'/K of ramification degree 6. Let $\chi: I_K/I_{\text{wild}} \rightarrow \bar{\mathbb{Q}}_l^\times$ be an order 6 character. By Theorem 6.4, as an I_K -module ($\text{char } k \neq 2, 3$),

$$H_{\text{ét}}^1(C_{\bar{K}}, \mathbb{Q}_l) \cong (\chi \oplus \chi^{-1} \oplus \chi^2 \oplus \chi^{-2}) \oplus (\chi^2 \oplus \chi^{-2}) \otimes \text{Sp}_2.$$

Theorem 6.7 (Reduction). *Suppose C/K is Δ_v -regular, of genus ≥ 1 . Then*

- (1) C is Δ_v -regular over every finite tame extension K'/K .
- (2) C has good reduction $\Leftrightarrow \Delta(\mathbb{Z}) = F(\mathbb{Z})$ for some v -face F with $\delta_F = 1$.
- (3) C is semistable \Leftrightarrow every principal v -face F has $\delta_F = 1$.
- (4) C is tame \Leftrightarrow every principal v -face F has $p \nmid \delta_F$ ($p = \text{char } k$).

In this case, C is Δ_v -regular over every finite extension K'/K .

Suppose k is perfect, and let J be the Jacobian of C . Then

- (5) J has good reduction $\Leftrightarrow \Delta(\mathbb{Z}) = \Delta(\mathbb{Z})_{\mathbb{Z}}^F$.
 (6) J is semistable $\Leftrightarrow \Delta(\mathbb{Z}) = \Delta(\mathbb{Z})_{\mathbb{Z}}$.
 (7) J is tame $\Leftrightarrow \Delta(\mathbb{Z}) = \Delta(\mathbb{Z})_{\mathbb{Z}_p}$. In this case,
 J has potentially good reduction $\Leftrightarrow \Delta(\mathbb{Z})^L = \emptyset$, and
 J has potentially totally toric reduction $\Leftrightarrow \Delta(\mathbb{Z})^F = \emptyset$.

Proof. Let \mathcal{C}/O_K be the m.r.n.c. model of C/K from Theorems 3.13, 5.7, with special fibre \mathcal{C}_k .

(1), (4, second claim) This is proved in Remark 3.11.

(3, \Leftarrow) For every principal v -face F , we have $\delta_L = 1$ for all $L \subset \partial F$ (inner or outer) since $\delta_F = 1$. It follows from 3.13, 5.7 and 3.15 that \mathcal{C} is a semistable model of C .

(3, \Rightarrow) If $\delta_F > 1$ for some principal v -face, then \mathcal{C} is not semistable as it has a multiple fibre. By the uniqueness of the m.r.n.c. model, C is not semistable.

(2, \Leftarrow) All faces but F are removable and \mathcal{C}_k has one smooth component, of multiplicity 1. So C has good reduction.

(2, \Rightarrow) Conversely, the m.r.n.c. model has one component, and so (i) there is only one principal face F , (ii) $\delta_F = 1$ as C is semistable, and (iii) the dual graph Γ of \mathcal{C}_k is a tree by 6.1, so $\Delta(\mathbb{Z})^L = \emptyset$.

(6) For an abelian variety A/K semistable reduction is equivalent to inertia I_K acting unipotently on $V_l A$ [SGA7_I, IX, 3.5/3.8], [BLR, 7.4.6], so the claim follows from 6.4.

(5) Semistable reduction is good if and only if the toric part of $\text{Pic}^0 \mathcal{C}_k$ is trivial, so this follows from 6.2.

(4, first claim), (7). By (1), C/K remains Δ_v -regular in tame extensions K'/K , and the function $v : \Delta \rightarrow \mathbb{R}$ gets multiplied by $e_{K'/K}$. So (3), (6) and 6.2(1,2) give the claims. \square

Remark 6.8. Let C be a Δ_v -regular curve, and suppose k is perfect. Then 6.7 and 5.5 show that the following conditions are equivalent:

- (1a) $\text{Jac } C$ is semistable (equivalently, I_K acts unipotently on $V_l \text{Jac } C$),
 (1b) C is either semistable of genus ≥ 1 , or a genus 1 curve with $C(K) \neq \emptyset$ and special fibre consisting of one non-reduced genus 1 component,

and the following are equivalent as well:

- (2a) C has tame reduction,
 (2b) Principal components of the special fibre of the some r.n.c. model have multiplicities prime to p .

Thus, 6.4 extends theorems of Saito [Sai, 3, 3.8] and Halle [Hal, 7.1] for Δ_v -regular curves, describing explicitly the wild inertia invariants.

7. ÉTALE COHOMOLOGY OVER LOCAL FIELDS

Suppose K is complete and k is finite, so K is a local field. We write $I_{\text{wild}} \triangleleft I_K \triangleleft G_K$ for the wild inertia and inertia. Recall that a choice of a *Frobenius element* $\Phi \in G_K$ (acting as $x \mapsto x^{|k|}$ on \bar{k}) identifies $G_K = I_K \rtimes \hat{\mathbb{Z}}$, and the *Weil group* $W_K = I_K \rtimes \mathbb{Z}$, with $\hat{\mathbb{Z}}, \mathbb{Z}$ generated by Φ . Fix a prime $l \neq \text{char } k$; for a scheme Y/F ($F \in \{k, K\}$) write $H^i(Y) = H_{\text{ét}}^i(Y_{\bar{F}}, \mathbb{Q}_l)$.

We note that 6.2 gives the *local polynomial* $\det(1 - \Phi^{-1}T \mid H^1(C)^{I_K})$,

$$(7.1) \quad L(T) = \det(1 - \Phi^{-1}T \mid H_1(\Gamma(\mathcal{C}_k), \mathbb{Z})) \prod_{v\text{-faces } F} \det(1 - \Phi^{-1}T \mid \text{Pic}^0 \bar{X}_F).$$

The first factor has roots of absolute value 1. The second is a product of ζ -functions of \bar{X}_F with trivial factors $(1-T)(1-|k|T)$ removed, and has roots of absolute value $|k|^{-1/2}$.

Theorem 7.2 (Tame decomposition). *Let C/K be a Δ_v -regular curve. For v -edges/faces λ of Δ , define O_K -schemes $X_\lambda^{\text{tame}}: f|_{\bar{\lambda}(\mathbb{Z})_{\mathbb{Z}_p}} = 0$. Then*

$$H^1(C)^{I_{\text{wild}}} \oplus \mathbf{1} \cong \bigoplus_{\substack{F \\ v\text{-faces}}} (H^1(X_F^{\text{tame}}) \oplus \mathbf{1}) \oplus \bigoplus_{\substack{L \text{ inner} \\ v\text{-edges}}} (H^0(X_L^{\text{tame}}) \otimes \text{Sp}_2).$$

Proof. Choose an embedding $\bar{\mathbb{Q}}_l \hookrightarrow \mathbb{C}$ and view LHS and RHS as complex Weil-Deligne representations of W_K . From (7.1), 6.2 (3) and (B.2), we find that $\text{LHS}^{I_K} \cong \text{RHS}^{I_K}$ as G_k -modules. Generally, if K'/K is a finite tame extension, then $\text{LHS}^{I_{K'}} \cong \text{RHS}^{I_{K'}}$ as $G_{K'}/I_{K'}$ -modules, using the same argument as in Theorem 6.4 and that Δ_v -regularity is stable in tame extensions. As LHS and RHS are tame Frobenius-semisimple Weil representations that have the same local polynomial in all finite tame extensions of K , they are isomorphic, by [D2w, Thm 1.1]. (This is shown in [D2w] without the two words ‘tame’, but the tame case is an easy consequence.) Finally, LHS and RHS are Weil-Deligne representations of the form $\text{Weil} \oplus \text{Weil} \otimes \text{Sp}_2$, so their semisimplifications determine the representations themselves, since Frobenius has different weights on the two constituents. \square

Theorem 7.3 (One tame face). *Suppose $C: \sum a_{ij}x^i y^j = 0$ is Δ_v -regular, K is local with $k = \mathbb{F}_q$, Δ_v has only one v -face F , and $p \nmid \delta_F = e$. Then*

- (1) C has good reduction over $K(\sqrt[e]{\pi})$.
- (2) The action $G_K \curvearrowright H^1(C)$ factors through $G = \text{Gal}(K^{nr}(\sqrt[e]{\pi})/K)$. It is faithful on G , and is uniquely determined by (3) and (4) below.

For $u \in \bar{\mathbb{F}}_q^\times$ define a complete smooth curve $\bar{C}_u/\mathbb{F}_q(u)$ by an affine equation

$$C_u: \sum_{i,j} \bar{a}_{ij} u^{v(i,j)} x^i y^j = 0, \quad \text{where } \bar{a}_P = \begin{cases} 0 & \text{if } v(P) \notin \mathbb{Z} \\ \frac{a_P}{\pi^{v(P)}} \pmod{\pi} & \text{if } v(P) \in \mathbb{Z} \end{cases}$$

Write $I \cong \frac{\mathbb{Z}}{e\mathbb{Z}} < G$ for the inertia group, and $\Phi \in G$ for the Frobenius element of $K(\sqrt[e]{\pi})$.

- (3) For every 1-dimensional character χ of I of order n , the multiplicity of χ in $H^1(C)$ viewed as an I -module is

$$\frac{2}{\phi(n)} |\{P \in \Delta(\mathbb{Z}) \mid v(P) \text{ has denominator } n\}|.$$

- (4) Every conjugacy class of the Weil group $W = I \rtimes \mathbb{Z} < G$ which is not in I contains either an element $\sigma\Phi^d$ or $(\sigma\Phi^d)^{-1}$ with $d > 0$, $\sigma \in I$ and $\frac{\sigma(\sqrt[e]{\pi})}{\sqrt[e]{\pi}}$ reducing to $u \frac{q^d - 1}{e}$ for some $u \in \bar{\mathbb{F}}_{q^d}^\times$. We have

$$\text{Tr}((\sigma\Phi^d)^{-1} \mid H^1(C)) = q^d + 1 - |\bar{C}_u(\mathbb{F}_{q^d})|.$$

Proof. (1) Recall from Remark 3.11(3) that C stays Δ_v -regular in finite tame extensions of K . This applies to $K' = K(\sqrt[e]{\pi})$, where δ_F becomes 1, and then C/K' has good reduction by Theorem 3.13.

(2) Theorem 3.13 also shows that the reduction is bad over $K(\sqrt[d]{\pi})$ for $d < e$. By the Néron-Ogg-Shafarevich criterion for $\text{Jac } C$ (see [ST, §2]), G_K acts on $H^1(C)$ through G , and faithfully on I ; the determinant of this action is a power of the cyclotomic character, so it is faithful on all of G . Finally, the traces of $g \curvearrowright H^1(C)$ for $g \in G$ determine $H^1(C)$ (as it is semisimple), so it is enough to compute them for $g \in G \setminus I$ and to describe $H^1(C)|_I$.

(3) This is a special case of Theorem 6.4 and the fact that $H^1(C)|_I$ is a rational representation.

(4) Let $\Psi \in W \setminus I$, and let $U = \langle \Psi \rangle < W$ be the subgroup it generates. Then Ψ has infinite order, $U \cap I = \{1\}$, and the image of U in $W/I \cong \mathbb{Z}$ is of some finite index d . The closure of U in G_K cuts out a finite extension K''/K , of ramification degree e , residue degree d , and Ψ is a Frobenius element of K'' or its inverse. Inverting Ψ if necessary, suppose from now on that it is the former. So $\Psi = \sigma\Phi^d$, for some $\sigma \in I$.

Let $K \subset K' \subset K''$ be the degree d unramified extension of K . Pick uniformisers π'' of K'' and π' of K' with $(\pi'')^e = \pi'$. Then $(\pi'')^e = \tilde{u}\pi$ with $\tilde{u} \in K'$ a unit, whose reduction we call $u \in \mathbb{F}_{q^d}^\times$. Thus, $K'' = K(\zeta_{q^d}, \sqrt[e]{\tilde{u}\pi})$.

Now, Ψ fixes $\sqrt[e]{\tilde{u}\pi}$, and sends $\sqrt[e]{\tilde{u}}$ to $(\sqrt[e]{\tilde{u}})^{q^d}$ modulo higher order terms, whence $\frac{\Psi(\sqrt[e]{\tilde{u}\pi})}{\sqrt[e]{\tilde{u}\pi}}$ reduces to $u^{\frac{q^d-1}{e}}$. Therefore $\sigma = \Psi\Phi^{-d}$ acts on $\sqrt[e]{\tilde{u}\pi}$ in the same way (as Φ fixes it). This proves the first assertion.

Finally, as in (1), C/K'' has good reduction and the reduced curve is exactly $\bar{C}_u/\mathbb{F}_{q^d}$. As $\Psi \subset H^1(C)$ and $\text{Frob} \subset H^1(\bar{C}_u)$ act in the same way, the Lefschetz trace formula gives the second claim in (4). \square

Example 7.4. Consider the two curves

$$\begin{aligned} y^2 &= x^3 + 7^2 & \text{over } K = \mathbb{Q}_7, & \quad \pi = 7 \\ y^2 &= x^3 + 5^2 & \text{over } K = \mathbb{Q}_5, & \quad \pi = 5. \end{aligned}$$

Both are Δ_v -regular, with one v -face F that has $e = \delta_F = 3$.

Tables 3 and 4 list, for various u and d the reduced curve $\bar{C}_u/\mathbb{F}_{q^d}$, its number of points and the conclusion of Theorem 7.3 (4) in that case. These determine (with an overkill) the action of $G = \frac{\mathbb{Z}}{3\mathbb{Z}} \rtimes \hat{\mathbb{Z}}$ on $H^1(C)$, shown at the bottom of the tables. In the first case, $I = \frac{\mathbb{Z}}{3\mathbb{Z}}$ commutes with $\hat{\mathbb{Z}}$ and the action is diagonal; here $\tau \in I$ is chosen to be the generator that multiplies $\sqrt[3]{7}$ by a 3rd root of unity congruent to 2 mod 7. In the second case, $\hat{\mathbb{Z}}$ acts on I by inversion and G is non-abelian; here $\tau \in I$ is any generator (the two are conjugate in G). In both tables, Φ is the Frobenius element of $K(\sqrt[3]{\pi})$, as in the theorem.

u	d	\bar{C}_u	$ \bar{C}_u(\mathbb{F}_{q^d}) $	Tr
± 1	1	$y^2 = x^3 + 1$	12	$\text{Tr}(\Phi^{-1}) = 7 + 1 - 12 = -4$
	2		48	$\text{Tr}(\Phi^{-2}) = 49 + 1 - 48 = 2$
± 2	1	$y^2 = x^3 + 4$	3	$\text{Tr}(\Phi^{-1}\tau) = 7 + 1 - 3 = 5$
	2		39	$\text{Tr}(\Phi^{-2}\tau^2) = 49 + 1 - 39 = 11$
± 3	1	$y^2 = x^3 + 2$	9	$\text{Tr}(\Phi^{-1}\tau^2) = 7 + 1 - 9 = -1$
	2		63	$\text{Tr}(\Phi^{-2}\tau) = 49 + 1 - 63 = -13$
$\Phi^{-1} : \begin{pmatrix} -2+\sqrt{-3} & 0 \\ 0 & -2-\sqrt{-3} \end{pmatrix} \quad \tau : \begin{pmatrix} \frac{-1-\sqrt{-3}}{2} & 0 \\ 0 & \frac{-1+\sqrt{-3}}{2} \end{pmatrix} \quad \Phi\tau = \tau\Phi$				

TABLE 3. $G = \langle \tau, \Phi \rangle = C_3 \times \hat{\mathbb{Z}} \subset H^1(C)$ for $C/\mathbb{Q}_7 : y^2 = x^3 + 7^2$

Example 7.5. Take any tame ‘2-variable Eisenstein equation’ over \mathbb{Q}_p ,

$$C : y^n + x^m + p \sum_{i,j} a_{ij} x^i y^j = 0, \quad p \nmid m n a_{00},$$

with (i, j) confined to the triangle of exponents of y^n , x^m and 1. Then C is Δ_v -regular, with one v -face F . We find that

- C has good reduction over $K(\sqrt[e]{\pi})$, with $e = \delta_F = \text{lcm}(m, n)$ (tame).

u	d	\bar{C}_u	$ \bar{C}_u(\mathbb{F}_{q^d}) $	Tr
$\in \mathbb{F}_5^\times$	1	$y^2 = x^3 + u^2$	6	$\text{Tr}(\Phi^{-1}) = 5 + 1 - 6 = 0$ (= $\text{Tr}(\Phi^{-1}\tau) = \text{Tr}(\Phi^{-1}\tau^2)$, same class)
	2		36	$\text{Tr}(\Phi^{-2}) = 25 + 1 - 36 = -10$
ζ_{24}	2	$y^2 = x^3 + \zeta_{12}$	21	$\text{Tr}(\Phi^{-2}\tau^2) = 25 + 1 - 21 = 5$
ζ_{12}	2	$y^2 = x^3 + \zeta_6$	21	$\text{Tr}(\Phi^{-2}\tau) = 25 + 1 - 21 = 5$
$\Phi^{-1} : \begin{pmatrix} 0 & \sqrt{-5} \\ \sqrt{-5} & 0 \end{pmatrix}$		$\tau : \begin{pmatrix} \frac{-1-\sqrt{-3}}{2} & 0 \\ 0 & \frac{-1+\sqrt{-3}}{2} \end{pmatrix}$		$\Phi\tau\Phi^{-1} = \tau^{-1}$

TABLE 4. $G = \langle \tau, \Phi \rangle = C_3 \rtimes \hat{\mathbb{Z}} \subset H^1(C)$ for $C/\mathbb{Q}_5 : y^2 = x^3 + 5^2$

- $H^1(C)|_{I_{\mathbb{Q}_p}}$ does not depend on C . It factors through the unique C_e -quotient of $I_{\mathbb{Q}_p}$, and is the permutation representation (see Thm. 6.4)

$$H^1(C)|_{I_{\mathbb{Q}_p}} \cong (\mathbb{Q}_l[C_e/C_{e/m}] \oplus \mathbb{Q}_l) \otimes (\mathbb{Q}_l[C_e/C_{e/n}] \oplus \mathbb{Q}_l).$$

- $H^1(C)$ depends only on $a_{00} \bmod p$ (see Thm. 7.3).

Remark 7.6. Generally, by a theorem of Kisin [Kis], l -adic representations are locally constant in p -adic families of varieties. Thus, 6.4 and 7.3 may be viewed as an explicit version of this statement for tame Δ_v -regular curves.

Remark 7.7. We end by noting that 7.2, 7.3 and 7.5 all fail without the tameness assumption. There is one Kodaira type for elliptic curves, namely I_n^* , when the regular model does not determine whether the reduction is potentially good or potentially multiplicative (see [LoM]). So Theorem 6.4 (and its refinements 7.2, 7.3) do not have an obvious analogue for the description of $I_K \subset H^1(C)$ in the wild case just in terms of Δ_v .

Similarly, the condition $\text{char } k \nmid mn$ in Example 7.5 is necessary: for instance, the two elliptic curves over \mathbb{Q}

$$y^2 = x^3 + 3, \quad y^2 = x^3 + 3x + 3$$

have conductors $2^4 3^5$ and $2^4 3^3 13$, so their Galois representations at 3 are certainly not the same (while the regular model is the same, of type II.)

8. DIFFERENTIALS

Let $v : K^\times \twoheadrightarrow \mathbb{Z}$, O_K , π , k , $p = \text{char } k$ be, as usual, a discretely valued field and the associated invariants. Let $C : f = 0$ be a Δ_v -regular curve; we choose variables as in §2 so that $f'_y \neq 0$. Recall that 2.2 (4) gives a basis of differentials for C/K . Here we aim to modify it to give a basis of the global sections of the relative dualising sheaf for \mathcal{C}_Δ/O_K (Theorem 8.12).

We refer to §4 for the charts for the components \bar{X}_F , $X_L \times \Gamma_L$ of the special fibre of \mathcal{C}_Δ , defined in 3.7.

Proposition 8.1. *Let F be a v -face of Δ , and $F^* : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ be the unique affine function that equals $-\delta_{Fv}$ on $\bar{F}(\mathbb{Z})$. Then*

$$\begin{aligned} \text{ord}_{\bar{X}_F} x^i y^j &= F^*(i, j) - F^*(0, 0), \\ \text{ord}_{\bar{X}_F} x^i y^j \frac{dx}{\pi x y f'_y} &= F^*(i, j) - 1. \end{aligned} \quad \text{for } (i, j) \in \mathbb{Z}^2.$$

Proof. 8.2. *Coordinate transformation.* Write $S = \text{Spec } O_K$, and let

$$(X, Y, Z) = (x, y, \pi) \bullet M, \quad M = (m_{ij}), \quad M^{-1} = (\tilde{m}_{ij})$$

be a coordinate transformation as in 4.1-4, associated to F and some v -edge L of F . Recall from the formula (4.1) for $M^{-1} \in \text{SL}_3(\mathbb{Z})$ that

$$\tilde{m}_{13} = k_{i+1}d_i - k_id_{i+1}, \quad \tilde{m}_{23} = \delta_L d_{i+1}, \quad \tilde{m}_{33} = \delta_L d_i.$$

If $\text{char } k = p > 0$ and $p \mid \delta_L$, then $\gcd(\tilde{m}_{13}, \tilde{m}_{23}, \tilde{m}_{33}) = 1$ forces $p \nmid \tilde{m}_{13}$; if $p \nmid \delta_L$, $p \nmid d_i$, $p \mid \tilde{m}_{23}$, we choose k_{i+1} (defined modulo δ) so that $p \nmid \tilde{m}_{13}$. Thus, $(\tilde{m}_{13}, \tilde{m}_{23}) \neq (0, 0) \in K^2$. Equivalently,

$$(8.3) \quad (m_{13}, m_{23}) \neq (0, 0) \in K^2,$$

since M preserves $(0, 0, 1) \in K^3$ if and only if M^{-1} does.

Also, the first two columns of M span the plane orthogonal to $\ker F^*$, so this plane is parallel to $(\tilde{m}_{31}, \tilde{m}_{32}, \tilde{m}_{33})^\perp$. As $\delta_F = \tilde{m}_{33}$, we find

$$(8.4) \quad F^*(i, j) = \tilde{m}_{31}i + \tilde{m}_{32}j + F^*(0, 0).$$

8.5. *Transformed equation.* Write $G(X, Y, Z) = f((X, Y, Z) \bullet M^{-1})$, $G_0(X, Y, Z) = Z^{F^*(0,0)}G(X, Y, Z)$ and $H(X, Y, Z) = X^{\tilde{m}_{13}}Y^{\tilde{m}_{23}}Z^{\tilde{m}_{33}}$, a transformed version of f and π to the X, Y, Z -chart. Then

$$U: G_0 = H = 0$$

defines a complete intersection in \mathbb{A}_S^3 , and restricted to $\mathbb{A}_S^3 \setminus \{XY = 0\}$ it gives an open subset $U: Z = 0$ of $\bar{X}_F \subset C_k^{\text{red}}$. By [Liu, 6/4.14], the sheaf $\omega_{C/S}$ is generated on \bar{X}_F by $dZ / \left| \begin{smallmatrix} (G_0)'_X & (G_0)'_Y \\ H'_X & H'_Y \end{smallmatrix} \right|$ if this determinant is non-zero. We will show that it is indeed non-zero (8.6, 8.8), and that

$$Z^{F^*(0,0)} dZ / \left| \begin{smallmatrix} (G_0)'_X & (G_0)'_Y \\ H'_X & H'_Y \end{smallmatrix} \right| = \frac{\pi^{-1}XY dZ}{\tilde{m}_{23}XG'_X - \tilde{m}_{13}YG'_Y} \stackrel{(8.10)}{=} -\pi^{-1}XY Z \frac{dx}{xyf'_y}.$$

As X, Y are units on U , and Z vanishes to order 1, we get the claim for $(i, j) = (0, 0)$. As $\text{ord}_U x = \tilde{m}_{31}$, $\text{ord}_U y = \tilde{m}_{32}$, the theorem follows from (8.4).

8.6. *Relation $f' \leftrightarrow G'$.* As in Lemma A.3, from the chain rule we get

$$\begin{pmatrix} xf'_x \\ yf'_y \\ \pi f'_\pi \end{pmatrix} = \begin{pmatrix} XG'_X \\ YG'_Y \\ ZG'_Z \end{pmatrix} M^t \Rightarrow \begin{cases} xf'_x = m_{11}XG'_X + m_{12}YG'_Y + m_{13}ZG'_Z, \\ yf'_y = m_{21}XG'_X + m_{22}YG'_Y + m_{23}ZG'_Z. \end{cases}$$

Therefore,

$$(8.7) \quad \begin{aligned} m_{23}xf'_x - m_{13}yf'_y &= (m_{23}m_{11} - m_{13}m_{21})XG'_X + \\ & (m_{23}m_{12} - m_{13}m_{22})YG'_Y + 0 \cdot ZG'_Z = \tilde{m}_{23}XG'_X - \tilde{m}_{13}YG'_Y. \end{aligned}$$

8.8. *Non-vanishing.* We claim that $m_{23}xf'_x - m_{13}yf'_y \neq 0$ in $K(C)$. If not, there is a linear relation in $K[x, y]$, non-trivial by (8.3),

$$c_1xf'_x + c_2yf'_y + c_3f = 0, \quad c_i \in K.$$

(The coefficient c_3 must be constant, and not a higher degree polynomial in x and y by degree considerations.) Then $\sum a_{ij}(c_1i + c_2j + c_3)x^i y^j$ is identically zero, so all monomial exponents are in the kernel of a non-trivial linear form. But this contradicts $\text{vol}(\Delta) > 0$.

8.9. $dx/f'_y \leftrightarrow dZ/\det$. Recall that $Z = x^{m_{13}}y^{m_{23}}\pi^{m_{33}}$. From the relations $\frac{dZ}{Z} = m_{13}\frac{dx}{x} + m_{23}\frac{dy}{y}$ and $df = f'_x dx + f'_y dy = 0$ in $\Omega_{C/K}$, we get

$$\frac{dZ}{Z} = \left(\frac{m_{13}}{x} - \frac{m_{23}}{y} \frac{f'_x}{f'_y} \right) dx = \frac{dx}{xyf'_y} (m_{13}yf'_y - m_{23}xf'_x).$$

Combined with (8.7) this yields

$$(8.10) \quad \frac{dZ}{\tilde{m}_{23}XG'_X - \tilde{m}_{13}YG'_Y} = -Z \frac{dx}{xyf'_y}.$$

□

Remark 8.11. In the notation of Theorem 2.2 and Proposition 8.1, $\pi^n \omega_P$ is regular on \bar{X}_F if and only if $n \geq \lfloor v_F(P) \rfloor$, and regular non-vanishing on \bar{X}_F if and only if $v_F(P) = n + 1 - \frac{1}{\delta_F}$.

Theorem 8.12. *If C is Δ_v -regular, then the differentials*

$$\omega_{ij}^v = \pi^{\lfloor v(i,j) \rfloor} x^{i-1} y^{j-1} \frac{dx}{f'_y}, \quad (i, j) \in \Delta(\mathbb{Z})$$

form an O_K -basis of global sections of the relative dualising sheaf $\omega_{\mathcal{C}_\Delta/O_K}$.

Proof. By Proposition 8.1, each $\pi^{\lfloor v(i,j) \rfloor} \omega_{ij}$ has order ≥ 0 at every component of \mathcal{C}_k . So they are global sections, linearly independent by Baker's theorem, and it remains to prove that the lattice they span is saturated in the global sections. Suppose not. Then there is a combination of the form

$$(8.13) \quad \frac{1}{\pi} \sum_{(i,j) \in \Sigma} u_{ij} \omega_{ij}^v \quad (\emptyset \neq \Sigma \subset \Delta(\mathbb{Z}), u_{ij} \in O_K^\times),$$

which is regular along every component. Pick $P \in \Sigma$ with $\epsilon = v(P) - \lfloor v(P) \rfloor$ maximal, and a v -face F with $P \in \bar{F}(\mathbb{Z})$. Write

$$m = \text{ord}_{\bar{X}_F} \frac{1}{\pi} \pi^{\lfloor v(P) \rfloor} \omega_P; \quad m \stackrel{8.1}{=} -\epsilon \delta_F - 1 < 0.$$

As (8.13) is supposed to be regular along \bar{X}_F , and all its terms have order $\geq m$ on \bar{X}_F by maximality of ϵ and lower convexity of Δ_v , those of order m must cancel along \bar{X}_F . Let $\Sigma_m \subset \Sigma$ be their indices.

First, note that $\Sigma_m \subset \bar{F}(\mathbb{Z})$. Indeed, else there is $P' \in \Sigma_m$ on some v -face $F' \neq F$ of Δ for which $v(P') - \lfloor v(P') \rfloor > \epsilon$, contradiction. Now, from Remark 2.3 it follows that Σ_m contains all vertices of F . This allows us to replace F by any of the neighbouring v -faces, changing P if necessary but not changing m , and it also shows that $\bar{F} \cap \partial\Delta = \emptyset$ (as $\Sigma \cap \partial\Delta = \emptyset$). Proceeding inductively, we find that (8.13) cannot exist at all (cf. [O'C]). □

Example 8.14. Let $C : y^2 = \pi x^4 + \pi^3$. If $\text{char } k \neq 2$, then C is Δ_v -regular, and its Δ_v and the special fibre of the m.r.n.c. model are as follows:

$$\begin{array}{c} 0 \\ | \\ 3/2 \quad 1 \quad 1/2 \\ | \\ 3 \quad -5/2 \quad -2 \quad -3/2 \quad 1 \end{array} \quad \xrightarrow{\quad 2 \text{ g1} \quad}$$

It is a genus 1 curve with $C(K) = \emptyset$ by 3.19, so it is not an elliptic curve; it has bad reduction but its Jacobian has good reduction, by 6.7(2,5). The differential $\pi dx/y$ spans the global sections of the relative dualising sheaf $\omega_{\mathcal{C}_\Delta/O_K}$ by the above theorem, but it vanishes

along the unique (reduced) component Γ of the special fibre, by 8.1. Hence, $\omega_{\mathcal{C}_\Delta/O_K} (=O(\Gamma))$ is not generated by global sections. In contrast, $\omega_{\mathcal{C}_\Delta/O_K}$ is trivial for genus 1 curves *with a rational point*, see [Liu, 9.4, Exc. 4.16].


Example 8.15. Consider the genus 2 curve ‘188’ from [FLS3W] at $p = 2$,

$$C: y^2 = x^5 - x^4 + x^3 + x^2 - 2x + 1.$$

Letting $x \rightarrow \frac{2}{x-1}$, $y \rightarrow \frac{y+1}{(x-1)^3} + \frac{x}{x-1}$, the equation becomes

$$y^2 + 2(x^3 - 2x^2 + x + 1)y + x(2x^2 - 5x + 4)^2 = 0.$$

This is Δ_v -regular, with special fibre of the m.r.n.c. model as follows:



The differentials dx/y , xdx/y form a basis of the relative dualising sheaf.

9. EXAMPLE: ELLIPTIC CURVES

As an application, we recover Tate’s algorithm for elliptic curves, and some immediate consequences for the Galois representation and the Néron component group. All of this is well-known: see [TaA, Sil2, §IV.9]+ ϵ [D2t] for Tate’s algorithm and the valuations of the coefficients of the minimal Weierstrass model, [Kra, Roh, ST] for the Galois representation, and [Liu, §10.2, Exc. 2.2] for r.n.c. models of elliptic curves.

As always, we have $v : K^\times \rightarrow \mathbb{Z}$, O_K , π , k and $p = \text{char } k$. Recall that an elliptic curve E/K has a Weierstrass equation,

$$y^2 + a_1xy + a_3y = x^3 + a_2x^2 + a_4x + a_6, \quad a_i \in K, \Delta_E \neq 0. \quad (\mathcal{W})$$

Here Δ_E is the discriminant of E , defined via

$$b_2 = a_1^2 + 4a_2, \quad b_4 = 2a_4 + a_1a_3, \quad b_6 = a_3^2 + 4a_6, \\ b_8 = a_1^2a_6 + 4a_2a_6 - a_1a_3a_4 + a_2a_3^2 - a_4^2, \quad \Delta_E = -b_2^2b_8 - 8b_4^3 - 27b_6^2 + 9b_2b_4b_6.$$

We write f for LHS-RHS of (\mathcal{W}) , so that $f = 0$ is the equation defining E .












Theorem 9.1. *Assume¹⁵ k is perfect or $p > 3$. An elliptic curve E/K has a Δ_v -regular Weierstrass equation (\mathcal{W}) with Δ_v as in Table 5, column 2, up to removable faces. The Kodaira type of E , special fibre of the m.r.n.c. model, condition for E to have tame reduction, inertia action on $H_{\acute{e}t}^1(E/\bar{K}, \mathbb{Q}_l)$ if E is tame and k is perfect, and the Néron component group $E(K)/E_0(K)$ are as in columns 1, 3–6.*

Proof. Start with a Weierstrass equation (\mathcal{W}) . As $x = \pi^{-2}x', y = \pi^{-3}y'$ changes $a_i \mapsto \pi^i a_i$, we can assume that all $v(a_i) \geq 0$. Write $P = (1, 1) \in \mathbb{Z}^2$.

A transformation $x \mapsto x + r, y \mapsto y + sx + t$ with $r, s, t \in O_K$ keeps Δ_E unchanged, and Δ confined to the triangle $(0,0)$ - $(3,0)$ - $(0,2)$. As long as (\mathcal{W}) is not Δ_v -regular, we will see that there is such a transformation that increases $v(P)$. Note that vertices of any v -face inside the triangle span an affine lattice of index 12 in \mathbb{Z}^2 , so $12v(P) \in \mathbb{Z}$. But $v(a_i) \geq i v(P)$ for all i , so $12v(P) \leq v(\Delta_E)$ and the algorithm terminates.

There are four reasons why f could be non- Δ_v -regular:

¹⁵Otherwise there are indeed more reduction types, classified in [Szy]; the proof of 9.1 uses that a multiple root of a polynomial of degree 2 or 3 over k is always k -rational.

Type	Δ_v	Model	tame \Leftrightarrow	Inertia if tame	$[E(K):E_0(K)]$
I_0	 Ex: $y^2 = x^3 - x$ ($p \neq 2$), $y^2 + y = x^3$ ($p \neq 3$)	$\frac{1 \text{ g1}}{F}$		C_1	$\{0\}$
II	 Ex: $y^2 = x^3 - \pi$	$\frac{3 \mid 2 \mid 1 \mid 6}{F}$	$p \neq 2, 3$	C_6	$\{0\}$
III	 Ex: $y^2 = x^3 - \pi x$	$\frac{1 \mid 2 \mid 1 \mid 4}{F}$	$p \neq 2$	C_4	$\mathbb{Z}/2\mathbb{Z}$
IV	 Ex: $y^2 + \pi y = x^3$	$\frac{1 \mid 1 \mid 1 \mid 3}{F}$	$p \neq 3$	C_3	$\mathbb{Z}/3\mathbb{Z}$ if $l=2$ $\{0\}$ if $l=0$
IO^*	 Ex: $y^2 = x^3 - \pi^2 x$ ($p \neq 2$), $y^2 = x^3 - \pi^3$ ($p \neq 3$)	$\frac{1 \mid 1 \mid 1 \mid 2}{F}$	$p \neq 2$	C_2	$(\mathbb{Z}/2\mathbb{Z})^2$ if $b=3$ $\mathbb{Z}/2\mathbb{Z}$ if $b=1$ $\{0\}$ if $b=0$
IV^*	 Ex: $y^2 + \pi^2 y = x^3$	$\frac{1 \mid 2 \mid 1 \mid 2 \mid 3}{F}$	$p \neq 3$	C_3	$\mathbb{Z}/3\mathbb{Z}$ if $l=2$ $\{0\}$ if $l=0$
III^*	 Ex: $y^2 = x^3 - \pi^3 x$	$\frac{1 \mid 2 \mid 3 \mid 2 \mid 1 \mid 2 \mid 3 \mid 4}{F}$	$p \neq 2$	C_4	$\mathbb{Z}/2\mathbb{Z}$
II^*	 Ex: $y^2 = x^3 - \pi^5$	$\frac{3 \mid 2 \mid 4 \mid 3 \mid 2 \mid 1 \mid 5 \mid 6}{F}$	$p \neq 2, 3$	C_6	$\{0\}$
I_n	 Ex: $y^2 + xy = x^3 - \pi^n$	$\frac{1 \dots 1}{1 \mid 1 \mid 1}$		$\begin{pmatrix} 1 & \tau \\ 0 & 1 \end{pmatrix}$	$\mathbb{Z}/n\mathbb{Z}$ if $s=2$ $\{0\}$ if $s=0, 2 \nmid n$ $\mathbb{Z}/2\mathbb{Z}$ if $s=0, 2 \mid n$
I_{2n}^*	 Ex: $y^2 = x^3 + \pi x^2 + \pi^{n+2} x$	$\frac{1 \mid 1 \mid 2}{F_1}$ $\frac{2 \mid \dots \mid 2}{F_2}$	$p \neq 2$	$\pm \begin{pmatrix} 1 & \tau \\ 0 & 1 \end{pmatrix}$	$(\mathbb{Z}/2\mathbb{Z})^2$ if $b=3$ $\mathbb{Z}/2\mathbb{Z}$ if $b=1$
I_{2n+1}^*	 Ex: $y^2 + \pi^{n+2} y = x^3 + \pi x^2$	$\frac{1 \mid 1 \mid 2}{F_1}$ $\frac{2 \mid \dots \mid 2}{F_2}$	$p \neq 2$	$\pm \begin{pmatrix} 1 & \tau \\ 0 & 1 \end{pmatrix}$	$\mathbb{Z}/4\mathbb{Z}$ if $l=2$ $\mathbb{Z}/2\mathbb{Z}$ if $l=0$

l = number of \hat{K} -roots of $t^2 + a_3 t - a_6$ (left 1-face)
 b = number of \hat{K} -roots of $t^3 + a_2 t^2 + a_4 t + a_6$ (bottom 1-face)
 s = number of \hat{K} -roots of $t^2 + a_1 t - a_2$ (skew 1-face)
 p = residue characteristic, π uniformiser, \hat{K} completion, τ tame character of I_K

TABLE 5. Reduction types of elliptic curves (Theorem 9.1)

- (1) Δ_v has v -edge λ (left, (0,0)-(0,2)) with $\overline{f_\lambda}$ non-squarefree, and bounding a non-removable face.
- (2) Δ_v has v -edge σ (skew, (0,2)-(2,0)) with $\overline{f_\sigma}$ non-squarefree.
- (3) Δ_v has v -edge β (bottom, (0,0)-(2,0), (0,0)-(3,0) or (1,0)-(3,0)) with $\overline{f_\beta}$ non-squarefree, and bounding a non-removable face.
- (4) Δ_v has a principal v -face F , with P in its interior, and $\overline{f_F} = 0$ is a geometrically singular subscheme of \mathbb{G}_m^2 .

In all four cases, the offending 1- or 2-face has denominator 1, for otherwise the corresponding reduction is linear and cannot define a singular scheme; for the same reason, in (1) and (3) the face is assumed to be non-removable.

In case (1)-(3), the reduction $\overline{f_*}$ ($*$ $\in \{\lambda, \sigma, \beta\}$) has a unique multiple root $\alpha \in k$, and a shift $y \mapsto y + \alpha\pi^d$, $y \mapsto y + \alpha\pi^d x$ or $x \mapsto x + \alpha\pi^d$ with d defined by the slope of $*$ shifts the root to 0, breaking the 1-face $*$. Let us call this change of variables the ‘shift along $*$ ’.

We refer to cases, i.e. table rows, by Kodaira types (first column).

Step 1. (I_n) If P is a v -vertex of Δ , then $v(1, 0) > v(P)$, $v(0, 1) > v(P)$, and $x \mapsto x - \frac{a_3}{a_1}$, $y \mapsto y + \frac{a_4}{a_1}$ makes $a_3 = a_4 = 0$; as $\Delta_E \neq 0$, we have $a_6 \neq 0$. Now all v -faces are contractible, with linear reductions, and f is always Δ_v -regular. The reduction type is I_n (cf. Remark 5.4), split in this case.

Step 2. If Δ_v has the v -edge $\sigma : (0,2)-(2,0)$, there are 3 possibilities.

If $\delta_\sigma = 1$ and $\overline{f_\sigma}$ is not-squarefree, then a shift along σ increases $v(P)$, and we go back to Step 1.

(I_n) If $\delta_\sigma = 1$ and $\overline{f_\sigma}$ is squarefree, we are again in the I_n -case, possibly non-split (with picture as in the table, up to possible removable faces).

(I_n^*) Otherwise $\delta_\sigma = 2$, and $v(P) = \frac{m}{2}$, $v(2, 0) = m$ for some odd m . The only possibly singular faces are the v -edges λ and β . If one of them is, shift along it. This increases the (smallest) slope of a face left of P . Since

$$v(a_3), v(a_4) \geq t; v(a_6) \geq 2t \Rightarrow v(b_4) \geq t; v(b_6), v(b_8) \geq 2t \Rightarrow v(\Delta_E) \geq 2t,$$

this process terminates in a Δ_v -regular model, of Type I_n^* .

Step 3. Finally suppose P is neither v -vertex nor lies on a v -edge; in other words, it lies inside a unique principal v -face, say F .

If $\{\overline{f_F} = 0\} \subset \mathbb{G}_m^2$ is singular (Case (4)), it has arithmetic genus 1, implying $\delta_F = 1$, and the singular point is unique and k -rational. Use a shift in x and y to translate it to (0,0). If $v(P)$ increases or P is not interior in a v -face any more, go back to Step 1. Otherwise, the only possibly singular faces now are the v -edges λ and β . If one of them is, shift along it; this necessarily increases $v(P)$ and we go back to Step 1.

If we reached this stage, we now have a Δ_v -regular model, with one non-removable face. Depending on $v(P) = \text{integer} + 0, \frac{1}{6}, \frac{1}{4}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{5}{6}$ (every face with vertices (3,0),(0,2) and somewhere else in Δ has $\delta_F | 4$ or $| 6$), we get types $I_0, II, III, IV, I_0^*, IV^*, III^*, II^*$ after a rescaling.

Finally, the Galois action on the components of multiplicity 1 follows from Theorem 3.13(2), and the ‘tame \Leftrightarrow ’ and ‘inertia if tame’ columns in the table by Theorems 6.7(4) and 6.4. \square

Remark 9.2. One also recovers the behaviour of minimal models of elliptic curves in tame extensions, see [D2t] Thm 3 (1)-(3).

10. EXAMPLE: FERMAT CURVES

Tate’s algorithm relies on Weierstrass models having a unique singularity that can be shifted to the origin. In higher genus, there may be multiple ones, and the algorithm ‘repeatedly shift to resolve a singularity’ of 9.1 often works but only *locally*. Thus, 3.14 may be applied for several choices of generators $x, y \in K(C)$ and the resulting schemes merged to a r.n.c. model.


For example, consider a genus 2 hyperelliptic curve

$$C/K: \pi Y^2 = X^3(X - 1)^2 + \pi \quad (\text{char } k > 3).$$

The polynomial on the right has a triple root $X \equiv 0$ and a double root $X \equiv 1 \pmod{\pi}$, so the equation is a model $\mathcal{C}_{\text{sing}}/O_K$ with two singular points

$$P_1 = (X, Y, \pi) \quad \text{and} \quad P_2 = (X - 1, Y, \pi).$$

Applying Theorem 3.14 to $(x_1, y_1) = (X, Y)$ and $(x_2, y_2) = (X - 1, Y)$ yields two schemes (with special fibres left and centre), both still singular:



Shifts and glueing; red = singular point on the model

The left one successfully resolves P_1 (dashed line and above) but not P_2 (red dot) and the centre one P_2 (dashed line and above) but not P_1 (red dot). They agree on two components — thick one of multiplicity 1 coming from the thick v -edge of Δ above it, and a thick one of multiplicity 2 coming from the shaded v -face of Δ above it, with P_1, P_2 removed. Glueing them together gives a r.n.c. model, bottom right. Here, one *could* resolve both P_1 and P_2 simultaneously by a Möbius transformation that sends their x -coordinates to 0 and ∞ respectively (top right), but that is generally not possible.


As an application, we construct regular models of Fermat curves

$$x^p + y^p = 1, \quad p \geq 3$$

over \mathbb{Z}_p (and thus over its tame extensions as well, see Remark 3.11). These are known over $\mathbb{Z}_p[\mu_p]$ [McC, CM], but seemingly not over \mathbb{Z}_p . Let $x \mapsto x - y + 1$, so that the new equation is

$$(x - y + 1)^p + y^p = 1.$$

The five monomials $x^p, px, -py, -py^{p-1}, -pxy^{p-1}$ determine Δ and Δ_v , as all the others have valuations ≥ 1 and lie inside the Newton polygon.



Example: Δ_v for $(x - y + 1)^p + y^p = 1$ with $p = 7$.

The Δ_v -regularity conditions are automatic except at the leftmost v -edge L , where the reduced equation is non-linear, namely

$$\bar{f}_L = \frac{1}{p}((1 - y)^p + y^p - 1) \in \mathbb{F}_p[y].$$


As shown in [CM, Lem. 3.1] (or [McC, p. 59]), considering the derivatives

$$\bar{f}'_L = -(1 - y)^{p-1} + y^{p-1}, \quad \bar{f}''_L = (p - 1)(1 - y)^{p-2} + (p - 1)y^{p-2}$$

and noting that the roots of \bar{f}'_L are $2, 3, \dots, p - 1$, we see that all multiple roots of \bar{f}_L are double and \mathbb{F}_p -rational¹⁶. Let ρ be their number, and lift them to $r_1, \dots, r_\rho \in \mathbb{Z}$. Pick one, say r_i , move it to zero with a shift $y \mapsto y + r_i$, and apply Theorem 3.14 to the shifted equation

$$(x - y + 1 - r_i)^p + (y + r_i)^p = 1.$$

The monomials defining its Δ_v are $x^p, -py^{p-1}, -pxy^{p-1}, (r - 1)^{p-1}px, c_2y^2$, and possibly c_1y, c_0 . Here $c_0, c_1, c_2 \in \mathbb{Z}$, and $p \mid c_2, p^2 \mid c_1, p^2 \mid c_0$ by definition of a double root. The picture for $p = 7$ is below (left):



Example: resolving one of the singularities for $p = 7$

We see that Δ_v has two principal v -faces, F_1 from $-py^{p-1}, x^p, c_2y^2$ of denominator p , and F_2 from $(r - 1)^{p-1}px, x^p, c_2y^2$ of denominator $2(p - 1)$. Theorem 3.14 now produces a model \mathcal{C}_i , whose only singularities still come from the v -edge L . So we have resolved one of the singular points.

Glueing the models \mathcal{C}_i together as in the first example above, we get a m.r.n.c. model \mathcal{C}/\mathbb{Z}_p , which is also minimal regular. The special fibre \mathcal{C}_k has the following components, all \mathbb{P}^1 s meeting transversally:

- Component X_p of multiplicity p ,
- $p - 2\rho$ chains from X_p with multiplicities $p - 1, p - 2, \dots, 2, 1$,
- Components Y_1, \dots, Y_ρ of multiplicity $2p - 2$, meeting X_p ,
- A chain with multiplicities $2p - 2, 2p - 1, \dots, 2, 1$ from each Y_i ,
- A component of multiplicity $p - 1$ from each Y_i .

¹⁶they are solutions to $(1 - y)^p \equiv 1 - y^p \pmod{p^2}$, viewed as elements of $\mathbb{F}_p \setminus \{0, 1\}$.

When $p = 3$, we have $\rho = 0$ and \mathcal{C}_k is a type IV* elliptic curve, see Table 5.

The multiplicities on the chains come from the slopes. For example, to see why Y_i meets X_p transversally, consider \mathcal{C}_i and compute the slopes at the v -edge L' where F_1 meets F_2 . In the notation of 3.12, let $P = (\frac{p-1}{2}, 1)$. Then

$$s_1^{L'} = \frac{p}{2p-2}, \quad s_2^{L'} = \frac{(p+1)/2}{p}, \quad \left| \frac{p}{2p-2} \frac{(p+1)/2}{p} \right| = 1.$$

The chain between Y_i and X_p is therefore empty; it continues to be empty in tame extensions of \mathbb{Z}_p with $e|(2p-2)$.

Remark 10.1. Theorem 3.14 (and glueing as above) also gives an alternative approach to the construction of regular models for semistable hyperelliptic curves in odd residue characteristic [M2D2].

APPENDIX A. ELEMENTARY FACTS ABOUT LATTICE POLYGONS IN \mathbb{R}^2

We start with equivalent characterisations of δ_F for a v -face F , and verify a statement that reflects rationality of étale cohomology (Thm. 6.4).

Lemma A.1. *Let $F \subset \mathbb{R}^2$ be a convex lattice polygon, and $v : F \rightarrow \mathbb{R}$ an affine function, \mathbb{Z} -valued on the vertices of F .*

(1) *If F has positive volume, then the following three numbers are equal:*

- $A =$ index of the lattice $v^{-1}(\mathbb{Z})$ in \mathbb{Z}^2 .
- $B =$ common denominator of $v(P)$ for $P \in \mathbb{Z}^2$.
- $C =$ common denominator of $v(P)$ for $P \in \bar{F}(\mathbb{Z})$.

(2) *Fix $d \in \mathbb{N}_{\geq 2}$. Define two counting functions: for $n \in (\mathbb{Z}/d\mathbb{Z})^\times$ let*

$$N\left(\frac{n}{d}\right) = |\Delta(\mathbb{Z})_{\frac{n}{d}+\mathbb{Z}}|, \quad \bar{N}\left(\frac{n}{d}\right) = |\bar{\Delta}(\mathbb{Z})_{\frac{n}{d}+\mathbb{Z}}|.$$

(i) $n \mapsto N\left(\frac{n}{d}\right) + N\left(\frac{-n}{d}\right)$, $n \mapsto \bar{N}\left(\frac{n}{d}\right) + \bar{N}\left(\frac{-n}{d}\right)$ are constant on $(\mathbb{Z}/d\mathbb{Z})^\times$.

(ii) *If F is a line segment or a parallelogram, then N, \bar{N} are constant.*

Proof. (1) $A = B, C|B$: clear from definitions. $B|C$: Replace F by any lattice triangle $T \subset F$. Now reflect T , completing it to a parallelogram, and tile the plane with it.

(2i) Suppose F is a line segment. Translate an endpoints of F to $(0, 0)$, and replace $v \mapsto v - v(0, 0)$. This does not affect the statement, but now $v : V(\mathbb{Z}) \rightarrow \mathbb{Q}$ is linear, where V is the line generated by F . It descends to $v : \bar{F}(\mathbb{Z}) \setminus \{\text{endpoint}\} \rightarrow \mathbb{Q}/\mathbb{Z}$, and all reduced fractions $\frac{n}{d}$ have preimage sizes independent of n . The same argument works for a parallelogram.

(2ii) If F is a triangle, reflect it in one of its edges L , completing it to a parallelogram $R = F \cup F' \cup L$. The values of v on F' are *minus* those on F , so the claim follows from (2i) for L and for R . In general, break F into triangles and use (2i) on the edges. \square

Lemma A.2. *Let $\Delta \subset \mathbb{R}^2$ be a convex lattice polygon with positive volume, and $\mathcal{L} = \{1\text{-dim faces of } \Delta\}$. For $L \in \mathcal{L}$ let $L^* : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ be the unique affine function vanishing on L and non-negative on Δ . Then for every $(i, j) \in \mathbb{Z}^2$,*

$$\sum_{L \in \mathcal{L}} (|\bar{L}(\mathbb{Z})| - 1) (L^*(i, j) - 1) = 2|\Delta(\mathbb{Z})| - 2. \quad (*_{\Delta})$$

Proof. Induction on $|\bar{\Delta}(\mathbb{Z})|$. If $|\bar{\Delta}(\mathbb{Z})|=3$, then $\text{LHS}(*_{\Delta})=\text{RHS}(*_{\Delta})=0$. Otherwise cut Δ into two lattice polygons A, B ; say $L_0 = A \cap B$. Then $(*_A), (*_B)$ hold by induction and we claim

that adding them up gives $(*_\Delta)$ minus $2(|L_0(\mathbb{Z})| - 1)$ in both sides. For the right-hand side,

$$2|A(\mathbb{Z})| - 2 + 2|B(\mathbb{Z})| - 2 = 2(|\Delta(\mathbb{Z})| - |L_0(\mathbb{Z})|) - 4.$$

For the left-hand side, the terms of $(*_A)$, $(*_B)$ add up to those of $(*_\Delta)$ except the contribution from L_0 . But L_0^* with respect to A is minus that for B , so the two contributions also add up to -2 times $|L_0(\mathbb{Z})| - 1$. \square

Finally, the proof of Baker's theorem 2.2 used that the differentials ω_{ij} behave well under $\text{Aut } \mathbb{G}_m^2$:

Lemma A.3. *Let $C : \{f(x, y) = 0\} \subset \mathbb{G}_{m, K}^2$ be a smooth curve. Consider a transformation of $\mathbb{G}_{m, K}^2$*

$$\begin{aligned} x &= X^a Y^b & X &= x^d y^{-b} \\ y &= X^c Y^d & Y &= x^{-c} y^a \end{aligned} \quad \text{with} \quad \begin{vmatrix} a & b \\ c & d \end{vmatrix} = 1.$$

Let $F(X, Y) = f(X^a Y^b, X^c Y^d)$, and assume that f'_y and F'_Y are not identically zero. Then

$$\frac{1}{xy} \frac{dx}{f'_y} = \frac{1}{XY} \frac{dX}{F'_Y}.$$

Proof. From the chain rule for $f(x, y) = F(x^d y^{-b}, x^{-c} y^a)$, we find

$$\begin{pmatrix} x f'_x \\ y f'_y \end{pmatrix} = \begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \begin{pmatrix} X F'_X \\ Y F'_Y \end{pmatrix}. \quad (*)$$

On the other hand, using $\frac{(gh)'}{gh} = \frac{g'}{g} + \frac{h'}{h}$ and the relation $(dF =) F'_X dX + F'_Y dY = 0$ on C , we have

$$\begin{pmatrix} \frac{dx}{x} \\ \frac{dy}{y} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \frac{dX}{X} \\ \frac{dY}{Y} \end{pmatrix} = \frac{1}{XY} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} Y F'_Y \\ X F'_X \end{pmatrix} \frac{dX}{F'_Y}. \quad (**)$$

Combining $(*)$ (2nd row) and $(**)$ (first row), the formula follows. \square

APPENDIX B. INERTIA INVARIANTS ON ÉTALE COHOMOLOGY

Let O_K be a complete DVR, with field of fractions K and perfect residue field k of characteristic $p \geq 0$.

Theorem B.1. *Let \mathcal{C}/O_K be a proper regular model of a smooth projective geometrically connected curve C/K . Write*

- \mathcal{A} = Néron model of $\text{Jac } C$ over O_K ,
- $\mathcal{C}_{\bar{k}}, \mathcal{A}_{\bar{k}}$ = special fibres of \mathcal{C} and \mathcal{A} base changed to \bar{k} ,
- d = gcd(multiplicities of components of $\mathcal{C}_{\bar{k}}$),
- Φ = component group $\mathcal{A}_{\bar{k}}/\mathcal{A}_{\bar{k}}^0$.

For every prime $l \neq \text{char } k$ and $n \geq 1$, there is a natural G_k -homomorphism

$$\text{Pic}(\mathcal{C}_{\bar{k}})[l^n] \longrightarrow \text{Pic}(C_{\bar{K}})[l^n]^{I_K},$$

with kernel killed by d and cokernel killed by $d|\Phi|$. In the limit,

$$(B.2) \quad V_l(\mathcal{A}_{\bar{k}}^0) \cong V_l \text{Pic}(\mathcal{C}_{\bar{k}}) \cong V_l \text{Pic}(C_K)^{I_K}$$

as G_k -modules; when $l \nmid d|\Phi|$, the same is true with \mathbb{Z}_l -coefficients.

Proof. This is essentially well-known, and the proof follows [BW, Prop 2.6] and [Liu, §10.4.2] closely. (The claim is proved in [BW] when $d = 1$.)

Write $S_0 = \text{Spec } O_K$, $S = \text{Spec } O_{K^{\text{nr}}}$ and $X = \mathcal{C} \times_{S_0} S$. Denote by X_s and X_η the special fibre and the generic fibre of X , and $X_{\bar{\eta}} = X_\eta/K^s$.

Let $\Gamma_1, \dots, \Gamma_n$ be components of X_s , with multiplicities d_1, \dots, d_n , so that $d = \text{gcd}_i(d_i)$. Write D for the free abelian group on the D_i and D^\vee for its dual, with dual basis $\Gamma_1^*, \dots, \Gamma_n^*$. There are natural maps

$$D \xrightarrow[\mathcal{Z} \mapsto [O_X(\mathcal{Z})]]{\lambda} \text{Pic } X \xrightarrow[\mathcal{L} \mapsto \sum (\text{deg } \mathcal{L}|_{\Gamma_i}) \Gamma_i^*]{\phi} D^\vee \quad D \xrightarrow[\mathcal{Z} \mapsto \sum (Z \cdot \Gamma_i) \Gamma_i^*]{\alpha = \phi \circ \lambda} D^\vee$$

with $\ker \alpha = d^{-1} X_s \mathbb{Z}$, and a commutative diagram with exact rows

$$(B.3) \quad \begin{array}{ccccc} \ker s & \hookrightarrow & \lambda(D) & \twoheadrightarrow & \alpha(D) \\ \downarrow & & \downarrow & & \downarrow \\ \ker \phi & \hookrightarrow & \text{Pic } X & \xrightarrow{\phi} & D^\vee \\ \downarrow^s & & \downarrow & & \downarrow \\ \ker \psi & \hookrightarrow & \text{Pic } X_\eta & \xrightarrow{\psi} & \frac{D^\vee}{\alpha(D)}. \end{array}$$

Two bottom rows are as in [Liu, Proof of 10.4.17], and the top row comes from the associated kernel-cokernel exact sequence, with zero cokernels. As proved in [Liu], $\ker s = \ker(\phi : \lambda(D) \rightarrow \alpha(D))$ is killed by d because $\ker \alpha = d^{-1} X_s \mathbb{Z}$ and $X_s \in \ker \lambda$. The multiplication by l^n map applied to the left column of (B.3) gives a kernel-cokernel exact sequence

$$0 \longrightarrow \ker s[l^n] \longrightarrow \ker \phi[l^n] \longrightarrow \ker \psi[l^n] \longrightarrow \frac{\ker s}{l^n \ker s}.$$

Now $\ker \phi[l^n] \cong \text{Pic } X[l^n]$ from the middle row of (B.3), as D^\vee is torsion-free, and $\text{Pic } X[l^n] \cong \text{Pic } X_s[l^n]$ by [Liu, 10/4.14-15]. As for $\ker \psi$, from the bottom row of (B.3)

$$0 \longrightarrow \ker \psi[l^n] \longrightarrow \text{Pic } X_\eta[l^n] \longrightarrow \frac{D^\vee}{\alpha(D)}[l^n] = \Phi[l^n],$$

the last equality by [Liu, 10/4.12]. Putting everything together, we get

$$(B.4) \quad \text{Pic } X_s[l^n] \xrightarrow{\cong} \text{Pic } X[l^n] \xrightarrow{\cong} \ker \phi[l^n] \xrightarrow{(1)} \ker \psi[l^n] \xrightarrow{(2)} \text{Pic } X_\eta[l^n]$$

with kernel and cokernel of (1) killed by d and cokernel of (2) by $|\Phi|$.

Finally, $\text{Pic } X_\eta = (\text{Pic } X_{\bar{\eta}})^{I_K}$ from the exact sequence (for any variety)

$$H^1(G, K^s[X_{\bar{\eta}}]^\times) \longrightarrow \text{Pic } X_\eta \longrightarrow (\text{Pic } X_{\bar{\eta}})^{I_K} \longrightarrow H^2(G, K^s[X_{\bar{\eta}}]^\times),$$

see [CTS, 1.5.0, p.386]. As X_η is proper and geometrically integral, $K^s[X_{\bar{\eta}}]^\times = (K^s)^\times$, so the first term is 0 by Hilbert 90, and the last one is the Brauer group, also 0 for a separably closed field.

This proves the first claim, and passing to the limit and tensoring with \mathbb{Q}_l gives the other two; also, $\mathcal{A}_\eta[l^n]^{I_K} \cong \mathcal{A}_s^0[l^n]$ by [ST, Lemma 2]. \square

Acknowledgements. I would like to thank Vladimir Dokchitser, Jack Thorne, Michael Harrison, Bartosz Naskrecki, Celine Maistret, Adam Morgan, David Holmes, Qing Liu, Raymond van Bommel, Natalia Afanassieva, Bjorn Poonen, Stefan Wewers, Christopher Doris, Michael Stoll, Dino Lorenzini, Pip Goodman, Simone Muselli and the referee for their help with the paper.

REFERENCES

- [Ark] J. R. Arkininstall, Minimal requirements for Minkowski's theorem in the plane I, *Bull. Aus. Math. Soc.* 22 (1980), 259–274.
- [Bak] H. F. Baker, Examples of applications of Newton's polygon to the theory of singular points of algebraic functions, *Trans. Cam. Phil. Soc.* 15 (1893) 403–450.
- [Bat] V. Batyrev, Variations of the mixed Hodge structure of affine hypersurfaces in algebraic tori, *Duke Math. J.* 69 (1993), no. 2, 349–409.
- [Bee] P. Beelen, A generalization of Baker's theorem, *Fin. Fields and Appl.* 15 (2009), 558–568.
- [BP] P. Beelen, R. Pellikaan, The Newton polygon of plane curves with many rational points, *Des. Codes Cryptogr.* 21 (2000), no. 1–3, 41–67.
- [BKK] D. N. Bernshtein, A. G. Kushnirenko, A. G. Khovanskii, Newton polyhedra, *Uspekhi Mat. Nauk*, 31 Issue 3(189) (1976), 201–202.
- [BLR] S. Bosch, W. Lütkebohmert, M. Raynaud, *Néron Models*, *Erg. Math.*, vol. 21, Springer, Berlin, 1990.
- [BW] I. Bouw, S. Wewers, Computing L -functions and semistable reduction of superelliptic curves, *Glasgow Math. J.* 59(1) (2017), 77–108.
- [Cas] W. Castryck, Smooth curves in toric surfaces, *Hab. Dir. Rech.*, Univ. de Lille-1, 2017.
- [CDV] W. Castryck, J. Denef, F. Vercauteren, Computing zeta functions of nondegenerate curves, *Int. Math. Res. Papers* vol. 1 (2006), 1–57.
- [CV] W. Castryck, J. Voight, On nondegeneracy of curves, *Algebra & Number Theory* 3 (3), 255–281.
- [CES] B. Conrad, B. Edixhoven, W. Stein, $J_1(p)$ has connected fibers, *Doc. Math.* 8 (2003), 331–408.
- [CTS] J.-L. Colliot-Thélène, J.-J. Sansuc, La descente sur les variétés rationnelles, II, *Duke Math. J.* 54, no 2 (1987), 375–492.
- [CM] C. Curilla, J. S. Müller, The minimal regular model of a Fermat curve of odd squarefree exponent and its dualizing sheaf, *arXiv:1605.04548*.
- [DM] P. Deligne, D. Mumford, The irreducibility of the space of curves of given genus, *Publ. Math. IHÉS*, Tome 36 (1969), 75–109.
- [DL] J. Denef, F. Loeser, Weights of exponential sums, intersection cohomology, and Newton polyhedra, *Invent. Math.* 106 (1991), 275–294.
- [D²r] T. V. Dokchitser, Regulator constants and the parity conjecture, *Invent. Math.* 178, no. 1 (2009), 23–71.
- [D²h] T. V. Dokchitser, Quotients of hyperelliptic curves and étale cohomology, to appear in *Quarterly J. Math.*
- [D²t] T. V. Dokchitser, A remark on Tate's algorithm and Kodaira types, *Acta Arith.* 160 (2013), 95–100.
- [D²w] T. V. Dokchitser, Euler factors determine local Weil representations, *Crelle* 717 (2016), 35–46.
- [FLS^W] E. V. Flynn, F. Leprevost, E. Schaefer, W. Stein, M. Stoll, J. Wetherell, Empirical evidence for the Birch and Swinnerton-Dyer conjectures for modular Jacobians of genus 2 curves, *Math. Comp.* 70 (2001), 1675–1697.
- [Ful] W. Fulton, *Introduction to toric varieties*, Princeton University Press, Princeton, 1993.
- [GRS²] R. Greenberg, K. Rubin, A. Silverberg, M. Stoll, On elliptic curves with an isogeny of degree 7, *Amer. J. Math.*, 136(1) (2014), 77–109.
- [Hal] L. H. Halle, Stable reduction of curves and tame ramification, *Math. Z.* 265, issue 3 (2010), 529–550.
- [HW] G. H. Hardy, E. M. Wright, *An introduction to the theory of numbers*, 4th ed., Oxford Univ. Press, 1960.
- [Har] R. Hartshorne, *Algebraic geometry*, GTM 52, Springer-Verlag, 1977.
- [K²MS] G. Kempf, F. Knudsen, D. Mumford, B. Saint-Donat, *Toroidal embeddings I*, LNM 339, Springer, 1973.
- [Kho] A. G. Khovanskii, Newton polyhedra and toroidal varieties, *Funktsional. Anal. i Prilozhen.* 11:4 (1977), 56–64; *Funct. Anal. Appl.* 11:4 (1977), 289–296.
- [Kis] M. Kisin, Local constancy in p -adic families of Galois representations, *Math. Z.*, 230 (1999), 569–593.
- [Kra] A. Kraus, Sur le défaut de semi-stabilité des courbes elliptiques à réduction additive, *Manuscripta Math.* 69 (1990), no. 4, 353–385.
- [Koe] R. Koelman, The number of moduli of families of curves on toric surfaces, PhD thesis, KU Nijmegen, 1991.
- [Kou] A. G. Kouchnirenko, Polyèdres de Newton et nombres de Milnor, *Inv. Math.* 32 (1976), 1–31.
- [KWZ] A. Kresch, J. Wetherell, and M. Zieve, Curves of every genus with many points, I: Abelian and toric families, *J. Algebra* 250 (2002), no. 1, 353–370.
- [Liu] Q. Liu, *Algebraic geometry and arithmetic curves*, Oxford Univ. Press, 2002.
- [LiM] Q. Liu, Modèles minimaux des courbes de genre deux, *J. Reine Angew. Math.* 453 (1994), 137–164.
- [LoD] D. Lorenzini, Dual graphs of degenerating curves, *Math. Ann.* 287 (1990), 135–150.
- [LoM] D. Lorenzini, Models of curves and wild ramification, *Pure and Appl. Math. Quart.* 6 (2010), 41–82.
- [M²D²] T. V. Dokchitser, C. Maistret, A. Morgan, Arithmetic of hyperelliptic curves over local fields, in preparation.
- [McC] W. G. McCallum, The degenerate fibre of the Fermat curve, in: *Number theory related to Fermat's last theorem* (Cambridge, Mass., 1981). *Progr. Math.* 26. Birkhäuser Boston, Mass., (1982), 57–70.
- [O'C] D. J. O'Connor, Pragmatic Paradoxes, *Mind*, 57 (1948), 358–359.
- [PSS] B. Poonen, E. F. Schaefer, M. Stoll, Twists of $X(7)$ and primitive solutions to $x^2 + y^3 = z^7$, *Duke Math. J.* 137, no. 1 (2007), 103–158.
- [Rab] S. Rabinowitz, A census of convex lattice polygons with at most one interior lattice point, *Ars Comb.* 28 (1989), 83–96.
- [Roh] D. Rohrlich, Galois Theory, elliptic curves, and root numbers, *Compos. Math.* 100 (1996), 311–349.
- [Sai] T. Saito, Vanishing cycles and geometry of curves over a DVR, *Amer. J. Math.* 109(6) (1987), 1043–1085.
- [SGA₇] A. Grothendieck, Modèles de Néron et monodromie, LNM 288, Springer, 1972.
- [Sil2] J. H. Silverman, *Advanced Topics in the Arithmetic of Elliptic Curves*, GTM 151, Springer-Verlag, 1994.
- [ST] J.-P. Serre, J. Tate, Good reduction of abelian varieties, *Annals of Math.* 68 (1968), 492–517.
- [Szy] M. Szydlowski, Elliptic fibers over non-perfect residue fields, *J. Number Theory* 104, Issue 1 (2004), 75–99.
- [TaA] J. Tate, Algorithm for determining the type of a singular fiber in an elliptic pencil, in: *Modular Functions of One Variable I*, LNM 476, B. J. Birch and W. Kuyk, eds., Springer-Verlag, Berlin, 1975, 33–52.
- [TaN] J. Tate, Number theoretic background, in: *Automorphic forms, representations and L-functions, Part 2* (ed. A. Borel and W. Casselman), *Proc. Symp. in Pure Math.* 33 (AMS, Providence, RI, 1979) 3–26.
- [Te] J. Tevelev, Compactifications of subvarieties of tori, *Amer. J. Math.* 129 no 4. (2007), 1087–1104.
- [Vie] E. Viehweg, Invarianten der degenerierten Fasern in lokalen Familien von Kurven, *Crelle* 293/294 (1977) 284–308.
- [Xi] G. Xiao, On the stable reduction of pencils of curves, *Math. Z.* 203(3) (1990), 379–389.