$N^{3/4}$ LAW IN THE CUBIC LATTICE

EDOARDO MAININI, PAOLO PIOVANO, BERND SCHMIDT, AND ULISSE STEFANELLI

ABSTRACT. We investigate the Edge-Isoperimetric Problem (EIP) for sets with n elements of the cubic lattice by emphasizing its relation with the emergence of the Wulff shape in the crystallization problem. Minimizers M_n of the edge perimeter are shown to deviate from a corresponding cubic Wulff configuration with respect to their symmetric difference by at most $O(n^{3/4})$ elements. The exponent 3/4 is optimal. This extends to the cubic lattice analogous results that have already been established for the triangular, the hexagonal, and the square lattice in two space dimensions.

1. INTRODUCTION

In this contribution we consider the Edge-Isoperimetric Problem (EIP) in

$$
\mathbb{Z}^3 := \{k_1e_1 + k_2e_2 + k_3e_3 : k_i \in \mathbb{Z} \text{ for } i = 1, 2, 3\}
$$

where $e_1 := (1, 0, 0), e_2 := (0, 1, 0)$ and $e_3 := (0, 0, 1)$. For any set C_n made of n elements of \mathbb{Z}^3 , we denote by $\Theta(C_n)$ the *edge boundary* of C_n , i.e.,

$$
\Theta(C_n) := \{ (x, y) \in \mathbb{Z}^3 \times \mathbb{Z}^3 \; : \; |x - y| = 1, \, x \in C_n \text{ and } y \in \mathbb{Z}^3 \setminus C_n \} \tag{1}
$$

and we refer to its cardinality $\#\Theta(C_n)$ as the *edge perimeter* of C_n . Given the family \mathcal{C}_n of all sets $C_n \subset \mathbb{Z}^3$ with n elements, the Edge-Isoperimetric Problem over \mathcal{C}_n consists in considering the minimum problem

$$
\theta_n := \min_{C_n \in \mathcal{C}_n} \# \Theta(C_n),\tag{2}
$$

which we denote by EIP_n , and in characterizing the EIP_n solutions. The EIP is a classical combinatorial problem and a review on the results in Combinatorics can be found in [\[2,](#page-16-0) [8\]](#page-16-1). Beyond its relevance in pure combinatorics, the EIP (and corresponding problems for similar notions of perimeter) plays a decisive role in a number of applied problems, ranging from machine learning (see [\[15\]](#page-16-2) and references therein) to the Crystallization Problem (CP). We refer the reader to [\[6\]](#page-16-3) for the relation between the EIP in the triangular lattice and the CP with respect to a two-body interatomic energy characterized by the *sticky-disc* interaction potential (see [\[9,](#page-16-4) [13\]](#page-16-5) for more details).

Our main objective is to prove that any minimizer of the EIP_n –after a suitable translation– differs from a fixed cubic configuration

$$
W_n := [0, \ell_n]^3 \cap \mathbb{Z}^3 \quad \text{with} \quad \ell_n := \lfloor \sqrt[3]{n} \rfloor. \tag{3}
$$

²⁰¹⁰ Mathematics Subject Classification. 82D25.

Key words and phrases. Wulff shape, $N^{3/4}$ law, cubic lattice, fluctuations, edge perimeter. 1

(with respect to the cardinality of their symmetric difference) by at most

$$
K n^{3/4} + o(n^{3/4}) \tag{4}
$$

elements of \mathbb{Z}^3 for some universal positive constant $K > 0$ (see Theorem [1.1\)](#page-1-0), and to show that this estimate is sharp for infinitely many n . In particular, the exponent $3/4$ of the leading term cannot be lowered in general (see Theorem [1.2\)](#page-1-1). In the following we refer to the cubic configuration W_n as the Wulff shape because of the analogy to the crystallization problem.

We first show that [\(4\)](#page-1-2) is an *upper bound* for every minimizer of EIP_n .

Theorem 1.1 (Upper Bound). There exists a constant $K_1 > 0$ independent of n such that

$$
\min_{a \in \mathbb{Z}^3} \#(M_n \triangle (a + W_n)) \le K_1 n^{3/4} + o(n^{3/4})
$$
\n(5)

for every $n \in \mathbb{N}$ and every minimizer M_n of the EIP_n .

Our second result shows that the exponent $3/4$ in (4) is optimal.

Theorem 1.2 (Lower Bound). There exists a sequence of minimizers M_{n_i} with a diverging number $n_i \in \mathbb{N}$ of particles such that

$$
\min_{a \in \mathbb{Z}^3} \#(M_{n_i} \triangle (a + W_{n_i})) \ge K_2 n_i^{3/4} + o(n_i^{3/4})
$$
\n(6)

for some constant $K_2 > 0$ (not depending on n_i).

We will prove Theorem [1.1](#page-1-0) in Section [3](#page-6-0) and Theorem [1.2](#page-1-1) in Section [4.](#page-12-0) By setting $K := \limsup_{n \to \infty} n^{-3/4} \max_{M_n} \min_{a \in \mathbb{Z}^3} \#(M_n \Delta(a + W_n)),$ where the maximum is taken among all configurations M_n that are EIP_n minimizers, by Theorems [1.1](#page-1-0) and [1.2](#page-1-1) we have $K \in (0, +\infty)$. We see that it is possible to choose $K_1 = K_2 = K$ in [\(5\)](#page-1-3) and [\(6\)](#page-1-4).

These results for \mathbb{Z}^3 are the first ones related to fluctuations of minimizers in three dimensions. Analogous results in two dimensions have been established in [\[6,](#page-16-3) [14\]](#page-16-6) for the triangular lattice (see also [\[1\]](#page-16-7)), in [\[11,](#page-16-8) [12\]](#page-16-9) for the square lattice, and finally in [\[5\]](#page-16-10) for the hexagonal lattice. The methods have been based on rearrangements techniques [\[14\]](#page-16-6) and on the isoperimetric characterization of the minimizers (with respect to suitable notions of perimeter P and area A of configurations) which also allows to find the optimal constants for relations of the type of (5) (see $[5, 6]$ $[5, 6]$).

2. Mathematical Setting

In this section we introduce the main definitions and notations used throughout the paper.

We first recall a useful characterization of EIP_n minimizers that we shall often exploit. The number of (unit) bonds of a configuration $C_n \in \mathcal{C}_n$ is

$$
b(C_n) := \frac{1}{2} \# \{ (x, y) \in C_n \times C_n : |x - y| = 1 \}.
$$

Then the elementary relation $\#\Theta(C_n) + 2b(C_n) = 6n$ shows that C_n is a minimizer for the EIP_n if and only if it maximizes the number of unit bonds.

We also introduce the 2-dimensional analogon of [\(2\)](#page-0-0) which we denote here as EIP_d^2 for $d \in \mathbb{N}$, i.e.,

$$
\eta_d := \min_{E_d \in \mathcal{C}_d^2} \# \Theta_2(E_d),\tag{7}
$$

where \mathcal{C}_d^2 is the family of subsets of the square lattice \mathbb{Z}^2 with d elements and

$$
\Theta_2(E_d) := \{ (x, y) \in \mathbb{Z}^2 \times \mathbb{Z}^2 \ : \ |x - y| = 1, \ x \in E_d \text{ and } y \in \mathbb{Z}^2 \setminus E_d \}. \tag{8}
$$

We recall from [\[11\]](#page-16-8) that E_d solves [\(7\)](#page-2-0) if and only if the number of unit bonds of E_d (i.e., $\frac{1}{2}$ #{(x, y) $\in E_d \times E_d$: $|x - y| = 1$ }) is equal to $\lfloor 2d - 2\sqrt{d} \rfloor$. This is equivalent to $\# \Theta_2(E_d) = 4d - 2\left[2d - 2\sqrt{d}\right]$, i.e., to

$$
\#\Theta_2(E_d) = 2\lceil 2\sqrt{d}\rceil. \tag{9}
$$

We also recall from [\[11\]](#page-16-8) that for any $d \in \mathbb{N}$ there exists a minimizer D_d of EIP_d^2 of the type

$$
D_d := R(s, s') \cup L_e \tag{10}
$$

for some $s, s' \in \mathbb{N}$ and $e \in \mathbb{N} \cup \{0\}$ such that $s' \in \{s, s+1\}, s \cdot s' + e = d$, and $e < s'$, where

$$
R(s, s') := \mathbb{Z}^2 \cap ([1, s] \times [1, s']) \tag{11}
$$

and

$$
L_e := \begin{cases} \mathbb{Z}^2 \cap ((0, e] \times \{s' + 1\}) & \text{if } s' = s, \\ \mathbb{Z}^2 \cap (\{s + 1\} \times (0, e]) & \text{if } s' = s + 1. \end{cases}
$$
(12)

Notice that if $e = 0$, then $L_e = \emptyset$. We refer to these 2-dimensional minimizers as daisies (as already done in [\[11\]](#page-16-8)) and to the integers $s-1$ (resp. $s'-1$) as the minimal (resp. maximal) side length of the rectangle (11) of the daisy.

Let us also introduce the notion of *minimal rectangle* associated to a 2-dimensional configuration C_n in \mathbb{Z}^3 .

Definition 2.1. Given a configuration

$$
C_n \subset \{k_1e_1 + k_2e_2 + ze_3 : k_i \in \mathbb{Z} \text{ for } i = 1, 2\}
$$

for some $z \in \mathbb{Z}$, we denote by $R(C_n)$ the closure of the minimal rectangle containing C_n , i.e., the minimal rectangle R with respect to set inclusion in

$$
\mathbb{R}_z^2 := \{(z_1, z_2, z_3) \in \mathbb{R}^3 : z_3 = z\}
$$

with sides parallel to e_i for $i = 1, 2$, and that satisfies $C_n \subset R$.

Moving ahead to 3-dimensional configurations in \mathbb{Z}^3 we introduce here a discrete rearrangement procedure, which we call *cuboidification*. Notice that the cuboidification is the 3-dimensional analogue of the 2-dimensional rearrangement introduced in [\[11\]](#page-16-8) and denoted rectangularization (see also [\[4,](#page-16-11) [7,](#page-16-12) [8\]](#page-16-1)), even though here we define such rearrangement only for EIP_n minimizers and not for a general configuration. To this end, let us introduce for every $z \in \mathbb{Z}$ the notion of z-levels of a configuration $C_n \subset \mathbb{Z}^3$ in the direction $i = 1, 2, 3$, i.e., the 2-dimensional configurations defined by

$$
C_n(z, \cdot, \cdot) := C_n \cap \{ze_1 + k_2e_2 + k_3e_3 : k_i \in \mathbb{Z} \text{ for } i = 2, 3\},
$$

\n
$$
C_n(\cdot, z, \cdot) := C_n \cap \{k_1e_1 + ze_2 + k_3e_3 : k_i \in \mathbb{Z} \text{ for } i = 1, 3\},
$$
 and
\n
$$
C_n(\cdot, \cdot, z) := C_n \cap \{k_1e_1 + k_2e_2 + ze_3 : k_i \in \mathbb{Z} \text{ for } i = 1, 2\},
$$

respectively. In the following, we also denote by $Q(a_1, a_2, a_3)$ for some $a_i \in \mathbb{N}$ the closed cuboid

$$
Q(a_1, a_2, a_3) := [1, a_1] \times [1, a_2] \times [1, a_3].
$$

Furthermore, we define the 3-vacancies of a configuration $C_n \subset \mathbb{Z}^3$ as the elements of $\mathbb{Z}^3 \setminus C_n$ that would activate three bonds if added to C_n , i.e., those elements of $\mathbb{Z}^3 \setminus C_n$ which have a distance 1 to exactly three different elements of C_n .

Definition 2.2. We define the *cuboidification* $\mathcal{Q}(M_n)$ (in the direction e_3) of a minimizer M_n of the EIP_n as the configuration resulting from rearranging the particles of M_n according to the following three steps.

(i) For every $z \in \mathbb{Z}$, let $d_z := \#M_n(\cdot, \cdot, z)$ and consider the 2-dimensional daisy D_{d_z} (which has been defined in [\(10\)](#page-2-2)), order the elements of the family $(D_{d_z})_{d_z\neq0}$ decreasingly with respect to their cardinality, say $(D^{(k)})_{k=1,\dots,f}$ with $f := \#\{z \in \mathbb{Z} : d_z \neq 0\}$, and consider the configuration M'_n characterized by

$$
M'_n(\cdot,\cdot,k) = D^{(k)} + k\mathbf{e}_3
$$

for $k = \{1, \ldots, f\}$ and $M'_n(\cdot, \cdot, k) = \emptyset$ if $k \notin \{1, \ldots, f\}$, see Figure [1.](#page-3-0)

By [\(10\)](#page-2-2) there exist $s, s' \in \mathbb{N}$ and $e \in \mathbb{N} \cup \{0\}$ with $s \cdot s' + e = d, s' \in \{s, s+1\},\$ and $e < s'$, such that $D^{(1)} = R(s, s') \cup L_e$ for $R(s, s')$ and L_e defined as in (11) and (12) , respectively.

It is clear that M'_n is still an EIP_n minimizer. Also, if $f \leq 2$ the cuboidification algorithm ends here. Otherwise, we proceed to the next steps.

FIGURE 1. Configuration M'_n . A caveat: in favor of illustrative clarity, proportions in this and the following figures do not correspond to the actual ones of a ground state.

(ii) Consecutively move the elements from $M'_n(\cdot, \cdot, f)$ with at most 3 bonds (there is always at least one of them) to fill the 3-vacancies in $M'_n \setminus M'_n(\cdot, \cdot, f)$. This allows to obtain a configuration M''_n whose levels $M''_n(\cdot, \cdot, k)$ for $k = 2, \ldots, f-1$ are rectangles (with possibly the extra segment $L_e + ke_3$), i.e.,

$$
M_n''(\cdot, \cdot, k) \setminus (L_e + ke_3) = R(a_k, b_k)
$$
\n(13)

for some $a_k, b_k \in \mathbb{N} \cup \{0\}$ which are decreasing in k, see Figure [2.](#page-4-0) We also notice that the $(z = 1)$ -level remains unchanged, i.e.,

$$
M''_n(\cdot,\cdot,1) = M'_n(\cdot,\cdot,1) = D^{(1)} + e_3,
$$

and we assume, without loss of generality, that $M''_n(\cdot, \cdot, f)$ is a daisy.

We stress that three-vacancies, if any, are filled one by one, first at the level $z = 2$, and then at the levels $z = 3, \ldots, f - 1$ (in this order), in such a way that all the z-levels up to $f - 1$ are still daisies. Moreover, each of these daisies is either coinciding with $M''_n(\cdot, \cdot, 1)$, or it is a rectangle: that is, if [\(13\)](#page-3-1) holds and $a_k < s$ or $b_k < s'$, then actually $M''_n(a_k, b_k) = R(a_k, b_k)$. M''_n is also an EIP_n minimizer.

FIGURE 2. Configuration M''_n .

- (iii) We now construct a configuration M''_n by iteratively performing the following procedure P_k for $k = 2, ..., f - 1$. The procedure P_k consists of performing the following two substeps:
	- 1) If $a_k = s$ we directly pass to substep 2). If instead $a_k < s$, then we move an entire external edge from the f-level (an f-level edge smaller or equal to b_k exists by Step (ii)), attach it at the k-level so that each of its atoms is bonded both to an atom that was already at the k-level and to one atom at the $(k-1)$ -level, and, if any 3-vacancies at the k-level appeared, then we repeat Step (ii) in order to fill them. This can be performed so that the k-level of the obtained configuration is $R(a_k+1, b_k)$. By iterating this Substep $a = s - a_k$ times, the k-level of the resulting configuration is $R(s, b_k)$.
	- 2) If $b_k = s'$, then the procedure \mathcal{P}_k is finished. If instead $b_k < s'$, then we move an entire external edge from the f -level, attach it at the k -level so that each of its atoms is bonded both to an atom that was already at the k-level and to one atom at the $(k-1)$ -level, and possibly remove any 3vacancies by repeating Step (ii). We then iterate this Substep $b = s' - b_k$ times, so that the k-level of the final configuration is $R(s, s')$.

For each $k = 2, \ldots, f - 1$, the output of this step is a k-level of the form $R(s, s')$, plus possibly an extra-line $L_e + k\mathbf{e}_3$ (which can be there only if $a_k = s$ and $b_k = s'$, that is, only if the procedure \mathcal{P}_k is empty), see Figure [3.](#page-5-0)

We notice that, if we denote by $\mathcal{P}_k(M''_n)$ the configuration obtained by iteratively performing P_i for $i = 2, \ldots, k$, we have that

$$
\mathcal{P}_k(M''_n)(\cdot,\cdot,i) \setminus (L_e + ie_3) = R(s,s')
$$

FIGURE 3. Configuration M_n''' .

for every
$$
i = 2, ..., k
$$
, and that

$$
M_n''' := \mathcal{P}_{f-1}(M_n'') = (\mathbb{Z}^3 \cap Q(s, s', f-1)) \cup F_1 \cup F_2
$$

where $F_1 := M_n'''(\cdot, \cdot, f)$ is rearranged as a daisy and

$$
F_2 := \begin{cases} M_n'''(\cdot, s + 1, \cdot) & \text{if } s' = s, \\ M_n'''(s + 1, \cdot, \cdot) & \text{if } s' = s + 1. \end{cases}
$$

We notice that $F_2 \setminus F_1$ is a rectangle $R(e, s'')$ for some $e \in \{0, \ldots, s'-1\}$, $s'' \in \{1, \ldots f-1\}$ and that, if $e = 0$, then $F_2 = \emptyset$. Without loss of generality we assume that $F_2 = M_n'''(s+1,\cdot,\cdot)$, as we can move the whole F_2 (hence also the extra-line of the daisy F_1 , if such line is contained in F_2) on that side of $Q(s, s', f)$ since $s' \geq s$. Similarly, also in case $F_1 \cap F_2 = \emptyset$ moving a line of atoms in F_1 yields F_1 which forms a square $\mathbb{Z}^3 \cap (R(a_f, b_f) \times \{f\})$ on the f-level with an extra line of atoms in $\{a_f + 1\} \times (0, e_f] \times \{f\}.$

The configuration $\mathcal{Q}(M_n) := M_n^{\prime\prime\prime}$ is still an EIP_n minimizer and it is the output of the cuboidification.

Remark 2.3. We stress that the recursive application of steps (ii) and (iii) in the above definition can never exhaust the upper face $M'_n(\cdot, \cdot, f)$ nor break its minimality for the two-dimensional EIP before a configuration of the form of M''_n is created, otherwise M_n would not be a minimizer of the EIP_n .

We conclude this section with two more definitions.

Definition 2.4. We say that a configuration C_n is *quasicubic*, or a *quasicube*, if there exist $s, s', s_3 \in \mathbb{N}$ with $s' \in \{s, s + 1\}$ such that (up to translation, relabeling, and reorienting the coordinate axes)

$$
C_n = (\mathbb{Z}^3 \cap Q(s, s', s_3 - 1)) \cup F_{d_1}^1 \cup F_{d_2}^2
$$

where $F_{d_i}^i$, $i = 1, 2$, are configurations with cardinality $d_i := \#F_{d_i}^i$ such that

$$
F_{d_1}^1 \subset \mathbb{Z}^3 \cap ([1, s+1] \times [1, s'] \times \{s_3\})
$$

and

$$
F_{d_2}^2 \subset \mathbb{Z}^3 \cap (\{s+1\} \times [1, s'-1] \times [1, s_3]).
$$

We observe that by Definition [2.2](#page-3-2) the cuboidification $\mathcal{Q}(M_n)$ of an EIP_n minimizer M_n is a quasicube with $s_3 = #\{z : M_n(\cdot, \cdot, z) \neq \emptyset\}$ and $s - 1$ the smallest side length of the rectangle $[1, s] \times [1, s']$ of the daisy with $\max_z \#M_n(\cdot, \cdot, z)$ elements.

Finally, we define the minimal cuboid of a configuration $C_n \subset \mathbb{Z}^3$.

Definition 2.5. Given a configuration $C_n \subset \mathbb{Z}^3$ we denote by $Q(C_n)$ the closure of the minimal rectangular cuboid containing C_n , i.e., the smallest rectangular cuboid Q with respect to set inclusion in \mathbb{R}^3 that has sides parallel to e_i for $i = 1, 2, 3$, and such that $C_n \subset Q$.

3. Upper bound: Proof of Theorem [1.1](#page-1-0)

We exploit the cuboidification algorithm from Section [2](#page-1-5) to obtain the proof of our first main result.

Proof of Theorem [1.1.](#page-1-0) Fix $n \in \mathbb{N}$ and let M_n be a minimizer of EIP_n. In the following, without loss of generality (up to a translation and rotation of the coordinate system), we assume that

$$
Q(M_n) = Q(\ell_1 + 1, \ell_2 + 1, \ell_3 + 1) \tag{14}
$$

for some $\ell_1, \ell_2, \ell_3 \in \mathbb{N} \cup \{0\}$ with

$$
0 \le \ell_1 \le \ell_2 \le \ell_3,\tag{15}
$$

where $Q(M_n)$ is the minimal cuboid of M_n (see Definition [2.5\)](#page-6-1) and ℓ_i , $i = 1, 2, 3$ its side lengths. Notice also that $Q(W_n) = Q(\ell_n + 1, \ell_n + 1, \ell_n + 1) - (1, 1, 1)$ for ℓ_n defined in (3) .

We now claim that there exists a constant $K > 0$ (which does not depend on n and M_n) such that

$$
\max_{i=1,2,3} |\ell_i - \ell_n| \le Kn^{1/12} + o(n^{1/12}).\tag{16}
$$

Once this is proved, Theorem [1.1](#page-1-0) follows since $\ell_n = n^{1/3} + O(1)$ by [\(3\)](#page-0-1) and hence each external face of $Q(\ell_1 + 1, \ell_2 + 1, \ell_3 + 1)$ intersected with \mathbb{Z}^3 has cardinality $n^{2/3} + o(n^{2/3})$. Thus we obtain

$$
\min_{a \in \mathbb{Z}^3} \#(Q(M_n) \triangle (a + W_n)) \le 3Kn^{3/4} + o(n^{3/4}).
$$

Since by [\(16\)](#page-6-2) moreover

$$
#(Q(M_n) \setminus M_n) = (\ell_1 + 1)(\ell_2 + 1)(\ell_3 + 1) - n
$$

\n
$$
\leq (n^{1/3} + Kn^{1/12} + O(1))^3 - n = 3Kn^{3/4} + o(n^{3/4}),
$$

[\(5\)](#page-1-3) follows.

In order to prove [\(16\)](#page-6-2) we proceed in 5 steps.

Step 1. In this step we show that by rearranging the elements of M_n we can construct another minimizer \overline{M}_n of the EIP_n that is quasicubic, i.e., there exists $\ell, \ell', \ell_3 \in \mathbb{N} \cup \{0\}$ with $0 \leq \ell \leq \ell_3$, $\ell' \in \{\ell, \ell+1\}$, and configurations $F_{d_1}^1$ and $F_{d_2}^2$ as in Definition [2.4](#page-5-1) such that

$$
\overline{M}_n = \left(\mathbb{Z}^3 \cap Q(\ell+1, \ell'+1, \ell_3)\right) \cup F_{d_1}^1 \cup F_{d_2}^2.
$$
\n
$$
(17)
$$

This assertion follows by choosing $\overline{M}_n = \mathcal{Q}(M_n)$ and by observing that [\(17\)](#page-6-3) is satisfied with ℓ being the smallest side length of the rectangle of the daisy D_m (see [\(10\)](#page-2-2)) where m is the maximal cardinality of the z-levels $M_n(\cdot, \cdot, z)$ of M_n for $z = 1, \ldots, \ell_3 + 1$, i.e.,

$$
m := \max_{z=1,\ldots,\ell_3+1} \#M_n(\cdot,\cdot,z).
$$

We notice that

$$
\ell_3 \ge \ell. \tag{18}
$$

Indeed, by [\(14\)](#page-6-4) we have $m \leq (\ell_1 + 1)(\ell_2 + 1)$. On the other hand, by definition of daisy, since ℓ is the minimal side length of the rectangle of the daisy D_m , it clearly satisfies $\ell + 1 \leq \sqrt{m}$. Therefore

$$
\ell + 1 \le \sqrt{(\ell_1 + 1)(\ell_2 + 1)},\tag{19}
$$

and with (15) we obtain (18) .

Step 2. We now further rearrange \overline{M}_n to "get rid of" the face $F_{d_2}^2$ and obtain a new EIP_n minimizer which we denote \overline{M}_n . To this end, recall from Definition [2.4](#page-5-1) and Definition [2.2](#page-3-2) that

$$
F_{d_1}^1 = \mathbb{Z}^3 \cap \left(([1, a_f] \times [1, b_f] \times \{\ell_3 + 1\}) \cup (\{a_f + 1\} \times (0, e_f] \times \{\ell_3 + 1\}) \right)
$$

for some $a_f \in \{1, ..., \ell+1\}$, $b_f \in \{a_f, a_f+1\}$, $e_f \in \{0, ..., b_f-1\}$ and that $F_{d_2}^2 \setminus F_{d_1}^1$ is a rectangle $R(e, s'')$ with $e \in \{0, 1, ..., \ell'\}$ and $s'' \in \{1, ..., \ell_3\}$. If $e = 0$ we set $\overline{M}_n := \overline{M}_n^1$, where

$$
\overline{M}_n^1 := \overline{M}_n = (\mathbb{Z}^3 \cap Q(\ell+1, \ell'+1, \ell_3)) \cup F^1 \tag{20}
$$

for $F^1 := F^1_{d_1} = \overline{M}_n(\cdot, \cdot, \ell_3 + 1).$

For $e > 0$ and $F_{d_1}^1 \cap F_{d_2}^2 = \emptyset$ we define \overline{M}_n by distinguishing 3 cases: $a_f \ge e \vee s''$, $a_f < e$ and $a_f < s''$. Later we will see how to treat the case $F_{d_1}^1 \cap F_{d_2}^2 \neq \emptyset$

1. If $a_f \ge e \vee s''$, then we move $F_{d_2}^2$ on top of $F_{d_1}^1$, and we consider the cuboidification of such configuration, which has the form

$$
\overline{M}_n^2 := (\mathbb{Z}^3 \cap Q(\ell+1, \ell'+1, \ell_3+1)) \cup F^2
$$
\n(21)

for some $F^2 := \overline{M}_n^2$ $n(n, \cdot, \ell_3 + 2)$ (which we considered rearranged as a daisy). We set $\overline{M}_n := \overline{M}_n^2$.

2. Let $a_f < e$. We can assume without loss of generality that $s'' = \ell_3$. In fact, if $s'' \nless \ell_3$, then we perform for $j = 1, \ldots, \ell_3 - s''$ the following transformation \mathcal{T}_j^1 : Move an edge with length less than e from $F_{d_1}^1$ onto $F_{d_2}^2$, so that $F_{d_2}^2$ becomes a rectangle $R(e, s'' + j)$ after removing (as done in Step (ii) of Definition [2.2\)](#page-3-2) any 3-vacancy which might have been created. The obtained configuration is

$$
M_n^b := (\mathbb{Z}^3 \cap Q(\ell + 1, \ell' + 1, \ell_3)) \cup F_1^b \cup F_2^b
$$

where $F_1^b := M_n^b(\cdot, \cdot, \ell_3 + 1)$ and F_2^b is the rectangle $R(e, \ell_3)$. We then iterate the following transformation \mathcal{T}_p^2 : for every element $p \in B_e$ where

$$
B_e := \mathbb{Z}^3 \cap (\{\ell + 2\} \times (e, \ell' + 1] \times \{1\})
$$
 (22)

remove an edge of F_1^b , attach it to F_2^b by first rotating it in order to make it parallel to e_3 and then translating it in such a way that one of its endpoints coincides with p , and then remove all 3-vacancies possibly created as in Step (ii) of Definition [2.2,](#page-3-2) so that F_2^b becomes $R(e+1, \ell_3)$. We notice that we can perform \mathcal{T}_p^2 since $a_f < e \leq \ell' \leq \ell + 1 \leq \ell_3 + 1$ by [\(18\)](#page-7-0), thus $a_f \leq \ell_3$. The configuration obtained after performing \mathcal{T}_p^2 for every $p \in B_e$ is

$$
\overline{M}_n^3 := (\mathbb{Z}^3 \cap Q(\ell+2, \ell'+1, \ell_3)) \cup F^3 \tag{23}
$$

where $F^3 := \overline{M}_n^3$ $\frac{3}{m}(\cdot, \cdot, \ell_3 + 1)$ can be rearranged as a daisy. We set $\overline{M}_n := \overline{M}_n^3$ $\frac{1}{n}$.

3. Let $a_f < s''$. We first perform the transformation \mathcal{T}_p^2 for every $p \in B_e$ to obtain the configuration

$$
M_n^c := (\mathbb{Z}^3 \cap Q(\ell + 1, \ell' + 1, \ell_3)) \cup F_1^c \cup F_2^c
$$

where $F_1^c := M_n^c(\cdot, \cdot, \ell_3 + 1)$ and F_2^c is a rectangle $R(\ell' + 1, s'')$. Then, if $s'' < \ell_3$ we perform \mathcal{T}_j^1 for every $j = 1, \ldots, \ell_3 - s''$ (without losing bonds since $a_f \leq \ell + 1 \leq \ell' + 1$ and we obtain also a configuration of the type \overline{M}_n^3 $\frac{1}{n}$. Therefore, also in this case we set $\overline{M}_n := \overline{M}_n^3$.

Let us now consider, for $e > 0$, the case $F_{d_1}^1 \cap F_{d_2}^2 \neq \emptyset$, which is possible according to Definition [2.2.](#page-3-2) The latter definition implies in such case $s'' = \ell_3$, $a_f = \ell + 1$, and

$$
F_{d_1}^1 \cap F_{d_2}^2 = \mathbb{Z}^3 \cap (\{\ell+2\} \times [1, \ldots, e_f] \times \{\ell_3 + 1\})
$$

with $1 \leq e_f \leq e$. Since $e \leq \ell' \leq \ell + 1$, we have $e_f \leq \ell + 1$, and thanks to [\(18\)](#page-7-0) we obtain $e_f \leq \ell_3 + 1$. If $e_f \leq \ell_3 = s''$, we may move the e_f points of $F_{d_1}^1 \cap F_{d_2}^2$ to $\mathbb{Z}^3 \cap (\{\ell+2\} \times \{e+1\} \times [1, e_f])$ and preserve the number of bonds (since $e_f \leq$ s''). This produces a new configuration, with the same structure of \overline{M}_n , but with $F_{d_1}^1 \cap F_{d_2}^2 = \emptyset$, and starting from such configuration we can proceed as above with the three cases. Else if $e_f = \ell_3 + 1$, then by $e_f \leq \ell + 1$ and by [\(18\)](#page-7-0) we get $\ell = \ell_3$, hence $a_f = \ell + 1 = \ell_3 + 1 = s'' + 1$. On the other hand, $e \leq \ell' \leq \ell + 1 = a_f \leq b_f$. Therefore it is possible to remove the entire $F_{d_2}^2$ and place it above the rectangle $[1, a_f] \times [1, b_f]$ of $F_{d_1}^1$ and conclude by arguing as in Case 1 above.

We observe that $\overline{\overline{M}}_n \in {\overline{M}}^i : i = 1, 2, 3$ where \overline{M}^i are defined for $i = 1, 2, 3$ in $(20), (21),$ $(20), (21),$ $(20), (21),$ $(20), (21),$ and $(23),$ respectively, and hence,

$$
\overline{\overline{M}}_n := (\mathbb{Z}^3 \cap Q(a, \ell' + 1, c)) \cup F_d,
$$
\n(24)

where $a \in \{\ell+1,\ell+2\}, c \in \{\ell_3,\ell_3+1\},\$ and $F_d := \overline{M}(\cdot,\cdot,c+1)$ with $d := \#F_d$. From here on, we assume that $a = \ell + 1$. The rest of the proof for the case $a = \ell + 2$ is essentially the same and we omit the details.

Step 3. In this step we show that

$$
\ell_3 - \ell = \sqrt{6\alpha^{1/4} \ell^{1/4}} + o(\ell^{1/4}),
$$

where $\alpha \in [0, 1)$ is specified later on in [\(31\)](#page-10-0) and depends on ℓ and ℓ_3 only.

Assume without loss of generality that $\ell_3 - \ell \geq 4$. Then there exists $k \in \mathbb{N}$ such that

$$
(c-1) - \ell = 3k + r \tag{25}
$$

for some $r \in \{0, 1, 2\}$. We further rearrange \overline{M}_n in a new minimizer \widetilde{M}_n . In order to define \widetilde{M}_n we consider the following subsets of $\overline{\overline{M}}_n$

$$
S_1 := \bigcup_{z=z_1+1,\dots,z_2} \overline{M}_n(\cdot,\cdot,z),
$$

\n
$$
S_2 := \bigcup_{z=z_2+1,\dots,z_3} \overline{M}_n(\cdot,\cdot,z),
$$

\n
$$
S_3 := F_d \cup \left(\bigcup_{z=z_3+1,\dots,z_4} \overline{M}_n(\cdot,\cdot,z)\right),
$$

\nand
$$
R := \bigcup_{z=\ell+2,\dots,z_1} \overline{M}_n(\cdot,\cdot,z),
$$
 (26)

where $z_1 := \ell + 1 + r$, $z_2 := \ell + 1 + r + k$, $z_3 := \ell + 1 + r + 2k$, and $z_4 := \ell + 1 + r + 3k = c$. Notice that the configuration

$$
G:=\overline{\overline{M}}_n\setminus\left[R\cup\left(\bigcup_{k=1,2,3}S_k\right)\right]
$$

is such that $G(\cdot, \cdot, z) := R(\ell + 1, \ell' + 1)$ for all $1 \le z \le \ell + 1$ (and $G(\cdot, \cdot, z) := \emptyset$ for $z > \ell + 1$.

We then define \widetilde{M}_n as the configuration resulting by performing the following transformations:

1. Move S_3 altogether in such a way that each element on $\overline{M}_n(\cdot,\cdot,z_3+1)$ loses its bond with $\overline{M}_n(\cdot, \cdot, z_3)$ and gains a bond with

$$
G \cap \{k_1e_1 + e_2 + k_3e_3 : k_i \in \mathbb{Z} \text{ for } i = 1,3\}.
$$

Note that this first rearrangement does not change the total number of bonds of the minimizer \overline{M}_n .

2. We then move S_2 altogether in such a way that each element on $\overline{M}_n(\cdot, \cdot, z_2+1)$ loses its bond with $\overline{M}_n(\cdot, \cdot, z_2)$ and gains a bond with

$$
G \cap \{e_1 + k_2e_2 + k_3e_3 : k_i \in \mathbb{Z} \text{ for } i = 2,3\}.
$$

Again, this rearrangement does not change the total number of bonds of the minimizer.

3. Observe that, after Transformation 2 and a translation of $+ke_1 + ke_2$ the resulting configuration, which we denote by T_n , is contained in the cuboid $[1, \ell+1+k] \times [0, \ell'+1+k] \times [1, \ell+1+r+k]$, (0 in the second factor is due to the new placement of F_d after moving S_3). However, T_n does not contain the points of $V := \mathbb{Z}^3 \cap (V_1 \cup V_2 \cup V_3)$, where

$$
V_1 := [k+1, \ell+1+k] \times [1, k] \times [\ell+2, \ell+r+k]
$$

\n
$$
V_2 := [1, k] \times [k+1, \ell+k+1] \times [\ell+2, \ell+r+k]
$$

\n
$$
V_3 = [1, k] \times [1, k] \times [1, \ell+1].
$$

Notice that $V_1 = V_2 = \emptyset$ if $k = 1$ and $r = 0$. We now fill-in the set V by subsequently moving edges with length ℓ from the side aligned in the direction

 e_1 of $T_n(\cdot, \cdot, z_2)$ (each contains $\ell+1$ points). We call the resulting configuration \widetilde{M}_n and we denote its (remaining) upper face $\widetilde{M}_n(\cdot, \cdot, z_2)$ by \widetilde{F}_m with

$$
m := #\widetilde{F}_m = (\ell+1)(\ell'+1) - #V
$$

= (\ell+1)(\ell'+1) - (\ell+1)[k^2 + 2k(r+k-1)]. (27)

Notice that in all the steps the total number of bonds of the configuration remains the same as in M_n , and so along these transformations the edges with length ℓ are not exhausted before V is filled (since this would contradict minimality of \overline{M}_n for the EIP_n). Hence, \widetilde{M}_n is an EIP_n minimizer as well.

By (27) we have

$$
m = (\ell + 1)(\ell' + 1) - #V = (\ell + 1)(\ell' + 1) - (\ell + 1)[k^2 + 2k(k + r - 1)]
$$

= (\ell + 1)(\ell' + 1 - (k^2 + 2k(k + r - 1))
= (\ell + 1)(\ell - 3k^2 - 2s_1k + s_2 + 1)
= \ell^2 - \ell(3k^2 + 2s_1k - s_2 - 2) - 3k^2 - 2s_1k + s_2 + 1 \t\t(28)

where $s_1 := r - 1 \in \{-1, 0, 1\}$ and $s_2 := \ell' - \ell \in \{0, 1\}.$

Furthermore, as M_n is a minimizer of the EIP_n , it is not possible to gain any bond by rearranging the elements of F_m over $M_n \setminus F_m$. Therefore, F_m is a minimizer (up to translation) of EIP_m^2 (see [\(7\)](#page-2-0)). By [\(9\)](#page-2-4),

$$
\Theta_2(\widetilde{F}_m) = 2\lceil 2\sqrt{m} \rceil. \tag{29}
$$

Since by the Transformation 3 the configuration \widetilde{F}_m is rectangular with side lengths ℓ and $\ell' - (k^2 + 2k(k + r - 1))$ its edge perimeter is simply

$$
\Theta_2(\widetilde{F}_m) = 2(\ell+1) + 2(\ell' - k^2 - 4k(k+r-1) + 1)
$$

= 4\ell - 6k^2 - 4s_1k + 2s_2 + 4. (30)

Therefore, by (28) , (29) , and (30) , we have that

$$
4\ell - 6k^2 - 4s_1k + 2s_2
$$

= $2[2\sqrt{\ell^2 - \ell(3k^2 + 2s_1k - s_2 - 2) - 3k^2 - 2s_1k + s_2 + 1}] - 4$,

which can be written as

$$
2\ell - 3k^2 - 2s_1k + s_2
$$

= $2\sqrt{\ell^2 - \ell(3k^2 + 2s_1k - s_2 - 2)} - 3k^2 - 2s_1k + s_2 + 1 - 2 + \alpha$

with

$$
\alpha := \lceil 2\sqrt{m} - 1 \rceil - (2\sqrt{m} - 1) = \lceil 2\sqrt{m} \rceil - 2\sqrt{m} \in [0, 1). \tag{31}
$$

By taking the square we obtain

$$
(2\ell - 3k^2 - 2s_1k + s_2 + 2 - \alpha)^2
$$

= $4\ell^2 - 4\ell(3k^2 + 2s_1k - s_2 - 2) - 12k^2 - 8s_1k + 4s_2 + 4$

from which it is straightforward to compute

$$
4\alpha\ell = 9k^4 + 12s_1k^3 + 2(2s_1^2 + 3\alpha - 3s_2)k^2 + 4s_1(\alpha - s_2)k + (\alpha - s_2)^2 - 4\alpha.
$$
 (32)

We now observe that (32) yields

$$
k = \frac{\sqrt{2} \,\alpha^{1/4}}{\sqrt{3}} \ell^{1/4} + \mathcal{O}(\ell^{1/4}).\tag{33}
$$

Therefore, from (25) and (33) we obtain

$$
\ell_3 - \ell = \sqrt{6\alpha^{1/4} \ell^{1/4}} + o(\ell^{1/4}).
$$
\n(34)

Step 4. In this step we show that (with ℓ_n as defined in [\(3\)](#page-0-1))

$$
\ell_n - \ell = \sqrt{\frac{2}{3}} \,\alpha^{1/4} \ell^{1/4} + \mathcal{O}(\ell^{1/4}).\tag{35}
$$

From [\(24\)](#page-8-2) we have that

$$
n = (\ell + 1) (\ell' + 1) (\ell_3 + s_3) + d = (\ell + 1) (\ell' + 1) (\ell + \ell_3 - \ell + s_3) + d
$$

= $\ell^3 + (\ell_3 - \ell) \ell^2 + O(\ell^2)$ (36)

where $s_2 := \ell' - \ell \in \{0, 1\}$ and $s_3 = c - \ell_3$, since $d = O(\ell^2)$ and since $\ell_3 - \ell = O(\ell^{1/4})$ by (34) . Then, (36) together with (3) yields

$$
\ell_n = \lfloor \sqrt[3]{n} \rfloor = \left\lfloor \ell \sqrt[3]{1 + \frac{\ell_3 - \ell}{\ell}} + O\left(\frac{1}{\ell}\right) \right\rfloor
$$

=
$$
\left\lfloor \ell \sqrt[3]{1 + \sqrt{6} \alpha^{1/4} \ell^{-3/4} + o(\ell^{-3/4})} \right\rfloor
$$

=
$$
\left\lfloor \ell \left(1 + \frac{\sqrt{6} \alpha^{1/4}}{3} \ell^{-3/4} + o(\ell^{-3/4})\right) \right\rfloor,
$$
 (37)

where in the second equality we used (34) . The assertion (35) follows now from (37) , since it implies

$$
\ell_n - \ell = \left\lfloor \sqrt{\frac{2}{3}} \, \alpha^{1/4} \ell^{1/4} + o(\ell^{1/4}) \right\rfloor. \tag{38}
$$

Step 5. In this step we conclude the proof of the estimate [\(16\)](#page-6-2).

Let us define $\varepsilon_i \in \mathbb{R}$ such that $\ell_i = \ell(1 + \varepsilon_i)$. We begin by observing that, as a consequence of Step 4. we have that

$$
\ell = n^{1/3} + o(n^{1/3}).\tag{39}
$$

Furthermore, from [\(34\)](#page-11-1) it follows that

$$
\varepsilon_3 = \frac{\ell_3 - \ell}{\ell} \le \sqrt{6} \,\alpha^{1/4} \ell^{-3/4} + \mathcal{O}(\ell^{-3/4}).\tag{40}
$$

By [\(15\)](#page-6-5) and [\(19\)](#page-7-3) we have $\ell_2 \geq \ell$. Therefore, by [\(40\)](#page-11-5) we obtain that

$$
0 \le \varepsilon_2 \le \varepsilon_3 \le \sqrt{6} \,\alpha^{1/4} \ell^{-3/4} + o(\ell^{-3/4}) \tag{41}
$$

as $\ell_2 \leq \ell_3$. If also $\ell_1 \geq \ell$, then the same reasoning yields that $0 \leq \varepsilon_1 \leq \varepsilon_3 \leq$ $\bar{6} \alpha^{1/4} \ell^{-3/4} + o(\ell^{-3/4})$. Therefore, it only remains to consider the case in which $\ell_1 < \ell$ and hence, $\varepsilon_1 < 0$. We have in such case, again by [\(15\)](#page-6-5) and [\(19\)](#page-7-3),

$$
\ell \leq \ell_1 \ell_2 \Rightarrow \ell^2 \leq \ell^2 (1 + \varepsilon_1)(1 + \varepsilon_2) \Rightarrow 0 \leq \varepsilon_1 + \varepsilon_2 + \varepsilon_1 \varepsilon_2
$$

$$
\Rightarrow -\varepsilon_1 \leq \varepsilon_2 + \varepsilon_1 \varepsilon_2
$$

so that, in particular

$$
0 \le -\varepsilon_1 \le \varepsilon_2. \tag{42}
$$

Therefore, by (40) , (41) , and (42) we conclude that

$$
|\varepsilon_i| \le \sqrt{6} \,\alpha^{1/4} \ell^{-3/4} + \mathcal{O}(\ell^{-3/4}).\tag{43}
$$

for $i = 1, 2, 3$. Finally, by Step 4. and (43) we observe that

$$
|\ell_i - \ell_n| \le |\ell_i - \ell| + |\ell - \ell_n|
$$

\n
$$
\le \ell |\varepsilon_i| + \sqrt{\frac{2}{3}} \alpha^{1/4} \ell^{1/4} + o(\ell^{1/4})
$$

\n
$$
\le \left(\sqrt{6} + \sqrt{\frac{2}{3}}\right) \alpha^{1/4} \ell^{1/4} + o(\ell^{1/4})
$$

for $i = 1, 2, 3$, which in turn by (39) yields estimate (16) with

$$
K:=\left(\sqrt{6}+\sqrt{\frac{2}{3}}\right)\alpha^{1/4}
$$

where we recall that $\alpha := \lceil 2\sqrt{m} \rceil - 2\sqrt{m} \in [0,1)$, see [\(31\)](#page-10-0), for m given by [\(27\)](#page-10-1). \Box

4. Lower bound: Proof of Theorem [1.2](#page-1-1)

We begin this section with an auxiliary lemma about solutions to the EIP in the two-dimensional square lattice. Indeed, for nonnegative integers s, p, q (with $s > p \vee q$), we consider configurations in \mathbb{Z}^2 of the form

$$
\mathcal{R}_{s,p,q} := R(s-p-1,s) \cup L_{s-q}^p,
$$

where $L_{s-q}^p := \mathbb{Z}^2 \cap (\{s-p\} \times [1, s-q])$. Note that $\#\mathcal{R}_{s,p,q} = s^2 - sp - q$.

Lemma 4.1. Let $s, p, q \in \mathbb{N} \cup \{0\}$ be such that $s \geq 1$, $p < s$, $q < s$. Then $\mathcal{R}_{s,p,q}$ is an EIP_n² minimizer (where $n = \#\mathcal{R}_{s,p,q} = s^2 - sp - q$) if and only if

$$
4(s - q) > (p + 1)^2.
$$

In particular, by choosing $p = \lfloor s^{1/2} \rfloor$ and $q = \lfloor s/4 \rfloor$, $\mathcal{R}_{s,p,q}$ is a EIP_n^2 minimizer for any $s \geq 2$.

Proof. We observe that the number of unit bonds in $\mathcal{R}_{s,p,q}$ is equal to

$$
(s-1)(s-p-1) + s(s-p-2) + 2(s-q) - 1.
$$

We use the fact that EIP_n^2 minimizers are characterized by a number of unit bonds equal to $\lfloor 2n - 2\sqrt{n} \rfloor$, as recalled in Section [2.](#page-1-5) As a consequence, $\mathcal{R}_{s,p,q}$ is an EIP_n^2 minimizer if and only if

$$
(s-1)(s-p-1)+s(s-p-2)+2(s-q)-1=\lfloor 2(s^2-sp-q)-2\sqrt{s^2-sp-q}\rfloor,
$$

which is equivalent to $\lfloor 2s - p - 2\sqrt{s^2 - sp - q} \rfloor = 0$, thus to

$$
0 \le 2s - p - 2\sqrt{s^2 - sp - q} < 1.
$$

As the first inequality is obvious, $\mathcal{R}_{s,p,q}$ is an EIP_n^2 minimizer if and only if

$$
2s - p - 1 < 2\sqrt{s^2 - sp - q},
$$

which is equivalent to $4(s - q) > (p + 1)^2$, as desired.

By choosing $p = \lfloor s^{1/2} \rfloor$ and $q = \lfloor s/4 \rfloor$, the latter is reduced to

$$
4s > 1 + 2\lfloor\sqrt{s}\rfloor + 4\lfloor s/4\rfloor + \lfloor\sqrt{s}\rfloor^2,
$$

which is implied by $2s > 1 + 2\sqrt{s}$ that is clearly true for $s \geq 2$.

A straightforward consequence of Lemma [4.1](#page-12-3) is the sharpness of the $N^{3/4}$ law in the two-dimensional square lattice, see [\[11\]](#page-16-8). Indeed, we might consider the sequence

$$
d_s := s^2 - s \lfloor s^{1/2} \rfloor - \lfloor s/4 \rfloor, \qquad s = 2, 3, \dots \tag{44}
$$

It is easy to check that d_s is a strictly increasing sequence. We have $d_s = \#\mathcal{R}_{s,p,q}$ with $p = \lfloor s^{1/2} \rfloor$ and $q = \lfloor s/4 \rfloor$, and $\mathcal{R}_{s,p,q}$ is an $\text{EIP}_{d_s}^2$ minimizer by Lemma [4.1.](#page-12-3) On the other hand,

$$
s - \lfloor d_s^{1/2} \rfloor = \left\lceil s - s\sqrt{1 - \frac{sp + q}{s^2}} \right\rceil \ge s - s\left(1 - \frac{sp + q}{2s^2}\right) \ge \frac{\lfloor s^{1/2} \rfloor}{2},
$$

so that we may compare the two-dimensional Wulff shape $W_{d_s}^2 := \left[1, \lfloor d_s^{1/2} \rfloor \right]^2 \cap \mathbb{Z}^2$ with $\mathcal{R}_{s,p,q}$ and get

$$
\min_{a \in \mathbb{Z}^2} \# (\mathcal{R}_{s,p,q} \bigtriangleup (a + W_{d_s}^2)) \ge \frac{1}{2} \lfloor s^{1/2} \rfloor (s - \lfloor s^{1/2} \rfloor - 1),
$$

for any $s \ge 2$. As [\(44\)](#page-13-0) implies $s = d_s^{1/2} + o(d_s^{1/2})$, we find

$$
\min_{a \in \mathbb{Z}^2} \# (\mathcal{R}_{s,p,q} \bigtriangleup (a + W_{d_s}^2)) \ge \frac{1}{2} d_s^{3/4} + o(d_s^{3/4}).
$$

We now proceed with the proof of the three-dimensional counterpart of this result.

Proof of Theorem [1.2.](#page-1-1) Let us consider the strictly increasing sequence

$$
n_s := s^3 + s^2 - s\lfloor s^{1/2} \rfloor - \lfloor s/4 \rfloor, \qquad s = 2, 3, \dots.
$$

For any integer $s \geq 2$, we may consider the configuration

$$
M_{n_s} := Q(s, s, s) \cup F_{d_s} \tag{45}
$$

where we have introduced a 2-dimensional configuration $F_{d_s} := M_{n_s}(\cdot, \cdot, s + 1)$ with $d_s := \# M_n(\cdot,\cdot,s+1).$ More precisely, we define the top face F_{d_s} as

$$
F_{d_s} := (\mathbb{Z}^3 \cap ([1, r] \times [1, s] \times \{s + 1\})) \cup L^s, \qquad r := s - \lfloor s^{1/2} \rfloor - 1, \qquad (46)
$$

where

$$
L^s := \mathbb{Z}^3 \cap (\{r+1\}, [1, s-q], \{s+1\}), \qquad q := \lfloor s/4 \rfloor. \tag{47}
$$

We see that

$$
d_s = s^2 - s\lfloor s^{1/2} \rfloor - \lfloor s/4 \rfloor < s^2 \tag{48}
$$

and that

$$
n_s = s^3 + d_s \tag{49}
$$

is indeed the number of points of M_{n_s} . Moreover, the top face F_{d_s} is an $EIP_{d_s}^2$ minimizer for any $s \geq 2$ by an application of Lemma [4.1.](#page-12-3) This ensures minimality of M_{n_s} for the EIP_{ns}, for any $s \geq 2$. We stress that

$$
s - r = \lfloor s^{1/2} \rfloor + 1. \tag{50}
$$

By (48) and (49) it follows that

$$
s = n_s^{1/3} + o(n_s^{1/3}),\tag{51}
$$

and hence

$$
r = n_s^{1/3} + o(n_s^{1/3})
$$
\n(52)

by (50) . Furthermore, by (48) and (51) we have

$$
d_s = n_s^{2/3} + o(n_s^{2/3}).
$$
\n(53)

We also refine (51) by recalling (3) and by claiming that

$$
s = \ell_{n_s}.\tag{54}
$$

To prove [\(54\)](#page-14-2) we observe that $s \leq \ell_{n_s}$ easily follows from [\(49\)](#page-13-2) and that

$$
\ell_{n_s} = \lfloor \sqrt[3]{n_s - d_s + d_s} \rfloor = \left\lfloor \sqrt[3]{n_s - d_s} \sqrt[3]{1 + \frac{d_s}{n_s - d_s}} \right\rfloor
$$

$$
= \left\lfloor s \sqrt[3]{1 + \frac{d_s}{s^3}} \right\rfloor \le \left\lfloor s \left(1 + \frac{1}{3} \frac{d_s}{s^3} \right) \right\rfloor = s \tag{55}
$$

where we used (3) in the first equality, (49) in the third one, (48) in the last one. The claim is proved.

We then proceed to construct another minimizer denoted by M''_{n_s} by performing the following two consecutive transformations on M_{n_s} :

1. Define the integer

$$
h_1 = \left\lfloor \frac{1}{3} n_s^{1/12} \right\rfloor. \tag{56}
$$

We translate by e_1 the top face F_{d_s} and we move altogether the edge $({s} \times [1, s] \times {s}) \cap \mathbb{Z}^3$ to the position $({1} \times [1, s] \times {s + 1}) \cap \mathbb{Z}^3$. We repeat then this procedure recursively for each line (parallel to e_2) with s elements of M_{n_s} which is included in the set

$$
H_1 := [s - h_1, s] \times [1, s] \times [s - h_1, s].
$$

This transformation gives the configuration M'_{n_s} , see Figure [4.](#page-15-0) Note that M'_{n_s} is an EIP_{n_s} minimizer for large enough s. In fact, the total number of moved lines is $(h_1 + 1)^2$. Hence, we can translate F_{d_s} by $(h_1 + 1)^2 e_1$ without losing any bond if $r + 1 + (h_1 + 1)^2 < s - h_1$, as such condition prevents the top face to reach the points above the 'hole' H_1 by this translation. The latter inequality is equivalent, by (50) , to

$$
h_1 + (h_1 + 1)^2 < \lfloor s^{1/2} \rfloor,\tag{57}
$$

which holds true for large enough s due to (54) and since the definition of h_1 entails

$$
(h_1 + 1)^2 = \frac{1}{9} n_s^{1/6} + o(n_s^{1/6}).
$$

FIGURE 4. Configuration M'_{n_s} .

2. Thanks to the previous step, there exists $s_0 \in \mathbb{N}$ such that, for any $s \geq s_0$, M'_{n_s} is an EIP_{n_s} minimizer. In particular, s_0 can be defined as the smallest integer such that [\(57\)](#page-14-3) hold for any $s \geq s_0$. For $s \geq s_0$ we move altogether the elements in the set

$$
H_2 := [1, s - h_1 - 1] \times [1, s] \times [s - h_1, s + 1]
$$

in such a way that each element of $M'_{n_s}(\cdot, \cdot, s - h_1)$ (loses the bond with $M'_{n_s}(\cdot, \cdot, s-h_1-1)$ and) gets bonded with an element of $M'_{n}(1,\cdot, \cdot) \setminus H_2$. We denote the resulting EIP_{n_s} minimizer by M''_{n_s} .

Thanks to the two steps above, for any $s \geq s_0$ the constructed configuration M''_{n_s} is a minimizer of the EIP_{n_s} problem and moreover we notice that

$$
\mathbb{Z}^3 \cap ([-h_1, s] \times [1, s] \times [1, s - h_1 - 1]) \subset M''_{n_s},\tag{58}
$$

therefore by (3) , (54) , (56) , (58) we conclude that

$$
\min_{a \in \mathbb{Z}^3} \#(M''_{n_s} \triangle (a + W_{n_s})) \ge \min_{a \in \mathbb{Z}^3} \#(M''_{n_s} \setminus (a + W_{n_s}))
$$

\n
$$
\ge s(s - h_1 - 1)(s + h_1 + 1 - \ell_{n_s})
$$

\n
$$
= s\left(s - \left\lfloor \frac{1}{3} n_s^{1/12} \right\rfloor - 1\right) \left(\left\lfloor \frac{1}{3} n_s^{1/12} \right\rfloor + 1\right).
$$

Hence,

$$
\min_{a \in \mathbb{Z}^3} \#(M''_{n_s} \triangle (a + W_{n_s})) \ge \frac{1}{3} n_s^{3/4} + o(n_s^{3/4})
$$

follows by (51) .

Acknowledgements

P. Piovano acknowledges support from the Austrian Science Fund (FWF) project P 29681 and by the Vienna Science and Technology Fund (WWTF), the City of Vienna, and Berndorf Privatstiftung through Project MA16-005. U. Stefanelli is

partially supported by the Vienna Science and Technology Fund (WWTF) through Project MA14-009 and by the Austrian Science Fund (FWF) projects F 65, P 27052, and I 2375.

REFERENCES

- [1] Y. Au Yeung, G. Friesecke, B. Schmidt. Minimizing atomic configurations of short range pair potentials in two dimensions: crystallization in the Wulff-shape, Calc. Var. Partial Differential Equations, 44 (2012), 81–100.
- [2] S.L. Bezrukov. Edge isoperimetric problems on graphs, in: Graph theory and combinatorial biology (Balatonlelle, 1996). Bolyai Soc. Math. Stud., 7 (1999), 157–197.
- [3] X. Blanc, M. Lewin. The crystallization conjecture: a review, *EMS Surv. Math. Sci.*, 2 (2015). 255–306.
- [4] B. Bollobas and I. Leader. Edge-isoperimetric inequalities in the grid, *Combinatorica*, 11 (1991), 4:299–314.
- [5] E. Davoli, P. Piovano, U. Stefanelli. Wulff shape emergence in graphene, Math. Models Methods Appl. Sci., 26 (2016), 2277–2310.
- [6] E. Davoli, P. Piovano, U. Stefanelli. Sharp $N^{3/4}$ law for the minimizers of the edge-isoperimetric problem on the triangular lattice, J. Nonlinear Sci., 27 (2017), 627-660.
- [7] F. Harary and H. Harborth. Extremal animals, J. Comb. Inf. Syst. Sci., 1 (1976), 1-8.
- [8] L.H. Harper, Global methods for combinatorial isoperimetric problems. Cambridge Studies in Advanced Mathematics, 90. Cambridge University Press, Cambridge, 2004.
- [9] R. Heitmann, C. Radin. Ground states for sticky disks, J. Stat. Phys., 22 (1980), 3:281–287.
- [10] E. Mainini, U. Stefanelli. Crystallization in carbon nanostructures, Comm. Math. Phys., 328 (2014), 2:545–571.
- [11] E. Mainini, P. Piovano, U. Stefanelli. Finite crystallization in the square lattice. Nonlinearity, 27 (2014), 717–737.
- [12] E. Mainini, P. Piovano, U. Stefanelli. Crystalline and isoperimetric square configurations, Proc. Appl. Math. Mech., 14 (2014), 1045–1048.
- [13] C. Radin. The ground state for soft disks, *J. Stat. Phys.*, **26** (1981), 2:365-373.
- [14] B. Schmidt. Ground states of the 2D sticky disc model: fine properties and $N^{3/4}$ law for the deviation from the asymptotic Wulff-shape. J. Stat. Phys., 153 (2013), 727-738.
- [15] N.G. Trillos, D. Slepcev. Continuum limit of total variation on point clouds, Arch. Ration. Mech. Anal., 220 (2016), 193–241.

(Edoardo Mainini) Dipartimento di Ingegneria meccanica, energetica, gestionale e dei TRASPORTI, UNIVERSITÀ DEGLI STUDI DI GENOVA, VIA ALL'OPERA PIA, 15 - 16145 GENOVA ITALY.

 $\it E\mbox{-}mail$ $address:$
 $\mbox{\tt mainini}$ @dime.unige.it

(Paolo Piovano) Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, A-1090 Vienna, Austria

 $\it E\mbox{-}mail\;address:$ paolo.piovano@univie.ac.at

URL: http://www.mat.univie.ac.at/∼piovano/

(Bernd Schmidt) INSTITUTE OF MATHEMATICS, UNIVERSITY OF AUGSBURG, UNIVERSITÄTSSTR. 14, 86159 Augsburg, Germany

E-mail address: bernd.schmidt@math.uni-augsburg.de

URL: http://www.math.uni-augsburg.de/ana/schmidt.html

(Ulisse Stefanelli) Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, A-1090 Vienna, Austria and Istituto di Matematica Applicata e Tecnologie In-FORMATICHE E. Magenes - CNR, VIA FERRATA 1, I-27100 PAVIA, ITALY

 $\it E\mbox{-}mail$ $address:$ u l isse.stefanelli@univie.ac.at

URL: http://www.mat.univie.ac.at/∼stefanelli