THE EQUIVALENCE OF SCHRÖDINGER AND HEISENBERG PICTURES IN QUANTUM CELLULAR AUTOMATA

ASIF SHAKEEL

ABSTRACT. Quantum cellular automata (QCA) are discrete models of space and time homogeneous quantum field theories (QFTs) and regarded as natural candidates for quantum simulation. Description of a QCA over the separable Hilbert space of finite, unbounded configurations (UFC Hilbert space) with unitary state evolution is the *Schrödinger template*, and over the incomplete infinite tensor product algebra (ITPA) with evolution by algebra automorphism is the *Heisenberg template*. Whether every Heisenberg template admits an equivalent Schrödinger template is a foundational question, and one that has persisted as an open problem. In the present paper we prove that for every Heisenberg template an equivalent Schrödinger template exists. We frame the question from a representation theory standpoint, using constructs and results from the representation theory of finite and countably infinite dimensional vector spaces and from category theory to answer it. With the previously known existence of a Heisenberg template for every Schrödinger template, our result establishes the equivalence of the templates.

1. INTRODUCTION

von Neumann [1] conceived of infinitely many systems interacting, replicating and producing complex patterns of behavior. He called them *cellular automata*. Likewise, quantum cellular automata (QCA) are models of infinitely many *quantum* systems interacting with each other. Proposed by Feynman [2], these are natural models of quantum systems that would simulate other quantum systems. Theory of QCA emerged from the initial work on QCA of Zeilinger [3], Durr and Santha [4], Watrous [5], and Meyer [6,7], the latter also bringing to fore an important multi-particle subclass of QCA, the quantum lattice gas automaton (QLGA). These papers establish conditions on QCA existence and unitary evolution, and reveal their dynamics and potential for simulating fundamental physics.

Axiomatic descriptions of QCA reflect the standard notions of quantum mechanics. Axioms of quantum mechanics [8,9] are stated in a Hilbert space version and a C^* algebra version. In the Hilbert space version, the state is a density operator, and the observables are bounded self-adjoint operators on the Hilbert space. In the C^* algebra version, the state is a normalized linear functional on the algebra. Within the Hilbert space formulation, of abiding interest are separable Hilbert spaces. They have played a primal role in quantum mechanics, and Gleason's crucial results [10] on quantum measurements also rest on them. von Neumann and Murray [11] develop rings of operators, von Neumann algebras, formalism of quantum mechanics. Along with C^* algebras from the ensuing work by Gelfand and Neumark [12], and Segal [13], these algebras maintain an enduring significance in algebraic quantum field theory (QFT) and infinite quantum systems.

E-mail address: asif.shakeel@gmail.com.

Schumacher and Werner [14] proposed an axiomatic model of a QCA in the operator formalism. A QCA resides on a lattice, the cells being identified with the points of the lattice. Each cell takes values in a finite dimensional C^* algebra of operators, the *cell algebra*, which is thus the endomorphisms of a finite dimensional Hilbert space. In it, the cell algebras are constituents of an *incomplete infinite tensor product algebra* [15–18] (ITPA), a *-aglebra of operators over the lattice. The system evolves at each time step through an automorphism or this *-algebra. To be a valid model of physics, such an automorphim is to be translation invariant, i.e., spatially homogeneous, acting identically on all cells. It must also obey *causality*, referring to a finite speed of information transfer, in that the image of a cell algebra under the automorphism is the identity operator on all but a finite *neighborhood* of the cell. This evolution of the operator algebra is deemed to be in the Heisenberg picture of quantum mechanics. It has led to a classification of one-dimensional QCA by an *index theory* in the work of Gross, Nesme, Vogts, and Werner [19]. We refer to this QCA model including its evolution as the *Heisenberg template* of QCA.

Arrighi, Nesme, and Werner [20] describe an axiomatic QCA in the Hilbert space formalism. Each cell takes values in the *states* of a finite dimensional Hilbert space, the *cell Hilbert space*. A typical basis element of the QCA Hilbert space consists of finitely many cells taking arbitrary values in the cell Hilbert space with a background of cells taking a designated identical quiescent value. The QCA Hilbert space is called the *Hilbert space of finite*, unbounded configurations (UFC Hilbert space). The state of the QCA evolves at each time step by a unitary, translation-invariant and *causal* operator. Causality, in this description, refers to a finite speed of information propagation at each step in the sense of evolution of the state: the state of each cell after a step of evolution depends on that of the cells in its finite neighborhood prior to the step. This QCA, because of unitary evolution, is in the Schrödinger picture of quantum mechanics. Among elegant results related to this formalism is the work of Arrighi, Nesme and Werner on localizabitity of quantum cellular automata (QCA) [21]. We refer to this QCA model including its evolution as the Schrödinger template of QCA.

Templates of QCA serve as primary points of investigations concerning properties, structures and classes of QCA that are potentially significant for quantum algorithms and simulations. Beyond classification of QCA structure, much work has gone into modeling physics with QCA, their multi-particle subclass the QLGA, and single-particle subclass, the quantum walks (QWs). Meyer [6] modeled Dirac equation in 1 + 1 dimensions with QW, and Arrgihi et al [22] in 2 + 1 dimensions. Boghosian and Taylor [23] developed simulation models for many particle Schrödinger equation based on QLGA. QWs incorporating curved spacetime appear in Arrighi et al [24], and Di Molfetta et al [25]. Bisio et al [26, 27] develop models of quantum field theory (QFT) as QCA. Arrgini et al [28] combine QCA with reversible causal graph dynamics as a framework for discrete time quantum gravity models. Meyer and Shakeel [29], seeking ways to simulate physics without particle description, combine QLGA to create QCA that no longer have a particle description at the time scale at which the dynamics are homogeneous. Schlingemann et al [30] introduced Clifford QCA (CQCA) and investigated their structure, properties and generalizations. Gütschow et al [31] studied CQCA further for their entanglement generation properties. Crossing over into several areas, CQCA are illustrative of QCA intersecting fundamental physics, quantum computation and quantum resource theory. Interestingly, CQCA, described in the operator formalism, do not readily have a particle description in the sense of, for instance, QLGA, or without particles in the sense of the QCA of [29]. Interested reader can find in [32] a comprehensive contemporary perspective and the significance of quantum cellular automata (QCA) as models of computation and simulation, their structural interpretations and analysis.

In their paper on QFT simulation, Jordan et al [33] calculate in polynomial time the scattering amplitudes of particles in ϕ^4 theory using a quantum gate array architecture. A question, raised in [29], is about the efficiency attainable in simulating physics without a particle description. QCA are contenders among those simulation models, and bringing the techniques available in operator formalism to bear on questions of this magnitude may be necessary. Physical considerations, especially implementation, would naturally demand that there be a separable Hilbert space description with a unitary state evolution. A result establishing that a Schrödinger template exists equivalent to any given Heisenberg template of a QCA links the two approaches.

It is a natural question to ask whether the templates are equivalent in the sense that each has a counterpart in the other with mirroring evolutionary dynamics. By this we mean that the measurement statistics are the same under evolution in either template, as in the Schrödinger and Heisenberg pictures of quantum mechanical systems. Given a Schrödinger template over any lattice, we can always find a Heisenberg template that has equivalent dynamics by previous results mainly from [20]. Also, given a Heisenberg template over a *finite* sized lattice, we can always find a Schrödinger template that yields equivalent dynamics, by the Skolem-Noether Theorem for finite dimensional simple algebras, the relevant version of which is cited as Scholium A.1 (Appendix A). There is no general theorem that directly gives this equivalence for countably infinite dimensional vector spaces, therefore, it was not known if, given a Heisenberg template over an infinite lattice, an equivalent Schrödinger template exists. In this paper we show that there is an equivalent Schrödinger template for every Heisenberg template. In fact, there is such a template for any UFC Hilbert space in whose bounded linear operators the ITPA of the Heisenberg template is naturally embedded.

This paper is organized as follows. Section 2 describes the Schrödinger and the Heisenberg templates of QCA. In Section 3 we first recall, without explicitly involving representation theory, the known result that given a Schrödinger template of a QCA, a dynamically equivalent Heisenberg template can be constructed. This leads to the formal definition of equivalence between the Schrödinger and Heisenberg templates of QCA in representation theoretic terms. Then we present the main result of this paper that given any Heisenberg template, an equivalent Schrödinger template exists for any UFC Hilbert space in whose bounded linear operators the ITPA of the Heisenberg template embeds. Section 4 examines the definition of a QLGA in [35], and its characterizing local condition, informed by the results of this paper. Section 5 is the conclusion.

2. QUANTUM CELLULAR AUTOMATON

We recall the definition of Quantum Cellular Automaton (QCA) as it is defined in [20] in the Hilbert space formulation, the Schrödinger template, and in the operator algebra formulation, the Heisenberg template.

The Schrödinger Template. The Schrödinger template in this paper is a slight generalization of the definition in [20,35] in terms of spatial homogeneity, i.e., translation invariance. Let us describe the Hilbert space of the QCA. Over each cell resides an identical finite dimensional complex Hilbert space, the *cell Hilbert space*, W. We designate a state $|q\rangle \in W$ as the *quiescent* state, and choose some orthonormal basis \mathcal{B} of W containing the quiescent state $|q\rangle \in \mathcal{B}$. Each basis element of the QCA Hilbert space is a tensor product, over the lattice, of basis elements from \mathcal{B} , in which except for a finitely many cells, the rest are in the quiescent state. This basis is called the *set of finite, unbounded configurations*, denoted by \mathcal{C} , defined to be

$$\mathcal{C} = \{\bigotimes_{x \in \mathbb{Z}^n} |b_x\rangle : |b_x\rangle \in \mathcal{B}, \text{ all but finite } |b_x\rangle = |q\rangle\}.$$
 (1)

Let the vector space spanned by \mathcal{C} be \mathbb{V} ,

$$\mathbb{V} = \operatorname{Span}(\mathcal{C}) . \tag{2}$$

We can obtain \mathbb{V} in other ways as well. Let an ascending chain of finite subsets be $\{D_k \subset \mathbb{Z}^n : |D_k| < \infty\}_{k \in \mathbb{N}}$, such that their union is the entire lattice, i.e.,

$$D_0 \subsetneq D_1 \subsetneq \dots \tag{3}$$

such that $\mathbb{Z}^n = \bigcup_{k \in \mathbb{N}} D_k$.

Let

$$W_{D_k} = \operatorname{Span}(\{\bigotimes_{x \in D_k} | b_x \rangle \otimes \bigotimes_{x \in \mathbb{Z}^n \setminus D_k} | q \rangle : | b_x \rangle \in \mathcal{B}\}).$$

Under the natural inclusion of W_{D_k} into $W_{D_{k+1}}$,

$$W_{D_k} \hookrightarrow W_{D_{k+1}}$$
.

 W_{D_k} can be regarded as a subspace of $W_{D_{k+1}}$. Then

$$\mathbb{V} = \cup_k W_{D_k} ,$$

Another precise construction of \mathbb{V} is as a direct limit in the manner of Guichardet [16]. The direct limit construction is in Appendix C, and is instructive for the rest of this paper. The *Hilbert space of finite, unbounded configurations* (UFC Hilbert space), denoted by \mathcal{H} , is the completion of \mathbb{V} under the inner product norm, i.e., under the pre-Hilbert structure on \mathbb{V} induced from the inner product on W.

The state of a QCA is a *density operator* on the UFC Hilbert space \mathcal{H} . A density operator is a positive trace class operator with trace 1. For the measurements that only concern a finite subset $D \subset \mathbb{Z}^n$ of cells, the density operator can be *restricted* to D. This restriction is obtained by a *partial trace* over cells (tensor factors) not in D. For that, let us write the \mathcal{H} as a tensor product of two Hilbert spaces, where completion in the inner product norm induced by those on the two tensor factors is assumed. We can write,

$$\mathbb{V} = \left(\bigotimes_{x \in D} W\right) \otimes \mathbb{V}_{\bar{D}} , \qquad (4)$$

where $\mathbb{V}_{\overline{D}}$ is the vector space complement to the tensor factors in D. Also,

$$\mathcal{H} = \left(\bigotimes_{x \in D} W\right) \otimes \mathcal{H}_{\bar{D}} ,$$

where $\mathcal{H}_{\bar{D}}$ is the Hilbert space completion of $\mathbb{V}_{\bar{D}}$. Let $\{|v_i\rangle\}$ be some orthonormal basis of $\mathcal{H}_{\bar{D}}$. The restricted density operator is calculated by first writing ρ as

$$\rho = \sum_{i,j} \rho_{i,j} \otimes |v_i\rangle \langle v_j| \,,$$

where $\rho_{i,j} \in \operatorname{End}(\bigotimes_{x \in D} W)$. Then ρ restricted to D is

$$ho|_D := \sum_i
ho_{i,i}$$
 .

Phase factors multiplying unitary evolution operators have no effect on measurements, so we are justified in the following definition.

Definition 2.1. Let R and R' be operators on a UFC Hilbert space. Then R and R' are equivalent if $R = e^{i\theta}R'$ for some $\theta \in \mathbb{R}$. The equivalence class of R is denoted by [R].

We will refer to [R] as an operator on a UFC Hilbert space when it is appropriate not to distinguish among the members of the class. When we say [R] is a unitary operator, we mean some (class) representative R of [R] is a unitary operator. By the definition of [R] then all representatives of [R] are unitary. In keeping with this equivalence, a concept that we will generalize from the model of QCA in [20,35] is that of spatial homogeneity, i.e., translation invariance, in the definition of the Schrödinger template.

The neighborhood of cell $z \in \mathbb{Z}^n$ is a finite set, denoted by $\mathcal{N}_z \subset \mathbb{Z}^n$. The state of the cell z (restriction of the state to the cell) after a step of evolution depends on the state of the neighborhood \mathcal{N}_z prior the step, in the sense of the Schrödinger template definition below. For spatially homogeneous evolution, it is given by translating a fixed finite subset $\mathcal{N} \subset \mathbb{Z}^n$ to the cell z,

$$\mathcal{N}_z = z + \mathcal{N} = \{ z + y : y \in \mathcal{N} \} .$$
(5)

Then \mathcal{N} is called the *neighborhood* for the QCA.

Definition 2.2 (Schrödinger template). Let W be a finite dimensional Hilbert space. The *Schrödinger template* of a QCA is a pair $(\mathcal{H}, [R])$, where \mathcal{H} is a UFC Hilbert space constructed from W, and [R] is a unitary operator on \mathcal{H} . [R] is required to be

(1) Translation invariant: A translation operator τ_z , for some $z \in \mathbb{Z}^n$, is a linear operator on \mathcal{H} , defined on a basis element $|c\rangle = \bigotimes_{x \in \mathbb{Z}^n} |b_x\rangle \in \mathcal{C}$ as:

$$\tau_z: \bigotimes_{x \in \mathbb{Z}^n} |b_x\rangle \mapsto \bigotimes_{x \in \mathbb{Z}^n} |b_{x+z}\rangle .$$

[R] is translation invariant if, for some representative R of [R], and for all $z \in \mathbb{Z}^n$, $\tau_z R \tau_z^{-1} = e^{i\theta_z} R$, where θ_z depends on z.

(2) Causal relative to some neighborhood \mathcal{N} : [R] is causal relative to the neighborhood \mathcal{N} if for some representative R of [R], all $z \in \mathbb{Z}^n$, and each pair ρ, ρ' of density operators on \mathcal{H} satisfying

$$\rho|_{\mathcal{N}_z} = \rho'|_{\mathcal{N}_z} ,$$

 $R\rho R^{\dagger}, R\rho' R^{\dagger}$ satisfy

$$R\rho R^{\dagger}|_{z} = R\rho' R^{\dagger}|_{z} \,.$$

The neighborhood \mathcal{N} is typically chosen to be the smallest set that satisfies the causality condition given in the Schrödinger template definition. [R] is called the *global evolution operator* of the QCA. If the state of QCA is ρ before a time step, then after the time step the state evolves to $R\rho R^{\dagger}$, where R is some representative of [R]. The Heisenberg Template. The Heisenberg template of a QCA is defined on an algebra called the *incomplete infinite tensor product algebra*. This algebra is constructed from algebras that are *local* on finite subsets of the lattice \mathbb{Z}^n , defined as follows.

Definition 2.3. Let W be a finite dimensional Hilbert space. For a finite subset $D \in \mathbb{Z}^n$, the algebra of operators local upon D is defined to be $\mathcal{A}_D = \bigotimes_{x \in D} \operatorname{End}(W) \otimes \bigotimes_{y \in \mathbb{Z}^n \setminus D} \mathbb{I}$.

Remark 2.4. We denote identity operator by \mathbb{I} for every vector space, to economize notation.

Remark 2.5. We write \mathcal{A}_z to mean $\mathcal{A}_{\{z\}}$, the *cell algebra* of operators local upon one tensor factor z.

Take an ascending chain of finite subsets as in (3). Under the obvious inclusion of \mathcal{A}_{D_k} into $\mathcal{A}_{D_{k+1}}$,

$$\mathcal{A}_{D_k} \hookrightarrow \mathcal{A}_{D_{k+1}}$$

 \mathcal{A}_{D_k} can be regarded as a subalgebra of $\mathcal{A}_{D_{k+1}}$. The incomplete infinite tensor product algebra (ITPA), which we denote by \mathcal{Z} , is

$$\mathcal{Z} = \bigcup_k \mathcal{A}_{D_k} \,. \tag{6}$$

Just as for the vector space of finite configurations (25), we can obtain \mathcal{Z} as direct limit of \mathcal{A}_{D_k} (see Guichardet [16]).

We note that \mathcal{Z} is a *-algebra since each of its constituent cell algebras, $\mathcal{A}_z = \operatorname{End}(W)$, is a *-algebra with the adjoint map on operators as its *-involution,

* :
$$\operatorname{End}(W) \to \operatorname{End}(W)$$

 $a \mapsto a^{\dagger}$.

This map extends to a *-involution on \mathcal{Z} .

Based on the constructs just introduced, we define the Heisenberg template of a QCA.

Definition 2.6 (Heisenberg template). Let W be a finite dimensional Hilbert space. The *Heisenberg template* of a QCA is the is a pair (\mathcal{Z}, γ) , where \mathcal{Z} is the ITPA (6), and γ is a *-automorphism of \mathcal{Z} . γ is required to be:

(1) translation invariant: A translation operator μ_z , for some $z \in \mathbb{Z}^n$, is a linear operator on \mathcal{Z} , defined on a basis element $b = \bigotimes_{x \in \mathbb{Z}^n} b_x \in \mathcal{Z}$ as:

$$\mu_z: \bigotimes_{x\in\mathbb{Z}^n} b_x \mapsto \bigotimes_{x\in\mathbb{Z}^n} b_{x+z} .$$

 γ is translation invariant if $\mu_z \gamma \mu_z^{-1} = \gamma$ for all $z \in \mathbb{Z}^n$.

(2) Causal relative to some neighborhood \mathcal{N} : For every element $A_z \in \mathcal{Z}$ local upon z, $\gamma(A_z)$ is local upon \mathcal{N}_z (see Definition 2.3).

At each time step of evolution, $b \in \mathbb{Z}$ evolves to $\gamma(b)$.

Just as in the Schrödinger template, Definition 2.2, the neighborhood \mathcal{N} is typically chosen to be the smallest subset of \mathbb{Z}^n satisfying the causality condition in the above definition.

3. Equivalence of the Schrödinger and Heisenberg Templates of QCA

In this section, we address the relation between the Schrödinger and Heisenberg templates. First, let us assume that we are given a Schrödinger template $(\mathcal{H}, [R])$, with neighborhood \mathcal{N} as in Definition 2.2, where \mathcal{H} is constructed from a finite dimensional Hilbert space W. Then the ITPA \mathcal{Z} constructed from $\operatorname{End}(W)$ is embeddeded in the algebra of bounded linear operators $B(\mathcal{H})$. The following useful result (Theorem 3.9 from [35]) says that conjugation of \mathcal{Z} by R is an automorphism of \mathcal{Z} .

Theorem 3.1. Let R be a unitary and causal operator on \mathcal{H} relative to some neighborhood \mathcal{N} . Then for every $x \in \mathbb{Z}^n$,

$$\mathcal{A}_x \subset \operatorname{span}(\prod_{y \in \mathcal{N}} R^{\dagger} \mathcal{A}_{x-y} R).$$

In particular $\mathcal{Z} = R^{\dagger} \mathcal{Z} R$.

Let the automorphism $\gamma_{[R]}$ be,

$$\gamma_{[R]} : \mathcal{Z} \longrightarrow \mathcal{Z}$$

$$b \mapsto R^{\dagger} b R ,$$

$$(7)$$

where R is some representative of [R]. $\gamma_{[R]}$ is a *-automorphism as it commutes with the *-involution of \mathcal{Z} (adjoint map).

We recall a theorem from [20] deriving the counterpart of causality in the Heisenberg template from that in the Schrödinger template. Let the *reflected neighborhood* of cell z be \mathcal{V}_z ,

$$\mathcal{V}_z = z - \mathcal{N}$$
 .

Theorem 3.2 (Structural Reversibility, Theorem 3 in [20]). Let $R : \mathcal{H} \longrightarrow \mathcal{H}$ be a unitary operator and \mathcal{N} a neighborhood. Then the following are equivalent:

- (1) R is causal relative to the neighborhood \mathcal{N} .
- (2) For every operator A_z local upon cell z, $R^{\dagger}A_zR$ is local upon \mathcal{N}_z .
- (3) R^{\dagger} is causal relative to the neighborhood \mathcal{V} .
- (4) For every operator A_z local upon cell z, RA_zR^{\dagger} is local upon \mathcal{V}_z .

Part 2 of this theorem implies that $\gamma_{[R]}$ (7) is causal in the Heisenberg template sense, with the same neighborhood \mathcal{N} as in the Schrödinger template $(\mathcal{H}, [R])$. We can also show that $\gamma_{[R]}$ is translation invariant in the Heisenberg template sense. This follows because [R]is translation invariant, and because for all $b \in \mathcal{Z}$,

$$\mu_z(b) = \tau_z b \tau_z^{-1},$$

where μ_z is the translation operator in the Heisenberg template, and τ_z is the translation operator in the Schrödinger template (Definition 2.2.1). Consequently, for any representative R of [R],

$$\begin{split} \mu_z \gamma_{[R]} \mu_z^{-1}(b) &= \tau_z R^{\dagger} \tau_z^{-1} b \tau_z R \tau_z^{-1} \\ &= e^{-i\theta_z} R^{\dagger} b e^{i\theta_z} R \\ &= R^{\dagger} b R = \gamma_{[R]}(b) \,. \end{split}$$

So $\gamma_{[R]}$ is translation invariant.

From the above discussion, it is clear that given a global evolution operator [R] on \mathcal{H} , a *-automorphism $\gamma_{[R]}$ of \mathcal{Z} always exists. This implies that given a Schrödinger template, we can always obtain a Heisenberg template (Definition 2.6) with equivalent dynamics, i.e., yielding the same measurement statistics as in the Schrödinger template.

Given a Heisenberg template, the condition that determines if a dynamically equivalent Schrödinger template exists is best phrased in the language of representations. The obvious irreducible action of \mathcal{Z} on $\mathbb{V} = \text{Span}(\mathcal{C})$ (2),

$$\mathcal{Z} \times \mathbb{V} \mapsto \mathbb{V} \\
\left(\bigotimes_{x \in \mathbb{Z}^n} A_x, \bigotimes_{x \in \mathbb{Z}^n} |b_x\rangle \right) \to \bigotimes_{x \in \mathbb{Z}^n} A_x(|b_x\rangle), \tag{8}$$

extends to an irreducible representation on \mathcal{H} . We denote this canonical representation (π, \mathcal{H}) . This representation can be obtained as a direct limit, as shown in Guichardet [16].

Before we state the definition of equivalence, we need to be certain that properties like translation invariance and causality would carry over from the Heisenberg template to the Schrödinger template.

Proposition 3.3. Let W be a finite dimensional Hilbert space, and (\mathcal{Z}, γ) be a Heisenberg template, where \mathcal{Z} is constructed from End(W). Suppose there exists a UFC Hilbert space \mathcal{H} constructed over W, and a unitary operator [R] on \mathcal{H} that intertwines the representations $(\pi\gamma, \mathcal{H})$ and (π, \mathcal{H}) of \mathcal{Z} , i.e., some representative R of [R] satisfies

$$R^{\dagger}\pi(b)R = \pi\left(\gamma(b)\right) \quad \forall b \in \mathcal{Z} \; .$$

Then

- (1) [R] is unique.
- (2) [R] is translation invariant.
- (3) If γ is causal relative to the neighborhood \mathcal{N} , then [R] is causal relative to the same neighborhood \mathcal{N} .
- *Proof.* (1) R is unique up to a phase factor by Schur's Lemma (Lemma A.2, Appendix A) as the representations (π, \mathcal{H}) and $(\pi\gamma, \mathcal{H})$ are irreducible; thus, [R] is unique.
 - (2) Observe that for all $b \in \mathbb{Z}$ and all $z \in \mathbb{Z}^n$,

$$\pi(\mu_z(b)) = \tau_z \pi(b) \tau_z^{-1}$$
,

which implies

$$\pi(\mu_z \gamma \mu_z^{-1}(b)) = \tau_z \pi(\gamma \mu_z^{-1}(b)) \tau_z^{-1}$$

= $\tau_z R^{\dagger} \pi(\mu_z^{-1}(b)) R \tau_z^{-1}$
= $\tau_z R^{\dagger} \tau_z^{-1} \pi(b) \tau_z R \tau_z^{-1}$.

Since γ is translation invariant,

$$\pi(\gamma(b)) = \pi(\mu_z \gamma \mu_z^{-1}(b))$$
$$= \tau_z R^{\dagger} \tau_z^{-1} \pi(b) \tau_z R \tau_z^{-1}$$

This implies that $\tau_z R \tau_z^{-1}$ intertwines the representations (π, \mathcal{H}) and $(\pi \gamma, \mathcal{H})$. But R is the unique intertwiner up to a phase factor, so

$$\tau_z R \tau_z^{-1} = e^{i\theta_z} R \,.$$

- for some $\theta_z \in \mathbb{R}$. Thus [R] is translation invariant.
- (3) This follows by the equivalence of Theorem 3.2 parts 1 and 2.

We are now justified in defining the equivalence of templates as follows.

Definition 3.4. Let W be a finite dimensional Hilbert space. A Heisenberg template (\mathcal{Z}, γ) , where \mathcal{Z} is constructed from End(W), and a Schrödinger template $(\mathcal{H}, [R])$, where \mathcal{H} is constructed from W, are *equivalent* if [R] intertwines the representations $(\pi\gamma, \mathcal{H})$ and (π, \mathcal{H}) of \mathcal{Z} , i.e., some representative R of [R] satisfies

$$R^{\dagger}\pi(b)R = \pi\left(\gamma(b)
ight) \quad \forall b \in \mathcal{Z}$$
 .

The following result shows that a Schrödinger template exists for any QCA given as a Heisenberg template and for any UFC Hilbert space in which the ITPA of the Heisenberg template is naturally embedded.

Theorem 3.5. Let W be a finite dimensional Hilbert space. Given a Heisenberg template (\mathcal{Z}, γ) , where \mathcal{Z} is constructed from $\operatorname{End}(W)$, an equivalent Schrödinger template exists for any UFC Hilbert space \mathcal{H} constructed from W.

Proof. We consider the vector space $\mathbb{V} = \text{Span}(\mathcal{C})$ (2), and show that there is an automorphism of \mathbb{V} that intertwines the \mathcal{Z} (ITPA) representations $(\pi\gamma, \mathbb{V})$ and (π, \mathbb{V}) . Then we extend it to the UFC Hilbert space \mathcal{H} .

Given that γ is an automorphism of \mathcal{Z} , and $\mathcal{A}_x \cap \mathcal{A}_y = \mathbb{CI}$ for $x \neq y$, implies that,

$$\gamma(\mathcal{A}_x) \cap \gamma(\mathcal{A}_y) = \mathbb{CI} \quad \text{for } x \neq y.$$

Consider, a general finite subset $D \subset \mathbb{Z}^n$. The image of \mathcal{A}_D under γ is $\gamma(\mathcal{A}_D)$. Let the commutant of $\gamma(\mathcal{A}_D)$ in \mathcal{Z} be $\gamma(\mathcal{A}_D)^C$,

$$\gamma(\mathcal{A}_D)^C = \{T \in \mathcal{Z} : \gamma(a)T = T\gamma(a) \text{ for all } a \in \mathcal{A}_D\}$$
$$= \bigcup_k \operatorname{Span}(\prod_{u \in D_k \setminus D} \gamma(\mathcal{A}_u))$$

 $\gamma(\mathcal{A}_D)$ is local upon (i.e., non-identity on, as in Definition 2.3) a finite set of tensor factor indices by the causality of γ , required in Definition 2.6 (1). Let $K \subset \mathbb{Z}^n$ be the smallest set such that $\gamma(\mathcal{A}_D) \subset \mathcal{A}_K = \bigotimes_{y \in K} \mathcal{A}_y$. By (3) again we are justified in writing

$$\mathcal{Z} = \operatorname{Span}(\gamma(\mathcal{A}_D)\gamma(\mathcal{A}_D)^C).$$

 $\mathcal{A}_K \subset \mathcal{Z}$, so

$$\mathcal{A}_{K} = \operatorname{Span}(\gamma(\mathcal{A}_{D})\gamma(\mathcal{A}_{D})^{C}) \cap \mathcal{A}_{K}$$
$$= \operatorname{Span}\left(\gamma(\mathcal{A}_{D})(\gamma(\mathcal{A}_{D})^{C} \cap \mathcal{A}_{K})\right), \qquad (9)$$

where the second equality follows since $\gamma(\mathcal{A}_D) \subset \mathcal{A}_K$. It allows us to consider $\gamma(\mathcal{A}_D)$ as a subalgebra of \mathcal{A}_K instead of \mathcal{Z} . We call the restriction of the representation $(\pi\gamma, \mathbb{V})$ to a subalgebra $\mathcal{A} \subset \mathcal{Z}$, the action of \mathcal{A} via $\pi\gamma$ on \mathbb{V} . The action of algebra $\mathcal{A}_D = \bigotimes_{x \in D} \mathcal{A}_x$ via $\pi\gamma$ on \mathbb{V} , because of this, can be restricted to $\bigotimes_{x \in K} W$, and extended trivially to the rest of \mathbb{V} . $\gamma(\mathcal{A}_D) \cong \bigotimes_{x \in D} \operatorname{End}(W) = \operatorname{End}(\bigotimes_{x \in D} W)$ is a simple, self-adjoint algebra. Let us denote the canonical representation of $\operatorname{End}(W)$ on W by (χ, W) . Up to equivalence, the finite dimensional irreducible module of $\mathcal{A}_D = \bigotimes_{x \in D} \operatorname{End}(W)$ occurring in $\bigotimes_{x \in K} W$ is the

outer tensor product (24) representation ($\bigotimes_{x \in D} \chi, \bigotimes_{x \in D} W$). By the Double-Commutant Theorem, Theorem A.5, there is an \mathcal{A}_D module isomorphism,

$$S_K: U \otimes \left(\bigotimes_{x \in D} W\right) \longrightarrow \bigotimes_{x \in K} W,$$

where $U = \operatorname{Hom}_{\mathcal{A}_D}(\bigotimes_{x \in D} \chi, \bigotimes_{x \in K} W)$ is the multiplicity space for \mathcal{A}_D action via $\pi \gamma$ on $\bigotimes_{x \in K} W$. Also, under this isomorphism,

$$\mathcal{A}_D \cong \mathbb{I} \otimes \operatorname{End} \left(\bigotimes_{x \in D} W \right)$$
.

This implies that the irreducible representation of \mathcal{A}_D in its action via $\pi\gamma$ on \mathbb{V} is the outer tensor product (24) representation $(\widehat{\bigotimes}_{x\in D}\chi, \bigotimes_{x\in D}W)$, which is locally irreducible. Therefore, by the primary decomposition (23), there is an \mathcal{A}_D module isomorphism,

$$S_D : E_D \otimes \left(\bigotimes_{x \in D} W\right) \cong \mathbb{V}$$

$$\epsilon_D \otimes \bigotimes_{x \in D} w_x \mapsto \epsilon_{D_n} \left(\bigotimes_{x \in D} w_x\right),$$
(10)

where $E_D = \operatorname{Hom}_{\mathcal{A}_D}(\bigotimes_{x \in D} \chi, \mathbb{V})$ is the multiplicity space for \mathcal{A}_D action via $\pi\gamma$. Denote by \mathcal{Z}^D the commutant of \mathcal{A}_D in \mathcal{Z} for action via $\pi\gamma$ on \mathbb{V} ,

$$\mathcal{Z}^{D} = \{T \in \mathcal{Z} : \pi\gamma(a)\pi\gamma(T) = \pi\gamma(T)\pi\gamma(a) \text{ for all } a \in \mathcal{A}_{D}\}$$
$$= \{T \in \mathcal{Z} : aT = Ta \text{ for all } a \in \mathcal{A}_{D}\}$$
$$= \bigcup_{k} \bigotimes_{y \in D_{k} \setminus D} \mathcal{A}_{y} .$$
(11)

 $T \in \mathcal{Z}^D$ (11) acts as $\gamma(T)$ on E_D by left multiplication (see the discussion in Appendix A after Proposition A.4). Let us denote by l the left multiplication by an element of $\gamma(\mathcal{Z}^D)$ on E_D . Since γ is an automorphism and \mathbb{V} is an irreducible \mathcal{Z} module, this implies $E_D \otimes (\bigotimes_{x \in D} W)$ is an irreducible $\mathcal{Z}^D \otimes \mathcal{A}_D$ module under the outer tensor product action $(l \circ \gamma) \otimes (\bigotimes_{x \in D} \chi)$. Thus, E_D is an irreducible \mathcal{Z}^D module under $l \circ \gamma$. We denote by V_D the vector space on the left side of equation (10),

$$V_D = E_D \otimes \left(\bigotimes_{x \in D} W\right).$$

Let us consider the canonical representation of \mathcal{A}_D on \mathbb{V} via π , i.e., restriction of π to \mathcal{A}_D . Then we have an \mathcal{A}_D module isomorphism

$$L_D: U_D \otimes \left(\bigotimes_{x \in D} W\right) \cong \mathbb{V}$$
$$u_D \otimes \bigotimes_{x \in D} w_x \mapsto u_D\left(\bigotimes_{x \in D} w_x\right),$$

where $U_D = \operatorname{Hom}_{\mathcal{A}_D}(\bigotimes_{x \in D} \chi, \mathbb{V})$ is the multiplicity space for \mathcal{A}_D action via π . We can identify the multiplicity space U_D in this case with $\mathbb{V}_{\overline{D}}$ (4), the vector space complement in \mathbb{V} to the tensor factors in D. We are implicitly identifying the indices of the modules $\bigotimes_{x \in D} W$ with those in \mathbb{V} for the \mathcal{A}_D actions via $\pi\gamma$ and π . Choose any map $F \in \text{Hom}(E_D, U_D)$. Then $L_D(\mathbb{I} \otimes F)S_D^{-1} \in \text{End}_{\mathcal{A}_D}(\mathbb{V})$, i.e., is an \mathcal{A}_D module homomorphism of V intertwining \mathcal{A}_D actions via $\pi\gamma$ and π . Let us denote by \mathcal{R}_D the vector space of \mathcal{A}_D module homomorphisms of V intertwining \mathcal{A}_D actions via $\pi\gamma$ and π ,

$$\mathcal{R}_D = \operatorname{End}_{\mathcal{A}_D}(\mathbb{V}) \,. \tag{12}$$

We have now set up the objects we need for a finite subset $D \subset \mathbb{Z}^n$.

Next, we specialize to the sets $\{D_k\}$ in the ascending chain (3) and relate the spaces and morphisms associated with them. Let $D_m \subset D_n \in \{D_k\}$. We construct a \mathcal{Z} module isomorphism between the spaces V_{D_m} and V_{D_n} ,

$$V_{D_m} = E_{D_m} \otimes \bigotimes_{x \in D_m} W,$$

$$V_{D_n} = E_{D_n} \otimes \bigotimes_{x \in D_n} W.$$
(13)

There is a canonical map,

$$J_{n,m}: E_{D_n} \otimes \bigotimes_{x \in D_n \setminus D_m} W \to E_{D_m} ,$$

that we need to describe first. Choose an orthonormal basis $\mathcal{B} \subset W$ containing the quiescent state $|q\rangle \in W$, as in the definition of the set of finite, unbounded configurations (1). Observe that an element $\epsilon_{D_n} \in E_{D_n}$ is a linear map

$$\epsilon_{D_n}: \bigotimes_{x \in D_n} W \to \mathbb{V}.$$

We can write $\bigotimes_{x \in D_n} W$ in the form

$$\bigotimes_{x\in D_n} W = \bigotimes_{x\in D_m} W \otimes \bigotimes_{y\in D_n\setminus D_m} W ,$$

expressing a basis element as

$$\bigotimes_{x\in D_n} b_x = \bigotimes_{x\in D_m} b_x \otimes \bigotimes_{y\in D_n\setminus D_m} b_y ,$$

where $b_x \in \mathcal{B}$. Then ϵ_{D_n} maps such an element to \mathbb{V} ,

$$\epsilon_{D_n} : \bigotimes_{x \in D_n} W \to \mathbb{V}$$
$$\bigotimes_{x \in D_m} b_x \otimes \bigotimes_{y \in D_n \setminus D_m} b_y \mapsto \epsilon_{D_n} \big(\bigotimes_{x \in D_m} b_x \otimes \bigotimes_{y \in D_n \setminus D_m} b_y \big) .$$

We obtain $J_{n,m}$ by evaluating elements of E_{D_n} at the components corresponding to the indices in $D_n \setminus D_m$,

$$J_{n,m}: E_{D_n} \otimes \bigotimes_{y \in D_n \setminus D_m} W \to E_{D_m}$$

$$\epsilon_{D_n} \otimes \bigotimes_{y \in D_n \setminus D_m} b_y \mapsto \epsilon_{D_m} \Big(\bigotimes_{x \in D_m} b_x \Big) = \epsilon_{D_n} \Big(\bigotimes_{x \in D_m} b_x \otimes \bigotimes_{y \in D_n \setminus D_m} b_y \Big) , \qquad (14)$$

and extending by linearity. We extend $J_{n,m}$ to V_{D_n} by identity on the indices in D_m , and denote this map by $\hat{J}_{n,m}$,

$$\hat{J}_{n,m}: E_{D_n} \otimes \bigotimes_{x \in D_n} W \to E_{D_m} \otimes \bigotimes_{x \in D_m} W$$

 $\hat{J}_{n,m}$ intertwines the irreducible actions of \mathcal{Z} on V_{D_n} and V_{D_m} (13). By Schur's Lemma (Lemma A.2), it is an isomorphism. Note that this further implies that $J_{n,m}$ (14) is an isomorphism. For $D_l \subset D_m \subset D_n$, we have that

$$\hat{J}_{n,l} = \hat{J}_{m,l} \circ \hat{J}_{n,m} \,.$$
 (15)

Also, $\hat{J}_{k,k} = \mathbb{I}$ for all k. Let

$$\hat{T}_{m,n} = \hat{J}_{n,m}^{-1} : V_{D_m} \to V_{D_n} .$$
 (16)

 $\{V_{D_m}, \hat{T}_{m,n}\}$ form a direct system (see Appendix B) of isomorphic \mathcal{Z} modules, i.e., for $D_l \subset D_m \subset D_n$, $\hat{T}_{l,n} = \hat{T}_{m,n} \circ \hat{T}_{l,m}$, which follows from (15). The direct limit of this system exists by Theorem B.1. Let $\hat{\mathbb{V}}$ be this direct limit,

$$\hat{\mathbb{V}} = \lim_{m \to \infty} V_{D_m}$$

Observe that \mathbb{V} also satisfies the universal property of a direct limit for the system $(V_{D_m}, \hat{T}_{m,n})$. To see this, without loss of generality, include the empty set in the ascending chain of sets (3), $D_0 = \emptyset \in \{D_k\}$. Then $V_{\emptyset} = \mathbb{V}$. Let $f^k = \hat{T}_{0,k}^{-1} = \hat{J}_{k,0}$ for D_k . Then $(\mathbb{V}, \{f^k\})$ is a target, since, $f^m = f^n \circ \hat{T}_{m,n}$ for $D_m \subset D_n$.

Let $(\mathbb{W}, \{g^k\})$ be another target. Let $g = g^0 : V_{\emptyset} = \mathbb{V} \to \mathbb{W}$. Then $g^k = g \circ f^k$. The map g is unique, i.e., if there is another map $h : V_{\emptyset} = \mathbb{V} \to \mathbb{W}$ satisfying $g^k = h \circ f^k$, then h = g, since f^k are isomorphisms. This shows that $(\mathbb{V}, \{f^k\})$ satisfies the universal property. Thus \mathbb{V} is isomorphic to $\hat{\mathbb{V}}$, which follows by the uniqueness up to unique isomorphism of the direct limit. Let us denote this isomorphism by \hat{S} ,

 $\hat{S}: \hat{\mathbb{V}} \to \mathbb{V},$

Let us denote by $(\hat{\pi}, \hat{\mathbb{V}})$ the representation of \mathcal{Z} on $\hat{\mathbb{V}}$. Note that the \mathcal{Z} isomorphisms $\hat{T}_{m,n}$ (16) are defined with respect to the canonical isomorphism (22), implying that the intertwining of actions is independent of the class representative in the direct system. $\hat{\mathbb{V}}$ is thus the natural representation of \mathcal{Z} on $\hat{\mathbb{V}}$ compatible with the tensor product structures of \mathcal{Z} and $\hat{\mathbb{V}}$, that of $\hat{\mathbb{V}}$ coming from $\hat{T}_{m,n}$ (16). \hat{S} intertwines the irreducible \mathcal{Z} representations $(\hat{\pi}, \widehat{\mathbb{V}})$ and $(\pi\gamma, \mathbb{V})$.

Next, we look for the \mathcal{Z} module isomorphism of \mathbb{V} intertwining the \mathcal{Z} representations $(\pi\gamma, \mathbb{V})$ and (π, \mathbb{V}) . Under the isomorphisms in (14), we have that

$$\operatorname{Hom}(E_{D_m}, U_{D_m}) \cong \operatorname{End}(\bigotimes_{x \in D_n \setminus D_m} W) \otimes \operatorname{Hom}(E_{D_n}, U_{D_n})$$

The trace, $\langle A, B \rangle = \operatorname{tr}(AB)$, for $A, B \in \operatorname{End}(\bigotimes_{x \in D_n \setminus D_m} W)$, is a non-degenerate bilinear form. Let $P_{m,n}$ be the linear operator

$$P_{m,n} : \operatorname{End}(\bigotimes_{x \in D_n \setminus D_m} W) \otimes \operatorname{Hom}(E_{D_n}, U_{D_n}) \to \operatorname{Hom}(E_{D_n}, U_{D_n})$$
$$\alpha \otimes \beta \mapsto \operatorname{tr}(\alpha)\beta .$$

 $P_{m,n}$ is a surjective homomorphism. By the discussion before (12), under the isomorphisms in (14), $P_{m,n}$ induces a surjective homomorphism of vector spaces,

$$M_{m,n}: \mathcal{R}_{D_m} = \operatorname{End}_{\mathcal{A}_{D_m}}(\mathbb{V}) \to \mathcal{R}_{D_n} = \operatorname{End}_{\mathcal{A}_{D_n}}(\mathbb{V})$$

For $\nu, \eta \in \text{End}(W)$, $\text{tr}(\nu \otimes \eta) = \text{tr}(\nu)\text{tr}(\eta)$. This implies, for $D_l \subset D_m \subset D_n$, $M_{l,n} = M_{m,n} \circ M_{l,m}$. Thus, $\{\mathcal{R}_{D_m}, M_{m,n}\}$ form a direct system. Let its direct limit, which exists by Theorem B.1, be

$$\mathcal{R} = arprojlim_{D_m}$$
 .

Remark 3.6. The vector space \mathbb{V} is a direct limit (25), the ITPA \mathcal{Z} and the canonical representation π can also be constructed as direct limits (Guichardet [16]). The direct limits in this proof should be viewed in that context.

Choose $\hat{R} \in \hat{\mathcal{R}}$,

$$\hat{R}: \mathbb{V} \to \mathbb{V}$$

Then \hat{R} intertwines the \mathcal{Z} representations $(\pi\gamma, \mathbb{V})$ and (π, \mathbb{V}) . By Schur's Lemma (Lemma A.2), \hat{R} is the unique \mathcal{Z} module isomorphism up to a phase factor. Since γ is a *-automorphism, \hat{R} is unitary on \mathbb{V} . Further, by the Bounded Linear Transformation (BLT) Theorem (Reed and Simon [34] Theorem 1.7, pg. 9), \hat{R} can be uniquely and unitarily extended to \mathcal{H} . We denote this extension to \mathcal{H} by R,

$$R: \mathcal{H} \to \mathcal{H},\tag{17}$$

By Proposition 3.3, the Schrödinger template given by $(\mathcal{H},[R])$ is equivalent to the given Heisenberg template (\mathcal{Z}, γ) .

4. Template equivalence and the quantum lattice gas automaton

A QLGA is a model of multiple particles propagating on a lattice and scattering at the lattice points (called *sites* in this context) through interactions with each other at the lattice points or through self-interactions. In [35], the authors studied the conditions that classify a QCA as a QLGA. The definition of a QLGA given in [35] is a Schrödinger template based on a particular UFC Hilbert space. The QLGA classifying condition, however, involves propagating cell algebras interacting at a cell, belonging to the Heisenberg template. Just as in the main theorem of [35], Theorem III.16 in that paper, we start from the QLGA classifying condition of a QCA, and construct the same Schrödinger template as in [35], albeit in a mode that reflects the approach of this paper. Through this construction, we will see restrictions being imposed beyond those stipulated by the classifying condition to reach the explicit form required by the QLGA definition of [35]. We then scrutinize these restrictions, now apprised of the more general Schrödinger template implied by Theorem 3.5.

Let us recall the definition of a QLGA from [35].

Definition 4.1. A QLGA is defined on a lattice \mathbb{Z}^n , and for a neighborhood \mathcal{N} , as follows:

- (1) The cell Hilbert space is $W = \bigotimes_{z \in \mathcal{N}} V_z$, for some finite dimensional vector spaces $\{V_z\}_{z \in \mathcal{N}}$.
- (2) The quiescent state $|q\rangle$, which is a simple product,

$$|q\rangle = \bigotimes_{z \in \mathcal{N}} |q_z\rangle$$
, where $|q_z\rangle \in V_z$

- (3) A UFC Hilbert space \mathcal{H} defined in terms of $W = \bigotimes_{z \in \mathcal{N}} V_z$ such that the quiescent states in all cells are identical $|q_x\rangle = |q\rangle$ for all $x \in \mathbb{Z}^n$.
- (4) A propagation operator relative to the neighborhood $\mathcal{N}, \sigma : \mathcal{H} \mapsto \mathcal{H}$, defined on the basis of \mathcal{H} to be

$$\sigma: \bigotimes_{x \in \mathbb{Z}^n} \bigotimes_{z \in \mathcal{N}} |\mathbf{k}_x(z)\rangle \mapsto \bigotimes_{x \in \mathbb{Z}^n} \bigotimes_{z \in \mathcal{N}} |\mathbf{k}_{x+z}(z)\rangle$$
(18)

(5) A local scattering operator F, which is a unitary operator on the site Hilbert space $W = \bigotimes_{z \in \mathcal{N}} V_z$, such that F fixes $|q\rangle$ (an eigenvector with eigenvalue one): $F|q\rangle = |q\rangle$. The global scattering operator $\tilde{F} : \mathcal{H} \mapsto \mathcal{H}$, is the application of F at every cell, defined as

$$\tilde{F} = \bigotimes_{x \in \mathbb{Z}^n} F \tag{19}$$

(6) A global evolution operator R consisting of propagation σ followed by the scattering \tilde{F} :

$$R = \tilde{F}\sigma$$

A state of the QLGA is a vector in the UFC Hilbert space.

First, let us recall the terminology of [35], but generalized to an arbitrary Heisenberg template automorphism γ . The patch of propagated image $\gamma(\mathcal{A}_z)$ on \mathcal{A}_x , is

$$\mathcal{D}_{z,x} = \gamma(\mathcal{A}_z) \cap \mathcal{A}_x.$$

We restate a version of Theorem III.10 of [35], as we will refer to it. The proof of this version would be very similar to that of Theorem III.10 in [35], except in the current context we have γ , whereas in [35] the proof was in the context of $\gamma_{[R]}$ (7).

Theorem 4.2 (Theorem III.10 in [35]). Let (\mathcal{Z}, γ) be the Heisenberg template of a QCA with neighborhood \mathcal{N} . Then $\mathcal{A}_x = \operatorname{span}(\prod_{y \in \mathcal{N}} \mathcal{D}_{x-y,x})$ if and only if there exists an isomorphism of vector spaces:

$$S: W \longrightarrow \bigotimes_{z \in \mathcal{N}} V_z \tag{20}$$

for some vector spaces $\{V_z\}_{z \in \mathcal{N}}$. Under the isomorphism S, for each $y \in \mathcal{N}$:

$$\mathcal{D}_{x-y,x} \cong \operatorname{End}(V_y) \otimes \bigotimes_{z \in \mathcal{N}, z \neq y} \mathbb{I}_{V_z}$$

The condition $\mathcal{A}_x = \operatorname{span}(\prod_{y \in \mathcal{N}} \mathcal{D}_{x-y,x})$ is the local characterization of the Schrödinger template of a QCA as a QLGA in Theorem III.16 in [35], although, it is manifestly a condition on γ , and hence on the Heisenberg template. Assuming this condition, we work toward a Schrödinger template in which the global evolution operator is explicit. First, we replace W with S(W), where S is the isomorphism in (20) and can be taken to be an isometric isomorphism under an appropriate inner product choice on S(W). Under this substitution, and abusing notation, we can assume that W has the tensor product structure implied in (20). Take the quiescent state $|q\rangle \in W = \bigotimes_{z \in \mathcal{N}} V_z$ to be any product vector in W,

$$|q\rangle = \bigotimes_{z \in \mathcal{N}} |q_z\rangle$$
, where $|q_z\rangle \in V_z$.

Using W and $|q\rangle$, we construct the UFC Hilbert space \mathcal{H} as described in Section 2. Next, define a propagation operator σ as in (18).

$$\sigma: \bigotimes_{x \in \mathbb{Z}^n} \bigotimes_{z \in \mathcal{N}} |\mathbf{k}_x(z)\rangle \mapsto \bigotimes_{x \in \mathbb{Z}^n} \bigotimes_{z \in \mathcal{N}} |\mathbf{k}_{x+z}(z)\rangle$$

Then $\sigma \pi \gamma(\mathcal{A}_x) \sigma^{-1} = \pi(\mathcal{A}_x)$, i.e., $\sigma \pi \gamma(\mathcal{A}_x) \sigma^{-1}$ is an automorphic image of $\pi(\mathcal{A}_x)$. But $\pi(\mathcal{A}_x) = \operatorname{End}(W)$. Therefore, there is a unitary map F on W (Schur's Lemma applied to \mathcal{A}_x action on W), such that

$$\sigma \pi \gamma(\mathcal{A}_x) \sigma^{-1} = F^{-1} \pi(\mathcal{A}_x) F \tag{21}$$

for each $x \in \mathbb{Z}^n$. By translation invariance of γ , the same F works for every cell. Note that F only depends on γ and not on the choice of $|q\rangle$. We say that F is associated with γ .

We state and prove the direction of the main theorem in [35], Theorem III.16, in which the local condition $\mathcal{A}_x = \operatorname{span}(\prod_{y \in \mathcal{N}} \mathcal{D}_{x-y,x})$ on the Heisenberg template of a QCA implies that its Schrödinger template is a QLGA (upto a global isomorphism). An eigenvector of an operator on a finite dimensional tensor product vector space will be called a *product eigenvector* if it is a simple product in the vector space.

Theorem 4.3 (See Theorem III.16 in [35]). Let (\mathcal{Z}, γ) be the Heisenberg template of a QCA with neighborhood \mathcal{N} , satisfying: $\mathcal{A}_x = \operatorname{span}(\prod_{y \in \mathcal{N}} \mathcal{D}_{x-y,x})$. Then it is a QLGA in the sense of Definition 4.1 if and only if F(21) associated with γ has a product eigenvector in W.

Proof. Suppose that F has a product eigenvector $|w\rangle \in W$. Since F is unitary, $F |w\rangle = e^{i\theta} |w\rangle$ for some $\theta \in \mathbb{R}$. We can replace F with $e^{-i\theta}F$, which fixes $|w\rangle$, and is still associated with γ in (21). Without loss of generality then, let us assume that F fixes $|w\rangle$. Let $|q\rangle = |w\rangle$. We can now construct a cell-wise automorphism \tilde{F} on \mathcal{H} as in (19),

$$\tilde{F} = \bigotimes_{x \in \mathbb{Z}^n} F,$$

As $\mathcal{A}_x, x \in \mathbb{Z}^n$, generate \mathcal{Z} ,

$$\sigma \pi \gamma(b) \sigma^{-1} = \tilde{F}^{-1} \pi(b) \tilde{F}$$

for all $b \in \mathcal{Z}$. Rewriting the above relation, we obtain

$$\pi\gamma(b) = \sigma^{-1}\tilde{F}^{-1}\pi(b)\tilde{F}\sigma.$$

Thus

$$R = \tilde{F}\sigma$$

intertwines (π, \mathcal{H}) and $(\pi\gamma, \mathcal{H})$. By Proposition 3.3, [R] is the unique such global evolution operator. Indeed it is a QLGA, as it is composed of the propagation operator σ followed by the scattering operator \tilde{F} .

Conversely, if F does not have a product eigenvector in W, that precludes defining a quiescent state $|q\rangle \in W$ from which to construct a UFC Hilbert space \mathcal{H} : both σ and \tilde{F} are required to have $\bigotimes_{x \in \mathbb{Z}^n} |q\rangle \in \mathcal{H}$ as an eigenvector by Lemma C.2 (Appendix C). Thus, it is not a QLGA in the sense of Definition 4.1.

Let us understand the contents of this derivation by way of the proof of Theorem 3.5. Instead of treating the general case, we take a simple special case that illustrates the ideas in Theorem 3.5 for the QLGA of Definition 4.1. We take a one-dimensional QLGA with neighborhood $\{0, 1\}$. Let $D_0 = \emptyset$ and $D_k = \{0, \ldots, k-1\}$ for k > 0, in the ascending chain of sets (3). Assume that the cell Hilbert space is $W = \mathbb{C}^2 \otimes \mathbb{C}^2$, with basis $\{|ij\rangle\}, i, j \in \{0, 1\}$. Let the quiescent symbol be $|q\rangle = |00\rangle$. Assume also that there is no scattering. Then, in the Heisenberg template, the automorphism γ is defined by

$$\dots (\mathbb{I} \otimes \mathbb{I}) \otimes (\mathbb{I} \otimes \mathbb{I}) \otimes \underbrace{(a \otimes b)}_{x} \otimes (\mathbb{I} \otimes \mathbb{I}) \otimes (\mathbb{I} \otimes \mathbb{I}) \dots \xrightarrow{\gamma} \dots (\mathbb{I} \otimes \mathbb{I}) \otimes (\mathbb{I} \otimes \mathbb{I}) \otimes \underbrace{(a \otimes \mathbb{I})}_{x} \otimes \underbrace{(\mathbb{I} \otimes b)}_{x+1} \otimes (\mathbb{I} \otimes \mathbb{I}) \otimes (\mathbb{I} \otimes \mathbb{I}) \dots$$

Now consider the Schrödinger template that Theorem 3.5 implies. We show some of the basis elements of \mathcal{H} and their transformation by R (17).

$$\dots |00\rangle |00\rangle \underbrace{|i0\rangle}_{x} \underbrace{|0j\rangle}_{x+1} |00\rangle |00\rangle \dots \xrightarrow{R} \dots |00\rangle |00\rangle \underbrace{|ij\rangle}_{x} |00\rangle |00\rangle \dots,$$

$$\dots |00\rangle |00\rangle \underbrace{|i0\rangle}_{x} \underbrace{|kj\rangle}_{x+1} \underbrace{|01\rangle}_{x+2} |00\rangle |00\rangle \dots \xrightarrow{R} \dots |00\rangle |00\rangle \underbrace{|ij\rangle}_{x} \underbrace{|kl\rangle}_{x+1} |00\rangle |00\rangle \dots.$$

The intertwining of the representations $(\pi\gamma, \mathcal{H})$ and (π, \mathcal{H}) by R is clear. We observe that R behaves as the propagation operator σ in Definition 4.1, except that we interpret it as the isomorphism arising from Theorem 3.5. This also provides a concrete interpretation of Theorem 3.5 when an explicit form of the intertwining operator R, the global evolution operator, can be constructed.

We can also consider how the general case of Theorem 3.5 compares with the QLGA definition, Definition 4.1. The local condition in Theorem 4.2 on the Heisenberg template has the direct consequence that the cell Hilbert space has a tensor product structure. This is a condition that the Heisenberg template requires and would carry over to the Schrödinger template in Theorem 3.5. In order to create the QLGA of Definition 4.1, further restrictions were imposed: the associated local scattering operator F associated with γ has at least one product eigenvector, and the quiescent state $|q\rangle$ is one of these eigenvectors. Theorem 3.5 circumvents these at the expense of the form of the global evolution operator.

5. CONCLUSION

The discussion in this paper is focused on the relation between the Schrödinger and Heisenberg templates of QCA, within a borader context of representations of von Neumann algebras. ITPA is encountered in discussions of hyperfinite II_1 factor, such as in [15, 17]. In this paper the representations of ITPA being alluded to are the *-algebra representations in the specific context of QCA, which in the Schrödinger template are defined on a separable Hilbert space, constituting a I_{∞} factor. We show that each Heisenberg template has an equivalent Schrödinger template. Moreover, an equivalent Schrödinger template exists for any UFC Hilbert space in whose space of bounded linear operators the ITPA of the Heisenberg template is naturally embedded.

Revisiting the case of QLGA, a subclass of QCA studied in [35], we found that the definition of QLGA in [35] was explicit but restrictive in the sense of the framework and results in this paper. That is precisely because the characterizing local condition of a QLGA belongs to the Heisenberg template. A priori, it does not embody those restrictions, which are imposed through the need to conform with the QLGA definition of [35], given as a particular Schrödinger template.

Theorem 3.5, does not provide the explicit form of a Schrödinger template for a QCA given in the Heisenberg template. Nevertheless, it aids in the pursuit of devising QCA through a Heisenberg template by assuring us of the existence of the Schrödinger template. Our result extends the reach of QCA further in the realms of quantum computation, information, and simulation, and enhances their versatility as models of quantum field theories and quantum gravity.

Acknowledgments

The author would like to thank Nolan Wallach, David Meyer and members of his research group, and Jiří Lebl for productive and edifying discussions.

References

- J. von Neumann, Theory of Self-Reproducing Automata (A. Burks, ed.), University of Illinois Press, Champaign, IL, USA, 1966.
- [2] R. Feynman, Simulating physics with computers, Int. J. Theor. Phys. 21 (1982), 467–488. doi:10.1007/BF02650179.
- [3] G. Grossing and A. Zeilinger, Quantum cellular automata, Complex Systems 2 (1988), 197–208.
- [4] C. Durr, H. LeThanh, and M. Santha, A decision procedure for well-formed linear quantum cellular automata, 1996. doi:10.1007/3-540-60922-9_24.
- [5] J. Watrous, On One-Dimensional Quantum Cellular Automata, Society Press, 1995.
- [6] D. Meyer, From quantum cellular automata to quantum lattice gases, J. Stat. Phys. 85 (1996), 551–574. doi:10.1007/BF02199356.
- [7] _____, Unitarity in one dimensional nonlinear quantum cellular automata, preprint (1996). arXiv:quant-ph/9605023.
- [8] P. Dirac, The Principles of Quantum Mechanics, Clarendon Press, Oxford, 1930.
- [9] J. von Neumann, Mathematische Grundlagen der Quantenmechanik, J. Springer, Berlin, 1932.
- [10] A. Gleason, Measures on the closed subspaces of a Hilbert space, Journal of Mathematics and Mechanics 6 (1957), no. 6, 885–893.
- [11] J. von Neumann and A. Taub, Collected Works, Volume III: Rings of Operators, Pergamon Press, New York, 1961.
- [12] I. Gelfand and M. Neumark, On the imbedding of normed rings into the ring of operators in Hilbert space, Rec. Math. [Mat. Sbornik] N.S. 12(54) (1943), no. 2, 197–217.
- [13] I. Segal, Irreducible representations of operator algebras, Bull. Amer. Math. Soc. 53 (1947), no. 2, 73–88.
- [14] B. Schumacher and R. Werner, *Reversible quantum cellular automata*, preprint (2004). arXiv:quant-ph/0405174.
- [15] O. Bratelli and D. Robinson, Operator Algebras and Quantum Statistical Mechanics 1, 2nd ed., Springer, New York, 2002.
- [16] A. Guichardet, Symmetric Hilbert Spaces and Related Topics: Infinitely Divisible Positive Definite Functions. Continuous Products and Tensor Products. Gaussian and Poissonian Stochastic Processes, Lecture Notes in Mathematics, vol. 261, Springer, New York.
- [17] V. Jones and V.S. Sunder, Introduction to Subfactors, London Mathematical Society, vol. 234, Cambridge University Press, Cambridge, United Kingdom, 1997.
- [18] J. von Neumann, On infinite direct products, Composito Mathematica 6 (1939), 1–77.
- [19] D. Gross, V. Nesme, H. Vogts, and R. Werner, Index Theory of One Dimensional Quantum Walks and Cellular Automata, Comm. Math. Phys. 310 (2012), 419–454. doi:10.1007/s00220-012-1423-1.
- [20] P. Arrighi, N. Nesme, and R. Werner, One-Dimensional Quantum Cellular Automata over Finite, Unbounded Configurations, 2008. doi:10.1007/978-3-540-88282-4_8.
- [21] _____, Unitarity plus causality implies localizability., J. Comput. Syst. Sci. **77** (2011), 372–378. doi:10.1016/j.jcss.2010.05.004.
- [22] P. Arrighi, G. Di Molfetta, I. Márquez-Martín, and A. Pérez, Dirac equation as a quantum walk over the honeycomb and triangular lattices, Phys. Rev. A 97 (2018), 062111. doi:10.1103/PhysRevA.97.062111.
- [23] B. Boghosian and W. Taylor, Quantum lattice-gas model for the many-particle Schrödinger equation in d dimensions, Phys. Rev. E 57 (1998), 54 –66. doi:10.1103/PhysRevE.57.54.

- [24] P. Arrighi, G. Di Molfetta, and S. Facchini, Quantum walking in curved spacetime: discrete metric, {Quantum} 2 (2018), 84. doi:10.22331/q2018082284.
- [25] G. Di Molfetta, M. Brachet, and F. Debbasch, Quantum walks as massless Dirac fermions in curved space-time, Phys. Rev. A 88 (2013), 042301. doi:10.1103/PhysRevA.88.042301.
- [26] A. Bisio, G. D'Ariano, and A. Tosini, Quantum field as a quantum cellular automaton: The Dirac free evolution in one dimension, Ann. Phys. 354 (2015), 244–264. doi:10.1016/j.aop.2014.12.016.
- [27] A. Bisio, G. D'Ariano, P. Perinotti, and A. Tosini, *Thirring quantum cellular automaton*, Phys. Rev. A 97 (2018), 032132. doi:10.1103/PhysRevA.97.032132.
- [28] P. Arrighi and S. Martiel, Quantum causal graph dynamics, Phys. Rev. D 96 (2017), no. 2, 024026. doi:10.1103/PhysRevD.96.024026.
- [29] D. Meyer and A. Shakeel, Quantum cellular automata without particles, Phys. Rev. A 93 (2016), 012333. doi:10.1103/PhysRevA.93.012333.
- [30] D. Schlingemann, H. Vogts, and R. Werner, On the structure of Clifford quantum cellular automata, J. Math. Phys. 49 (2008), 112104. doi:10.1063/1.3005565.
- [31] J. Gütschow, S. Uphoff, R. Werner, and Z. Zimborás, Time asymptotics and entanglement generation of Clifford quantum cellular automata, J. Math. Phys. 51 (2010), no. 1, 015203. doi:10.1063/1.3278513.
- [32] P. Arrighi, An overview of Quantum Cellular Automata, preprint (2019). arXiv:quant-ph/1904.12956.
- [33] S. Jordan, K. Lee, and J. Preskill, Quantum Algorithms for Quantum Field Theories, Science 336 (2012), no. 6085, 1130–1133. doi:10.1126/science.1217069.
- [34] M. Reed and B. Simon, Methods of Modern Mathematical Physics 1. Functional Analysis, Academic Press Inc., San Diego, California, USA, 1980.
- [35] A. Shakeel and P. Love, When is a quantum cellular automaton (QCA) a quantum lattice gas automaton (QLGA)?, J. Math. Phys. 54 (2013), no. 9, 092203. doi:10.1063/1.4821640.
- [36] R. Goodman and N. Wallach, Symmetry, Representations and Invariants, Graduate Texts in Mathematics, vol. 255, Springer, New York, 2010.
- [37] _____, Representations and Invariants of the Classical Groups, Encyclopedia of mathematics and its applications, vol. 68, Cambridge University Press, Cambridge, United Kingdom, 1999.
- [38] C. Procesi, Lie Groups: An Approach through Invariants and Representations, Universitext, Springer-Verlag, New York, 2007.
- [39] J. Rotman, An Introduction to Homological Algebra, Universitext, Springer-Verlag, New York, 2009.

APPENDIX A. RELEVANT REPRESENTATION THEORY

The material on representation theory is adapted from Goodman and Wallach [36]. The following result is Scholium 3.3.2 in Goodman and Wallach [37] pg. 135.

Scholium A.1. Let $\phi \in Aut(End(V))$, where V is a finite dimensional vector space. Then there exists $G \in GL(V)$ such that $\phi(x) = gxg^{-1}$ for all $x \in End(V)$.

Let (γ, V) , (μ, W) be two representations of an associative algebra \mathcal{A} . Let $\operatorname{Hom}(V, W)$ be the space of \mathbb{C} -linear maps from V to W. Denote by $\operatorname{Hom}_{\mathcal{A}}(V, W)$ the set of all $T \in \operatorname{Hom}(V, W)$ such that $T\gamma(a) = \mu(a)T$ for all $a \in \mathcal{A}$. Such a map is called an *intertwining operator* between the two representations. Two representations (γ, V) , (μ, W) are *equivalent* if there exists an invertible intertwining operator between the two representations.

Lemma A.2 (Schur's Lemma, Lemma 4.1.4, pg. 180, Goodman and Wallach [36]). Let (γ, V) and (μ, W) be irreducible representations of an associative algebra \mathcal{A} . Assume that V and W have at most countable dimensions over \mathbb{C} . Then

dim Hom_{$$\mathcal{A}$$} $(V, W) = \begin{cases} 1, & if(\gamma, V) \cong (\mu, W) \\ 0, & otherwise \end{cases}$

Let V be a vector space. The dimension of V is not assumed to be finite. Let \mathcal{A} be a finite dimensional semisimple associative subalgebra, and let $\widehat{\mathcal{A}}$ be the set of all equivalence classes

of finite dimensional irreducible representations of \mathcal{A} . Let $\widehat{\mathcal{A}}$ be the set of all equivalence classes of finite dimensional irreducible representations of \mathcal{A} . For each $\lambda \in \widehat{\mathcal{A}}$, fix a module $(\pi^{\lambda}, F^{\lambda})$.

Define the map:

$$s_{\lambda} : \operatorname{Hom}_{\mathcal{A}}(F^{\lambda}, V) \otimes F^{\lambda} \longrightarrow V$$

$$u \otimes v \mapsto u(v)$$

$$(22)$$

Then s_{λ} is an intertwining operator with $\operatorname{Hom}_{\mathcal{A}}(F^{\lambda}, V) \otimes F^{\lambda}$ an \mathcal{A} -module under the action $a.(u \otimes v) = u \otimes (av)$ for $a \in \mathcal{A}$.

For $\lambda \in \widehat{\mathcal{A}}$, define the λ -isotypic component:

$$V_{(\lambda)}:\sum_{W\subset V:W\sim\lambda}W$$

Definition A.3. The A-module V is locally completely reducible if the cyclic A- submodule Av is finite dimensional and completely reducible for every $v \in V$.

Proposition A.4. Let V be a locally completely reducible A-module. Then the map s_{λ} gives an A-module isomorphism

$$\operatorname{Hom}_{\mathcal{A}}(F^{\lambda}, V) \otimes F^{\lambda} \cong V_{(\lambda)}$$

Furthermore,

$$V = \bigoplus_{\lambda \in \widehat{\mathcal{A}}} V_{(\lambda)} \ (Algebraic \ direct \ sum)$$

This is called the *primary decomposition* of V.

Let $\mathcal{R} \subset \operatorname{End}(V)$ be a subalgebra that acts irreducibly on V. Let $\mathcal{A} \subset \mathcal{R}$ be a subalgebra, acting locally completely reducibly on V. Let $\operatorname{Spec}(\mathcal{A})$ be the set of irreducible representations of \mathcal{A} that occur in the primary decomposition of V. For each $\lambda \in \operatorname{Spec}(\mathcal{A})$, fix a module $(\pi^{\lambda}, F^{\lambda})$.

Let $\mathcal{R}^{\mathcal{A}} = \{T \in \mathcal{R} : aT = Ta \text{ for all } a \in \mathcal{A}\}.$

Let $E^{\lambda} = \operatorname{Hom}_{\mathcal{A}}(F^{\lambda}, V)$ for $\lambda \in \operatorname{Spec}(\mathcal{A})$. Then E^{λ} is a module for $\mathcal{R}^{\mathcal{A}}$ in a natural way by left multiplication, since

$$Tu(\pi^{\lambda}(a)v) = T(a(u(v))) = a(Tu(v))$$

for $T \in \mathcal{R}^{\mathcal{A}}$, $u \in E^{\lambda}$, $a \in \mathcal{A}$, and $v \in F^{\lambda}$. As $\mathcal{R}^{\mathcal{A}}$ commutes with \mathcal{A} , there is a representation of $\mathcal{R}^{\mathcal{A}} \otimes \mathcal{A}$ on V. As a module for the algebra $\mathcal{R}^{\mathcal{A}} \otimes \mathcal{A}$ the space V decomposes, by Proposition A.4, as

$$V \cong \bigoplus_{\lambda \in \operatorname{Spec}(\mathcal{A})} E^{\lambda} \otimes F^{\lambda}$$
(23)

In Proposition A.4 an operator $T \in \mathcal{R}^{\mathcal{A}}$ acts by $T \otimes I$ on the summand of type λ . E^{λ} is called the *multiplicity space* of λ .

We state the semisimple algebra version of the Double Commutant Theorem, in Goodman and Wallach [37] pg. 137. Let V be a finite dimensional vector space.

Theorem A.5 (Double Commutant Theorem for Semisimple Algebras). Suppose $\mathcal{A} \subset$ End(V) is a semisimple subalgebra containing the identity operator. Then the algebra $\mathcal{B} =$

 $\operatorname{Comm}(\mathcal{A})$ is semisimple and $\mathcal{A} = \operatorname{Comm}(\mathcal{B})$. Furthermore, there exists an \mathcal{A} -module isomorphism:

$$S_{\mathcal{A}} : \bigoplus_{j=1}^{r} U_{j} \otimes V_{j} \longrightarrow V$$
$$\sum_{j=1}^{r} u_{j} \otimes v_{j} \mapsto \sum_{j=1}^{r} u_{j}(v_{j})$$

)

where V_j is an irreducible \mathcal{A} -module, and $U_j = \operatorname{Hom}_{\mathcal{A}}(V_j, V)$. Under this isomorphism:

$$\mathcal{A} \cong \bigoplus_{j=1}^{r} \mathbb{I}_{U_j} \otimes \operatorname{End}(V_j)$$

and:

$$\mathcal{B} \cong \bigoplus_{j=1}^r \operatorname{End}(U_j) \otimes \mathbb{I}_{V_j}$$

Related Results from Representations of Self-Adjoint Algebras. We state a few results on representations of self-adjoint algebras. Let V be a finite dimensional Hilbert space, with an inner product $\langle \cdot | \cdot \rangle$, and $\mathcal{A} \subset \operatorname{End}(V)$ a subalgebra. Let the adjoint of an algebra $\mathcal{A} \subset \operatorname{End}(V)$ be $\mathcal{A}^{\dagger} = \{A^{\dagger} : A \in \mathcal{A}\}$. An algebra \mathcal{A} is self-adjoint if $\mathcal{A} = \mathcal{A}^{\dagger}$. From the results in this section, Lemma A.6, and Proposition A.7 appear in [38] (pg. 145).

Lemma A.6. Let $\mathcal{A} \subset \operatorname{End}(V)$ be a self-adjoint subalgebra. If $W \in V$ is an \mathcal{A} -invariant subspace, then $W^{\perp} = \{v \in V : \langle v | w \rangle = 0 \forall w \in W\}$ is \mathcal{A} -invariant.

Proof. Let $w \in W$, $v \in W^{\perp}$, $A \in \mathcal{A}$. Then $A^{\dagger} \in \mathcal{A}$, which implies $A^{\dagger}w \in W \implies \langle Av|w \rangle = \langle v|A^{\dagger}w \rangle = 0 \implies Av \in W^{\perp}$.

Proposition A.7. Let $\mathcal{A} \subset \operatorname{End}(V)$ be a self-adjoint subalgebra. Then V is an orthogonal direct sum of irreducible \mathcal{A} -modules. In particular, V is a completely reducible \mathcal{A} -module.

Proof. Let $W \subset V$ be an \mathcal{A} -invariant subspace of minimal dimension. Then it is by definition irreducible. Since $\mathcal{A} = \mathcal{A}^{\dagger}$, by Lemma A.6, $V = W \oplus W^{\perp}$ is an orthogonal direct sum of \mathcal{A} -modules. The conclusion follows by induction on dimension. \Box

Corollary A.8. Let $\mathcal{A} \subset \operatorname{End}(V)$ be a self-adjoint subalgebra containing the identity operator. Then \mathcal{A} is semisimple.

Proof. By Proposition A.7, V is a completely reducible \mathcal{A} -module. Then Proposition A.4 implies that \mathcal{A} is semisimple.

Corollary A.9. Let $\mathcal{A} \subset \operatorname{End}(V)$ be a self-adjoint subalgebra. Then there exist for every $\lambda \in \widehat{\mathcal{A}}$, a set of intertwining operators $\{u_j^{\lambda}\} \subset U^{\lambda}$, such that the λ -isotypic component can be written as an orthogonal direct sum: $V^{[\lambda]} = \bigoplus_j u_j^{\lambda}(V^{\lambda})$.

Proof. By Proposition A.7, V is a completely reducible \mathcal{A} -module. Thus by Proposition (A.4), s_{λ} in (22) is an \mathcal{A} -module isomorphism:

$$s_{\lambda}: U^{\lambda} \otimes V^{\lambda} \longrightarrow V^{[\lambda]}$$

Again by Proposition A.7, the λ -isotypic component,

$$V^{[\lambda]} = \bigoplus_{W_j \subset V: W_j \sim \lambda} W_j$$

is an orthogonal direct sum. Choose u_i such that $u_i(V^{\lambda}) = W_i$.

Outer Tensor Product Representation. Let \mathcal{A}, \mathcal{B} be associative algebras. Let (σ, V) and (τ, W) be finite dimensional representations of \mathcal{A}, \mathcal{B} respectively. The *outer tensor product* representation $(\sigma \widehat{\otimes} \tau, V \otimes W)$ of the tensor product $\mathcal{A} \otimes \mathcal{B}$ is defined as follows. On a basis element $a \otimes b \in \mathcal{A} \otimes \mathcal{B}$:

$$(\sigma \widehat{\otimes} \tau)(a \otimes b) = \sigma(a) \otimes \tau(b) \tag{24}$$

Extend by linearity to a representation of $\mathcal{A} \otimes \mathcal{B}$. If (σ, V) and (τ, W) are irreducible representations, then $(\sigma \widehat{\otimes} \tau, V \otimes W)$ is an irreducible representation.

APPENDIX B. RELEVANT DIRECT LIMIT BACKGROUND

The reader is referred to Rotman [39] for background material on direct limits. Let (I, \leq) be a *directed* set, i.e., a set with a preorder \leq (partial order is a special case), with the following property: for any pair of elements $a, b \in I$, there exists an element $c \in I$ such that $a \leq c$ and $b \leq c$.

Let C be a category. Let a family of objects in C be indexed by $I: \{C_i\}, i \in I$. Let $\{f_j^i\}, i, j \in I, i \leq j$, be a directed family of morphisms, by which we mean $f_j^i \in Mor(C_i, C_j)$, such that $f_i^i = id$, and $f_k^j \circ f_j^i = f_k^i$ for $i \leq j \leq k$. Then the pair $((C_i), (f_j^i))$, abbreviated $\{C_i, f_j^i\}$ is called a direct system (over I).

Consider the following category of pairs, each such pair being a *target*: $(C, \{f^i\}_{i \in I})$, where $C \in \mathcal{C}$ and each f^i is an *insertion* morphism $f^i : C_i \to C$, such that $f^i = f^j \circ f^i_j$ for $i \leq j$.

A target $(A, \{h^i\}_{i \in I})$ is a direct limit, denoted A_i , if it satisfies the following universal property: for every target $(G, \{g^i\}_{i \in I})$ there exists a unique morphism $\phi : A \to G$, such that $g^i = \phi \circ h^i$. Direct limit may not exist, but if it does, it is unique up to unique isomorphism, i.e., if A is a direct limit, then for any other direct limit A', there exists a unique isomorphism $A' \to A$ that commutes with the insertion morphisms.

Let R be a ring.

Theorem B.1 (Proposition 5.23, Rotman [39]). The direct limit of any direct system (M_i, f_j^i) of left R- modules over a partially ordered index set I exists.

Appendix C. Relevant QCA Related Results

Hilbert space of finite, unbounded configurations as a direct limit. For any subset $D \subset \mathbb{Z}^n$, let

$$W_D = \bigotimes_{x \in D} W \; .$$

Consider the ascending chain of subsets (3). For $D_m \subset D_n$, define a linear, isometric map

$$f_{m,n}: W_{D_m} \to W_{D_n}$$

by first expressing

$$W_{D_n} = \bigotimes_{x \in D_m} W \otimes \bigotimes_{y \in D_n \setminus D_m} W.$$

and then defining $f_{m,n}$ as

$$f_{m,n}(|v\rangle) = |v\rangle \otimes \bigotimes_{y \in D_n \setminus D_m} |q\rangle, \quad \forall |v\rangle \in W_{D_m}.$$

For $D_l \subset D_m \subset D_n$, $f_{n,l} = f_{m,n} \circ f_{l,m}$. Thus $\{W_{D_m}, f_{m,n}\}$ form a direct system (see Appendix B). Then the direct limit of this system $\varinjlim W_{D_m}$ exists by Theorem B.1, with a pre-Hilbert structure induced from the inner product on W. Let the canonical insertion map for W_{D_m} be $h_m : W_{D_m} \to \varinjlim W_{D_m}$. Under h_m, W_{D_m} can be considered a subspace of $\varinjlim W_{D_m}$, so $\varinjlim W_{D_m}$ is the union of $\{W_{D_m}\}$. Thus we see that

$$\mathbb{V} = \lim_{m \to \infty} W_{D_m} \,. \tag{25}$$

The Hilbert space of finite, unbounded configurations (UFC Hilbert space), denoted by \mathcal{H} , is the completion of \mathbb{V} under the inner product norm (the pre-Hilbert structure) on \mathbb{V} induced from the inner product on W.

Remark C.1. For the direct system above, we do not need to use an ascending chain of sets (3) as the directed set on which the system is defined. We can simply use set inclusion as a partial order on finite subsets of \mathbb{Z}^n to form the direct system. This is true for all the direct systems and limits in this paper.

$\bigotimes_{x\in\mathbb{Z}^n} |q angle$ as an eigenvector of translation-invariant unitary operator.

Lemma C.2 (Lemma III.5 in [35]). An invertible and translation invariant operator M on a Hilbert space of finite configurations \mathcal{H} has $\bigotimes_{x \in \mathbb{Z}^n} |q\rangle$ as an eigenvector:

$$M\Big(\bigotimes_{x\in\mathbb{Z}^n}|q\rangle\Big)=\lambda\bigotimes_{x\in\mathbb{Z}^n}|q\rangle$$

for some $\lambda \in \mathbb{C}^{\times}$. In particular, if M is unitary and translation invariant, then $\lambda = e^{i\theta}$ for some $\theta \in \mathbb{R}$.