Partitioning Vectors into Quadruples: Worst-Case Analysis of a Matching-Based Algorithm

Annette M. C. Ficker* Thomas Erlebach† Matúš Mihalák‡ Frits C.R. Spieksma§ ${\rm July}\ 6,\ 2018$

Abstract

Consider a problem where 4k given vectors need to be partitioned into k clusters of four vectors each. A cluster of four vectors is called a quad, and the cost of a quad is the sum of the component-wise maxima of the four vectors in the quad. The problem is to partition the given 4k vectors into k quads with minimum total cost. We analyze a straightforward matching-based algorithm, and prove that this algorithm is a $\frac{3}{2}$ -approximation algorithm for this problem. We further analyze the performance of this algorithm on a hierarchy of special cases of the problem, and prove that, in one particular case, the algorithm is a $\frac{5}{4}$ -approximation algorithm. Our analysis is tight in all cases except one.

1 Introduction

Partitioning Vectors into Quadruples (PQ) is the problem of partitioning 4k given nonnegative vectors v_1, \ldots, v_{4k} , each consisting of n components, into k clusters, each containing exactly four vectors. We refer to such a cluster of four vectors as a quadruple or a quad for short. The cost of a quad $Q = \{v_{i_1}, v_{i_2}, v_{i_3}, v_{i_4}\}$ is the sum of the component-wise maxima of the four vectors in the quad. The goal of the problem is to find a partition of the 4k vectors into k quads such that the total cost of all quads is minimum.

We will analyze the following matching-based algorithm, called algorithm A, that finds a solution to problem PQ by proceeding in two phases. In the first phase, algorithm A builds a complete, edgeweighted graph G = (V, E) that has a node in V for each vector in the instance (hence |V| = 4k). The weight of an edge equals the sum of the component-wise maxima of the two vectors whose corresponding nodes span the edge. Now, algorithm A computes a minimum-cost perfect matching M in the complete graph G, yielding 2k vector pairs. Let p_1, \ldots, p_{2k} be the 2k matched vector pairs corresponding to the computed matching M.

In the second phase, algorithm A builds a complete, edge-weighted graph G' = (V', E') that has a node in V' for each vector pair p_i found in the first phase (i = 1, ..., 2k; |V'| = 2k). The weight of an edge equals the sum of the component-wise maxima of the two vector pairs whose corresponding nodes span the edge. Now, algorithm A computes a minimum-cost perfect matching

^{*}Faculty of Economics and Business, KU Leuven, Leuven, Belgium. E-mail: annette.ficker@kuleuven.be.

[†]Department of Informatics, University of Leicester, Leicester, UK. E-mail: t.erlebach@leicester.ac.uk. Supported by a study leave granted by University of Leicester.

[‡]Department of Data Science and Knowledge Engineering, Maastricht University, Maastricht, The Netherlands. E-mail: matus.mihalak@maastrichtuniversity.nl.

[§]Department of Mathematics and Computer Science, Eindhoven University of Technology, Eindhoven, The Netherlands. E-mail: f.c.r.spieksma@tue.nl.

M' in the complete graph G'. Each of the k edges of M' matches two vector pairs, which naturally induces a quad. The k quads induced by the edges of M' constitute a solution to the problem. Clearly, A is a polynomial-time algorithm. A rigorous description can be found in Section 2. It is not hard to see that algorithm A may fail to find an optimum solution for an instance of the problem, i.e., A is not exact, and we are interested in analyzing how far off algorithm A's output can be from an optimum solution.

In this paper we show that A is a $\frac{3}{2}$ -approximation algorithm for problem PQ, and that this bound is tight. We also show that algorithm A has better approximation guarantees for various special cases of problem PQ. In particular, we show that A is a $\frac{5}{4}$ -approximation algorithm for the special case of PQ where each vector is a $\{0,1\}$ -vector containing exactly two ones, and the vectors, when seen as expressing the vertex/edge incidence matrix of a graph, correspond to edges of a simple, connected graph. We give a precise overview of our results in Section 2.3.

The remainder of this section introduces some terminology and discusses related work that motivates our research. Section 2 discusses preliminaries and also states our results. The proofs of the upper bounds on the worst-case ratio of algorithm A for the problem PQ and its special cases can be found in Section 3, while Section 4 contains the lower bound results. We conclude in Section 5.

1.1 Terminology and related literature

Worst-case analysis is a well-established tool to analyze the quality of solutions found by heuristics. We refer to books by Vazirani [13] and Williamson and Shmoys [14] for a thorough introduction to the field. We use the following, standard terminology that applies to minimization problems. In the next definition, A(I) stands for the value of the solution to instance I found by algorithm A, while OPT(I) stands for the value of an optimum solution to instance I.

Definition 1. Algorithm A is an α -approximation algorithm for a minimization problem P if for every instance I of problem P: (i) algorithm A runs in polynomial-time, and (ii) $A(I) \leq \alpha \cdot OPT(I)$. We refer to α as an upper bound on the worst-case ratio of algorithm A.

Different problems in various fields are related to problem PQ, and share some of its characteristics. In addition, algorithm A can often be adjusted to work in a particular setting. We now review related literature and provide a number of such examples.

Onn and Schulman [10] consider a problem where a given set of vectors in n-dimensional space needs to be partitioned in a given number of clusters. The number of vectors in a cluster (its size) is not specified, and in addition, they assume that the objective function, which is to be maximized, is convex in the sum of the vectors in the same cluster. Their framework contains many different problems with diverse applications, and they show, for their setting, strongly-polynomial time, exact algorithms. This is in contrast to our problem which is NP-hard (cf. Section 2.1).

Another problem, distinct from, yet related to, our problem, comes from computational biology, and is described in Figuero et al. [6]. Here, a component of a vector is a 0 or a 1 or an "N". In this setting neither the size of a cluster, nor the number of clusters is fixed; the goal is to find a partition of the set of vectors into a minimum number of clusters while satisfying the condition that a pair of vectors that is in the same cluster can only differ at a component where at least one of them has the value N. They prove hardness of this problem, and analyze the approximation behavior of heuristics for this problem.

Hochbaum and Levin [7] describe a problem in the design of optical networks that is related to our special case where each vector is a $\{0,1\}$ -vector containing two ones. In essence, their problem is to cover the edges of a given bipartite graph by a minimum number of 4-cycles. They observe

that this problem is a special case of unweighted 4-set cover; they give a $(\frac{13}{10} + \epsilon)$ -approximation algorithm (using local search), and analyze the performance of a greedy algorithm for a more general version of the problem. Our problem differs from theirs in the sense that we deal with a partitioning problem, where there is a weight for each set; in addition, our problem does not necessarily have a bipartite structure, nor do our quads need to correspond to 4-cycles.

Our problem is also intimately related to a problem occurring in wafer-to-wafer yield optimization (see, e.g., Reda et al. [11] for a description). Central in this application is the production of so-called waferstacks, which can be seen as a set of superimposed wafers. In our context, a wafer can be represented by a vector. A wafer consists of many dies, each of which can be in two states: either functioning, i.e., good (which corresponds to a component in the vector with value '0'), or malfunctioning, i.e., bad (which corresponds to a component in the vector with value '1'). The quality of a waferstack is measured by simply counting the number of components that have only 0's in the wafers contained in the waferstack. The goal is to partition the set of wafers into waferstacks (clusters) such that total quality is as high as possible. In this application, however, there are different types of wafers, and a waferstack needs to consist of one wafer of each type. This would correspond to an a priori given partition of the vectors. In addition, a typical waferstack consists in practice of many, i.e., more than 4, wafers. Dokka et al. [4] analyze the worst-case behavior of different algorithms that have as a common feature solving assignment problems repeatedly. The case where there are three types of wafers, and the problem is to find waferstacks that are triples containing one wafer of each type is investigated in Dokka et al. [3]; for a particular objective function, they describe a $\frac{4}{3}$ -approximation algorithm.

A special, yet very relevant special case of our problem is one where the edges of a given graph need to be partitioned into subsets each containing four edges (see Section 2 for a precise description). Indeed, from a graph-theoretical perspective, there is quite some interest and literature in partitioning the edge-set of a graph, i.e., to find an edge-decomposition. In fact, edge-decompositions where each cluster has prescribed size have already been studied in e.g. Jünger et al. [9]. Thomassen [12] studies the existence of edge-decompositions into paths of length 4, and Barat and Gerbner [1] even study edge-decompositions where each cluster is isomorphic to a tree consisting of 4 edges.

2 Preliminaries

2.1 About problem PQ: special cases and complexity

We first observe that, for the analysis of algorithm A, we can restrict ourselves to instances of problem PQ where the 4k vectors are $\{0,1\}$ vectors. Notice that we call a vector *nonnegative* when each of its entries is nonnegative.

Lemma 2. Each instance of problem PQ with arbitrary (rational) nonnegative vectors can be reduced to an instance of problem PQ with $\{0,1\}$ vectors.

Proof. If the components of the input vectors are non-negative integers, we can reduce the problem to the problem with $\{0,1\}$ -vectors as follows: Consider any component $i, 1 \le i \le n$, and let M_i be the largest value in the i-th component of any input vector. Then replace in each vector component i by M_i components, and a vector with value x in the i-th component is replaced by a vector that has x ones followed by $M_i - x$ zeros in the M_i components that replace component i. The resulting vectors are $\{0,1\}$ -vectors, and the cost of any set of original vectors is the same as the cost of the corresponding set of modified vectors. If the input vectors have non-negative rational values, we

first multiply all vectors by the lowest common denominator of all the rational numbers to make all vector components integers, and then use the reduction described above. \Box

This shows that for a worst-case analysis of algorithm A, it is sufficient to consider $\{0,1\}$ -vectors only. Indeed, any worst-case ratio of A shown to hold for $\{0,1\}$ -vectors holds, using the argument of Lemma 2, for arbitrary rational nonnegative vectors. However, the reduction described in the proof of Lemma 2 is not polynomial. We only need the lemma for the purpose of the analysis; and of course, algorithm A can work directly with the original input vectors.

Thus, from hereon we restrict ourselves, without loss of generality, to the case of binary vectors. There are various special cases of PQ that are of independent interest. We will describe the particular special case in brackets following 'PQ'; we distinguish the following special cases.

- Problem $PQ(\#1 \in \{1,2\})$. The case where each vector contains either one or two 1's; all other components have value 0. It will turn out that, at least in terms of the worst-case behavior of algorithm A, this special case displays the same behavior as the general problem PQ.
- Problem PQ(#1=2). The case where each binary vector contains exactly two 1's. Instances of this type can be represented by a multi-graph F with n nodes, each node corresponding to a component of a vector. Each vector is then represented by an edge spanning the two nodes that correspond to components with value 1. Of course, now a quad can be seen as a set of four edges, and its cost equals the number of nodes in the subgraph induced by these four edges.
- Problem PQ(#1 = 2, distinct). The case where the graph F is a simple graph. Equivalently, this means that each vector contains exactly two 1's and the vectors are pairwise distinct.
- Problem PQ(#1 = 2, distinct, connected). We distinguish a further special case by demanding that the graph F is also connected.

Clearly, the special cases are ordered, in the sense that each next one is a special case of its predecessor.

Although our interest is on the worst-case behavior of algorithm A, it is relevant to establish the computational complexity of problem PQ. We prove that even its special case PQ(#1 = 2, distinct, connected) is NP-hard. This fact shows that no polynomial-time algorithm for problem PQ can be exact, unless P=NP.

Theorem 3. PQ(#1 = 2, distinct, connected) is NP-hard.

Proof. As noted by Hochbaum and Levin [7], the following problem was shown to be NP-complete by Holyer [8]. Given a connected, bipartite graph G = (V, E), where |E| = 4k for some $k \in \mathcal{N}$, does there exist a partition of the edge-set E such that each set is isomorphic to a cycle on 4 nodes, i.e., a C_4 ? We refer to this decision problem as EPC_4 .

Given an instance of EPC_4 , we build the following instance of PQ(#1 = 2, distinct, connected). There are 4k vectors, each of length |V|. The |V| components of each vector correspond to the nodes in V. Each edge in E gives rise to a vector whose entries are 0, except in the two components that correspond to the nodes spanning the edge; these components have value 1. This specifies all 4k vectors. The question is: does there exist a solution of this instance of PQ(#1 = 2, distinct, connected) with cost at most 4k?

We claim that an instance of EPC_4 is a yes-instance if and only if there exists a solution to PQ(#1=2, distinct, connected) with cost at most 4k. Indeed, if the instance of EPC_4 is a yes-instance, the four edges of each C_4 directly correspond to four vectors making up a quadruple with cost of 4, leading to a total cost of 4k.

Consider now a solution to PQ(#1=2, distinct, connected), i.e., a set of k quads, with total cost 4k. Since any four vectors are pairwise distinct, it follows that the four edges corresponding to each quadruple must span at least four vertices, i.e., each quadruple must have cost at least 4. And since the total cost equals 4k, it follows that each quadruple must have cost exactly 4. Finally, since the only possibility for four edges to span four nodes in a simple bipartite graph is a C_4 , it follows that a partition into C_4 's must exist.

2.2 About algorithm A: notation and properties

Recall that, in our analysis, we may assume that all vectors are $\{0,1\}$ -vectors. Let $v_i \vee v_j$ denote the vector that is the component-wise maximum of the two vectors v_i and v_j , i.e.:

$$v_i \vee v_j = (\max(v_{i,1}, v_{j,1}), \max(v_{i,2}, v_{j,2}), \dots, \max(v_{i,n}, v_{j,n})).$$

Here, $v_{i,\ell}$ denotes the ℓ -th component of vector v_i ($\ell = 1, ..., n$). We use $|v_i|$ to denote the number of ones in vector v_i ($1 \le i \le 4k$), i.e.:

$$|v_i| = \sum_{\ell=1}^n v_{i,\ell} .$$

The cost of a quad $Q = \{v_1, v_2, v_3, v_4\}$ is then $cost(Q) = |v_1 \lor v_2 \lor v_3 \lor v_4|$. For a pair $p = \{v_1, v_2\}$ of vectors, we set $cost(p) = |v_1 \lor v_2|$.

For two vectors v_i and v_j , let sav (v_i, v_j) (the "savings" made by combining v_i and v_j) denote the number of common ones in v_i and v_j , i.e.:

$$sav(v_i, v_j) = \sum_{\ell=1}^n \min(v_{i,\ell}, v_{j,\ell}).$$

If $p = \{v_1, v_2\}$ and $p' = \{v_3, v_4\}$ are pairs of vectors, we also write sav(p, p') for $sav(v_1 \lor v_2, v_3 \lor v_4)$. The following observation concerning two $\{0, 1\}$ -vectors u and v is immediate:

Observation 4. $|u| + |v| = \text{sav}(u, v) + |u \vee v|$.

Proof. Recall that we may assume that all vectors are $\{0,1\}$ -vectors (Lemma 2). Let us partition the set of components that make up the vectors u and v into four sets:

- Those with a '1' in u, and a '0' in v: say there are $k_{u,\bar{v}}$ of them.
- Those with a '0' in u, and a '1' in v: say there are $k_{\bar{u},v}$ of them.
- Those with a '1' in u, and a '1' in v: say there are $k_{u,v}$ of them.
- Those with a '0' in u, and a '0' in v: say there are $k_{\bar{u},\bar{v}}$ of them.

Obviously, since $|u| = k_{u,\bar{v}} + k_{u,v}$, $|v| = k_{\bar{u},v} + k_{u,v}$, sav $(u,v) = k_{u,v}$, and $|u \vee v| = k_{u,\bar{v}} + k_{\bar{u},v} + k_{u,v}$, the claim follows.

Let us revisit the description of Algorithm A. In the first phase, it computes a minimum-cost perfect matching M in the complete graph G on the given 4k vectors, where the weight of the edge between vectors v_i and v_j is set to $|v_i \vee v_j|$. Let p_1, \ldots, p_{2k} be the 2k matched vector pairs corresponding to the computed matching M, and let cost(M) denote the cost of the matching M. For $1 \leq i \leq 2k$, let v_i^1 and v_i^2 be the two vectors in the vector pair p_i , and let $v_i' = v_i^1 \vee v_i^2$.

In the second phase, Algorithm A computes a minimum-cost perfect matching M' in the complete graph G' on the 2k vector pairs, where the weight of the edge between pairs p_i and p_j is set to $|v_i' \vee v_j'|$. The quads corresponding to M' are output as a solution. Let cost(M') be the cost of matching M'.

Observation 5. $A(I) = \cos(M')$ and $\cos(M') \le \cos(M)$.

Lemma 6. In the first phase of algorithm A, we can equivalently set the weight of the edge between v_i and v_j to be $-\text{sav}(v_i, v_j)$. Similarly, in the second phase of algorithm A, we can set the weight of the edge between p_i and p_j to be $-\text{sav}(v'_i, v'_j)$.

Proof. For the first phase, it follows from Observation 4 that the cost of any perfect matching M can be written as:

$$cost(M) = \sum_{(v_i, v_j) \in M} |v_i \vee v_j| = \sum_{i=1}^{4k} |v_i| - \sum_{(v_i, v_j) \in M} sav(v_i, v_j).$$

Hence, finding a matching M that minimizes $\sum_{(v_i,v_j)\in M} |v_i\vee v_j|$ is equivalent to finding a matching M that minimizes $\sum_{(v_i,v_j)\in M} -\text{sav}(v_i,v_j)$.

In the second phase, the cost of any perfect matching M' is

$$cost(M') = \sum_{(v'_i, v'_j) \in M'} |v'_i \lor v'_j| = \sum_{(v'_i, v'_j) \in M'} (|v'_i| + |v'_j| - sav(v'_i, v'_j)) =
\sum_i |v'_i| - \sum_{(v'_i, v'_j) \in M'} sav(v'_i, v'_j) = cost(M) - \sum_{(v'_i, v'_j) \in M'} sav(v'_i, v'_j).$$
(1)

Therefore, finding a matching M' that minimizes $\sum_{(v_i',v_j')\in M'}|v_i'\vee v_j'|$ is equivalent to finding a matching M' that minimizes $\sum_{(v_i',v_j')\in M'}-\operatorname{sav}(v_i',v_j')$.

Let weight (M') denote the total savings of the perfect matching M', i.e.,

weight(
$$M'$$
) = $\sum_{(v'_i, v'_j) \in M'} \operatorname{sav}(v'_i, v'_j)$.

Then, it follows from Equation (1) that

$$cost(M') = cost(M) - \sum_{(v_i', v_j') \in M'} sav(v_i', v_j') = cost(M) - weight(M').$$

$$(2)$$

Observation 5 and Equation (2) imply:

Corollary 7. A(I) = cost(M) - weight(M').

In view of this corollary, it follows that if we can show that $cost(M) \leq B$ and $cost(M') \geq S$ for some bounds B and S, we can conclude that $A(I) \leq B - S$.

Two vectors u and v are *identical* when u = v, and a pair of identical vectors is called an *identical pair*. In the following we show that among the set of minimum-cost perfect matchings, there is one that contains a maximum number of identical pairs.

Lemma 8. There is a minimum-cost perfect matching in G, as well as in G', that contains a maximum number of identical pairs.

Proof. Assume that, in a minimum-cost perfect matching, there are two identical vectors u and v that are not matched to each other; instead, let u be matched to some vector a, and v be matched to some vector b. We need to prove that, when u = v, the cost of the pairs $\{u, a\}$ and $\{v, b\}$ is at least as large as the cost of the pairs $\{u, v\}$ and $\{a, b\}$.

$$|u \vee a| + |v \vee b| = \operatorname{sav}((u \vee a), (v \vee b)) + |(u \vee a) \vee (v \vee b)|$$

$$\geq |u| + |u \vee a \vee b|$$

$$\geq |u| + |a \vee b|.$$

The first equality holds by Observation 4, the first inequality follows from u = v, and the second inequality is trivial.

Thus, in the implementation of our algorithm A, we can first greedily match pairs of identical vectors as long as they exist, and then use any standard minimum-cost perfect matching algorithm to compute a perfect matching of the remaining vectors.

2.3 Our results

In this paper, we show the following bounds on the worst-case ratio of algorithm A (see Table 1 for a summary).

Theorem 9. Algorithm A is a $\frac{3}{2}$ -approximation algorithm for problem PQ, and this bound is tight.

Theorem 10. Algorithm A is a $\frac{3}{2}$ -approximation algorithm for problem $PQ(\#1 \in \{1,2\})$, and this bound is tight.

Theorem 11. Algorithm A is a $\frac{4}{3}$ -approximation algorithm for problem PQ(#1=2), and this bound is tight.

Theorem 12. Algorithm A is a $\frac{13}{10}$ -approximation algorithm for problem PQ(#1=2, distinct), and its worst-case ratio is at least $\frac{5}{4}$.

Theorem 13. Algorithm A is a $\frac{5}{4}$ -approximation algorithm for problem $PQ(\#1=2,\ distinct,\ connected)$, and this bound is tight.

Proofs of the theorems are presented in the following sections: The upper bound proofs (Lemmas 14-20) are given in Section 3, and the lower bound results (Observations 22-24) in Section 4. As an aside, in Section 4.4 we also give instances that show that the worst-case ratio of a natural greedy algorithm is worse than the worst-case ratio of algorithm A, both for problem PQ and for problem PQ(#1 = 2, distinct, connected).

Problem name	Lower Bound	Upper Bound
PQ	$\frac{3}{2}$	$\frac{3}{2}$ (Lemma 14)
$PQ(\#1 \in \{1,2\})$	$\frac{3}{2}$ (Observation 22)	$\frac{3}{2}$
PQ(#1=2)	$\frac{4}{3}$ (Observation 23)	$\frac{4}{3}$ (Lemma 18)
PQ(#1 = 2, distinct)	$\frac{5}{4}$	$\frac{13}{10}$ (Lemma 19)
PQ(#1 = 2, distinct, connected)	$\frac{5}{4}$ (Observation 24)	$\frac{5}{4}$ (Lemma 20)

Table 1: Overview of bounds on the worst-case ratio of algorithm A

3 Upper bound proofs

In this section, we prove the upper bounds for the worst-case ratios of algorithm A for problem PQ and its special cases. In Section 3.1 we prove the upper bound $\frac{3}{2}$ for the worst-case ratio of Problem PQ, in Section 3.2 we prove the upper bound $\frac{4}{3}$ for the worst-case ratio of Problem PQ(#1 = 2), and in Section 3.3 we prove the upper bound $\frac{13}{10}$ for the worst-case ratio of Problem PQ(#1 = 2, distinct). Finally, in Section 3.4 we prove the upper bound $\frac{5}{4}$ for the worst-case ratio of Problem PQ(#1 = 2, distinct, connected).

3.1 Approximation analysis for PQ

Lemma 14. The worst-case ratio of algorithm A for PQ is at most $\frac{3}{2}$.

Proof. We use the terminology from Section 2, where M refers to the minimum-cost perfect matching found by A in the first phase based on the costs $|v_i \vee v_j|$, and M' refers to the maximum-weight perfect matching found in the second phase based on the savings $\text{sav}(v_i', v_j')$. As described in Corollary 7, we can express the cost of the solution found by algorithm A as follows:

$$A(I) = cost(M) - weight(M').$$

Consider the quads in an optimum solution. By specifying two vector pairs in each quad from the optimum solution, we obtain a matching \hat{M} that we can compare to the matching M found by A. Clearly, by the optimality of the first phase's matching of algorithm A, we have:

$$cost(M) \le cost(\hat{M})$$
, for any possible choice of \hat{M} . (3)

Further, we will identify potential matches between vector pairs in M with corresponding savings that algorithm A could make in the second phase. These potential matches are represented as edges in an auxiliary graph H whose vertex set is the set of vector pairs resulting from the algorithm's matching of the first phase. Thus we will construct a graph $H = (V', E_1 \cup E_2)$ where the edge-sets E_1 and E_2 will be described in detail. The weight of each edge e in the graph H, called w(e), represents the savings that algorithm A would realize in the second phase if it were to match the vector pairs that are the endpoints of e. The graph H can be seen as a "proxy" for the graph G' that is used in the second phase of algorithm A; it will allow us to argue that a certain amount of savings is guaranteed to exist in an optimum second phase matching.

The graph H will be bipartite and have maximum degree 2, implying that every cycle in H must have even length. Furthermore, each edge of H will connect two vertices with the same degree. We use E_1 to denote the edges in H whose both endpoints have degree one, with S_1 their total weight, i.e., $S_1 = \sum_{e \in E_1} w(e)$. We use E_2 to denote the remaining edges in H, with S_2 representing their total weight, i.e., $S_2 = \sum_{e \in E_2} w(e)$. Our construction will ensure that E_2 is a collection of even-length cycles.

Claim 15. Consider an undirected, edge-weighted graph H that is bipartite, and has maximum degree 2. Further, assume that every edge connects two vertices of the same degree. Let S_1 (S_2) be the total weight of edges between nodes with degree 1 (degree 2), and let M_H be a maximum-weight matching in H. Then

weight
$$(M_H) \geq S_1 + \frac{1}{2}S_2$$
.

Proof. The claim follows because there exists a matching in H with that weight that can be obtained by taking all edges from E_1 and partitioning E_2 into two matchings and taking the one with maximum weight.

We claim that the matching M' that algorithm A finds in G' in the second phase has total savings at least $S_1 + \frac{1}{2}S_2$. This follows from Claim 15 because H is a subgraph of G' and we can obtain a perfect matching of G' by taking a maximum-weight matching of H and matching any remaining vector pairs arbitrarily. Thus, we get:

$$weight(M') \ge S_1 + \frac{1}{2}S_2. \tag{4}$$

Inequalities (3) and (4) imply:

$$A(I) = \operatorname{cost}(M) - \operatorname{weight}(M') \le \operatorname{cost}(\hat{M}) - (S_1 + \frac{1}{2}S_2).$$
 (5)

Consider the quantity $\operatorname{cost}(\hat{M}) - (S_1 + \frac{1}{2}S_2)$, which - according to (5) - serves as an upper bound for the cost of the solution found by algorithm A. Informally speaking, we are going to distribute this quantity over the quads from the optimum solution: for each quad Q in the optimum solution, we will define its corresponding "share" of $\operatorname{cost}(\hat{M}) - (S_1 + \frac{1}{2}S_2)$ by ϕ_Q ; we will refer to ϕ_Q as the contribution reserved for Q. This contribution ϕ_Q consists of terms reflecting the contribution to \hat{M} , and terms reflecting the contribution to the total savings $S_1 + \frac{1}{2}S_2$. We will show that the choice of ϕ_Q satisfies, for each Q from the optimum solution:

$$\phi_Q \le \frac{3}{2} \mathrm{cost}(Q). \tag{6}$$

This leads to the following:

$$A(I) \le \cot(\hat{M}) - (S_1 + \frac{1}{2}S_2) = \sum_{Q} \phi_Q \le \sum_{Q} \frac{3}{2} \cot(Q) = \frac{3}{2} OPT(I).$$

The remainder of the proof is devoted to proving the above relationship. Thus, correctness hinges upon proving that

- (i) the graph H that we will construct is bipartite, and has maximum degree 2, and that our choice of ϕ_Q satisfies
- (ii) $\phi_Q \leq \frac{3}{2} \text{cost}(Q)$ for each Q from the optimum solution, and

(iii)
$$\sum_{Q} \phi_{Q} = \cos(\hat{M}) - (S_1 + \frac{1}{2}S_2).$$

We now prove (i), (ii), and (iii).

Proving that the graph H is bipartite, and has maximum degree 2

Definition 16. A lucky pair is a pair of vectors that are in the same quad in the optimum solution and that are matched by algorithm A in the first phase.

Let us first construct the edge-set E_1 of the graph H. Consider an optimal quad Q that contains two lucky pairs p_1 and p_2 . By definition, p_1 and p_2 correspond to nodes in H, and we add the edge $e = (p_1, p_2)$ to E_1 , and we set its weight equal to the corresponding savings: $w(e) = \text{sav}(p_1, p_2)$. Notice that neither node p_1 nor p_2 will be incident to any other edge in H.

Let us now proceed with the edge-set E_2 of the graph H. The edge set E_2 of H will be constructed as follows. Consider an auxiliary multi-graph K whose vertices are the quads of the optimal solution that contain at most one lucky pair. For every vector pair (v_1, v_2) that is matched by the algorithm in the first phase, add an edge (Q_1, Q_2) to K, where Q_i is the optimal quad that contains the vector v_i for i = 1, 2. We say that this edge (Q_1, Q_2) corresponds to the pair (v_1, v_2) . If (v_1, v_2) is a lucky pair, the edge added to K is a self-loop at the node corresponding to the quad that contains v_1 and v_2 . Every edge of K corresponds to a vertex of the auxiliary graph H, as the vertices of H are the pairs of vectors matched by the algorithm in the first phase. Note that each vertex in K has degree four, where a self-loop contributes 2 to the degree of the vertex to which it is attached. As every vertex of K has even degree, every connected component of K admits an Eulerian cycle. Note that each Eulerian cycle of a connected component of K has an even number of edges as each vertex in the component has degree 4 and the number of edges in a multi-graph with self-loops is equal to half the sum of the vertex degrees.

Pick an arbitrary Eulerian cycle (possibly including self-loops) in each component of K. We will use these Eulerian cycles to determine edges to be added to E_2 in H in such a way that H is bipartite and has maximum degree 2. Orient each Eulerian cycle in an arbitrary way into a directed cycle. For every pair of consecutive edges (Q_i, Q_{i+1}) and (Q_{i+1}, Q_{i+2}) on such a cycle, where (Q_i, Q_{i+1}) corresponds to $p_1 = \{v_i, v_{i+1}\}$ and (Q_{i+1}, Q_{i+2}) to $p_2 = \{v'_{i+1}, v_{i+2}\}$, add the edge $e = (p_1, p_2)$ to H, and set its weight to $w(e) = \text{sav}(p_1, p_2)$. The edge (p_1, p_2) is considered to be added for the optimal quad Q_{i+1} , and its savings are used only in the analysis of that optimal quad. See Figure 1 for an example of graph K and H.


Claim 17. The graph H is bipartite, and has maximum degree 2.

Proof. Observe that, for every component of K, the edges added to H form a single cycle consisting of an even number of edges. This is true because the vertices of H correspond to edges of K and the edges of H connect consecutive edges of an Eulerian cycle of K, which has even length. \square


Clearly, when building H as described above, different choices can be made, since there might be different Eulerian cycles possible in K in the case of a quad not containing a lucky pair. We need to be explicit about these different possibilities. Indeed, consider any quad $Q = \{v_1, v_2, v_3, v_4\}$ that does not contain a lucky pair. Let p_i be the vector pair containing v_i that was matched by the algorithm in the first phase, for $1 \le i \le 4$. Let (p_1, p_3) and (p_2, p_4) be the edges added to H, and observe that they lie on a single even-length cycle C in H (namely, the cycle created in H from the Eulerian cycle of the component of K that contains Q). Assume, without loss of generality, that after removing the edges (p_1, p_3) and (p_2, p_4) from H, the cycle C splits into two paths in H, one between node p_1 and node p_2 , and one between node p_3 and node p_4 . Define $\{\{p_1, p_2\}, \{p_3, p_4\}\}$ to be the good partition associated with Q.

Observe now that replacing the edges (p_1, p_3) and (p_2, p_4) in H by the edges (p_1, p_4) and (p_2, p_3) in H replaces the cycle C by another cycle C' consisting of the same number of edges. This means that any one of the two possible combinations of two independent edges between a vertex on one

OPT quads and first phase matching M:



Graph K:



Two possible graphs H:




Figure 1: Example of graphs K and H with p_0 and p_5 as lucky pairs

side of the good partition and a vertex on the other side of the good partition can be chosen for inclusion in E_2 , while maintaining the property that E_2 consists of even-length cycles. For example, in Figure 1, we show the two possible cycles of even length for graph H.

Observe that the discussion following Claim 17 has identified a collection of graphs H, each satisfying the conditions of Claim 17.

Proving that $\phi_Q \leq \frac{3}{2} \mathrm{cost}(Q)$ for each quad Q Now we are ready to analyse the contribution to ϕ of each quad Q from the optimum solution. There are three types of quads in an optimum solution:

- those quads that contain two lucky pairs; let us refer to this set of quads as O_2 ,
- those quads that contain one lucky pair; let us refer to this set of quads as O_1 ,
- those quads that contain no lucky pairs; let us refer to this set of quads as O_0 .

Let us first consider the quads from the set O_2 . Let $Q \in O_2$ equal $\{p_1, p_2\}$. We set

$$\phi_Q = \cos(p_1) + \cos(p_2) - \sin(p_1, p_2). \tag{7}$$

Since $cost(Q) = cost(p_1) + cost(p_2) - sav(p_1, p_2) = \phi_Q$, it trivially follows that:

$$\phi_Q \le \frac{3}{2} \text{cost}(Q) \text{ for each quad } Q \in O_2.$$
 (8)

Next, we consider the quads containing a single lucky pair, i.e., the quads from O_1 . Thus, with $Q = \{u_1, u_2, v_1, v_2\}$, let the lucky pair be $p = \{u_1, u_2\}$, and let $p_1 = \{v_1, w_1\}$ and $p_2 = \{v_2, w_2\}$ be the other nodes in H that contain the vectors from quad Q. Clearly, the edge-set E_2 contains the edge (p, p_1) as well as (p, p_2) , with weights respectively $\text{sav}(p, p_1)$ and $\text{sav}(p, p_2)$. Observe that there are no other edges in H incident to node p.

For each $Q = \{u_1, u_2, v_1, v_2\} \in O_1$, we set:

$$\phi_Q = \cos(p) + |v_1 \vee v_2| - \frac{1}{2}(\sin(p, p_1) + \sin(p, p_2)). \tag{9}$$

The cost of Q is:

$$cost(Q) = cost(p) + |v_1 \vee v_2| - sav(p, v_1 \vee v_2).$$
(10)

Notice that $cost(Q) \ge cost(p)$ and $cost(Q) \ge |v_1 \lor v_2|$, and therefore:

$$cost(Q) \ge \frac{1}{2}(cost(p) + |v_1 \lor v_2|). \tag{11}$$

Further, we have:

$$sav(p, v_1 \lor v_2) \le sav(p, v_1) + sav(p, v_2) \le sav(p, p_1) + sav(p, p_2).$$
 (12)

Combining (9), (11), and (12) gives, for each quad $Q \in O_1$:

$$\phi_{Q} = \cot(p) + |v_{1} \vee v_{2}| - \frac{1}{2}(\operatorname{sav}(p, p_{1}) + \operatorname{sav}(p, p_{2}))$$

$$\leq \cot(p) + |v_{1} \vee v_{2}| - \frac{1}{2}\operatorname{sav}(p, v_{1} \vee v_{2})$$

$$= \frac{1}{2}(\cot(p) + |v_{1} \vee v_{2}|) + \frac{1}{2}(\cot(p) + |v_{1} \vee v_{2}| - \operatorname{sav}(p, v_{1} \vee v_{2}))$$

$$\leq \cot(Q) + \frac{1}{2} \cdot \cot(Q) = \frac{3}{2} \cdot \cot(Q). \tag{13}$$

Now, consider a quad $Q = \{v_1, v_2, v_3, v_4\}$ from O_0 , i.e., a quad with no lucky pairs. As mentioned before, the term ϕ_Q consists of terms reflecting the contribution to \hat{M} , and terms reflecting the contribution to the savings. As there are three ways to choose two vector pairs from Q, the contribution to the matching \hat{M} can be realized by any of the three expressions $|v_1 \vee v_2| + |v_3 \vee v_4|$, $|v_1 \vee v_3| + |v_2 \vee v_4|$, $|v_1 \vee v_4| + |v_2 \vee v_3|$.

Let p_i be the vector pair matched by the algorithm in the first phase that contains v_i , for $1 \le i \le 4$. Assume that $\{\{p_1, p_2\}, \{p_3, p_4\}\}$ is the good partition associated with Q; it follows that we can choose for inclusion in H either the edges (p_1, p_3) and (p_2, p_4) or the edges (p_1, p_4) and (p_2, p_3) . Depending on which pair of edges we choose for inclusion in H, the term in ϕ_Q that reflects the contribution to the savings equals either $\frac{1}{2}(\text{sav}(p_1, p_3) + \text{sav}(p_2, p_4))$ or $\frac{1}{2}(\text{sav}(p_1, p_4) + \text{sav}(p_2, p_3))$. Observe that we have:

$$sav(p_1, p_3) + sav(p_2, p_4) \ge sav(v_1, v_3) + sav(v_2, v_4)$$
(14)

$$\operatorname{sav}(p_1, p_4) + \operatorname{sav}(p_2, p_3) \ge \operatorname{sav}(v_1, v_4) + \operatorname{sav}(v_2, v_3). \tag{15}$$

Summarizing, depending on which pairs of vectors from Q we put in \hat{M} (there are three possibilities), and which two edges we add to H (there are two possibilities), we can get (using (14)-(15)) any of the following six bounds on the contribution ϕ_Q of Q:

$$\phi_Q \leq |v_1 \vee v_2| + |v_3 \vee v_4| - \frac{1}{2} (\operatorname{sav}(v_1, v_3) + \operatorname{sav}(v_2, v_4)), \tag{16}$$

$$\phi_Q \leq |v_1 \vee v_2| + |v_3 \vee v_4| - \frac{1}{2} (\operatorname{sav}(v_1, v_4) + \operatorname{sav}(v_2, v_3)), \tag{17}$$

$$\phi_Q \leq |v_1 \vee v_3| + |v_2 \vee v_4| - \frac{1}{2} (\operatorname{sav}(v_1, v_3) + \operatorname{sav}(v_2, v_4)), \tag{18}$$

$$\phi_Q \leq |v_1 \vee v_3| + |v_2 \vee v_4| - \frac{1}{2} (\operatorname{sav}(v_1, v_4) + \operatorname{sav}(v_2, v_3)), \tag{19}$$

$$\phi_Q \leq |v_1 \vee v_4| + |v_2 \vee v_3| - \frac{1}{2} (\operatorname{sav}(v_1, v_3) + \operatorname{sav}(v_2, v_4)), \tag{20}$$

$$\phi_Q \leq |v_1 \vee v_4| + |v_2 \vee v_3| - \frac{1}{2} (\operatorname{sav}(v_1, v_4) + \operatorname{sav}(v_2, v_3)). \tag{21}$$

We can choose the pairs of vectors for \hat{M} and the two edges we add to H for Q in such a way that the smallest of these six bounds becomes an upper bound on the contribution reserved for Q. For that choice, all right-hand sides (16)-(21) are upper bounds on the contribution ϕ_Q .

Since we may assume that each vector v_i is a $\{0,1\}$ -vector, and since the right-hand sides of (16)-(21) involve the four vectors of quad Q, there are $2^4 = 16$ possible configurations for the 4 values of a particular component in the four vectors. Thus, for each $j = 0, 1, \ldots, 15$, we can write its binary expansion as $j = 8b_{j,1} + 4b_{j,2} + 2b_{j,3} + b_{j,4}$. We denote by n_j the number of components of the vectors of the quad whose values equal the binary expansion of j, i.e., the number of components r with $v_{1,r} = b_{j,1}$, $v_{2,r} = b_{j,2}$, $v_{3,r} = b_{j,3}$, $v_{4,r} = b_{j,4}$.

We can now express each of the relevant quantities as linear functions of the n_i :

$$cost(Q) = n_1 + n_2 + \dots + n_{15}, \tag{22}$$

$$|v_1 \lor v_2| = n_4 + n_5 + n_6 + n_7 + n_8 + n_9 + n_{10} + n_{11} + n_{12} + n_{13} + n_{14} + n_{15},$$

$$+ n_{11} + n_{12} + n_{13} + n_{14} + n_{15},$$

$$|v_1 \lor v_3| = n_2 + n_3 + n_6 + n_7 + n_8 + n_9 + n_{10}$$
(23)

$$+ n_{11} + n_{12} + n_{13} + n_{14} + n_{15}, (24)$$

$$|v_1 \lor v_4| = n_1 + n_3 + n_5 + n_7 + n_8 + n_9 + n_{10} + n_{11} + n_{12} + n_{13} + n_{14} + n_{15},$$
(25)

$$|v_2 \vee v_3| = n_2 + n_3 + n_4 + n_5 + n_6 + n_7 + n_{10} + n_{11} + n_{12} + n_{13} + n_{14} + n_{15},$$
(26)

$$|v_2 \vee v_4| = n_1 + n_3 + n_4 + n_5 + n_6 + n_7 + n_9 + n_{11} + n_{12} + n_{13} + n_{14} + n_{15},$$
(27)

$$|v_3 \lor v_4| = n_1 + n_2 + n_3 + n_5 + n_6 + n_7 + n_9 + n_{10} + n_{11} + n_{13} + n_{14} + n_{15},$$
(28)

$$sav(v_1, v_3) = n_{10} + n_{11} + n_{14} + n_{15}, (29)$$

$$sav(v_2, v_4) = n_5 + n_7 + n_{13} + n_{15}, (30)$$

$$sav(v_1, v_4) = n_9 + n_{11} + n_{13} + n_{15}, (31)$$

$$sav(v_2, v_3) = n_6 + n_7 + n_{14} + n_{15}. (32)$$

Using the identities in (22)–(32), we can write the inequalities (16), (19), and (21) as follows:

$$\phi_{Q} \leq n_{1} + n_{2} + n_{3} + n_{4} + \frac{3}{2}n_{5} + 2n_{6} + \frac{3}{2}n_{7} + n_{8}
+ 2n_{9} + \frac{3}{2}n_{10} + \frac{3}{2}n_{11} + n_{12} + \frac{3}{2}n_{13} + \frac{3}{2}n_{14} + n_{15},$$

$$\phi_{Q} \leq n_{1} + n_{2} + 2n_{3} + n_{4} + n_{5} + \frac{3}{2}n_{6} + \frac{3}{2}n_{7} + n_{8}
+ \frac{3}{2}n_{9} + n_{10} + \frac{3}{2}n_{11} + 2n_{12} + \frac{3}{2}n_{13} + \frac{3}{2}n_{14} + n_{15},$$

$$\phi_{Q} \leq n_{1} + n_{2} + 2n_{3} + n_{4} + 2n_{5} + \frac{1}{2}n_{6} + \frac{3}{2}n_{7} + n_{8}
+ \frac{1}{2}n_{9} + 2n_{10} + \frac{3}{2}n_{11} + 2n_{12} + \frac{3}{2}n_{13} + \frac{3}{2}n_{14} + n_{15}.$$
(33)

Multiplying (33) by $\frac{1}{2}$, (34) by $\frac{1}{4}$, and (35) by $\frac{1}{4}$, and then adding the three inequalities, we get, for each quad $Q \in O_0$:

$$\phi_{Q} \leq n_{1} + n_{2} + \frac{3}{2}n_{3} + n_{4} + \frac{3}{2}n_{5} + \frac{3}{2}n_{6} + \frac{3}{2}n_{7} + n_{8}$$

$$+ \frac{3}{2}n_{9} + \frac{3}{2}n_{10} + \frac{3}{2}n_{11} + \frac{3}{2}n_{12} + \frac{3}{2}n_{13} + \frac{3}{2}n_{14} + n_{15}$$

$$= \frac{3}{2} \cdot \sum_{j=1}^{15} n_{j} - \frac{1}{2}n_{1} - \frac{1}{2}n_{2} - \frac{1}{2}n_{4} - \frac{1}{2}n_{8} - \frac{1}{2}n_{15}$$

$$= \frac{3}{2} \cdot \cot(Q) - \frac{1}{2}n_{1} - \frac{1}{2}n_{2} - \frac{1}{2}n_{4} - \frac{1}{2}n_{8} - \frac{1}{2}n_{15}$$

$$\leq \frac{3}{2} \cdot \cot(Q). \tag{36}$$

Thus, (8), (13), and (36) show that indeed:

 $\phi_Q \leq \frac{3}{2} \text{cost}(Q)$ for each Q from the optimum solution.

Proving that $\sum_{Q} \phi_{Q} = \mathbf{cost}(\hat{M}) - (S_{1} + \frac{1}{2}S_{2})$ Since a quad contains either zero, one, or two lucky pairs, it follows that the expressions for ϕ_{Q} given in (7), (9), (16)-(21) contain the terms that jointly sum up to $\mathbf{cost}(\hat{M})$. Additionally, it is not difficult to verify that the construction of the graph H is such that the savings on the edges of H (as defined in (7), (9), (16)-(21)) sum up to $S_{1} + \frac{1}{2}S_{2}$.

The proof is complete. \Box

3.2 Approximation analysis for PQ(#1 = 2)

Lemma 18. The worst-case ratio of algorithm A for PQ(#1=2) is at most $\frac{4}{3}$.

Proof. Recall that an instance of PQ(#1=2) is a multi-graph G with 4k edges. We assume that the algorithm chooses in the first phase a matching that matches as many pairs of duplicates as possible (see Lemma 8).

Our analysis follows the structure of the proof of Lemma 14, but we obtain a better bound on the ratio between ϕ_Q and $\cos(Q)$ by exploiting the restricted set of configurations that are possible for a quad Q in the optimal solution if every vector in the given instance of the problem has exactly two ones.

By specifying two edge pairs in each quad from the optimal solution, we obtain again a matching \hat{M} whose cost is an upper bound on the cost of the matching M computed by the algorithm in the first phase, i.e. (cf. (3)):

$$cost(M) \le cost(\hat{M})$$
, for any possible choice of \hat{M} . (37)

Furthermore, we again construct an auxiliary graph $H = (V', E_1 \cup E_2)$ that represents potential matches between vector pairs in M with corresponding savings that algorithm A could make in the second phase. The weight w(e) of each edge e in the graph H represents the savings that algorithm A would realize in the second phase if it were to match the vector pairs that are the endpoints of e. The graph H is again bipartite (we will use the concept of good partitions introduced in the proof of Lemma 14 to ensure this) and has maximum degree 2, and the edges in E_1 connect vertices of degree 1 and the edges in E_2 connect vertices of degree 2. Letting S_1 denote the total weight of E_1 , and S_2 the total weight of E_2 , it follows from Claim 15 that the maximum-weight matching M_H in H satisfies weight M_H in M_H satisfies weight M_H and M_H finds in the second phase satisfies:

$$weight(M') \ge S_1 + \frac{1}{2}S_2. \tag{38}$$

Thus, we again have the relationship

$$A(I) \le \cot(\hat{M}) - (S_1 + \frac{1}{2}S_2)$$

and distribute the value on the right-hand side over the quads of the optimal solution, with each quad Q receiving a share ϕ_Q that is referred to as the contribution reserved for Q. We will show that $\phi_Q \leq \frac{4}{3} \text{cost}(Q)$ holds for all quads Q of the optimal solution, implying that:

$$A(I) \le \cot(\hat{M}) - (S_1 + \frac{1}{2}S_2) = \sum_{Q} \phi_Q \le \sum_{Q} \frac{4}{3}\cot(Q) = \frac{4}{3}OPT(I)$$

Thus, correctness hinges upon proving that

- (i) $\phi_Q \leq \frac{4}{3} \text{cost}(Q)$ for each Q from the optimum solution,
- (ii) $\sum_{Q} \phi_{Q} = \cos(\hat{M}) (S_{1} + \frac{1}{2}S_{2})$, and
- (iii) the graph H that we construct is bipartite and has maximum degree 2.

Proving that $\phi_Q \leq \frac{4}{3}\mathbf{cost}(Q)$ for each Q from the optimum solution Note that a quad contains four edges of multi-graph G and its cost equals the number of vertices in the subgraph induced by these four edges. Thus, in this proof we write quad Q from the optimum solution as a set of four edges of G, i.e., $Q = (e_1, e_2, e_3, e_4)$. We distinguish the following cases for the quads depending on the value of $\mathbf{cost}(Q)$:

• cost(Q) = 2. This means all four edges are identical, see Figure 2. As we may assume that the algorithm matches as many duplicate vectors (edges of G) as possible (Lemma 8), we can say that the algorithm matches $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ in the first phase.




Figure 2: cost(Q) = 2; 1 and 2 are nodes in G

We select both p_1 and p_2 to be part of matching \hat{M} , with total cost 2+2=4. If the algorithm, in the second phase, matches p_1 and p_2 , the algorithm will make a saving of 2. Thus, we add edge (p_1, p_2) with savings 2 to edge-set E_1 in H, as both p_1 and p_2 do not appear in any other quads. Hence,

$$\phi_Q = \cos(p_1) + \cos(p_2) - \sin(p_1, p_2) = 2 < \frac{4}{3} \cos(Q).$$

• cost(Q) = 3. The edges in G belonging to Q can have one of the following three structures, see Figure 3:




Figure 3: Quads with cost(Q) = 3; 1, 2 and 3 are nodes in G

(i) If quad Q contains $e_1 = (1,2)$, $e_2 = (1,2)$, $e_3 = (2,3)$ and $e_4 = (2,3)$, we select both $p_1 = \{(1,2),(1,2)\}$ and $p_2 = \{(2,3),(2,3)\}$ to be part of matching \hat{M} , with a total cost of 2+2=4. Furthermore, we add the edge (p_1,p_2) with savings 1 to the edge-set E_1 of H. Hence,

$$\phi_Q \le \cos(p_1) + \cos(p_2) - \sin(p_1, p_2) = 3 < \frac{4}{3} \cos(Q).$$

(ii) If quad Q contains $e_1 = (1,2)$, $e_2 = (1,2)$, $e_3 = (1,2)$ and $e_4 = (2,3)$, we may assume that the algorithm has matched $p_1 = \{(1,2), (1,2)\}$ in the first phase. Furthermore, we select p_1 and $p_2 = \{e_3, e_4\}$ to be part of matching \hat{M} , with total cost 2 + 3 = 5.

In a worst case scenario, p_2 is not contained in M, hence the algorithm has matched e_3 to another vector (edge in G), say x_3 , and e_4 to say x_4 . We define $p_3 = \{e_3, x_3\}$ and $p_4 = \{e_4, x_4\}$, which are nodes in the auxiliary graph H. We add edges (p_1, p_3) and (p_1, p_4) to edge set E_2 , with total savings at least 2 + 1 = 3. Hence,

$$\phi_Q = \cos(p_1) + \cos(p_2) - \frac{1}{2} \left(\sin(p_1, p_3) + \sin(p_1, p_4) \right) \le 5 - 1.5 < \frac{4}{3} \operatorname{cost}(Q).$$

(iii) If quad Q contains $e_1 = (1, 2)$, $e_2 = (1, 2)$, $e_3 = (1, 3)$ and $e_4 = (2, 3)$, we use a similar argument as above. We may assume that the algorithm has matched $p_1 = \{(1, 2), (1, 2)\}$ in the first phase and select p_1 and $p_2 = \{e_3, e_4\}$ to be part of matching \hat{M} , with total cost 2 + 3 = 5.

In a worst case scenario, p_2 is not contained in M, hence the algorithm has matched e_3 to say x, and e_4 to say y. We define $p_3 = \{e_3, x\}$ and $p_4 = \{e_4, y\}$, which are nodes in auxiliary graph H. We add edges (p_1, p_3) and (p_1, p_4) to edge set E_2 , with total savings at least 1 + 1 = 2. Hence,

$$\phi_Q = \cos(p_1) + \cos(p_2) - \frac{1}{2} \left(\sin(p_1, p_3) + \sin(p_1, p_4) \right) \le 4 \le \frac{4}{3} \cot(Q).$$

• cost(Q) = 4. The edges in G belonging to Q can have one of the following six structures, see Figure 4:




Figure 4: Quads with cost(Q) = 4; 1, 2, 3 and 4 are nodes in G

(i) If quad Q contains $e_1 = (1,2)$, $e_2 = (1,2)$, $e_3 = (3,4)$ and $e_4 = (3,4)$, we select both $p_1 = \{(1,2),(1,2)\}$ and $p_2 = \{(3,4),(3,4)\}$ to be part of matching \hat{M} , with a total cost of 2+2=4. We add the edge (p_1,p_2) with savings 0 to the edge-set E_1 of H. Hence,

$$\phi_Q \le \cot(p_1) + \cot(p_2) = 4 < \frac{4}{3}\cot(Q).$$

(ii)&(iii) If the graph induced by quad Q contains two identical edges, e_1 and e_2 , and two adjacent edges, e_3 and e_4 , then select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of matching M, with a total cost of 2 + 3 = 5. We can assume that algorithm A has matched e_1 and e_2 in the first phase. In the worst case, the algorithm has matched e_3 to another edge x_3 and e_4 to another edge x_4 . Let $p_3 = (e_3, x_3)$ and $p_4 = (e_4, x_4)$, and add the edges (p_1, p_3) and (p_1, p_4) with total savings at least 1 + 0 = 1 to the edge-set E_2 of H. Hence,

$$\phi_Q \le \cos(p_1) + \cos(p_2) - \frac{1}{2} = 4.5 < \frac{4}{3} \cos(Q).$$

(iv)&(v) If the graph induced by quad Q contains two identical edges, e_1 and e_2 , and two non-adjacent edges, e_3 and e_4 , then we select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of matching \hat{M} , with a total cost of 2 + 4 = 6.

In a worst case scenario, p_2 is not contained in M, hence the algorithm has matched e_3 to another vector (edge in G), say x_3 , and e_4 to say x_4 . We define that $p_3 = \{e_3, x_3\}$ and $p_4 = \{e_4, x_4\}$, which are nodes in auxiliary graph H. We add edges (p_1, p_3) and (p_1, p_4) to edge set E_2 in H. Either e_3 is a duplicate of e_1 (case (iv)), in which case the total saving is at least 2 + 0 = 2, or both edges are adjacent to both e_1 and e_2 (case (v)), in which case the total saving is at least 1 + 1 = 2. Hence,

$$\phi_Q = \cos(p_1) + \cos(p_2) - \frac{1}{2} \left(\sin(p_1, p_3) + \sin(p_1, p_4) \right) \le 5 < \frac{4}{3} \operatorname{cost}(Q).$$

(vi) Suppose quad Q contains $e_1 = (1,2)$, $e_2 = (1,3)$, $e_3 = (2,4)$ and $e_4 = (3,4)$ (if Q contains $e_1 = (2,3)$, $e_2 = (1,3)$, $e_3 = (2,4)$ and $e_4 = (3,4)$ we use the exact same argument). We select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of matching \hat{M} , with total cost 3 + 3 = 6.

In a worst case scenario, no edge pair of G in Q is contained in M, hence we say the algorithm has matched e_i to x_i and define $p_{e_i} = \{e_i, x_i\}$ for $i \in \{1, ..., 4\}$. Note that each of the p_{e_i} is a node in auxiliary graph H. If the good partition associated with Q (as defined in the proof of Lemma 14) is $\{\{p_{e_1}, p_{e_2}\}, \{p_{e_3}, p_{e_4}\}\}$, we add edges (p_{e_1}, p_{e_3}) and (p_{e_2}, p_{e_4}) to edge-set E_2 of H, with total savings at least 1 + 1 = 2. Otherwise, we add edges (p_{e_1}, p_{e_2}) and (p_{e_3}, p_{e_4}) to edge set E_2 , with total savings at least 1 + 1 = 2. Hence,

$$\phi_Q \le \cot(p_1) + \cot(p_2) - 1 = 5 < \frac{4}{3}\cot(Q).$$

• cost(Q) = 5. If Q contains two identical edges, say e_1 and e_2 , we select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of \hat{M} , with a total cost of at most 2+4=6. We can assume that the algorithm has matched (e_1, e_2) in M. In the worst case, the algorithm has not matched e_3 to e_4 in M. Let $p_{e_3} = \{e_3, x_3\}$ and $p_{e_4} = \{e_4, x_4\}$ be the matched pairs in M that contain e_3 and e_4 , respectively. We add the edges (p_1, p_{e_3}) and (p_1, p_{e_4}) to the edge-set E_2 of H. Even without taking the savings of those edges into account, we have

$$\phi_Q \le \cos(p_1) + \cos(p_2) = 6 < \frac{4}{3}\cos(Q).$$

Assume now that Q does not contain any two identical edges. Then the graph induced by $Q = (e_1, e_2, e_3, e_4)$ must fall into one of the four following cases, see Figure 5:




Figure 5: Quads with cost(Q) = 5; 1, 2, 3, 4 and 5 are nodes in G

(i) Q is a triangle of edges e_1 , e_2 , e_3 plus a disjoint edge e_4 . We select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of \hat{M} , with a total cost of 3+4=7. In the worst case, Q does not contain a lucky pair. For $1 \le i \le 4$, let $p_{e_i} = \{e_i, x_i\}$ be the pair of matched edges in M that includes e_i . If the good partition associated with Q is $\{\{p_{e_1}, p_{e_2}\}, \{p_{e_3}, p_{e_4}\}\}$, we add the edges (p_{e_1}, p_{e_3}) and (p_{e_2}, p_{e_4}) to E_2 with savings at least 1+0=1. Otherwise, add the edges (p_{e_1}, p_{e_2}) and (p_{e_3}, p_{e_4}) to E_2 , again with savings at least 1+0=1. We have

$$\phi_Q \le \cos(p_1) + \cos(p_2) - \frac{1}{2} = 6.5 < \frac{4}{3} \cot(Q).$$

(ii) Q is a star (i.e., all four edges of Q are incident with the same same vertex). We select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of \hat{M} , with a total cost of 3+3=6. In the worst case, Q does not contain a lucky pair. For $1 \leq i \leq 4$, let $p_{e_i} = \{e_i, x_i\}$ be the pair of matched edges in M that includes e_i . If the good partition associated with Q is $\{\{p_{e_1}, p_{e_2}\}, \{p_{e_3}, p_{e_4}\}\}$, add the edges (p_{e_1}, p_{e_3}) and (p_{e_2}, p_{e_4}) to E_2 , with savings at least 1+1=2. Otherwise, add the edges (p_{e_1}, p_{e_2}) and (p_{e_3}, p_{e_4}) to E_2 , again with savings at least 1+1=2. We have

$$\phi_Q \le \cos(p_1) + \cos(p_2) - 1 = 5 < \frac{4}{3} \cos(Q).$$

(iii) Q is a path of four edges. Assume that the edges appear on the path in the order (e_1, e_2, e_3, e_4) . We select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of \hat{M} , with a total cost of 3+3=6. In the worst case, Q does not contain a lucky pair. For $1 \leq i \leq 4$, let $p_{e_i} = \{e_i, x_i\}$ be the pair of matched edges in M that includes e_i . If the good partition associated with Q is $\{\{p_{e_1}, p_{e_2}\}, \{p_{e_3}, p_{e_4}\}\}$, add the edges (p_{e_1}, p_{e_4}) and (p_{e_2}, p_{e_3}) to E_2 , with savings at least 0+1=1. Otherwise, add the edges (p_{e_1}, p_{e_2}) and (p_{e_3}, p_{e_4}) to E_2 , with savings at least 1+1=2. We have

$$\phi_Q \le \cos(p_1) + \cos(p_2) - \frac{1}{2} = 5.5 < \frac{4}{3} \cot(Q).$$

(iv) Q is a tree of diameter 3. Assume that $e_1 = (1,2)$, $e_2 = (2,3)$, $e_3 = (3,4)$, $e_4 = (3,5)$. We select $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of \hat{M} , with a total cost of 3+3=6. In the worst case, Q does not contain a lucky pair. For $1 \le i \le 4$, let $p_{e_i} = \{e_i, x_i\}$ be the pair of matched edges in M that includes e_i . If the good partition associated with Q is $\{\{p_{e_1}, p_{e_2}\}, \{p_{e_3}, p_{e_4}\}\}$, add the edges (p_{e_1}, p_{e_4}) and (p_{e_2}, p_{e_3}) to E_2 , with savings at least 0+1=1. Otherwise, add the edges (p_{e_1}, p_{e_2}) and (p_{e_3}, p_{e_4}) to E_2 , with savings at least 1+1=2. We have

$$\phi_Q \le \cos(p_1) + \cos(p_2) - \frac{1}{2} = 5.5 < \frac{4}{3} \cot(Q).$$

• $cost(Q) \ge 6$. If we partition Q into two edge pairs p_1 and p_2 for matching \hat{M} , their total cost will be at most 4 + 4 = 8. Even if the savings of the edges that we add to the edge-set E_2 of H (while taking into account the good partition associated with Q to ensure that H is bipartite, of course) are zero, we have

$$\phi_Q \le \operatorname{cost}(p_1) + \operatorname{cost}(p_2) = 8 \le \frac{4}{3} \operatorname{cost}(Q).$$

Proving that H is a simple, bipartite graph with maximum degree 2 An edge has been added to the edge-set E_1 of H when considering a quad Q only if Q is a quad with two lucky pairs, and hence those pairs indeed become vertices of degree 1 in H. If $p = (e_1, e_2)$ is a lucky pair that is in a quad Q with two other edges e_3 , e_4 that do not form a lucky pair, it becomes the endpoint of exactly two edges in E_2 : The edges (p, p') and (p, p''), where p' is the pair of edges containing e_3 that is matched in M, and p'' is the pair of edges containing e_4 that is matched in M. Finally, every edge pair $p = (e_1, e_2)$ that has been matched by algorithm A in M and is not a lucky pair becomes the endpoint of exactly two edges in E_2 : An edge (p, p') added for quad Q_1 and an edge

(p, p'') added for quad Q_2 , where Q_1 is the optimal quad containing e_1 and Q_2 the optimal quad containing e_2 . This shows that H is a graph with maximum degree 2 in which each edge connects two vertices of the same degree. Furthermore, by ensuring for each quad Q that the edges added to E_2 are compatible with the good partition associated with Q if Q does not have a lucky pair, we have ensured that H is simple and bipartite (recall that the cycles in H correspond to Eulerian cycles of the connected components of K, which have an even number of edges, as discussed in the proof of Lemma 14.

Proving that $\sum_{Q} \phi_{Q} = \cot(\hat{M}) - (S_{1} + \frac{1}{2}S_{2})$ For each quad Q we have selected two pairs of edges in Q for inclusion in \hat{M} , so the costs of those edge pairs clearly add up to $\cot(\hat{M})$. Furthermore, each edge of H was added to H by some quad Q. Furthermore, no two quads could have added the same edge to H (as H is a simple graph), so the sum of the savings of the added edges is indeed $S_{1} + \frac{1}{2}S_{2}$.

The proof is complete. \Box

3.3 Approximation analysis for PQ(#1 = 2, distinct)

Lemma 19. The worst-case ratio of algorithm A for PQ(#1=2, distinct) is at most $\frac{13}{10}$.

Proof. Recall that an instance of PQ(#1=2, distinct) is nothing else but a simple graph F with 4k edges. Note that the cost of every optimal quad is at least 4 since 4 edges in a simple graph touch at least 4 different vertices. Hence we can repeat the arguments in the proof for Lemma 18 for quads of cost 4 and higher.

Then, we get the following results:

- For cost(Q) = 4, we have $\phi_Q \le 5 = \frac{5}{4}cost(Q) < \frac{13}{10} \cdot cost(Q)$,
- For cost(Q) = 5, we have $\phi_Q \le 7 \frac{1}{2} = \frac{6.5}{5} cost(Q) = \frac{13}{10} \cdot cost(Q)$,
- For cost(Q) = 6, note that the average degree of the subgraph of G induced by Q is $\frac{8}{6} > 1$, so there must exist a vertex of degree at least 2. This means that Q contains two adjacent edges e_1 and e_2 . Denote the remaining edges by e_3 and e_4 . We can select the pairs $p_1 = \{e_1, e_2\}$ and $p_2 = \{e_3, e_4\}$ to be part of \hat{M} , giving $\phi_Q \leq cost(p_1) + cost(p_2) \leq 3 + 4 = 7 = \frac{7}{6}cost(Q) < \frac{13}{10} \cdot cost(Q)$.

• For $cost(Q) \ge 7$, we have $\phi_Q \le 8 \le \frac{8}{7} cost(Q) < \frac{13}{10} \cdot cost(Q)$.

3.4 Approximation analysis for PQ(#1 = 2, distinct, connected)

Lemma 20. Algorithm A is a $\frac{5}{4}$ -approximation algorithm for PQ(#1=2, distinct, connected).

Proof. Recall that an instance of PQ(#1=2, distinct, connected) can be viewed as a simple, connected graph F with 4k edges, and that the cost of a quad is the number of vertices spanned by the edges in the quad. Note that the cost of every optimal quad is at least 4 since 4 edges in a simple graph touch at least 4 different vertices. Hence, $OPT \ge 4k$. Furthermore, if we can show that there are z quads in the optimal solution that have cost at least 5, we get that $OPT \ge 4(k-z) + 5z = 4k + z$.

Observation 21. cost(M) = 6k.

Proof. The line graph of a connected graph with an even number of edges admits a perfect matching (Jünger et al. [9], Dong et al. [5]). Thus, the minimum-cost perfect matching M pairs adjacent edges of the graph. Hence, every pair in M has cost 3, and thus the cost of M is $2k \cdot 3 = 6k$.

Let p_1, \ldots, p_{2k} be the pairs corresponding to M. Consider the auxiliary graph H with vertex set $V' = \{p_1, \ldots, p_{2k}\}$ in which an edge is added between p_i and p_j if p_i and p_j have at least one common vertex (implying that matching p_i to p_j in the matching M' that A computes in the second phase would create a saving of at least one). Note that H is connected as F is connected. Let μ be the size of a maximum matching in H, $1 \le \mu \le k$. Note that the maximum matching of H can be extended to a perfect matching of V' that makes savings at least μ . Therefore, we have

$$A(I) \le 6k - \mu$$
.

If H contains a perfect matching, we have $\mu = k$ and hence $A(I) \leq 5k$, implying that $A(I)/OPT(I) \leq 5k/(4k) = \frac{5}{4}$. It remains to consider the case $\mu < k$.

If a maximum matching in H has size $\mu < k$, the number of unmatched vertices is $2k - 2\mu$. We will show that the optimal solution then contains at least $k - \mu$ quads with cost at least 5, and hence we have $OPT(I) \ge 4k + (k - \mu) = 5k - \mu$. Therefore,

$$\frac{A(I)}{OPT(I)} \le \frac{6k - \mu}{5k - \mu} \le \frac{5}{4},$$

where the last inequality follows because $(6k - \mu)/(5k - \mu)$ is maximized if μ takes its maximum possible value, $\mu = k$.

It remains to show that the optimal solution contains at least $k - \mu$ quads with cost at least 5. Recall that a maximum matching in H leaves $2k - 2\mu$ vertices unmatched. By the Tutte-Berge formula [2], the number of unmatched vertices of a maximum matching in H is equal to

$$\max_{X \subset V'} (\operatorname{odd}(H - X) - |X|),$$

where odd(H-X) is the number of connected components of H-X that have an odd number of vertices (H-X) is the graph that results when the nodes in X, and their incident edges, are removed from H). Hence, there exists a set $X \subseteq V'$ such that $odd(H-X) - |X| = 2k - 2\mu$. Let d = odd(H-X), and let O_1, O_2, \ldots, O_d denote the d odd components of H-X. We have

$$2k - 2\mu = d - |X|.$$

For a subgraph S of H, let $E_F(S)$ denote the set of edges of F that are contained in the edge pairs that form the vertex set of S (recall that the vertices of H are pairs of edges from F). Note that $|E_F(O_i)| \mod 4 = 2$ for $1 \le i \le d$ as O_i contains an odd number of edge pairs. Therefore, each $E_F(O_i)$ contains at least two edges that are contained in optimal quads that do not only contain edges from $E_F(O_i)$. If such a quad contains three edges from $E_F(O_i)$, note that there must be at least one other optimal quad that contains at most three edges from $E_F(O_i)$ as $(|E_F(O_i)| - 3) \mod 4 = 3$.

For each optimal quad that contains one or two edges from $E_F(O_i)$, define these one or two edges to be *special* edges. For each optimal quad that contains three edges from $E_F(O_i)$, select one of these three edges arbitrarily and define it to be a *special* edge. There are at least two special edges in each $E_F(O_i)$, $1 \le i \le d$, and hence at least 2d special edges in total. More precisely, we

refer to these special edges as the edge-set SE, and partition it into two subsets: those special edges occurring in a quad with cost 4 (the set SE4), and those special edges occurring in a quad with cost at least 5 (the set SE5). Clearly:

$$2d \le |SE4| + |SE5|. \tag{39}$$

Consider a quad with cost 4 from the optimum solution. It consists of four edges of F. Since F is a connected simple graph there are only two possible subgraphs induced by Q, as depicted in Figure 6. These four edges can be in the sets $E_F(O_i)$ for some $1 \le i \le d$, the set $E_F(X)$, and the sets $E_F(C)$ for even components C of H - X. We now define types of quads of cost 4 depending on how many edges are in which set.



Figure 6: Quads with cost(Q) = 4

Note that an edge from $E_F(O_i)$ cannot be incident to the same vertex as an edge from $E_F(O_j)$ for $j \neq i$ because otherwise H would contain an edge between O_i and O_j . Similarly, an edge from $E_F(O_i)$ cannot be incident to the same vertex as an edge from $E_F(C)$ where C is an even component of H - X. The only edges that can share endpoints with edges in $E_F(O_i)$ are those in $E_F(X)$.

We tabulate the different types of quads with cost 4 in Table 2. Thus, a quad with cost 4 with a special edge must be of type 1, 2, 3, 4 or 5. For each of these types, the number of edges from $E_F(X)$ is at least the number of special edges in the quad. Thus,

$$|E_F(X)| \ge |SE4|. \tag{40}$$

Type of	Number of edges			Cost	Number of
quad	in $E_F(O_i)$	in $E_F(X)$	in $E_F(C)$		special edges
1	3	1		4	1
2	2	2		4	2
3	1, 1	2		4	2
4	1	2	1	4	1
5	1	3		4	1

Table 2: Overview of different types of quads with cost 4, containing at least 1 edge from $E_F(O_i)$. The entry "1,1" for quad type 3 means that there is one edge from $E_F(O_i)$ and one edge from $E_F(O_{i'})$ for $i \neq i'$

Further, since $|E_F(X)| = 2|X|$, it follows from (40) and (39) that $|SE5| \ge 2d - 2|X|$. Thus, the number of quads of cost at least 5 is at least $\frac{2d-2|X|}{4} = \frac{1}{2}(d-|X|) = k - \mu$.

4 Bad instances

In this section, we give the instances that provide the lower bound results for problem PQ and its special cases, as announced in Table 1: Section 4.1 presents the instance of problem $PQ(\#1 \in$

 $\{1,2\}$), Section 4.2 the instance of problem PQ(#1=2), and Section 4.3 the instance of problem PQ(#1=2, distinct, connected). Furthermore, we illustrate in Section 4.4 that a natural greedy algorithm (that can be seen as an alternative for algorithm A) has a worst-case ratio that is worse than the worst-case ratio of algorithm A.

4.1 An instance of $PQ(\#1 \in \{1, 2\})$

Consider the instance I consisting of the following 8 vectors, v_1, \ldots, v_8 :

$$\begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix}.$$

Since each vector contains either one or two 1's, this is an instance of $PQ(\#1 \in \{1,2\})$. Clearly, the optimum solution consists of the quads $\{v_1, v_2, v_5, v_6\}$ and $\{v_3, v_4, v_7, v_8\}$, with a total cost of OPT(I) = 4; algorithm A however, may find, as an optimum matching in the first phase, the pairs $\{v_1, v_3\}$, $\{v_2, v_4\}$, $\{v_5, v_6\}$ and $\{v_7, v_8\}$, leading to a final solution with cost A(I) = 6. Thus, we arrive at the following observation.

Observation 22. For the instance depicted above, $cost(A) = \frac{3}{2}OPT$.

Theorems 9 and 10 now follow from Lemma 14 and Observation 22.

As a remark, if we would allow all-zero vectors in the input (which we do not allow in $PQ(\#1 \in \{1,2\})$), we can get an even smaller example with ratio $\frac{3}{2}$. Let I consist of the following 8 vectors:

$$\left(\begin{array}{c}1\\0\end{array}\right),\left(\begin{array}{c}1\\0\end{array}\right),\left(\begin{array}{c}1\\0\end{array}\right),\left(\begin{array}{c}0\\0\end{array}\right),\left(\begin{array}{c}0\\1\end{array}\right),\left(\begin{array}{c}0\\1\end{array}\right),\left(\begin{array}{c}0\\1\end{array}\right),\left(\begin{array}{c}0\\0\end{array}\right).$$

The optimal cost is 2; algorithm A however, may match the two all-zero vectors in the first phase and get a solution of cost 3.

4.2 An instance of PQ(#1=2)

Consider the instance I consisting of the following 8 vectors, v_1, \ldots, v_8 .

Since each vector contains two 1's, this is an instance of PQ(#1 = 2). This instance can be represented by the graph shown in Figure 7.

The optimal solution for this instance has cost 6, with the two quads

$${v_1, v_2, v_3, v_4} = {(1, 2), (1, 2), (1, 3), (2, 3)},$$

$${v_5, v_6, v_7, v_8} = {(3,4), (3,5), (4,5), (4,5)}.$$




Figure 7: An instance of PQ(#1 = 2)

The algorithm may in the first phase construct a matching with cost 10 consisting of the following pairs:

$$\{v_1, v_2\} = \{(1, 2), (1, 2)\}, \ \{v_3, v_5\} = \{(1, 3), (3, 4)\},$$

 $\{v_4, v_6\} = \{(2, 3), (3, 5)\}, \ \{v_7, v_8\} = \{(4, 5), (4, 5)\}.$

Any two pairs share at most 1 node. Hence, the total savings that can be made in the second matching are at most 2, so by Corollary 7 we have $A(I) \ge 8$. Hence, the worst-case approximation ratio of A is at least 8/6 = 4/3.

Observation 23. For the instance depicted in Figure 7, $cost(A) = \frac{4}{3}OPT$.

Theorem 11 now follows from Lemma 18 and Observation 23.

4.3 An instance of PQ(#1 = 2, distinct, connected)

Consider the instance I consisting of the following 8 vectors v_1, \ldots, v_8 .

Since each vector contains two 1's, the vectors are pairwise distinct, and the induced graph is connected, this is an instance of PQ(#1=2), distinct, connected). The instance can be represented by the graph shown in Figure 8.

The optimal solution for this instance has cost 8, with the two quads

$$\{v_1, v_2, v_3, v_4\} = \{(1, 2), (1, 3), (2, 4), (3, 4)\},\$$
$$\{v_5, v_6, v_7, v_8\} = \{(4, 5), (4, 6), (5, 7), (6, 7)\}.$$

Algorithm A may, in the first phase, construct a matching with cost 12 consisting of the following pairs:

$$\{v_1, v_2\} = \{(1, 2), (1, 3)\}, \{v_3, v_5\} = \{(2, 4), (4, 5)\},$$

$$\{v_4, v_6\} = \{(3, 4), (4, 6)\}, \{v_7, v_8\} = \{(5, 7), (6, 7)\}.$$




Figure 8: An instance of PQ(#1 = 2, distinct, connected)

Any two pairs share at most 1 node. Hence, the total savings that can be made in the second matching are at most 2, so by Corollary 7 we have $A(I) \ge 10$. Hence, the worst-case ratio of A is at least 10/8 = 5/4.

Observation 24. For the instance depicted in Figure 8, $cost(A) = \frac{5}{4}OPT$.

Theorem 13 now follows from Lemma 20 and Observation 24.

4.4 Bad instances for a natural greedy algorithm

In this section, we show that the worst-case ratio of a natural greedy algorithm is worse than the worst-case ratio of algorithm A.

An informal description of the greedy algorithm for problem PQ (and its special cases) is as follows: repeatedly select, among all possible quads, a quad with lowest cost, and remove the vectors in the selected quad from the instance; stop when no more vectors remain.

Below we present instances of problem PQ, as well as of its special case PQ(#1 = 2, distinct, connected), showing that the worst-case performance of this greedy algorithm is worse than the worst-case performance of algorithm A.

An instance of PQ Consider the following instance I of PQ consisting of the following 12 vectors, v_1, \ldots, v_{12} :

$$\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}.$$

Clearly, an optimum solution consists of the quads $\{v_1, v_4, v_5, v_6\}$, $\{v_2, v_7, v_8, v_9\}$ and $\{v_3, v_{10}, v_{11}, v_{12}\}$, with a total cost of OPT(I) = 3. The greedy algorithm however, may first select quad $\{v_1, v_2, v_3, v_4\}$ with cost 1. Then the cheapest possible quad is one of cost 2 and the greedy algorithm may select $\{v_7, v_8, v_{11}, v_{12}\}$. The remaining vectors form a quad of cost 3. Thus, the greedy algorithm finds a solution of cost $6 = 2 \cdot OPT(I)$.

An instance of PQ(#1 = 2, distinct, connected) Consider the following instance I of PQ(#1 = 2, distinct, connected) consisting of 8 vectors represented in a graph shown in Figure 9 (recall that a vector in PQ(#1 = 2, distinct,

connected) corresponds to an edge in a simple graph).




Figure 9: An instance of PQ(#1 = 2, distinct, connected)

An optimal solution for this instance has cost 10, with the two quads $\{(1,2),(2,5),(3,5),(3,4)\}$ and $\{(6,7),(5,7),(5,8),(8,9)\}$, each having cost 5.

Since the instance features no quad with cost 4, the greedy algorithm may first select the following quad with cost 5: $\{(2,5),(3,5),(5,7),(5,8)\}$. Next, what remains is a quad of cost 8: $\{(1,2),(3,4),(6,7),(8,9)\}$.

Hence, the worst-case ratio of the greedy algorithm is at least 13/10, which is larger than the 5/4 approximation guarantee for algorithm A.

5 Conclusion

We have studied the worst-case behavior of a natural algorithm for partitioning a given set of vectors into quadruples. Informally, by running a matching algorithm once, we find pairs, and by running it one more time, we match the pairs into quadruples. Under the specific cost-structure studied here, we have shown the precise worst-case behavior of this method for all cases except PQ(#1 = 2, distinct), where a small gap remains.

It is a natural question to study an extension where we form clusters consisting of 2^s vectors for some given integer $s \geq 2$. Indeed, if we form groups of size 2^s by running s rounds of matching, the worst-case ratio is easily seen to be bounded by 2^{s-1} . To explain this, let M be the minimum-cost matching of the first round. Then $A(I) \leq \cos(M)$ and $OPT(I) \geq \cos(M)/2^{s-1}$ as the cost of the optimum (viewed as being constructed in s rounds) is at least $\cos(M)$ after the first round and could then halve in each further round. Moreover, since we have shown that the cost of the algorithm after two rounds is at most $\frac{3}{2}$ times the optimal cost after two rounds, we get a ratio of $\frac{3}{2} \times 2^{s-2} = 3 \times 2^{s-3}$. We leave the question of finding the worst-case ratio for arbitrary s as an open problem.

References

- [1] J. Barát and D. Gerbner. Edge-decomposition of graphs into copies of a tree with four edges. The Electronic Journal of Combinatorics, 21(1):1-55, 2014.
- [2] C. Berge. Sur le couplage maximum d'un graphe. Comptes Rendus de l'Académie des Sciences, 247:258 259, 1958.

- [3] T. Dokka, M. Bougeret, V. Boudet, R. Giroudeau, and F.C.R. Spieksma. Approximation algorithms for the wafer to wafer integration problem. In *Proceedings of the 10th International Workshop on Approximation and Online Algorithms (WAOA 2012)*, volume 7846 of *LNCS*, pages 286 297. Springer, 2013.
- [4] T. Dokka, Y. Crama, and F.C.R. Spieksma. Multi-dimensional vector assignment problems. Discrete Optimization, 14:111 – 125, 2014.
- [5] F. Dong, W. Yan, and F. Zhang. On the number of perfect matchings of line graphs. *Discrete Applied Mathematics*, 161(6):794 801, 2013.
- [6] A. Figueroa, A. Goldstein, T. Jiang, M. Kurowski, A. Lingas, and M. Persson. Approximate clustering of fingerprint vectors with missing values. In *Proceedings of the 2005 Australasian* Symposium on Theory of Computing (CATS 2005), volume 41 of CRPIT, pages 57 – 60. Australian Computer Society, 2005.
- [7] D.S. Hochbaum and A. Levin. Covering the edges of bipartite graphs using $K_{2,2}$ graphs. Theoretical Computer Science, 411(1):1-9, 2010.
- [8] I. Holyer. The NP-completeness of some edge-partition problems. SIAM Journal on Computing, 10(4):713 717, 1981.
- [9] M. Jünger, G. Reinelt, and W.R. Pulleyblank. On partitioning the edges of graphs into connected subgraphs. *Journal of Graph Theory*, 9(4):539 549, 1985.
- [10] S. Onn and L.J. Schulman. The vector partition problem for convex objective functions. Mathematics of Operations Research, 26(3):583 – 590, 2001.
- [11] S. Reda, G. Smith, and L. Smith. Maximizing the functional yield of wafer-to-wafer 3-d integration. *IEEE Transactions on Very Large Scale Integration (VLSI) Systems*, 17(9):1357 1362, 2009.
- [12] C. Thomassen. Edge-decompositions of highly connected graphs into paths. In *Abhandlungen* aus dem Mathematischen Seminar der Universität Hamburg, volume 78, pages 17 26. Springer, 2008.
- [13] V.V. Vazirani. Approximation Algorithms. Springer-Verlag, Inc., New York, USA, 2001.
- [14] D.P. Williamson and D.B. Shmoys. *The Design of Approximation Algorithms*. Cambridge University Press, New York, USA, 1st edition, 2011.