A Hilbert space approach to difference equations

Konrad Kitzing¹, Rainer Picard^{1,2}, Stefan Siegmund^{1,3}, Sascha Trostorff¹, and Marcus Waurick²

¹Technische Universität Dresden, Faculty of Mathematics, Institute of Analysis, Zellescher Weg 12-14, 01069 Dresden, Germany

²Department of Mathematics and Statistics, University of Strathclyde, 26 Richmond Street, Glasgow G1 1XH, Scotland

³Technische Universität Dresden, Faculty of Mathematics, Center for Dynamics, Zellescher Weg 12-14, 01069 Dresden, Germany

March 9, 2022

Abstract

We consider general difference equations $u_{n+1} = F(u)_n$ for $n \in \mathbb{Z}$ on exponentially weighted ℓ_2 spaces of two-sided Hilbert space valued sequences u and discuss initial value problems. As an application of the Hilbert space approach, we characterize exponential stability of linear equations and prove a stable manifold theorem for causal nonlinear difference equations.

1 Introduction

Let H be a separable Hilbert space over \mathbb{C} with scalar product $\langle \cdot, \cdot \rangle_H$, linear in the second and conjugate linear in the first argument, let $F: H^{\mathbb{Z}} \to H^{\mathbb{Z}}$. In this paper, we discuss difference equations

$$u_{n+1} = F(u)_n \qquad (n \in \mathbb{Z}) \tag{1.1}$$

and their solutions $u \in \ell$ on exponentially weighted sequence spaces $\ell \subseteq H^{\mathbb{Z}}$ (see Lemma 2.1). With the shift $\tau \colon H^{\mathbb{Z}} \to H^{\mathbb{Z}}$, $(u_n)_{n \in \mathbb{Z}} \mapsto (u_{n+1})_{n \in \mathbb{Z}}$, (see Lemma 2.6), equation (1.1) on ℓ can be reformulated equivalently as

$$\tau u = F(u) \qquad (u \in \ell). \tag{1.2}$$

Since τ is invertible with inverse $\tau^{-1} \colon H^{\mathbb{Z}} \to H^{\mathbb{Z}}$, $(u_n)_{n \in \mathbb{Z}} \mapsto (u_{n-1})_{n \in \mathbb{Z}}$, each solution $u \in \ell$ of (1.2) satisfies $u = \tau^{-1}F(u)$. If $\ell \subseteq H^{\mathbb{Z}}$ is a Banach space and $\tau^{-1} \circ F \colon \ell \to \ell$ a contraction, Banach's fixed point theorem yields a unique solution to (1.2) (see Theorem 2.7).

For example, the search for bounded solutions of an implicit (backward) Euler scheme approximation $u_{n+1} = u_n + hf(u_{n+1}), n \in \mathbb{Z}, h > 0$, of a differential equation with right-hand side $f : \mathbb{R}^d \to \mathbb{R}^d$, is of the form (1.2) with $F(u)_n = u_n + hf(u_{n+1})$ for $n \in \mathbb{Z}$ on the sequence space $\ell = \ell_{\infty}(\mathbb{Z}; \mathbb{R}^d) = \{x : \mathbb{Z} \to \mathbb{R}^d; \sup_{k \in \mathbb{Z}} |x_k|_{\mathbb{R}^d} < \infty\}$. On the other hand, if the implicit Euler scheme $u_{n+1} = F(u)_n$ is considered on $\mathbb{N} := \mathbb{Z}_{\geq 0}$ with an initial condition $u_0 = x \in \mathbb{R}^n$, it is well-known that for each u_n the implicit equation $u_{n+1} = u_n + hf(u_{n+1})$ needs to be solved iteratively, e.g. with the Newton method, to compute u_{n+1} from u_n . We will prove in Theorem 2.8 that initial value problems can be equivalently reformulated into (1.1) on \mathbb{Z} by adding the initial value x as an impulse $\delta_k x := (\delta_{k,n} x)_{n \in \mathbb{Z}} \in H^{\mathbb{Z}}$ at k = -1, where $\delta_{k,n}$ denotes the Kronecker delta symbol for $k, n \in \mathbb{Z}$, if $F(u)_n$ only depends on u_k for $k \in \mathbb{Z}_{\leq n}$, i.e. for so-called causal F (Definition 4.2). Note that for the implicit Euler scheme, $F(u)_n$ depends on u_{n+1} and F is not causal.

Let L(H) denote the space of bounded linear operators from H to H. In Section 3 we characterize exponential stability of linear difference equations $u_{n+1} = Au_n$ for $A \in L(H)$ (Theorem 3.9), by utilizing the \mathscr{Z} transform (Theorem 3.2). In Section 4 we prove a stable manifold theorem (Theorem 4.11) for causal (Definitions 3.6 & 4.2) difference equations $u_{n+1} = Au_n + F(u)_n$ with F(0) = 0. Section 2 is devoted to basic properties of sequence spaces and existence of solutions.

In this paper we follow an approach which goes back to [6] (see also [3, 5] and the references therein).

2 Difference equations on sequence spaces

Lemma 2.1 (Exponentially weighted ℓ_p spaces). Let $1 \le p < \infty$, $\rho > 0$. Define

$$\ell_{p,\varrho}(\mathbb{Z};H) \coloneqq \{ x \in H^{\mathbb{Z}}; \sum_{k \in \mathbb{Z}} |x_k|_H^p \varrho^{-pk} < \infty \},\$$
$$\ell_{\infty,\varrho}(\mathbb{Z};H) \coloneqq \{ x \in H^{\mathbb{Z}}; \sup_{k \in \mathbb{Z}} |x_k|_H \varrho^{-k} < \infty \}.$$

Then $\ell_{p,\varrho}(\mathbb{Z}; H)$ and $\ell_{\infty,\varrho}(\mathbb{Z}; H)$ are Banach spaces with norms

$$|x|_{\ell_{p,\varrho}(\mathbb{Z};H)} \coloneqq \left(\sum_{k \in \mathbb{Z}} |x_k|_H^p \varrho^{-pk}\right)^{\frac{1}{p}} \qquad (x \in \ell_{p,\varrho}(\mathbb{Z};H))$$

and

$$|x|_{\ell_{\infty,\varrho}(\mathbb{Z};H)} \coloneqq \sup_{k \in \mathbb{Z}} |x_k|_H \varrho^{-k} \qquad (x \in \ell_{\infty,\varrho}(\mathbb{Z};H)),$$

respectively. Moreover, $\ell_{2,\rho}(\mathbb{Z};H)$ is a Hilbert space with the inner product

$$\langle x, y \rangle_{\ell_{2,\varrho}(\mathbb{Z};H)} \coloneqq \sum_{k \in \mathbb{Z}} \langle x_k, y_k \rangle_H \varrho^{-2k} \qquad (x, y \in \ell_{2,\varrho}(\mathbb{Z};H)).$$

Write $\ell_p(\mathbb{Z}; H) \coloneqq \ell_{p,1}(\mathbb{Z}; H)$ for $1 \le p \le \infty$.

Remark 2.2 (Intersections of weighted ℓ_p spaces). Let $1 \leq p \leq \infty$, $0 < \varrho_1 < \varrho_2$. Then

$$\ell_{p,\varrho_1}(\mathbb{Z};H) \cap \ell_{p,\varrho_2}(\mathbb{Z};H) = \bigcap_{\varrho \in [\varrho_1,\varrho_2]} \ell_{p,\varrho}(\mathbb{Z};H).$$

Lemma 2.3 (Natural embedding of one-sided sequences). Let $1 \le p \le \infty$, $\rho > 0$, $a \in \mathbb{Z}$. Define

$$\ell_{p,\varrho}(\mathbb{Z}_{\geq a};H) \coloneqq \{x \upharpoonright_{\mathbb{Z}_{\geq a}}; x \in \ell_{p,\varrho}(\mathbb{Z};H)\},\$$

and for $x \in H^{\mathbb{Z}_{\geq a}}$ define $\iota x \in H^{\mathbb{Z}}$ by

$$(\iota x)_k := \begin{cases} 0 & \text{if } k < a, \\ x_k & \text{if } k \ge a. \end{cases}$$

Then $\ell_{p,\varrho}(\mathbb{Z}_{\geq a}; H)$ is a Banach space with norm $|\cdot|_{\ell_{p,\varrho}(\mathbb{Z}_{\geq a}; H)} \coloneqq |\iota \cdot |_{\ell_{p,\varrho}(\mathbb{Z}; H)}$, and

$$\iota \colon \ell_{p,\varrho}(\mathbb{Z}_{\geq a};H) \hookrightarrow \ell_{p,\varrho}(\mathbb{Z};H)$$

is an isometric embedding. Write $\ell_{p,\varrho}(\mathbb{Z}_{\geq a}; H) \subseteq \ell_{p,\varrho}(\mathbb{Z}; H)$.

Lemma 2.4 (Scale of one-sided weighted ℓ_p spaces). Let $1 \leq p < q \leq \infty$, $\rho, \varepsilon > 0$, $a \in \mathbb{Z}$. Then

(a) $\ell_{p,\varrho}(\mathbb{Z}_{\geq a}; H) \subsetneq \ell_{q,\varrho}(\mathbb{Z}_{\geq a}; H),$ (b) $\ell_{q,\varrho}(\mathbb{Z}_{\geq a}; H) \subsetneq \ell_{p,\varrho+\varepsilon}(\mathbb{Z}_{\geq a}; H).$

Proof. (a): Let $x \in \ell_{p,\varrho}(\mathbb{Z}_{\geq a}; H)$, i.e. $\sum_{k=a}^{\infty} |x_k|_H^p \varrho^{-pk} < \infty$. Then $\exists k_0 \in \mathbb{Z}_{\geq a}, C \in [0,1)$:

$$|x_k|_H^p \varrho^{-pk} \le C \qquad (k \ge k_0).$$

Hence, $\sup_{k\geq k_0} |x_k|_H \varrho^{-k} \leq C^{\frac{1}{p}}$, proving that $x \in \ell_{\infty,\varrho}(\mathbb{Z}_{\geq a}; H)$. If $q < \infty$, then $(|x_k|_H \varrho^{-k})^{q-p} < 1$ for $k \geq k_0$ and therefore

$$|x_k|_H^q \varrho^{-kq} = |x_k|_H^p \varrho^{-kp} (|x_k|_H \varrho^{-k})^{q-p} < |x_k|_H^p \varrho^{-kp} \qquad (k \ge k_0),$$

proving that $x \in \ell_{q,\varrho}(\mathbb{Z}_{\geq a}; H)$.

(b): Let $x \in \ell_{q,\varrho}(\mathbb{Z}_{\geq a}; H)$. Show that $x \in \ell_{1,\varrho+\varepsilon}(\mathbb{Z}_{\geq a}; H)$. By (a), $x \in \ell_{p,\varrho+\varepsilon}(\mathbb{Z}_{\geq a}; H)$. If $q = \infty$, using the fact that $|x_k|_H \varrho^{-k} \leq |x|_{\ell_{\infty,\varrho}(\mathbb{Z}_{\geq a}; H)}$ for $k \in \mathbb{Z}_{\geq a}$,

$$\sum_{k=a}^{\infty} |x_k|_H (\varrho + \varepsilon)^{-k} = \sum_{k=a}^{\infty} |x_k|_H \varrho^{-k} (\frac{\varrho}{\varrho + \varepsilon})^k \le |x|_{\ell_{\infty,\varrho}(\mathbb{Z}_{\ge a};H)} (\frac{\varrho}{\varrho + \varepsilon})^a (\frac{\varrho + \varepsilon}{\varepsilon}),$$

proving that $x \in \ell_{1,\varrho+\varepsilon}(\mathbb{Z}_{\geq a}; H)$. If $q < \infty$, let $q' \in (0,\infty)$ with $\frac{1}{q} + \frac{1}{q'} = 1$. Using Hölder's inequality,

$$\begin{split} \sum_{k=a}^{\infty} |x_k|_H (\varrho + \varepsilon)^{-k} &= \sum_{k=a}^{\infty} |x_k|_H \varrho^{-k} (\frac{\varrho}{\varrho + \varepsilon})^k \le \Big(\sum_{k=a}^{\infty} |x_k|_H^q \varrho^{-kq} \Big)^{\frac{1}{q}} \Big(\sum_{k=a}^{\infty} \left(\frac{\varrho}{\varrho + \varepsilon} \right)^{kq'} \Big)^{\frac{1}{q'}} \\ &= |x|_{\ell_{q,\varrho}(\mathbb{Z}_{\ge a};H)} \Big(\frac{\varrho}{\varrho + \varepsilon} \Big)^{aq'} \Big(1 - \Big(\frac{\varrho}{\varrho + \varepsilon} \Big)^{q'} \Big)^{-\frac{1}{q'}}, \end{split}$$

proving that $x \in \ell_{1,\varrho+\varepsilon}(\mathbb{Z}_{\geq a}; H)$.

Example 2.5. Solution to scalar equation with Kronecker delta impulse Let $x \in \mathbb{C}$, $a \in \mathbb{C} \setminus \{0\}$. We consider the scalar difference equation

$$u_{n+1} = au_n + \delta_{-1,n}x \qquad (n \in \mathbb{Z}).$$

A sequence $u \in \mathbb{C}^{\mathbb{Z}}$ satisfies the preceding difference equation if and only if

$$u_n = \begin{cases} a^{n+1}c & \text{if } n \in \mathbb{Z}_{\leq -1}, \\ a^{n+1}c + a^n x & \text{if } n \in \mathbb{Z}_{\geq 0}, \end{cases}$$

for some $c \in \mathbb{C}$. In dependence on c the support of u satisfies

$$\operatorname{supp} u \subseteq \begin{cases} \mathbb{Z}_{\leq -1} & \text{if } c = -x/a, \\ \mathbb{Z}_{\geq 0} & \text{if } c = 0, \end{cases}$$

and

$$\operatorname{supp} u = \mathbb{Z} \quad \text{if } c \neq 0, c \neq -x/a.$$

Let $\rho > 0$ and let $1 \le p \le \infty$. If $\rho \ne |a|$, we can select a unique solution by requiring that $u \in \ell_{p,\varrho}(\mathbb{Z};\mathbb{C})$. Indeed if $\rho \ne |a|$ then

$$u \in \ell_{p,\varrho}(\mathbb{Z};\mathbb{C}) \quad \Leftrightarrow \quad \begin{cases} (\varrho < |a| \land \operatorname{supp} u \subseteq \mathbb{Z}_{\leq -1}) \lor \\ (\varrho > |a| \land \operatorname{supp} u \subseteq \mathbb{Z}_{\geq 0}) \end{cases}$$
$$\Leftrightarrow \quad \begin{cases} (\varrho < |a| \land c = -x/a) \lor \\ (\varrho > |a| \land c = 0). \end{cases}$$

Lemma 2.6 (Shift operator). Let $1 \le p \le \infty$, $\rho > 0$. Then

$$\tau \colon \ell_{p,\varrho}(\mathbb{Z}; H) \to \ell_{p,\varrho}(\mathbb{Z}; H),$$
$$(x_k)_{k \in \mathbb{Z}} \mapsto (x_{k+1})_{k \in \mathbb{Z}},$$

is linear, bounded, invertible, and

$$\|\tau^n\|_{L(\ell_{p,\varrho}(\mathbb{Z};H))} = \varrho^n \qquad (n \in \mathbb{Z}).$$

Proof. Let $1 \le p \le \infty$, $\rho > 0$, $x \in \ell_{p,\rho}(\mathbb{Z}; H)$, $n \in \mathbb{Z}$. Using the fact that

$$(\tau^n x)_k = x_{n+k} \qquad (k \in \mathbb{Z}),$$

if $p < \infty$,

$$|\tau^{n}x|_{\ell_{p,\varrho}(\mathbb{Z};H)}^{p} = \sum_{k\in\mathbb{Z}} |x_{n+k}|_{H}^{p} \varrho^{-pk} = \sum_{k\in\mathbb{Z}} |x_{n+k}|_{H}^{p} \varrho^{-p(n+k)} \varrho^{pn} = \varrho^{pn} |x|_{\ell_{p,\varrho}(\mathbb{Z};H)}^{p},$$

proving that $|\tau^n x|_{\ell_{p,\varrho}(\mathbb{Z};H)} = \varrho^n |x|_{\ell_{p,\varrho}(\mathbb{Z};H)}$, and, if $p = \infty$,

$$|\tau^n x|_{\ell_{\infty,\varrho}(\mathbb{Z};H)} = \sup_{k \in \mathbb{Z}} |x_{n+k}|_H \varrho^{-k} = \sup_{k \in \mathbb{Z}} |x_{n+k}|_H \varrho^{-(n+k)} \varrho^n = \varrho^n |x|_{\ell_{\infty,\varrho}(\mathbb{Z};H)}. \quad \Box$$

Theorem 2.7 (Existence and uniqueness of solutions). Let $1 \leq p \leq \infty$, $\rho > 0$, $F: \ell_{p,\varrho}(\mathbb{Z}; H) \to \ell_{p,\varrho}(\mathbb{Z}; H)$ with

$$|F|_{\operatorname{Lip}(\ell_{p,\varrho}(\mathbb{Z};H))} \coloneqq \sup_{x,y \in \ell_{p,\varrho}(\mathbb{Z};H), x \neq y} \frac{|F(x) - F(y)|_{\ell_{p,\varrho}(\mathbb{Z};H)}}{|x - y|_{\ell_{p,\varrho}(\mathbb{Z};H)}} < \varrho.$$

Then the difference equation

$$\tau u = F(u) \qquad (u \in \ell_{p,\varrho}(\mathbb{Z}; H))$$

has a unique solution.

Proof. Then

$$\mathscr{T} \colon \ell_{p,\varrho}(\mathbb{Z}; H) \mapsto \ell_{p,\varrho}(\mathbb{Z}; H)$$
$$u \mapsto \tau^{-1} F(u)$$

is a contraction, since, by Lemma 2.6, for $x, y \in \ell_{p,\varrho}(\mathbb{Z}; H)$,

$$|\mathscr{T}(x) - \mathscr{T}(y)| \le |\tau^{-1}|_{L(\ell_{p,\varrho}(\mathbb{Z};H))}|F(x) - F(y)|_{\ell_{p,\varrho}(\mathbb{Z};H)} = \frac{1}{\varrho}|F|_{\mathrm{Lip}(\ell_{p,\varrho}(\mathbb{Z};H))} < 1.$$

Its unique fixed point $u = \mathscr{T}(u)$ satisfies $\tau u = F(u)$.

Explicit difference equations [1] of the form $u_{n+1} = G(u_n)$ on $\mathbb{Z}_{\geq 0}$ with initial condition $u_0 \in \mathbb{R}^n$ have a unique solution $u: \mathbb{Z}_{\geq 0} \to \mathbb{R}^n$ for arbitrary functions $G: \mathbb{R}^n \to \mathbb{R}^n$. In the following theorem, solutions of more general initial value problems are characterized as one-sided sequences which solve a difference equation on \mathbb{Z} with an impulse corresponding to the initial condition.

Theorem 2.8 (Initial value problems). Let $A \in L(H)$, $F: H^{\mathbb{Z}} \to H^{\mathbb{Z}}$, $u \in H^{\mathbb{Z}}$, spt $u \subseteq \mathbb{Z}_{\geq 0}$, spt $F(u) \subseteq \mathbb{Z}_{\geq 0}$, $x \in H$. Then the following statements are equivalent.

(i)
$$u_{n+1} = Au_n + F(u)_n$$
, $u_0 = x$ $(n \in \mathbb{Z}_{\geq 0})$. (Initial value problem)
(ii) $u_n = A^n x + \sum_{k=0}^{n-1} A^{n-k-1} F(u)_k$ $(n \in \mathbb{Z}_{\geq 0})$. (Variation of constants)
(iii) $\tau u = Au + F(u) + \delta_{-1} x$. (Initial condition as impulse)

Proof. (i) \Rightarrow (ii): By induction for $n \in \mathbb{Z}_{\geq 0}$, $u_n = A^n x + \sum_{k=0}^{n-1} A^{n-k-1} F(u)_k$, since

$$u_{n+1} = Au_n + F(u)_n = A\left(A^n x + \sum_{k=0}^{n-1} A^{n-k-1} F(u)_k\right) + F(u)_n$$
$$= A^{n+1} x + \sum_{k=0}^n A^{n-k} F(u)_k.$$

(ii) \Rightarrow (iii): $u_0 = x$, and for $n \in \mathbb{Z}_{\geq 0}$, $u_{n+1} = Au_n + F(u)_n$, since

$$u_{n+1} = A^{n+1}x + \sum_{k=0}^{n} A^{n-k}F(u)_k = A\left(A^nx + \sum_{k=0}^{n-1} A^{n-k-1}F(u)_k\right) + F(u)_n$$
$$= Au_n + F(u)_n.$$

Using spt u, spt $F(u) \subseteq \mathbb{Z}_{\geq 0}$, for $n \in \mathbb{Z}$,

$$(Au + F(u) + \delta_{-1}x)_n = Au_n + F(u)_n + \delta_{-1,n}x = \begin{cases} 0 & \text{if } n \le -2, \\ x & \text{if } n = -1, \\ Au_n + F(u)_n & \text{if } n \ge 0, \end{cases}$$

hence $(\tau u)_n = u_{n+1} = (Au + F(u) + \delta_{-1}x)_n$, proving (iii).

(iii) \Rightarrow (i): For $n \in \mathbb{Z}$, $u_{n+1} = (\tau u)_n = (Au + F(u) + \delta_{-1}x)_n = Au_n + F(u)_n + \delta_{-1,n}x$. Using the fact that spt $u \subseteq \mathbb{Z}_{\geq 0}$, spt $F(u) \subseteq \mathbb{Z}_{\geq 0}$, we see that $u_0 = Au_{-1} + F(u)_{-1} + x = x$, implying (i).

Remark 2.9 (Classical initial value problems). Let $G \colon \mathbb{R}^n \to \mathbb{R}^n$ and

$$F(u)_n \coloneqq \begin{cases} G(u_n) & \text{if } n \in \mathbb{Z}_{\geq 0}, \\ 0 & \text{if } n \in \mathbb{Z}_{\leq -1} \end{cases}$$

for $u: \mathbb{Z} \to \mathbb{R}^n$. Then every solution $u: \mathbb{Z} \to \mathbb{R}^n$ of $\tau u = Au + F(u) + \delta_{-1}x$ with spt $u \subseteq \mathbb{Z}_{\geq 0}$ gives rise to a solution of the initial value problem $u_{n+1} = G(u_n)$ on $\mathbb{Z}_{\geq 0}$ with initial condition $u_0 = x \in \mathbb{R}^n$ and vice versa.

3 The \mathscr{Z} transform and the operator $(\tau - A)^{-1}$

Let $A \in L(H)$, $\rho > 0$, $F : \ell_{2,\rho}(\mathbb{Z}; H) \to \ell_{2,\rho}(\mathbb{Z}; H)$. As in Theorem 2.8, for $x \in H$, consider initial value problems of the form

$$\tau u = Au + F(u) + \delta_{-1}x \qquad (u \in \ell_{2,\rho}(\mathbb{Z}; H)).$$

In this section we reformulate this problem equivalently as a fixed point problem

$$u = (\tau - A)^{-1} (F(u) + \delta_{-1} x)$$

by discussing the operator $(\tau - A)^{-1}$ on $\ell_{2,\varrho}(\mathbb{Z}; H)$. Our main tool is the \mathscr{Z} transform.

Lemma 3.1 (L_2 space on circle and orthonormal basis). Let $\varrho > 0$, $S_{\varrho} := \{z \in \mathbb{C}; |z| = \varrho\}$. Define

$$L_2(S_{\varrho};H) := \{ f \colon S_{\varrho} \to H; \int_{S_{\varrho}} |f(z)|_H^2 \frac{\mathrm{d}z}{|z|} < \infty \}.$$

Then $L_2(S_{\varrho}; H)$ is a Hilbert space with the inner product

$$\langle f,g\rangle_{L_2(S_\varrho;H)} \coloneqq \frac{1}{2\pi} \int_{S_\varrho} \left\langle f(z),g(z)\right\rangle_H \frac{\mathrm{d}z}{|z|} \qquad (f,g\in L_2(S_\varrho;H)).$$

Moreover, let $(\psi_n)_{n\in\mathbb{Z}}$ be an orthonormal basis in H. Then $(p_{k,n})_{k,n\in\mathbb{Z}}$ with

$$p_{k,n}(z) \coloneqq \varrho^k z^{-k} \psi_n \qquad (z \in S_{\varrho})$$

is an orthonormal basis in $L_2(S_{\varrho}; H)$.

Proof. Let $L_2((-\pi,\pi); H)$ denote the Hilbert space of square integrable functions from $(-\pi,\pi)$ to H with the scalar product

$$\langle f,g \rangle_{L_2((-\pi,\pi);H)} \coloneqq \frac{1}{2\pi} \int_{-\pi}^{\pi} \langle f(z),g(z) \rangle_H \, dz \qquad (f,g \in L_2((-\pi,\pi);H)).$$

The isometric isomorphism

$$\Phi: L_2(S_{\varrho}; H) \to L_2((-\pi, \pi); H), \quad f \mapsto \left(\varphi \mapsto f(\varrho e^{i\varphi})\right),$$

maps $(p_{k,n})_{k,n\in\mathbb{Z}}$ onto the orthonormal basis $(\varphi \mapsto e^{-ik\varphi}\psi_n)_{k,n\in\mathbb{Z}}$ of $L_2((-\pi,\pi);H)$, proving that $(\Phi(p_{k,n}))_{k,n\in\mathbb{Z}}$ is an orthonormal basis of $L_2(S_\varrho;H)$. **Theorem 3.2** (\mathscr{Z} transform). Let $\rho > 0$. The operator

$$\mathscr{Z}_{\varrho} \colon \ell_{2,\varrho}(\mathbb{Z}; H) \to L_2(S_{\varrho}; H),$$
$$x \mapsto \Big(z \mapsto \sum_{k \in \mathbb{Z}} \langle \psi_n, \varrho^{-k} x_k \rangle_H p_{k,n}(z) \Big),$$

is well-defined and unitary. For $x \in \ell_{1,\varrho}(\mathbb{Z}; H) \subseteq \ell_{2,\varrho}(\mathbb{Z}; H)$ we have

$$\mathscr{Z}_{\varrho}(x) = \left(z \mapsto \sum_{k \in \mathbb{Z}} x_k z^{-k}\right).$$

Remark 3.3 (\mathscr{Z} transform of $x \in \ell_{2,\varrho}(\mathbb{Z}; H) \setminus \ell_{1,\varrho}(\mathbb{Z}; H)$). Let $\varrho > 0, x \in \ell_{2,\varrho}(\mathbb{Z}; H) \setminus \ell_{1,\varrho}(\mathbb{Z}; H)$. Then

$$\sum_{k \in \mathbb{Z}} x_k z^{-k}$$

does not necessarily converge for all $z \in S_{\varrho}$. For example if $H = \mathbb{C}$, $x \in \ell_{2,\varrho}(\mathbb{Z}; H) \setminus \ell_{1,\varrho}(\mathbb{Z}; H)$ with $x_k := \frac{\varrho^k}{k}$ and $z = \varrho$.

Proof of Theorem 3.2. Let $\varrho > 0$, $x \in \ell_{2,\varrho}(\mathbb{Z}; H)$. We prove that \mathscr{Z}_{ϱ} is well-defined, i.e. that $\mathscr{Z}_{\varrho}(x)$ exists as an element in $L_2(S_{\varrho}; H)$. We compute

$$\begin{split} \sum_{k,n\in\mathbb{Z}} \langle \psi_n, \varrho^{-k} x_k \rangle_H p_{k,n} \big|_{L_2(S_\varrho;H)}^2 &= \frac{1}{2\pi} \int_{S_\varrho} \big| \sum_{k,n\in\mathbb{Z}} \langle \psi_n, \varrho^{-k} x_k \rangle_H p_{k,n}(z) \big|_H^2 \frac{\mathrm{d}z}{|z|} \\ &= \frac{1}{2\pi} \int_{S_\varrho} \big| \sum_{k,n\in\mathbb{Z}} \langle \psi_n, \varrho^{-k} x_k \rangle_H \varrho^k z^{-k} \psi_n \big|_H^2 \frac{\mathrm{d}z}{|z|} \\ &= \frac{1}{2\pi} \int_{S_\varrho} \big| \sum_{k,n\in\mathbb{Z}} \langle \psi_n, x_k \rangle_H z^{-k} \psi_n \big|_H^2 \frac{\mathrm{d}z}{|z|} \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \big| \sum_{k,n\in\mathbb{Z}} \langle \psi_n, x_k \rangle_H (\varrho^{e^{i\varphi}})^{-k} \psi_n \big|_H^2 \mathrm{d}\varphi \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \big| \sum_{k\in\mathbb{Z}} \varrho^{-k} e^{-i\varphi k} \sum_{n\in\mathbb{Z}} \langle \psi_n, x_k \rangle_H \psi_n \big|_H^2 \mathrm{d}\varphi \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \big| \sum_{k\in\mathbb{Z}} \varrho^{-k} e^{-i\varphi k} x_k \big|_H^2 \mathrm{d}\varphi \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{k,\ell\in\mathbb{Z}} \varrho^{-k} e^{-i\varphi k} x_k \sum_{\ell\in\mathbb{Z}} \varrho^{-\ell} e^{-i\varphi \ell} x_\ell \rangle_H \mathrm{d}\varphi \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{k,\ell\in\mathbb{Z}} \varrho^{-k-\ell} e^{-i\varphi(\ell-k)} \langle x_k, x_\ell \rangle_H \mathrm{d}\varphi \\ &= \frac{1}{2\pi} \sum_{k,\ell\in\mathbb{Z}} \varrho^{-k-\ell} \langle x_k, x_\ell \rangle_H \int_{-\pi}^{\pi} e^{-i\varphi(\ell-k)} \mathrm{d}\varphi \end{split}$$

$$= \sum_{k \in \mathbb{Z}} |x_k|_H^2 \varrho^{-2k}$$
$$= |x|_{\ell_{2,\varrho}(\mathbb{Z};H)}^2,$$

proving that \mathscr{Z}_{ϱ} is well-defined and an isometry. To prove that \mathscr{Z}_{ϱ} is unitary, it remains to show that \mathscr{Z}_{ϱ} is surjective. Let $f \in L_2(S_{\varrho}; H)$. Define

$$x_k \coloneqq \frac{1}{2\pi} \int_{S_{\varrho}} f(z) \varrho^{2k} \overline{z}^{-k} \frac{\mathrm{d}z}{|z|} \qquad (k \in \mathbb{Z}).$$

Then $\mathscr{Z}_{\varrho} x = f$, since for $k, n \in \mathbb{Z}$,

<

$$\begin{aligned} \mathscr{X}_{\varrho}x, p_{k,n}\rangle_{L_{2}(S_{\varrho};H)} &= \frac{1}{2\pi} \int_{S_{\varrho}} \left\langle \sum_{\ell,m\in\mathbb{Z}} \langle \psi_{m}, \varrho^{-\ell}x_{\ell} \rangle_{H} p_{\ell,m}(z), p_{k,n}(z) \rangle_{H} \frac{\mathrm{d}z}{|z|} \right. \\ &= \frac{1}{2\pi} \int_{S_{\varrho}} \sum_{\ell,m\in\mathbb{Z}} \langle \varrho^{-\ell}x_{\ell}, \psi_{m} \rangle_{H} \left\langle p_{\ell,m}(z), p_{k,n}(z) \rangle_{H} \frac{\mathrm{d}z}{|z|} \right. \\ &= \sum_{\ell,m\in\mathbb{Z}} \langle \varrho^{-\ell}x_{\ell}, \psi_{m} \rangle_{H} \frac{1}{2\pi} \int_{S_{\varrho}} \left\langle p_{\ell,m}(z), p_{k,n}(z) \rangle_{H} \frac{\mathrm{d}z}{|z|} \right. \\ &= \left\langle \varrho^{-k}x_{k}, \psi_{n} \right\rangle_{H} \\ &= \left\langle \frac{1}{2\pi} \int_{S_{\varrho}} f(z) \varrho^{2k}(\overline{z})^{-k} \frac{\mathrm{d}z}{|z|}, \varrho^{-k}\psi_{n} \right\rangle_{H} \\ &= \frac{1}{2\pi} \int_{S_{\varrho}} \langle f(z), \varrho^{k}z^{-k}\psi_{n} \rangle_{H} \frac{\mathrm{d}z}{|z|} \\ &= \left\langle f, p_{k,n} \right\rangle_{L_{2}(S_{\varrho};H)}. \end{aligned}$$

For $x \in \ell_{2,\varrho}(\mathbb{Z}; H) \cap \ell_{1,\varrho}(\mathbb{Z}; H)$ and $z \in S_{\varrho}$ we compute

$$\sum_{k,n\in\mathbb{Z}} \langle \psi_n, \varrho^{-k} x_k \rangle_H p_{k,n}(z) = \sum_{k,n\in\mathbb{Z}} \langle \psi_n, \varrho^{-k} x_k \rangle_H \varrho^k z^{-k} \psi_n$$
$$= \sum_{k,n\in\mathbb{Z}} \langle \psi_n, x_k z^{-k} \rangle_H \psi_n$$
$$= \sum_{k\in\mathbb{Z}} x_k z^{-k}.$$

Lemma 3.4 (Shift is unitarily equivalent to multiplication). Let $\rho > 0$. Then

$$\mathscr{Z}_{\varrho}\tau\mathscr{Z}_{\varrho}^{*}=\mathrm{m},$$

where m is the multiplication-by-the-argument operator acting in $L_2(S_{\varrho}; H)$, i.e.,

$$\begin{split} \mathbf{m} \colon L_2(S_\varrho;H) &\to L_2(S_\varrho;H) \\ f \mapsto (z \mapsto zf(z)). \end{split}$$

Proof. By the boundedness of the operators involved and the unitarity of \mathscr{Z}_{ϱ} , it suffices to prove

$$\mathbf{m}\mathscr{Z}_{\rho}x = \mathscr{Z}_{\rho}\tau x$$

for $x \in H^{\mathbb{Z}}$ with compact support. For this, we compute

$$(\mathbf{m}\mathscr{Z}_{\varrho}x)(z) = z\mathscr{Z}_{\varrho}x(z)$$

= $z\sum_{k\in\mathbb{Z}} x_k z^{-k}$
= $\sum_{k\in\mathbb{Z}} x_k z^{-k+1}$
= $\sum_{k\in\mathbb{Z}} x_k z^{-(k-1)} = \sum_{k\in\mathbb{Z}} (\tau x)_k z^{-k} = (\mathscr{Z}_{\varrho}\tau x)(z).$

Next, we present a Payley–Wiener type result for the \mathscr{Z} transform.

Lemma 3.5 (Characterization of positive support). Let $\varrho > 0$, $x \in \ell_{2,\varrho}(\mathbb{Z}; H)$. Then the following statements are equivalent: (i) spt $x \in \mathbb{Z}_{>0}$

(i) spt
$$x \subseteq \mathbb{Z}_{\geq 0}$$
,
(ii) $z \mapsto \sum_{k \in \mathbb{Z}} x_k z^{-k}$ is analytic on $\mathbb{C}_{|\cdot| > \varrho}$ and

$$\sup_{\mu > \varrho} \int_{S_{\mu}} \left| \sum_{k \in \mathbb{Z}} x_k z^{-k} \right|_H^2 \frac{\mathrm{d}z}{|z|} < \infty.$$
(3.1)

Proof. (i) \Rightarrow (ii): Let $\varepsilon > 0$. Then for $z \in \mathbb{C}$ with $|z| \ge \varrho + \varepsilon$ and $k \in \mathbb{Z}_{\ge 0}$, $|z|^{-k} \le (\varrho + \varepsilon)^{-k}$, and

$$\begin{split} \sum_{k=0}^{\infty} |x_k z^{-k}|_H &= \sum_{k=0}^{\infty} |x_k|_H |z^{-k}| \\ &= \sum_{k=0}^{\infty} |x_k|_H \varrho^{-k} \varrho^k |z^{-k}| \\ &\leq \sum_{k=0}^{\infty} |x_k|_H \varrho^{-k} \varrho^k (\varrho + \varepsilon)^{-k} \\ &\leq \left(\sum_{k=0}^{\infty} |x_k|_H^2 \varrho^{-2k}\right)^{\frac{1}{2}} \left(\sum_{k=0}^{\infty} \left(\frac{\varrho}{\varrho + \varepsilon}\right)^{2k}\right)^{\frac{1}{2}} \\ &= |x|_{\ell_{2,\varrho}(\mathbb{Z};H)} \left(1 - \left(\frac{\varrho}{\varrho + \varepsilon}\right)^2\right)^{-\frac{1}{2}}. \end{split}$$

Hence $z \mapsto \sum_{k \in \mathbb{Z}} x_k z^{-k}$ is analytic on $\mathbb{C}_{|\cdot| > \varrho}$. Moreover, for $\mu > \varrho$,

$$|x|_{\ell_{2,\mu}(\mathbb{Z};H)}^{2} = \sum_{k=0}^{\infty} |x_{k}|_{H}^{2} \mu^{-2k}$$

$$\leq \sum_{k=0}^{\infty} |x_k|_H^2 \varrho^{-2k}$$
$$= |x|_{\ell_{2,\varrho}(\mathbb{Z};H)}^2.$$

Using the fact that \mathscr{Z}_{μ} is unitary,

$$\frac{1}{2\pi} \int_{S_{\mu}} \left| \sum_{k \in \mathbb{Z}} x_k z^{-k} \right|_H^2 \frac{\mathrm{d}z}{|z|} = |\mathscr{Z}_{\mu} x|_{L_2(S_{\mu};H)}^2 = |x|_{\ell_{2,\mu}(\mathbb{Z};H)}^2 \le |x|_{\ell_{2,\varrho}(\mathbb{Z};H)}^2,$$

proving (3.1).

(ii) \Rightarrow (i): Let $k \in \mathbb{Z}_{<0}$. Define $e^{(k)} \in \mathbb{C}^{\mathbb{Z}}$ by $e_{\ell}^{(k)} := \delta_{k,\ell}$ for $\ell \in \mathbb{Z}$, then $e^{(k)} \in \ell_2(\mathbb{Z};\mathbb{C})$. Let $n \in \mathbb{Z}$. Then

$$\begin{aligned} |\langle x_k, \psi_n \rangle_H| &= \Big| \sum_{\ell \in \mathbb{Z}} \langle x_\ell, \psi_n \rangle_H e_\ell^{(k)} \Big| \\ &= \Big| \sum_{\ell \in \mathbb{Z}} \langle x_\ell, \mu^{2\ell} \psi_n e_\ell^{(k)} \rangle_H \mu^{-2\ell} \Big| \\ &= |\langle x, \mu^{2\cdot} \psi_n e_{\cdot}^{(k)} \rangle_{\ell_{2,\mu}(\mathbb{Z};H)} | \\ &= |\langle \mathscr{Z}_\mu x, \mathscr{Z}_\mu \mu^{2\cdot} \psi_n e_{\cdot}^{(k)} \rangle_{L_2(S_\mu;H)} | \\ &\leq |\mathscr{Z}_\mu x|_{L_2(S_\mu;H)} |\mathscr{Z}_\mu \mu^{2\cdot} \psi_n e_{\cdot}^{(k)}|_{L_2(S_\mu;H)} \\ &\leq |\mathscr{Z}_\mu x|_{L_2(S_\mu;H)} \mu^k |\psi_n|. \end{aligned}$$

Since $(\mathscr{Z}_{\mu}\mu^{2\cdot}\psi_{n}e^{(k)}_{\cdot})(z) = \sum_{\ell\in\mathbb{Z}}\mu^{2\ell}\psi_{n}e^{(k)}_{\ell}z^{-\ell} = \mu^{2k}\psi_{n}z^{-k}$ for $z\in S_{\mu}$, $|\langle x_{k},\psi_{n}\rangle_{H}| \leq |\mathscr{Z}_{\mu}x|_{L_{2}(S_{\mu};H)}\mu^{k}|\psi_{n}|.$

By (3.1), that is, $\sup_{\mu>\varrho} |\mathscr{Z}_{\mu}x|_{L_2(S_{\mu};H)} < \infty$, and since k < 0, the right-hand side tends to 0 as $\mu \to \infty$, proving that $x_k = 0$.

Definition 3.6 (Causal linear operator). We call a linear operator $B: \ell_{2,\varrho}(\mathbb{Z}; H) \to \ell_{2,\varrho}(\mathbb{Z}; H)$ causal, if for all $a \in \mathbb{Z}$, $f \in \ell_{2,\varrho}(\mathbb{Z}; H)$, we have

 $\operatorname{spt} f \subseteq \mathbb{Z}_{\geq a} \quad \Rightarrow \quad \operatorname{spt} Bf \subseteq \mathbb{Z}_{\geq a}.$

Note, that if an operator $B: \ell_{2,\varrho}(\mathbb{Z}; H) \to \ell_{2,\varrho}(\mathbb{Z}; H)$ commutes with the shift operator τ , the operator B is causal, if and only if for all $f \in \ell_{2,\varrho}(\mathbb{Z}; H)$ we have

$$\operatorname{spt} f \subseteq \mathbb{Z}_{\geq 0} \quad \Rightarrow \quad \operatorname{spt} Bf \subseteq \mathbb{Z}_{\geq 0}.$$

The next major result is Theorem 3.7 which characterizes the causality of the operator $(\tau - A)^{-1}$ on $\ell_{2,\varrho}(\mathbb{Z}; H)$ by the spectral radius of A. Recall [4, VIII.3.6, p. 222] that for $A \in L(H)$ with spectrum $\sigma(A)$, the spectral radius

$$r(A) \coloneqq \sup\{|z|; z \in \sigma(A)\}$$

of A satisfies

$$r(A) = \lim_{n \to \infty} |A^n|_{L(H)}^{1/n}$$

Let $A \in L(H)$ and $\varrho > 0$. We denote the operators $\ell_{2,\varrho}(\mathbb{Z}, H) \to \ell_{2,\varrho}(\mathbb{Z}, H)$, $x \mapsto (Ax_k)_k$, and $L_2(S_{\varrho}, H) \to L_2(S_{\varrho}, H)$, $f \mapsto (z \mapsto Af(z))$, which have the same operator norm as A, again by A.

Theorem 3.7 (Characterization of causality of $(\tau - A)^{-1}$ by spectral radius). Let $\rho > 0, A \in L(H)$. Then the following statements are equivalent:

(i)
$$(\tau - A)^{-1} \in L(\ell_{2,\varrho}(\mathbb{Z}; H))$$
 is causal,
(ii) $(\tau - A)^{-1} \in L(\ell_{2,\varrho}(\mathbb{Z}; H))$ and $\operatorname{spt}(\tau - A)^{-1}\delta_{-1}x \subseteq \mathbb{Z}_{\geq 0}$ for all $x \in H$,
(iii) $\varrho > r(A)$.

Proof. (i) \Rightarrow (ii): By assumption $(\tau - A)^{-1} \in L(\ell_{2,\varrho}(\mathbb{Z}; H))$. Let $u \coloneqq (\tau - A)^{-1}\delta_{-1}x$. By causality of $(\tau - A)^{-1}$, it follows that $\operatorname{spt} u \subseteq \mathbb{Z}_{\geq -1}$. Using the fact that $\tau u = Au + \delta_{-1}x$, we get

$$u_{-1} = Au_{-2} + \delta_{-1,-2}x = 0,$$

proving spt $u \subseteq \mathbb{Z}_{>0}$.

(ii) \Rightarrow (iii): We have $(\tau - A)^{-1} \in L(\ell_{2,\varrho}(\mathbb{Z}; H))$. Hence, by Lemma 3.4, $(m - A)^{-1} \in L(L_2(S_{\varrho}; H))$. By the definition of m, this implies that S_{ϱ} along with a neighborhood of S_{ϱ} is contained in the resolvent set of $A \in L(H)$. Therefore the spectral radius cannot be equal to ϱ . Assume $\varrho < r(A)$. By the formula for the spectral radius, there exists $\varepsilon > 0$, $n_0 \in \mathbb{N}$ with $(\varrho + \varepsilon)^n \leq |A^n|_{L(H)}$ for all $n \geq n_0$. Hence,

$$\left|\frac{A^n}{\varrho^n}\right|_{L(H)} \to \infty \qquad (n \to \infty).$$

By the uniform boundedness principle there exists $x \in H$ with

$$\frac{|A^n x|_H}{\varrho^n |x|_H} \to \infty \qquad (n \to \infty),$$

i.e., there exists $n_1 \in \mathbb{N}$ such that $|A^n x|_H \ge \varrho^n |x|_H$ for all $n \ge n_1$. By hypothesis we obtain $u \coloneqq (\tau - A)^{-1} \delta_{-1} x \in \ell_{2,\varrho}(\mathbb{Z}; H)$ and

spt
$$u \subseteq \mathbb{Z}_{\geq 0}$$
.

Since $\tau u = Au + \delta_{-1}x$, it follows that

$$u_n = \begin{cases} 0 & \text{if } n \le -1, \\ A^n x & \text{if } n \ge 0. \end{cases}$$

Consequently, we obtain

$$|u|^2_{\ell_{2,\varrho}(\mathbb{Z};H)} = \sum_{k \in \mathbb{Z}} |u_k|^2_H \varrho^{-2k}$$

$$= \sum_{k=0}^{\infty} |A^k x|_H^2 \varrho^{-2k}$$
$$\ge \sum_{k=n_1}^{\infty} |x|_H \varrho^{2k} \varrho^{-2k} = \infty$$

which is a contradiction. Thus, $\rho > r(A)$.

(iii) \Rightarrow (i): Let $\rho > r(A)$. Then $(m - A)^{-1} \in L(L_2(S_{\rho}; H))$ and hence $(\tau - A)^{-1} \in L(\ell_{2,\rho}(\mathbb{Z}; H))$. To show (i), let $f \in \ell_{2,\rho}(\mathbb{Z}; H)$, spt $f \subseteq \mathbb{Z}_{\geq 0}$, and prove that $u \coloneqq (\tau - A)^{-1}f$ satisfies spt $u \subseteq \mathbb{Z}_{\geq 0}$. By Lemma 3.5, $z \mapsto \sum_{k \in \mathbb{Z}} f_k z^{-k}$ is analytic on $\mathbb{C}_{|\cdot| > \rho}$. Hence, also

$$z \mapsto \sum_{k \in \mathbb{Z}} u_k z^{-k} = (z - A)^{-1} \sum_{k \in \mathbb{Z}} f_k z^{-k} \quad \text{is analytic on } \mathbb{C}_{|\cdot| > \varrho}.$$

Next, by Lemma 3.5, $\sup_{\mu>\varrho} \int_{S_{\mu}} \left| \sum_{k\in\mathbb{Z}} f_k z^{-k} \right|_H^2 \frac{\mathrm{d}z}{|z|} < \infty$. Moreover, we have

$$\sup_{z\in\mathbb{C}_{|\cdot|>\varrho}}\left|(z-A)^{-1}\right|_{L(H)}<\infty.$$

Indeed, this follows from $(z-A)^{-1} = \frac{1}{z}(1-z^{-1}A)^{-1} = \frac{1}{z}\sum_{k=0}^{\infty}(\frac{1}{z}A)^k$ for $z \in \mathbb{C}_{|\cdot|>\varrho}$ and $\rho > r(A)$. For $\mu > \rho$

$$\int_{S_{\mu}} \left| \sum_{k \in \mathbb{Z}} u_{k} z^{-k} \right|_{H}^{2} \frac{\mathrm{d}z}{|z|} = \int_{S_{\mu}} \left| (z - A)^{-1} \sum_{k \in \mathbb{Z}} f_{k} z^{-k} \right|_{H}^{2} \frac{\mathrm{d}z}{|z|}$$
$$\leq \sup_{z \in \mathbb{C}_{|\cdot| > \varrho}} \left| (z - A)^{-1} \right|_{L(H)}^{2} \int_{S_{\mu}} \left| \sum_{k \in \mathbb{Z}} f_{k} z^{-k} \right|_{H}^{2} \frac{\mathrm{d}z}{|z|},$$

hence $\sup_{\mu>\varrho} \int_{S_{\mu}} \left| \sum_{k\in\mathbb{Z}} u_k z^{-k} \right|_{H}^2 \frac{dz}{|z|} < \infty$. By Lemma 3.5, spt $u \subseteq \mathbb{Z}_{\geq 0}$.

Next, we address linear initial value problems. In particular, we show that for $x \in H$ the problem

$$\tau u = Au + \delta_{-1}x$$

has exactly one solution $u \in H^{\mathbb{Z}}$ satisfying spt $u \subseteq \mathbb{Z}_{\geq 0}$ and this solution u solves the initial value problem

$$u_{n+1} = Au_n, \quad u_0 = x \qquad (n \in \mathbb{Z}_{\ge 0}).$$

Proposition 3.8 (Linear initial value problem). Let $A \in L(H)$, $x \in H$ and $u \in H^{\mathbb{Z}}$. Then the following statements are equivalent:

(i) $\tau u = Au + \delta_{-1}x$ and $\operatorname{spt}(u) \subseteq \mathbb{Z}_{>0}$. (*ii*) $u_{n+1} = Au_n$ on $\mathbb{Z}_{>0}$ and $u_0 = x$ and $u_n = 0$ on $\mathbb{Z}_{<0}$.

(iii)
$$u_n = \begin{cases} 0 & \text{if } n < 0, \\ A^n x & \text{if } n \ge 0. \end{cases}$$

Moreover, if $\rho > r(A)$ then u satisfying (i), (ii), (iii) is unique in $\ell_{2,\rho}(\mathbb{Z}; H)$. I.e., if $\rho > r(A)$ then (i), (ii), (iii) are equivalent to (iv) and (v).

- (iv) $\tau u = Au + \delta_{-1}x$ and $u \in \ell_{2,\varrho}(\mathbb{Z}; H)$.
- (v) $u = (\tau A)^{-1} \delta_{-1} x \in \ell_{2,\varrho}(\mathbb{Z}; H).$

Proof. The equivalence of (i), (ii) and (iii) are reformulations in the spirit of Theorem 2.8. By Theorem 3.7, the operator $(\tau - A)^{-1} \in L(\ell_{2,\varrho}(\mathbb{R}; H))$ is well-defined and causal. This shows $(iv) \Leftrightarrow (v)$ and $(v) \Leftrightarrow (i)$.

We present a characterization of exponential stability next.

Proposition 3.9 (Characterization of exponential stability of linear equations). Let $A \in L(H)$. Then the following statements are equivalent:

- (*i*) r(A) < 1.
- (ii) There exists $\varrho \in (0,1)$ such that for all $x \in H$ the unique solution $u \in H^{\mathbb{Z}}$ of

$$\tau u = Au + \delta_{-1}x$$

with spt $u \subseteq \mathbb{Z}_{>0}$ (c.f. Proposition 3.8 (i) \Leftrightarrow (iii)) satisfies $u \in \ell_{2,\rho}(\mathbb{Z}; H)$.

(iii) There exists $\varrho \in (0,1)$ such that for all $x \in H$ for the unique solution $u \in H^{\mathbb{Z}}$ of

$$\tau u = Au + \delta_{-1}x$$

with spt $u \subseteq \mathbb{Z}_{\geq 0}$ there is M > 0 with

$$|u_n|_H \le M\varrho^n \qquad (n \in \mathbb{Z}),$$

i.e., $u \in \ell_{\infty,\varrho}(\mathbb{Z}; H)$.

Proof. $(i) \Rightarrow (ii)$: From Theorem 3.7 we deduce that for every $r \in (r(A), 1) \neq \emptyset$ we have $u = (\tau - A)^{-1} \delta_{-1} x \in \ell_{2,\varrho}(\mathbb{Z}; H)$.

 $(ii) \Rightarrow (iii)$: This follows as $\ell_{2,\varrho}(\mathbb{Z}_{\geq 0}; H) \subseteq \ell_{\infty,\varrho}(\mathbb{Z}_{\geq 0}; H)$ for all $\varrho > 0$.

 $(iii) \Rightarrow (i)$: Let $\varrho \in (0,1)$, $x \in H$ and $u \in \ell_{\infty,\varrho}(\mathbb{Z}; H)$ as in (iii). Let $\tilde{\varrho} > \varrho$. Applying Lemma 2.4(ii) we obtain $u \in \ell_{1,\tilde{\varrho}}(\mathbb{Z}; H)$. Also $u_n = A^n x$ for $n \in \mathbb{Z}_{\geq 0}$ by Proposition 3.8. We observe that the closed graph theorem implies that the mapping

$$H \ni x \mapsto \left(A^n x\right)_{n \in \mathbb{Z}_{>0}} \in \ell_{1,\tilde{\varrho}}(\mathbb{Z}_{\geq 0}; H)$$

is continuous. We have

$$\sup_{x \in H, |x|_H \le 1} \sum_{n=0}^{\infty} |A^n x|_H \tilde{\varrho}^{-n} < \infty.$$

For $n \in \mathbb{Z}_{\geq 0}$ we find $x_n \in H$, $|x_n|_H = 1$ such that $|A^n x_n|_H \geq \frac{1}{2} |A^n|_{L(H)}$. Thus, by the monotone convergence theorem, we deduce

$$\sum_{n=0}^{\infty} |(A/\tilde{\varrho})^n|_{L(H)} \le 2\sum_{n=0}^{\infty} \sup_{k \in \mathbb{N}} |A^n x_k|_H \tilde{\varrho}^{-n}$$

$$= 2 \sup_{k \in \mathbb{N}} \sum_{n=0}^{\infty} |A^n x_k|_H \tilde{\varrho}^{-n}$$
$$\leq 2 \sup_{x \in H, |x|_H \leq 1} \sum_{n=0}^{\infty} |A^n x|_H \tilde{\varrho}^{-n} < \infty$$

Using the Neumann series, we obtain $(1-(A/\tilde{\varrho}))^{-1} \in L(H)$ and therefore $(\tilde{\varrho}-A)^{-1} \in L(H)$. We can repeat the argument for θA in place of A for all $\theta \in S_1$. Thus, $(z-A)^{-1} \in L(H)$ for all $z \in S_{\tilde{\varrho}}$. As $\varrho < 1$, we obtain r(A) < 1.

4 Stable manifolds

Stable manifolds of difference equations $u_{n+1} = Au_n + f(u_n)$, $n \in \mathbb{Z}_{\geq 0}$, with hyperbolic linear part A and f(0) = 0, can be constructed e.g. with the graph transform method or the Lyapunov–Perron method (see e.g. [1] and the references therein). In this section we extend the Lyapunov–Perron method to general difference equations $\tau u = Au + F(u)$ (Theorem 4.11).

Theorem 4.1 (Existence and uniqueness of solution in $\ell_{2,\varrho}(\mathbb{Z}; H)$ with ϱ in spectral gap). Let $A \in L(H)$, $\varrho > 0$ with $\sigma(A) \cap S_{\varrho} = \emptyset$, $F \colon \ell_{2,\varrho}(\mathbb{Z}; H) \to \ell_{2,\varrho}(\mathbb{Z}; H)$ with

$$|F|_{\operatorname{Lip}(\ell_{2,\varrho}(\mathbb{Z};H))} < 1/M_{\varrho}$$

for $M_{\varrho} := \sup_{z \in S_{\varrho}} |(z - A)^{-1}|_{L(H)} < \infty$. Then for each $x \in H$ there exists a unique $u :: \mathscr{S}_{\varrho}(x) \in \ell_{2,\varrho}(\mathbb{Z}; H)$ with

$$\tau u = Au + F(u) + \delta_{-1}x.$$

Proof. By Theorem 3.7, $(\tau - A)^{-1} \in L(\ell_{2,\varrho}(\mathbb{Z}; H))$. Using Lemma 3.4,

$$|(\tau - A)^{-1}|_{L(\ell_{2,\varrho}(\mathbb{Z};H))} = \sup_{z \in S_{\varrho}} |(z - A)^{-1}|_{L(H)} = M_{\varrho}.$$

Let $x \in H$. Then

$$\ell_{2,\varrho}(\mathbb{Z};H) \to \ell_{2,\varrho}(\mathbb{Z};H)$$
$$u \mapsto (\tau - A)^{-1} F(u) + (\tau - A)^{-1} \delta_{-1} x$$

is a contraction with unique fixed point $\mathscr{S}_{\varrho}(x) \in \ell_{2,\varrho}(\mathbb{Z}; H)$, since for $u, v \in \ell_{2,\varrho}(\mathbb{Z}; H)$

$$\left| \left[(\tau - A)^{-1} F(u) + (\tau - A)^{-1} \delta_{-1} x \right] - \left[(\tau - A)^{-1} F(v) + (\tau - A)^{-1} \delta_{-1} x \right] \right|_{\ell_{2,\varrho}(\mathbb{Z};H)}$$

$$\leq \left| (\tau - A)^{-1} \right|_{L(\ell_{2,\varrho}(\mathbb{Z};H))} |F(u) - F(v)|_{\ell_{2,\varrho}(\mathbb{Z};H)} \leq M_{\varrho} |F|_{\operatorname{Lip}(\ell_{2,\varrho}(\mathbb{Z};H))} |u - v|_{\ell_{2,\varrho}(\mathbb{Z};H)}.$$

Let $cc = cc(\mathbb{Z}; H) := \{x \in H^{\mathbb{Z}}; x_k \neq 0 \text{ for only finitely many } k \in \mathbb{Z}\}$ denote the vector space of sequences in $H^{\mathbb{Z}}$ with compact support. $cc \subseteq \ell_{p,\varrho}(\mathbb{Z}; H)$ for $1 \leq p \leq \infty, \ \varrho > 0.$

Definition 4.2 (Causality). A mapping $F: cc(\mathbb{Z}; H) \to H^{\mathbb{Z}}$ is called *causal*, if for all $u, v \in cc, a \in \mathbb{Z}$,

$$\operatorname{spt}(u-v) \subseteq \mathbb{Z}_{\geq a} \quad \Rightarrow \quad \operatorname{spt}(F(u)-F(v)) \subseteq \mathbb{Z}_{\geq a}$$

Corollary 4.3 (Solution operator for initial value problems). Let $A \in L(H)$, $\rho > r(A)$, $F: \ell_{2,\rho}(\mathbb{Z}; H) \to \ell_{2,\rho}(\mathbb{Z}; H)$ with F(0) = 0, $F|_{cc}$ causal, and

$$|F|_{\operatorname{Lip}(\ell_{2,\varrho}(\mathbb{Z};H))} < 1/M_{\varrho}$$

for $M_{\varrho} := \sup_{z \in S_{\varrho}} |(z - A)^{-1}|_{L(H)} < \infty$. Then for each $x \in H$ there exists a unique $u :: \mathscr{S}_{\varrho}(x) \in \ell_{2,\varrho}(\mathbb{Z}; H)$ with

$$\tau u = Au + F(u) + \delta_{-1}x$$

and the solution operator \mathscr{S}_{ϱ} satisfies

$$\operatorname{spt} \mathscr{S}_{\varrho}(x) \subseteq \mathbb{Z}_{\geq 0}.$$

Proof. Step 1: Show that $F: \ell_{2,\varrho}(\mathbb{Z}; H) \to \ell_{2,\varrho}(\mathbb{Z}; H)$ is causal, i.e.

$$\forall u, v \in \ell_{2,\varrho}(\mathbb{Z}; H), a \in \mathbb{Z} \colon \operatorname{spt}(u - v) \subseteq \mathbb{Z}_{\geq a} \ \Rightarrow \ \operatorname{spt}(F(u) - F(v)) \subseteq \mathbb{Z}_{\geq a}.$$

To this end, let $u, v \in \ell_{2,\varrho}(\mathbb{Z}; H)$ and $a \in \mathbb{Z}$ such that $\operatorname{spt}(u - v) \subseteq \mathbb{Z}_{\geq a}$. We define sequences $(u^{(k)})_{k \in \mathbb{N}}, (v^{(k)})_{k \in \mathbb{N}}$ in cc by

$$u_j^{(k)} \coloneqq \begin{cases} u_j & \text{if } j \in [-k,k], \\ 0 & \text{if } j \in \mathbb{Z} \setminus [-k,k], \end{cases} \quad (k \in \mathbb{N}),$$

and similarly for $v_j^{(k)}$. Then $u^{(k)} \to u$, $v^{(k)} \to v$ in $\ell_{2,\varrho}(\mathbb{Z}; H)$ and for all $k \in \mathbb{N}$ we have $\operatorname{spt}(u^{(k)} - v^{(k)}) \subseteq \mathbb{Z}_{\geq a}$. Using the fact that $F|_{\operatorname{cc}}$ is causal,

$$\operatorname{spt}(F(u^{(k)}) - F(v^{(k)})) \subseteq \mathbb{Z}_{\geq a} \qquad (k \in \mathbb{N}).$$

F is Lipschitz continuous and hence the limit $\lim_{k\to\infty} (F(u^{(k)}) - F(v^{(k)})) = F(u) - F(v)$ also satisfies $\operatorname{spt}(F(u) - F(v)) \subseteq \mathbb{Z}_{\geq a}$.

Step 2: Let $x \in H$. As in the proof of Theorem 4.1 we see that $\mathscr{S}_{\varrho}(x)$ is the unique fixed point of the contraction

$$\ell_{2,\varrho}(\mathbb{Z};H) \to \ell_{2,\varrho}(\mathbb{Z};H),$$
$$u \mapsto (\tau - A)^{-1} F(u) + (\tau - A)^{-1} \delta_{-1} x.$$

To show that spt $\mathscr{S}_{\varrho}(x) \subseteq \mathbb{Z}_{\geq 0}$, let $u \in \ell_{2,\varrho}(\mathbb{Z}_{\geq 0}; H)$ and show that the fixed point iteration preserves the support of u, i.e.

spt
$$((\tau - A)^{-1}F(u) + (\tau - A)^{-1}\delta_{-1}x) \subseteq \mathbb{Z}_{\geq 0}.$$
 (4.1)

By Theorem 3.7(*ii*) we know that $\operatorname{spt}(\tau - A)^{-1}\delta_{-1}x \subseteq \mathbb{Z}_{\geq 0}$. We have $\operatorname{spt}(u - 0) \subseteq \mathbb{Z}_{\geq 0}$ and as F(0) = 0 we have seen in Step 1 that $\operatorname{spt} F(u) = \operatorname{spt}(F(u) - F(0)) \subseteq \mathbb{Z}_{\geq 0}$. Since $(\tau - A)^{-1}$ is causal by Theorem 3.7(*i*), we deduce that $\operatorname{spt}(\tau - A)^{-1}F(u) \subseteq \mathbb{Z}_{\geq 0}$, proving (4.1).

Remark 4.4 (Riesz projection [2, Proposition 6.9]). Let $A \in L(H)$, $\gamma \in \mathbb{R}_{>0}$. If $\sigma(A) \cap S_{\gamma} = \emptyset$, then the Riesz projections

$$P_{\gamma}^{+} \coloneqq \frac{1}{2\pi i} \int_{S_{\gamma}} (z - A)^{-1} \, \mathrm{d}z \in L(H) \quad \text{and} \quad P_{\gamma}^{-} \coloneqq I - P_{\gamma}^{+} \in L(H)$$

satisfy

- (i) $(P_{\gamma}^{\pm})^2 = P_{\gamma}^{\pm}, P_{\gamma}^{+}[H] \oplus P_{\gamma}^{-}[H] = H.$
- (ii) $P_{\gamma}^{\pm}A = AP_{\gamma}^{\pm}, AP_{\gamma}^{\pm}[H] = P_{\gamma}^{\pm}[H]$, i.e. the subspaces $P_{\gamma}^{\pm}[H]$ are invariant under A.

(iii)
$$\sigma(AP_{\gamma}^+) = \sigma(A) \cap B(0,\gamma), \ \sigma(AP_{\gamma}^-) = \sigma(A) \setminus B(0,\gamma).$$

Moreover, for $\gamma_1, \gamma_2 \in \mathbb{R}_{>0}$ with $\gamma_1 < \gamma_2$,

(iv)
$$\forall \gamma \in [\gamma_1, \gamma_2] : \sigma(A) \cap S_{\gamma} = \emptyset \Rightarrow P_{\gamma_1}^{\pm} = P_{\gamma_2}^{\pm}.$$

Theorem 4.5 (Lyapunov–Perron operator). Under the assumptions of Theorem 4.1, the Lyapunov–Perron operator

$$\begin{aligned} \mathscr{L}_{\varrho} \colon H \times \ell_{2,\varrho}(\mathbb{Z}; H) &\to \ell_{2,\varrho}(\mathbb{Z}; H) \\ (x, u) &\mapsto \chi_{\mathbb{Z}_{\geq 0}}(\tau - A)^{-1}(F(u) + \delta_{-1}x) \end{aligned}$$

is well-defined and for $x \in H$, $\mathscr{L}_{\rho}(x, \cdot)$ is a contraction.

Proof. The proof is along the lines of Theorem 4.1.

Definition 4.6 (Hyperbolic). Let $A \in L(H)$. The operator A is called *hyperbolic*, if $\sigma(A) \cap S_1 = \emptyset$.

Remark 4.7. Let $A \in L(H)$ such that $\sigma(A) \subseteq \mathbb{C} \setminus B[0,1]$. Then by definition of the spectrum, $A^{-1} \in L(H)$ exists. Also $\sigma(A^{-1}) \subseteq B(0,1)$ and as the spectrum is closed, $r(A^{-1}) < 1$. Let $x \in H$ such that $(A^n x)_{n \in \mathbb{Z}_{\geq 0}} \in \ell_2(\mathbb{Z}_{\geq 0}; H)$. It follows that x = 0. Indeed there is an $N \in \mathbb{Z}_{\geq 0}$ such that for every n > N

$$|x|_{H} = |A^{-n}A^{n}x|_{H} \le |A^{-n}|_{H}|A^{n}x|_{H} \le |A^{n}x|_{H}$$

by the definition of the spectral radius. However, for $(A^n x)_{n \in \mathbb{Z}_{\geq 0}}$ to be square summable, we necessarily have $\lim_{n \to \infty} |A^n x|_H = 0$.

Lemma 4.8. Let $A \in L(H)$ be hyperbolic. We write $P := P_1^+$, $Q := P_1^-$, for the Riesz-projections. Then for every $v \in \ell_2(\mathbb{Z}; H)$ we have for $n \in \mathbb{Z}$

$$(\tau - PAP)^{-1}v = \left(\sum_{k=-\infty}^{n-1} (PAP)^{n-1-k}v_k\right)_n, (\tau - QAQ)^{-1}v = \left(-\sum_{k=n}^{\infty} (QAQ)^{n-1-k}v_k\right)_n.$$

Proof. Note that $PAP = AP = A|_{P[H]}$ on P[H] and $\sigma(A|_{P[H]}) = \sigma(A) \cap B(0,1)$ by the Riesz-projection theorem. Therefore $r(A|_{P[H]}) < 1$. Similarly $QAQ = AQ = A|_{Q[H]}$ on Q[H] and $\sigma(A|_{Q[H]}) = \sigma(A) \setminus B[0,1]$. In particular, 0 is in the resolvent set of $A|_{Q[H]}$ and $r((A|_{Q[H]})^{-1}) < 1$. By definition of the spectral radius both sums of Lemma 4.8 exist for all $v \in \ell_2(\mathbb{Z}; H)$.

For every $v \in \ell_2(\mathbb{Z}; H)$ we compute for $n \in \mathbb{Z}$

$$(\tau - PAP) \left(\sum_{k=-\infty}^{n-1} (PAP)^{n-1-k} v_k\right)_n$$
$$= \left(\sum_{k=-\infty}^n (PAP)^{n-k} v_k - \sum_{k=-\infty}^{n-1} (PAP)^{n-k} v_k\right)_n = v$$

and

$$(\tau - QAQ) \left(\sum_{k=n}^{\infty} (QAQ)^{n-1-k} v_k \right)_n$$
$$= \left(\sum_{k=n+1}^{\infty} (QAQ)^{n-k} v_k - \sum_{k=n}^{\infty} (QAQ)^{n-k} v_k \right)_n = -v.$$

The assertion follows by rearranging the terms.

Theorem 4.9 (Characterization of Lyapunov–Perron fixed point). Let $A \in L(H)$ be hyperbolic, $F: \ell_2(\mathbb{Z}; H) \to \ell_2(\mathbb{Z}; H)$ with

$$|F|_{\operatorname{Lip}(\ell_2(\mathbb{Z};H))} < 1/M_1$$

for $M_1 := \sup_{z \in S_1} |(z - A)^{-1}|_{L(H)} < \infty$. Let $\xi \in P[H]$ and $u \in \ell_2(\mathbb{Z}; H)$. Then the following statements are equivalent:

(i) u is the unique fixed point of the Lyapunov–Perron operator $\mathscr{L}_1(\xi, \cdot)$. (ii) spt $u \subseteq \mathbb{Z}_{\geq 0}$ and for $n \in \mathbb{Z}_{\geq 0}$

$$Pu_{n} = (PAP)^{n}\xi + \sum_{k=-\infty}^{n-1} (PAP)^{n-1-k} PF(u)_{k},$$
$$Qu_{n} = -\sum_{k=n}^{\infty} (QAQ)^{n-1-k} QF(u)_{k}.$$

Proof. $(i) \Rightarrow (ii)$: Let $u \in \ell_2(\mathbb{Z}; H)$ be the unique fixed point. By definition of the Lyapunov–Perron operator, it follows that $\operatorname{spt}(u) \subseteq \mathbb{Z}_{\geq 0}$. Let $n \geq 0$. We have $((\tau - A)^{-1}\delta_{-1}\xi)_n = A^n\xi$ and $P\xi = \xi$ and $Q\xi = 0$. Using the equivalence $(iii) \Leftrightarrow (v)$ of Proposition 3.8

$$Pu_n = (P\chi_{\mathbb{Z}_{\geq 0}}(\tau - A)^{-1}(F(u) + \delta_{-1}\xi))_n = (\chi_{\mathbb{Z}_{\geq 0}}(\tau - A)^{-1}PF(u))_n + (PAP)^n\xi,$$

$$Qu_n = (Q\chi_{\mathbb{Z}_{\geq 0}}(\tau - A)^{-1}(F(u) + \delta_{-1}\xi))_n = (\chi_{\mathbb{Z}_{\geq 0}}(\tau - A)^{-1}QF(u))_n,$$

as P and Q commute with all linear operators involved. Thus we have to compute $(\tau - A)^{-1}PF(u)$ and $(\tau - A)^{-1}QF(u)$ to show the first and second equation of (ii). Using the properties of the Riesz-projections, we compute

$$P(\tau - A)^{-1} = (\tau - A)^{-1}P = (\tau - PAP)^{-1}P,$$

$$Q(\tau - A)^{-1} = (\tau - A)^{-1}Q = (\tau - QAQ)^{-1}Q.$$

Therefore (*ii*) follows when setting either v = PF(u) or v = QF(u) in Lemma 4.8. (*ii*) \Rightarrow (*i*): For every $n \ge 0$ we compute

$$\begin{split} u_n &= Pu_n + Qu_n \\ &= (PAP)^n \xi + \sum_{k=-\infty}^{n-1} (PAP)^{n-1-k} PF(u)_k - \sum_{k=n}^{\infty} (QAQ)^{n-1-k} QF(u)_k \\ &= A^n \xi + \sum_{k=0}^{n-1} (PAP)^{n-1-k} PF(u)_k + \sum_{k=0}^{n-1} (QAQ)^{n-1-k} QF(u)_k \\ &+ \sum_{k=-\infty}^{-1} (PAP)^{n-1-k} PF(u)_k - \sum_{k=0}^{\infty} (QAQ)^{n-1-k} QF(u)_k \\ &= A^n \xi + \sum_{k=0}^{n-1} (PA^{n-1-k} F(u)_k + QA^{n-1-k} F(u)_k) \\ &+ A^n \left(\sum_{k=-\infty}^{-1} (PAP)^{-1-k} PF(u)_k - \sum_{k=0}^{\infty} (QAQ)^{-1-k} QF(u)_k \right) \\ &= A^n \left(\xi + \sum_{k=-\infty}^{-1} (PAP)^{-1-k} PF(u)_k - \sum_{k=0}^{\infty} (QAQ)^{-1-k} QF(u)_k \right) \\ &+ \sum_{k=0}^{n-1} A^{n-1-k} F(u)_k. \end{split}$$

By variation of constants we see that $u_{n+1} = Au_n + F(u)_n = Au_n + F(u)_n + \delta_{-1,n}\xi$ for every $n \ge 0$. That is $u_n = ((\tau - A)^{-1}(F(u) + \delta_{-1}\xi))_n$ for all n > 0. Furthermore we have

$$\left((\tau - A)^{-1}(F(u) + \delta_{-1}\xi)\right)_0 = \left((\tau - A)^{-1}(P + Q)(F(u) + \delta_{-1}\xi)\right)_0$$

$$= \left((\tau - A)^{-1} PF(u) \right)_0 + \xi + \left((\tau - A)^{-1} QF(u) \right)_0.$$

We have computed the expressions $((\tau - A)^{-1}PF(u))_0$ and $((\tau - A)^{-1}QF(u))_0$ in Lemma 4.8 and they coincide with those of u_0 . Therefore $u = \chi_{\mathbb{Z}_{\geq 0}}(\tau - A)^{-1}(F(u) + \delta_{-1}\xi)$, since $\operatorname{spt}(u) \subseteq \mathbb{Z}_{\geq 0}$.

Definition 4.10 (Admissibility condition). Let $\varrho_1, \varrho_2 > 0$. Then $F: \operatorname{cc}(\mathbb{Z}; H) \to \ell_{2,\varrho_1}(\mathbb{Z}; H) \cap \ell_{2,\varrho_2}(\mathbb{Z}; H)$ is called (ϱ_1, ϱ_2) -admissible if

$$F: \operatorname{cc}(\mathbb{Z}; H) \subseteq \ell_{2,\varrho_i}(\mathbb{Z}; H) \to \ell_{2,\varrho_i}(\mathbb{Z}; H)$$

is Lipschitz for $i \in \{1, 2\}$.

We denote the unique Lipschitz extensions of F to $\ell_{2,\varrho_i}(\mathbb{Z}; H)$ for $i \in \{1, 2\}$ again by F.

We are now in a position to prove the existence of a stable manifold of the trivial solution u = 0.

Theorem 4.11 (Stable manifold). Let $A \in L(H)$ be hyperbolic, $\rho > r(A)$. Let F be $(1, \rho)$ -admissible, such that

$$|F|_{\operatorname{Lip}(\ell_2(\mathbb{Z};H))} < 1/M_1$$
 and $|F|_{\operatorname{Lip}(\ell_{2,\varrho}(\mathbb{Z};H))} < 1/M_{\varrho}.$

We also assume that F is causal and F(0) = 0. Let $P := P_1^+$, $Q := P_1^-$ denote the Riesz projections, $T(\xi) \in \ell_2(\mathbb{Z}; H)$ the unique fixed point of the Lyapunov–Perron operator $\mathscr{L}_1(\xi, \cdot)$ for $\xi \in P[H]$. Then the stable set

$$W^s \coloneqq \{ x \in H; \mathscr{S}_o(x) \in \ell_2(\mathbb{Z}; H) \}$$

is isomorphic to the graph of $w^s \colon P[H] \to Q[H], \xi \mapsto Q(T(\xi)(0)),$

$$W^{s} = \{\xi + w^{s}(\xi) \in H; \xi \in P[H]\},\$$

i.e. W^s is a manifold, called stable manifold (of $\tau u = Au + F(u) + \delta_{-1}x$ in $\ell_{2,\varrho}(\mathbb{Z}; H)$ at the fixed point 0).

Proof. An important observation is that as F is causal and F(0) = 0, we deduce that for all $u \in \ell_2(\mathbb{Z}; H)$ with $\operatorname{spt} u = \operatorname{spt}(u - 0) \subseteq \mathbb{Z}_{\geq 0}$ we have $\operatorname{spt} F(u) = \operatorname{spt}(F(u) - F(0)) \subseteq \mathbb{Z}_{\geq 0}$. This simplifies the first sum in Theorem 4.9(*ii*).

 (\subseteq) : Let $x \in W^s$. We set $\xi \coloneqq P(x) \in P[H]$ and $\eta \coloneqq Q(x) \in Q[H]$, such that $x = \xi + \eta$. We will prove that $\eta = w^s(\xi) = Q(T(\xi)(0))$ and even more that $u \coloneqq \mathscr{S}_{\varrho}(x) = T(\xi)$ (note that $(\mathscr{S}_{\varrho}(x))_0 = x = \xi + \eta$). By Corollary 4.3 we have $\operatorname{spt}(u) \subseteq \mathbb{Z}_{\geq 0}$, and as $x \in W^s$ we know that $u \in \ell_2(\mathbb{Z}; H)$.

We prove the two equalities of Theorem 4.9(*ii*). Let $n \ge 0$. By Theorem 2.8 we have the variation of constants formula

$$u_n = A^n x + \sum_{k=0}^{n-1} A^{n-1-k} F(u)_k.$$
(4.2)

Multiplying (4.2) with P results in

$$Pu_n = (PAP)^n \xi + \sum_{k=0}^{n-1} (PAP)^{n-1-k} PF(u)_k$$
$$= (PAP)^n \xi + \sum_{k=-\infty}^{n-1} (PAP)^{n-1-k} PF(u)_k.$$

By multiplying (4.2) with Q, we compute

$$Qu_n = (QAQ)^n \eta + \sum_{k=0}^{n-1} (QAQ)^{n-1-k} QF(u)_k$$

= $(QAQ)^n \eta + \left(\sum_{k=0}^{\infty} (QAQ)^{n-1-k} QF(u)_k - \sum_{k=n}^{\infty} (QAQ)^{n-1-k} QF(u)_k\right)$
= $(QAQ)^n \left(\eta + \sum_{k=0}^{\infty} (QAQ)^{-1-k} QF(u)_k\right) - \sum_{k=n}^{\infty} (QAQ)^{n-1-k} QF(u)_k).$

By Lemma 4.8 we know that $\left(\sum_{k=n}^{\infty}(QAQ)^{n-1-k}QF(u)_k)\right)_n=(\tau-QAQ)^{-1}QF(u)\in \mathbb{R}$ $\ell_2(\mathbb{Z}; H)$ and so

$$\left((QAQ)^n \left(\eta + \sum_{k=0}^{\infty} (QAQ)^{-1-k} QF(u)_k \right) \right)_n \in \ell_2(\mathbb{Z}; H).$$

In Remark 4.7 we have noted that necessarily $\eta + \sum_{k=0}^{\infty} (QAQ)^{-1-k} QF(u)_k = 0$. By Theorem 4.9, u is the unique fixed point of the Lyapunov–Perron operator $\mathscr{L}_1(\xi, \cdot)$.

 $(\supseteq): \text{ Let } (\xi,\eta) \in \text{graph}(w^s), \text{ that is } \xi \in P[H] \text{ and } Q(T(\xi)_0) = -\sum_{k=0}^{\infty} (QAQ)^{-1-k}.$ $QF(u)_k = \eta \text{ by Theorem 4.9}(ii). \text{ We will verify that } \xi + \eta \in W^s, \text{ that is } \mathscr{S}_{\varrho}(\xi + \eta) \in QF(U).$ $\ell_2(\mathbb{Z}; H)$. To this end we show that $u := T(\xi) = \mathscr{S}_{\rho}(\xi + \eta)$. By the variation of constants formula (4.2) we know that

$$u = \mathscr{S}_{\varrho}(\xi + \eta) \quad \Leftrightarrow \quad u \in \ell_{2,\varrho}(\mathbb{Z}; H) \text{ and } \operatorname{spt}(u) \subseteq \mathbb{Z}_{\geq 0}$$

and $u_n = A^n(\xi + \eta) + \sum_{k=0}^{n-1} A^{n-1-k} F(u)_k$

By Theorem 4.9(*ii*) we have $spt(u) \subseteq \mathbb{Z}_{\geq 0}$. Therefore if $\rho > 1$ we see that $u \in \mathbb{Z}_{\geq 0}$ $\ell_2(\mathbb{Z}_{\geq 0}; H) \subseteq \ell_{2,\varrho}(\mathbb{Z}_{\geq 0}; H)$. If $\varrho < 1$ the statement of the theorem is trivial as $W^s = H$ on the one hand and P = I and Q = 0 on the other hand. It remains to establish the formula for u_n . For this, we see that Theorem 4.9(*ii*) also yields

$$u_n = Pu_n + Qu_n$$

= $(PAP)^n \xi + \sum_{k=-\infty}^{n-1} (PAP)^{n-1-k} PF(u)_k - \sum_{k=n}^{\infty} (QAQ)^{n-1-k} QF(u)_k$

ъ

$$\begin{split} &= A^{n}\xi + P\sum_{k=0}^{n-1} A^{n-1-k}F(u)_{k} + Q\sum_{k=0}^{n-1} A^{n-1-k}F(u)_{k} - \sum_{k=0}^{\infty} (QAQ)^{n-1-k}QF(u)_{k} \\ &= A^{n}\xi + \sum_{k=0}^{n-1} A^{n-1-k}F(u)_{k} - A^{n}\sum_{k=0}^{\infty} (QAQ)^{-1-k}QF(u)_{k} \\ &= A^{n}\left(\xi - \sum_{k=0}^{\infty} (QAQ)^{-1-k}QF(u)_{k}\right) + \sum_{k=0}^{n-1} A^{n-1-k}F(u)_{k} \\ &= A^{n}(\xi + \eta) + \sum_{k=0}^{n-1} A^{n-1-k}F(u)_{k}. \end{split}$$

References

- [1] S. Elaydi. An Introduction to Difference Equations. Undergraduate Texts in Mathematics. Springer New York, third edition, 2005.
- [2] P. Hislop and I. Sigal. Introduction to spectral theory. With applications to Schrödinger operators. New York, NY: Springer-Verlag, 1996.
- [3] A. Kalauch, R. Picard, S. Siegmund, S. Trostorff, and M. Waurick. A Hilbert space perspective on ordinary differential equations with memory term. *Journal* of Dynamics and Differential Equations, 26(2):369–399, 2014.
- [4] Y. Katznelson. An introduction to harmonic analysis. Cambridge: Cambridge University Press, 3rd edition, 2004.
- [5] R. Picard, S. Trostorff, and M. Waurick. A functional analytic perspective to delay differential equations. Operators and Matrices, 8(1):217–236, 2014.
- [6] R. H. Picard and D. F. McGhee. Partial Differential Equations: A Unified Hilbert Space Approach. De Gruyter expositions in mathematics. De Gruyter, 2011.