Syzygies for periodic orbits in the restricted three-body problem

Robert Nicholls

Abstract

In this paper we show the existence of syzygies for all periodic orbits inside the bounded Hill's region of the planer circular restricted three-body problem with energy below the second critical value. The proof will follow some ideas of Birkhoff [2] to compute the roots of partial derivatives of the effective potential. Birkhoff's methods are extended to higher energies and a new base case is created and shown to fulfil the requirements. An other step from Birkhoff is scrutinized to continue the statement to all mass ratios. The final step is achieved by integrating over periodic orbits. Applying the same methods to Hill's lunar problem delivers similar results in that setting as well.

1 Introduction

Going back to Poincaré, the planar circular restricted three-body problem (PCR3BP) is one of the oldest and most studied simplifications of an N-body problem. This paper presents a proof that under certain conditions all periodic orbits have syzygies. In view of Birkhoff's conjecture that the retrograde bounds a disc-like global surface of section, one would like to show that the retrograde is the systole. The result from this work proves the non-existence of periodic orbits without syzygies and thus reduces the complexity of the search for further periodic orbits by one dimension.

An other possible application, which this paper could be a starting point for, is the development of symbolic dynamics for the PCR3BP. Symbolic dynamics in three symbols has been successfully applied to the Euler problem of two centres in [3] and the existence of an infinite sequence of syzygies has been proven for every—exept Lagrange's—solution in the general three-body problem with zero angular momentum in [9] and [10]. As soon as one adds angular momentum there are periodic non-collisional Langrangian solutions without syzygies. However, if the angular momentum is small enough then sequences of syzygies derived by free homotopy classes on the reduced configurations space are realized by periodic solutions [8] and symbolic dynamics has at least numerically been constructed for special types of orbits in a certain PCR3BP situation by [13] and [14]. For the general case of the PCR3BP the existence of syzygies for periodic orbits is in general not true. In [2] Birkhoff proved a very helpful statement about the roots of the partial derivatives of the effective potential, which we will use in this work. We will also extend his main argument to higher energies and finally prove the existence of syzygies for periodic orbits within the bounded Hill's region below the second critical value. In the preparation for the proof we will mainly use notations and arguments from [4], which were slightly adapted and summarized in [11], to also cover the proof of the statement below the first critical level.

The structure of this work will be the following: First we will go through a short introduction to the PCR3BP in section 2 and then in section 3 recall some general definitions and facts about the PCR3BP as preparation for the proof. Since we will be using some elementary Morse theory to formalize certain steps from [2], we will also state the required definitions and lemmata. From there on, we will go through two base cases in section 4: The first basically comes directly from [2] and only achieves the weaker statement below the first critical value, while the second case is new and its continuation covers all energies below the second critical value (in particular it also covers the first case). The continuation of these bases cases in section 5 will cover all mass ratios and all stated energies to prove the main result

Theorem 1:

Every periodic orbit of the PCR3BP inside the bounded Hill's region \Re^b_c for an energy $c < H(L_2), H(L_3)$ below the second critical value has at least two distinct syzygies during each period.

2 The planar circular restricted 3-body problem

The restricted problem of three bodies has been widely studied, so we will only do a very brief introduction here. For further and more thorough inspection of the equations refer to works such as [5] or [12].

The PCR3BP is the dynamics of a particle—in our case the moon—attracted by and moving in the same plane as the two primaries—here called the sun and the earth—which we assume to have circular motion around their common centre of mass as a Keplerian solution. We normalize the masses to be $0 < \mu < 1$ for the earth and $1 - \mu$ for the sun. By introducing rotating coordinates the positions of the primaries become stationary at $(-\mu, 0) = s$ and $(1 - \mu, 0) = e$ and the Hamiltonian becomes autonomous:

$$H(q,p) = \frac{1}{2}((p_1 + q_2)^2 + (p_2 - q_1)^2) \underbrace{-\frac{\mu}{|q-e|} - \frac{1-\mu}{|q-s|} - \frac{1}{2}q^2}_{=:V(q)}$$
(1)

Here the effective potential energy, as the part independent of momentum, is denoted as V(q). By Hamilton's equations of motion one gets the following second order ODEs

$$\ddot{q}_1 = 2\dot{q}_2 - \frac{\partial V}{\partial q_1} \tag{2}$$

$$\ddot{q}_2 = -2\dot{q}_1 - \frac{\partial V}{\partial q_2}.\tag{3}$$

In [2] Birkhoff uses the alternative potential function

$$\Omega(q) := \frac{1}{2} \left((1-\mu) \cdot |q-s|^2 + \mu \cdot |q-e|^2 \right) + \frac{\mu}{|q-e|} + \frac{1-\mu}{|q-s|},\tag{4}$$

which only depends on the distances of the moon to sun and earth. This potential function only differs from V by sign and a constant: $\Omega(q) = -V(q) + \mu(1-\mu)/2$, so adjusting the signs in the ODEs will render the same dynamics. It has the advantage that it makes it easy to find critical points through a transformation we will use in section 3.1.

3 Preparations for the proof

In this section we will first recall some notations for periodic solutions in general, compute critical points and Hill's regions of the PCR3BP and then prove some elementary lemmata from Morse theory.

In a symplectic manifold (M, ω) with an autonomous Hamiltonian H a periodic orbit is a solution $x \in C^{\infty}(\mathbb{R}, M)$ to the ODE

$$\frac{\mathrm{d}}{\mathrm{d}t}x(t) = X_H(x(t)) \qquad \forall t \in \mathbb{R},$$

such that there exists a period T > 0 for which x(T + t) = x(t) for all $t \in \mathbb{R}$. Here X_H is the Hamiltonian vectorfield defined implicitly by $dH = \omega(\cdot, X_H)$. For every non-trivial—that is non-constant—periodic orbit x there exists a minimal period $T_x := \min\{T > 0 \mid x(T + t) = x(t) \quad \forall t \in \mathbb{R}\}$ and every period $T = nT_x$ is just a natural multiple. In our case—the symplectic manifold arises as the cotangent bundle of an open subset of \mathbb{R}^2 endowed with the standard symplectic form—a periodic orbit is a smooth map $x \in C^{\infty}(\mathbb{R}, \mathbb{R}^2 \setminus \{e, s\})$ solving (2) and (3) such that x(T + t) = x(t) and thus also $\dot{x}(T + t) = \dot{x}(t)$ for all $t \in \mathbb{R}$. Since the value of the autonomous Hamiltonian is constant along its flow we can assert an energy H(x) to every periodic orbit.



Figure 1: Transformation of the upper half-plane (adapted from [1])

3.1 Critical points and Hill's regions in the PCR3BP

We will now compute the critical points and values of the PCR3BP as we will need to refer back to some steps during the main proof later on. This part will mainly follow [1] and [4]. Critical points of the effective potential and thus of the Hamiltonian are the five Lagrange points, which I will denote by

$$\pi_{|\operatorname{crit}(H)}(L_i) =: \ell_i \quad \text{for} \quad i = 1, \dots 5,$$

where

$$\pi_{|\operatorname{crit}(H)} \colon \operatorname{crit}(H) \to \operatorname{crit}(V) \quad \text{with} \\ \left(\pi_{|\operatorname{crit}(H)}\right)^{-1} (q_1, q_2) = ((q_1, q_2), (-q_2, q_1)) \in \operatorname{crit}(H).$$
(5)

is a bijection of critical points of H and V, given by the footpoint projection of the cotangent bundle. In this 1 to 1 correspondence also the critical values and the Morse indices (the number of negative eigenvalues of the Hessian) of critical points coincide, as the twisted momentum only adds two positive eigenvalues to the Hessian at this point. Using the symmetry of Ω with respect to the q_1 -axis to our advantage, we will first search for critical points on the upper half plane and then on the q_1 -axis by restricting the effective potential.

On the upper half plane we use a transformation to the half strip (see figure 1) by

$$\begin{split} \Phi \colon \mathbb{H}_+ &\to \Theta \\ q \mapsto \left(|q-s|, |q-e| \right), \end{split}$$

where Θ is the diagonal half-strip in the first quadrant of \mathbb{R}^2 :

$$\Theta := \left\{ (\sigma, \rho) \in (0, \infty)^2 \mid \sigma + \rho > 1, |\sigma - \rho| < 1 \right\}$$

We will now look at our effective potential in the new coordinates (σ, ρ) . Using the alternative form Ω of the effective potential from (4), this representation is made very easy as it becomes

$$\begin{split} U &:= \Omega \circ \Phi^{-1} \colon \Theta \to \mathbb{R} \\ U(\sigma, \rho) &= \frac{1}{2} \left((1-\mu)\sigma^2 + \mu \rho^2 \right) + \frac{\mu}{\rho} + \frac{1-\mu}{\sigma} \end{split}$$

Critical points of U can now be found by differentiating:

$$\mathrm{d}U(\sigma,\rho) = \frac{(1-\mu)\left(\sigma^3-1\right)}{\sigma^2}\,\mathrm{d}\sigma + \frac{\mu\left(\rho^3-1\right)}{\rho^2}\,\mathrm{d}\rho \stackrel{!}{=} 0,$$

thus the only critical point of U is (1,1). Computing the Hessian of U we get

$$\operatorname{Hess}_{U}(1,1) = \begin{pmatrix} 3(1-\mu) & 0\\ 0 & 3\mu \end{pmatrix}$$

and our single critical point of U is a local minimum.

Transforming back to coordinates in the plane, the critical point has the same distance $r_e = r_s = 1$ from sun and earth and thus forms an equilateral triangle with the primaries. We therefore get the critical point

$$\ell_4 := \Phi^{-1}(1,1) = \left(\frac{1}{2} - \mu, \frac{\sqrt{3}}{2}\right),$$

which is a local minimum of Ω . Recall, that $V(q) = -\Omega(q) + \mu(1-\mu)/2$, so ℓ_4 is a local maximum, i. e. a point with maximal Morse index 2 of the effective potential for all $\mu \in (0, 1)$. The Morse index is invariant under the bijection (5), so L_4 becomes a critical point of Morse index 2 of H, i. e. only a saddle point.

By reflection on the q_1 -axis, we get another local maximum of V at

$$\ell_5 := \left(\frac{1}{2} - \mu, -\frac{\sqrt{3}}{2}\right).$$

The Hamiltonian at these critical points takes the same critical value as on V, i. e.

$$H(L_4) = H(L_5) = V(\ell_4) = V(\ell_5) = \frac{\mu(1-\mu) - 3}{2}$$

The three remaining collinear Lagrange points are attained by restricting the effective potential V to the q_1 -axis. Here we will use the original effective potential V. Critical points of this one-dimensionally restricted function are then also critical points of the general function as V is symmetric with respect to the q_1 -axis. Therefore, let

$$u := V|_{\mathbb{R} \setminus \{s,e\}} : \mathbb{R} \setminus \{-\mu, 1-\mu\} \to \mathbb{R}$$
$$x \mapsto -\frac{\mu}{|x-(1-\mu)|} - \frac{1-\mu}{|x+\mu|} - \frac{x^2}{2}.$$
(6)

Finding explicit formulas for the critical points would mean solving quintic equations dependent on μ . These can be found in Chapter 10 of [1]. For our purposes though, it suffices to know in which of the three open intervals they lie and how their energies compare. By differentiating utwice, we get

$$u''(x) = -\frac{2\mu}{|x - (1 - \mu)|^3} - \frac{2(1 - \mu)}{|x + \mu|^3} - 1 < 0,$$

i. e. u is strictly concave. As u tends towards $-\infty$ for $x \to -\infty, -\mu, 1-\mu$ as well as ∞ , we can state that there exist exactly three local maxima of u

$$\ell_3 \in (-\infty, -\mu)$$
 $\ell_1 \in (-\mu, 1-\mu)$ $\ell_2 \in (1-\mu, \infty),$

lying in each of the connected components of the domain as can be seen in figure 2.

These critical points of V are saddle points, i. e. have Morse index 1, as proven by topological arguments in [4, chapter 5, lemma 4.2] or by computing the Hessian of Ω in [1, chapter 10.2]. Furthermore, this proves, that the pair of critical points ℓ_4 and ℓ_5 are global maxima of the effective potential, in view of V tending towards negative infinity at e, s and for large |q|.

As we have not specified the exact coordinates of the critical points, we can also not state the exact critical energies. However, we will show in which way they are ordered by following [4]. For a point $-\mu < x < 1 - \mu$ denote by

$$\rho := \rho(x) := (1 - \mu) - x = |x - e| \in (0, 1)$$

the distance of this point to earth again. We will compute the values of the effective potential at the points x and $x' := (1 - \mu) + \rho$, which lies symmetrically to x on the other side of e, by



Figure 2: The critical points of the effective potential when restricted to the q_1 -axis

using the restricted function u from (6):

$$u(x) - u(x') = -\frac{2(1-\mu)\rho^3}{1-\rho^2} < 0$$

So the values of u and thus of V in between s and e are always smaller than symmetrically on the other side of e. This holds especially for the point ℓ_1 and its point opposite of earth $\ell'_1 := (1 - \mu) + \rho(\ell_1)$, so their values are ordered by

$$V(\ell_1) = u(\ell_1) < u(\ell_1) \le u(\ell_2) = V(\ell_2),$$

as ℓ_2 was the maximum of u for all points larger than $1 - \mu$. Analogously by estimating the difference of u for a point $x \in (-\mu, 1 - \mu)$ and its opposite to the sun, we get

$$V(\ell_1) = u(\ell_1) < u(\ell_3) = V(\ell_3).$$

Via the identification (5) of critical points of V and H retaining the values we have

$$H(L_1) < H(L_2), H(L_3).$$

One can furthermore show, that if the sun is strictly heavier than earth, then $H(L_2)$ is strictly less than $H(L_3)$ and they are equal exactly at $\mu = 1/2$. We summarise the results so far in the following lemma:

Lemma 2:

For all $\mu \in (0, 1)$ there are five critical points of the Hamiltonian H. They are all saddle points of H. The pair of symmetric critical points ℓ_4 and ℓ_5 are global maxima of V, i. e. have Morseindex 2 with coordinates

$$\ell_4 = \left(\frac{1}{2} - \mu, \frac{\sqrt{3}}{2}\right) \quad \ell_5 = \left(\frac{1}{2} - \mu, -\frac{\sqrt{3}}{2}\right) \quad and \quad H(L_4) = H(L_5) = \frac{\mu(1-\mu) - 3}{2}.$$

The remaining three critical points lie along the q_1 -axis at

$$\ell_3 < -\mu, \quad -\mu < \ell_1 < 1 - \mu \quad and \quad \ell_2 > 1 - \mu,$$

and are saddle points of V, i. e. their Morse indices are 1 and their energies are ordered by

$$H(L_1) < H(L_2), H(L_3) < H(L_4) = H(L_5).$$

With this information, we can now state how the space of all accessible positions changes as the energy decreases from infinity. This set is known as the Hill's region and is defined as the projection of the energy level set $\Sigma_c := H^{-1}(c) \subset T^*(\mathbb{R}^2 \setminus \{e, s\})$ from phase space onto



Figure 3: Ovals of zero velocity and critical points in the restricted three-body problem

configuration space:

$$\mathfrak{K}_c := \pi(\Sigma_c) = \{ q \in \mathbb{R}^2 \setminus \{e, s\} \mid V(q) \le c \}$$

From (1) we see that the Hill's region corresponds to the sublevel set of the effective potential. In Birkhoff's work [2] the notion of the oval of zero velocity is used, which denotes the boundary of the Hill's region excluding the primaries or equivalently the level set of the effective potential:

$$\mathfrak{O}_c := \partial \mathfrak{K}_c \setminus \{e, s\} = V^{-1}(c)$$

As the gradient $\nabla V(q)$ is a normal vector on the level set \mathfrak{O}_c for every $q \in V^{-1}(c)$ and regular value c, the tangent to the oval of zero velocity is spanned by

$$\left\langle \left(\frac{\partial V}{\partial q_2}(q), -\frac{\partial V}{\partial q_1}(q)\right) \right\rangle_{\mathbb{R}} = \nabla V(q)^{\perp}.$$

Since H is symmetric with respect to the q_1 -axis, so is \mathfrak{K}_c and \mathfrak{O}_c , and from the type and location of critical points of V we can state the changes in topology of the Hill's region as the energy varies. Maxima of V are attained at ℓ_4 and ℓ_5 , so energies $c \ge V(\ell_4) = V(\ell_5) = (\mu(1-\mu)-3)/2$ above the common critical value result in the Hill's region being all of $\mathbb{R}^2 \setminus \{e, s\}$, i. e. movement is possible everywhere apart from collisions with the primaries. At energies below this value, holes will appear around ℓ_4 and ℓ_5 . These will grow larger with decreasing energy until at the next critical value, they connect to a single hole on the far side of the heavier primary. If the sun is strictly heavier than earth, i. e. $0 < \mu < 1/2$ then this will be at ℓ_3 . This horseshoe shaped hole will grow larger, as c declines further until its ends meet at the critical point on the far side of the lighter primary, in our case at ℓ_2 . At this point, we now have two connected components: One bounded peanut-shaped component $\mathfrak{K}^b_c = \pi(\Sigma^b_c)$ containing e and s in its closure and one unbounded outer component $\mathfrak{K}^{u}_{c} = \pi(\Sigma^{u}_{c})$. As the energy declines further yet, the bounded component narrows between the two primaries until they disconnect after surpassing the final critical value of ℓ_1 . Finally, for all energies below $V(\ell_1)$ the Hill's region is divided into three connected components: A punched disc-like component surrounds each one of the two primaries and will be denoted by $\mathfrak{K}_c^e = \pi(\Sigma_c^e)$ and $\mathfrak{K}_c^s = \pi(\Sigma_c^s)$, whereas the unbounded part will remain \mathfrak{K}_c^u . These changes of the corresponding ovals of zero velocity are visualized in figure 3 for $\mu = 0.2$. Notations of the oval describing the bounded and unbounded parts will be analogous to the Hill's regions.

3.2 Some elementary statements about Morse functions

A smooth function is called Morse if all critical points are non-degenerate, i. e. if the Hessian at this point is invertible. As mentioned before the Morse index of a critical point is the number of negative eigenvalues of the Hessian and both non-degeneracy and Morse index of critical points are invariant under coordinate change, thus can be defined for general smooth functions $f \in C^{\infty}(M, \mathbb{R})$ on manifolds. From these simple definitions we get that all critical points are isolated, which can be easily seen using the Morse Lemma, which states that there exist local coordinates u around a critical point p, such that u(0) = p and

$$f(u) = f(p) - \sum_{i=1}^{\mu_p} u_i^2 + \sum_{i=\mu_p+1}^n u_i^2,$$

where μ_p is the Morse index of p. Obviously the only critical point in this chart is p.

Lemma 3:

Let M be a smooth manifold of dimension n and $f: M \to \mathbb{R}$ a Morse function. Then all critical points of f are isolated, i. e. for every $p \in \operatorname{crit} f$ there exists an open neighbourhood U of p, such that $U \cap \operatorname{crit} f = \{p\}$.

As a direct corollary we get that Morse functions on closed Manifolds have finitely many critical points:

Corollary 4:

Let M be a closed differentiable manifold and f a Morse function, then the number of critical points of f is finite.

The last lemma we will need as preparation for the theorem, is essentially a step in Birkhoff's proof [2, chapter 17], which he claimed to be obvious. In the situation of the proof, it is indeed very apparent. A rigorous proof of a more general statement can be achieved by using the implicit function theorem:

Lemma 5:

Let M be a closed Manifold and $f : \mathbb{R} \times M \to \mathbb{R}$ a smooth 1-parameter family of Morse functions. Then the number of critical points of $f_r := f(r, \cdot)$ is constant.

Proof: We show, that the function $c \colon \mathbb{R} \to \mathbb{N}$ whereas $c(r) := \# \operatorname{crit} f_r$, is locally constant:

Let $r_0 \in \mathbb{R}$ be a real number, then the number of critical points $\#\operatorname{crit} f_{r_0} =: N < \infty$ is finite by corollary 4, because f_{r_0} is Morse and M closed. We will call theses critical points $\{p_1, \ldots, p_N\} =$ $\operatorname{crit} f_{r_0}$. The proof, that there exists an open neighbourhood of r_0 in \mathbb{R} on which the number of critical points is exactly N, will be split into two steps: First, we will show, that in an open neighbourhood the number of critical points is at least N. After that, we precede to prove, that c(r) is at most N close to r_0 .

For both steps we will use the implicit function theorem, therefore, we will first apply it to our situation: Working in a chart again, we know, that all critical points p_i of f_{r_0} satisfy $Df_{r_0}(p_i) = 0$. Since f_{r_0} is Morse, we also know, that the Hessian matrix $H_{f_{r_0}}(p_i)$ is invertible for all $i = 1, \ldots, N$, hence the requirements of the implicit function theorem are fulfilled, such that we can state:

There exist open neighbourhoods U_i of p_i , positive $\epsilon_i > 0$ and unique maps

$$y_i \colon (r_0 - \epsilon_i, r_0 + \epsilon_i) \to U_i, \quad \text{such that} \\ y_i(r_0) = p_i \quad \text{and} \\ \forall r \in (r_0 - \epsilon_i, r_0 + \epsilon_i), p \in U_i \colon y_i(r) = p \iff Df_{r_0}(p) = 0.$$

Without loss of generality we can assume $U_i \cap U_j = \emptyset$ if $i \neq j$. Step 1 $(\exists \epsilon > 0 : \forall r \in (r_0 - \epsilon, r_0 + \epsilon) : \# \operatorname{crit} f_r \geq N)$:

Set $\epsilon := \min\{\epsilon_1, \ldots, \epsilon_N\}$, then there are N distinguished critical points at $y_i(r) \in U_i$, since the open sets U_i are disjoint.

Step 2 $(\exists \epsilon > 0 : \forall r \in (r_0 - \epsilon, r_0 + \epsilon) : \# \operatorname{crit} f_r \leq N)$:

Suppose not, then for all $n \in \mathbb{N}$ there exists some $r_n \in (r_0 - 1/n, r_0 + 1/n)$ such that there are strictly more than N critical points of f_{r_n} on M. Since the functions y_i are unique, we can choose a critical point $p^{(n)}$ for every $n \in \mathbb{N}$, such that $p^{(n)} \notin \bigcup_{i=1}^N U_i$. This gives us a sequence of critical points in $M' := M \setminus \bigcup_{i=1}^N U_i$. M and therefore also M' are compact, so there exists a converging subsequence $p^{(n_k)} \to p_0 \in M'$, i. e. $p_0 \neq p_i$ for $1 \leq i \leq N$. We can now compute the limit

$$0 = \lim_{k \to \infty} \underbrace{Df_{r_{n_k}}\left(p^{(n_k)}\right)}_{=0} = Df_{r_0}\left(p_0\right)$$

and get another critical point p_0 of f_{r_0} , contradicting the assumption, that there were N critical points to start with. This concludes the proof, that $\#\operatorname{crit} f_r$ is locally constant, implying it is constant on all of \mathbb{R} .

With this Lemma, we are now ready to start to prove the main theorem of this work.

4 Base cases

The first base case is for an energy below the first critical value and comes directly from Birkhoff [2]. It is included here in order to complete the review of Birkhoff's original proof. The second base case extends this proof to all energies below the second critical value and thus enables a stronger statement with the possibility of more interesting syzygy sequences. What we want to show for each base case is that there are exactly two vertical tangents to the bounded part of the oval of zero velocity and them being the ones induced by symmetry along the q_1 -axis. Recall from the preparation that a vertical tangent corresponds to the derivative $V_{q_2} := \partial V/\partial q_2$ of the effective potential with respect to q_2 vanishing while $V_{q_1} := \partial V/\partial q_1$ remains non-zero.

4.1 Below the first critical value

Here we consider the limiting cases $\mu = 0$ and 1, i. e. where one of the primaries has zero mass. In this situation the PCR3BP turns into the rotating Kepler problem, i. e. the Kepler problem in rotating coordinates:

$$H(q,p) = \frac{1}{2} \left((p_1 + q_2)^2 + (p_2 - q_1)^2 \right) - \frac{1}{|q|} - \frac{1}{2} q^2$$

= $\frac{1}{2} p^2 - \frac{1}{|q|} + q_2 p_1 - q_1 p_2$ (7)

Note, that this Hamiltonian is no longer a Morse function, since the critical set is the unit circle. In this simple case, the Hill's regions are either all of the plane minus the origin for energies above the critical value c = -1.5, or a punched disc for the bounded Hill's region and the plane minus a larger disc as the unbounded part below the critical value. Vertical tangents of the ovals of zero velocity obviously only lie along the q_1 -axis since $V_{q_2} = q_2(1/|q|^3 - 1)$ vanishes at $q_2 = 0$ and the bounded component has |q| < 1 making $(1/|q|^3 - 1)$ strictly positive. V_{q_1} can not vanish simultaneously as there are no other critical points.

4.2 Between the first and the second critical value

The second base case will use the symmetry for $\mu = 1/2$ and an energy just above the first critical value. Here, the Hill's region is still reasonably small and we can show, that we can enclose the Hills region in a small enough neighbourhood of the origin, such that within this region the partial derivative V_{q_2} can not vanish.

At $\mu = 1/2$ the effective potential becomes

$$V(q) = -\frac{1}{2|q - (\frac{1}{2}, 0)|} - \frac{1}{2|q + (\frac{1}{2}, 0)|} - \frac{q^2}{2}$$



Figure 4: Estimates of |q - e|, |q - s| and |q|

which is additionally symmetric with respect to the q_2 -axis. Hence, it suffices to show that on the first quadrant the partial derivative V_{q_2} does not vanish outside of $\{q_2 = 0\}$. We compute the derivative

$$\mathrm{d}V(q) = \left(\frac{q_1 - \frac{1}{2}}{2|q - e|^3} + \frac{q_1 + \frac{1}{2}}{2|q - s|^3} - q_1\right)\mathrm{d}q_1 + q_2\left(\frac{1}{2|q - e|^3} + \frac{1}{2|q - s|^3} - 1\right)\mathrm{d}q_2$$

to check that (0,0) is the critical point ℓ_1 and takes the value V(0) = -2. Next, we estimate the value of V on the unit circle by using the symmetry and restricting to the first quadrant as shown in figure 4a:

$$V(q) \ge -\frac{1}{2\frac{1}{2}} - \frac{1}{2\frac{\sqrt{5}}{2}} - \frac{1}{2} > -2 \qquad \forall |q| = 1 \ q_1, q_2 \ge 0$$

Since e and s are contained in $B_1(0)$, there exists some $\epsilon_0 > 0$ such that the single bounded component of the Hill's region $\mathfrak{K}^b_{-2+\epsilon_0} \subset \{V(q) \leq -2 + \epsilon_0\}$, which has e and s in its closure, is contained within $B_1(0)$. Additionally, we compute that inside the unit ball the q_2 -values can be restricted to 2/3 by using lower estimates of |q - e| and |q - s|, and an upper estimate of |q| as shown in figure 4b:

$$V(q) > -\frac{1}{2 \cdot \frac{2}{3}} - \frac{1}{2 \cdot \frac{5}{6}} - \frac{1}{2}$$

= $-\frac{37}{20} > -2$ $\forall |q| < 1, q_2 \ge \frac{2}{3}, q_1 \ge 0$

So, there exists some $\epsilon_0 \geq \epsilon > 0$ such that $\Re^b_{-2+\epsilon}$ is contained in $B_1(0) \cap \{|q_2| < 2/3\}$. This concludes the step for restricting the Hill's region and we continue by excluding the set of zeros of the partial derivative V_{q_2} from this region.

Here, we estimate as in figure 4c, again using symmetry to restrict to the first quadrant, to get

$$\frac{1}{2|q-e|^3} + \frac{1}{2|q-s|^3} > \frac{1}{2\frac{125}{216}} + \frac{1}{2\frac{27}{8}} = \frac{3416}{3375} > 1.$$

We can conclude that inside of $B_1(0) \cap \{|q_2| < 2/3\}$ the partial derivative V_{q_2} only vanishes along the q_1 -axis. Summarizing the results of this section we state:

Lemma 6:

There exists some $\epsilon > 0$ such that inside the bounded Hill's region $\Re^{b}_{-2+\epsilon}$ for $\mu = 1/2$ the partial derivative V_{q_2} only vanishes along the q_1 -axis.

5 Continuation to all mass ratios

The final part of the proof is to continue the base case(s) to all mass ratios and to all energies below the second critical value. We will discuss Birkhoff's continuation below the first critical value along the way, but focus on the proof of the main (and stronger) statement, which is:

Lemma 7:

Inside the bounded Hill's region \mathfrak{K}_c^b for energy $c < H(L_2), H(L_3)$ below the second critical value and for all mass ratios $0 < \mu < 1$, the roots of $V_{q_2}(q)$ are precisely the points on the q_1 -axis.

Proof: Obviously all points of $\{q_2 = 0\}$ are zeros of

$$V_{q_2,\mu}(q) = q_2 \left(\frac{\mu}{|q-e|^3} + \frac{1-\mu}{|q-s|^3} - 1\right) =: q_2 W_{\mu}(q)$$
(8)

It remains to show, that inside of \mathfrak{K}^b_c there are no roots of W_{μ} .

The first claim is that 0 is a regular value: We prove this by computing where critical points lie and comparing that to the values attained in this region. The differential of W_{μ} is

$$\mathrm{d}W_{\mu}(q) = -3\left(\frac{\mu(q_1-1+\mu)}{|q-e|^5} + \frac{(1-\mu)(q_1+\mu)}{|q-s|^5}\right)\mathrm{d}q_1 - 3q_2\left(\frac{\mu}{|q-e|^5} + \frac{1-\mu}{|q-s|^5}\right)\mathrm{d}q_2$$

and thus a critical point requires q_2 and simultaneously $\mu(q_1-1+\mu)/|q-e|^5+(1-\mu)(q_1+\mu)/|q-s|^5$ to vanish. This does not happen if $|q_1| \ge 1$, since if $q_1 \ge 1$ then the latter term is bound from below by

$$\frac{\mu^2}{|q_1-1+\mu|^5} + \frac{1-\mu^2}{|q_1+\mu|^5} > 0$$

which is strictly positive. Analogously if $q_1 \leq -1$ then the later term is bound from above by

$$\frac{\mu(-2+\mu)}{|q_1-1+\mu|^5} - \frac{(1-\mu)^2}{|q_1+\mu|^5} < 0$$

So, critical values can only exist if $|q_1| < 1$ and $q_2 = 0$ for all mass ratios $\mu \in [0, 1]$. Estimating the values of W_{μ} in this region gives

$$\frac{\mu}{|q_1 - 1 + \mu|^3} + \frac{1 - \mu}{|q_1 + \mu|^3} - 1 > 0.$$

This follows by checking the cases $-1 < q_1 < -\mu$, $-\mu < q_1 < 1 - \mu$ and $1 - \mu < q_1 < 1$. So the claim is proven that 0 is a regular value of W_{μ} for all μ and thus the preimage $W_{\mu}^{-1}(0)$ is a smooth 1-manifold diffeomorphic to $W_0^{-1}(0) = S^1$.

smooth 1-manifold diffeomorphic to $W_0^{-1}(0) = S^1$. Suppose for some $\mu \in (0,1)$ and $c < V_{\mu}(\ell_2)$ that $\mathfrak{K}^b_{\mu,c} \cap W^{-1}_{\mu}(0) \neq \emptyset$, i. e. for some point inside the bounded Hill's region the factor W_{μ} of $V_{q_2,\mu}$ vanishes. Choose a smooth path (figure 5)

$$\begin{split} \gamma \colon [0,1] \to (0,1) \times \mathbb{R} \\ t \mapsto (\mu(t),c(t)) \end{split}$$

from $\gamma(0) = (1/2, -2 + \epsilon)$ to $\gamma(1) = (\mu, c_1)$ such that $c_1 \ge c$ and for all $t \in [0, 1]$

$$H_{\mu(t)}\left(L_{1}^{\mu(t)}\right) < c(t) < H_{\mu(t)}\left(L_{2}^{\mu(t)}\right), H_{\mu(t)}\left(L_{3}^{\mu(t)}\right),$$

i. e. a path between the first and the second critical value from the base case to the mass ratio and (possibly higher) energy of the supposed problematic case. Note that $\mathfrak{K}^{b}_{c,\mu} \subset \mathfrak{K}^{b}_{c',\mu}$ for $c \leq c'$, so we get $\mathfrak{K}^{b}_{\mu,c_{1}} \cap W^{-1}_{\mu}(0) \neq \emptyset$. Since $\mathfrak{K}^{b}_{\mu(0),c(0)} \cap W^{-1}_{\mu(0)}(0) = \emptyset$ by Lemma 6 but $\mathfrak{K}^{b}_{\mu(1),c(1)} \cap W^{-1}_{\mu(1)}(0) \neq \emptyset$ by assumption and everything deforms smoothly along the path, there must be some $t_{0} \in (0,1]$ such that $\partial \mathfrak{K}^{b}_{\mu(t_{0}),c(t_{0})} \cap W^{-1}_{\mu(t_{0})}(0) \neq \emptyset$, i. e. the oval of zero velocity



Figure 5: Path γ of mass ratios and energies between critical values

intersects the zero level set of W. Using the symmetry of both $\mathfrak{K}^b_{\mu,c}$ and $W^{-1}_{\mu}(0)$ we can assume this intersection appears outside of $\{q_2 = 0\}$, say $q^0 = (q_1^0, q_2^0) \in \partial \mathfrak{K}^b_{\mu(t_0), c(t_0)} \cap W^{-1}_{\mu(t_0)}(0)$ and $q_2^0 \neq 0$. This implies $V_{q_2,\mu(t_0)}(q^0) = q_2^0 W_{\mu(t_0)}(q^0) = 0$, i. e. $\mathfrak{O}^b_{\mu(t_0),c(0)}$ has an additional vertical tangent away from the q_1 -axis at q^0 .

To continue the first base case below the first critical value one would have to find a smooth path to either $\mu = 0$ (as originally in [2]; left dottet path in figure 5) or to $\mu = 1$. In order to simply be able to fix the energy (second dotted path) one would additionally have to prove the convexity of the first critical energy level. From there on one gets an additional vertical tangent to the oval of zero velocity using a similar argument as above.

In order to show that there can not be an additional vertical tangent of the oval of zero velocity to the obvious ones at the intersection with the q_1 -axis, we write down the smooth dependence of the ovals along the path γ as a smooth 1-parameter family of diffeomorphisms

$$\phi_t \colon S^1 \to \mathfrak{O}^b_{c(t),\mu(t)}$$

So, we can define a smooth 1-parameter family of real valued functions

$$f: [0,1] \times S^1 \to \mathbb{R}$$
$$(t,x) \mapsto f_t(x) := \pi_1 \left(\phi_t(x) \right),$$

where $\pi_1: (q_1, q_2) \mapsto q_1$ is the projection along the second coordinate.

We will show next that f_t is Morse for all $t \in [0, 1]$ in order to apply lemma 5. For that, we have to check that every critical point of f_t is nondegenerate. Critical points here are points where the oval of zero velocity has vertical tangent, i. e. where the partial derivative V_{q_2} vanishes. A critical point is degenerate if it is also a point of inflexion, i. e. if the second derivative $\partial^2 V/\partial q_2^2 =: V_{q_2^2}$ vanishes as well. The proof that there are no vertical points will be split into several cases, depending on the location of the point (as also done in [2]) <u>Case 1</u> ($q_2 = 0$):

If $q_2 = 0$ then $V_{q_2^2}$ reduces to

$$\frac{\mu}{|q_1 - (1 - \mu)|^3} + \frac{1 - \mu}{|q_1 + \mu|^3} - 1 \stackrel{!}{=} 0.$$
(9)

<u>Case 1.1</u> $(-\mu < q_1 < 1 - \mu)$:

So, if q_1 lies between e and s, the denominators in the two fractions of (9) are both strictly smaller than 1, making the left hand side strictly negative and especially non-zero.

<u>Case 1.2</u> $(q_1 > 1 - \mu)$:

On the other hand, if $q_1 > 1 - \mu$ we compute the partial derivative of V by q_1 to be

$$V_{q_1}(q_1,0) = \frac{\mu \left(q_1 - (1-\mu)\right)}{|q_1 - (1-\mu)|^3} + \frac{(1-\mu)\left(q_1 + \mu\right)}{|q_1 + \mu|^3} - q_1$$

= $(q_1 - (1-\mu))\left(\frac{\mu}{|q_1 - (1-\mu)|^3} + \frac{1-\mu}{|q_1 + \mu|^3} - 1\right) + (1-\mu)\left(\frac{1}{|q_1 + \mu|^3} - 1\right)$

on the q_1 -axis. For vertical points of inflexion, equation (9) gives us

$$u'(q_1) = V_{q_1}(q_1, 0) \stackrel{!}{=} (1 - \mu) \left(\frac{1}{|q_1 + \mu|^3} - 1\right) < 0.$$

However, this is exactly the first derivative of the function $u = V|_{\mathbb{R}\setminus\{e,s\}}$ from (6), which was defined earlier and shown to be strictly monotonically increasing for $q_1 > 1 - \mu$ until it reaches the critical point ℓ_2 and only then starts to decline again (see figure 2). By symmetry of the Hill's regions and the discussion in chapter 3.1 about the shape of the Hill's regions there is no part of the bounded Hill's region beyond this critical point since $c < H(L_2)$. So $u'(q_1)$ must be strictly positive and so contradicting the assumption that there is a vertical point of inflexion in this case.

<u>Case 1.3</u> $(q_1 < -\mu)$:

This case works analogously to the latter case, as V_{q_1} becomes

$$\begin{aligned} V_{q_1}(q_1,0) &= \frac{\mu \left(q_1 - (1-\mu)\right)}{|q_1 - (1-\mu)|^3} + \frac{(1-\mu) \left(q_1 + \mu\right)}{|q_1 + \mu|^3} - q_1 \\ &= (q_1 + \mu) \left(\frac{\mu}{|q_1 - (1-\mu)|^3} + \frac{1-\mu}{|q_1 + \mu|^3} - 1\right) - \mu \left(\frac{1}{|q_1 - (1-\mu)|^3} - 1\right) \end{aligned}$$

and for vertical points of inflexion we would have

$$u'(q_1) = V_{q_1}(q_1, 0) \stackrel{!}{=} -\mu\left(\frac{1}{|q_1 - (1 - \mu)|^3} - 1\right) > 0.$$

As in case 1.2 this implies that the restricted potential increases, which is only the case up until ℓ_3 . By $c < H(L_3)$ these points are not reached by any of the bounded Hill's regions \mathfrak{K}^b_c .

From these first cases we can conclude, that there can be no vertical point of inflexion of the oval of zero velocity along the q_1 -axis inside of \Re^b_c .

Next, we show that a vertical point of inflexion can also not lie at any other point inside the bounded Hill's region.

<u>Case 2</u> $(q_2 \neq 0)$:

Here, a vertical point of inflexion implies, that by the quotient rule

$$\frac{\partial}{\partial q_2} \left(\frac{V_{q_2}}{q_2} \right) = \frac{q_2 V_{q_2^2} - V_{q_2}}{q_2^2} \stackrel{!}{=} 0.$$

But V_{q_2}/q_2 is just the coefficient of q_2 in (8) and after differentiating there remains

$$\frac{\partial}{\partial q_2} \left(\frac{V_{q_2}}{q_2} \right) = \frac{\partial}{\partial q_2} \left(\frac{\mu}{|q-e|^3} + \frac{1-\mu}{|q-s|^3} - 1 \right) = -3q_2 \left(\frac{\mu}{|q-e|^5} + \frac{(1-\mu)}{|q-s|^5} \right) \neq 0.$$

This final contradiction concludes the statement that there can not be a vertical point of inflexion inside the bounded Hill's region \mathfrak{K}^b_c and thus all critical points of f_t are non-degenerate, i. e. Morse for all t. Therefore, we can apply lemma 5 and we obtain, that the number of critical values of f_t is constant for all $t \in [0, 1]$. Explicitly the number of critical points is exactly two, since $\mathfrak{O}^b_{-2+\epsilon,1/2}$ is a symmetric and connected 1-submanifold of \mathbb{R} and critical points only lie on the q_1 -axis. In particular there can not be a vertical tangent of $\mathfrak{O}^b_{\mu(t_0),c(t_0)}$ at $q^0 \notin \{q_2 = 0\}$.

Using lemma 7, one can now conclude that every periodic orbit inside the bounded Hill's region



(a) Periodic orbits without quadratures

Figure 6: Hill's lunar problem

has at least two syzygies: Let x be a non-trivial orbit in Σ_c^e and $T_x > 0$ its minimal period time the only trivial periodic orbit is stationary at ℓ_1 , which lies on the q_1 -axis, so the statement also holds here. Then by integrating the second equation (3) of the equations of motion, we get

$$\int_{0}^{T_{x}} \ddot{x}_{2}(t) dt = -2 \int_{0}^{T_{x}} \dot{x}_{1}(t) dt - \int_{0}^{T_{x}} V_{q_{2}}(x(t)) dt$$

i. e. $0 = \int_{0}^{T_{x}} V_{q_{2}}(x(t)) dt.$

Since the integral over a smooth function can only be zero if the function itself passes zero we need $V_{q_2}(x(t_0)) = 0$ for some $0 < t_0 < T_x$. By lemma 7 this is only the case if $x_2(t_0) = 0$, i. e. if x intersects the q_1 -axis. We can integrate the same equation again from t_0 to $t_0 + T_x$ to get a second intersection during one period. Note, that these intersections must be transverse, as the sign of V_{q_2} must change at these syzygies.

6 Another (easier) application of the same method: Hill's lunar problem

Hill's lunar problem is a limit case of the PCR3BP by letting the sun get infinitely heavy and at same time infinitely far away (see [6]). The Hamiltonian is given by

$$H(q,p) = \frac{1}{2}|p|^2 + q_2p_1 - q_1p_2 - \frac{1}{|q|} - q_1^2 + \frac{1}{2}q_2^2$$
$$= \frac{1}{2}\left((p_1 + q_2)^2 + (p_2 - q_1)^2\right) - \frac{1}{|q|} - \frac{3}{2}q_1^2$$

and the corresponding second order differential equations are

$$\ddot{q}_1 = 2\dot{q}_2 - \frac{q_1}{|q|^3} + 3q_1 \tag{10}$$

$$\ddot{q}_2 = -2\dot{q}_1 - \frac{q_2}{|q|^3}.\tag{11}$$



Figure 7: Set of zeros of V_{q_1}

We can now do the same integration over (11) as in the final step in chapter 5 to get

$$\begin{split} \int_0^T \ddot{x}_2(t) \, \mathrm{d}t &= -2 \int_0^T \dot{x}_1(t) \, \mathrm{d}t - \int_0^T \frac{x_2(t)}{|x(t)|^3} \, \mathrm{d}t \\ 0 &= \int_0^T \frac{x_2(t)}{|x(t)|^3} \, \mathrm{d}t \end{split}$$

which only happens if the non-trivial periodic orbit x passes the q_1 -axis at some point t_0 . By integrating from t_0 to $t_0 + T$ there is a second syzygy in each periodic orbit of Hill's lunar problem and both syzygies are again transverse. There are two critical points at $(\pm 3^{-1/3}, 0)$ so the statement also holds for these stationary orbits.

By integrating (10) we can also prove the existence of quadratures. However since the two critical points generate families of periodic orbits away from the q_2 -axis not all closed orbits have quadratures (figure 6a). These are families a and c as classified in [7]. One can show—as done in [11] that the bounded Hill's region, i. e. the bounded component of possible positions for energies below the common critical value $-3^{4/3}/2$, is contained within the ball of radius $3^{-1/3}$ around the origin. So all periodic orbits of Hill's lunar problem with energy below the first critical value have quadratures. Again one can see the correspondence that additional—in this case horizontal—tangents of the ovals of zero velocity prevent an extension of the statement to higher energies (figure 6b).

7 Remarks

Since in Hill's lunar problem the horizontal tangents originate in the critical points which lie away from the q_2 -axis there are direct counterexamples to show the statement can not be extended to a higher energy. In the PCR3BP however the additional vertical tangents, preventing the proof to go through for higher energies, originate in critical points lying on the q_1 -axis themselves. So there are no direct and obvious obstructions for all periodic orbits having syzygies for energies $c < H(L_4) = H(L_5)$.

Similarly for the unbounded Hill's region, the proof as shown in this paper does not work as there can be vertical points of inflexion and so the constructed 1-parameter family of maps on S^1 are not Morse. Indeed, vertical points of inflexion generate additional vertical tangents below the second critical value, which disables our argument of integration. However, for small enough energies obviously roots of V_{q_2} only lie along the q_1 -axis also in the unbounded part and we get syzygies in the same way as before.

However for energies $c > H(L_5)$ there is no hope to prove the existence of syzygies for all periodic orbits since the Lyapunov orbits around the triangular Lagrange points L_4 and L_5 provide direct counterexamples.

Opposed to Hill's lunar problem, in the PCR3BP quadratures with respect to one of the primaries can not be predicted in this manner, since in general neither $\{q_1 = -\mu\}$ nor $\{q_1 = 1-\mu\}$ makes the partial derivative V_{q_1} vanish (although the set of roots comes arbitrarily close for small

enough radii; see figure 7 and [2]).

Further interesting questions which can not be answered by this method is whether periodic orbits in the bounded Hill's region below the first critical value have a non-trivial winding number with respect to the respective primary or even if they have syzygies on both sides (i. e. if there always exist both solar and lunar eclipse). Equally unknown is what we can say about sequences of syzygy types for periodic or even arbitrary orbits in the bounded Hill's region below the second critical value.

References

- R. Abraham and J. E. Marsden. Foundation of Mechanics. Addison-Wesley Publishing Company, Inc., 2nd edition, 1987.
- [2] G. D. Birkhoff. The restricted problem of three bodies. In *Rendiconti del Circolo Matematico di Palermo*, volume 39, pages 265 334, 1915.
- [3] H. R. Dullin and R. Montgomery. Syzygies in the two center problem. Nonlinearity, 29(4):1212, 2016.
- [4] U. Frauenfelder and O. van Koert. The restricted three body problem and holomorphic curves. Pathways in Mathematics. Birkhäuser Basel, 2018.
- [5] Y. Hagihara. Dynamical principles and transformation theory, volume 1 of Celestial Mechanics. MIT Press, 1970.
- [6] G. W. Hill. Researches in the lunar theory. American Journal of Mathematics, 1(1):5–26, 1878.
- [7] M. Hénon. Numerical Exploration of the Restricted Problem. V. Astronomy and Astrophysics, 1:223 – 238, 1969.
- [8] R. Moeckel and R. Montgomery. Realizing all reduced syzygy sequences in the planar threebody problem. *Nonlinearity*, 28(6), 2015.
- [9] R. Montgomery. Infinitely many syzygies. Archive for Rational Mechanics and Analysis, 164(4):311–340, 2002.
- [10] R. Montgomery. The zero angular momentum, three-body problem: All but one solution has syzygies. *Ergodic Theory and Dynamical Systems*, 27(6):1933–1946, 2007.
- [11] R. Nicholls. Syzygies of periodic orbits in the restricted 3-body problem. Master's thesis, University of Augsburg, 2017.
- [12] V. Szebehely. Theory of orbits, the restricted problem of three bodies. Academic Press, 1967.
- [13] D. Wilczak and P. Zgliczyński. Heteroclinic connections between periodic orbits in planar restricted circular three-body problem – a computer assisted proof. *Communications in Mathematical Physics*, 234(1):37–75, 2003.
- [14] D. Wilczak and P. Zgliczyński. Heteroclinic connections between periodic orbits in planar restricted circular three body problem. part ii. *Communications in Mathematical Physics*, 259(3):561–576, 2005.