A THEIL-LIKE CLASS OF INEQUALITY MEASURES, ITS ASYMPTOTIC NORMALITY THEORY AND APPLICATIONS

 † PAPE DJIBY MERGANE, †† TCHILABALO ABOZOU KPANZOU, ††† DIAM BA, AND †††† GANE SAMB LO

ABSTRACT. In this paper, we consider a coherent theory about the asymptotic representations for a family of inequality indices called Theil-Like Inequality Measures (TLIM), within a Gaussian field. The theory uses the functional empirical process approach. We provide the finite-distribution and uniform asymptotic normality of the elements of the TLIM class in a unified approach rather than in a case by case one. The results are then applied to some UE-MOA countries databases.

[†] Pape Djiby Mergane LERSTAD, Gaston Berger University, Saint-Louis, Sénégal. Email : mergane@gmail.com.

^{††} Tchilabalo Abozou Kpanzou (corresponding author). Kara University, Kara, Togo. Email : kpanzout@gmail.com

^{†††} Diam Ba LERSTAD, Gaston Berger University, Saint-Louis, Sénégal. Email : diamba79@gmail.com.

^{††††} Gane Samb Lo.

LERSTAD, Gaston Berger University, Saint-Louis, Sénégal (main affiliation). LSTA, Pierre and Marie Curie University, Paris VI, France. AUST - African University of Sciences and Technology, Abuja, Nigeria gane-samb.lo@edu.ugb.sn, gslo@aust.edu.ng, ganesamblo@ganesamblo.net Permanent address : 1178 Evanston Dr NW T3P 0J9,Calgary, Alberta, Canada.

keywords and phrases. Inequality measures; Asymptotic behaviour; Asymptotic representations; functional empirical proces.

AMS 2010 Mathematics Subject Classification : 62G05; 62G20; 62G07; 91B82; 62P20.

2 PAPE DJIBY MERGANE, †† TCHILABALO ABOZOU KPANZOU, ††† DIAM BA, AND †††† GANE SAMB LO

Résumé. (French) Dans cet article, nous présentons une théorie cohérente de représentations asymptotiques d'une famille de mesures d'ingégalité dénommée TLIM dans un champ gaussien précis. Notre méthode est fondée sur le processus empirique fonctionnel. Nous tirons de la représentation asymptotique les limites en distribution des estimateurs plug-in des membres de la famille en dimension finie. Les résultats sont ensuite appliqués à des données issues des pays de l'UEMOA.

1. INTRODUCTION

In this paper, we deal with a modern weak theory for some large class of inequality indices that, further, will allow to handle easy comparison studies with different kinds of statistics.

According to earlier economists, inequality indices are functional relations between the income and the economic welfare (see Dalton (1920)). This explains, among others, the wide variety of such indices in the literature (See, e.g., Cowell (1980a,b, 2000)).

Such statistics, of course, have been widely studied with respect to a great variety of interests, including statistical characterizations and asymptotic properties (See Davidson and Duclos (2000), Barrett and Donald (2009), for recent studies).

Recently, Greselin et al. (2009) provided a mathematical investigation of these indices in a modern setting including Vervaat processes, Lstatistics and empirical processes.

Having in mind the necessity of comparing inequality measures with different kind of statistics such as growth statistics, we aim at providing a coherent asymptotic weak theory for some class of inequality measures. Indeed we propose the functional empirical process setting (see Van der Vaart and Wellner (1996)) which provide natural Gaussian field in which many statistics used in Economics may be represented in.

Our best achievement consists of the asymptotic representations for the elements of our class of inequality measures, in terms of the above mentioned Gaussian field. The results are illustrated in data driven applications, on Senegalese data for instance. The class on which we focus here is a functional family of inequality measures which gathers various ones around the central Theil measure. This class named after the Theil-Like Inequality Measure (TLIM) will be the central point of our study. It includes the Generalized Entropy Measure, the Mean Logarithmic Deviation (Cowell (2003); Theil (1967); Cowell (1980a)), the different inequality measures of Atkinson (1970), Champernowne and Cowell (1998), Kolm (1976a), and the divergence of Renyi (1961).

This means that, here, we will not discuss other inequality statistics such as the Gini, the Generalized Gini, the S-Gini, the E-Gini (See Barrett and Donald (2009)). Those statistics and similar ones will be treated in separate papers.

Now we are going to introduce our family. For that, let X denote the income (or expense) random variable related to a given population. We assume that X and its independent observations are defined on the probability space $(\Omega, \mathcal{A}, \mathbb{P})$ and take their values in on some interval $\mathcal{V}_X \subset \mathbb{R}^*_+$ and have common cumulative distribution function (cdf), $F(x), x \in \mathcal{V}_X$. In this paper, we only use Lebesgue-Stieljes integrals and for any measurable function $\ell : \mathbb{R} \to \mathbb{R}$, we have, whenever it makes sense,

$$\mathbb{E}(\ell(X)) = \int \ell(x) \ dF(x) \equiv \int_{\mathcal{V}_X} \ell \ d\mathbb{P}_X,$$

where $\mathbb{P}_X = \mathbb{P}X^{-1}$ is the measure image of \mathbb{P} by X, but is also Lebesgue-Stieljes probability measure characterized by: $\mathbb{P}_X([a,b]) = F(b) - F(a)$ for any $-\infty \leq a \leq b \leq +\infty$.

Now, consider a sample of $n \geq 1$ individuals or households of that population and observe their income X_1, X_2, \dots, X_n . We define the following family of inequality indices, indexed by $\phi = (\tau, h, h_1, h_2) \in \mathcal{P}_0$ as follows

(1.1)
$$T_n(\phi, X) = \tau \left(\frac{1}{n} \sum_{j=1}^n \left(\frac{h(X_j)}{h_1(\mu_n)} - h_2(\mu_n) \right) \right), \ h_1(\mu_n) \neq 0,$$

where $\mu_n = \frac{1}{n} \sum_{j=1}^n X_j$ is the empirical mean while h(x), $h_1(y)$, $h_2(z)$, and $\tau(t)$ are real and measurable functions of $x, y, z \in \mathcal{V}_X$ and $t \in \mathbb{R}$. The exact form of \mathcal{P}_0 is not important here, in opposite to the conditions on the functions τ , h, h_1 and h_2 under which the results are valid. In a future paper on the uniform limits in ϕ , that class will be crucial.

We will see below that T_n under specific hypotheses on τ , h, h_1 , h_2 and μ_n , converges to the exact inequality measure

(1.2)
$$T(\phi, X) = \tau \left(\frac{1}{h_1(\mu)} \int_{\mathcal{V}_X} h(x) dF(x) - h_2(\mu)\right), \ h_1(\mu) \neq 0,$$

where $\mu = \mathbb{E}(X)$ is the mathematical expectation of X that we suppose finite here. We will come back later on the function classes \mathcal{F}_1 , \mathcal{F}_2 , \mathcal{F}_3 and \mathcal{F}_4 in which h, τ , h_1 and h_2 are supposed to lie.

Each measure of this Theil-like family has its own particular properties, that are derived from the combination of different concepts. One may mention the concept of welfare criteria (Atkinson (1970), Sen (1973)), that of the analogy with analysis of risks (Harsanyi (1953), Harsanyi (1955), Rothschild and Stiglitz (1973)), that of the complaints approach (Temkin (1993)) etc. The Theil inequality itself finds all its interest in the information-theoretic idea following that of main components (Kullback 1959). It is based on the three following axioms: Zero-valuation of certainty, Diminishing-valuation of probability, Additivity of independent events. A deep review of such of individual properties for a number inequality measures can be found in Cowell (Cowell (1980a,b, 2000)) for instance.

It is worth mentioning that the TLIM presented here, is rather a mathematical form gathering a number of different measures.

The rest of our paper is organized as follows. In Section 2, we describe the TLIM family and show how the particular indices are derived from it. In Section 3, we briefly recall the functional empirical processes setting. In section 4, we deal with the asymptotic theory of the TLIM, state and describe our main results and demonstrate them. Section 5 is devoted to datadriven applications. We finish by a conclusion in Section 7.

2. Description of the TLIM

This inequality measures mentioned above are derived from (1.1) with the particular values of the mesurable functions τ , h, h_1 and h_2 as described below for all s > 0.

2.1. Generalized Entropy.

$$GE_{n,\alpha}(X) = \frac{1}{n\alpha(\alpha-1)} \sum_{j=1}^{n} \left(\left(\frac{X_j}{\mu_n} \right)^{\alpha} - 1 \right);$$

$$\alpha \neq 0, \ \alpha \neq 1, \ \tau(s) = \frac{s-1}{\alpha (\alpha - 1)}, \ h(s) = h_1(s) = s^{\alpha}, \ h_2(s) \equiv 0.$$

2.2. Theil's measure.

$$Th_n(X) = \frac{1}{n} \sum_{j=1}^n \frac{X_j}{\mu_n} \log \frac{X_j}{\mu_n};$$

$$\tau(s) = s, \ h(s) = s \ \log(s), \ h_1(s) = s, \ h_2(s) = \log(s).$$

2.3. Mean Logarithmic Deviation.

$$MLD_n(X) = \frac{1}{n} \sum_{j=1}^n \log\left(\frac{X_j}{\mu_n}\right)^{-1};$$

$$\tau(s) = s, \ h(s) = h_2(s) = \log(s^{-1}), \ h_1(s) \equiv 1.$$

2.4. Atkinson's measure.

$$Atk_{n,\alpha}(X) = 1 - \frac{1}{\mu_n} \left(\frac{1}{n} \sum_{j=1}^n X_j^{\alpha}\right)^{1/\alpha};$$

 $\alpha < 1$ and $\alpha \neq 0$, $\tau(s) = 1 - s^{1/\alpha}$, $h(s) = h_1(s) = s^{\alpha}$, $h_2(s) \equiv 0$.

2.5. Champernowne's measure.

$$Ch_n(X) = 1 - \frac{1}{\exp\left(-\frac{1}{n}\sum_{j=1}^n \log\frac{X_j}{\mu_n}\right)};$$

$$\tau(s) = 1 - \exp(s), \ h(s) = h_2(s) = \log(s), \ h_1(s) \equiv 1.$$

2.6. Kolm's measure.

$$Ko_{n,\alpha}(X) = \log\left(\frac{1}{n}\sum_{j=1}^{n}\exp\left(-\alpha(X_j - \mu_n)\right)\right)^{1/\alpha};$$

$$\alpha > 0, \ \tau(s) = \frac{1}{\alpha} \log(s), \ h(s) = h_1(s) = \exp(-\alpha s), \ h_2(s) \equiv 0.$$

2.7. Divergence of Renyi.

$$DR_{n,\alpha}(X) = \frac{1}{\alpha - 1} \log \left(\frac{1}{n} \sum_{j=1}^{n} \left(\frac{X_j}{\mu_n} \right)^{\alpha} \right);$$

$$\alpha \in \mathbb{R}_+ \setminus \{1\}, \ \tau(s) = \frac{1}{\alpha - 1} \log(s), \ h(s) = h_1(s) = s^{\alpha}, \ h_2(s) \equiv 0.$$

3. The functional empirical process

Let Z_1, Z_2, \ldots, Z_n be a sequence of independent and identically distributed random elements defined on the probability space $(\Omega, \mathcal{A}, \mathbb{P})$, with values in some metric space (S, d). Given a collection \mathcal{F} of measurable functions $f: S \to \mathbb{R}$ satisfying

$$\sup_{f\in\mathcal{F}}|f(z)-\mathbb{P}(f)|<\infty, \text{ for every } z,$$

where $\mathbb{P}(f) = \mathbb{E}(f(Z))$ is the mathematical expectation of f(Z), the functional empirical process (*FEP*) based on the $(Z_j)_{j=1,\dots,n}$ and indexed by \mathcal{F} is defined by:

$$\forall f \in \mathcal{F}, \mathbb{G}_n(f) = \frac{1}{\sqrt{n}} \sum_{j=1}^n \left(f(Z_j) - \mathbb{P}(f) \right).$$

This process is widely studied in Van der Vaart and Wellner (1996) for instance. It is readily derived from the real Law of Larges Numbers (*LLN*) and the real Central Limit Theorem (*CLT*) that $\mathbb{P}_n(f) = \frac{1}{n} \sum_{j=1}^n f(Z_j) \to \mathbb{P}(f)$ a.s. and that $\mathbb{G}_n(f) \to \mathcal{N}(0, \sigma_f^2)$, where

(3.1)
$$\sigma_f^2 = \mathbb{P}\left(\left(f - \mathbb{P}\left(f\right)\right)^2\right) < \infty,$$

whenever $\mathbb{E}(f(Z)^2) < \infty$.

When using the FEP, we are often interested in uniform LLN's and weak limits of the FEP considered as stochastic processes. This gives the so important results on Glivenko-Cantelli classes and Donsker ones. Let us define them here (for more details see Van der Vaart and Wellner (1996)).

Since we may deal with non measurable sequences of random elements, we generally use the outer almost sure convergence defined as follows: a sequence U_n converges outer almost surely to zero, denoted by $U_n \rightarrow 0 as^*$, whenever there is a measurable sequence of measurable random variables V_n such that

- (1) $\forall n, |U_n| \leq V_n,$
- (2) $V_n \longrightarrow 0 \ a.s.$

The weak convergence generally holds in $\ell^{\infty}(\mathcal{F})$, the space of all bounded real functions defined on \mathcal{F} , equipped with the supremum norm $||x||_{\mathcal{F}} = \sup_{f \in \mathcal{F}} |x(f)|$.

Definition 1. $\mathcal{F} \subset L_1(\mathbb{P})$ is called a Glivenko-Cantelli class for \mathbb{P} , if

$$\lim_{n \to \infty} \left\| \frac{1}{n} \sum_{j=1}^n \left(f(Z_j) - \mathbb{P}(f) \right) \right\|_{\mathcal{F}} = 0 \ a.s.^*$$

Definition 2. $\mathcal{F} \subset L_2(\mathbb{P})$ is called a Donsker class for \mathbb{P} , or \mathbb{P} -Donsker class if $\{\mathbb{G}_n(f); f \in \mathcal{F}\}$ converges in $l^{\infty}(\mathcal{F})$ to a centered Gaussian process $\{\mathbb{G}(f); f \in \mathcal{F}\}$ with covariance function

$$\Gamma(f,g) = \int_{\mathbb{R}} \left(f(z) - \mathbb{P}(f) \right) \left(g(z) - \mathbb{P}(g) \right) \, d\mathbb{P}_{Z}(z) \, ; \, \forall f,g \in \mathcal{F}.$$

Remark 1. When $S = \mathbb{R}$ and $\mathcal{F} = \{f_t = \mathbf{1}_{(-\infty,t]}, t \in \mathbb{R}\}, \mathbb{G}_n \text{ is called real empirical process and is denoted by } \alpha_n$.

In this paper, we only use finite-dimensional forms of the FEP, that is $(\mathbb{G}_n(f_i), i = 1, ..., k)$. And then, any family $\{f_i, i = 1, ..., k\}$ of measurable functions satisfying (3.1), is a Glivenko-Cantelli and a Donsker class, and hence

$$\left(\mathbb{G}_{n}\left(f_{i}\right), i=1,\ldots,k\right) \stackrel{a}{\rightarrow} \left(\mathbb{G}\left(f_{1}\right),\mathbb{G}\left(f_{2}\right),\ldots,\mathbb{G}\left(f_{k}\right)\right),$$

where \mathbb{G} is the Gaussian process, defined in Definition 2. We will make use of the linearity property of both \mathbb{G}_n and \mathbb{G} . Let f_1, \ldots, f_k be measurable functions satisfying (3.1) and $a_i \in \mathbb{R}, i = 1, \ldots, k$, then

$$\sum_{j=1}^{k} a_{j} \mathbb{G}_{n} (f_{j}) = \mathbb{G}_{n} \left(\sum_{j=1}^{k} a_{j} f_{j} \right) \stackrel{d}{\to} \mathbb{G} \left(\sum_{j=1}^{k} a_{j} f_{j} \right).$$

The materials defined here, when used in a smart way, lead to a simple handling of the problem which is addressed here.

4. Our results

Let us introduce some notation.

$$B_{h,n} = \frac{1}{n} \sum_{j=1}^{n} h(X_j), \ B_h = \int_{\mathcal{V}_X} h(x) \, dF(x);$$
$$K_{\phi} = \tau' \left(\frac{B_h}{h_1(\mu)} - h_2(\mu)\right) \neq 0;$$

for all $x \in \mathcal{V}_X$, we define the following function

$$F_{\phi}(x) = K_{\phi} \left(\frac{1}{h_1(\mu)} h(x) - \left(\frac{B_h h_1'(\mu)}{h_1^2(\mu)} + h_2'(\mu) \right) I_d(x) \right)$$

with $I_d(x) = x$, and τ' is the derivative of the function τ .

The following general condition will be assumed in all the paper:

(C) h_1 is not null in a neighborhood of μ . Here are our main results.

4.1. Pointwise asymptotic laws. Consider the following hypotheses based on the functions h, τ , h_1 , h_2 . The A1.x series concern the almost-sure limits and the A2.x the asymptotic normality.

(A1.1) $\mathbb{E}h(X) < \infty$; (A1.2) τ is a continuous function on \mathcal{V}_X ; (A1.3) for $i \in \{1, 2\}$, $h_i(\mu) < \infty$ and h_i is continuous on \mathcal{V}_X .

(A2.1) $\mathbb{E}h^2(X) < \infty$, $\mathbb{E}(X h(X)) < \infty$; (A2.2) τ is continuously differentiable such that $\tau' \neq 0$; (A2.3) $\forall i \in \{1, 2\}, h_i(\mu) < \infty, h_i$ is continuously differentiable at μ .

We have :

Theorem 1. Suppose that the conditions (C), (A1.1), (A1.2) and (A1.3) are satisfied, then T_n converges almost surely to T.

Theorem 2. Suppose that the conditions (C), (A2.1), (A2.2) and (A2.3) are satisfied, and K_{ϕ} is finite. Then

(a) we have the following asymptotic representation in the empirical functional process

$$\sqrt{n}\left(\tau(I_n) - \tau(I)\right) = \mathbb{G}_n\left(F_\phi\right) + o_{\mathbb{P}}(1), \ as \ n \to +\infty,$$

where

$$F_{\phi} = \tau' \left(\frac{\mathcal{E}h(Y)}{h_1(\mu)} - h_2(\mu) \right) \left(\frac{1}{h_1(\mu)} h - \left(\frac{\mathcal{E}h(Y) h_1'(\mu)}{h_1^2(\mu)} + h_2'(\mu) \right) I_d \right)$$

(b) and we have the convergence in distribution, as n tends to infinity, of $\sqrt{n}(T_n(\phi, X) - T(\phi, X))$ to centered normal Gaussian law:

$$\sqrt{n}(T_n(\phi, X) - T(\phi, X)) \rightsquigarrow \mathcal{N}(0, \sigma_{\phi}^2),$$

where

$$\sigma_{\phi}^{2} = \int \left(F_{\phi}(x) - \int F_{\phi}(x) \ d\mathbb{P}_{X}(x) \right)^{2} \ d\mathbb{P}_{X}(x)$$

$$= a_{\phi}^{2} \mathbb{E} \left(h(X) - \mathbb{E}h(X) \right)^{2} + b_{\phi}^{2} \mathbb{E} \left(X - \mu \right)^{2}$$

$$- 2a_{\phi}b_{\phi} \mathbb{E} \left(h(X) - \mathbb{E}h(X) \right) \left(X - \mu \right),$$

with

$$a_{\phi} = \frac{K_{\phi}}{h_1(\mu)} \text{ and } b_{\phi} = K_{\phi} \left(\frac{B_h h_1'(\mu)}{h_1^2(\mu)} + h_2'(\mu) \right).$$

Remark. The main result is the one given in Point (a). From it, Point (b) is deduced in a straightforward way.

The results above cover all the TLIM class. They should be particularized for the practitioner who would pick one of the elements of that class for analyzing data. Here are then the details for each case. $\dagger \Phi \text{APE}$ DJIBY MERGANE, †† TCHILABALO ABOZOU KPANZOU, ††† DIAM BA, AND †††† GANE SAMB LO

4.2. Particular cases for pointwise results.

a. The Theil's measure

The empirical form of Theil measure is defined as follows

$$Th_n = \frac{1}{\mu_n} \frac{1}{n} \sum_{j=1}^n X_j \log X_j - \log \mu_n,$$

 $\forall s > 0, \ \tau(s) = s, \ h(s) = s \ \log(s), \ h_1(s) = s, \ h_2(s) = \log(s),$

Denote by

$$Th = \frac{1}{\mu} \int_{\mathcal{V}_X} x \log x \, dF(x) - \log \mu$$

the continuous form of the Theil measure.

All these functions are continuous on \mathcal{V}_X , then the assumptions defined above become for the *a.s.* requires that $\mathbb{E}X \log X$ is finite and $0 < \mu < \infty$. As for the asymptotic normality, we need that

$$\mathbb{E} |X|^2$$
, $\mathbb{E} |X \log X|^2 \mathbb{E} |X^2 \log X|^2$ are finite.
And we have $K_{\phi} = 1$, $B_h = \mathbb{E} (X \log X)$. We conclude that

$$\sqrt{n} \left(Th_n - Th \right) \rightsquigarrow \mathcal{N}(0, \sigma_{Theil}^2)$$

with

$$\sigma_{Theil}^{2} = \frac{\mathbb{E}\left(X\log X\right)^{2}}{\mu^{2}} + \frac{\mathbb{E}X^{2}}{\mu^{2}} \left(\frac{B_{h}}{\mu} + 1\right)^{2} - \frac{2\mathbb{E}\left(X^{2}\log X\right)}{\mu^{2}} \left(\frac{B_{h}}{\mu} + 1\right) - 1.$$

b. The Mean Logarithmic Deviation

Let

$$MLD_n = \frac{1}{n} \sum_{j=1}^n \log X^{-1} - \log \mu_n^{-1}$$

be the empirical form of the Mean Logarithmic Deviation. Its theoretical form is given as follows

$$MLD = \int_{\mathcal{V}_X} \log x^{-1} \, dF(x) - \log \mu^{-1}.$$

These specific functions are given by:

$$\forall s > 0, \ \tau(s) = s, \ h(s) = h_2(s) = \log s^{-1}, \ h_1(s) \equiv 1.$$

The consistency requires that $\mathbb{E} \log X < \infty$ and that $0 < \mu < \infty$ while the normality is got when

$$\mathbb{E}|X|^2$$
, $\mathbb{E}|\log X|^2$ and $\mathbb{E}|X\log X|$ are finite.

In that case, we find easily that $K_{\phi} = 1, B_h = \mathbb{E} \log X^{-1}$ and

$$\sqrt{n} \left(MLD_n - MLD \right) \rightsquigarrow \mathcal{N}(0, \sigma_{MLD}^2)$$

where

$$\sigma_{MLD}^2 = \frac{\mathbb{E}(X^2)}{\mu^2} + \mathbb{E}\left(\log^2 X\right) - \frac{2}{\mu}\mathbb{E}\left(X\log X\right) - \left(B_h + 1\right)^2.$$

c. The Champernowne's measure

In this case, the specific functions are given by:

$$\tau(s) = 1 - \exp(s), \ h(s) = h_2(s) = \log(s), \ h_1(s) \equiv 1.$$

And, the various forms are:

$$Ch_n = 1 - \exp\left(\frac{1}{n}\sum_{j=1}^n \log\frac{X_j}{\mu_n}\right);$$
$$Ch = 1 - \exp\left(\int \mathcal{V}_X \log\frac{x}{\mu} dF(x)\right).$$

We find that $Ch_n = \tau (-MLD_n)$ and $Ch = \tau (-MLD)$, where MLD is the Mean Logarithmic Deviation. As τ is continuous on \mathcal{V}_X , we consider the same hypotheses as in the case of Mean Logarithmic Deviation.

The function τ is continuously differentiable, we put $B_h = \mathbb{E} \log X^{-1}$ and $K_{\phi} = \frac{\exp(-B_h)}{\mu}$, then we have

$$\sqrt{n} \left(Ch_n - Ch \right) \rightsquigarrow \mathcal{N}(0, \sigma_{Ch}^2)$$

with

$$\sigma_{Ch}^2 = K_\phi^2 \, \sigma_{MLD}^2.$$

d. Cases of the Generalized Entropy $(\alpha \neq 0, \alpha \neq 1)$; the Atkinson's measure $(\alpha < 1, \alpha \neq 0)$; the Divergence of Renyi $(\alpha > 0, \alpha \neq 1)$.

We may gather these indices into one subclass by giving different values to the function τ and to the parameter α , with this common expression

$$\forall s > 0, h(s) = h_1(s) = s^{\alpha} \text{ and } h_2 \equiv 0,$$

and then give a general description of the results. For that, Let $I_{n,\alpha} =$ $\mathbb{P}_n(h)/h_1(\mu_n)$ and $I_\alpha = \int_{\mathcal{V}_X} \frac{h(x)}{h_1(\mu)} dF(x).$

We require for consistency that $\mathbb{E}X^{\alpha} < \infty$ and that $\mu \neq 0$ and, for asymptotic normality that

 $\mathbb{E} |X|^{2\alpha} < \infty, \ \mathbb{E} |X|^2 < \infty \text{ and } \mathbb{E} |X|^{\alpha+1} < \infty.$ Further, let $B_h = \mathbb{E} X^{\alpha}$ and $K_{\phi} = \tau'(I_{\alpha})$. Then we get

$$\sqrt{n}\left(I_{n,\alpha} - I_{\alpha}\right) = \mathbb{G}_n\left(\frac{h}{\mu^{\alpha}} - \frac{\alpha \mathbb{E}X^{\alpha}}{\mu^{\alpha} + 1}I_d\right) + o_{\mathbb{P}}^*(1),$$

which tends towards a centered Gaussian process with variance

$$\sigma_{I_{\alpha}}^{2} = \frac{1}{\mu^{2\alpha}} \left(\mathbb{E}X^{2\alpha} + \frac{(\alpha B_{h})^{2}}{\mu^{2}} \mathbb{E}X^{2} - \frac{2\alpha B_{h}}{\mu} \mathbb{E}X^{\alpha+1} \right) - \frac{B_{h}^{2}}{\mu^{2\alpha}} \left(1 - \alpha\right)^{2}.$$

Now, we may return to the individual cases.

d.1. Generalized Entropy

We find $K_{\phi} = 1/(\alpha(\alpha - 1))$, from there, we get the variance

$$\sigma_{GE_{\alpha}}^2 = K_{\phi}^2 \sigma_{I_{\alpha}}^2$$
, where $\sigma_{I_{\alpha}}^2$ is given in Equation (4.1).

d.2. Atkinson's measure Put $K_{\phi} = (\mathbb{E}X^{\alpha})^{(1/\alpha-1)}/\alpha$. We similarly get that

$$\sigma_{Atk_{\alpha}}^{2} = \frac{1}{\alpha^{2}} \left(\mathbb{E} X^{\alpha} \right)^{\left(\frac{1-\alpha}{\alpha}\right)^{2}} \sigma_{I_{\alpha}}^{2}.$$

d.3. Divergence of Renyi

By taking $K_{\phi} = ((\alpha - 1)\mathbb{E}X^{\alpha})^{-1}$, we obtain by the same way, that

$$\sigma_{DR_{\alpha}}^{2} = \frac{\sigma_{I_{\alpha}}^{2}}{\left(\left(\alpha - 1\right)\mathbb{E}X^{\alpha}\right)^{2}}$$

where $\sigma_{I_{\alpha}}^2$ is given in (4.1).

e. Case of the Kolm's measure

This index is defined for $\alpha > 0$, and its specific functions are:

$$\tau(s) = \frac{1}{\alpha} \log(s), \ h(s) = h_1(s) = \exp(-\alpha s), \ h_2(s) \equiv 0, \ \forall s > 0.$$

Its empirical form is given by

$$Ko_{n,\alpha} = \log\left(\frac{1}{n}\sum_{j=1}^{n}\exp\left(-\alpha(X_j - \mu_n)\right)\right)^{1/\alpha};$$

and its theoretical form is defined as follows

$$Ko_{\alpha} = \log\left(\int_{\mathcal{V}_X} \left(\frac{e^{-x}}{e^{-\mu}}\right)^{\alpha} dF(x)\right)^{1/\alpha}.$$

We need for consistency that $\mu < \infty$ and that $\mathbb{E} \exp(-\alpha X) < \infty$ and, for asymptotic normality that

$$\mathbb{E}(|X|^2)$$
, $\mathbb{E}(|e^{-\alpha X}|)$, $\mathbb{E}(|e^{-2\alpha X}|)$ and $\mathbb{E}(|Xe^{-\alpha X}|)$ are finite.

Then we have $B_h = \mathbb{E}\left(e^{-\alpha X}\right)$ and $K_{\phi} = \left(\alpha B_h e^{\alpha \mu}\right)^{-1}$.

Put

$$I_{n,\alpha} = \frac{1}{e^{-\alpha\mu_n}} \frac{1}{n} \sum_{j=1}^n e^{-\alpha X_j} \text{ and } I_\alpha = \frac{1}{e^{-\alpha\mu}} \int_{\mathcal{V}_X} e^{-\alpha x} dF(x).$$

Then

$$\sqrt{n}\left(I_{n,\alpha}-I_{\alpha}\right) = \mathbb{G}_n\left(e^{\alpha\mu}\left(h+\alpha B_h I_d\right)\right) + o_{\mathbb{P}}^*(1).$$

Since $Ko_{n,\alpha} = \tau(I_{n,\alpha})$, we deduce that

$$\sigma_{Ko_{\alpha}}^{2} = \frac{\mathbb{E}e^{-2\alpha X}}{\left(\alpha B_{h}\right)^{2}} + \mathbb{E}X^{2} + \frac{2}{\alpha B_{h}}\mathbb{E}\left(Xe^{\alpha X}\right) - \left(\frac{1}{\alpha} + \mu\right)^{2}.$$

Finally, we summarize the used abbreviations in Table 1, and, for each index, the expression of the function F_{ϕ} and $\mathbb{P}(F_{\phi})$ in Table 2 where we can find the expressions of a_{ϕ} and b_{ϕ} .

Notations	Indices	
$GE(\alpha), \alpha \neq 0, 1$	Generalized Entropy with parameter α	
THEIL	Theil	
MLD	Mean Logarithmic Deviation	
ATK(α), $\alpha < 1$ and $\alpha \neq 0$	Atkinson with parameter α	
CHAMP	Champernowne	
$\operatorname{KOLM}(\alpha)\alpha > 0$	Kolm with parameter α	
$\mathrm{DR}(\alpha)\alpha \ge 0, \alpha \ne 1$	Divergence of Renyi with parameter α	

 $\ddagger \bullet \mathsf{PAPE} \, \mathsf{DJIBY} \, \mathsf{MERGANE}, \dagger^\dagger \, \mathsf{TCHILABALO} \, \mathsf{ABOZOU} \, \mathsf{KPANZOU}, \dagger^{\dagger\dagger} \, \mathsf{DIAM} \, \mathsf{BA}, \, \mathsf{AND} \, \dagger^{\dagger\dagger\dagger} \, \mathsf{GANE} \, \mathsf{SAMB} \, \mathsf{LO}$

TABLE 1. Notations of the indices

Indices	B_h	$F_{\phi}(x), \ \forall x \in \mathcal{V}_X$	$\mathbb{P}(F_{\phi})$
$GE(\alpha)$	$\int_{\mathcal{V}_X} x^{\alpha} dF(x)$	$\frac{1}{\alpha(\alpha-1)}\frac{1}{\mu^{\alpha}}\left(x^{\alpha}-\frac{\alpha B_{h}}{\mu}x\right)$	$\frac{-B_h}{\alpha\mu^{\alpha}}$
THEIL	$\int_{\mathcal{V}_X} x \log x dF(x)$	$\frac{1}{\mu} \left(x \log x - \left(\frac{B_h}{\mu} + 1 \right) x \right)$	-1
MLD	$\int_{\mathcal{V}_X} \log x^{-1} dF(x)$	$\frac{1}{\mu}x - \log x$	$1 + B_h$
$\operatorname{ATK}(\alpha)$	$\int_{\mathcal{V}_X} x^{\alpha} dF(x)$	$\frac{B_h^{1/\alpha}}{\mu} \left(\frac{1}{\mu}x - \frac{B_h^{-1}}{\alpha}x^\alpha\right)$	$\left(1-\frac{1}{\alpha}\right)\frac{B_{h}^{1/\alpha}}{\mu}$
CHAMP	$\int_{\mathcal{V}_X} \log x dF(x)$	$\left(\frac{1}{\mu}x - \log x\right) \frac{\exp(B_h)}{\mu}$	$\frac{1-B_h}{\mu}\exp(B_h)$
$\operatorname{KOLM}(\alpha)$	$\int_{\mathcal{V}_X} \exp\left(-\alpha x\right) dF(x)$	$x + \frac{1}{\alpha B_h} \exp\left(-\alpha x\right)$	$\mu + \frac{1}{\alpha}$
$\overline{\mathrm{DR}}(\alpha)$	$\int_{\mathcal{V}_X} x^{\alpha} dF(x)$	$\frac{1}{\alpha - 1} \left(\frac{1}{B_h} x^{\alpha} - \frac{\alpha}{\mu} x \right)$	-1

TABLE 2. Summary of the functions F for each index

5. Proof of Theorems 1 and 2

Proof of Theorem 1.

On one hand, denote by

(5.1)
$$I_n = \frac{\mathbb{P}_n(h)}{h_1(\mu_n)} - h_2(\mu_n) \text{ and } I = \frac{\mathbb{P}(h)}{h_1(\mu)} - h_2(\mu),$$

by decomposing the difference of I_n and I, we get the next equality

$$I_n - I = \frac{\left(\mathbb{P}_n - \mathbb{P}\right)(h)}{h_1(\mu_n)} - \frac{\mathbb{P}(h)}{h_1(\mu)h_1(\mu_n)} \left(h_1(\mu_n) - h_1(\mu)\right) - \left(h_2(\mu_n) - h_2(\mu)\right).$$

As for all i = 1, 2; the function h_i is continuous on \mathcal{V}_X and using the fact that μ_n converges almost surely to μ , then we have when n tends to infinity

(5.2)
$$h_i(\mu_n) \xrightarrow{as} h_i(\mu) < \infty.$$

We have also

$$\left(\mathbb{P}_n - \mathbb{P}\right)(h) = \frac{1}{n} \sum_{j=1}^n \left(h(X_j) - \mathbb{E}h(X_j)\right).$$

Or the sequence of the random variables $\{h(X_j)\}_{j=1,\dots,n}$ is independent and identically distributed, and as $\mathbb{E}h(X) < \infty$ by the hypothesis (A1.1), then the Law of Large Numbers implies that

(5.3) $(\mathbb{P}_n - \mathbb{P})(h) \xrightarrow{as} 0.$

Finally, using (5.2) and (5.3), we get

$$I_n \xrightarrow{as} I$$
, when $n \to \infty$.

On the other hand, as τ satisfies the hypothesis (A1.2), then we deduce that

$$T_n \xrightarrow{as} T$$
, when $n \to \infty$. \Box

Proof of Theorem 2.

Using the equation (5.1), we have

 $\dagger \Phi \text{APE}$ DJIBY MERGANE, †† TCHILABALO ABOZOU KPANZOU, ††† DIAM BA, AND †††† GANE SAMB LO

$$I_n - I = \frac{\left(\mathbb{P}_n - \mathbb{P}\right)(h)}{h_1(\mu_n)} - \frac{B_h}{h_1(\mu)h_1(\mu_n)} \left(h_1(\mu_n) - h_1(\mu)\right) - \left(h_2(\mu_n) - h_2(\mu)\right).$$

Since h_i is continuously differentiable at μ for i = 1, 2, we get

$$h_i(\mu_n) - h_i(\mu) = h'_i(\mu) \ (\mathbb{P}_n - \mathbb{P}) \ (I_d) + o_{\mathbb{P}}(n^{-\frac{1}{2}}).$$

Then

$$I_{n} - I = \frac{\left(\mathbb{P}_{n} - \mathbb{P}\right)(h)}{h_{1}(\mu_{n})} - \frac{B_{h}}{h_{1}(\mu) h_{1}(\mu_{n})} \left(h_{1}'(\mu) \left(\mathbb{P}_{n} - \mathbb{P}\right)(I_{d}) + o_{\mathbb{P}}^{*}(n^{-\frac{1}{2}})\right) - h_{2}'(\mu) \left(\mathbb{P}_{n} - \mathbb{P}\right)(I_{d}) + o_{\mathbb{P}}^{*}(n^{-\frac{1}{2}}).$$

But

$$\frac{B_h}{h_1(\mu) h_1(\mu_n)} o_{\mathbb{P}}(n^{-\frac{1}{2}}) + o_{\mathbb{P}}^*(n^{-\frac{1}{2}}) = o_{\mathbb{P}}(n^{-\frac{1}{2}}),$$

then we get the next expression

$$I_n - I = \frac{(\mathbb{P}_n - \mathbb{P})(h)}{h_1(\mu)} - \left(\frac{B_h h_1'(\mu)}{h_1^2(\mu)} + h_2'(\mu)\right) (\mathbb{P}_n - \mathbb{P})(I_d) + o_{\mathbb{P}}(n^{-\frac{1}{2}}).$$

Then

$$\sqrt{n}(I_n - I) = \frac{1}{h_1(\mu)} \mathbb{G}_n(h) - \left(\frac{B_h h'_1(\mu)}{h_1^2(\mu)} + h'_2(\mu)\right) \mathbb{G}_n(I_d) + o_{\mathbb{P}}(1).$$

By the linearity property of \mathbb{G}_n , we get

$$\sqrt{n} (I_n - I) = \mathbb{G}_n \left(\frac{1}{h_1(\mu)} h - \left(\frac{B_h h'_1(\mu)}{h_1^2(\mu)} + h'_2(\mu) \right) I_d \right) + o_{\mathbb{P}}(1).$$

Since K_{ϕ} is finite by assumption, we apply a gain the delta-method to the function τ to have

$$\sqrt{n} \left(\tau(I_n) - \tau(I) \right) = \mathbb{G}_n \left(K_\phi \left(\frac{1}{h_1(\mu)} h - \left(\frac{B_h h_1'(\mu)}{h_1^2(\mu)} + h_2'(\mu) \right) I_d \right) \right) + o_{\mathbb{P}}(1).$$

Using the notations of the equation (2), therefore

$$\sqrt{n}\left(\tau(I_n) - \tau(I)\right) = \mathbb{G}_n(a_{\phi}h - b_{\phi}I_d) + o_{\mathbb{P}}^*(1) = \mathbb{G}_n(F_{\phi}) + o_{\mathbb{P}}(1)$$

and we easily obtain by (3.1) the variance σ_{ϕ}^2 . This ends the proof of Theorem 2.

6. Data driven applications and variance computations

We here give data driven applications to show how our results work. We consider the ESAM2 (Enquête Sénégalaise auprès des Ménages, $2^{\text{ème}}$ édition) and the ESPS (Enquête de Suivi de la Pauvreté au Sénégal) databases respectively collected in 2001-2002 and in 2005-2006. (See ANSD SENEGAL (2001-2006)). Sénégal is a member of UEMOA. In both databases, we consider expense variables aggregated at the level of Heads households as indicators of welfare.

We present the data in Table 3.

Data	Years of data collection	Number of households	Mean of the expenses
ESAM2	2001-2002	6565	995.20
ESPS	2005-2006	13568	898.70

TABLE 3. Descriptive Statistics for the Distribution

We proceeded to the computations of the inequality measures and the corresponding variances using R Software (2009). We obtained the results in Table 4.

ESAM2				
TLIM	T(in $%)$	σ_{ϕ}^2		ſ
GE(.5)	36.362	1.643		0
GE(2)	100.984	148.274		0
THEIL	43.102	4.371		ſ
MLD	34.286	1.024		Ν
ATK(.5)	17.355	0.339		A
ATK(5)	37.497	0.532		A
CHAMP	4.846	1.636		(
DR(.5)	19.061	0.339		Ι
DR(2)	110.515	65.043		Ι

ESPS			
TLIM	T(in $%)$	σ_{ϕ}^2	
GE(.5)	22.684	0.238	
GE(2)	34.206	3.709	
THEIL	24.007	0.411	
MLD	23.060	0.202	
ATK(.5)	11.021	0.053	
ATK(5)	29.591	0.280	
CHAMP	3.334	0.519	
DR(.5)	11.677	0.053	
DR(2)	52.124	5.231	

TABLE 4. Results of the variances computations

7. CONCLUSION

The family we introduced allows a flexible and unified approach in the asymptotic theory of a class of inequality indices. In parallel, the computer packages also may be presented in more compact forms. We illustrated both aspects (theoretical and computational) in the paper. Hence the practitioner has all he needs about these indices in one place. But we only studied the finite dimensional limits. In a future paper, we will try to present uniform asymptotic laws of the family index by the parameter $\phi = (\tau, h, h_1, h_2)$.

References

- Agence Nationale de la Statistique et de la Démographie, Dakar, Sénégal. *www.ansd.sn.* Atkinson, A.B. (1970). On the Measurement of Inequality, *Journal of Economic Theory*, **2**, 244-263.
- Barrett, G., & Donald, S. (2009). Statistical inference with generalized Gini indices of inequality, poverty, and welfare. J. Bus. Econom. Statist., 27(1), 1-17. http://dx.doi.org/10.1198/jbes.2009.0001
- Bishop, J.A., Chow, K.V. and Zheng, B. (1995). Statistical Inference and Decomposable Poverty Measures. Bulletin of Economic Research, 47, 329-340.
- Bishop, J.A., Formby, J.P. and Zheng, B. (1997). Statistical Inference and the Sen Index of Poverty. *International Economic Review*, **38** (2), 381-387.
- Chakravarty, S.R. (1983). A new Poverty Index. Mathematical Social Science, 6, 307-313.
- Champernowne, D.G. and Cowell, F. A. (1998). Economic inequality and income distribution,. Cambridge: Cambridge University Press.
- Clark, S., Hemming, R. and Ulph, D. (1981). On Indices for the Measurement of Poverty. *Economic Journal*, 91, 525-526.
- Cowell, F.A. (1980a). Generalized entropy and the measurement of distributional change. European Economic Review, 13, 147-159.
- Cowell, F.A. (1980b). On the structure of additive in equality measures. *Review of Economic Studies*, 47, 521-531.
- Cowell, F.A. (2000). Measurement of inequality. In A.B. Atkinson and F. Bourguignon (Eds.), *Handbook of Income Distribution*, Chapter 2, pp.87-166. Amsterdam: North Holland.
- Cowell, Frank A. (2003). Theil, Inequality and the Structure of Income Distribution. London School of Economics and Political Sciences. available at: http://eprints.lse.ac.uk/2288/.
- Cowell, F.A. and Kuga, K. (1981a). Additivity and the entropy concept: Anaxiomatic approach to inequality measure. *Journal of Economic Theory* **25**, 131-143.
- Cowell, F.A. and Kuga, K. (1981b). Inequality measurement: anaxiomatic approach. European Economic Review, 15, 287-305.
- Dalton H. (1920). The Measurement of Inequality of Incomes, *Economic Journal*, vol. 30, 348-361.
- Davidson, R. and Duclos, J.Y. (2000). Statistical Inference for Stochastic Dominance and for the Measurement of Poverty and Inequality. *Econometrica*, 68 (6), 1435-1464.
- Foster, J.E., Greer, J. and Thorbecke, E. (1984) "A class of Decomposable Poverty Measures" *Econometrica*, 52, 761-766.
- Giné, E. and Zinn, J. (1980). Boostrapping general empirical processes. Annals of Probability, 18 (2), 851-869.
- Greselin, F., Puri, M.L. and Zitikis, R. (2009). L-functions, processes, and statistics in measuring economic inequality and actuarial risks. *Statistics and Its Interface*, 2, 227-245.
- Harsanyi, J. C. (1953). Cardinal utility in welfare economics of concentration. Journal of the Royal Statistical Society, Series A 123, 423-34.

- Harsanyi, J. C. (1955). Cardinal welfare, individualistic ethics and interpersonal comparisons of utility. *Journal of Political Economy* 63, 309-321.
- Kakwani, N. (1980). On a Class of Poverty Measures. Econometrica, 48, 437-446. (MR0560520)
- http://dx.doi.org/10.2307/1911106.
- Kakwani, N. (1993). Statistical Inference in the Measurement of Poverty. The Review of Economics and Statistics, Vol. 75, (No. 4), 632-639.
- Kakwani, N. (2003). Issues on Setting Absolute Poverty Line. Poverty and Social Developpement, Papers, n⁰3, June, Asian Bank of Development Bank (ADB).
- Kolm S. (1976a): Unequal Inequalities I, Journal of Economic Theory, 12, 416-442.
- Kolm S. (1976b): Unequal Inequalities II, Journal of Economic Theory, 13, 82-111.
- Kolm S-C. (1969). The Optimal production of justice, In Public Economics, ed. By J. Margolis, and S. Guitton, London. MacMillan.
- Kullback, S. (1959). Inference Theory and Statistics. New-York: John Wiley.
- Lo G.S. (2010). A simple note on some empirical stochastic process as a tool in uniform L-statistics weak laws. *Afrika Statistika*, 5, 437-446. (MR2920301)
- Lo G.S. and Sall, S.T. (2010). Asymptotic Representation Theorems for Poverty Indices. *Afrika Statistika*, **5**, 238-244. (MR2920300)
- Lo, G.S., Sall, S.T. and Seck, C.T. (2006). Une Théorie asymptotique des indicateurs de pauvreté. C. R. Math. Acad. Sci. Soc. R. Can., **31** (2009), no. 2, 45-52. (MR2535867), (2010m:91167)
- R Development Core Team (2009) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL http://www.R-project.org.
- Renyi A. (1961). On measures of entropy and information. In 4th Berkeley Symposium on Mathematical Statistics and Probability, 1, 547-561.
- Rongve, I. and Beach, C.M. (1997). Estimation and inference for normative inequality indices of distributions in the continuum. *International Economic Review*, **38** (1), 83-96.
- Rothschild, M. and Stiglitz, J. E. (1973). Some further results on the measurement of inequality. *Journal of Economic Theory*, 6, 188-203.
- Sall, S.T. and Lo, G.S. (2007). The Asymptotic Theory of the Poverty Intensity in View of Extreme Values Theory For Two Simple Cases. *Afrika Statistika*, **2**, (1), 41-55
- Sall, S.T. and Lo, G.S. (2010). Uniform Weak Convergence of the time-dependent poverty Measure for Continuous Longitudinal Data. *Brazilian Journal of Probability and Statistics*, 24, (3), 457-467. (MR2719696) http://dx.doi.org/10.1214/08-BJPS101
- Seck, C.T. and Lo, G.S. (2008). Uniform Weak Convergence of the Nonweithed Poverty measure. To appear in *Communication in Statistics*, A.
- Sen, A.K. (1973). on Economic Inequality. Oxford: Clarendon Press.
- Sen, A.K. (1976). Poverty: An Ordinal Approach to Measurement. *Econometrica*, 44, 219-231.
- Shorack, G.R. and Wellner, J.A. (1986). Empirical Processes with Applications to Statistics, wiley-Interscience, New-York.

http://dx.doi.org/10.1137/1.9780898719017

- Shorrocks, A. (1995). Revisiting the Sen Poverty Index. *Econometrica*, **63**, 1225-1230. (doi:10.2307/2171728)
- Temkin, L. (1993). Inequality. Oxford University Press.
- Theil, H. (1967). Economics and Information Theory, Amsterdam, North Holland.
- Van der Vaart, A.W. and Wellner, J.A. (1996). Weak Convergence and Empirical Processes: With Applications to Statistics, Springer-Verlag New-York.
- Zheng, B. (1997). Aggregate Poverty Measures. Journal of Economic Surveys, 11 (2), 123-162.