

DELAYED PERTURBATION OF MITTAG-LEFFLER FUNCTIONS AND THEIR APPLICATIONS TO FRACTIONAL LINEAR DELAY DIFFERENTIAL EQUATIONS

N. I . MAHMUDOV

ABSTRACT. In this paper, we propose a delayed perturbation of Mittag-Leffler type matrix function, which is an extension of the classical Mittag-Leffler type matrix function and delayed Mittag-Leffler type matrix function. With the help of the delayed perturbation of Mittag-Leffler type matrix function, we give an explicit formula of solutions to linear nonhomogeneous fractional delay differential equations.

1. INTRODUCTION

It is known that a solution of a linear system $y'(t) = Ay(t)$, $t \geq 0$ has the form $y(t) = e^{At}y(0)$, where exponential matrix e^{At} is also called fundamental matrix. However, it becomes more complex for seeking a fundamental matrix for linear delay system

$$(1.1) \quad \begin{aligned} y'(t) &= Ay(t) + By(t-h), \quad t \geq 0, h > 0, \\ y(t) &= \varphi(t), \quad -h \leq t \leq 0, \end{aligned}$$

where A, B are two constant square matrices. Under the assumptions that A and B are permutation matrices, Khusainov & Shuklin [4] give a representation of a solution of a linear homogeneous system with delay by introducing the concept of delayed matrix exponential e_h^{Bt} corresponding to delay h and matrix B . They proved that fundamental matrix of linear delay system (1.1) (delayed perturbation of exponential matrix e^{At}) can be given by $e^{At}e_h^{B_1(t-h)}$, $B_1 = e^{-Ah}B$. Notice that the fractional analogue of the same problem was considered by Li and Wang [10] in the case $A = \Theta$. For more recent contributions on oscillating system with pure delay, relative controllability of system with pure delay, asymptotic stability of nonlinear multidelay differential equations, finite time stability of differential equations, one can refer to [1]-[15] and reference therein.

Motivated by Khusainov & Shuklin [4], Li and Wang [10], we extend to consider representation of solutions of a fractional delay differential equation of the form by introducing delayed perturbation of Mittag-Leffler function

$$(1.2) \quad \begin{aligned} ({}^C D_{-h+}^\alpha y)(t) &= Ay(t) + By(t-h) + f(t), \quad t \in (0, T], h > 0, \\ y(t) &= \varphi(t), \quad -h \leq t \leq 0, \end{aligned}$$

where $({}^C D_{-h+}^\alpha y)(\cdot)$ is the Caputo derivative of order $\alpha \in (0, 1)$, $A, B \in R^{n \times n}$ denotes constant matrix, and $\varphi : [-h, 0] \rightarrow R^n$ is an arbitrary Caputo differentiable vector function, $f \in C([-h, T], R^n)$, $t = lh$ for a fixed natural number l .

To end this section, we would like to state the main contribution as follows:

(i) We propose delayed perturbation $X_{h,\alpha,\beta}^{A,B}(t)$ of Mittag-Leffler type functions, by means of the matrix equations (1.4). We show that for $B = \Theta$ the function $X_{h,\alpha,\beta}^{A,B}(t)$ coincide with Mittag-Leffler type

Date: December 26, 1997.

2000 Mathematics Subject Classification. Primary 34A08; Secondary 39B05 .

Key words and phrases. fractional linear systems, delayed perturbation, Mittag-Leffler type matrix function.

function of two parameters $t^{\alpha-1}E_{\alpha,\beta}(At^\alpha)$. For $A = \Theta$ $X_{h,\alpha,\beta}^{A,B}(t)$ coincide with delayed Mittag-Leffler type matrix function of two parameters $E_{h,\alpha,\beta}^B(t-h)$.

(ii) We explicitly represent the solution of fractional delay linear system (1.2) via delayed perturbation of Mittag-Leffler type function.

Definition 1. *Mittag-Leffler type matrix function of two parameters $\Phi_{\alpha,\beta}(A, z) : R \rightarrow R^{n \times n}$ is defined by*

$$\Phi_{\alpha,\beta}(A, z) := z^{\beta-1}E_{\alpha,\beta}(Az^\alpha) := z^{\beta-1} \sum_{k=0}^{\infty} \frac{A^k z^{\alpha k}}{\Gamma(k\alpha + \beta)}, \quad \alpha, \beta > 0, z \in R.$$

Definition 2. *Delayed Mittag-Leffler type matrix function of two parameters $E_{h,\alpha,\beta}^B(t) : R \rightarrow R^{n \times n}$ is defined by*

(1.3)

$$E_{h,\alpha,\beta}^B(t) := \begin{cases} \Theta, & -\infty < t \leq -h, \\ I \frac{(h+t)^{\beta-1}}{\Gamma(\beta)}, & -h < t \leq 0, \\ I \frac{(h+t)^{\beta-1}}{\Gamma(\beta)} + B \frac{t^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + B^2 \frac{(t-h)^{2\alpha+\beta-1}}{\Gamma(2\alpha+\beta)} + \dots + B^k \frac{(t-(k-1)h)^{k\alpha+\beta-1}}{\Gamma(k\alpha+\beta)}, & (k-1)h < t \leq kh. \end{cases}$$

In order to define delayed perturbation of Mittag-Leffler type matrix functions, we introduce the following matrix equation for $Q_k(s)$, $k = 1, 2, \dots$

$$(1.4) \quad \begin{aligned} Q_{k+1}(s) &= A Q_k(s) + B Q_k(s-h), \\ Q_0(s) &= Q_k(-h) = \Theta, \quad Q_1(0) = I, \\ k &= 0, 1, 2, \dots, s = 0, h, 2h, \dots \end{aligned}$$

Simple calculations show that

	$s = 0$	$s = h$	$s = 2h$	$s = 3h$	\dots	$s = ph$
$Q_1(s)$	I	Θ	Θ	Θ	\dots	Θ
$Q_2(s)$	A	B	Θ	Θ	\dots	Θ
$Q_3(s)$	A^2	$AB + BA$	B^2	Θ	\dots	Θ
$Q_4(s)$	A^3	$A(AB + BA) + BA^2$	$AB^2 + B(AB + BA)$	B^3	\dots	Θ
\dots	\dots	\dots	\dots	\dots	\dots	Θ
$Q_{p+1}(s)$	A^p	\dots	\dots	\dots	\dots	B^p

Definition 3. *Delayed perturbation of two parameter Mittag-Leffler type matrix function $X_{h,\alpha,\beta}^{A,B}$ generated by A, B is defined by*

$$(1.5) \quad X_{h,\alpha,\beta}^{A,B}(t) := \begin{cases} \Theta, & -h \leq t < 0, \\ I, & t = 0, \\ \sum_{i=0}^{\infty} \sum_{j=0}^{p-1} Q_{i+1}(jh) \frac{(t-jh)^{i\alpha+\beta-1}}{\Gamma(i\alpha + \beta)}, & (p-1)h < t \leq ph. \end{cases}$$

Lemma 1. *Let $X_{h,\alpha,\beta}^{A,B}(t)$ be defined by (1.5). Then the following holds true:*

- (i): if $A = \Theta$ then $X_{h,\alpha,\beta}^{A,B}(t) = E_{h,\alpha,\beta}^B(t-h)$, $(p-1)h \leq t-h \leq ph$,
- (ii): if $B = \Theta$ then $X_{h,\alpha,\beta}^{A,B}(t) = t^{\beta-1}E_{\alpha,\beta}(At^\alpha)$,
- (iii): if $\alpha = \beta = 1$ and $AB = BA$ then $X_{h,1,1}^{A,B}(t) = e^{At} e_h^{B(t-h)}$, $(p-1)h < t \leq ph$.

Proof. (i) If $A = \Theta$, then

$$Q_{i+1}(jh) = \begin{cases} \Theta, & i \neq j, \\ B^i, & i = j, \end{cases}$$

and $X_{h,\alpha,\beta}^{A,B}(t)$ coincides with $E_{h,\alpha,\beta}^B(t-h)$:

$$\begin{aligned} X_{h,\alpha,\beta}^{A,B}(t) &= \sum_{i=0}^p B^i \frac{(t-ih)^{i\alpha+\beta-1}}{\Gamma(i\alpha+\beta)} = \frac{t^{\beta-1}}{\Gamma(\beta)} + B \frac{(t-h)^{\alpha+\beta-1}}{\Gamma(\alpha+\beta)} + \dots + B^p \frac{(t-ph)^{p\alpha+\beta-1}}{\Gamma(p\alpha+\beta)} \\ &= E_{h,\alpha,\beta}^B(t-h), \quad (p-1)h < t-h \leq ph. \end{aligned}$$

(ii) Trivially, from definition of $X_{h,\alpha,\beta}^{A,B}(t)$ we have: if $B = \Theta$, then

$$X_{h,\alpha,\beta}^{A,B}(t) = \sum_{i=0}^{\infty} A^i \frac{t^{i\alpha+\beta-1}}{\Gamma(i\alpha+\beta)} = t^{\beta-1} E_{\alpha,\beta}(At^\alpha).$$

(iii) It can be easily shown that $Q_{i+1}(jh) = \binom{i}{j} A^{i-j} B^j$. So, for $(p-1)h < t \leq ph$ and $B_1 = e^{-Ah}B$ we have

$$\begin{aligned} X_{h,1,1}^{A,B}(t) &= \sum_{i=0}^{\infty} Q_{i+1}(0) \frac{t^i}{i!} + \sum_{i=1}^{\infty} Q_{i+1}(h) \frac{(t-h)^i}{i!} + \dots + \sum_{i=1}^{\infty} Q_{i+1}((p-1)h) \frac{(t-(p-1)h)^i}{i!} \\ &= \sum_{i=0}^{\infty} A^i \frac{t^i}{i!} + \sum_{i=1}^{\infty} \binom{i}{1} A^{i-1} B \frac{(t-h)^i}{i!} + \dots + \sum_{i=p-1}^{\infty} \binom{i}{p-1} A^{i-p+1} B^{p-1} \frac{(t-(p-1)h)^i}{i!} \\ &= e^{At} + e^{A(t-h)} B(t-h) + \dots + \sum_{i=0}^{\infty} \binom{i+p-1}{p-1} A^i B^{p-1} \frac{(t-(p-1)h)^{i+p-1}}{(i+p-1)!} \\ &= e^{At} + e^{A(t-h)} B(t-h) + \dots + e^{A(t-(p-1)h)} B^{p-1} \frac{1}{(p-1)!} (t-(p-1)h)^{p-1} \\ &= e^{At} \left(I + e^{-Ah} B(t-h) + \dots + e^{-A(p-1)h} B^{p-1} \frac{1}{(p-1)!} (t-(p-1)h)^{p-1} \right) = e^{At} e_h^{B_1(t-h)}. \end{aligned}$$

□

It turns out that $X_{h,\alpha,\beta}^{A,B}(t)$ is a delayed perturbation of the fundamental matrix of the equation (1.2) with $f = 0$.

Lemma 2. $X_{h,\alpha,\alpha}^{A,B} : R \rightarrow R^n$ is a solution of

$$(1.6) \quad {}^C D_{-h+}^\alpha X_{h,\alpha,\alpha}^{A,B}(t) = AX_{h,\alpha,\alpha}^{A,B}(t) + BX_{h,\alpha,\alpha}^{A,B}(t-h).$$

Proof. We verify that $X_{h,\alpha,\alpha}^{A,B}(t)$ satisfies differential equation (1.6) for $t \in (t_p, t_{p+1}]$. We adopt mathematical induction to prove our result.

(i) For $p = 0$, $0 < t \leq h$, we have

$$\begin{aligned} X_{h,\alpha,\alpha}^{A,B}(t) &= t^{\alpha-1} E_{\alpha,\alpha}(At), \quad X_{h,\alpha,\alpha}^{A,B}(t-h) = \Theta, \\ {}^C D_{-h+}^\alpha X_{h,\alpha,\alpha}^{A,B}(t) &= {}^C D_{-h+}^\alpha \sum_{i=1}^{\infty} A^i \frac{t^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} = AX_{h,\alpha,\alpha}^{A,B}(t) = AX_{h,\alpha,\alpha}^{A,B}(t) + BX_{h,\alpha,\alpha}^{A,B}(t-h). \end{aligned}$$

(ii) Suppose $p = n$, $(n-1)h < t \leq nh$ the following relation holds:

$$X_{h,\alpha,\alpha}^{A,B}(t) = \sum_{i=0}^{\infty} \sum_{j=0}^{n-1} Q_{i+1}(jh) \frac{(t-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)}.$$

Next, for $p = n + 1$, $nh < t \leq (n + 1)h$, by elementary computation, one obtains

$$\begin{aligned}
{}^C D_{-h^+}^\alpha X_{h,\alpha,\alpha}^{A,B}(t) &= \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+1}(jh) \frac{\Gamma((i+1)\alpha)}{\Gamma(i\alpha)} \frac{(t-jh)^{i\alpha-1}}{\Gamma((i+1)\alpha)} \\
&= \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+1}(jh) \frac{(t-jh)^{i\alpha-1}}{\Gamma(i\alpha)} = \sum_{i=0}^{\infty} \sum_{j=0}^n (AQ_i(jh) + BQ_i(jh-h)) \frac{(t-jh)^{i\alpha-1}}{\Gamma(i\alpha)} \\
&= \sum_{i=0}^{\infty} \sum_{j=0}^n AQ_{i+1}(jh) \frac{(t-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} + \sum_{i=0}^{\infty} \sum_{j=0}^{n-1} BQ_{i+1}(jh) \frac{(t-h-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} \\
&= AX_{h,\alpha,\alpha}^{A,B}(t) + BX_{h,\alpha,\alpha}^{A,B}(t-h).
\end{aligned}$$

This ends the proof. \square

Lemma 3. Let $(k-1)h < t \leq kh$, $-h \leq s \leq t$. We have

$$\int_s^t (t-r)^{-\alpha} X_{h,\alpha,\alpha}^{A,B}(r-s) dr = (t-s-jh)^{-\alpha+i\alpha+\beta} \sum_{i=0}^{\infty} \sum_{j=0}^{p-1} Q_{i+1}(jh) \frac{\Gamma(1-\alpha)}{\Gamma(i\alpha+\beta+1-\alpha)}.$$

Theorem 1. The solution $y(t)$ of (1.2) satisfying zero initial condition, has a form

$$y(t) = \int_{-h}^t X_{h,\alpha,\alpha}^{A,B}(t-s) f(s) ds, \quad t \geq 0.$$

Proof. By using the method of variation of constants, any solution of nonhomogeneous system $y(t)$ should be satisfied in the form

$$(1.7) \quad y(t) = \int_{-h}^t X_{h,\alpha,\alpha}^{A,B}(t-s) c(s) ds, \quad t \geq 0,$$

where $c(s)$, $0 \leq s \leq t$ is an unknown vector function and $y(0) = 0$. Having Caputo fractional differentiation on both sides of (1.7), we obtain the following cases:

(i) For $0 < t \leq h$ we have

$$\begin{aligned}
({}^C D_{-h^+}^\alpha y)(t) &= Ay(t) + By(t-h) + f(t) \\
&= A \int_{-h}^t X_{h,\alpha,\alpha}^{A,B}(t-s) c(s) ds + \int_{-h}^{t-h} X_{h,\alpha,\alpha}^{A,B}(t-h-s) c(s) ds + f(t) \\
&= A \int_{-h}^t X_{h,\alpha,\alpha}^{A,B}(t-s) c(s) ds + f(t).
\end{aligned}$$

According to Lemma 3, we have

$$\begin{aligned}
({}^C D_{-h^+}^\alpha y)(t) &= \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_{-h}^t (t-r)^{-\alpha} \left(\int_{-h}^r X_{h,\alpha,\alpha}^{A,B}(r-s) c(s) ds \right) dr \\
&= \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_{-h}^t c(s) \int_s^t (t-r)^{-\alpha} \sum_{i=0}^{\infty} A^i \frac{(r-s)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} dr ds \\
&= \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_{-h}^t c(s) \int_s^t (t-r)^{-\alpha} \frac{(r-s)^{\alpha-1}}{\Gamma(\alpha)} dr ds \\
&+ \frac{1}{\Gamma(1-\alpha)} \sum_{i=1}^{\infty} A^i \frac{d}{dt} \int_{-h}^t c(s) \int_s^t (t-r)^{-\alpha} \frac{(r-s)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} dr ds
\end{aligned}$$

$$\begin{aligned}
 &= c(t) + \sum_{i=1}^{\infty} \frac{1}{\Gamma(1-\alpha)\Gamma((i+1)\alpha)} A^i \frac{d}{dt} \int_{-h}^t c(s) (t-s)^{i\alpha-1} B(\alpha(i+1), 1-\alpha) ds \\
 &= c(t) + \sum_{i=1}^{\infty} \frac{\Gamma(1-\alpha)\Gamma((i+1)\alpha)}{\Gamma(1-\alpha)\Gamma((i+1)\alpha)\Gamma(1+i\alpha)} A^i \frac{d}{dt} \int_{-h}^t c(s) (t-s)^{i\alpha} ds \\
 &= c(t) + \sum_{i=1}^{\infty} A^i \frac{1}{\Gamma(1+i\alpha)} \frac{d}{dt} \int_{-h}^t c(s) (t-s)^{i\alpha} ds \\
 &= c(t) + \sum_{i=1}^{\infty} A^i \frac{\alpha i}{\Gamma(1+i\alpha)} \int_{-h}^t c(s) (t-s)^{i\alpha-1} ds = c(t) + \int_{-h}^t \sum_{i=1}^{\infty} A^i \frac{1}{\Gamma(i\alpha)} (t-s)^{i\alpha-1} c(s) ds \\
 &= c(t) + A \int_{-h}^t \sum_{i=0}^{\infty} A^i \frac{(t-s)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} c(s) ds = c(t) + A \int_{-h}^t X_{h,\alpha,\beta}^{A,B}(t-s) c(s) ds.
 \end{aligned}$$

Hence, we obtain $c(t) = f(t)$.

(ii) For $kh < t \leq (k+1)h$, according to (1.2), we have

$$\begin{aligned}
 ({}^C D_{-h^+}^\alpha y)(t) &= Ay(t) + By(t-h) + f(t) \\
 &= A \int_{-h}^t X_{h,\alpha,\alpha}^{A,B}(t-s) c(s) ds + B \int_{-h}^{t-h} X_{h,\alpha,\alpha}^{A,B}(t-h-s) c(s) ds + f(t) \\
 &= A \int_{-h}^t \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+1}(jh) \frac{(t-s-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} c(s) ds \\
 &\quad + B \int_{-h}^{t-h} \sum_{i=0}^{\infty} \sum_{j=0}^{n-1} Q_{i+1}(jh) \frac{(t-s-h-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} c(s) ds + f(t) \\
 &= A \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+1}(jh) \int_{-h}^{t-jh} \frac{(t-s-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} c(s) ds \\
 &\quad + B \sum_{i=0}^{\infty} \sum_{j=1}^n Q_{i+1}(jh-h) \int_{-h}^{t-jh} \frac{(t-s-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} c(s) ds + f(t) \\
 &= \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+2}(jh) \int_{-h}^{t-jh} \frac{(t-s-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} c(s) ds + f(t).
 \end{aligned}$$

According to Lemma 3, we have

$$\begin{aligned}
 &({}^C D_{-h^+}^\alpha y)(t) \\
 &= \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_{-h}^t (t-r)^{-\alpha} \left(\int_{-h}^r X_{h,\alpha,\alpha}^{A,B}(r-s) c(s) ds \right) dr \\
 &= \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_{-h}^t c(s) \int_s^t (t-r)^{-\alpha} X_{h,\alpha,\alpha}^{A,B}(r-s) dr ds \\
 &= \frac{1}{\Gamma(1-\alpha)} \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+1}(jh) \frac{d}{dt} \int_{-h}^t c(s) \int_s^t (t-r)^{-\alpha} \frac{(r-s-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} dr ds
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\Gamma(1-\alpha)} \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+1}(jh) \frac{d}{dt} \int_{-h}^{t-jh} c(s) (t-s-jh)^{i\alpha} \frac{\Gamma(1-\alpha)}{\Gamma(i\alpha+1)} ds \\
&= \sum_{j=0}^n Q_1(jh) \frac{d}{dt} \int_{-h}^{t-jh} c(s) ds + \sum_{i=1}^{\infty} \sum_{j=0}^n Q_{i+1}(jh) \int_{-h}^{t-jh} c(s) (t-s-jh)^{i\alpha-1} \frac{1}{\Gamma(i\alpha)} ds \\
&= c(t) + \sum_{i=0}^{\infty} \sum_{j=0}^n Q_{i+2}(jh) \int_{-h}^{t-jh} \frac{(t-s-jh)^{(i+1)\alpha-1}}{\Gamma((i+1)\alpha)} c(s) ds.
\end{aligned}$$

Hence, we obtain $c(t) = f(t)$. The proof is completed. \square

Theorem 2. Let $p = 0, 1, \dots, l$. A solution $y \in C(((p-1)h, ph], R^n)$ of (1.2) has a form

$$y(t) = X_{h,\alpha,1}^{A,B}(t+h) \varphi(0) + \int_{-h}^0 X_{h,\alpha,\alpha}^{A,B}(t-s) (({}^C D_{-h+}^\alpha \varphi)(s) - A\varphi(s)) ds.$$

Proof. We looking for a solution of the form

$$y(t) = X_{h,\alpha,1}^{A,B}(t+h) c + \int_{-h}^0 X_{h,\alpha,\alpha}^{A,B}(t-s) g(s) ds,$$

where c is an unknown constants, $g(t)$ is an unknown continuously differentiable function. Moreover, it satisfies initial condition $y(t) = \varphi(t)$, $-h \leq t \leq 0$, i.e.,

$$y(t) = X_{h,\alpha,1}^{A,B}(t+h) c + \int_{-h}^0 X_{h,\alpha,\alpha}^{A,B}(t-s) g(s) ds := \varphi(t), \quad -h \leq t \leq 0.$$

Let $t = -h$ we have

$$X_{h,\alpha,1}^{A,B}(-h-s) = \begin{cases} \Theta, & -h < s \leq 0, \\ I, & s = -h. \end{cases}$$

Thus $c = \varphi(-h)$. Since $-h \leq t \leq 0$, one obtains

$$X_{h,\alpha,1}^{A,B}(t-s) = \begin{cases} \Theta, & t < s \leq 0, \\ E_{\alpha,1}(A(t-s)^\alpha), & -h \leq s \leq t, 0 \leq t-s \leq t+h \leq h. \end{cases}$$

Thus on interval $-h \leq t \leq 0$, one can derive that

$$\begin{aligned}
(1.8) \quad \varphi(t) &= X_{h,\alpha,1}^{A,B}(t+h) \varphi(-h) + \int_{-h}^0 X_{h,\alpha,\alpha}^{A,B}(t-s) g(s) ds \\
&= X_{h,\alpha,1}^{A,B}(t+h) \varphi(-h) + \int_{-h}^t X_{h,\alpha,\alpha}^{A,B}(t-s) g(s) ds + \int_t^0 X_{h,\alpha,\alpha}^{A,B}(t-s) g(s) ds \\
&= E_{\alpha,1}(A(t+h)^\alpha) \varphi(-h) + \int_{-h}^t (t-s)^{\alpha-1} E_{\alpha,\alpha}(A(t-s)^\alpha) g(s) ds.
\end{aligned}$$

Having differentiated (1.8), we obtain

$$\begin{aligned}
({}^C D_{-h+}^\alpha \varphi)(t) &= A \sum_{k=0}^{\infty} \frac{A^k (t+h)^{\alpha k}}{\Gamma(1+k\alpha)} \varphi(-h) + \int_{-h}^t \sum_{k=1}^{\infty} \frac{A^k (t-s)^{\alpha k-1}}{\Gamma(k\alpha)} g(s) ds + g(t) \\
&= A E_{\alpha,1}(A(t+h)^\alpha) \varphi(-h) + A \int_{-h}^t E_{\alpha,\alpha}(A(t-s)^\alpha) g(s) ds + g(t) \\
&= A\varphi(t) + g(t).
\end{aligned}$$

Therefore, $g(t) = ({}^C D_{-h+}^\alpha \varphi)(t) - A\varphi(t)$ and the desired result holds. \square

Combining Theorems 1 and 2, we have the following result.

Corollary 1. *A solution $y \in C([-h, T] \cap ((p-1)h, ph], R^n)$ of (1.2) has a form*

$$y(t) = X_{h,\alpha,1}^{A,B}(t+h)\varphi(-h) + \int_{-h}^0 X_{h,\alpha,\alpha}^{A,B}(t-s) [({}^C D_{-h+}^\alpha \varphi)(s) - A\varphi(s)] ds \\ + \int_0^t X_{h,\alpha,\alpha}^{A,B}(t-s)f(s) ds.$$

REFERENCES

- [1] J. Diblík, M. Fečkan, M. Pospíšil, Representation of a solution of the Cauchy problem for an oscillating system with two delays and permutable matrices, *Ukrainian Math. J.* 65 (2013) 58–69.
- [2] J. Diblík, D.Ya. Khusainov, J. Baštinec, A.S.Sirenko, Exponential stability of linear discrete systems with constant coefficients and single delay. *Appl. Math. Lett.*, 51, 68–73. (2016)
- [3] J. Diblík, M. Fečkan, M. Pospíšil, Representation of a solution of the cauchy problem for an oscillating system with two delays and permutable matrices, *Ukrainian Math. J.* 65 (2013) 58–69.
- [4] D.Ya. Khusainov, G.V. Shuklin, Linear autonomous time-delay system with permutation matrices solving, *Stud. Univ.Žilina* 17 (2003) 101–108.
- [5] D.Ya. Khusainov and G.V. Shuklin, Relative controllability in systems with pure delay, *Int. J. Appl. Math.* 2 (2005) 210–221.
- [6] M. Medveď, M. Pospíšil, and L. Škripková, Stability and the nonexistence of blowing-up solutions of nonlinear delay systems with linear parts defined by permutable matrices, *Nonlinear Anal.* 74 (2011) 3903–3911.
- [7] M. Medved’ and M. Pospíšil, Sufficient conditions for the asymptotic stability of nonlinear multidelay differential equations with linear parts defined by pairwise permutable matrices, *Nonlinear Anal.* 75 (2012) 3348–3363.
- [8] M. Medved’ and M. Pospíšil, Representation of solutions of systems linear differential equations with multiple delays and linear parts given by nonpermutable matrices, *Journal of Mathematical Sciences*, Vol. 228, No. 3, January, 2018
- [9] M. Li, J. Wang, Finite time stability of fractional delay differential equations. *Appl. Math. Lett.* 64 (2017), 170–176.
- [10] M. Li, J. Wang, Exploring delayed Mittag-Leffler type matrix functions to study finite time stability of fractional delay differential equations, *Appl. Math. Comput.* 324 (2018) 254–265.
- [11] M. Li, A. Debbouche, J. Wang, Relative controllability in fractional differential equations with pure delay, *Mathematical Methods in the Applied Sciences*, DOI: 10.1002/mma.4651
- [12] C. Liang, J. Wang, D. O’Regan, Representation of solution of a fractional linear system with pure delay, *Appl. Math. Lett.* 77 (2018) 72–78.
- [13] Luo, Z. & Wang, J. (2016) Finite time stability analysis of systems based on delayed exponential matrix. *J. Appl. Math. Comput.*, doi:10.1007/s12190-016-1039-2.
- [14] M. Pospíšil, Representation and stability of solutions of systems of functional differential equations with multiple delays. *Electronic Journal of Qualitative Theory of Differential Equations* 2012: 54.(2012).
- [15] M. Pospíšil, Representation of solutions of delayed difference equations with linear parts given by pairwise permutable matrices via Z-transform. *Appl. Math. Comput.*, 294, 180–194.(2017)

DEPARTMENT OF MATHEMATICS, EASTERN MEDITERRANEAN UNIVERSITY, FAMAGUSTA, T.R. NORTH CYPRUS, VIA MERSIN 10, TURKEY

E-mail address: nazim.mahmudov@emu.edu.tr