

Quantum Distributions for the Plane Rotator

Marius Grigorescu

Quantum phase-space distributions (Wigner functions) for the plane rotator are defined using wave functions expressed in both angle and angular momentum representations, with emphasis on the quantum superposition between the Fourier dual variable and the canonically conjugate coordinate. The standard quantization condition for angular momentum appears as necessary for consistency. It is shown that at finite temperature the time dependence of the quantum wave functions may provide classical sound waves. Non-thermal quantum entropy is associated with localization along the orbit.

1 Introduction

The action-angle coordinates $\{(J_i, \varphi_i), i = 1, N/J_i \in \mathbb{R}, \varphi_i \in [-\pi, \pi]\}$ on the phase-space M arise in the description of the integrable Hamiltonian systems with periodic orbits [1]. In these variables¹ the Hamilton function depends only on $\mathbf{J} \equiv \{J_i, i = 1, N\}$, such that a submanifold $\Sigma_{\mathbf{J}}$ of constant \mathbf{J} is a torus parameterized by $\{\varphi_i, i = 1, N\}$. In the old quantum mechanics J_i takes only a discrete set of values, $J_i = n_i \hbar$, $n_i \in \mathbb{Z}$, $2\pi\hbar = h = 6.626 \times 10^{-34}$ J·s, such that the corresponding Lagrangian submanifolds $\Sigma_{\mathbf{n}\hbar}$ provide a partition of M in cells $b_{\mathbf{n}}$ of volume h^N . However, these cells are not ordered along a complete set of local coordinates on M , and in the limit $h \rightarrow 0$ they become singular submanifolds of M , rather than points.

Probability distributions of particles on M may arise from thermal fluctuations, or from an intrinsic "quantum structure", resembling the partition in cells $b_{\mathbf{n}}$ of finite volume. The quantum structure on $M = T^*\mathbb{R}^N$ is usually associated with the Wigner transform [2, 3] $f_{\psi} \in \mathcal{F}(M)$ of the quantum "wave function" $\psi \in L^2(\mathbb{R}^N)$, defined in Cartesian coordinates. For integrable systems the quantum distributions f_{ψ_n} provided by the eigenstates ψ_n of the Hamiltonian operator show an increased localization probability on $\Sigma_{\mathbf{n}\hbar} \subset M$ [4], but despite constant effort, a direct definition of f_{ψ} in terms of the action-angle variables is faced with difficulties. Various aspects of the problem are presented in [5, 6, 7, 8].

In this work the quantum distributions for the action-angle variables are discussed on the representative example of the plane rotator ($M = T^*S^1$). The treatment is similar to the one applied before to the rigid rotator [9], but instead of discretization here the emphasis is on the quantum superposition between the symplectic dual (canonically conjugate) and Fourier dual variables. The basic elements of the formalism are presented in Section 2, followed in Section 3 by applications to the Wigner functions f_{ψ} of the plane rotator. Finite temperature effects, beyond the single particle coherence time, are discussed in Section 4. Concluding remarks are summarized in Section 5.

2 The partial Fourier transform as Hermitian operator

Let $f(x, y)$ be a real integrable function of $x, y \in \mathbb{R}$ and $\tilde{f}(x, k)$ the partial Fourier transform of f only with respect to y ,

$$\tilde{f}(x, k) = \int dy e^{iky} f(x, y) . \quad (1)$$

Because $\tilde{f}(x, k)^* = \tilde{f}(x, -k)$, we may consider $\tilde{f}(x, k)$ as matrix element of a Hermitian operator \hat{f} on $L^2(\mathbb{R})$, having x and k as indices not along rows and columns, but along the diagonals [10]. In the case of $M = T^*\mathbb{R}$ parameterized by the canonical variables (q, p) , the Fourier transform in momentum $\hat{f}(q, k)$ of $f \in \mathcal{F}(M)$ (the set of smooth functions on M), provides a matrix element $\hat{f}_{ab} \equiv h^{-1} \tilde{f}((a+b)/2, (a-b)/\hbar)$ (the "Weyl

¹In the standard approach J_i are considered as coordinates and φ_i as momenta. For systems with symmetry J_i are provided by the momentum mapping and φ_i are group coordinates.

quantization" of f) with the row and column indices $a = q + \hbar k/2$, $b = q - \hbar k/2$ defined using \hbar as a conversion factor from k to q -scale. Thus, if $f_1, f_2 \in \mathcal{F}(M)$ then [10]

$$(f_1, f_2) = \int_M dq dp f_1(q, p) f_2(q, p) = \hbar \int da \int db \hat{f}_{1ab} \hat{f}_{2ba} \equiv \hbar \text{Tr}(\hat{f}_1 \hat{f}_2) . \quad (2)$$

The change of integration variables from (q, p) to (a, b) is completely formal and it does not change the physics (classical or quantum) of the observables f_1, f_2 . However, it distinguishes between a pure quantum distribution $\mathbf{f}_\psi \in \mathcal{F}(M)$ and other observables by reducing $\hbar \hat{\mathbf{f}}_\psi$ to a projection operator, $\hbar(\hat{\mathbf{f}}_\psi)_{ab} = \psi_a \psi_b^*$, $\psi \in L^2(\mathbb{R})$, $\|\psi\| = 1$. In this case the expectation value of $A \in \mathcal{F}(M)$ with respect to \mathbf{f}_ψ is

$$\langle A \rangle_{\mathbf{f}_\psi} = (\mathbf{f}_\psi, A) = \hbar \text{Tr}(\hat{\mathbf{f}}_\psi \hat{A}) = \langle \psi | \hat{A} | \psi \rangle .$$

Similar results can be obtained using the "momentum representation", defined by the Fourier transform in coordinate,

$$\tilde{f}'(k', p) = \int dq e^{ik'q} f(q, p) , \quad (3)$$

such that $\hat{f}'_{b'a'} \equiv \hbar^{-1} \tilde{f}'((a' - b')/\hbar, (a' + b')/2)$, with $a' = p + \hbar k'/2$, $b' = p - \hbar k'/2$. It can be shown that if $\hbar \hat{\mathbf{f}}$ is separable as $\hbar \hat{\mathbf{f}}_{ab} = \psi_a \psi_b^*$, then $\hbar \hat{\mathbf{f}}'$ is also separable, $\hbar \hat{\mathbf{f}}'_{b'a'} = \psi'_{b'} (\psi'_{a'})^*$, with

$$\psi'_p = \frac{1}{\sqrt{2\pi\hbar}} \int dq e^{-ipq/\hbar} \psi_q .$$

This result ensures that both marginal distributions are positive definite,

$$F_\psi^{cs}(q) \equiv \int dp \mathbf{f}_\psi(q, p) = \psi_q \psi_q^* , \quad F_\psi^{ms}(p) \equiv \int dq \mathbf{f}_\psi(q, p) = \psi'_p \psi'^*_p ,$$

and that we may consider ψ_q and ψ'_p as components of the same "state vector" $|\psi\rangle$ in dual bases, $|q\rangle$ ($\equiv |\hbar k\rangle$), and $|p\rangle$, formally related by Fourier transform,

$$|p\rangle = \frac{1}{\sqrt{2\pi\hbar}} \int dq e^{ipq/\hbar} |q\rangle .$$

It is interesting to note that the ordering of the matrix indices a, b does not always follow the one of the variables q, k . Thus, for variations $\delta a > 0$, $\delta b > 0$ we get also $\delta q > 0$, but $\delta k > 0$ only if $\delta a > \delta b$. A related aspect is the sensitivity of \mathbf{f}_ψ to the local inversion symmetry of ψ , as $\mathbf{f}_\psi(q, 0)$ has large negative values at the points q_n where $\psi(q_n + \delta q) = -\psi(q_n - \delta q)$.

Because $F_\psi^{cs}(q) > 0$ and $F_\psi^{ms}(p) > 0$, the negative values of $\mathbf{f}_\psi(q, p)$ indicate that in a quantum distribution the canonical coordinates (q, p) are not independent, but correlated by the implicit dependence of \mathbf{f}_ψ on the Fourier dual variables, k or k' . A measure of these correlations is given by the function $C_\psi(q, p) = \mathbf{f}_\psi(q, p) - F_\psi^{cs}(q) F_\psi^{ms}(p)$.

3 Distributions for the plane rotator

A distribution function $f(\varphi, J)$ of angle (φ) and orbital angular momentum ($J \equiv L_z$) on $M = T^*S^1 \simeq S^1 \times \mathbb{R}$ may describe an ensemble of beads on a circle, and can be regarded as a constrained distribution on $T^*\mathbb{R}^2$ (Appendix 1). Along the lines of Section 2 we may also start with the partial Fourier transform

$$\tilde{f}(\varphi, k) = \int dJ e^{ikJ} f(\varphi, J) . \quad (4)$$

To proceed towards the one-particle quantum distributions one should note that if we let $k \in \mathbb{R}$ and $\varphi \in [-\pi, \pi]$ then $\alpha = \varphi + \hbar k/2$, $\beta = \varphi - \hbar k/2$ are not well defined as indices for a matrix element $\hat{f}_{\alpha\beta} = \hbar^{-1} \tilde{f}((\alpha + \beta)/2, (\alpha - \beta)/\hbar)$ of a Hermitian operator \hat{f} on the quantum Hilbert space $\mathcal{H} = L^2(S^1)$. Therefore, following the example of the rigid rotator [9], quantum distributions f_ψ can be defined using a separable expression $\tilde{f}_\psi(\varphi, k) \equiv \psi_\alpha \psi_\beta^*$, only if the range of $\gamma = \hbar k$ is restricted to the first "Brillouin zone", $\gamma \in [-\pi, \pi]$. In this case one obtains

$$f_\psi(\varphi, J) = \frac{1}{2\pi\hbar} \int_{-\pi}^{\pi} d\gamma e^{-i\gamma J/\hbar} \psi(\varphi + \frac{\gamma}{2}) \psi^*(\varphi - \frac{\gamma}{2}) , \quad (5)$$

in agreement with $V_\psi(\theta, p)$ of [5]. The overlap between two such functions is

$$(f_{\psi_1}, f_{\psi_2}) = \int_{-\infty}^{\infty} dJ \int_{-\pi}^{\pi} d\varphi f_{\psi_1} f_{\psi_2} = \hbar \text{Tr}(\hat{f}_{\psi_1} \hat{f}_{\psi_2}) = \frac{|\langle \psi_1 | \psi_2 \rangle|^2}{2\pi\hbar} , \quad (6)$$

where $\langle \psi_1 | \psi_2 \rangle \equiv \oint d\varphi \psi_1^* \psi_2$ is the scalar product between ψ_1 and ψ_2 as elements of \mathcal{H} . The marginal distributions provided by (5) are

$$F_\psi^{cs}(\varphi) = \int dJ f_\psi(\varphi, J) = \psi_\varphi \psi_\varphi^* , \quad (7)$$

positive definite, and

$$F_\psi^{ms}(J) = \oint d\varphi f_\psi(\varphi, J) = \frac{1}{\hbar} \langle \psi | \hat{P}_J | \psi \rangle , \quad (8)$$

where $(\partial_\varphi \equiv \partial/\partial\varphi)$,

$$\hat{P}_J = \frac{1}{2\pi} \oint d\gamma e^{i\gamma(\hat{J}-J)/\hbar} , \quad \hat{J} = -i\hbar\partial_\varphi . \quad (9)$$

If $\psi_n(\varphi) = e^{in\varphi}/\sqrt{2\pi}$, $n \in \mathbb{Z}$, is an "integral" eigenstate² of \hat{J} , ($\hat{J}\psi_n = n\hbar\psi_n$), then $\langle \psi_n | \hat{P}_J | \psi_n \rangle = j_0(\pi(n - J/\hbar))$, $j_0(x) = \sin(x)/x$. This shows that $F_\psi^{ms}(J)$ is not positive definite if $J/\hbar \in \mathbb{R}$, but if $J/\hbar \in \mathbb{Z}$ then $\hat{P}_{n\hbar}$, $n \in \mathbb{Z}$ becomes a projection operator on

²For the harmonic oscillator Hamiltonian H on $T^*\mathbb{R}$ the action variable $J = H/\omega$ is positive, $(\sqrt{2J}, \varphi)$ are polar coordinates on $T^*\mathbb{R}$, $(dp \wedge dq = dJ \wedge d\varphi)$, and the eigenstates of \hat{J} are real.

ψ_n , $\hat{P}_{n\hbar} = |\psi_n\rangle\langle\psi_n|$, and $F_\psi^{ms}(n\hbar) = |\langle\psi|\psi_n\rangle|^2/\hbar \geq 0$. Moreover, if $J/\hbar \in \mathbb{Z}$ and ψ is a function of good parity, $\psi(\varphi + \pi) = \pm\psi(\varphi)$, then the integral (5) becomes intrinsic on S^1 , namely invariant to a change of parameter $\gamma \rightarrow \gamma + 2\pi$.

To obtain quantum distributions in the angular momentum representation the approach is similar, but also faced with difficulties. Because φ has a finite range a function $f \in \mathcal{F}(M)$ can be expanded in a Fourier series,

$$f(\varphi, J) = \frac{1}{2\pi\hbar} \sum_{m \in \mathbb{Z}} e^{-im\varphi} \tilde{f}'(m, J/\hbar) \quad , \quad (10)$$

but with $J/\hbar \in \mathbb{R}$ and $m \in \mathbb{Z}$ we cannot take $a = J/\hbar + m/2$ and $b = J/\hbar - m/2$ as indices of a matrix element. However, for $f_\psi(\varphi, J)$ of the form (5), if $J/\hbar \in \mathbb{Z}$ and ψ is of good parity³, then $a, b \in \mathbb{Z}$ too, and in the expansion (10) $\tilde{f}'_{\psi'}(m, J/\hbar) = \psi'_b(\psi'_a)^*$ with

$$\psi'_n = \frac{1}{\sqrt{2\pi}} \oint d\varphi e^{-in\varphi} \psi_\varphi \quad . \quad (11)$$

In this representation

$$F_{\psi'}^{cs}(\varphi) = \hbar \sum_{n \in \mathbb{Z}} f_{\psi'}(\varphi, n\hbar) = \psi_\varphi \psi_\varphi^* \quad , \quad (12)$$

(considering $\int dJ = \hbar \sum_{n=J/\hbar}$), and

$$F_{\psi'}^{ms}(J) = \oint d\varphi f_{\psi'}(\varphi, J) = \psi'_{J/\hbar}(\psi'_{J/\hbar})^* \quad . \quad (13)$$

Moreover, for $f_1, f_2 \in \mathcal{F}(M)$ we get $(f_1, f_2) = \hbar Tr'(f_1 \hat{f}'_2)$, where $Tr' \hat{A}' \equiv \sum_{b \in \mathbb{Z}} \hat{A}'_{bb}$ and $\hat{f}'_{ba} = \tilde{f}'(m, J/\hbar)/\hbar$, $m = a - b$, $J = \hbar(a + b)/2$. In particular, $\hat{1}'_{ba} = \delta_{ba}$, $\hat{J}'_{ba} = \hbar a \delta_{ba}$, and

$$\hat{\varphi}'_{ba} = -\frac{i}{a-b} (-1)^{a-b} (1 - \delta_{ab}) \quad . \quad (14)$$

The angle operator $\hat{\varphi}'$ coincides with $(\hat{\phi}_{-\pi})_p$ from [11], and corresponds to the series expansion

$$\varphi = - \sum_{m \neq 0} \frac{(-1)^m}{m} \sin m\varphi \quad .$$

One should consider though φ only as a local coordinate, because at the points of discontinuity $\varphi = \pm\pi$ this series contains $\pm \sin m\pi = 0$, while instead of π , as is $\lim_{n \rightarrow \infty} (\pi - \varphi/n)$, the limit

$$\lim_{n \rightarrow \infty} 2 \sum_{m=1}^n \frac{\sin(m\varphi/n)}{m}$$

yields 1.08949π (the "Gibbs phenomenon").

³ This means that m is an even integer. The odd values of m enlarge the domain $\{a \in \mathbb{Z}\}$ of ψ'_a by new points, $a = n + 1/2$, $n \in \mathbb{Z}$.

4 Coherence properties and temperature effects

Similarly to the case of the free particle on the \mathbb{R} -axis [12], also for the free plane rotator the quantum distribution $f_\psi(\varphi, J)$ is coherent, in the sense that if f_ψ is a solution of the classical Liouville equation,

$$\partial_t f_\psi + \frac{J}{I} \partial_\varphi f_\psi = 0 \quad , \quad (15)$$

then ψ is a solution of the Schrödinger equation, $i\hbar\partial_t\psi = \hat{H}\psi$, $\hat{H} = \hat{J}^2/2I$, by I denoting the moment of inertia.

At a finite temperature T we may consider the thermal average over quantum distributions of the form (10),

$$f_T(\varphi, J) = \frac{1}{2\pi\hbar} \sum_{m \in \mathbb{Z}} e^{-im\varphi} \tilde{f}'_T(m, J/\hbar) \quad , \quad (16)$$

where $\tilde{f}'_T(m, J/\hbar) = \sum_{s \in \mathcal{S}} w_{s,T} \psi_b^s(\psi_a^s)^*$, $a = J/\hbar + m/2$, $b = J/\hbar - m/2$, and $w_{s,T}$ is the thermal distribution function, (e.g. $w_{s,T} \sim e^{-E_s/k_B T}$), over a set \mathcal{S} of one-particle states s with energy E_s and average angular momentum J_s .

At thermal equilibrium, during a small single-particle coherence time τ [13], a quantum wave function $\psi^s(J/\hbar + \mu, t)$, $\mu = \pm m/2$, will become

$$\psi^s(J/\hbar + \mu, t + \tau) = e^{-\frac{i}{\hbar}(E_s - \mu\hbar J_s/I)\tau} \psi^s(J/\hbar + \mu, t) \quad ,$$

such that \tilde{f}'_T changes into

$$\tilde{f}'_T(m, J/\hbar, t + \tau) = \sum_{s \in \mathcal{S}} w_{s,T} e^{-i\tau m J_s/I} \psi_b^s(\psi_a^s)^*|_t \quad .$$

Presuming that in the sum above we can approximate $w_{s,T}(J_s^2 - \langle J^2 \rangle_T) \approx 0$, with $\langle J^2 \rangle_T = \sum_s w_{s,T} J_s^2$, we get

$$\partial_t^2 \tilde{f}'_T(m, J/\hbar, t) = \partial_\tau^2 \tilde{f}'_T(m, J/\hbar, t + \tau)|_{\tau=0} \approx -m^2 \Omega_T^2 \tilde{f}'_T(m, J/\hbar, t) \quad ,$$

where $\Omega_T^2 = \langle J^2 \rangle_T / I^2$. In this approximation we find the transition expected in [12], from the complex wave functions ψ^s to real classical waves, (thermal noise), because for a time $t \gg \tau$, f_T of (16) is a solution of the classical wave equation $\partial_t^2 f_T = \Omega_T^2 \partial_\varphi^2 f_T$. The result is independent of \hbar and should hold also during a macroscopic perturbation, with the condition of constructive interference along the circle providing a discrete spectrum of "angular wavelengths".

5 Concluding remarks

Quasiprobability distributions (Wigner functions) for the angle (φ) and orbital angular momentum (J) of the plane rotator have been defined using quantum wave functions expressed in both representations, ψ_φ and $\psi'_{J/\hbar}$. It is shown that the "quantum" identification between the symplectic dual and the Fourier dual ($\times\hbar$) introduces constraints, and the integrality condition $J/\hbar \in \mathbb{Z}$ appears as necessary for consistency.

It is interesting to note that the Titius-Bode law for the planetary system suggests a constraint resembling a form of "entropy quantization", such as

$$\log_2\left(\frac{J}{J_G}\right)^3 = n \quad , \quad n = 0, 1, 2, \dots \quad (17)$$

where $J_G = McR_G$ with $R_G = 2\gamma_G M_o/c^2$ denoting the Schwarzschild radius of the central body (the Sun for the planets or Jupiter for its satellites, Appendix 2). At the atomic scale a constraint of this type is unlikely, but for the high circular Rydberg levels under active investigation [14, 15], a mixed state with a Poisson distribution (19) could be attributed to a stage of localization along the orbit.

Appendix 1: Rotational coherent states

Let us consider a particle of mass M , in uniform rotation with the angular frequency $\omega > 0$, around the Z -axis, on a circle of radius R in the XY plane. Thus, if $\mathbf{u} = (u_x, u_y)$ and $\mathbf{v} = (v_x, v_y)$ are the position and momentum vectors, then $\mathbf{u} = (R \cos \varphi, R \sin \varphi)$, $\mathbf{v} = (-P \sin \varphi, P \cos \varphi)$, with $P = M\omega R$ and $\varphi = \varphi_0 + \omega t$. It can be shown that a Gaussian distribution on $T^*\mathbb{R}^2$, centered on \mathbf{u} and \mathbf{v} , of the form

$$f_{u,v}(\mathbf{q}, \mathbf{p}) = \frac{1}{\pi^2 \hbar^2} e^{-(\mathbf{q}-\mathbf{u})^2/a - (\mathbf{p}-\mathbf{v})^2/b} \quad , \quad a = \hbar^2/b = \hbar/M\omega$$

can be obtained by a standard Wigner transform of the rotational coherent state ("symmetry breaking vacuum"),

$$|z\rangle = e^{z\hat{b}_u^\dagger - z^*\hat{b}_u} |0\rangle \quad , \quad (18)$$

where $z = \sqrt{J/\hbar} e^{-i\varphi}$, $J = M\omega R^2$, and $\hat{b}_u^\dagger = (\hat{b}_x^\dagger + i\hat{b}_y^\dagger)/\sqrt{2}$, with $\hat{b}_q = \sqrt{M\omega/2\hbar}(\hat{q} + i\hat{p}_q/M\omega)$, $\hat{b}_q|0\rangle = 0$, $q = x, y$. Moreover, the average of $f_{u,v}$ over $\varphi \in [-\pi, \pi]$ at constant J is the Wigner transform of the density operator

$$\hat{\rho}_w = \sum_{n=0}^{\infty} w_n |n\rangle \langle n| \quad , \quad |n\rangle = \frac{1}{\sqrt{n!}} (\hat{b}_u^\dagger)^n |0\rangle \quad , \quad (19)$$

expressed by a Poisson (non-thermal) distribution $w_n = e^{-J/\hbar} (J/\hbar)^n / n!$ of quantum entropy $S_q = -\sum_n w_n \ln w_n$ [10], over the eigenstates $|n\rangle$ of the angular momentum operator $\hat{L}_z = \hbar(\hat{b}_u^\dagger \hat{b}_u - \hat{b}_d^\dagger \hat{b}_d)$, $\hat{b}_d^\dagger = (\hat{b}_x^\dagger - i\hat{b}_y^\dagger)/\sqrt{2}$. Worth noting, a Gaussian distribution on \mathbb{R} with the same mean and variance as $\{w_n\}$ has the entropy $S_J = [1 + \ln(2\pi J/\hbar)]/2$.

Appendix 2: Comparison with astronomical data

The condition (17) and the third law of J. Kepler (May 15 1618) yield for the n 'th circular orbit the radius $r_n = R_G 2^{1+2n/3}$, $n = 0, 1, 2, \dots$. For the Sun $R_G = 2.95$ km, and with $n \geq 35$ the calculated values r_n are close to the astronomical data for all planets (e.g. the Earth's orbit has an average radius of $0.95r_{37}$), excepting Jupiter, whose average orbital radius is $0.96(r_{40} + r_{41})/2$.

In the case of Jupiter $R_G = 2.82$ m, and for $n = 39, 40, 41, 42$ (Table 1), we get values⁴ (r_n) close to the ones observed (r_{obs}) for its largest satellites: Io, Europa, Ganimede, Callisto, discovered by G. Galilei in 1610.

Apart from the scale factor R_G , a peculiar common trait for these systems is the weighted average \bar{n} of the orbital number,

$$\bar{n} = \frac{\sum_n n M_n}{\sum_n M_n}, \quad (20)$$

where M_n denotes the observed mass of the body assigned to the radius r_n . Thus, one obtains $\bar{n} = 41$ for the planetary system (considering $M_{40} = M_{41} = 0.5M_{Jupiter}$), and $\bar{n} = 40.7$ for the satellites of Table 1.

In a historical perspective, the Titius-Bode law (1766-72) was anticipated by Kepler's geometric model of the solar system, (in *Mysterium Cosmographicum*, Tübingen, 1596), where the intervals between the planetary orbits are not arbitrary, but determined by "perfect" solids. Less known is that in fact, as shown in [16], the standard quantization of angular momenta can be related to these solids.

Table 1. Comparison between the observed orbital radius (r_{obs}) of the Jupiter satellites and the calculated value (r_n).

Satellite/n	Io/39	Eu/40	Ga/41	Ca/42
r_{obs} (10^3 km)	421.6	670.8	1070	1882
r_n (10^3 km)	378.5	600.8	953.7	1514
r_{obs}/r_n	1.11	1.12	1.12	1.24

References

- [1] R. Abraham and J. E. Marsden, *Foundations of Mechanics*, Benjamin/Cummings, (1978), p. 396.
- [2] E. P. Wigner, *On the quantum correction for thermodynamic equilibrium*, Phys. Rev. **40** 749 (1932).
- [3] J. G. Kirkwood, *Quantum statistics of almost classical assemblies*, Phys. Rev. **44** 31 (1933).

⁴ $r_{35} = 59609$ km corresponds to the "internal" orbit close to the 1 bar surface of radius 71492 km.

- [4] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics*, Springer-Verlag, New York, (1990), p. 244.
- [5] H. A. Kastrup, *Wigner functions for the pair angle and orbital angular momentum*, Phys. Rev A **94** 062113 (2016).
- [6] J. Tosiek, R. Cordero and F. J. Turrubiates, *The Wentzel-Kramers-Brillouin approximation method applied to Wigner functions*, J. Math. Phys. **57** 062103 (2016).
- [7] M. Przanowski and J. Tosiek, *From the discrete Weyl-Wigner formalism for symmetric ordering to a number-phase Wigner function*, J. Math. Phys. **58** 102106 (2016).
- [8] P. Busch, J. Kiukas and R. F. Werner, *Sharp uncertainty relations for number and angle*, J. Math. Phys. **59** 042102 (2018).
- [9] M. Grigorescu, *Coherent distributions for the rigid rotator*, J. Math. Phys. **57** 062702 (2016).
- [10] M. Grigorescu, *Coherent distributions and quantization*, arXiv:1410.1338v4.
- [11] S. M. Barnett and D. T. Pegg, *Quantum theory of rotation angles*, Phys. Rev. A **41** 3427 (1990).
- [12] M. Grigorescu, *Classical probability waves*, Physica A **387** 6497-6504 (2008).
- [13] D. Pines, *Elementary Excitations in Solids; Lectures on Phonons, Electrons and Plasmons*, W. A. Benjamin, New York (1964), p. 101.
- [14] R. Blümel *et al.*, *Microwave excitations of Rydberg atoms in the presence of noise*, Phys. Rev. Lett. **62** 341 (1989).
- [15] A. Signoles *et al.*, *Coherent transfer between low angular momentum and circular Rydberg states*, Phys. Rev. Lett. **118** 253603 (2017).
- [16] G. S. Anagnostatos, *The geometry of the quantization of angular momenta (l,s,j) in fields of central symmetry*, Lett. Nuovo Cimento **28** 573 (1980).