SEMI-CLASSICAL GREEN FUNCTIONS

A.ANIKIN¹, S.DOBROKHOTOV¹, V.NAZAIKINSKII¹ ℰ M.ROULEUX²

¹ Ishlinski Institute for Problems of Mechanics Moscow Institute of Physics and Technology, Moscow, Russia; anikin83@inbox.ru; dobr@ipmnet.ru; nazay@ipmnet.ru

² Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France; rouleux@univ-tln.fr

Abstract: Let $H(x,p) \sim H_0(x,p) + hH_1(x,p) + \cdots$ be a semi-classical Hamiltonian on $T^*\mathbf{R}^n$, and $\Sigma_E = \{H_0(x,p) = E\}$ a non critical energy surface. Consider f_h a semi-classical distribution (the "source") microlocalized on a Lagrangian manifold Λ which intersects cleanly the flow-out Λ_+ of the Hamilton vector field X_{H_0} in Σ_E . Using Maslov canonical operator, we look for a semi-classical distribution u_h satisfying the limiting absorption principle and $H^w(x, hD_x)u_h = f_h$ (semi-classical Green function). In this report, we elaborate (still at an early stage) on some results announced in [AnDoNaRo1] and provide some examples, in particular from the theory of wave beams.

1. Introduction

Our general motivation is to solve equations like $(H - E)u_h = f_h$, with $f_h(x) = h^{-n}f(\frac{x}{h})$, $f \in L^2(\mathbf{R}^n)$, a "localized function" at $x_0 = 0$. In case $H = -h^2\Delta$, we apply h-Fourier transform

$$\mathcal{F}_h u_h(\xi) = \int e^{-ix\xi/h} u_h(x) \, dx$$

to get $\mathcal{F}_h u_h(\xi) = \frac{\mathcal{F}_1 f(\xi)}{\xi^2 - E}$. If E < 0, this is an elliptic equation, and u_h has the same form as f_h . If E > 0 however,

(1)
$$u_{+}(x;E,h) = (2\pi h)^{-n} \int e^{ix\xi/h} \frac{\mathcal{F}_{1}f(\xi)}{\xi^{2} - E - i0} d\xi$$

is defined through regularisation, so to satisfy the limiting absorption principle. Actually,

$$u_{+}(x; E, h) = E_{+}f_{h}, \quad E_{+} = \frac{i}{h} \int_{0}^{\infty} e^{-it(H-E)/h} dt$$

is the forward parametrix. When $f_h(x)$ is replaced by $\delta(x-x_0)$, u_+ is the Green function. Such a distribution is very singular since there are infinitely many classical trajectories linking the "source" x_0 to x in time t; so it is suitable to consider instead $\mathcal{F}_1 f$ with compact support.

This problem has of course received considerable attention [Ke], [Ba], [Ku]; our special purpose is to reformulate it in terms of Maslov canonical operator, and describe the solution globally, including unfolding of Lagrangian singularities; this is of special importance in the context of wave propagation.

More generally, let $f_h = f(\cdot; h)$ be a semi-classical Lagrangian distribution (or oscillatory integral) defined on the manifold M (for all practical purposes, we shall assume here $M = \mathbf{R}^n$), i.e. locally

$$f(x;h) = (2\pi h)^{-n/2} \int e^{i\varphi(x,\theta)/h} a(x,\theta;h) d\theta$$

where $\varphi(x,\theta)$ is a non-degenerate phase function in the sense of [Hö], and $a(x,\theta;h) = a_0(x,\theta) + ha_1(x,\theta) + \cdots$ an amplitude. With f we associate the critical set $C_{\varphi} = \{(x,\theta) \in M \times \mathbf{R}^N : \partial_{\theta}\varphi = 0\}$ and $\iota_{\varphi} : C_{\varphi} \to T^*M$ with image the Lagrangian submanifold $\Lambda_{\varphi} = \{(x,\partial_x\varphi) : (x,\theta) \in C_{\varphi}\}$. Choose local coordinates $\xi \in \mathbf{R}^d$ on Λ_{φ} and define the half-density in the local chart $(C_{\varphi}, \iota_{\varphi})$ by $\sqrt{d\mu_{\varphi}} = |\det \varphi''|^{-1/2}|d\xi|^{1/2}$. The (oscillating) principal symbol of f in Λ_{φ} is then defined (up to the factor $\sqrt{d\mu_{\varphi}}$) as

$$e^{i\phi(\xi)/h}A_0(\xi) = e^{i\phi(\xi)/h}e^{i\pi\operatorname{sgn}\varphi''/4}a_0(x(\xi),\theta(\xi))$$

where ϕ is a "reduced phase function". Conversely, assume $\iota:\Lambda\to T^*M$ is a smooth Lagrangian immersion, with a smooth positive half-density $\sqrt{d\mu}$, it can be parametrized locally by phase functions φ in canonical charts $\iota_{\varphi}:U_{\varphi}\to T^*M$. These phases can be chosen coherently, and define a class of "reduced phase functions" ϕ , parametrizing ι locally. This gives the fibre bundle of phases \mathcal{L}_h , including Maslov indices, equipped with transition functions. We are also given local smooth half-densities $|d\mu_{\varphi}|^{1/2}$ on Λ , defining the fibre bundle of half-densities $\Omega^{1/2}$, equipped with transition functions. The collection of these objects make a fibre bundle $\Omega^{1/2}\otimes\mathcal{L}_h$ over Λ . A section of $\Omega^{1/2}\otimes\mathcal{L}_h$ will be written as

$$f(x;h) = \left[K_{(\Lambda,\mu)}^h A\right](x;h)$$

where $K_{(\Lambda,\mu)}^h$ is called Maslov canonical operator. The "reduced phase function" and the "principal symbol" of f are defined invariantly. See [M], [Du], [Iv], [DoZh], [DNS] for details.

So our general aim is to construct, in term of Maslov canonical operators, a representation of E_+f . Here are some examples of f (expressed in a single chart):

(1) $\Lambda = \{(x, \partial_x \varphi) : x \in M\}$. WKB functions $f(x, h) = e^{i\varphi(x)/h} a(x, h)$ or in Fourier representation

(2)
$$f(x;h) = \frac{e^{i\pi n/4}}{(2\pi h)^{n/2}} \int e^{i(xp+S(p))/h} A(p;h) dp$$

(such an integral conveniently normalized including a phase factor, will be written $\int_{-\infty}^{\infty} (\cdots)$; here $\operatorname{sgn}(x \cdot p)'' = -n$).

- (2) Gaussian functions $f(x;h) = \frac{1}{h^n} \exp(-\frac{\omega^2 \cdot x^2}{2h})$ or more general (superposition of) coherent states.
 - (3) A conormal distribution with $\Lambda = T_N^* \mathbf{R}^n$, $N = \{x_n = 0\}$

(3)
$$f(x;h) = \int_{-\infty}^{\infty} e^{ix_n p_n/h} A(x', p_n) dp_n$$

(4) $\Lambda = \{x = X(\varphi, \psi) = \varphi\omega(\psi), p = P(\varphi, \psi) = \omega(\psi), \varphi \in \mathbf{R}\}, \ \omega \in \mathbf{S}^{n-1}, \ f \text{ identifies with a "Bessel beam", see Sect. 4.}$

2. Hypotheses and main result

Let $\Lambda_0 \to T^*M$, $\Lambda_1 \to T^*M$ be smooth embedded Lagrangian manifolds, Λ_1 with a smooth boundary $\partial \Lambda_1$ (isotropic manifold). Following [MelUh], we say that (Λ_0, Λ_1) is an *intersecting pair* iff $\Lambda_0 \cap \Lambda_1 = \partial \Lambda_1$ and the intersection is *clean*, i.e.

$$\forall \rho \in \partial \Lambda_1 \quad T_{\rho} \Lambda_0 \cap T_{\rho} \Lambda_1 = T_{\rho} \partial \Lambda_1$$

In particular, near each $\rho \in T^*M$, there is a canonical transform mapping Λ_0 to $T_0^*\mathbf{R}^n = \{x = 0\}$ and Λ_1 to

$$\Lambda_{+}^{0} = \{(x,\xi) : x = (0,x_n), \xi = (\xi',0), x_n \ge 0\}$$

i.e. the flow-out of $T_0^* \mathbf{R}^n$ by the Hamilton vector field $X_{\xi_n} = ((0, \dots, 0, 1), 0)$ of ξ_n , passing through $x = 0, \xi_0 = (\xi', 0)$.

Let now $H(x, p; h) = H_0(x, p) + hH_1(x, p) + \cdots$ be a symbol of class $S^0(m)$ and $H^w(x, hD_x; h)$ its Weyl quantization, as a bounded operator on $L^2(M)$. Assume

- (A.1) The energy surface $\Sigma_E = H_0^{-1}(E)$ is non critical.
- (A.2) The set $L = \Sigma_E \cap \Lambda$ is compact.
- (A.3) The Hamiltonian vector field X_{H_0} is transverse to Λ along L.

For $t \geq 0$, let $g_{H_0}^t = \exp t X_{H_0}$, $\Lambda_t = g_{H_0}^t(\Lambda)$, and $\Lambda_+ = \bigcup_{t \geq 0} g_{H_0}^t(L)$. The pair (Λ, Λ_+) define a Lagrangian intersection, and $L = \partial \Lambda_+$. We assume also

- (A.4) The Hamilton flow $g_{H_0}^t|_L$ extends for all t > 0.
- (A.5) $|\pi_x(g_{H_0}^t|_L)| \to \infty$ as $t \to \infty$ (non trapping condition).

Let ψ be a system of coordinates on L, completed (locally near L) to coordinates (ψ, τ) on Λ and to coordinates (ψ, t) on Λ_+ , so that the positive measure on Λ (square of half-density) takes the form $d\mu = |d\tau \wedge d\psi|$. This implies the measures $d\mu_t = (g_{H_0}^{t*})^{-1}(d\mu)$ on Λ_t and $d\mu_+ = |dt \wedge d\psi|$ on Λ_+ .

We shall consider a partition of unity subordinated to a covering of the pair (Λ, Λ_+) , with cut-off functions (of small enough support) $\widetilde{\chi}_0 \in C_0^{\infty}(\Lambda)$, that depends only on τ (the "radial variable" on Λ), with $\widetilde{\chi}_0(\tau) \equiv 1$ near $L = \{\tau = 0\}$, and $\chi_0 \in C_0^{\infty}(\Lambda_+)$, that depends only on t, $\chi_0(t) \equiv 1$ near t = 0. Let also $\chi_T \in C^{\infty}(\mathbf{R}^+)$, $\chi_T \equiv 1$ for $t \leq T/2$, $\chi_T \equiv 0$ for $t \geq T/2$ where T is so large that $x \notin \pi_x(g_{H_0}^t|_L)$ if $t \geq T$ (using the non-trapping condition (A.5)).

For simplicity, we state our main result [AnDoNaRo1] in the case $\Lambda = T_0^*M$, so that all solutions of Hamilton equations start from some (0,p), with $p \in \operatorname{supp} A$. Representing E_+ as a sum of Maslov canonical operators associated with the pair (Λ, Λ_+) in particular we shall retrieve the theorem on propagation of singularities for wave-front sets $\operatorname{WF}_h u \subset \operatorname{WF}_h(Hu) \cup \operatorname{Char} H$.

Theorem 1: Let $\Omega \subset M$ be a bounded domain. Under hypotheses (A.1)-(A.5) above, there is $u = u_h$, solution of $H^w(x, hD_x; h)u_h = f_h$, satisfying the limiting absorption principle, which can generically be cast in the following form

$$u(x,h) = \int_{0}^{*} \frac{(1-\tilde{\chi}_{0}(\tau(p))A(p))}{H(0,p)} e^{ipx/h} dp$$

$$+ \int_{0}^{\varepsilon_{0}} dt \int_{0}^{*} e^{i\Phi(t,x,\eta)/h} e^{-i\Theta(t,x,\eta)} |\det J(t,x,\eta)|^{-1/2} \chi_{T}(t)\chi_{0}(t)\rho(\tau(\eta))A(\eta) d\eta dt$$

$$+ K_{(\Lambda_{+},d\mu_{+})}^{h} [\chi_{T}(\cdot)(1-\chi_{0}(\cdot))e^{-i\Theta}A_{+}]$$

up to $\mathcal{O}(h)$ terms; the summands on the right hand side are boundary, transient and wave part, respectively. In Eq.(4), $\Phi(t, x; \eta)$ solves Hamilton-Jacobi equation (see Sect.3), and $\Theta(t, x, \eta) = \int_0^t H_1 \circ H_1 \circ H_2 \circ H_2 \circ H_3 \circ H_3$

 $\exp(sX_{H_0}(0,\eta))$ ds is the integral of the sub-principal 1-form, where we recall H_1 is the sub-principal symbol of H. Moreover $J(t,x,\eta)$ is a Jacobian computed from Hamilton-Jacobi equation.

The case when Λ is slightly "tilted" with respect to the vertical plane, i.e. has generating function $\phi(x,\xi) = x\xi + S(\xi)$ can be formulated in a very similar way (see Sect.3). For more general f's, the boundary part should be written as $\left[K_{(\Lambda,\mu)}^h \frac{(1-\widetilde{\chi}_0)A}{H}\right](x;h)$, and the transient part as the integral over t of a phase factor times $\left[K_{(\Lambda_t,\mu_t)}^h \chi_T(\cdot)\widetilde{\chi}_0(\cdot)\chi_0(\cdot)A_t\right](x;h)$, A_t being the solution of the transport equation along the projection of $\exp tX_{H_0}$ with initial data $A(\eta)$.

The boundary part is microlocalized on Λ , the wave part on Λ_+ . Let us make first a few comments:

- 1) (4) can be interpreted as "integrated" Van Vleck Formula [CdV], which expresses the semiclassical propagator $e^{-itH/h}$ acting on a localized function.
- 2) The wave part contributes generally at $x \in \Omega$ only if $t \mapsto \Phi(t, x, \eta)$ has a non-degenerate critical point for all $\eta \in \text{supp } f$: let hD_{x_n} be the "model" operator, and $f(x; h) = \int^* e^{ixp/h} A(p) dp$. Then [MelUh]

$$u_h(x) = \frac{i}{h} \int_0^\infty \chi_T(t) dt \int_0^* e^{i(x'\xi' + (x_n - t)\xi_n)/h} A(\xi) d\xi$$

verifies the limiting absorption principle and $hD_{x_n}u(x,h) = f(x,h) + \mathcal{O}(h^{\infty})$ for $x_n \leq T/2$; still u_h has no "wave part".

- 3) Formula (4) is also valid near a focal point $x \in \Omega$, or more generally when x is linked to x_0 through a trajectory containing several focal points. The wave-part simplifies outside the focal points to a WKB form, involving non trivial Maslov indices passing the first focal point.
- 4) The phase function $\Phi(t, x, \eta)$ can be also replaced by a Lagrangian action, in the spirit of [DNS]. This will be discussed in detail in a future work.

We illustrate Theorem 1 by computing u_h explicitely in the 2-D case for Helmholtz operator with constant coefficient as in (1), but f with compact support. Let f also be radially symmetric; its Fourier transform $g = \mathcal{F}_1 f$ is again of the form g(p) = g(|p|) = g(r) and extends holomorphically to \mathbb{C}^2 . For $E = k^2$, k > 0, we rewrite (1) as $u_h(x) = u(x) = u_0(x) + u_1(x)$ with

$$u_0(x) = \frac{k + i\varepsilon}{(2\pi h)^2} \int_0^{2\pi} d\theta \int_0^{\infty} \exp[i|x|r\cos\theta/h] \frac{g(r)}{r^2 - (k + i\varepsilon)^2} dr$$

$$u_1(x) = \frac{1}{(2\pi h)^2} \int_0^{2\pi} d\theta \int_0^{\infty} \exp[i|x|r\cos\theta/h] \frac{g(r)}{r + k + i\varepsilon} dr$$

To compute u_0 we use contour integrals. When $\theta \in]-\frac{\pi}{2},\frac{\pi}{2}[$, we shift the contour of integration to the positive imaginary axis and get by the residues formula

(5)
$$\int_{0}^{\infty} \exp[i|x|r\cos\theta/h] \frac{g(r)}{r^{2} - (k+i\varepsilon)^{2}} dr + \int_{0}^{\infty} \exp[-|x|r\cos\theta/h] \frac{g(ir)}{r^{2} + (k+i\varepsilon)^{2}} idr = 2i\pi \frac{g(k+i\varepsilon)}{2(k+i\varepsilon)} \exp[i|x|(k+i\varepsilon)\cos\theta/h]$$

while for $\theta \in]\frac{\pi}{2}, \frac{3\pi}{2}[,$

(6)
$$\int_0^\infty \exp[i|x|r\cos\theta/h] \frac{g(r)}{r^2 - (k+i\varepsilon)^2} dr - \int_0^\infty \exp[|x|r\cos\theta/h] \frac{g(-ir)}{r^2 + (k+i\varepsilon)^2} i dr = 0$$

Summing up (5) and (6), integrating over $\theta \in]0,2\pi[$ and letting $\varepsilon \to 0$, we obtain

$$u_0(x) = \frac{i\pi g(k)}{(2\pi h)^2} \int_{-\pi/2}^{\pi/2} \exp[i|x|k\cos\theta/h] d\theta + \int_0^\infty \frac{dr}{r^2 + k^2} \left[\int_{-\pi/2}^{\pi/2} g(ir) - \int_{\pi/2}^{3\pi/2} g(-ir) \right] \exp[-|x||\cos\theta/h] d\theta$$

Since g(ir) = g(-ir), the latter integral vanishes, so we end up with

$$u_0(x) = \frac{i\pi g(k)}{(2\pi h)^2} \int_{-\pi/2}^{\pi/2} \exp[i|x|k\cos\theta/h] \, d\theta$$

It is readily seen that

WF_h
$$u_0 \subset \{x = 0\} \cup \{(x, k \frac{x}{|x|}), x \neq 0\} = \Lambda \cup \Lambda_+$$

Consider now u_1 . We let $\varepsilon \to 0$ and set $\widetilde{g}(r) = \frac{g(r)}{r(r+k)}$. Since $\widetilde{g}(r)\sqrt{r} \in L^1(\mathbf{R}_+)$, we have $u_1(x) = H_0(\widetilde{g})(\frac{|x|}{h})$, where H_0 denotes Hankel transform of order 0.

Let $\chi \in C_0^{\infty}(\mathbf{R}^2)$ be radially symmetric, and equal to 1 near 0, since WF_h $f_h = \{x = 0\}$, we have

$$g = \mathcal{F}_h(\chi f_h) + \mathcal{O}(h^{\infty}) = (2\pi h)^{-2} \mathcal{F}_h(\chi) * g + \mathcal{O}(h^{\infty})$$

so in the expression for u_1 we may replace mod $\mathcal{O}(h^{\infty})$, $\widetilde{g}(r)$ by a constant times $\widehat{g}(r) = \frac{(\mathcal{F}_h(\chi)*g)(r)}{r(r+m)}$ (see [Bad] for 2-D convolution and Fourier transform in polar coordinates). To estimate WF_h u_1 , we compute again the Fourier transform of $(1-\widetilde{\chi})\widehat{g}$ where $\widetilde{\chi}$ is a cut-off equal to 1 near 0, and we find it is again $\mathcal{O}(h^{\infty})$ if $\chi \equiv 1$ on supp $\widetilde{\chi}$. This shows that WF_h $u_1 \subset \{x=0\}$. See also [MelUh], Prop. 2.3. Note that the decomposition $u_0 + u_1$ is directly related with the corresponding one in Theorem 1 as the sum of the boundary (u_1) , and the wave part (u_0) . We conjecture that the transient part in Theorem 1 can be removed (taking a limit supp $\chi_0 \to \emptyset$) when the Hamiltonian flow X_{H_0} enjoys some non-degeneracy properties (see Sect.3), as in the case of a geodesic flow.

3. Maslov canonical operators associated with (Λ, Λ_+)

Proof of Theorem 1 consists first in looking at the propagator $e^{-itH/h}$ acting on Lagrangian distribution f_h as above, i.e. the solution of the Cauchy problem $(hD_t+H^w(x,hD_x))v=0, v|_{t=0}=f_h$. Next step is to integrate with respect to t after introducing the partition of unity above.

In this report, we shall content to construct the phase functions by solving Hamilton-Jacobi equation. First we recall from [Hö], Thm 6.4.5 the following:

Theorem 2: Denote the variable in $T^*\mathbf{R}^d$ by $(y,\eta) = (y',y_d;\eta',\eta_d)$. Let \mathcal{H} be a real valued, smooth Hamiltonian near $(0,\eta)$ such that $\mathcal{H}(0,\eta) = 0$, $\partial_{\eta_d}\mathcal{H}(0,\eta) \neq 0$, and let ϕ be a real valued, smooth function on \mathbf{R}^{d-1} such that $\partial_{y'}\phi = \eta'$. Then there exists in a neighborhood of $0 \in \mathbf{R}^d$ a unique real valued solution $\Phi(y;\eta)$ of Hamilton-Jacobi equation $\mathcal{H}(y,\partial_y\Phi) = 0$ satisfying the boundary condition

$$\Phi(y', 0; \eta) = \phi(y'), \quad \frac{\partial \Phi}{\partial y}(0; \eta) = \eta$$

We consider here the case of a "maximally singular" chart U for Λ where $\Lambda = \Lambda_{\phi}$ has generating function $\phi(x,\xi) = S(\xi) + x\xi$. Even if Λ is a plane, the outgoing manifold Λ_+ may be very complicated far away from x_0 , but changing the canonical charts, we can proceed step by step.

Without loss of generality we can also assume here E=0, S(0)=0, $\partial_{\xi}S(0)=0$. Let η, τ such that $\tau + H(0, \eta) = 0$, by Theorem 2 (after slightly changing the notations), there exists $\Phi(x, t; \eta, \tau)$ such that

(7)
$$\frac{\partial \Phi}{\partial t} + H(x, \frac{\partial \Phi}{\partial x}) = 0$$

$$\Phi(x, 0; \eta, \tau) = S(\eta) + x\eta$$

$$\frac{\partial \Phi}{\partial x}(0, 0; \eta, \tau) = \eta, \quad \frac{\partial \Phi}{\partial t}(0, 0; \eta, \tau) = \tau$$

Moreover, when τ is small enough, the intersecting pair (Λ, Λ_+) for energy level H = 0 extends to a smooth family of intersecting pairs $(\Lambda, \Lambda_+(\tau))$ for energy levels $H = -\tau$

$$\Lambda_{+}(\tau) = \{(x,\xi) \in T^{*}M, \exists t \geq 0, \exists (y,\eta) \in \Lambda, (x,\xi) = \exp tX_{H}(y,\eta), \tau + H(y,\eta) = 0\}$$

intersecting along $L(\tau)$ given by $\eta_n = \widetilde{\eta}_n(\eta', \tau)$, with $\widetilde{\eta}_n(\xi_0', 0) = 0$.

So near $x = x_0$ we can assume, possibly after permuting the ξ -coordinates, that $H(0; \xi'_0, 0) = 0$, $\frac{\partial H}{\partial \xi_n}(0; \xi'_0, 0) \neq 0$, and for τ small enough, the equation $\tau + H(x, \xi) = 0$ is equivalent to $\xi_n = \widetilde{\xi}_n(x, \xi', \tau)$, with $\widetilde{\xi}_n(0, \xi'_0, 0) = 0$.

Now we want to set $\tau=0$ and therefore, given (x,η) , solve the equation $\frac{\partial \Phi}{\partial t}=0$ for $t\geq 0$. As we have seen, uniqueness of solutions doesn't always hold, as shows the "model case" hD_{x_n} . Namely, the phase function given by Hamilton-Jacobi theory is $\Phi(x,t;\eta)=\phi(x',x_n-t,\eta)$, so $\partial_t\Phi=0$ iff $\eta_n=0$, for all t. However the phase parametrizing Λ_+ is given for small t by Taylor expansion

$$\Phi(x,t;\eta) = x\eta + S(\eta) - tH(x,\eta) + \frac{t^2}{2}\partial_{\xi}H(x,\eta)\partial_xH(x,\eta) + \cdots$$

The following assumption in turn ensures existence of a finite number of solutions $t = t(x; \eta) > 0$ of $\frac{\partial \Phi}{\partial t} = 0$. Denote by $\operatorname{Exp}_{x_0}^t \eta = \pi_x(\exp tX_{H_0}(x_0, \eta))$ the projection of the bicharacteristic of H starting from (x_0, η) near (x_0, ξ_0) at time 0. Assume:

 (A'_0) $\exists \Omega_{\xi} \subset \mathbf{R}^n$ open (small) neighborhood of ξ_0 , such that if $x = \operatorname{Exp}_{x_0}^t \eta$ for some t > 0, and $\eta \in \Omega_{\xi}$, then the map $\xi \mapsto \operatorname{Exp}_{x_0}^t \xi$ is a local diffeomorphism near η ; in other terms, x_0 and x are not conjugated along any trajectory that links them together within time t, with initial momentum ξ .

The set of such x is an open set Ω_x .

Proposition 1: Under hypothesis (A'_0) , for all $(x, \eta) \in \Omega_x \times \Omega_\xi$ there is a finite number of $t_j = t_j(x, \eta) > 0$ $(1 \le j \le N)$ solutions of $\partial_t \Phi(x, t; \eta) = 0$ and t_j are non degenerate critical points.

Moreover (A'_0) is generically fulfilled before occurrence of the first focal point, as show the following examples.

Let H_0 be a geodesic flow, E > 0, f conormal to $N = \{x_0\}$; Ω_x is a small enough ball centered at x_0 , (geodesic ball), or Ω_x is a neighborhood of a minimal geodesic $\{y = \operatorname{Exp}_{x_0}^s \eta, 0 \le s \le t_0\}$ for η

in some small neighborhood of η_0 . This applies when f is as in (2) and supp A is localized near η_0 . The same holds with f as in (3) conormal to the hypersurface $N = \{x_n = 0\}$ (or more generally to a surface N of positive codimension in M). For instance when $N = \{x_n = 0\}$, (A'_0) holds with Ω_{ξ} an open (small) neighborhood of $\eta = (0, \dots, 1)$, and Ω_x a neighborhood of a minimal geodesic from N to some x_1 . This follows from a well-known property of minimal geodesics (see e.g. [HeSj], Proposition 6.3). In case of a Schrödinger operator with principal symbol $H_0(x,\xi) = \xi^2 + V(x) - E$ (where E is a scattering energy) we use Maupertuis-Jacobi principle to reduce again to a Riemannian metric d_E given by $ds = (E - V(x))^{1/2} |dx|$. So we arrive at the same conclusions, the metric d_E being conformal to the standard metric. We proved the following:

Proposition 2: Under the minimality assumptions above, (A'_0) holds. For $\eta \in \Omega_{\xi}$ and $x = Exp_{x_0}^t(\eta) \in \Omega_x$, $0 < t < t_0$, the Lagrangian manifold $\Lambda_t = \exp tX_H(\Lambda_{\phi})$ has same rank as Λ , and is of the form $\Lambda_{\Phi(t,\cdot)}$. In particular

$$\exp tX_{H_0}(\Lambda_\phi) = \{(x,\partial_x \Phi(t,x;\eta)): \partial_\eta \Phi(t,x;\eta) = 0\} \subset \{\exp tX_{H_0}(x,\eta), (x,\eta) \in T^*N^\perp\}$$

Moreover for all $(x, \eta) \in \Omega_x \times \Omega_\xi$ there is a unique $t(x, \eta) > 0$ solution of $\partial_t \Phi(x, t; \eta) = 0$ and $t(x, \eta)$ is a non degenerate critical point.

Extending the geodesic γ beyond the first focal point occuring in some Λ_t for some $t=t_0$. we only need another representation of the phase function. Using the mixed representation for Lagrangian manifolds, we know that for any $k=0,\cdots,n$, (k=0 corresponds to a maximally singular chart, k=n to a regular chart, they are mapped onto each other by Fourier transform), there exists a partition of variables $x=(x',x'')\in \mathbf{R}^k\times\mathbf{R}^{n-k}$ and $\xi=(\xi',\xi'')$, such that if $\widetilde{\pi}:\mathbf{R}^{2n}\to\mathbf{R}^n$, $(x,\xi)\mapsto (x',\xi'')$, then rank $d\widetilde{\pi}=n$. In such a chart, the generating function for Λ_t takes the form $\phi(x,\eta)=x''\eta''+S(x',\eta'')$. We can reformulate Hamilton-Jacobi equations as in (7) in these coordinates, which has again a unique solution for small $t-t_0$. Generically (i.e. under an assumption (A'_k) modeled after (A'_0) in the new (x,ξ) coordinates), one still obtains a non degenerate phase function, which contributes to the fibre bundle \mathcal{L}_h over Λ_+ together with Maslov index of γ . More explicit formulae will be given elswewhere. Of course, γ may contribute a finite number of times in the expression for the Green function at x, and a finite number of γ contribute to the wave part of the phase function.

4. Using eikonal coordinates.

The computations above can be simplified using special coordinates adapted to Λ , called *eikonal* coordinates.

Let $\iota: \Lambda \to T^*M$ be a smooth embedded Lagrangian manifold. The 1-form $p\,dx$ is closed on Λ , so locally exact, and $p\,dx = dS$ on any simply connected domain U. Such a S is called an eikonal (or action) and is defined up to a constant. Assume $dS \neq 0$ on Λ , then S can be chosen as a coordinate on U, which we complete by smooth functions $\psi \in \mathbf{R}^{n-1}$.

We use here eikonal coordinates to construct a phase function solving Hamilton-Jacobi equation in case of a positively homogeneous Hamiltonian of degree m with respect to p. To fix ideas, we take

n=2 (for simplicity) m=1 and

(8)
$$H(x,p) = \frac{|p|}{n(x)}$$

Example 1: $\Lambda = \{x = 0\}$ intersects the energy surface H = 1 along L. Let us compute the eikonal S on Λ_+ . Integrating Hamilton equations we have $x = X(t, \psi), p = P(t, \psi)$ where $\psi \in \mathbf{R}^2$, hence $dS = p \, dx|_{\Lambda_+} = \langle P(t, \psi), dX(t, \psi) \rangle$. Since dx = 0 on Λ , we have $S(0, \psi) = \text{Const.} = S_0$, and

(9)
$$S(t,\psi) = S(0,\psi) + \int_{(0,\psi)}^{(t,\psi)} p \, dx |_{\Lambda_+} = S_0 + \int_0^t \langle P(s,\psi), \dot{X}(s,\psi) \rangle \, ds$$

By Hamilton equations and Euler identity, we have on H=1.

$$\langle P(s,\psi), \dot{X}(s,\psi) \rangle = \langle P(s,\psi), \partial_p H(X,P) \rangle = mH(X,P) = m$$

and $S(t, \psi) = S_0 + mt$ is the action on Λ_+ . So

$$m dt = \langle P(t, \psi), dX(t, \psi) \rangle = \langle P(t, \psi), \dot{X}(t, \psi) \rangle dt + \langle P(t, \psi), X_{\psi}(t, \psi) \rangle d\psi$$

It follows that $\langle P(t, \psi), \dot{X}(t, \psi) \rangle = m$ and $\langle P(t, \psi), \partial_{\psi} X(t, \psi) \rangle = 0$. Now we complete the coordinate system ψ on Λ by a smooth function r such that L is given by r = 1, and set

$$\Phi(x, (t, \psi, r)) = mt + r\langle P(t, \psi), x - X(t, \psi) \rangle$$

where r can be interpreted as a Lagrange multiplier. Let us check that Φ satisfies Hamilton-Jacobi equation. We have

(10)
$$\partial_t \Phi = \dot{\Phi} = m + r \langle \dot{P}, x - X(t, \psi) \rangle - r \langle P, \dot{X} \rangle = m(1 - r) + r \langle \dot{P}(t, \psi), x - X(t, \psi) \rangle$$
$$\partial_r \Phi = \langle P(t, \psi), x - X(t, \psi) \rangle$$
$$\partial_\psi \Phi = r \langle \partial_\psi P(t, \psi), x - X(t, \psi) \rangle$$

Last 2 equations in $(\dot{\Phi}, \partial_r \Phi, \partial_\psi \Phi) = 0$ give an homogeneous linear system with determinant $\det(P, P_\psi)$. On x = 0 we get |p| = n(0) > 0, and in dimension n = 2, $p = n(0)^t(\cos\psi, \sin\psi)$. It follows that det $(P, P_\psi) = |n(0)|^2$ so for small t, we get $x - X(t, \psi) = 0$, so the phase is critical with respect to (ψ, r) for $x = X(t, \psi)$. Substituting into the last equation (10) we get $\dot{\Phi} = 0$ when r = 1. We complete the proof that Φ is a generating function for Λ_+ by checking $d\partial_t \phi, d\partial_\psi \Phi, d\partial_r \Phi$ are linearly independent on the set $x = X(t, \psi)$. This can be done by examining the variational system associated with Hamilton equations.

Moreover we can reduce this generating function by eliminating t by stationary phase. Again, of course, this holds only for small t > 0, before unfolding of Lagrangian singularities.

Example 2: We take H as in (8) with n(x) = n(|x|), n = 2 for simplicity and

$$\Lambda = \{ x = X(\varphi, \psi) = \varphi \omega(\psi), p = P(\varphi, \psi) = \omega(\psi), \ \varphi \in \mathbf{R}, \omega \in \mathbf{S}^{n-1} \}$$

 ψ being the usual angles parametrizing $\omega \in \mathbf{S}^{n-1}$. When n=2 this is the wave-front set of Bessel function $f_h(x;h) = J_0(\frac{|x|}{h})$; such functions arise in the wave beam theory (see [Ki], [DoMaNa] and references therein), so we call f_h a "Bessel beam".

Computing the action we find $p dx|_{\Lambda} = d\varphi$ so coordinate φ will play the role of x in he previous Example. We have $n(|x|) = n(\varphi)$, and $\varphi = \text{Const.}$ on $L = \Lambda \cap \{H = 1\}$. The argument above extends readily to this setting, in particular as in (9)

$$S(t,\varphi,\psi) = S(0,\varphi,\psi) + \int_{(0,\varphi,\psi)}^{(t,\varphi,\psi)} p \, dx|_{\Lambda_{+}} = S(0,\varphi,\psi) + \int_{(0,\varphi,\psi)}^{(t,\varphi,\psi)} P(s,\varphi,\psi) \, dX(s,\varphi,\psi) =$$

$$S_{0} + \int_{0}^{t} \langle P(s,\varphi,\psi), \dot{X}(s,\varphi,\psi) \rangle \, ds$$

and

$$\langle P(t,\varphi,\psi), \partial_{\psi} X(t,\varphi,\psi) \rangle = \langle P, \partial_{\varphi} X \rangle = 0$$

Now we can apply the results of [DNS], Sect.2.2 on eikonal coordinates for a Lagrangian manifold in a general position, with $X_{\widetilde{\psi}}(\widetilde{\varphi},\widetilde{\psi})$ of rank k. Here $\widetilde{\varphi},\widetilde{\psi}$ are coordinates in a local chart of Λ_+ in the extended phase space. More specifically, we take $\widetilde{\psi}=(t,\psi), \ \widetilde{\varphi}=\varphi$, try to make a change of variables $\varphi=\varphi(x,t,\psi)$, and seek for a generating function of Λ_+ in the form $\Phi(x,(t,\psi,r))$ as in the previous Example. Details will be given elsewhere.

5. More examples and perspectives.

The methods above apply in a number of situations as:

- (1) The water-wave Hamiltonian $H(x, hD_x; h)$ with $H_0(x, p) = |p| \tanh(|p|D(x)) E$, together with $\Lambda = \{x = x_0\}$ has been discussed in [DoNa], [AnDoNaRo2], in relationship with (11) or Helmhotz operator $H(x, hD_x) = -h^2\Delta (n(x))^2$. The localized function f can be a Gaussian (even in x) or a Gaussian times a linear function (odd in x), or can be of antenna type, i.e. its Fourier transform localized in a cone in p.
- (2) The kinetic part of Hamiltonian is of Lorenzian type (as $-p_0^2 + p_1^2$), and f a localized function (Gaussian) supported on $\Lambda = \{x = x_0\}$, so that the semi-classical Green function u_h is the linear response to f localized on Kelvin angle (or Mach cone).

It should also be possible to construct semi-classical Green functions in case the pair (Λ, Λ_+) is no longer intersecting cleanly, but glancing. We then need second-microlocalization, and introducing so called 2-phases, see e.g. [LaWi] in the standard (polyhomogeneous) calculus.

Acknowledgements: This work was supported by Grant PRC No. 1556 CNRS-RFBR 2017-2019 "Multi-dimensional semi-classical problems of Condensed Matter Physics and Quantum Mechanics", and RFBR Grant No. 17-51-150006.

References

[AnDoNaRo1] A.Anikin, S.Dobrokhotov, V.Nazaikinskii, M.Rouleux. Maslov's canonical operator on a pair of Lagrangian manifolds and asymptotic solutions of stationary equations with localized right-hand sides. Doklady Akad. Nauk, Vol. 76, No1, p.1-5, 2017.

[AnDoNaRo2] A.Anikin, S.Dobrokhotov, V.Nazaikinski, M.Rouleux. Asymptotics of Green function for the linear waves equations in a domain with a non-uniform bottom. Proceedings "Days of Diffraction 2017", Saint-Petersburg, IEEE p.18-23.

[Ba] V.M.Babich, On the short-wave asymptotic behavior of Green's function for the Helmholtz equation, Math. Sb. 65(107) (4), p.576-630, 1964.

[Bad] N.Baddour, Operational and convolution properties of two-dimensional Fourier transforms in polar coordinates, J. Opt. Soc. Am. A, Vol.26, p.1767-1777, 2009.

[CdV] Y.Colin de Verdière. Méthodes semi-classiques et théorie spectrale. https://www-fourier.ujf-grenoble.fr/ycolver/ All-Articles/93b.pdf

[DoNa] S.Dobrokhotov, V.Nazaikinskii. Punctured Lagrangian manifolds and asymptotic solutions of the linear water-wave equations with localized initial solutions. Math. Notes, 101, No.6, p.130-137, 2017.

[DoMaNa] S.Dobrokhotov, G. Makrakis, V.Nazaikinskii. Fourier Integrals and a new representation of Maslov's canonical operator near caustics. Amer. Math. Soc. Trans. Vol. 233, p.95-115, 2014.

[DNS] S.Dobrokhotov, V.Nazaikinskii, A.Shafarevich. New integral representations of Maslov canonical operators in singular charts. Izvestiya Math. 81(2), p.286-328, 2017.

[DoZh] S.Dobrokhotov, P.Zhevandrov. Asymptotic expansions and the Maslov canonical operator in the linear theory of water-waves I. Main constructions and equations for surface gravity waves. Russian J. Math. Phys. Vol.10 No.1, p.1-31, 2003.

[Du] J.J.Duistermaat. Oscillatory integrals, Lagrangian immersions and unfolding of singularities, Commun. Pure Applied Math., Vol.27, p.207-281, 1974.

[HeSj] B.Helffer, J.Sjöstrand. Multiple wells in the semi-classical limit I. Comm. Part. Diff. Eqn. 9(4) p.337-408, 1984.

[Hö] L.Hörmander. The Analysis of Linear Partial Differential Operators I, IV. Springer, 1985.

[Iv] V.Ivrii. Microlocal Analysis and Precise Spectral Asymptotics. Springer-Verlag, Berlin, 1998.

[Ke] J.B. Keller, Geometrical Theory of Diffraction, J. Opt. Soc. Am. 52, p.116-130, 1962.

[Ki] A.P.Kiselev. Localized light waves: Paraxial and exact solutions of the wave equation (a review). Optics and Spectroscopy, Vol.102 (4), p.603-622, 2007.

[Ku] V.Kucherenko. Quasi-classical asymptotics of a point source function for the stationary Schrödinger equation. Teor. i Mat. Fiz. Vol.1, No.3, p.384-406. Transl. Consultants Bureau, 1970.

[LaWi] P.Laubin, B.Willems. Distributions associated to a 2-microlocal pair of Lagrangian manifolds. Comm. Part. Diff. Eq. 19(9& 10), p.1581-1610, 1994.

[M] V.P.Maslov. Théorie des Perturbations et Méthodes Asymptotiques. Dunod, Paris, 1972.

[MelUh] R.Melrose and G.Uhlmann, Lagrangian intersection and the Cauchy problem, Comm. Pure Appl. Math. 32 (4), p.483-519, 1979.