

Self-interacting Brownian motion*

András Sütő
 Wigner Research Centre for Physics
 Hungarian Academy of Sciences
 P. O. B. 49, H-1525 Budapest, Hungary
 E-mail: suto@szfki.hu

August 2, 2018

Abstract

We prove a property of Brownian bridges whose certain time-equidistant sequences of points are pairwise coupled by an interaction. Roughly saying, if the total time span t of the bridge tends to infinity while the distance of its end points is fixed or increases slower than \sqrt{t} , the process asymptotically forgets this distance, just as in the absence of interaction. The conclusion remains valid if the bridge interacts in a similar way also with another set of trajectories. The main example for the interaction is the Coulomb potential.

1 Introduction

The problem we consider in this paper arises in Quantum Statistical Physics when a system of interacting particles is investigated by utilizing the Feynman-Kac formula [F1-2, G1-4]. It is in relation with Bose-Einstein condensation that we do not discuss here. Let $P_{\mathbf{0}\mathbf{x}}^\beta(d\boldsymbol{\omega})$ denote the Wiener measure for Brownian paths $\boldsymbol{\omega} : [0, \beta] \rightarrow \mathbb{R}^\nu$ with $\boldsymbol{\omega}(0) = \mathbf{0}$, $\boldsymbol{\omega}(\beta) = \mathbf{x}$. It is generated by the Gaussian functions

$$\psi_t(\mathbf{x}) = \lambda_t^{-\nu} e^{-\pi \mathbf{x}^2 / \lambda_t^2}, \quad 0 \leq t \leq \beta. \quad (1.1)$$

In physics λ_t is the thermal de Broglie wave length at inverse temperature t , $\lambda_t = \sqrt{2\pi\hbar^2 t/m}$, where m is the mass of the particle. In particular,

$$\int P_{\mathbf{0}\mathbf{x}}^\beta(d\boldsymbol{\omega}) = \lambda_\beta^{-\nu} e^{-\pi \mathbf{x}^2 / \lambda_\beta^2}. \quad (1.2)$$

Our starting observation is that

$$\lim_{\mathbf{x}^2 / \lambda_\beta^2 \rightarrow 0} \frac{\int P_{\mathbf{0}\mathbf{x}}^\beta(d\boldsymbol{\omega})}{\int P_{\mathbf{0}\mathbf{0}}^\beta(d\boldsymbol{\omega})} = 1, \quad (1.3)$$

and we wish to prove a similar result in the case of interaction. For this we take a real function $u(\mathbf{x})$ on \mathbb{R}^ν that depends only on $|\mathbf{x}|$; the interaction between points \mathbf{x} and \mathbf{y} is $u(\mathbf{x} - \mathbf{y})$. The time span of the bridge is $n\beta$, where $\beta > 0$ and n is a positive integer. Given a trajectory $\boldsymbol{\omega} : [0, n\beta] \rightarrow \mathbb{R}^\nu$, its self-interaction is defined as

$$U(\boldsymbol{\omega}) = \frac{1}{\beta} \sum_{0 \leq k < l \leq n-1} \int_0^\beta u(\boldsymbol{\omega}(l\beta + t) - \boldsymbol{\omega}(k\beta + t)) dt, \quad (1.4)$$

*Work supported by OTKA Grant No. K109577.

and our interest is to give a lower bound on

$$\int P_{\mathbf{0}\mathbf{x}}^\beta(d\boldsymbol{\omega})e^{-\beta U(\boldsymbol{\omega})} \Big/ \int P_{\mathbf{0}\mathbf{0}}^\beta(d\boldsymbol{\omega})e^{-\beta U(\boldsymbol{\omega})}. \quad (1.5)$$

As to the physical background, let

$$H_n = -\frac{\hbar^2}{2m} \sum_{i=1}^n \Delta_{\mathbf{x}_i} + \sum_{1 \leq i < j \leq n} u(\mathbf{x}_j - \mathbf{x}_i), \quad (1.6)$$

the energy operator of n interacting particles of mass m . For suitable choices of u , $e^{-\beta H_n}$ has an integral kernel, and by the Feynman-Kac formula we have

$$\int e^{-\beta H_n}(\mathbf{0}, \mathbf{x}_2, \dots, \mathbf{x}_n; \mathbf{x}_2, \dots, \mathbf{x}_n, \mathbf{x}) d\mathbf{x}_2 \dots d\mathbf{x}_n = \int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\boldsymbol{\omega})e^{-\beta U(\boldsymbol{\omega})}. \quad (1.7)$$

In addition, we shall consider also the problem when, apart from a Brownian bridge $\boldsymbol{\omega}_0$ of time span $n\beta$ and self-interaction U_1 generated by a pair potential u_1 , there is also a set of M other trajectories $\boldsymbol{\omega}_i : [0, \beta] \rightarrow \mathbb{R}^\nu$

$$\boldsymbol{\omega}^M = (\boldsymbol{\omega}_1, \dots, \boldsymbol{\omega}_M), \quad \boldsymbol{\omega}_i(0) = \mathbf{x}_i, \quad \boldsymbol{\omega}_i(\beta) = \mathbf{y}_i. \quad (1.8)$$

These may interact among themselves; their potential energy is then

$$U_2(\boldsymbol{\omega}^M) = \frac{1}{\beta} \sum_{1 \leq i < j \leq M} \int_0^\beta u_2(\boldsymbol{\omega}_i(t) - \boldsymbol{\omega}_j(t)) dt, \quad (1.9)$$

with a pair potential u_2 . The inclusion of this interaction is optional; what is important is an interaction between $\boldsymbol{\omega}_0$ and $\boldsymbol{\omega}^M$,

$$V(\boldsymbol{\omega}_0, \boldsymbol{\omega}^M) = \frac{1}{\beta} \sum_{k=0}^{n-1} \sum_{i=1}^M \int_0^\beta u_3(\boldsymbol{\omega}_0(k\beta + t) - \boldsymbol{\omega}_i(t)) dt, \quad (1.10)$$

acting as an external field for $\boldsymbol{\omega}_0$. Both $u_2(\mathbf{x})$ and $u_3(\mathbf{x})$ depend only on $|\mathbf{x}|$. The underlying energy operator in this case is that of n particles of mass m_1 with pair interaction u_1 and M particles of mass m_2 with pair interaction u_2 , the two sets pairwise interacting via u_3 :

$$H_{n,M} = -\frac{\hbar^2}{2m_1} \sum_{i=1}^n \Delta_{\mathbf{x}_i} + \sum_{1 \leq i < j \leq n} u_1(\mathbf{x}_j - \mathbf{x}_i) + \sum_{i=1}^n \sum_{j=1}^M u_3(\mathbf{x}_i - \mathbf{y}_j) - \frac{\hbar^2}{2m_2} \sum_{i=1}^M \Delta_{\mathbf{y}_i} + \sum_{1 \leq i < j \leq M} u_2(\mathbf{y}_j - \mathbf{y}_i). \quad (1.11)$$

Some special cases of the external field will be presented in Section 3.

2 Isolated Brownian motion

Here we treat a Brownian bridge of time span $n\beta$ under self-interaction alone.

Theorem 2.1 *Let u be twice differentiable outside $\mathbf{0}$. If $\Delta u \leq 0$, then*

$$\lim_{n \rightarrow \infty, \mathbf{x}^2/n \rightarrow 0} \frac{\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\boldsymbol{\omega})e^{-\beta U(\boldsymbol{\omega})}}{\int P_{\mathbf{0}\mathbf{0}}^{n\beta}(d\boldsymbol{\omega})e^{-\beta U(\boldsymbol{\omega})}} \geq 1. \quad (2.1)$$

Proof. We start with the identity

$$\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\boldsymbol{\omega})\phi(\boldsymbol{\omega}) = \exp\left\{-\frac{\pi\mathbf{x}^2}{n\lambda_\beta^2}\right\} \int P_{\mathbf{0}\mathbf{0}}^{n\beta}(d\boldsymbol{\omega})\phi(\tilde{\boldsymbol{\omega}}) \quad (2.2)$$

where

$$\tilde{\omega}(t) = \omega(t) + \frac{t}{n\beta}\mathbf{x}, \quad (2.3)$$

see [G1]. Applying this to $\phi(\omega) = \exp -\beta U(\omega)$ and using the notation

$$E_{\omega}(\mathbf{x}) = \sum_{0 \leq k < l \leq n-1} \int_0^{\beta} u \left(\omega(l\beta + t) - \omega(k\beta + t) + \frac{l-k}{n}\mathbf{x} \right) dt \quad (2.4)$$

we obtain

$$\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(\mathrm{d}\omega) e^{-\beta U(\omega)} = \exp \left\{ -\frac{\pi \mathbf{x}^2}{n\lambda_{\beta}^2} \right\} \int P_{\mathbf{0}\mathbf{0}}^{n\beta}(\mathrm{d}\omega) e^{-E_{\omega}(\mathbf{x})}. \quad (2.5)$$

Now

$$\lim_{n \rightarrow \infty, \mathbf{x}^2/n \rightarrow 0} \exp \left\{ -\frac{\pi \mathbf{x}^2}{n\lambda_{\beta}^2} \right\} = 1, \quad (2.6)$$

so for

$$I^{n\beta}(\mathbf{x}) := \int P_{\mathbf{0}\mathbf{0}}^{n\beta}(\mathrm{d}\omega) e^{-E_{\omega}(\mathbf{x})} \quad (2.7)$$

we must prove that

$$\lim_{n \rightarrow \infty, \mathbf{x}^2/n \rightarrow 0} \frac{I^{n\beta}(\mathbf{x})}{I^{n\beta}(\mathbf{0})} \geq 1. \quad (2.8)$$

Actually, the stronger result

$$\frac{I^{n\beta}(\mathbf{x})}{I^{n\beta}(\mathbf{0})} \geq 1, \quad (2.9)$$

without taking the limit, also holds. Note that $I^{n\beta}(\mathbf{x})$ is a spherical function, thus it has an extremum at $\mathbf{0}$. We show that $I^{n\beta}(\mathbf{0})$ is a global minimum by proving that $\Delta I^{n\beta}(\mathbf{x}) \geq 0$ everywhere. Indeed, this implies that $I^{n\beta}$ is a convex even function of \mathbf{x} along every axis passing through $\mathbf{0}$. Now

$$\nabla I^{n\beta}(\mathbf{x}) = - \int P_{\mathbf{0}\mathbf{0}}^{n\beta}(\mathrm{d}\omega) e^{-E_{\omega}(\mathbf{x})} \nabla E_{\omega}(\mathbf{x}), \quad (2.10)$$

$$\Delta I^{n\beta}(\mathbf{x}) = \int P_{\mathbf{0}\mathbf{0}}^{n\beta}(\mathrm{d}\omega) e^{-E_{\omega}(\mathbf{x})} \left[|\nabla E_{\omega}(\mathbf{x})|^2 - \Delta E_{\omega}(\mathbf{x}) \right]. \quad (2.11)$$

So $\Delta I^{n\beta}(\mathbf{x}) \geq 0$ if $\Delta E_{\omega}(\mathbf{x}) \leq 0$, and the result follows from

$$\begin{aligned} \Delta E_{\omega}(\mathbf{x}) &= \sum_{k < l} \int_0^{\beta} \Delta_{\mathbf{x}} u \left(\omega(l\beta + t) - \omega(k\beta + t) + \frac{l-k}{n}\mathbf{x} \right) dt \\ &= \sum_{k < l} \left(\frac{l-k}{n} \right)^2 \int_0^{\beta} \Delta u \left(\omega(l\beta + t) - \omega(k\beta + t) + \frac{l-k}{n}\mathbf{x} \right) dt. \end{aligned} \quad (2.12)$$

The general solution of $\Delta u \leq 0$ is easily obtained. Define $f : [0, \infty) \rightarrow \mathbb{R} \cup \{\infty\}$ by $u(\mathbf{y}) = f(\mathbf{y}^2)$. Because

$$\Delta u(\mathbf{y}) = 2 \left[\nu f'(\mathbf{y}^2) + 2\mathbf{y}^2 f''(\mathbf{y}^2) \right], \quad (2.13)$$

one must find those f satisfying

$$\nu f'(s) + 2s f''(s) \leq 0 \quad \text{for any } s > 0. \quad (2.14)$$

Let $g(s)$ be a nonnegative function, and consider the differential equation

$$\nu f'(s) + 2s f''(s) = -g(s). \quad (2.15)$$

Choose some $a, b > 0$, then for the boundary conditions

$$f(a) = c_1, \quad f'(b) = c_2 b^{-\nu/2} \quad (2.16)$$

the solution of Eq. (2.15) is

$$f(s) = c_1 - \frac{1}{2} \int_a^s dx x^{-\nu/2} \int_b^x dt g(t) t^{\nu/2-1} + c_2 \begin{cases} \frac{s^{1-\nu/2} - a^{1-\nu/2}}{1-\nu/2}, & \nu \neq 2 \\ (\ln s - \ln a), & \nu = 2. \end{cases} \quad \square \quad (2.17)$$

Examples. For $g = 0$ we can obtain the ν -dimensional Coulomb potential of both signs. In general, $f(s) = s^{-\alpha}$ is a solution of Eq. (2.15) for $g(s) = \alpha(\nu - 2 - 2\alpha)s^{-\alpha-1}$ which is nonnegative if $\alpha \leq \nu/2 - 1$. Another interesting solution is obtained in dimensions $\nu \geq 3$ if

$$a = b = +\infty, \quad c_1 = c_2 = 0, \quad g(s) = s^{-\alpha}, \quad \alpha > \nu/2 > 1. \quad (2.18)$$

It reads

$$f(s) = \frac{-s^{-\alpha+1}}{(\alpha-1)(2\alpha-\nu)} \quad \text{or} \quad u(\mathbf{x}) = \frac{-|\mathbf{x}|^{-2\alpha+2}}{(\alpha-1)(2\alpha-\nu)}. \quad (2.19)$$

In three dimensions with $\alpha = 4$ this is the form of the induced dipole-dipole interaction which appears in the Lennard-Jones potential.

3 Self-interacting Brownian bridge in an external field

The external field is created by a set of other trajectories, and is given by Eq. (1.10). Let

$$\begin{aligned} E_{\omega_0}(\mathbf{x}) &= \sum_{0 \leq k < l \leq n-1} \int_0^\beta u_1 \left(\omega_0(l\beta + t) - \omega_0(k\beta + t) + \frac{l-k}{n} \mathbf{x} \right) dt, \\ E_{\omega_0, \omega^M}(\mathbf{x}) &= \sum_{k=0}^{n-1} \sum_{i=1}^M \int_0^\beta u_3 \left(\omega_0(k\beta + t) - \omega_i(t) + \frac{k\beta + t}{n\beta} \mathbf{x} \right) dt, \end{aligned} \quad (3.1)$$

and

$$I_{\omega^M}^{n\beta}(\mathbf{x}) = \int P_{\mathbf{0}\mathbf{0}}^{n\beta}(d\omega_0) e^{-E_{\omega_0}(\mathbf{x}) - E_{\omega_0, \omega^M}(\mathbf{x}) - \beta U_2(\omega^M)}. \quad (3.2)$$

As before,

$$\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\omega_0) e^{-\beta[U_1(\omega_0) + V(\omega_0, \omega^M) + U_2(\omega^M)]} = e^{-\pi \mathbf{x}^2 / \lambda_{n\beta}^2} I_{\omega^M}^{n\beta}(\mathbf{x}), \quad (3.3)$$

where $\lambda_{n\beta} = \sqrt{2\pi\hbar^2 n\beta / m_1}$.

Proposition 3.1 *If u_1 and u_3 are twice differentiable outside the origin and $\Delta u_1 \leq 0$, $\Delta u_3 \leq 0$, then $\Delta I_{\omega^M}^{n\beta} \geq 0$.*

Proof. Again, a direct computation gives

$$\begin{aligned} \Delta I_{\omega^M}^{n\beta}(\mathbf{x}) &= \int P_{\mathbf{0}\mathbf{0}}^{n\beta}(d\omega_0) e^{-E_{\omega_0}(\mathbf{x}) - E_{\omega_0, \omega^M}(\mathbf{x}) - \beta U_2(\omega^M)} \\ &\quad \times \left[|\nabla E_{\omega_0}(\mathbf{x}) + \nabla E_{\omega_0, \omega^M}(\mathbf{x})|^2 - \Delta E_{\omega_0}(\mathbf{x}) - \Delta E_{\omega_0, \omega^M}(\mathbf{x}) \right]. \end{aligned} \quad (3.4)$$

Recalling the identity (2.12) and

$$\Delta E_{\omega_0, \omega^M}(\mathbf{x}) = \sum_{k=0}^{n-1} \sum_{i=1}^M \int_0^\beta \left(\frac{k\beta + t}{n\beta} \right)^2 \Delta u_3 \left(\omega_0(k\beta + t) - \omega_i(t) + \frac{k\beta + t}{n\beta} \mathbf{x} \right) dt \quad (3.5)$$

the result follows. \square

$I_{\omega^M}^{n\beta}(\mathbf{x})$ is not a spherical function, so it may not be minimal at $\mathbf{x} = \mathbf{0}$. If, for example,

$$\text{dist}(\mathbf{x}, \{\mathbf{x}_i\}_{i=1}^M) \ll \text{dist}(\mathbf{0}, \{\mathbf{x}_i\}_{i=1}^M) \quad (3.6)$$

then possibly

$$E_{\omega_0, \omega^M}(\mathbf{x}) \gg E_{\omega_0, \omega^M}(\mathbf{0}) \quad (3.7)$$

and therefore $I_{\omega^M}^{n\beta}(\mathbf{x}) < I_{\omega^M}^{n\beta}(\mathbf{0})$.

Let $\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{x})$ denote a spherical symmetrization of $I_{\omega^M}^{n\beta}(\mathbf{x})$, obtained by integration over a set of ω^M . Examples will be given below. As a function of \mathbf{x} , $\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{x})$ depends only on $|\mathbf{x}|$, and may preserve some dependence on ω^M . From Eq. (3.3),

$$\mathcal{S}\left(\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\omega_0)e^{-\beta[U_1(\omega_0)+V(\omega_0, \omega^M)+U_2(\omega^M)]}\right)(\mathbf{x}) = e^{-\pi\mathbf{x}^2/\lambda_{n\beta}^2}\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{x}). \quad (3.8)$$

One can differentiate inside the integral over ω^M , so

$$\Delta\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{x}) = \mathcal{S}(\Delta I_{\omega^M}^{n\beta})(\mathbf{x}). \quad (3.9)$$

Theorem 3.1 For u_1 and u_3 as in Proposition 3.1,

$$\lim_{n \rightarrow \infty, \mathbf{x}^2/n \rightarrow 0} \frac{\mathcal{S}\left(\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\omega_0)e^{-\beta[U_1(\omega_0)+V(\omega_0, \omega^M)+U_2(\omega^M)]}\right)(\mathbf{x})}{\mathcal{S}\left(\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\omega_0)e^{-\beta[U_1(\omega_0)+V(\omega_0, \omega^M)+U_2(\omega^M)]}\right)(\mathbf{0})} \geq 1. \quad (3.10)$$

Proof. $\Delta I_{\omega^M}^{n\beta} \geq 0$ implies $\Delta\mathcal{S}(I_{\omega^M}^{n\beta}) \geq 0$ and, because $\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{x})$ is spherical,

$$\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{x}) \geq \mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{0}).$$

Then,

$$\lim_{n \rightarrow \infty, \mathbf{x}^2/n \rightarrow 0} \frac{\mathcal{S}\left(\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\omega_0)e^{-\beta[U_1(\omega_0)+V(\omega_0, \omega^M)+U_2(\omega^M)]}\right)(\mathbf{x})}{\mathcal{S}\left(\int P_{\mathbf{0}\mathbf{x}}^{n\beta}(d\omega_0)e^{-\beta[U_1(\omega_0)+V(\omega_0, \omega^M)+U_2(\omega^M)]}\right)(\mathbf{0})} = \lim_{n \rightarrow \infty, \mathbf{x}^2/n \rightarrow 0} \frac{\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{x})}{\mathcal{S}(I_{\omega^M}^{n\beta})(\mathbf{0})} \geq 1. \quad (3.11)$$

\square

Examples.

(i) Fix some $L > 0$ and define

$$\mathcal{S}(I^{n\beta})(\mathbf{x}) = \sum_{\pi \in S_M} \prod_{i=1}^M \int_{|\mathbf{x}_i| < L} d\mathbf{x}_i \int P_{\mathbf{x}_i \mathbf{x}_{\pi(i)}}^{\beta}(d\omega_i) I_{\omega^M}^{n\beta}(\mathbf{x}) \quad (3.12)$$

where S_M is the set of permutations of $\{1, \dots, M\}$. This is the typical choice when the particles associated with ω^M are bosons. The physically interesting case is $n \propto M \propto L^\nu$ and $n \rightarrow \infty$ replaced by $L \rightarrow \infty$. If $n \propto L^\nu$, for $\mathbf{x}^2/n \rightarrow 0$ to hold one must have $|\mathbf{x}| = o(L^{\nu/2})$. In $\nu \geq 3$ dimensions all \mathbf{x} of length $|\mathbf{x}| = O(L)$ satisfy this condition. When $n = M$, $u_1 = u_2 = -u_3 = u$ with $\Delta u = 0$, the system is a neutral two-component plasma.

(ii) The second set of particles can be treated classically. This is the $m_2 = \infty$ limit in which both the kinetic and the potential energy of these particles (the last two terms of the Hamiltonian (1.11)) can be dropped. Now $\omega_i(t) \equiv \mathbf{x}_i$, $\omega^M = \mathbf{x}^M = (\mathbf{x}_1, \dots, \mathbf{x}_M)$, and the interaction between the two sets becomes

$$V(\omega_0, \mathbf{x}^M) = \frac{1}{\beta} \sum_{k=0}^{n-1} \sum_{i=1}^M \int_0^\beta u_3(\omega_0(k\beta + t) - \mathbf{x}_i) dt. \quad (3.13)$$

Spherical symmetrization can be done by keeping all $|\mathbf{x}_i - \mathbf{x}_j|$ unchanged. Let g denote a general element of the rotation group $SO(\nu)$, and let μ be the Haar measure on $SO(\nu)$. Define

$$\mathcal{S}\left(I_{(\mathbf{x}_1, \dots, \mathbf{x}_M)}^{n, \beta}\right)(\mathbf{x}) = \int I_{(g\mathbf{x}_1, \dots, g\mathbf{x}_M)}^{n, \beta}(\mathbf{x}) \mu(dg). \quad (3.14)$$

Because u_1 and u_3 are spherical functions and the measure $P_{00}^{n, \beta}$ is rotation invariant, one actually has

$$\mathcal{S}\left(I_{(\mathbf{x}_1, \dots, \mathbf{x}_M)}^{n, \beta}\right)(\mathbf{x}) = \int I_{(\mathbf{x}_1, \dots, \mathbf{x}_M)}^{n, \beta}(g\mathbf{x}) \mu(dg). \quad (3.15)$$

When $n = M$, $u_1 = -u_3 = u$ with $\Delta u = 0$, we have a neutral system of light charged particles moving in the field of immobile heavy ions.

References

- [Fe1] Feynman R. P.: *Space-time approach to non-relativistic quantum mechanics*. Rev. Mod. Phys. **20**, 367-387 (1948).
- [Fe2] Feynman R. P.: *Atomic theory of the λ transition in helium*. Phys. Rev. **91**, 1291-1301 (1953).
- [G1] Ginibre J.: *Some applications of functional integration in Statistical Mechanics*. In: *Statistical Mechanics and Quantum Field Theory*, eds. C. De Witt and R. Stora, Gordon and Breach (New York 1971).
- [G2] Ginibre J.: *Reduced density matrices of quantum gases. I. Limit of infinite volume*. J. Math. Phys. **6**, 238-251 (1965).
- [G3] Ginibre J.: *Reduced density matrices of quantum gases. II. Cluster property*. J. Math. Phys. **6**, 252-262 (1965).
- [G4] Ginibre J.: *Reduced density matrices of quantum gases. III. Hard-core potentials*. J. Math. Phys. **6**, 1432-1446 (1965).