

ON THE SECOND BOUNDARY VALUE PROBLEM FOR LAGRANGIAN MEAN CURVATURE EQUATION

Chong Wang ¹ · Rongli Huang ² · Jiguang Bao ³

ABSTRACT. Considering the second boundary value problem of the Lagrangian mean curvature equation, we obtain the existence and uniqueness of the smooth uniformly convex solution, which generalizes the Brendle-Warren's theorem about minimal Lagrangian diffeomorphism in Euclidean metric space.

Mathematics Subject Classification 2000: 35J25 · 35J60 · 53A10

1. INTRODUCTION

The main aim of this article is to study the existence and uniqueness of the smooth uniformly convex solution for the second boundary value problem of the Lagrangian mean curvature equation

$$(1.1) \quad \begin{cases} F_\tau(\lambda(D^2u)) = \kappa \cdot x + c, & x \in \Omega, \\ Du(\Omega) = \tilde{\Omega}, \end{cases}$$

where Ω and $\tilde{\Omega}$ are two uniformly convex bounded domains with smooth boundary in \mathbb{R}^n , $\kappa \in \mathbb{R}^n$ is a constant vector, $\lambda(D^2u) = (\lambda_1, \dots, \lambda_n)$ are the eigenvalues of

The second author is supported by National Natural Science Foundation of China (No. 11771103 and 11871102) and Guangxi Natural Science Foundation (2017GXNSFFA198017). The third author is supported in part by the National Natural Science Foundation of China (11631002 and 11871102).

¹Fengtai School of the High School Affiliated to Renmin University of China, Beijing 100074, China. wch0229@mail.bnu.edu.cn

²The corresponding author. School of Mathematics and Statistics, Guangxi Normal University, Guangxi 541004, China. ronglihuangmath@gxnu.edu.cn

³School of Mathematical Sciences, Beijing Normal University, Laboratory of Mathematics and Complex Systems, Ministry of Education, Beijing 100875, China. jgbao@bnu.edu.cn

Hessian matrix D^2u , c is a constant to be determined and

$$(1.2) \quad F_\tau(\lambda) := \begin{cases} \frac{1}{n} \sum_{i=1}^n \ln \lambda_i, & \tau = 0, \\ \frac{\sqrt{a^2+1}}{2b} \sum_{i=1}^n \ln \frac{\lambda_i + a - b}{\lambda_i + a + b}, & 0 < \tau < \frac{\pi}{4}, \\ -\sqrt{2} \sum_{i=1}^n \frac{1}{1 + \lambda_i}, & \tau = \frac{\pi}{4}, \\ \frac{\sqrt{a^2+1}}{b} \sum_{i=1}^n \arctan \frac{\lambda_i + a - b}{\lambda_i + a + b}, & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ \sum_{i=1}^n \arctan \lambda_i, & \tau = \frac{\pi}{2}, \end{cases}$$

where $a = \cot \tau$, $b = \sqrt{|\cot^2 \tau - 1|}$.

Let

$$g_\tau = \sin \tau \delta_0 + \cos \tau g_0, \quad \tau \in \left[0, \frac{\pi}{2}\right]$$

be the linear combined metric of the standard Euclidean metric

$$\delta_0 = \sum_{i=1}^n dx_i \otimes dx_i + \sum_{j=1}^n dy_j \otimes dy_j$$

and the pseudo-Euclidean metric

$$g_0 = \frac{1}{2} \sum_{i=1}^n dx_i \otimes dy_i + \frac{1}{2} \sum_{j=1}^n dy_j \otimes dx_j$$

in $\mathbb{R}^n \times \mathbb{R}^n$.

In 2010, Warren [1] firstly obtained that if $(x, Du(x))$ is a minimal Lagrangian graph in $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$, then u satisfies

$$(1.3) \quad F_\tau(\lambda(D^2u)) = c,$$

which is a special case of (1.1) when $\kappa \equiv 0$.

If $\tau = 0$, (1.3) is the famous Monge-Ampère equation

$$\det D^2u = e^{2c},$$

which general form is

$$(1.4) \quad \det D^2u = f(x, u, Du).$$

If $\tau = \frac{\pi}{2}$, (1.3) becomes the special Lagrangian equation

$$(1.5) \quad \sum_{i=1}^n \arctan \lambda_i(D^2u) = c.$$

The special Lagrangian equation was first introduced by Harvey and Lawson in [2]. They proved that a Lagrangian graph $(x, Du(x))$ in $(\mathbb{R}^n \times \mathbb{R}^n, \delta_0)$ is minimal if and

only if the Lagrangian angle is a constant, that is, (1.5) holds. According to (1.5), several authors obtained the same Bernstein type theorems simultaneously using different techniques. Jost and Xin [3] used the properties of harmonic maps into convex subsets of Grassmannians. Yuan [4] used the geometric measure theory.

We will be considering the Lagrangian graphs of prescribed constant mean curvature κ in $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$ and Du is a diffeomorphism from Ω to $\tilde{\Omega}$. For $\kappa \equiv 0$, finding a minimal Lagrangian diffeomorphism between two uniformly convex bounded domains in $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$ is equivalent to solving (1.3) with second boundary condition

$$(1.6) \quad Du(\Omega) = \tilde{\Omega},$$

that is,

$$(1.7) \quad \begin{cases} F_\tau(\lambda(D^2u)) = c, & x \in \Omega, \\ Du(\Omega) = \tilde{\Omega}. \end{cases}$$

Here Du is a minimal Lagrangian diffeomorphism from Ω to $\tilde{\Omega}$ in $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$.

In dimension 2, Delanoë [5] obtained a unique smooth solution for the second boundary value problem of the Monge-Ampère equation for $\tau = 0$ in (1.7) if both domains are uniformly convex. Later the generalization of Delanoë's theorem to higher dimensions was given by Caffarelli [6] and Urbas [7]. Using the parabolic method, Schnürer and Smoczyk [8] also obtained the existence of solutions to (1.7) for $\tau = 0$.

As far as $\tau = \frac{\pi}{2}$ is concerned, Brendle and Warren [9] proved the existence and uniqueness of the solution by the elliptic method, and the second author [10] obtained the existence of solution by the parabolic method. Then by the elliptic and parabolic method, the second author with Ou [11], Ye [12] [13] and Chen [13] proved the existence and uniqueness of the solution to (1.7) for $0 < \tau < \frac{\pi}{2}$.

For a smooth function f , Chen, Zhang and the third author [14] proved that if u satisfies

$$(1.8) \quad \sum_{i=1}^n \arctan \lambda_i(D^2u) = f(x),$$

then $Df(x)$ is the mean curvature of the gradient graph $(x, Du(x))$ in $(\mathbb{R}^n \times \mathbb{R}^n, \delta_0)$. Motivated by the works of [1] and [14], for $\tau \in [0, \frac{\pi}{2}]$ we obtain a generalization of their results.

Proposition 1.1. *If u satisfies*

$$(1.9) \quad F_\tau(\lambda(D^2u)) = f(x),$$

Then $Df(x)$ is the mean curvature of the gradient graph $(x, Du(x))$ in $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$.

For $f(x) = \kappa \cdot x + c$, Proposition 1.1 becomes

Corollary 1.2. *If u satisfies (1.1), then κ is the constant mean curvature of the gradient graph $(x, Du(x))$ in $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$.*

By the continuity method, for the second boundary value problem (1.1), we have

Theorem 1.3. For $\tau \in (0, \frac{\pi}{2}]$, if $|\kappa|$ is sufficiently small, then there exist a uniformly convex solution $u \in C^\infty(\bar{\Omega})$ and a unique constant c solving (1.1), and u is unique up to a constant.

Remark 1.4. For $\tau = 0$, the problem (1.1) was solved by J. Urbas [7], O.C. Schnürer and K. Smoczyk [8].

The geometric meaning of this theorem is that if Ω and $\tilde{\Omega}$ are two uniformly convex bounded domains with smooth boundary in \mathbb{R}^n , then there exists a diffeomorphism $\psi = Du : \Omega \rightarrow \tilde{\Omega}$ such that

$$\Sigma := \{(x, \psi(x)) : x \in \Omega\}$$

is a Lagrangian submanifold, of which the mean curvature is κ in $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$.

In another paper, we shall point out by the parabolic method that c means the coefficient of time variable for the translating solution of the parabolic problem corresponding to (1.1), and therefore c can not be given in advance.

Theorem 1.3 presents an extension of the previous work on $\kappa = 0$ done by Brendle-Warren [9], Huang [10], Huang-Ou [11], Huang-Ye [12] and Chen-Huang-Ye [13].

The rest of this article is organized as follows. In Section 2, we give the proof of Proposition 1.1 and introduce a class of fully nonlinear elliptic equation containing (1.1). Then we present a theorem on the corresponding second boundary value problem, which is a generalization of Theorem 1.3. To prove this new theorem, we verify the strictly oblique estimate in Section 3, present the C^2 estimate in Section 4 and give the proof of this theorem by the continuity method in Section 5.

Throughout the following, Einstein's convention of summation over repeated indices will be adopted. We denote, for a smooth function u ,

$$u_i = \frac{\partial u}{\partial x_i}, \quad u_{ij} = \frac{\partial^2 u}{\partial x_i \partial x_j}, \quad u_{ijk} = \frac{\partial^3 u}{\partial x_i \partial x_j \partial x_k}, \dots$$

2. A GENERALIZATION OF THEOREM 1.3

We begin with the proof of Proposition 1.1.

Proof. Let $e_i = (0, \dots, 1, \dots, 0)$ be the i -th axis vector in $\mathbb{R}^n \times \mathbb{R}^n$, $i = 1, 2, \dots, 2n$. Then the tangential vector fields of $\vec{E} := (x, Du(x))$ —which is an n -dimensional submanifold of $(\mathbb{R}^n \times \mathbb{R}^n, g_\tau)$ —are

$$E_i = e_i + u_{ij}e_{n+j}, \quad i = 1, \dots, n.$$

Therefore,

$$\bar{\nabla}_{E_j}^{E_i} = u_{ijk}e_{n+k},$$

where $\bar{\nabla}$ is the Levi-Civita connection of \mathbb{R}^{2n} . Then the induced metric on \vec{E} is given by

$$g_{ij} = \langle E_i, E_j \rangle = \langle e_i + u_{ik}e_{n+k}, e_j + u_{jl}e_{n+l} \rangle = \sin \tau (\delta_{ij} + u_{ik}u_{kj}) + 2 \cos \tau u_{ij}.$$

Denote $(g^{ij}) = (g_{ij})^{-1}$, then the normal vector fields of \vec{E} are

$$M_\alpha = M_\alpha^i e_i + \tilde{M}_\alpha^j e_{n+j}, \quad \alpha = 1, \dots, n.$$

Consequently, the part of the mean curvature vector

$$\vec{H} = g^{ij} \left(\overline{\nabla}_{E_i}^{E_j} \right)^\perp$$

on M_α is

$$\begin{aligned} H_\alpha &:= \langle \vec{H}, M_\alpha \rangle \\ &= g^{ij} \langle \overline{\nabla}_{E_i}^{E_j}, M_\alpha \rangle \\ (2.1) \quad &= g^{ij} \langle u_{ijk} e_{n+k}, M_\alpha^l e_l + \tilde{M}_\alpha^p e_{n+p} \rangle \\ &= g^{ij} \left(\cos \tau u_{ijk} M_\alpha^k + \sin \tau u_{ijk} \tilde{M}_\alpha^k \right). \end{aligned}$$

Let $\vec{F} = (Df)^\perp$, then

$$\begin{aligned} F_\alpha &:= \langle Df^\perp, M_\alpha \rangle \\ (2.2) \quad &= \langle f_l e_{n+l}, M_\alpha^i e_i + \tilde{M}_\alpha^j e_{n+j} \rangle \\ &= \cos \tau f_l M_\alpha^l + \sin \tau f_l \tilde{M}_\alpha^l. \end{aligned}$$

Comparing (2.1) with (2.2), we know that if for any $k = 1, \dots, n$ we have

$$(2.3) \quad g^{ij} u_{ijk} = f_k,$$

then $\vec{H} = (Df)^\perp$.

In conclusion, we complete the proof of Proposition 1.1. \square

In order to prove Theorem 1.3 in four cases of all together, we reduce it to a more general form. For the convenience, we introduce some notations.

It is obvious that $F_\tau(\lambda_1, \dots, \lambda_n)$, $\tau \in (0, \frac{\pi}{2}]$ is a smooth symmetric function defined on Γ_n^+ , where

$$\Gamma_n^+ := \{(\lambda_1, \dots, \lambda_n) \in \mathbb{R}^n : \lambda_i > 0, i = 1, \dots, n\}.$$

By direct calculation, we get

$$F_\tau(0, \dots, 0) = \begin{cases} \frac{n\sqrt{a^2+1}}{2b} \ln \frac{a-b}{a+b}, & 0 < \tau < \frac{\pi}{4}, \\ -\sqrt{2}n, & \tau = \frac{\pi}{4}, \\ \frac{n\sqrt{a^2+1}}{b} \arctan \frac{a-b}{a+b}, & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ 0, & \tau = \frac{\pi}{2}, \end{cases}$$

$$F_\tau(+\infty, \dots, +\infty) = \begin{cases} 0, & 0 < \tau < \frac{\pi}{4}, \\ 0, & \tau = \frac{\pi}{4}, \\ \frac{n\pi\sqrt{a^2+1}}{4b}, & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ \frac{n\pi}{2}, & \tau = \frac{\pi}{2}, \end{cases}$$

$$\frac{\partial F_\tau}{\partial \lambda_i} = \begin{cases} \frac{\sqrt{a^2+1}}{(\lambda_i+a)^2 - b^2}, & 0 < \tau < \frac{\pi}{4}, \\ \frac{\sqrt{2}}{(1+\lambda_i)^2}, & \tau = \frac{\pi}{4}, \\ \frac{\sqrt{a^2+1}}{(\lambda_i+a)^2 + b^2}, & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ \frac{1}{1+\lambda_i^2}, & \tau = \frac{\pi}{2}, \end{cases}$$

and

$$\frac{\partial^2 F_\tau}{\partial \lambda_i \partial \lambda_j} = \begin{cases} -\frac{2\sqrt{a^2+1}(\lambda_j+a)\delta_{ij}}{[(\lambda_i+a)^2 - b^2]^2}, & 0 < \tau < \frac{\pi}{4}, \\ -\frac{2\sqrt{2}\delta_{ij}}{(1+\lambda_i)^3}, & \tau = \frac{\pi}{4}, \\ -\frac{2\sqrt{a^2+1}(\lambda_j+a)\delta_{ij}}{[(\lambda_i+a)^2 + b^2]^2}, & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ -\frac{2\lambda_j\delta_{ij}}{(1+\lambda_i^2)^2}, & \tau = \frac{\pi}{2}, \end{cases}$$

for $i, j = 1, \dots, n$. Then

$$(2.4) \quad -\infty < F_\tau(0, \dots, 0) < F_\tau(+\infty, \dots, +\infty) < +\infty, \quad \tau \in \left(0, \frac{\pi}{2}\right],$$

$$(2.5) \quad \frac{\partial F_\tau}{\partial \lambda_i} > 0, \quad 1 \leq i \leq n \quad \text{on } \Gamma_n^+,$$

and

$$(2.6) \quad \left(\frac{\partial^2 F_\tau}{\partial \lambda_i \partial \lambda_j}\right) \leq 0 \quad \text{on } \Gamma_n^+.$$

For any $s_1 > 0, s_2 > 0$, define

$$\Gamma_{s_1, s_2}^+ = \{(\lambda_1, \dots, \lambda_n) \in \Gamma_n^+ : 0 \leq \min_{1 \leq i \leq n} \lambda_i \leq s_1, \max_{1 \leq i \leq n} \lambda_i \geq s_2\}.$$

Then for any $(\lambda_1, \dots, \lambda_n) \in \Gamma_{]s_1, s_2[}^+$, we have

$$(2.7) \quad \sum_{i=1}^n \frac{\partial F_\tau}{\partial \lambda_i} \in \begin{cases} \left[\frac{\sqrt{a^2+1}}{(s_1+a)^2-b^2}, \frac{n\sqrt{a^2+1}}{a^2-b^2} \right], & 0 < \tau < \frac{\pi}{4}, \\ \left[\frac{\sqrt{2}}{(1+s_1)^2}, n\sqrt{2} \right], & \tau = \frac{\pi}{4}, \\ \left[\frac{\sqrt{a^2+1}}{(s_1+a)^2+b^2}, \frac{n\sqrt{a^2+1}}{a^2+b^2} \right], & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ \left[\frac{1}{1+s_1^2}, n \right], & \tau = \frac{\pi}{2}, \end{cases}$$

and

$$(2.8) \quad \sum_{i=1}^n \frac{\partial F_\tau}{\partial \lambda_i} \lambda_i^2 \in \begin{cases} \left[\frac{s_2^2 \sqrt{a^2+1}}{(s_2+a)^2-b^2}, n\sqrt{a^2+1} \right], & 0 < \tau < \frac{\pi}{4}, \\ \left[\frac{s_2^2 \sqrt{2}}{(1+s_2)^2}, n\sqrt{2} \right], & \tau = \frac{\pi}{4}, \\ \left[\frac{s_2^2 \sqrt{a^2+1}}{(s_2+a)^2+b^2}, n\sqrt{a^2+1} \right], & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ \left[\frac{s_2^2}{1+s_2^2}, n \right], & \tau = \frac{\pi}{2}. \end{cases}$$

For any $(\mu_1, \dots, \mu_n) \in \Gamma_n^+$, denote

$$\lambda_i = \frac{1}{\mu_i}, \quad 1 \leq i \leq n,$$

and

$$\tilde{F}_\tau(\mu_1, \dots, \mu_n) := -F_\tau(\lambda_1, \dots, \lambda_n).$$

Then

$$\frac{\partial \tilde{F}_\tau}{\partial \mu_i} = \lambda_i^2 \frac{\partial F_\tau}{\partial \lambda_i}, \quad \mu_i^2 \frac{\partial \tilde{F}_\tau}{\partial \mu_i} = \frac{\partial F_\tau}{\partial \lambda_i},$$

and

$$\begin{aligned} \frac{\partial^2 \tilde{F}_\tau}{\partial \mu_i \partial \mu_j} &= -\lambda_i^3 \left(\lambda_i \frac{\partial^2 F_\tau}{\partial \lambda_i^2} + 2 \frac{\partial F_\tau}{\partial \lambda_i} \right) \delta_{ij} \\ &= \begin{cases} -\frac{2\sqrt{a^2+1}(\mu_i+a)}{[(1+a\mu_i)^2-(b\mu_i)^2]^2} \delta_{ij}, & 0 < \tau < \frac{\pi}{4}, \\ -\frac{2\sqrt{2}\delta_{ij}}{(1+\mu_i)^3}, & \tau = \frac{\pi}{4}, \\ -\frac{2\sqrt{a^2+1}(\mu_i+a)}{[(1+a\mu_i)^2+(b\mu_i)^2]^2} \delta_{ij}, & \frac{\pi}{4} < \tau < \frac{\pi}{2}, \\ -\frac{2\mu_i\delta_{ij}}{(1+\mu_i^2)^2}, & \tau = \frac{\pi}{2}. \end{cases} \end{aligned}$$

Therefore, we have

$$\frac{\partial \tilde{F}_\tau}{\partial \mu_i} > 0, \quad 1 \leq i \leq n \quad \text{on } \Gamma_n^+,$$

and

$$(2.9) \quad \left(\frac{\partial^2 \tilde{F}_\tau}{\partial \mu_i \partial \mu_j} \right) \leq 0 \quad \text{on } \Gamma_n^+.$$

Motivated by (2.4)-(2.9), in order to prove Theorem 1.3, we introduce a class of nonlinear functions containing $F_\tau(\lambda)$, $\tau \in (0, \frac{\pi}{2}]$. Let $F(\lambda_1, \dots, \lambda_n)$ be a smooth symmetric function defined on Γ_n^+ , and satisfy

$$(2.10) \quad -\infty < F(0, \dots, 0) < F(+\infty, \dots, +\infty) < +\infty,$$

$$(2.11) \quad \frac{\partial F}{\partial \lambda_i} > 0, \quad 1 \leq i \leq n \quad \text{on } \Gamma_n^+,$$

and

$$(2.12) \quad \left(\frac{\partial^2 F}{\partial \lambda_i \partial \lambda_j} \right) \leq 0 \quad \text{on } \Gamma_n^+.$$

For any $(\mu_1, \dots, \mu_n) \in \Gamma_n^+$, denote

$$\lambda_i = \frac{1}{\mu_i}, \quad 1 \leq i \leq n,$$

and

$$\tilde{F}(\mu_1, \dots, \mu_n) := -F(\lambda_1, \dots, \lambda_n).$$

Assume that

$$(2.13) \quad \left(\frac{\partial^2 \tilde{F}}{\partial \mu_i \partial \mu_j} \right) \leq 0 \quad \text{on } \Gamma_n^+.$$

In addition, for $s_1, s_2 > 0$, we assume that there exist positive constants Λ_1 and Λ_2 , depending on s_1 and s_2 , such that for any $(\lambda_1, \dots, \lambda_n) \in \Gamma_{s_1, s_2}^+$,

$$(2.14) \quad \Lambda_1 \leq \sum_{i=1}^n \frac{\partial F}{\partial \lambda_i} \leq \Lambda_2,$$

and

$$(2.15) \quad \Lambda_1 \leq \sum_{i=1}^n \frac{\partial F}{\partial \lambda_i} \lambda_i^2 \leq \Lambda_2.$$

Remark 2.1. *Since*

$$\frac{\partial^2 \tilde{F}}{\partial \mu_i \partial \mu_j} = -\frac{\partial^2 F}{\partial \lambda_i \partial \lambda_j} \lambda_i^2 \lambda_j^2 - 2\lambda_i^3 \delta_{ij} \frac{\partial F}{\partial \lambda_i},$$

we cannot deduce (2.13) from (2.11) and (2.12).

By the discussion above, we have

Proposition 2.2. *The operator $F_\tau(\lambda)$, $\tau \in (0, \frac{\pi}{2}]$ satisfies the structure conditions (2.10)-(2.15).*

For $f(x) \in C^\infty(\bar{\Omega})$, we define

$$\text{osc}_\Omega(f) := \max_{x, y \in \Omega} |f(x) - f(y)|,$$

and

$$\mathcal{A}_\delta := \left\{ f(x) \in C^\infty(\bar{\Omega}) : f \text{ is concave, } \text{osc}_\Omega(f) \leq \delta \right\}.$$

The constant δ is any positive constant satisfying

$$\delta < \min \{ F(+\infty, \dots, +\infty) - F(\Theta_0, +\infty, \dots, +\infty), F(0, \dots, 0, \Theta_0) - F(0, \dots, 0) \},$$

where $\Theta_0 := \left(\frac{|\tilde{\Omega}|}{|\Omega|} \right)^{1/n}$.

Considering the more general second boundary value problem

$$(2.16) \quad \begin{cases} F(\lambda(D^2u)) = f(x) + c, & x \in \Omega, \\ Du(\Omega) = \tilde{\Omega}, \end{cases}$$

we claim that

Theorem 2.3. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If $|Df|$ is sufficiently small, then there exist a uniformly convex solution $u \in C^\infty(\bar{\Omega})$ and a unique constant c solving (2.16), and u is unique up to a constant.*

Remark 2.4. *It's not hard to deduce that if $|Df|$ is sufficiently small and f is concave, then $f \in \mathcal{A}_\delta$.*

Remark 2.5. Let $n \geq 2$, $f = \kappa_0 x_1$, $\Omega = \{x \in \mathbb{R}^n : |x| < 1\}$, and $\tilde{\Omega}$ be a uniformly convex domain in \mathbb{R}^n . If (u, c) satisfies

$$\begin{cases} \sum_{i=1}^n \arctan \lambda_i = f(x) + c, & x \in \Omega, \\ Du(\Omega) = \tilde{\Omega}. \end{cases}$$

Then

$$|Df| = |\kappa_0| = \frac{1}{n} \operatorname{osc}_{\tilde{\Omega}}(f + c) = \frac{1}{n} \operatorname{osc}_{\tilde{\Omega}} \left(\sum_{i=1}^n \arctan \lambda_i \right) < \frac{\pi}{2}.$$

Therefore, the range of $|Df|$ should be limited for the solvability of the problem (2.16).

By Proposition 2.2 we know that, once Theorem 2.3 is proved, Theorem 1.3 holds.

In the next three sections, we are going to prove Theorem 2.3 through the continuity method, which is based on the strictly oblique estimate and the C^2 estimate.

3. THE STRICT OBLIQUENESS ESTIMATE

To prove the strict obliqueness estimate, first we need

Lemma 3.1. Let $\lambda_1(x), \dots, \lambda_n(x)$ be the eigenvalues of D^2u at x . Suppose that (2.10) and (2.11) hold, if $\operatorname{osc}_{\tilde{\Omega}}(f) \leq \delta$ and $u \in C^\infty(\tilde{\Omega})$ is a uniformly convex solution of (2.16), then there exists $\mu > 0$ and $\omega > 0$ depending only on F , Θ_0 and δ such that

$$(3.1) \quad \min_{1 \leq i \leq n} \lambda_i(x) \leq \mu, \quad \max_{1 \leq i \leq n} \lambda_i(x) \geq \omega.$$

Proof. By $Du(\Omega) = \tilde{\Omega}$, we have

$$\int_{\Omega} \det D^2u(x) dx = |\tilde{\Omega}|.$$

Then we can find $\bar{x} \in \tilde{\Omega}$ such that

$$\prod_{i=1}^n \lambda_i(\bar{x}) = \det D^2u(\bar{x}) = \frac{|\tilde{\Omega}|}{|\Omega|} = \Theta_0^n.$$

Therefore,

$$\min_{1 \leq i \leq n} \lambda_i(\bar{x}) \leq \Theta_0 \leq \max_{1 \leq i \leq n} \lambda_i(\bar{x}).$$

By (2.16) and condition (2.11), we obtain

$$\begin{aligned} F \left(\min_{1 \leq i \leq n} \lambda_i(x), \dots, \min_{1 \leq i \leq n} \lambda_i(x) \right) &\leq F(\lambda_1(x), \dots, \lambda_n(x)) \\ &= F(\lambda_1(\bar{x}), \dots, \lambda_n(\bar{x})) + f(x) - f(\bar{x}) \\ &\leq F(\Theta_0, +\infty, \dots, +\infty) + \operatorname{osc}_{\tilde{\Omega}}(f) \\ &\leq F(\Theta_0, +\infty, \dots, +\infty) + \delta \\ &< F(+\infty, \dots, +\infty), \end{aligned}$$

and

$$\begin{aligned}
F\left(\max_{1 \leq i \leq n} \lambda_i(x), \dots, \max_{1 \leq i \leq n} \lambda_i(x)\right) &\geq F(\lambda_1(x), \dots, \lambda_n(x)) \\
&= F(\lambda_1(\bar{x}), \dots, \lambda_n(\bar{x})) + f(x) - f(\bar{x}) \\
&\geq F(0, \dots, 0, \Theta_0) - \underset{\Omega}{\text{osc}}(f) \\
&\geq F(0, \dots, 0, \Theta_0) - \delta \\
&> F(0, \dots, 0).
\end{aligned}$$

By the monotonicity of F and condition (2.10), we get the desired result. \square

By Lemma 3.1, the points $(\lambda_1, \dots, \lambda_n)$ are always in $\Gamma_{\mu, \omega}^+$ under the problem (2.16). Then there exist $\Lambda_1 > 0$ and $\Lambda_2 > 0$ depending only on F , Θ_0 and δ , such that F satisfies the structure conditions (2.14) and (2.15). In the following, we always assume that Λ_1 and Λ_2 are universal constants depending only on the known data.

For technical needs below, we introduce the Legendre transformation of u . For any $x \in \mathbb{R}^n$, define

$$\tilde{x}_i := \frac{\partial u}{\partial x_i}(x), \quad i = 1, 2, \dots, n,$$

and

$$\tilde{u}(\tilde{x}_1, \dots, \tilde{x}_n) := \sum_{i=1}^n x_i \frac{\partial u}{\partial x_i}(x) - u(x).$$

In terms of $\tilde{x}_1, \dots, \tilde{x}_n$ and $\tilde{u}(\tilde{x}_1, \dots, \tilde{x}_n)$, we can easily check that

$$\left(\frac{\partial^2 \tilde{u}}{\partial \tilde{x}_i \partial \tilde{x}_j}\right) = \left(\frac{\partial^2 u}{\partial x_i \partial x_j}\right)^{-1}.$$

Let μ_1, \dots, μ_n be the eigenvalues of $D^2 \tilde{u}$ at $\tilde{x} = Du(x)$. We denote

$$\mu_i = \lambda_i^{-1}, \quad i = 1, 2, \dots, n.$$

Then

$$\frac{\partial \tilde{F}}{\partial \mu_i} = \lambda_i^2 \frac{\partial F}{\partial \lambda_i}, \quad \mu_i^2 \frac{\partial \tilde{F}}{\partial \mu_i} = \frac{\partial F}{\partial \lambda_i}.$$

Moreover, it follows from (2.16) that

$$(3.2) \quad \begin{cases} \tilde{F}(\lambda(D^2 \tilde{u})) = -f(D\tilde{u}) - c, & \tilde{x} \in \tilde{\Omega}, \\ D\tilde{u}(\tilde{\Omega}) = \Omega. \end{cases}$$

Remark 3.2. By Lemma 3.1, if u is a smooth uniformly convex solution of (2.16), then the eigenvalues of $D^2 u$ and $D^2 \tilde{u}$ must be in $\Gamma_{\mu, \omega}^+$ and $\Gamma_{\omega^{-1}, \mu^{-1}}^+$ respectively. Therefore, \tilde{F} also satisfies the structure conditions (2.14) and (2.15).

Next, we will carry out the strictly oblique estimate. Let \mathcal{P}_n be the set of positive definite symmetric $n \times n$ matrices, and $\lambda_1(A), \dots, \lambda_n(A)$ be the eigenvalues of A . For $A = (a_{ij}) \in \mathcal{P}_n$, denote

$$F[A] := F(\lambda_1(A), \dots, \lambda_n(A))$$

and

$$(a^{ij}) = (a_{ij})^{-1}, \quad F^{ij} = \frac{\partial F}{\partial a_{ij}}, \quad F^{ij,rs} = \frac{\partial^2 F}{\partial a_{ij} \partial a_{rs}}.$$

Definition 3.3. A smooth function $h : \mathbb{R}^n \rightarrow \mathbb{R}$ is called the defining function of $\tilde{\Omega}$ if

$$\tilde{\Omega} = \{p \in \mathbb{R}^n : h(p) > 0\}, \quad |Dh|_{\partial\tilde{\Omega}} = 1,$$

and there exists $\theta > 0$ such that for any $p = (p_1, \dots, p_n) \in \tilde{\Omega}$ and $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$,

$$\frac{\partial^2 h}{\partial p_i \partial p_j} \xi_i \xi_j \leq -\theta |\xi|^2.$$

Therefore, the diffeomorphism condition $Du(\Omega) = \tilde{\Omega}$ in (2.16) is equivalent to

$$(3.3) \quad h(Du) = 0, \quad x \in \partial\Omega.$$

Then (2.16) can be rewritten as

$$(3.4) \quad \begin{cases} F[D^2u] = f(x) + c, & x \in \Omega, \\ h(Du) = 0, & x \in \partial\Omega. \end{cases}$$

This is an oblique boundary value problem of second order fully nonlinear elliptic equation. We also denote $\beta = (\beta^1, \dots, \beta^n)$ with $\beta^i := h_{p_i}(Du)$, and $\nu = (\nu_1, \dots, \nu_n)$ as the unit inward normal vector at $x \in \partial\Omega$. The expression of the inner product is

$$\langle \beta, \nu \rangle = \beta^i \nu_i.$$

Lemma 3.4. (See J. Urbas [7].) Let $\nu = (\nu_1, \nu_2, \dots, \nu_n)$ be the unit inward normal vector of $\partial\Omega$. If $u \in C^2(\tilde{\Omega})$ with $D^2u \geq 0$, then there holds $h_{p_k}(Du)\nu_k \geq 0$.

Now, we can present

Lemma 3.5. Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If u is a uniformly smooth convex solution of (2.16) and $|Df|$ is sufficiently small, then the strict obliqueness estimate

$$(3.5) \quad \langle \beta, \nu \rangle \geq \frac{1}{C_1} > 0$$

holds on $\partial\Omega$ for some universal constant C_1 , which depends only on F , Θ_0 , Ω , $\tilde{\Omega}$ and δ .

Proof. Define

$$v = \langle \beta, \nu \rangle + h(Du).$$

Let $x_0 \in \partial\Omega$ such that

$$\langle \beta, \nu \rangle(x_0) = h_{p_k}(Du(x_0))\nu_k(x_0) = \min_{\partial\Omega} \langle \beta, \nu \rangle.$$

By rotation, we may assume that $\nu(x_0) = (0, \dots, 0, 1) =: e_n$. Using the above assumptions and the boundary condition, we obtain

$$v(x_0) = \min_{\partial\Omega} v = h_{p_n}(Du(x_0)).$$

By the convexity of Ω and its smoothness, we extend ν smoothly to a tubular neighborhood of $\partial\Omega$ such that in the matrix sense

$$(3.6) \quad (\nu_{kl}) := (D_k \nu_l) \leq -\frac{1}{C} \text{diag}(1, \dots, 1, 0),$$

where C is a positive constant. By Lemma 3.4, we see that $h_{p_n}(Du(x_0)) \geq 0$.

At x_0 we have

$$(3.7) \quad 0 = v_r = h_{p_n p_k} u_{kr} + h_{p_k} \nu_{kr} + h_{p_k} u_{kr}, \quad 1 \leq r \leq n-1.$$

We assume that the following key estimate

$$(3.8) \quad v_n(x_0) > -C$$

holds which will be proved later, where C is a constant depending only on Ω , h , \tilde{h} and δ .

It's not hard to check that (3.8) can be rewritten as

$$(3.9) \quad h_{p_n p_k} u_{kn} + h_{p_k} \nu_{kn} + h_{p_k} u_{kn} > -C.$$

Multiplying (3.9) with h_{p_n} and (3.7) with h_{p_r} respectively, and summing up together, we obtain

$$(3.10) \quad h_{p_k} h_{p_l} u_{kl} \geq -C h_{p_n} - h_{p_k} h_{p_l} \nu_{kl} - h_{p_k} h_{p_n p_l} u_{kl}.$$

Using (3.6) and

$$1 \leq r \leq n-1, \quad h_{p_k} u_{kr} = \frac{\partial h(Du)}{\partial x_r} = 0, \quad h_{p_k} u_{kn} = \frac{\partial h(Du)}{\partial x_n} \geq 0, \quad -h_{p_n p_n} \geq 0,$$

we have

$$h_{p_k} h_{p_l} u_{kl} \geq -C h_{p_n} + \frac{1}{C} |Dh|^2 - \frac{1}{C} h_{p_n}^2 = -C h_{p_n} + \frac{1}{C} - \frac{1}{C} h_{p_n}^2.$$

For the last term of the above inequality, we distinguish two cases at x_0 .

Case (i). If

$$-C h_{p_n} + \frac{1}{C} - \frac{1}{C} h_{p_n}^2 \leq \frac{1}{2C},$$

then

$$h_{p_k}(Du)\nu_k = h_{p_n} \geq \sqrt{\frac{1}{2} + \frac{C^4}{4}} - \frac{C^2}{2}.$$

It shows that there is a uniform positive lower bound for the quantity $\min_{\partial\Omega} \langle \beta, \nu \rangle$.

Case (ii). If

$$-C h_{p_n} + \frac{1}{C} - \frac{1}{C} h_{p_n}^2 > \frac{1}{2C},$$

then we obtain a positive lower bound of $h_{p_k} h_{p_l} u_{kl}$.

Let \tilde{u} be the Legendre transformation of u , then \tilde{u} satisfies

$$(3.11) \quad \begin{cases} \tilde{F} [D^2\tilde{u}] = -f(D\tilde{u}) - c, & \tilde{x} \in \tilde{\Omega}, \\ \tilde{h}(D\tilde{u}) = 0, & \tilde{x} \in \partial\tilde{\Omega}, \end{cases}$$

where \tilde{h} is the defining function of Ω . That is,

$$\Omega = \{\tilde{p} \in \mathbb{R}^n : \tilde{h}(\tilde{p}) > 0\}, \quad |D\tilde{h}|_{\partial\Omega} = 1, \quad D^2\tilde{h} \leq -\tilde{\theta}I,$$

where $\tilde{\theta}$ is some positive constant. The unit inward normal vector of $\partial\Omega$ can be expressed by $\nu = D\tilde{h}$. For the same reason, $\tilde{\nu} = D\tilde{h}$, where $\tilde{\nu} = (\tilde{\nu}_1, \tilde{\nu}_2, \dots, \tilde{\nu}_n)$ is the unit inward normal vector of $\partial\tilde{\Omega}$.

Let $\tilde{\beta} = (\tilde{\beta}^1, \dots, \tilde{\beta}^n)$ with $\tilde{\beta}^k := \tilde{h}_{p_k}(D\tilde{u})$. We also define

$$\tilde{\nu} = \langle \tilde{\beta}, \tilde{\nu} \rangle + \tilde{h}(D\tilde{u}),$$

in which

$$\langle \tilde{\beta}, \tilde{\nu} \rangle = \langle \beta, \nu \rangle.$$

Denote $\tilde{x}_0 = Du(x_0)$. Then $\tilde{v}(\tilde{x}_0) = v(x_0) = \min_{\partial\tilde{\Omega}} \tilde{v}$. Using the same methods, under the assumption of

$$(3.12) \quad \tilde{v}_n(\tilde{x}_0) \geq -C,$$

we obtain the positive lower bounds of $\tilde{h}_{p_k}\tilde{h}_{p_l}\tilde{u}_{kl}$, or

$$h_{p_k}(Du)\nu_k = \tilde{h}_{p_k}(D\tilde{u})\tilde{\nu}_k = \tilde{h}_{p_n} \geq \sqrt{\frac{1}{2} + \frac{C^4}{4}} - \frac{C^2}{2}.$$

We notice that

$$\tilde{h}_{p_k}\tilde{h}_{p_l}\tilde{u}_{kl} = \nu_i\nu_j u^{ij}.$$

Then by the positive lower bounds of $h_{p_k}h_{p_l}u_{kl}$ and $\tilde{h}_{p_k}\tilde{h}_{p_l}\tilde{u}_{kl}$, the lemma follows from

$$(3.13) \quad \langle \beta, \nu \rangle = \sqrt{h_{p_k}h_{p_l}u_{kl}u^{ij}\nu_i\nu_j},$$

which is proved in [7].

It remains to prove the key estimate (3.8) and (3.12). We prove (3.8) first. By $D^2\tilde{h} \leq -\tilde{\theta}I$ and (2.14) we have

$$(3.14) \quad L\tilde{h} \leq -\tilde{\theta} \sum_{i=1}^n F^{ii},$$

where $L := F^{ij}\partial_{ij}$. On the other hand,

$$(3.15) \quad \begin{aligned} Lv &= h_{p_k p_l p_m} \nu_k F^{ij} u_{li} u_{mj} + 2h_{p_k p_l} F^{ij} \nu_{kj} u_{li} \\ &\quad + h_{p_k p_l} F^{ij} u_{lj} u_{ki} + h_{p_k p_l} \nu_k L u_l + h_{p_k} L \nu_k + h_{p_k} L u_k. \end{aligned}$$

At first we estimate the first term on the right hand side of (3.15). By the diagonal basis and (2.15), we have

$$|h_{p_k p_l p_m} \nu_k F^{ij} u_{li} u_{mj}| \leq C \sum_{i=1}^n \frac{\partial F}{\partial \lambda_i} \lambda_i^2 \leq C,$$

where C is a constant depending only on $h, \Omega, \Lambda_1, \Lambda_2$ and δ . Similarly, we also get

$$|h_{p_k p_l} F^{ij} u_l u_{ki}| \leq C \sum_{i=1}^n \frac{\partial F}{\partial \lambda_i} \lambda_i^2 \leq C.$$

For the second term, by Cauchy inequality, we obtain

$$\begin{aligned} |2h_{p_k p_l} F^{ij} \nu_{kj} u_{li}| &\leq C \sum_{i=1}^n \frac{\partial F}{\partial \lambda_i} \lambda_i = C \sum_{i=1}^n \sqrt{\frac{\partial F}{\partial \lambda_i}} \sqrt{\frac{\partial F}{\partial \lambda_i}} \lambda_i \\ &\leq C \left(\sum_{i=1}^n \frac{\partial F}{\partial \lambda_i} \right) \left(\sum_{i=1}^n \frac{\partial F}{\partial \lambda_i} \lambda_i^2 \right) \\ &\leq C. \end{aligned}$$

By (2.16) we have $Lu_l = f_l$. Then we get

$$|h_{p_k p_l} \nu_k Lu_l| \leq C, \quad |h_{p_k} Lu_k| \leq C.$$

It follows from (2.14) that

$$|h_{p_k} L\nu_k| \leq C \sum_{i=1}^n F^{ii}.$$

Inserting these into (3.15) and using (2.14), it is immediate to check that there exists a positive constant C depending only on $h, \Omega, \Lambda_1, \Lambda_2, \Theta_0$ and δ , such that

$$(3.16) \quad Lv \leq C \sum_{i=1}^n F^{ii}.$$

Denote a neighborhood of x_0 in Ω by

$$\Omega_\rho := \Omega \cap B_\rho(x_0),$$

where ρ is a positive constant such that ν is well defined in Ω_ρ . To obtain the desired results, it suffices to consider the function

$$\Phi(x) := v(x) - v(x_0) + C_0 \tilde{h}(x) + A|x - x_0|^2,$$

where C_0 and A are positive constants to be determined. On $\partial\Omega$, it is clear that $\Phi \geq 0$. Since v is bounded, we can choose A large enough such that on $\Omega \cap \partial B_\rho(x_0)$

$$\Phi(x) = v(x) - v(x_0) + C_0 \tilde{h}(x) + A\rho^2 > 0.$$

It follows from (3.14) that

$$L(C_0 \tilde{h}(x) + A|x - x_0|^2) \leq (-C_0 \tilde{\theta} + 2A) \sum_{i=1}^n F^{ii}.$$

Then using (3.16) and choosing $C_0 \gg A$ we have

$$L\Phi(x) \leq 0.$$

Therefore,

$$(3.17) \quad \begin{cases} L\Phi \leq 0, & x \in \Omega_\rho, \\ \Phi \geq 0, & x \in \partial\Omega_\rho. \end{cases}$$

We apply the maximum principle to get

$$\Phi|_{\Omega_\rho} \geq \min_{\partial\Omega_\rho} \Phi \geq 0.$$

Combining it with $\Phi(x_0) = 0$, we obtain $\partial_n \Phi(x_0) \geq 0$, which gives the desired estimate (3.8).

Finally, we prove (3.12). The proof of (3.12) is similar to the one of (3.8). Define

$$\tilde{L} := \tilde{F}^{ij} \partial_{ij} + f_{p_i} \partial_i.$$

By (3.11), we see that $\tilde{L}\tilde{u}_l = 0$, and thus

$$\tilde{L}\tilde{v} = \tilde{F}^{ij} \tilde{u}_{mj} \tilde{u}_{li} \tilde{h}_{p_k p_l p_m} \tilde{v}_k + 2\tilde{h}_{p_k p_l} \tilde{F}^{ij} \tilde{u}_{li} \tilde{v}_{kj} + \tilde{F}^{ij} \tilde{h}_{p_k} \tilde{v}_{kij} + \tilde{h}_{p_k p_l} \tilde{F}^{ij} \tilde{u}_{lj} \tilde{u}_{ki} + \tilde{h}_{p_k} f_{p_i} \tilde{v}_{ki}.$$

By making use of the following identities

$$\frac{\partial \tilde{F}}{\partial \mu_i} = \lambda_i^2 \frac{\partial F}{\partial \lambda_i}, \quad \mu_i^2 \frac{\partial \tilde{F}}{\partial \mu_i} = \frac{\partial F}{\partial \lambda_i}.$$

we deduce that \tilde{F} satisfies the structure conditions (2.10)-(2.15). Repeating the proof of (3.16), we have

$$(3.18) \quad \tilde{L}\tilde{v} \leq C \sum_{i=1}^n \tilde{F}^{ii},$$

where C depends only on Ω , $\tilde{\Omega}$, Θ_0 and δ .

Denote a neighborhood of \tilde{x}_0 in $\tilde{\Omega}$ by

$$\tilde{\Omega}_r := \tilde{\Omega} \cap B_r(\tilde{x}_0),$$

where r is a positive constant such that \tilde{v} is well defined in $\tilde{\Omega}_r$. Consider

$$\tilde{\Phi}(y) := \tilde{v}(y) - \tilde{v}(\tilde{x}_0) + \tilde{C}_0 h(y) + \tilde{A} |y - \tilde{x}_0|^2,$$

where \tilde{C}_0 and \tilde{A} are positive constants to be determined. It is clear that $\tilde{\Phi} \geq 0$ on $\partial\tilde{\Omega}$. Since \tilde{v} is bounded, we can choose \tilde{A} large enough such that on $\tilde{\Omega} \cap \partial B_r(\tilde{x}_0)$

$$\tilde{\Phi}(y) = \tilde{v}(y) - \tilde{v}(\tilde{x}_0) + \tilde{C}_0 h(y) + \tilde{A} r^2 > 0.$$

By (2.15) and (3.18), it is not difficult to show that

$$\tilde{L}\tilde{\Phi}(y) \leq \left(C - \frac{\tilde{C}_0 \theta}{2} + 2\tilde{A} \right) \sum_{i=1}^n \tilde{F}^{ii} + 2\tilde{A} f_{p_i}(y_i - \tilde{x}_{0i}) - \tilde{C}_0 \left(\frac{\theta}{2} \sum_{i=1}^n \tilde{F}^{ii} - f_{p_i} \partial_i h \right).$$

In order to make

$$\tilde{L}\tilde{\Phi}(y) \leq 0,$$

we only need to choose $\tilde{C}_0 \gg \tilde{A}$ and

$$|Df| \leq \frac{\theta \Lambda_1}{2} \cdot \frac{1}{\max_{\bar{\Omega}} |Dh|}.$$

Consequently,

$$(3.19) \quad \begin{cases} \tilde{L}\tilde{\Phi} \leq 0, & y \in \tilde{\Omega}_r, \\ \tilde{\Phi} \geq 0, & y \in \partial\tilde{\Omega}_r. \end{cases}$$

Therefore, we get (3.12) as same as the argument in (3.8). Thus the proof of (3.5) is completed. \square

4. THE C^2 ESTIMATE

The following definition provides a basic connection between (3.11) and (2.16) and will be used frequently in the sequel.

Definition 4.1. *We say that \tilde{u} in (3.11) is a dual solution to (2.16).*

We now proceed to carry out the C^2 estimate. The strategy is to reduce the C^2 global estimate of u and \tilde{u} to the boundary.

Lemma 4.2. *If u is a smooth uniformly convex solution of (2.16) and there hold (2.11), (2.12) and (2.14), then there exists a positive constant C depending only on n , Ω , $\tilde{\Omega}$, Λ_1 and $\text{diam}(\Omega)$, such that*

$$(4.1) \quad \sup_{\Omega} |D^2 u| \leq \max_{\partial\Omega} |D^2 u| + C \sup_{\Omega} |D^2 f|.$$

Proof. Without loss of generality, we may assume that Ω lies in cube $[0, d]^n$. Let

$$L := F^{ij} \partial_{ij}.$$

For any unit vector ξ , differentiating the equation in (2.16) twice in direction ξ gives

$$Lu_{\xi\xi} + F^{ij,rs} u_{ij\xi} u_{rs\xi} = f_{\xi\xi}.$$

Then by the concavity of F on Γ_n^+ , we have

$$(4.2) \quad Lu_{\xi\xi} = -F^{ij,rs} u_{ij\xi} u_{rs\xi} + f_{\xi\xi} \geq f_{\xi\xi}.$$

Let

$$v = \sup_{\partial\Omega} u_{\xi\xi} + \frac{1}{\Lambda_1} \left(ne^d - \sum_{i=1}^n e^{x_i} \right) \sup_{\Omega} |f_{\xi\xi}|.$$

By direct calculation and (2.14), we obtain

$$(4.3) \quad \begin{aligned} Lv &= -\frac{1}{\Lambda_1} \sup_{\Omega} |f_{\xi\xi}| \left(\sum_{i=1}^n e^{x_i} F^{ii} \right) \\ &\leq -\frac{1}{\Lambda_1} \sup_{\Omega} |f_{\xi\xi}| \left(\sum_{i=1}^n F^{ii} \right) \\ &\leq -\sup_{\Omega} |f_{\xi\xi}|. \end{aligned}$$

Combining (4.2) with (4.3), we have

$$L(v - u_{\xi\xi}) \leq -\left(\sup_{\Omega} |f_{\xi\xi}| + f_{\xi\xi} \right) \leq 0.$$

It is obvious that $v - u_{\xi\xi} \geq 0$ on $\partial\Omega$. Then by the maximum principle we obtain

$$\sup_{\Omega} u_{\xi\xi} \leq \sup_{\Omega} v \leq \sup_{\partial\Omega} u_{\xi\xi} + \frac{ne^d}{\Lambda_1} \sup_{\Omega} |f_{\xi\xi}|.$$

This completes the proof of (4.1). \square

Next, we estimate the second order derivative on the boundary. By differentiating the boundary condition $h(Du) = 0$ in any tangential direction ς , we have

$$(4.4) \quad u_{\beta\varsigma} = h_{p_k}(Du)u_{k\varsigma} = 0.$$

The second order derivative of u on the boundary is controlled by $u_{\beta\varsigma}$, $u_{\beta\beta}$ and $u_{\varsigma\varsigma}$. In the following we give the arguments as in [7], one can see there for more details.

At $x \in \partial\Omega$, any unit vector ξ can be written in terms of a tangential component $\varsigma(\xi)$ and a component in the direction β by

$$\xi = \varsigma(\xi) + \frac{\langle \nu, \xi \rangle}{\langle \beta, \nu \rangle} \beta,$$

where

$$\varsigma(\xi) := \xi - \langle \nu, \xi \rangle \nu - \frac{\langle \nu, \xi \rangle}{\langle \beta, \nu \rangle} \beta^T,$$

and

$$\beta^T := \beta - \langle \beta, \nu \rangle \nu.$$

By the strict obliqueness estimate (3.5), we have

$$(4.5) \quad \begin{aligned} |\varsigma(\xi)|^2 &= 1 - \left(1 - \frac{|\beta^T|^2}{\langle \beta, \nu \rangle^2}\right) \langle \nu, \xi \rangle^2 - 2\langle \nu, \xi \rangle \frac{\langle \beta^T, \xi \rangle}{\langle \beta, \nu \rangle} \\ &\leq 1 + C\langle \nu, \xi \rangle^2 - 2\langle \nu, \xi \rangle \frac{\langle \beta^T, \xi \rangle}{\langle \beta, \nu \rangle} \\ &\leq C. \end{aligned}$$

Denote $\varsigma := \frac{\varsigma(\xi)}{|\varsigma(\xi)|}$, then by (4.5) and (3.5) we obtain

$$(4.6) \quad \begin{aligned} u_{\xi\xi} &= |\varsigma(\xi)|^2 u_{\varsigma\varsigma} + 2|\varsigma(\xi)| \frac{\langle \nu, \xi \rangle}{\langle \beta, \nu \rangle} u_{\beta\varsigma} + \frac{\langle \nu, \xi \rangle^2}{\langle \beta, \nu \rangle^2} u_{\beta\beta} \\ &= |\varsigma(\xi)|^2 u_{\varsigma\varsigma} + \frac{\langle \nu, \xi \rangle^2}{\langle \beta, \nu \rangle^2} u_{\beta\beta} \\ &\leq C(u_{\varsigma\varsigma} + u_{\beta\beta}), \end{aligned}$$

where C depends only on Ω , $\tilde{\Omega}$, Λ_1 , Λ_2 , δ and the constant C_1 in (3.5). Therefore, we only need to estimate $u_{\beta\beta}$ and $u_{\varsigma\varsigma}$ respectively.

First we have

Lemma 4.3. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If u is a smooth uniformly convex solution of (2.16), then there exists a positive constant C depending only on Ω , $\tilde{\Omega}$, Λ_1 , Λ_2 and δ , such that*

$$(4.7) \quad \max_{\partial\Omega} u_{\beta\beta} \leq C.$$

Proof. Let $x_0 \in \partial\Omega$ satisfy $u_{\beta\beta}(x_0) = \max_{\partial\Omega} u_{\beta\beta}$. Consider the barrier function

$$\Psi := -h(Du) + C_0\tilde{h} + A|x - x_0|^2.$$

For any $x \in \partial\Omega$, $Du(x) \in \partial\tilde{\Omega}$, then $h(Du) = 0$. It is clear that $\tilde{h} = 0$ on $\partial\Omega$. As the proof of (3.17), we can find the constants C_0 and A such that

$$(4.8) \quad \begin{cases} L\Psi \leq 0, & x \in \Omega_\rho, \\ \Psi \geq 0, & x \in \partial\Omega_\rho. \end{cases}$$

By the maximum principle, we get

$$\Psi(x) \geq 0, \quad x \in \Omega_\rho.$$

Combining it with $\Psi(x_0) = 0$ we obtain $\Psi_\beta(x_0) \geq 0$, which implies

$$\frac{\partial h}{\partial \beta}(Du(x_0)) \leq C_0.$$

On the other hand, we see that at x_0 ,

$$\frac{\partial h}{\partial \beta} = \langle Dh(Du), \beta \rangle = \frac{\partial h}{\partial p_k} u_{kl} \beta^l = \beta^k u_{kl} \beta^l = u_{\beta\beta}.$$

Therefore,

$$u_{\beta\beta} = \frac{\partial h}{\partial \beta} \leq C.$$

□

Next, we estimate the double tangential derivative.

Lemma 4.4. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If u is a smooth uniformly convex solution of (2.16), then there exists a positive constant C depending only on Ω , $\tilde{\Omega}$, Λ_1 , Λ_2 and δ , such that*

$$(4.9) \quad \max_{\partial\Omega} u_{\zeta\zeta} \leq C.$$

Proof. Assume that $u_{\zeta\zeta}|_{\partial\Omega}$ attains its maximum at $x_0 \in \partial\Omega$. Let

$$M := u_{11}(x_0) = \max_{\partial\Omega} u_{\zeta\zeta},$$

and e_n be the unit inward normal vector of $\partial\Omega$ at x_0 .

For any $x \in \partial\Omega$, we have by (4.5),

$$(4.10) \quad \begin{aligned} u_{\xi\xi} &= |\varsigma(\xi)|^2 u_{\zeta\zeta} + \frac{\langle \nu, \xi \rangle^2}{\langle \beta, \nu \rangle^2} u_{\beta\beta} \\ &\leq \left(1 + C \langle \nu, \xi \rangle^2 - 2 \langle \nu, \xi \rangle \frac{\langle \beta^T, \xi \rangle}{\langle \beta, \nu \rangle} \right) M + \frac{\langle \nu, \xi \rangle^2}{\langle \beta, \nu \rangle^2} u_{\beta\beta}. \end{aligned}$$

Without loss of generality, we assume that $M \geq 1$. Then by (3.5) and (4.7) we have

$$(4.11) \quad \frac{u_{\xi\xi}}{M} + 2 \langle \nu, \xi \rangle \frac{\langle \beta^T, \xi \rangle}{\langle \beta, \nu \rangle} \leq 1 + C \langle \nu, \xi \rangle^2.$$

Let $\xi = e_1$, then

$$(4.12) \quad \frac{u_{11}}{M} + 2 \langle \nu, e_1 \rangle \frac{\langle \beta^T, e_1 \rangle}{\langle \beta, \nu \rangle} \leq 1 + C \langle \nu, e_1 \rangle^2.$$

We see that the function

$$(4.13) \quad w := A|x - x_0|^2 - \frac{u_{11}}{M} - 2\langle \nu, e_1 \rangle \frac{\langle \beta^T, e_1 \rangle}{\langle \beta, \nu \rangle} + C\langle \nu, e_1 \rangle^2 + 1$$

satisfies

$$w|_{\partial\Omega} \geq 0, \quad w(x_0) = 0.$$

Then, it follows by (4.1) that we can choose the constant A large enough such that

$$w|_{\Omega \cap \partial B_\rho(x_0)} \geq 0.$$

By (2.16), we deduce that

$$Lw \leq C \sum_{i=1}^n F^{ii}.$$

As in the proof of Lemma 4.3, we consider the function

$$\Upsilon := w + C_0 \tilde{h}.$$

A standard barrier argument shows that

$$\Upsilon_\beta(x_0) \geq 0.$$

Therefore,

$$(4.14) \quad u_{11\beta}(x_0) \leq CM.$$

On the other hand, differentiating $h(Du)$ twice in the direction e_1 at x_0 , we have

$$h_{p_k} u_{k11} + h_{p_k p_l} u_{k1} u_{l1} = 0.$$

The concavity of h yields that

$$h_{p_k} u_{k11} = -h_{p_k p_l} u_{k1} u_{l1} \geq \theta M^2.$$

Combining it with $h_{p_k} u_{k11} = u_{11\beta}$, and using (4.14) we obtain

$$\theta M^2 \leq CM.$$

Then we get the upper bound of $M = u_{11}(x_0)$ and thus the desired result follows. \square

By Lemma 4.3, Lemma 4.4 and (4.6), we obtain the C^2 a-priori estimate on the boundary.

Lemma 4.5. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If u is a smooth uniformly convex solution of (2.16), then there exists a positive constant C depending only on Ω , $\tilde{\Omega}$, Λ_1 , Λ_2 and δ , such that*

$$(4.15) \quad \max_{\partial\Omega} |D^2 u| \leq C.$$

In terms of Lemma 4.2 and Lemma 4.5, we see that

Lemma 4.6. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If u is a smooth uniformly convex solution of (2.16), then there exists a positive constant C depending only on Ω , $\tilde{\Omega}$, Λ_1 , Λ_2 and δ , such that*

$$(4.16) \quad \max_{\tilde{\Omega}} |D^2 u| \leq C.$$

In the following, we describe the positive lower bound of D^2u . For (3.11), in consider of the Legendre transformation of u , define

$$\tilde{L} := \tilde{F}^{ij} \partial_{ij} + f_{p_i} \partial_i.$$

Then our goal is to show the upper bound of $D^2\tilde{u}$ and the argument is very similar to the one used in the proof of Lemma 4.6 by the concavity of f and the condition that $|Df|$ being sufficiently small. For the convenience of readers, we give the details.

At the beginning of the repeating procedure, we have

Lemma 4.7. *Suppose that f is concave on Ω . If \tilde{u} is a smooth uniformly convex solution of (3.11), then there holds*

$$(4.17) \quad \sup_{\tilde{\Omega}} |D^2\tilde{u}| \leq \max_{\partial\tilde{\Omega}} |D^2\tilde{u}|.$$

Proof. For any unit vector $\tilde{\xi}$, differentiating the equation in (3.11) twice in direction $\tilde{\xi}$ gives

$$\tilde{L}\tilde{u}_{\tilde{\xi}\tilde{\xi}} + \tilde{F}^{ij,rs}\tilde{u}_{ij\tilde{\xi}}\tilde{u}_{rs\tilde{\xi}} + \frac{\partial^2 f}{\partial p_i \partial p_j} \tilde{u}_{i\tilde{\xi}}\tilde{u}_{j\tilde{\xi}} = 0.$$

Then by the concavity of \tilde{F} on Γ_n^+ and f on Ω , we have

$$\tilde{L}\tilde{u}_{\tilde{\xi}\tilde{\xi}} = -\tilde{F}^{ij,rs}\tilde{u}_{ij\tilde{\xi}}\tilde{u}_{rs\tilde{\xi}} - \frac{\partial^2 f}{\partial p_i \partial p_j} \tilde{u}_{i\tilde{\xi}}\tilde{u}_{j\tilde{\xi}} \geq 0.$$

Then by the maximum principle we obtain

$$\sup_{\tilde{\Omega}} \tilde{u}_{\tilde{\xi}\tilde{\xi}} \leq \sup_{\partial\tilde{\Omega}} \tilde{u}_{\tilde{\xi}\tilde{\xi}}.$$

This completes the proof of (4.17). \square

Recall that $\tilde{\beta} = (\tilde{\beta}^1, \dots, \tilde{\beta}^n)$ with $\tilde{\beta}^k := \tilde{h}_{p_k}(D\tilde{u})$ and $\tilde{\nu} = (\tilde{\nu}_1, \tilde{\nu}_2, \dots, \tilde{\nu}_n)$ is the unit inward normal vector of $\partial\tilde{\Omega}$. Similar to the discussion of (4.4), (4.5) and (4.6), for any tangential direction $\tilde{\zeta}$, we have

$$(4.18) \quad u_{\tilde{\beta}\tilde{\zeta}} = \tilde{h}_{p_k}(D\tilde{u})\tilde{u}_{k\tilde{\zeta}} = 0.$$

Then the second order derivative of \tilde{u} on the boundary is also controlled by $u_{\tilde{\beta}\tilde{\zeta}}$, $u_{\tilde{\beta}\tilde{\beta}}$ and $u_{\tilde{\zeta}\tilde{\zeta}}$.

At $\tilde{x} \in \partial\tilde{\Omega}$, any unit vector $\tilde{\xi}$ can be written in terms of a tangential component $\tilde{\zeta}(\tilde{\xi})$ and a component in the direction $\tilde{\beta}$ by

$$\tilde{\xi} = \tilde{\zeta}(\tilde{\xi}) + \frac{\langle \tilde{\nu}, \tilde{\xi} \rangle}{\langle \tilde{\beta}, \tilde{\nu} \rangle} \tilde{\beta},$$

where

$$\tilde{\zeta}(\tilde{\xi}) := \tilde{\xi} - \langle \tilde{\nu}, \tilde{\xi} \rangle \tilde{\nu} - \frac{\langle \tilde{\nu}, \tilde{\xi} \rangle}{\langle \tilde{\beta}, \tilde{\nu} \rangle} \tilde{\beta}^T,$$

and

$$\tilde{\beta}^T := \tilde{\beta} - \langle \tilde{\beta}, \tilde{\nu} \rangle \tilde{\nu}.$$

Therefore,

$$(4.19) \quad |\tilde{\zeta}(\tilde{\xi})| \leq C,$$

and

$$(4.20) \quad u_{\tilde{\zeta}\tilde{\xi}} \leq C(u_{\zeta\bar{\zeta}} + u_{\tilde{\beta}\tilde{\beta}}),$$

where $\tilde{\zeta} := \frac{\tilde{\zeta}(\tilde{\xi})}{|\tilde{\zeta}(\tilde{\xi})|}$ and C depends only on Ω , $\tilde{\Omega}$, Λ_1 , Λ_2 , δ and the constant C_1 in (3.5). Then we also only need to estimate $u_{\tilde{\beta}\tilde{\beta}}$ and $u_{\zeta\bar{\zeta}}$ respectively.

Indeed, as shown by Lemma 4.3, we state

Lemma 4.8. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If \tilde{u} is a smooth uniformly convex solution of (3.11) and $|Df|$ is sufficiently small, then there exists a positive constant C depending only on Ω , $\tilde{\Omega}$, Λ_1 , Λ_2 and δ , such that*

$$(4.21) \quad \max_{\partial\Omega} \tilde{u}_{\tilde{\beta}\tilde{\beta}} \leq C.$$

Proof. Let $\tilde{x}_0 \in \partial\tilde{\Omega}$ satisfy $\tilde{u}_{\tilde{\beta}\tilde{\beta}}(\tilde{x}_0) = \max_{\partial\Omega} \tilde{u}_{\tilde{\beta}\tilde{\beta}}$. To estimate the upper bound of $\tilde{u}_{\tilde{\beta}\tilde{\beta}}$, we consider the barrier function

$$\tilde{\Psi} := -\tilde{h}(D\tilde{u}) + C_0h + A|y - \tilde{x}_0|^2.$$

For any $y \in \partial\tilde{\Omega}$, $D\tilde{u}(y) \in \partial\Omega$, then $\tilde{h}(D\tilde{u}) = 0$. It is clear that $h = 0$ on $\partial\tilde{\Omega}$. As the proof of (3.19) in terms of $|Df|$ being sufficiently small, we can find the constants C_0 and A such that

$$(4.22) \quad \begin{cases} \tilde{L}\tilde{\Psi} \leq 0, & y \in \tilde{\Omega}_r, \\ \tilde{\Psi} \geq 0, & y \in \partial\tilde{\Omega}_r. \end{cases}$$

By the maximum principle, we get

$$\tilde{\Psi}(y) \geq 0, \quad y \in \tilde{\Omega}_r.$$

Combining it with $\tilde{\Psi}(\tilde{x}_0) = 0$ we obtain $\tilde{\Psi}_{\tilde{\beta}}(\tilde{x}_0) \geq 0$, which implies

$$\frac{\partial\tilde{h}}{\partial\tilde{\beta}}(D\tilde{u}(\tilde{x}_0)) \leq C_0.$$

On the other hand, we see that at \tilde{x}_0 ,

$$\frac{\partial\tilde{h}}{\partial\tilde{\beta}} = \langle D\tilde{h}(D\tilde{u}), \tilde{\beta} \rangle = \frac{\partial\tilde{h}}{\partial p_k} \tilde{u}_{kl} \tilde{\beta}^l = \tilde{\beta}^k \tilde{u}_{kl} \tilde{\beta}^l = \tilde{u}_{\tilde{\beta}\tilde{\beta}}.$$

Therefore,

$$\tilde{u}_{\tilde{\beta}\tilde{\beta}} = \frac{\partial\tilde{h}}{\partial\tilde{\beta}} \leq C.$$

□

Next, we estimate the double tangential derivative of \tilde{u} .

Lemma 4.9. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If \tilde{u} is a smooth uniformly convex solution of (3.11) and $|Df|$ is sufficiently small, then there exists a positive constant C depending only on $\Omega, \tilde{\Omega}, \Lambda_1, \Lambda_2$ and δ , such that*

$$(4.23) \quad \max_{\partial\tilde{\Omega}} \tilde{u}_{\zeta\bar{\zeta}} \leq C.$$

Proof. Assume that $\tilde{u}_{\zeta\bar{\zeta}}|_{\partial\tilde{\Omega}}$ attains its maximum at $\tilde{x}_0 \in \partial\tilde{\Omega}$. Let

$$\tilde{M} := \tilde{u}_{11}(\tilde{x}_0) = \max_{\partial\tilde{\Omega}} \tilde{u}_{\zeta\bar{\zeta}},$$

and e_n be the unit inward normal vector of $\partial\tilde{\Omega}$ at \tilde{x}_0 .

For any $y \in \partial\tilde{\Omega}$, we have by (4.19),

$$(4.24) \quad \begin{aligned} \tilde{u}_{\xi\bar{\xi}} &= |\tilde{\zeta}(\tilde{\xi})|^2 \tilde{u}_{\zeta\bar{\zeta}} + \frac{\langle \tilde{\nu}, \tilde{\xi} \rangle^2}{\langle \tilde{\beta}, \tilde{\nu} \rangle^2} \tilde{u}_{\tilde{\beta}\bar{\beta}} \\ &\leq \left(1 + C \langle \tilde{\nu}, \tilde{\xi} \rangle^2 - 2 \langle \tilde{\nu}, \tilde{\xi} \rangle \frac{\langle \tilde{\beta}^T, \tilde{\xi} \rangle}{\langle \tilde{\beta}, \tilde{\nu} \rangle} \right) \tilde{M} + \frac{\langle \tilde{\nu}, \tilde{\xi} \rangle^2}{\langle \tilde{\beta}, \tilde{\nu} \rangle^2} \tilde{u}_{\tilde{\beta}\bar{\beta}}. \end{aligned}$$

Without loss of generality, we assume that $\tilde{M} \geq 1$. Then by (3.5) and (4.21) we have

$$(4.25) \quad \frac{\tilde{u}_{\xi\bar{\xi}}}{\tilde{M}} + 2 \langle \tilde{\nu}, \tilde{\xi} \rangle \frac{\langle \tilde{\beta}^T, \tilde{\xi} \rangle}{\langle \tilde{\beta}, \tilde{\nu} \rangle} \leq 1 + C \langle \tilde{\nu}, \tilde{\xi} \rangle^2.$$

Let $\tilde{\xi} = e_1$, then

$$(4.26) \quad \frac{\tilde{u}_{11}}{\tilde{M}} + 2 \langle \tilde{\nu}, e_1 \rangle \frac{\langle \tilde{\beta}^T, e_1 \rangle}{\langle \tilde{\beta}, \tilde{\nu} \rangle} \leq 1 + C \langle \tilde{\nu}, e_1 \rangle^2.$$

We see that the function

$$(4.27) \quad \tilde{w} := A|y - \tilde{x}_0|^2 - \frac{\tilde{u}_{11}}{\tilde{M}} - 2 \langle \tilde{\nu}, e_1 \rangle \frac{\langle \tilde{\beta}^T, e_1 \rangle}{\langle \tilde{\beta}, \tilde{\nu} \rangle} + C \langle \tilde{\nu}, e_1 \rangle^2 + 1$$

satisfies

$$\tilde{w}|_{\partial\tilde{\Omega}} \geq 0, \quad \tilde{w}(\tilde{x}_0) = 0.$$

Then, by (4.17) we can choose the constant A large enough such that

$$\tilde{w}|_{\tilde{\Omega} \cap \partial B_r(\tilde{x}_0)} \geq 0.$$

By (3.11), $f \in \mathcal{A}_\delta$ and $|Df|$ is sufficiently small, we can show that

$$\tilde{L}\tilde{w} \leq C \sum_{i=1}^n \tilde{F}^{ii}.$$

Since the proof of Lemma 4.8, let us define

$$\tilde{\Upsilon} := \tilde{w} + C_0 h.$$

A standard barrier argument makes conclusion of

$$\tilde{\Upsilon}_{\tilde{\beta}}(\tilde{x}_0) \geq 0.$$

Therefore,

$$(4.28) \quad \tilde{u}_{11\tilde{\beta}}(\tilde{x}_0) \leq C\tilde{M}.$$

On the other hand, differentiating $\tilde{h}(D\tilde{u})$ twice in the direction e_1 at \tilde{x}_0 , we have

$$\tilde{h}_{p_k} \tilde{u}_{k11} + \tilde{h}_{p_k p_l} \tilde{u}_{k1} \tilde{u}_{l1} = 0.$$

The concavity of \tilde{h} yields that

$$\tilde{h}_{p_k} \tilde{u}_{k11} = -\tilde{h}_{p_k p_l} \tilde{u}_{k1} \tilde{u}_{l1} \geq \tilde{\theta} \tilde{M}^2.$$

Combining it with $\tilde{h}_{p_k} \tilde{u}_{k11} = \tilde{u}_{11\tilde{\beta}}$, and using (4.28) we obtain

$$\tilde{\theta} \tilde{M}^2 \leq C\tilde{M}.$$

Then we get the upper bound of $\tilde{M} = \tilde{u}_{11}(\tilde{x}_0)$ and thus the desired result follows. \square

By Lemma 4.8, Lemma 4.9 and (4.20), we obtain the C^2 a-priori estimate of \tilde{u} on the boundary.

Lemma 4.10. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If \tilde{u} is a smooth uniformly convex solution of (3.11) and $|Df|$ is sufficiently small, then there exists a positive constant C depending only on $\Omega, \tilde{\Omega}, \Lambda_1, \Lambda_2$ and δ , such that*

$$(4.29) \quad \max_{\partial\tilde{\Omega}} |D^2\tilde{u}| \leq C.$$

By Lemma 4.7 and Lemma 4.10, one can see that

Lemma 4.11. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If \tilde{u} is a smooth uniformly convex solution of (3.11) and $|Df|$ is sufficiently small, then there exists a positive constant C depending only on $\Omega, \tilde{\Omega}, \Lambda_1, \Lambda_2$ and δ , such that*

$$(4.30) \quad \max_{\tilde{\Omega}} |D^2\tilde{u}| \leq C.$$

By Lemma 4.6 and Lemma 4.11, we conclude that

Lemma 4.12. *Let F satisfy the structure conditions (2.10)-(2.15) and $f \in \mathcal{A}_\delta$. If u is a smooth uniformly convex solution of (2.16) and $|Df|$ is sufficiently small, then there exists a positive constant C depending only on $\Omega, \tilde{\Omega}, \Lambda_1, \Lambda_2$ and δ , such that*

$$(4.31) \quad \frac{1}{C}I_n \leq D^2u(x) \leq CI_n, \quad x \in \tilde{\Omega},$$

where I_n is the $n \times n$ identity matrix.

5. PROOF OF THEOREM 2.3

In this section, we will prove Theorem 2.3. Two lemmas will be needed when we start the proof. Thanks to the discussion on the uniqueness in the fifth section of [9], we deduce that

Lemma 5.1. *If (2.11) holds and $u \in C^\infty(\tilde{\Omega})$ be uniformly convex solutions of (2.16), then u is unique up to a constant.*

Next, similar to Corollary 1.2 in [12], we obtain

Lemma 5.2. *If (2.11) holds and $u \in C^2(\bar{\Omega})$ is a uniformly convex solution of (2.16), then $u \in C^\infty(\bar{\Omega})$.*

Proof. By Evans-Krylov theorem, it's obvious that

$$u \in C^\infty(\Omega).$$

In the following, we describe the regularity of u on the boundary.

Suppose that $x_0 \in \partial\Omega$ and in the neighbourhood of x_0 ,

$$\partial\Omega = \{x_n = \varphi(x') : x' \in \Omega'\}$$

for some domain $\Omega' \subset \mathbb{R}^{n-1}$. Without loss of generality, we set

$$x_0 = (\theta, 0), \varphi(0) = 0, D\varphi(0) = \theta, D^2\varphi(0) > 0.$$

Let m_0 be small enough such that

$$\Omega_{m_0} := \Omega \cap \{x_n < m_0\} = \{(x', x_n) : x' \in \Omega'', \varphi(x') < x_n < m_0\}$$

for some domain $\Omega'' \subset \mathbb{R}^{n-1}$. And we denote Ω'' also by Ω' .

We take the coordinate transformation in the following

$$T : \Omega_{m_0} \rightarrow T(\Omega_{m_0}), \quad x = (x', x_n) \mapsto z = (z', z_n) = (x', x_n - \varphi(x')),$$

where

$$T(\Omega_{m_0}) := \{(z', z_n) : z' \in \Omega', \quad 0 < z_n < m_0 - \varphi(z')\}.$$

We set $w(z) := u(T^{-1}(z))$. Then $u(x) = w(x', x_n - \varphi(x'))$. Such that for $1 \leq i \leq n-1$, we see that $u_i = w_i - w_n \varphi_i$ and $u_n = w_n$. Furthermore, for $1 \leq k, l \leq n-1$, a direct computation shows that

$$\begin{aligned} u_{kl} &= w_{kl} - w_{kn} \varphi_l - w_{nl} \varphi_k + w_{nn} \varphi_k \varphi_l, \\ u_{kn} &= w_{kn} - w_{nn} \varphi_k, \\ u_{ln} &= w_{ln} - w_{nn} \varphi_l, \\ u_{nn} &= w_{nn}. \end{aligned}$$

It is obvious that by

$$\|D^2u\|_{C(\bar{\Omega})} + \|Du\|_{C(\bar{\Omega})} \leq C$$

we get

$$(5.1) \quad \|D^2w\|_{C(\overline{T(\Omega_{m_0})})} + \|Dw\|_{C(\overline{T(\Omega_{m_0})})} \leq C,$$

where C is constant depending only on the known data.

Since $w(z) = u(T^{-1}(z))$ and $x = T^{-1}(z)$, we obtain

$$D^2u = D(DT \cdot Dw),$$

$$D_x u = D_z w - w_n (D_{z'} \varphi, 0)^t,$$

where D_x, D_z and $D_{z'}$ denote the gradient according to the variables x, z and z' .

Then by (2.16), w satisfies

$$(5.2) \quad \begin{cases} F[\lambda(D(DT \cdot Dw))] = f(T^{-1}(z)) + c, & z \in T(\Omega_{m_0}), \\ h(D_z w - w_n (D_{z'} \varphi, 0)^t) = 0, & z \in \partial T(\Omega_{m_0}) \cap \{z_n = 0\}. \end{cases}$$

When $1 \leq s \leq n-1$, we set $v = \frac{\partial w}{\partial z_s}$. Let $a^{ij}(1 \leq i, j \leq n), \beta^i(1 \leq i \leq n)$ be the definitions in Section 3.

In the following, the repeated indices k, l denote the summation from 1 to $n-1$ and the repeated indices i, j present the summation from 1 to n . Define

$$[A^{ij}]_{1 \leq i, j \leq n} = \begin{cases} a^{kl}, & 1 \leq k, l \leq n-1, (i, j) = (k, l), \\ a^{kn} - a^{kl}\varphi_l, & 1 \leq k \leq n-1, (i, j) = (k, n), \\ a^{nn} + a^{kl}\varphi_k\varphi_l - 2a^{kn}\varphi_k, & (i, j) = (n, n). \end{cases}$$

Then differentiating the equation (5.2) with respect to z_s , we see that v solves the following linear equation with oblique boundary condition

$$(5.3) \quad \begin{cases} L_s v = \frac{\partial}{\partial z_s} f(T^{-1}(z)), & z \in T(\Omega_{m_0}), \\ \gamma^k v_k + \gamma^n v_n = \phi, & z \in \partial T(\Omega_{m_0}) \cap \{z_n = 0\}, \end{cases}$$

where

$$L_s = A^{ij} \frac{\partial^2}{\partial z_i \partial z_j},$$

$$\gamma^k = \beta^k (1 \leq k \leq n-1), \quad \gamma^n = \beta^n - \beta^k \varphi_k,$$

$$\phi = w_n \beta^k \varphi_{ks}.$$

By Theorem 1.1 in [16], we get $u \in C^{2,\alpha}(\bar{\Omega})$ for some $0 < \alpha < 1$, from which one deduces that

$$A^{ij} \in C^\alpha(T(\bar{\Omega}_{m_0})),$$

$$\gamma^i \in C^{1,\alpha}(T(\bar{\Omega}_{m_0})),$$

$$\phi \in C^{1,\alpha}(T(\bar{\Omega}_{m_0})).$$

It's obvious that the unit inward normal vector of $\partial\Omega$ at $x_0 = (\theta, 0)$ is $\nu = (0, 0, \dots, 0, 1)$. Using lemma 3.5, one can shows that

$$\beta^n|_{x_0} = \langle \beta, \nu \rangle|_{x_0} \geq \frac{1}{C_1}.$$

By $D\varphi(\theta) = 0$, taking $r > 0$ small enough such that for any $z' \in B'_r(\theta) \subset \Omega'$, we obtain

$$3n\Lambda_2 \geq [A^{ij}] \geq \frac{1}{3n}\Lambda_1.$$

and

$$\langle \gamma, \nu \rangle = \beta^n - \beta^k \varphi_k \geq \frac{1}{3C_1},$$

where $\gamma = (\gamma^1, \gamma^2, \dots, \gamma^n)$ and $\nu = (0, 0, \dots, 0, 1)$ is the unit inward normal vector of $\partial T(\Omega_{m_0}) \cap \{z_n = 0\}$. This implies that the equation (5.3) is uniformly elliptic with strict oblique boundary condition.

Set

$$\mathfrak{B}_r := \{(z', z_n) : z' \in B'_r(0), 0 < z_n < m_0 - \varphi(z')\}.$$

Then it follows by Theorem 6.30 and Theorem 6.3.1 in [15], for $1 \leq s \leq n-1$, we deduce that

$$\frac{\partial w}{\partial z_s} =: v \in C^{2,\alpha}(\overline{\mathfrak{G}_r}).$$

Let $Y = \frac{\partial w}{\partial z_n}$. Then the differentiation of the equation (5.2) to z_n gives

$$a^{nn}Y_{nn} + a^{kn}(Y_{kn} - Y_{nn}\varphi_k) + a^{ln}(Y_{ln} - Y_{nn}\varphi_l) + a^{kl}(Y_{kl} - Y_{kn}\varphi_l - Y_{nl}\varphi_k + Y_{nn}\varphi_k\varphi_l) = \frac{\partial f}{\partial z_n}.$$

It reads as

$$(5.4) \quad (a^{nn} + a^{kl}\varphi_k\varphi_l - a^{kn}\varphi_k - a^{ln}\varphi_l)Y_{nn} = R.H.S.,$$

where

$$R.H.S := -a^{kn}Y_{kn} - a^{ln}Y_{ln} - a^{kl}Y_{kl} + a^{kl}Y_{kn}\varphi_l + a^{kl}Y_{nl}\varphi_k + \frac{\partial f}{\partial z_n}.$$

By $v = \frac{\partial w}{\partial z_s} \in C^{2,\alpha}(\overline{\mathfrak{G}_r})$, we have $R.H.S \in C^\alpha(\overline{\mathfrak{G}_r})$.

Recalling the proof of $3n\Lambda_2 \geq [A^{ij}] \geq \frac{1}{3n}\Lambda_1$ and taking $r > 0$ small enough, we can also obtain

$$a^{nn} + a^{kl}\varphi_k\varphi_l - a^{kn}\varphi_k - a^{ln}\varphi_l \geq \frac{1}{3n}\Lambda_1.$$

Then by (5.4) and $(a^{nn} + a^{kl}\varphi_k\varphi_l - a^{kn}\varphi_k - a^{ln}\varphi_l) \in C^\alpha(\overline{\mathfrak{G}_r})$, we see that

$$Y_{nn} \in C^\alpha(\overline{\mathfrak{G}_r})$$

and therefore

$$w \in C^{3,\alpha}(\overline{\mathfrak{G}_r}).$$

Consequently, $u \in C^{3,\alpha}(\overline{\Omega}_{m_0})$.

By Finite Covering Theorem, we get $u \in C^{3,\alpha}(\overline{\Omega})$. Using bootstrap argument, we obtain $u \in C^\infty(\overline{\Omega})$. \square

Now, by the continuity method, we can show that:

Proof of Theorem 2.3. For each $t \in [0, 1]$, consider

$$(5.5) \quad \begin{cases} F[D^2u] = tf(x) + c(t), & x \in \Omega, \\ Du(\Omega) = \tilde{\Omega}, \end{cases}$$

which is equivalent to

$$(5.6) \quad \begin{cases} F[D^2u] = tf(x) + c(t), & x \in \Omega, \\ h(Du) = 0, & x \in \partial\Omega. \end{cases}$$

By the main results in [11] and [12], (5.5) is solvable if $t = 0$. By Lemma 5.2, denote the closed subset

$$X := \{u \in C^{2,\alpha}(\overline{\Omega}) : \int_{\Omega} u = 0\}$$

in $C^{2,\alpha}(\overline{\Omega})$ and

$$Y := C^\alpha(\overline{\Omega}) \times C^{1,\alpha}(\partial\Omega).$$

Define a map from $X \times \mathbb{R}$ to Y as

$$\mathfrak{F}^t := (F(D^2u) - tf(x) - c(t), h(Du)).$$

Then the linearized operator $D\mathfrak{F}_{(u,c)}^t : X \times \mathbb{R} \rightarrow Y$ is given by

$$D\mathfrak{F}_{(u,c)}^t(w, a) = (F^{ij}(D^2u)\partial_{ij}w - a, h_{p_i}(Du)\partial_iw).$$

Repeating the proof of Proposition 3.1 in [9], we know that $D\mathfrak{F}_{(u,c)}^t$ is invertible for any $t \in [0, 1]$ and (u, c) being the solution to (5.5).

Define the set

$$I := \{t \in [0, 1] : (5.5) \text{ has at least one convex solution}\}.$$

Since $0 \in I$, by Huang-Ou's theorem [11] or Huang-Ye's theorem [12], I is not empty. We claim that $I = [0, 1]$, which is equivalent to the fact that I is not only open, but also closed. It follows from Proposition 3.1 in [9] again and Theorem 17.6 in [15] that I is open. So we only need to prove that I is a closed subset of $[0, 1]$.

That I is closed is equivalent to the fact that for any sequence $\{t_k\} \subset I$, if $\lim_{k \rightarrow \infty} t_k = t_0$, then $t_0 \in I$. For t_k , denote $(u_k, c(t_k))$ solving

$$\begin{cases} F[D^2u_k] = t_k f(x) + c(t_k), & x \in \Omega, \\ Du_k(\Omega) = \tilde{\Omega}. \end{cases}$$

It follows from Lemma 4.12 that $\|u_k\|_{C^{2,\alpha}(\bar{\Omega})} \leq C$, where C is independent of t_k . Since

$$|c(t)| = |F[D^2u] - tf(x)| \leq F(+\infty, \dots, +\infty) + \max_{\bar{\Omega}} |f(x)|,$$

by Arzela-Ascoli Theorem we know that there exists $\hat{u} \in C^{2,\alpha}(\bar{\Omega})$, $\hat{c} \in \mathbb{R}$ and a subsequence of $\{t_k\}$, which is still denoted as $\{t_k\}$, such that letting $k \rightarrow \infty$,

$$\begin{cases} \|u_k - \hat{u}\|_{C^2(\bar{\Omega})} \rightarrow 0, \\ c(t_k) \rightarrow \hat{c}. \end{cases}$$

Since $(u_k, c(t_k))$ satisfies

$$\begin{cases} F[D^2u_k] = t_k f(x) + c(t_k), & x \in \Omega, \\ h(Du_k) = 0, & x \in \partial\Omega, \end{cases}$$

letting $k \rightarrow \infty$, we have

$$\begin{cases} F[D^2\hat{u}] = t_0 f(x) + \hat{c}, & x \in \Omega, \\ h(D\hat{u}) = 0, & x \in \partial\Omega. \end{cases}$$

Therefore, $t_0 \in I$, and thus I is closed. Consequently, $I = [0, 1]$. By Lemma 5.1 we know that the solution of (5.5) u is unique up to a constant.

Then we complete the proof of Theorem 2.3. \square

ACKNOWLEDGEMENTS

The authors would like to express deep gratitude to Professor Yuanlong Xin for his suggestions and constant encouragement.

REFERENCES

- [1] M. Warren, *Calibrations associated to Monge-Ampère equations*, Trans. Amer. Math. Soc. **362** (2010), 3947–3962.
- [2] R. Harvey, H.B. Lawson, *Calibrated geometry*. Acta Math. **148** (1982), 47–157.
- [3] J. Jost, Y.L. Xin, *A Bernstein theorem for special Lagrangian graphs*, Calc.Var.Partial Differential Equations. **15** (2002), 299–312.
- [4] Y. Yuan, *A Bernstein problem for special Lagrangian equations*, Invent. Math. **150** (2002), 117–125.
- [5] P. Delanoë, *Classical solvability in dimension two of the second boundary-value problem associated with the Monge-Ampère operator*, Ann. Inst. H. Poincaré Anal. Non Linéaire. **8** (1991), 443–457.
- [6] L. Caffarelli, *Boundary regularity of maps with convex potentials*, II, Ann. of Math. Stud. **144** (1996), 453–496.
- [7] J. Urbas, *On the second boundary value problems for equations of Monge-Ampère type*, J. Reine Angew. Math. **487** (1997), 115–124.
- [8] O.C. Schnürer, K. Smoczyk, *Neumann and second boundary value problems for Hessian and Gauss curvature flows*, Ann. Inst. H. Poincaré Anal. Non Linéaire. **20** (2003), 1043–1073.
- [9] S. Brendle, M. Warren, *A boundary value problem for minimal Lagrangian graphs*, J. Differential Geom. **84** (2010), 267–287.
- [10] R.L. Huang, *On the second boundary value problem for Lagrangian mean curvature flow*, J. Funct. Anal. **269** (2015), 1095–1114.
- [11] R.L. Huang, Q.Z. Ou, *On the second boundary value problem for a class of fully nonlinear equations*, J. Geom. Anal. **27** (2017), 2601–2617.
- [12] R.L. Huang, Y.H. Ye, *On the second boundary value problem for a class of fully nonlinear flows I*, to appear in International Mathematics Research Notices. (2017). DOI: 10.1093/imrn/rnx278.
- [13] J.J. Chen, R.L. Huang, Y.H. Ye, *On the second boundary value problem for a class of fully nonlinear flows II*, Archiv Der Mathematik. **111** (2018), 407–419.
- [14] J.G. Bao, J.Y. Chen, W. Zhang, *Prescribed mean curvature equation for Lagrangian graph*, manuscript.
- [15] D. Gilbarg, N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, 2nd ed., 1998.
- [16] G.M. Lieberman, N.S. Trudinger, *Nonlinear Oblique Boundary Value Problems for Nonlinear Elliptic Equations*, Trans. Amer. Math. Soc. **295** (1986), 509–546.