

# A DISPERSIVE ESTIMATE FOR THE MULTIDIMENSIONAL BURGERS EQUATION

LUIS SILVESTRE

ABSTRACT. We study the multi-dimensional Burgers equation  $u_t + uu_{x_1} + \dots + u^d u_{x_d} = 0$ . We prove that the  $L^\infty$  norm of entropy solutions of this equation decays polynomially as  $t \rightarrow \infty$  in terms of the  $L^1$  norm of the initial data only.

We consider entropy solutions of the equation

$$(1) \quad u_t + uu_{x_1} + \dots + u^d u_{x_d} = 0 \quad \text{for } t \in [0, \infty) \times x \in \mathbb{R}^d.$$

We prove the following estimate.

**Theorem 1.** *Let  $u_0 \in L^\infty \cap L^1(\mathbb{R}^d)$  and  $u \in L^\infty([0, \infty) \times \mathbb{R}^d) \cap C([0, \infty), L^1(\mathbb{R}^d))$  be the unique entropy solution to the equation (1) with initial data  $u_0$ . Then  $u$  satisfies the estimate*

$$\|u(t, \cdot)\|_{L^\infty(\mathbb{R}^d)} \leq C \|u_0\|_{L^1(\mathbb{R}^d)}^\gamma t^{-d\gamma_0},$$

where  $\gamma_0 = (1 + d(d+1)/2)^{-1}$ . The constant  $C$  depends on dimension only.

The multi-dimensional Burgers equation (1) is the canonical example of a multidimensional scalar conservation law that satisfies the genuine nonlinearity condition (originally stated in [2]). As such, it is often used as a test ground for regularity estimates for scalar conservation laws with the strongest possible genuine nonlinearity assumptions. See for example [1], [3] or [4].

Very recently, Denis Serre proved in [3] the following estimate for (1).

$$(2) \quad \|u(t, \cdot)\|_{L^{(d+1)^2/d}} \leq C \|u_0\|_{L^1}^\gamma t^{-\delta},$$

where

$$\gamma = \frac{d^2 + 2d + 2}{(1+d)(d^2 + d + 2)} \quad \text{and} \quad \delta = \frac{2d(d^2 + d + 1)}{(1+d)^2(d^2 + d + 2)}.$$

Note that the exponents in [3] are written in terms of the space-time dimension  $d+1$ . Here, we express them in terms of the spacial dimension and that is why the formulas look different. Using this estimate, it is explained in [3] how one can extend the notion of entropy solutions for unbounded initial data  $u_0 \in L^1$ .

The estimate (2) follows from Theorem 1 by interpolation since  $\|u(t, \cdot)\|_{L^1} \leq \|u_0\|_{L^1}$ .

In [4], we obtained the following result for general scalar conservation laws.

**Theorem 2** (Theorem 1.5 in [4]). *Let  $u$  be the entropy solution of a genuinely nonlinear scalar conservation law  $u_t + a(u) \cdot \nabla u = 0$ . Then, there is an exponent  $\gamma_0$  depending on the nonlinearity  $a$  (see [4] for a specific expression) such that for all  $\gamma \in (0, \gamma_0)$ , there exists a constant  $C = C(\gamma, d, \|u_0\|_{L^\infty})$  such that*

$$\|u(t, \cdot)\|_{L^\infty} \leq C \|u_0\|_{L^1}^\gamma t^{-d\gamma}.$$

In the case of the multidimensional Burgers equation, a more explicit form of the estimate can be obtained by scaling. This is done in Remark 8.4 in [4]. We obtain

$$(3) \quad \|u(t, \cdot)\|_{L^\infty} \leq C(\gamma, d) \|u_0\|_{L^\infty}^{1-\gamma(1+d(d+1)/2)} \|u_0\|_{L^1}^\gamma t^{-d\gamma},$$

for any  $\gamma < \gamma_0 := (1 + d(d+1)/2)^{-1}$ .

The estimate (3) is not good enough to define entropy solutions with unbounded initial data as in [3] because of its dependence on  $\|u_0\|_{L^\infty}$  on the right hand side. This drawback is pointed out in the comments after the main theorem in [3] together with the fact that the endpoint exponent  $\gamma = \gamma_0$  is not reachable in (3) since  $C(d, \gamma)$  may not stay bounded as  $\gamma \rightarrow \gamma_0$ . In this short note, we show a quick argument to overcome

---

*Date:* August 6, 2018.

LS was partially supported by NSF grants DMS-1254332 and DMS-1764285.

both drawbacks and obtain Theorem 1. Our proof is based on the application of (3) in dyadic subintervals of the form  $(t/2^{j+1}, t/2^j)$  for  $j = 0, 1, 2, 3, \dots$ . Compounding all these inequalities, we obtain Theorem 1.

The advantage of the multidimensional Burgers equation (1) over general scalar conservation laws is that there is an explicit global two-parameter scaling that is used in Remark 8.4 in [4] to obtain (3). It is easy to work out a version of Theorem 1 whenever a similar scaling property works (for example in the case of monomial equations as in Section 2.1 in [3]). It is currently unclear whether there is any form of Theorem 1 that applies to more general scalar conservation laws. In particular, we do not know yet if the dependence on  $\|u_0\|_{L^\infty}$  can be removed from the right hand side in Theorem 2 or if the decay exponent  $\gamma = \gamma_0$  is reachable in general.

**Acknowledgment.** I thank Denis Serre for pointing out this problem to me. This work started as a result of an email exchange between the author and Denis Serre after the paper [3] was completed.

*Proof of Theorem 1.* Let  $\gamma$  be any exponent in the range  $(0, \gamma_0)$  so that (3) applies. Let us define  $\theta := 1 - \gamma(1 + d(d+1)/2)$ . Note that  $\theta \in (0, 1)$ .

Because of the semigroup property, we can apply the estimate (3) to estimate  $\|u(t, \cdot)\|_{L^\infty}$  in terms of the  $L^\infty$  and  $L^1$  norms of  $u(t_1, 0)$  at any earlier time  $t_1$ . In particular, for  $t_1 = t/2$  we obtain

$$\begin{aligned} \|u(t, \cdot)\|_{L^\infty} &\leq C \left(\frac{t}{2}\right)^{-d\gamma} \|u(t/2, \cdot)\|_{L^1}^\gamma \|u(t/2, \cdot)\|_{L^\infty}^\theta, \\ &\leq C \left(\frac{t}{2}\right)^{-d\gamma} \|u_0\|_{L^1}^\gamma \|u(t/2, \cdot)\|_{L^\infty}^\theta \end{aligned}$$

We reapply (3) to estimate  $\|u(t/2, \cdot)\|_{L^\infty}$  in terms of  $\|u(t/4, \cdot)\|_{L^\infty}$ . Then we iteratively apply (3) to estimate  $\|u(t/2^j, \cdot)\|_{L^\infty}$  in terms of  $\|u(t/2^{j+1}, \cdot)\|_{L^\infty}$ . In each step, we get

$$\|u(t/2^j, \cdot)\|_{L^\infty} \leq C t^{-d\gamma} 2^{(j+1)d\gamma} \|u_0\|_{L^1}^\gamma \|u(t/2^{j+1}, \cdot)\|_{L^\infty}^\theta$$

Compounding all these inequalities, after  $k+1$  iterations we are left with

$$\|u(t, \cdot)\|_{L^\infty} \leq C (\sum_{j=0}^k \theta^j) \|u_0\|_{L^1}^{(\gamma \sum_{j=0}^k \theta^j)} t^{-(d\gamma \sum_{j=0}^k \theta^j)} 2^{(d\gamma \sum_{j=0}^k (j+1)\theta^j)} \|u(t/2^{k+1}, \cdot)\|_{L^\infty}^{\theta^{k+1}}.$$

Since  $\theta \in (0, 1)$  and entropy solutions are bounded, we can pass to the limit  $k \rightarrow \infty$  and obtain

$$\begin{aligned} \|u(t, \cdot)\|_{L^\infty} &\leq C^{1/(1-\theta)} \|u_0\|_{L^1}^{\gamma/(1-\theta)} t^{-d\gamma/(1-\theta)} 2^{d\gamma/(1-\theta)^2}, \\ &\leq \tilde{C} \|u_0\|_{L^1(\mathbb{R}^d)}^{\gamma_0} t^{-d\gamma_0}. \end{aligned}$$

□

## REFERENCES

- [1] Gianluca Crippa, Felix Otto, and Michael Westdickenberg. Regularizing effect of nonlinearity in multidimensional scalar conservation laws. In *Transport equations and multi-D hyperbolic conservation laws*, pages 77–128. Springer, 2008.
- [2] P-L Lions, Benoit Perthame, and Eitan Tadmor. A kinetic formulation of multidimensional scalar conservation laws and related equations. *Journal of the American Mathematical Society*, 7(1):169–191, 1994.
- [3] Denis Serre. Multi-dimensional scalar conservation laws with unbounded integrable initial data. *arXiv preprint arXiv:1807.10474*, 2018.
- [4] Luis Silvestre. Oscillation properties of scalar conservation laws. *arXiv preprint arXiv:1708.03401*, 2017.