

# Chevalley formula for anti-dominant weights in the equivariant $K$ -theory of semi-infinite flag manifolds\*

Satoshi Naito

Department of Mathematics, Tokyo Institute of Technology,  
2-12-1 Oh-okayama, Meguro-ku, Tokyo 152-8551, Japan  
(e-mail: [naito@math.titech.ac.jp](mailto:naito@math.titech.ac.jp))

Daniel Orr

Department of Mathematics (MC 0123), 460 McBryde Hall, Virginia Tech,  
225 Stanger St., Blacksburg, VA 24061, U. S. A. (e-mail: [dorr@vt.edu](mailto:dorr@vt.edu))

Daisuke Sagaki

Institute of Mathematics, University of Tsukuba,  
1-1-1 Tennodai, Tsukuba, Ibaraki 305-8571, Japan  
(e-mail: [sagaki@math.tsukuba.ac.jp](mailto:sagaki@math.tsukuba.ac.jp))

## Abstract

We prove a Chevalley formula for anti-dominant weights in the torus-equivariant  $K$ -group of semi-infinite flag manifolds, which is described explicitly in terms of semi-infinite Lakshmibai-Seshadri paths (or, equivalently, quantum Lakshmibai-Seshadri paths). Based on the isomorphism, recently established by Kato, between the (small) torus-equivariant quantum  $K$ -group of finite-dimensional flag manifolds and the torus-equivariant  $K$ -group of semi-infinite flag manifolds, these formulas give an explicit description of the structure constants for the quantum multiplication in the quantum  $K$ -group of finite-dimensional flag manifolds. Our proof of these formulas is based on standard monomial theory for semi-infinite Lakshmibai-Seshadri paths, which is established in our previous work, and also uses a string property of Demazure-like subsets of the crystal basis of a level-zero extremal weight module over a quantum affine algebra.

---

\*Key words and phrases: Chevalley formula, Monk formula, semi-infinite LS path, semi-infinite flag manifold, (quantum) Schubert calculus.

Mathematics Subject Classification 2010: Primary 17B37; Secondary 14N15, 14M15, 33D52, 81R10.

# 1 Introduction.

Let  $\mathbf{Q}_G^{\text{rat}}$  denote the (whole) semi-infinite flag manifold, which is the reduced ind-scheme whose set of  $\mathbb{C}$ -valued points is  $G(\mathbb{C}((z)))/(H \cdot N(\mathbb{C}((z)))$  (see [Kat2] for details), where  $G$  is a simply-connected simple algebraic group over  $\mathbb{C}$  with Borel subgroup  $B = HN$ ,  $H$  maximal torus and  $N$  unipotent radical. In this paper, we concentrate on the semi-infinite Schubert (sub)variety  $\mathbf{Q}_G := \mathbf{Q}_G(e) \subset \mathbf{Q}_G^{\text{rat}}$  associated to the identity element  $e$  of the affine Weyl group  $W_{\text{af}} = W \ltimes Q^\vee$ , with  $W = \langle s_i \mid i \in I \rangle$  the (finite) Weyl group and  $Q^\vee = \bigoplus_{i \in I} \mathbb{Z}\alpha_i^\vee$  the coroot lattice of  $G$ ; we also call  $\mathbf{Q}_G$  the semi-infinite flag manifold. In [KNS], we proposed a definition of an equivariant (with respect to the Iwahori subgroup)  $K$ -group of  $\mathbf{Q}_G$ , and also proved a Chevalley formula for dominant weights in this  $K$ -group; this formula describes, in terms of semi-infinite Lakshmibai-Seshadri (LS for short) paths, the tensor product of the class of the line bundle  $[\mathcal{O}_{\mathbf{Q}_G}(\lambda)]$  associated to a dominant weight  $\lambda$  with the class of the structure sheaf  $[\mathcal{O}_{\mathbf{Q}_G(x)}]$  of a semi-infinite Schubert (sub)variety  $\mathbf{Q}_G(x) \subset \mathbf{Q}_G$  for  $x = wt_\xi \in W_{\text{af}}^{\geq 0} := W \times Q^{\vee,+} \subset W_{\text{af}}$ , where  $Q^{\vee,+} := \sum_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i^\vee \subset Q^\vee$ ; note that the class of the structure sheaf  $[\mathcal{O}_{\mathbf{Q}_G(x)}]$  in this paper is denoted by  $[\overline{\mathcal{O}}_x]$  in [O]. The purpose of this paper is to give a Chevalley formula for anti-dominant weights in the  $H \times \mathbb{C}^*$ -equivariant  $K$ -group  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  of  $\mathbf{Q}_G$ , and also in an  $H$ -equivariant  $K$ -group  $K'_H(\mathbf{Q}_G)$ .

A breakthrough in the study of the equivariant  $K$ -group of  $\mathbf{Q}_G$  was achieved in [Kat1] and [Kat3], in which Kato established a  $\mathbb{C}[P]$ -module isomorphism from the (small)  $H$ -equivariant quantum  $K$ -group  $QK_H(G/B) := K_H(G/B) \otimes \mathbb{C}[Q^{\vee,+}]$  of the finite-dimensional flag manifold  $G/B$  onto an  $H$ -equivariant  $K$ -group  $K'_H(\mathbf{Q}_G)$  of  $\mathbf{Q}_G$ , where  $\mathbb{Z}[P] \subset \mathbb{C}[P]$  is the group ring of the weight lattice  $P = \bigoplus_{i \in I} \mathbb{Z}\varpi_i$  of  $G$ , identified with the representation ring of  $H$ , and  $\mathbb{C}[Q^{\vee,+}]$  is the polynomial (not formal power series) ring in the (Novikov) variables  $Q_i$ ,  $i \in I$ ; also, it is noteworthy that Kato verified conjectures in [L<sup>2</sup>MS] about the relationship between  $QK_H(G/B)$  and the (suitably localized)  $H$ -equivariant  $K$ -group of the affine Grassmannian associated to  $G$ . This  $\mathbb{C}[P]$ -module isomorphism sends the (opposite) Schubert class  $[\mathcal{O}^w]$  in  $QK_H(G/B)$  to the corresponding semi-infinite Schubert class  $[\mathcal{O}_{\mathbf{Q}_G(w)}]$  in  $K'_H(\mathbf{Q}_G)$  for each  $w \in W$ . Moreover, it respects the quantum multiplication  $\star$  in  $QK_H(G/B)$  and the tensor product in  $K'_H(\mathbf{Q}_G)$ ; to be more precise, it respects the quantum multiplication  $\star$  with the class of the line bundle  $[\mathcal{O}_{G/B}(-\varpi_i)]$  and the tensor product  $\otimes$  with the class of the line bundle  $[\mathcal{O}_{\mathbf{Q}_G}(w_\circ\varpi_i)]$  for each  $i \in I$ , where  $w_\circ$  is the longest element of the finite Weyl group  $W$ . Namely, the following diagram commutes:

$$\begin{array}{ccc} QK_H(G/B) & \xrightarrow{\cong} & K'_H(\mathbf{Q}_G) \\ \cdot \star [\mathcal{O}_{G/B}(-\varpi_i)] \downarrow & & \downarrow \cdot \otimes [\mathcal{O}_{\mathbf{Q}_G}(w_\circ\varpi_i)] \\ QK_H(G/B) & \xrightarrow[\cong]{} & K'_H(\mathbf{Q}_G); \end{array}$$

here we warn the reader that the convention in [Kat1] differs from that in [KNS] and in this paper by the twist coming from the involution  $-w_\circ$ , and the line bundles  $\mathcal{O}_{\mathbf{Q}_G}(\mu)$  for  $\mu \in P$  are normalized in such a way that the quantum multiplication with  $[\mathcal{O}_{G/B}(-\varpi_i)]$  corresponds to the tensor product with  $[\mathcal{O}_{\mathbf{Q}_G}(-\varpi_i)]$  under the  $\mathbb{C}[P]$ -module isomorphism

above. Note that the quantum multiplicative structure of  $QK_H(G/B)$  is determined by the quantum multiplication with the class of the line bundles  $[\mathcal{O}_{\mathbf{Q}_G}(-\varpi_i)]$ ,  $i \in I$ , and the  $\mathbb{C}[P]$ -module structure of  $QK_H(G/B)$ ; for the  $\mathbb{C}[P]$ -module structure, see [O].

The  $H \times \mathbb{C}^*$ -equivariant  $K$ -group  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  has a topological  $\mathbb{C}[q, q^{-1}][P]$ -basis of the semi-infinite Schubert classes  $[\mathcal{O}_{\mathbf{Q}_G(wt_\xi)}]$  for  $w \in W$  and  $\xi \in Q^{\vee,+} = \sum_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i^\vee \subset Q^\vee$ ; the  $H$ -equivariant  $K$ -group  $K'_H(\mathbf{Q}_G)$  above is, by definition, obtained by the specialization  $q = 1$  from the  $\mathbb{C}[q, q^{-1}][P]$ -submodule  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G) \subset K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  consisting of all finite  $\mathbb{C}[q, q^{-1}][P]$ -linear combinations of these semi-infinite Schubert classes. Hence a Chevalley formula for anti-dominant weights in  $K'_H(\mathbf{Q}_G)$  (and, in particular, a Chevalley formula for the quantum multiplication with  $[\mathcal{O}_{G/B}(-\varpi_i)]$ ,  $i \in I$ , in  $QK_T(G/B)$ ) is obtained from our Chevalley formula in  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  by the specialization  $q = 1$ .

Let us state our Chevalley formula more precisely. Denote by  $P^+ := \sum_{i \in I} \mathbb{Z}_{\geq 0} \varpi_i$  the set of dominant integral weights. For a dominant weight  $\mu \in P^+$ , we set  $J_\mu := \{i \in I \mid \langle \mu, \alpha_i^\vee \rangle = 0\}$ ; we denote by  $W^{J_\mu}$  the set of minimal-length coset representatives for the cosets in  $W/W_{J_\mu}$ , where  $W_{J_\mu} := \langle s_i \mid i \in J_\mu \rangle$  is the stabilizer of  $\mu$  in  $W$ . Let  $\text{QLS}(\mu)$  denote the set of quantum LS paths of shape  $\mu$ . Recall that an element  $\eta \in \text{QLS}(\mu)$  is a sequence of elements of  $W^{J_\mu}$  that satisfies a certain condition described in terms of the parabolic quantum Bruhat graph (see Definition 3.11), and we can endow the set  $\text{QLS}(\mu)$  with a crystal structure with weights in  $P$ . For  $\eta \in \text{QLS}(\mu)$  and  $v \in W$ , we define an element  $\kappa(\eta, v) \in W$  (called the final direction of  $\eta$  with respect to  $v$ ) and an element  $\zeta(\eta, v) \in Q^{\vee,+}$  in terms of the quantum version of the Deodhar lifts introduced in [LNS<sup>3</sup>1] and the (quantum) weight of a directed path in the quantum Bruhat graph (see (3.27) and (3.28)). Also, we denote by  $\text{deg} : \text{QLS}(\mu) \rightarrow \mathbb{Z}_{\leq 0}$  the (tail) degree function on  $\text{QLS}(\mu)$  introduced in [NS2] and [LNS<sup>3</sup>2], which is described in terms of the parabolic quantum Bruhat graph (see (3.26)).

One of the main results of this paper is the following Chevalley formula for anti-dominant weights in the  $H \times \mathbb{C}^*$ -equivariant  $K$ -group  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$ ; in fact, this is a formula in  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  because of its significant finiteness.

**Theorem 1** (Chevalley formula in  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G) \subset K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$ ). *Let  $\lambda \in P^+$  be a dominant integral weight, and let  $x = wt_\xi \in W_{\text{af}}^{\geq 0}$ . Then, in the  $H \times \mathbb{C}^*$ -equivariant  $K$ -group  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G) \subset K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$ , we have*

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G}(-\lambda)] \otimes [\mathcal{O}_{\mathbf{Q}_G(x)}] \\ &= \sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(-w_\circ \lambda) \\ \kappa(\eta, v) = w}} (-1)^{\ell(v) - \ell(w)} q^{-\text{deg}(\eta) + \langle -w_\circ \lambda, \xi \rangle} e^{-\text{wt}(\eta)} [\mathcal{O}_{\mathbf{Q}_G(vt_\xi + \zeta(\eta, v))}]. \end{aligned} \quad (1.1)$$

We remark that the sum on the right-hand side of (1.1) is clearly a finite sum, since  $W$  is a finite Weyl group and  $\text{QLS}(-w_\circ \lambda)$  is a finite set; in fact, the set  $\text{QLS}(-w_\circ \lambda)$  provides a realization of the crystal basis of the quantum version of a local Weyl module (which is finite-dimensional). This finiteness assertion plays an important role in the proof of the existence of an isomorphism between  $QK_H(G/B)$  and  $K'_H(\mathbf{Q}_G)$  (see [Kat3] and also [ACT]).

Now we explain how to prove our Chevalley formula for anti-dominant weights in  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$ . Denote by  $K_{H \times \mathbb{C}^*}^\dagger(\mathbf{Q}_G)$  the  $\mathbb{C}[q, q^{-1}][P]$ -module consisting of all (possibly infinite) linear combinations of the semi-infinite Schubert classes  $[\mathcal{O}_{\mathbf{Q}_G(x)}]$  with coefficients  $a_x \in \mathbb{C}[q, q^{-1}][P]$  for  $x \in W_{\text{af}}^{\geq 0} = W_{\text{af}} \times Q^{\vee,+}$  such that the sum  $\sum_{x \in W_{\text{af}}^{\geq 0}} |a_x|$  of the absolute values  $|a_x|$  (where the absolute value of the coefficient of each monomial appearing in  $a_x$  is taken) lie in  $\mathbb{C}((q^{-1})[P])$ . Also, let  $\text{Fun}_P(\mathbb{C}((q^{-1})[P]))$  denote the  $\mathbb{C}[q, q^{-1}][P]$ -module of all functions on  $P$  with values in  $\mathbb{C}((q^{-1})[P])$ , and set

$$\text{Fun}_P^{\text{ess}}(\mathbb{C}((q^{-1})[P])) := \text{Fun}_P(\mathbb{C}((q^{-1})[P])) / \text{Fun}_P^{\text{neg}}(\mathbb{C}((q^{-1})[P])),$$

where  $\text{Fun}_P^{\text{neg}}(\mathbb{C}((q^{-1})[P]))$  is the  $\mathbb{C}[q, q^{-1}][P]$ -submodule of  $\text{Fun}_P(\mathbb{C}((q^{-1})[P]))$  consisting of those  $f \in \text{Fun}_P(\mathbb{C}((q^{-1})[P]))$  such that there exists some  $\gamma \in P$  for which  $f(\mu) = 0$  for all  $\mu \in \gamma + P^+$ , with  $P^+ := \sum_{i \in I} \mathbb{Z}_{\geq 0} \varpi_i$ . Then, for each  $x \in W_{\text{af}}^{\geq 0}$ , the assignment

$$P \ni \mu \mapsto \text{gch } H^0(\mathbf{Q}_G, \mathcal{O}_{\mathbf{Q}_G(x)} \otimes \mathcal{O}_{\mathbf{Q}_G}(\mu)) \in \mathbb{C}((q^{-1})[P])$$

defines an element of  $\text{Fun}_P^{\text{ess}}(\mathbb{C}((q^{-1})[P]))$ , which we denote by  $f^x(\cdot)$ ; here we denote by  $\text{gch } H^0(\mathbf{Q}_G, \mathcal{O}_{\mathbf{Q}_G(x)} \otimes \mathcal{O}_{\mathbf{Q}_G}(\mu))$  the graded character of the  $H \times \mathbb{C}^*$ -module  $H^0(\mathbf{Q}_G, \mathcal{O}_{\mathbf{Q}_G(x)} \otimes \mathcal{O}_{\mathbf{Q}_G}(\mu))$ , which is (as proved in [KNS]) identical to the graded character of the Demazure submodule  $V_x^-(-w_\circ \mu)$  of the level-zero extremal weight module  $V(-w_\circ \mu)$  over the quantum affine algebra  $U_v(\mathfrak{g}_{\text{af}})$  if  $\mu \in P^+$ , and is zero if  $\mu \notin P^+$ , where  $\mathfrak{g}_{\text{af}}$  is the untwisted affine Lie algebra whose underlying finite-dimensional simple Lie algebra is the Lie algebra  $\mathfrak{g}$  of  $G$ . Thus we obtain a  $\mathbb{C}[q, q^{-1}][P]$ -linear map

$$\Phi^\dagger : K_{H \times \mathbb{C}^*}^\dagger(\mathbf{Q}_G) \rightarrow \text{Fun}_P^{\text{ess}}(\mathbb{C}((q^{-1})[P]))$$

given by  $\Phi^\dagger([\mathcal{O}_{\mathbf{Q}_G(x)}]) = f^x(\cdot)$  for each  $x \in W_{\text{af}}^{\geq 0}$ . Following [Kat3], we define an  $H \times \mathbb{C}^*$ -equivariant  $K$ -group  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  to be the quotient  $\mathbb{C}[q, q^{-1}][P]$ -module  $K_{H \times \mathbb{C}^*}^\dagger(\mathbf{Q}_G) / \text{Ker}(\Phi^\dagger)$ ; also, we denote by  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  the  $\mathbb{C}[q, q^{-1}][P]$ -submodule of  $K_{H \times \mathbb{C}^*}^\dagger(\mathbf{Q}_G)$  consisting of all finite  $\mathbb{C}[q, q^{-1}][P]$ -linear combinations of the semi-infinite Schubert classes  $[\mathcal{O}_{\mathbf{Q}_G(x)}]$ ,  $x \in W_{\text{af}}^{\geq 0}$ . Because the graded characters  $\text{gch } V_x^-(-w_\circ \mu)$ ,  $x \in W_{\text{af}}^{\geq 0}$ , are linearly independent over  $\mathbb{C}[q, q^{-1}][P]$  when they are regarded as functions of sufficiently large  $\mu \in P^+$ , the  $\mathbb{C}[q, q^{-1}][P]$ -linear map  $\Phi^\dagger$  restricted to  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  is injective, and hence we have an embedding of  $\mathbb{C}[q, q^{-1}][P]$ -modules:

$$K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G) \hookrightarrow K_{H \times \mathbb{C}^*}(\mathbf{Q}_G),$$

which is compatible with the  $\mathbb{C}[q, q^{-1}][P]$ -linear injective map

$$\Phi : K_{H \times \mathbb{C}^*}(\mathbf{Q}_G) \rightarrow \text{Fun}_P^{\text{ess}}(\mathbb{C}((q^{-1})[P]))$$

induced by the map  $\Phi^\dagger$  above.

From the explicit identities obtained in [KNS] (in the case of dominant weights) and in this paper (in the case of anti-dominant weights) for the graded characters of Demazure submodules of level-zero extremal weight modules, it can be shown (see [Kat3]) that

there exist  $\mathbb{C}[q, q^{-1}][P]$ -module endomorphisms  $\Xi(\lambda)$ ,  $\lambda \in P$ , of  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  such that  $\Xi(\lambda + \lambda') = \Xi(\lambda) \circ \Xi(\lambda')$  for  $\lambda, \lambda' \in P$  and such that the following diagram commutes for all  $\lambda \in P$ :

$$\begin{array}{ccc} K_{H \times \mathbb{C}^*}(\mathbf{Q}_G) & \xrightarrow{\Phi} & \text{Fun}_P^{\text{ess}}(\mathbb{C}((q^{-1}))[P]) \\ \Xi(\lambda) \downarrow & & \downarrow \Theta(\lambda) \\ K_{H \times \mathbb{C}^*}(\mathbf{Q}_G) & \xrightarrow[\Phi]{} & \text{Fun}_P^{\text{ess}}(\mathbb{C}((q^{-1}))[P]), \end{array}$$

where  $\Theta(\lambda)f(\cdot) = f(\cdot + \lambda)$  for  $f(\cdot) \in \text{Fun}_P^{\text{ess}}(\mathbb{C}((q^{-1}))[P])$ ; the  $\mathbb{C}[q, q^{-1}][P]$ -module endomorphism  $\Xi(\lambda)$  can be thought of as the tensor product  $\cdot \otimes [\mathcal{O}_{\mathbf{Q}_G}(\lambda)]$  with the class of the line bundle  $[\mathcal{O}_{\mathbf{Q}_G}(\lambda)]$  in  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  (see [KNS]). In view of the commutativity of the diagram above and the injectivity of the  $\mathbb{C}[q, q^{-1}][P]$ -linear map  $\Phi$ , the proof of our Chevalley formula for anti-dominant weights in  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  is reduced to the proof of the corresponding identity in Corollary 4 below for the graded characters of Demazure submodules of level-zero extremal weight modules; observe that the submodule  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G) \subset K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  is stabilized by the endomorphisms  $\Xi(-\lambda)$ ,  $\lambda \in P^+$ , because of the finiteness of the sum on the right-hand side of equation (1.6) in Corollary 4. Indeed, since the left-hand side of (1.1) can be written as  $\Xi(-\lambda)([\mathcal{O}_{\mathbf{Q}_G(x)}])$ , its image under  $\Phi$  is identical to  $\Theta(-\lambda)(\Phi([\mathcal{O}_{\mathbf{Q}_G(x)}]))$  by the commutativity of the diagram above; by the definitions, this is identical to the graded character  $\text{gch } V_x^-(-w_\circ(\mu - \lambda))$  (regarded as a function of  $\mu \in P$ ) if  $\mu - \lambda \in P^+$ , and is zero if  $\mu - \lambda \notin P^+$ . Also, the image under  $\Phi$  of the right-hand side of (1.1) is identical to:

$$\sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(-w_\circ \lambda) \\ \kappa(\eta) = w}} (-1)^{\ell(v) - \ell(w)} q^{-\deg(\eta) + \langle -w_\circ \lambda, \xi \rangle} \mathbf{e}^{-\text{wt}(\eta)} \text{gch } V_{vt_\xi + \zeta(\eta, v)}^-(-w_\circ \mu)$$

(regarded as a function of  $\mu \in P$ ) if  $\mu \in P^+$ ; note that this turns out to be zero if  $\mu - \lambda \notin P^+$ , as shown in Appendix B. Because these two functions of  $\mu$  coincide if  $\lambda, \mu \in P^+$  are such that  $\mu - \lambda \in P^+$  by Corollary 4 below, we deduce the equality (1.1) in  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  from the injectivity of the  $\mathbb{C}[q, q^{-1}][P]$ -linear map  $\Phi$ .

Our second main result of this paper is a Chevalley formula for anti-dominant weights in the torus-equivariant  $K$ -group  $K'_H(\mathbf{Q}_G)$ , which is obtained from  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G) \subset K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$  by the specialization  $q = 1$  (of coefficients). Since the equality in Theorem 1 above is, in fact, the one in  $K'_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$ , it immediately implies the following:

**Theorem 2** (Chevalley formula in  $K'_H(\mathbf{Q}_G)$ ). *Let  $\lambda \in P^+$  be a dominant weight, and let  $x = wt_\xi \in W_{\text{af}}^{\geq 0}$ . Then, in the torus-equivariant  $K$ -group  $K'_H(\mathbf{Q}_G)$ , we have*

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G}(-\lambda)] \otimes [\mathcal{O}_{\mathbf{Q}_G(x)}] \\ &= \sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(-w_\circ \lambda) \\ \kappa(\eta, v) = w}} (-1)^{\ell(v) - \ell(w)} \mathbf{e}^{-\text{wt}(\eta)} [\mathcal{O}_{\mathbf{Q}_G(vt_\xi + \zeta(\eta, v))}]. \end{aligned} \quad (1.2)$$

By applying (1.2) to the case that  $\lambda = -w_\circ \varpi_i$ ,  $i \in I$ , and  $w = e$  (see §5.2), we deduce that  $[\mathcal{O}_{\mathbf{Q}_G(s_i)}] = [\mathcal{O}_{\mathbf{Q}_G(e)}] - \mathbf{e}^{\varpi_i} [\mathcal{O}_{\mathbf{Q}_G(w_\circ \varpi_i)}]$  (which agrees with [Kat1, Lemma 1.14]; recall

the difference of the conventions mentioned above). By combining this equality and (1.2), we obtain the following formula, which describes the tensor product in  $K'_H(\mathbf{Q}_G)$  by the class  $[\mathcal{O}_{\mathbf{Q}_G(s_i)}]$ :

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G(s_i)}] \otimes [\mathcal{O}_{\mathbf{Q}_G(w)}] \\ &= [\mathcal{O}_{\mathbf{Q}_G(w)}] + \sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(\varpi_i) \\ \kappa(\eta, v) = w}} (-1)^{\ell(v) - \ell(w) + 1} \mathbf{e}^{\varpi_i - \text{wt}(\eta)} [\mathcal{O}_{\mathbf{Q}_G(vt_{\zeta(\eta, v)})}] \end{aligned} \quad (1.3)$$

for  $i \in I$  and  $w \in W$ . In particular, if  $\varpi_i$  is a minuscule weight, then (see §3.6)

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G(s_i)}] \otimes [\mathcal{O}_{\mathbf{Q}_G(w)}] \\ &= [\mathcal{O}_{\mathbf{Q}_G(w)}] - \mathbf{e}^{\varpi_i - w\varpi_i} \sum_{\substack{v \in W \\ \max(wW_{I \setminus \{i\}}, \leq_v^*) = w}} (-1)^{\ell(v) - \ell(w)} [\mathcal{O}_{\mathbf{Q}_G(vt_{\text{wt}(w \Rightarrow v)})}], \end{aligned} \quad (1.4)$$

where  $\max(wW_{I \setminus \{i\}}, \leq_v^*)$  denotes the quantum version of the Deodhar lift (see Proposition 2.25) and  $\text{wt}(w \Rightarrow v)$  is the (quantum) weight of a directed path from  $w$  to  $v$  in the quantum Bruhat graph (see (2.21)). Based on the  $\mathbb{C}[P]$ -module isomorphism (established in [Kat1] and [Kat3]) from  $QK_H(G/B)$  onto  $K'_H(\mathbf{Q}_G)$ , which respects the quantum multiplication in  $QK_H(G/B)$  and the tensor product in  $K'_H(\mathbf{Q}_G)$  and which sends each (opposite) Schubert class in  $QK_H(G/B)$  to the corresponding semi-infinite Schubert class in  $K'_H(\mathbf{Q}_G)$ , we immediately obtain from formula (1.3) the corresponding formula for the quantum multiplication in  $QK_H(G/B)$ . Moreover, a conjectural Monk formula ([LeP, Conjecture 17.1]) in  $QK_T(G/B)$ , which is described in terms of the quantum alcove model, follows from this formula (see [LNS] for details). Also, for the torus-equivariant quantum  $K$ -group of partial flag manifolds of minuscule type, we can deduce (see [KoNS]) from formula (1.4) an explicit formula describing the quantum multiplication (cf. [BCMP] in the case of cominuscule type).

Our formula (1.2) can also be thought of as a semi-infinite analog of the corresponding formula in [LeS] in the torus-equivariant  $K$ -theory for Kac-Moody thick flag manifolds (see also [GR] for the finite-dimensional case), though our proof is quite different from the one by them and is much more difficult; this is mainly because an ordinary induction argument using a string property of Demazure-like subsets does not suffice in our case in contrast to the case of Kac-Moody thick flag manifolds.

Let us explain the representation-theoretic (or, crystal-theoretic) aspect of our main results; as mentioned above, Theorems 1 and 2 follow from Theorem 3 and Corollary 4 below, which are proved by using crystal bases of level-zero extremal weight modules. Let  $\mu \in P^+$  be a dominant integral weight. We denote by  $(W^{J\mu})_{\text{af}}$  the set of Peterson's coset representatives for the cosets in  $W_{\text{af}}/(W_{J\mu})_{\text{af}}$  (see §2.2), with  $\Pi^{J\mu} : W_{\text{af}} \rightarrow (W^{J\mu})_{\text{af}}$  the canonical projection. Let  $\mathbb{B}^{\infty}(\mu)$  denote the set of semi-infinite LS paths of shape  $\mu$ . Recall that an element  $\pi \in \mathbb{B}^{\infty}(\mu)$  is a certain decreasing sequence of elements of  $(W^{J\mu})_{\text{af}} \subset W_{\text{af}}$  in the semi-infinite Bruhat order  $\succeq$  (see Definition 3.2), and we can endow the set  $\mathbb{B}^{\infty}(\mu)$  with a crystal structure with weights in  $P_{\text{af}}^0 = P \oplus \mathbb{Z}\delta$ , where  $\delta$  is the (primitive) null root of the untwisted affine Lie algebra  $\mathfrak{g}_{\text{af}}$ . If we denote by  $\kappa(\pi) \in (W^{J\mu})_{\text{af}}$  the final direction

of  $\pi \in \mathbb{B}_{\frac{\infty}{2}}(\mu)$ , then we know from [NS3] that for each  $z \in W_{\text{af}}$ , the graded character  $\text{gch } V_z^-(\mu)$  of the Demazure submodule  $V_z^-(\mu)$  of the level-zero extremal weight module  $V(\mu)$  over  $U_{\mathfrak{v}}(\mathfrak{g}_{\text{af}})$  is identical to the following sum:

$$\sum_{\substack{\pi \in \mathbb{B}_{\frac{\infty}{2}}(\mu) \\ \kappa(\pi) \succeq z}} q^{\text{nul}(\text{wt}(\pi))} \mathbf{e}^{\text{fin}(\text{wt}(\pi))},$$

where for  $\nu \in P_{\text{af}}^0 = P \oplus \mathbb{Z}\delta$ , we write  $\nu = \text{fin}(\nu) + \text{nul}(\nu)\delta$ . Based on this fact, we make essential use of standard monomial theory for semi-infinite LS paths (established in [KNS]) to prove the following theorem (see Theorem 3.9); we also need a string property of Demazure-like subsets of the crystal basis of an extremal weight module (see Proposition 6.8).

**Theorem 3.** *Let  $\lambda, \mu \in P^+$  be dominant integral weights such that  $\mu - \lambda \in P^+$ , and set  $\lambda_i := \langle \lambda, \alpha_i^\vee \rangle \in \mathbb{Z}_{\geq 0}$ ,  $\mu_i := \langle \mu, \alpha_i^\vee \rangle \in \mathbb{Z}_{\geq 0}$  for  $i \in I$ ; note that  $\mu_i - \lambda_i \in \mathbb{Z}_{\geq 0}$  for all  $i \in I$ . Then, for  $x \in W_{\text{af}}$ , we have*

$$\begin{aligned} & \prod_{i \in I} \prod_{k=\mu_i-\lambda_i+1}^{\mu_i} \frac{1}{1-q^{-k}} \text{gch } V_x^-(\mu - \lambda) \\ &= \sum_{\substack{y \in W_{\text{af}} \\ y \succeq x}} \left( \sum_{\substack{\eta \in \mathbb{B}_{\frac{\infty}{2}}(\lambda) \\ \iota(\eta) \leq \Pi^{J\lambda}(y), \kappa(\eta, y) = x}} (-1)^{\ell_{\frac{\infty}{2}}(y) - \ell_{\frac{\infty}{2}}(x)} q^{-\text{nul}(\text{wt}(\eta))} \mathbf{e}^{-\text{fin}(\text{wt}(\eta))} \right) \text{gch } V_y^-(\mu), \end{aligned} \quad (1.5)$$

where  $\ell_{\frac{\infty}{2}} : W_{\text{af}} \rightarrow \mathbb{Z}$  denotes the semi-infinite length function (see Definition 2.2), and  $\kappa(\eta, y) \in W_{\text{af}}$  denotes the final direction of  $\eta$  with respect to  $y$  (see (3.15)).

The canonical projection  $\text{cl} : W_{\text{af}} = W \rtimes Q^\vee \twoheadrightarrow W$  induces a surjective map from  $\mathbb{B}_{\frac{\infty}{2}}(\mu)$  onto  $\text{QLS}(\mu)$ , which is also denoted by  $\text{cl}$ . By using this surjective map, we can reformulate this theorem in terms of the (parabolic) quantum Bruhat graph as follows (see Corollary 3.15).

**Corollary 4.** *Let  $\lambda, \mu \in P^+$  be dominant integral weights such that  $\mu - \lambda \in P^+$ . Then, for  $x = wt_\xi \in W_{\text{af}}$ , we have*

$$\begin{aligned} & \text{gch } V_x^-(\mu - \lambda) \\ &= \sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(\lambda) \\ \kappa(\eta, v) = w}} (-1)^{\ell(v) - \ell(w)} q^{-\text{deg}_\lambda(\eta) + \langle \lambda, \xi \rangle} \mathbf{e}^{-\text{wt}(\eta)} \text{gch } V_{vt_\xi + \zeta(\eta, v)}^-(\mu). \end{aligned} \quad (1.6)$$

Here we remark that in Theorem 3 and Corollary 4 above, the right-hand sides of the equations (1.5) and (1.6) turn out to be zero unless  $\mu - \lambda \in P^+$  (see Appendix B).

This paper is organized as follows. In Section 2, we fix our notation for untwisted affine Lie algebras, and then review some basic facts about the (parabolic) semi-infinite

Bruhat graph, the (parabolic) quantum Bruhat graph, and analogs of the Deodhar lifts for these graphs. In Section 3, after recalling the notions of semi-infinite LS paths and quantum LS paths, we state the formulas (Theorem 3 and Corollary 4) above for the graded characters of Demazure submodules, from which our main results (Theorem 1 and Theorem 2) immediately follow on the basis of the results established in [KNS]. In Section 4, we show some results in standard monomial theory for semi-infinite LS paths, which will be needed in the proof of Theorem 3. In Section 5, we prove a special case of Theorem 3, in which the dominant integral weight  $\lambda \in P^+$  is a fundamental weight and  $w = e$ , the identity element, by using standard monomial theory for semi-infinite LS paths. In Section 6, we show a string property of certain Demazure-like subsets of the crystal of semi-infinite LS paths of a given shape; using this, in Section 7, we prove Theorem 3 in the case that  $\lambda$  is a fundamental weight and  $w$  is an arbitrary element of  $W$ . In Section 8, we complete the proof of Theorem 3 by induction on the positive integer  $\sum_{i \in I} \lambda_i$ ; the base case that  $\lambda$  is a fundamental weight is already established in Section 7. Also, we reformulate Theorem 3 in terms of the quantum Bruhat graph as Corollary 4 above. In Appendix A, we give examples of formula (1.4) in the case that  $\mathfrak{g}$  is of type  $A_2$  and  $\lambda = \varpi_1$ . In Appendix B, we show that the right-hand side of equation (1.6) is indeed zero if  $\mu - \lambda \notin P^+$ . In Appendix C, we give a rephrasing of [KNS, Theorem 5.10] in terms of quantum LS paths.

**Acknowledgments.** We would like to thank Syu Kato for related collaborations. S.N. was partially supported by JSPS Grant-in-Aid for Scientific Research (B) 16H03920. D.O. was supported by NSF grant DMS-1600653. D.S. was partially supported by JSPS Grant-in-Aid for Scientific Research (C) 15K04803, (C) 19K03415.

## 2 Semi-infinite Bruhat order and quantum Bruhat graph.

**2.1 Affine Lie algebras.** Let  $\mathfrak{g}$  be a finite-dimensional simple Lie algebra over  $\mathbb{C}$  with Cartan subalgebra  $\mathfrak{h}$ . Denote by  $\{\alpha_i^\vee\}_{i \in I}$  and  $\{\alpha_i\}_{i \in I}$  the set of simple coroots and simple roots of  $\mathfrak{g}$ , respectively, and set  $Q := \bigoplus_{i \in I} \mathbb{Z}\alpha_i$ ,  $Q^+ := \sum_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i$ , and  $Q^\vee := \bigoplus_{i \in I} \mathbb{Z}\alpha_i^\vee$ ,  $Q^{\vee,+} := \sum_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i^\vee$ ; for  $\xi, \zeta \in Q^\vee$ , we write  $\xi \geq \zeta$  if  $\xi - \zeta \in Q^{\vee,+}$ . Let  $\Delta$ ,  $\Delta^+$ , and  $\Delta^-$  be the set of roots, positive roots, and negative roots of  $\mathfrak{g}$ , respectively, with  $\theta \in \Delta^+$  the highest root of  $\mathfrak{g}$ ; we set  $\rho := (1/2)\sum_{\alpha \in \Delta^+} \alpha$ . Also, let  $\varpi_i$ ,  $i \in I$ , denote the fundamental weights for  $\mathfrak{g}$ , and set

$$P := \bigoplus_{i \in I} \mathbb{Z}\varpi_i, \quad P^+ := \sum_{i \in I} \mathbb{Z}_{\geq 0}\varpi_i. \quad (2.1)$$

Let  $\mathfrak{g}_{\text{af}} = (\mathbb{C}[z, z^{-1}] \otimes \mathfrak{g}) \oplus \mathbb{C}c \oplus \mathbb{C}d$  be the (untwisted) affine Lie algebra over  $\mathbb{C}$  associated to  $\mathfrak{g}$ , where  $c$  is the canonical central element and  $d$  is the scaling element (or degree operator), with Cartan subalgebra  $\mathfrak{h}_{\text{af}} = \mathfrak{h} \oplus \mathbb{C}c \oplus \mathbb{C}d$ . We regard an element  $\mu \in \mathfrak{h}^* := \text{Hom}_{\mathbb{C}}(\mathfrak{h}, \mathbb{C})$  as an element of  $\mathfrak{h}_{\text{af}}^*$  by setting  $\langle \mu, c \rangle = \langle \mu, d \rangle := 0$ , where  $\langle \cdot, \cdot \rangle : \mathfrak{h}_{\text{af}}^* \times \mathfrak{h}_{\text{af}} \rightarrow \mathbb{C}$  denotes the canonical pairing of  $\mathfrak{h}_{\text{af}}^* := \text{Hom}_{\mathbb{C}}(\mathfrak{h}_{\text{af}}, \mathbb{C})$  and  $\mathfrak{h}_{\text{af}}$ . Let  $\{\alpha_i^\vee\}_{i \in I_{\text{af}}} \subset \mathfrak{h}_{\text{af}}^*$  and  $\{\alpha_i\}_{i \in I_{\text{af}}} \subset \mathfrak{h}_{\text{af}}$  be the set of simple coroots and simple roots of  $\mathfrak{g}_{\text{af}}$ ,



respectively, where  $I_{\text{af}} := I \sqcup \{0\}$ ; note that  $\langle \alpha_i, c \rangle = 0$  and  $\langle \alpha_i, d \rangle = \delta_{i,0}$  for  $i \in I_{\text{af}}$ . Denote by  $\delta \in \mathfrak{h}_{\text{af}}^*$  the null root of  $\mathfrak{g}_{\text{af}}$ ; recall that  $\alpha_0 = \delta - \theta$ . Also, let  $\Lambda_i \in \mathfrak{h}_{\text{af}}^*$ ,  $i \in I_{\text{af}}$ , denote the fundamental weights for  $\mathfrak{g}_{\text{af}}$  such that  $\langle \Lambda_i, d \rangle = 0$ , and set

$$P_{\text{af}} := \left( \bigoplus_{i \in I_{\text{af}}} \mathbb{Z} \Lambda_i \right) \oplus \mathbb{Z} \delta \subset \mathfrak{h}^*, \quad P_{\text{af}}^0 := \{ \mu \in P_{\text{af}} \mid \langle \mu, c \rangle = 0 \}; \quad (2.2)$$

notice that  $P_{\text{af}}^0 = P \oplus \mathbb{Z} \delta$ , and that  $\langle \mu, \alpha_0^\vee \rangle = -\langle \mu, \theta^\vee \rangle$  for  $\mu \in P_{\text{af}}^0$ . We remark that for each  $i \in I$ ,  $\varpi_i$  is identical to  $\Lambda_i - \langle \Lambda_i, c \rangle \Lambda_0$ , which is called the level-zero fundamental weight in [Kas3].

Let  $W := \langle s_i \mid i \in I \rangle$  and  $W_{\text{af}} := \langle s_i \mid i \in I_{\text{af}} \rangle$  be the (finite) Weyl group of  $\mathfrak{g}$  and the (affine) Weyl group of  $\mathfrak{g}_{\text{af}}$ , respectively, where  $s_i$  is the simple reflection with respect to  $\alpha_i$  for  $i \in I_{\text{af}}$ . We denote by  $l : W_{\text{af}} \rightarrow \mathbb{Z}_{\geq 0}$  the length function on  $W_{\text{af}}$ , whose restriction to  $W$  agrees with the one on  $W$ , by  $e \in W \subset W_{\text{af}}$  the identity element, and by  $w_\circ \in W$  the longest element. For each  $\xi \in Q^\vee$ , let  $t_\xi \in W_{\text{af}}$  denote the translation in  $\mathfrak{h}_{\text{af}}^*$  by  $\xi$  (see [Kac, Sect. 6.5]); for  $\xi \in Q^\vee$ , we have

$$t_\xi \mu = \mu - \langle \mu, \xi \rangle \delta \quad \text{if } \mu \in \mathfrak{h}_{\text{af}}^* \text{ satisfies } \langle \mu, c \rangle = 0. \quad (2.3)$$

Then,  $\{t_\xi \mid \xi \in Q^\vee\}$  forms an abelian normal subgroup of  $W_{\text{af}}$ , in which  $t_\xi t_\zeta = t_{\xi+\zeta}$  holds for  $\xi, \zeta \in Q^\vee$ . Moreover, we know from [Kac, Proposition 6.5] that

$$W_{\text{af}} \cong W \rtimes \{t_\xi \mid \xi \in Q^\vee\} \cong W \rtimes Q^\vee.$$

Denote by  $\Delta_{\text{af}}$  the set of real roots of  $\mathfrak{g}_{\text{af}}$ , and by  $\Delta_{\text{af}}^+ \subset \Delta_{\text{af}}$  the set of positive real roots; we know from [Kac, Proposition 6.3] that  $\Delta_{\text{af}} = \{\alpha + n\delta \mid \alpha \in \Delta, n \in \mathbb{Z}\}$  and  $\Delta_{\text{af}}^+ = \Delta^+ \sqcup \{\alpha + n\delta \mid \alpha \in \Delta, n \in \mathbb{Z}_{>0}\}$ . For  $\beta \in \Delta_{\text{af}}$ , we denote by  $\beta^\vee \in \mathfrak{h}_{\text{af}}$  its dual root, and by  $s_\beta \in W_{\text{af}}$  the corresponding reflection; if  $\beta \in \Delta_{\text{af}}$  is of the form  $\beta = \alpha + n\delta$  with  $\alpha \in \Delta$  and  $n \in \mathbb{Z}$ , then  $s_\beta = s_\alpha t_{n\alpha^\vee} \in W \rtimes Q^\vee$ .

Finally, let  $U_\vee(\mathfrak{g}_{\text{af}})$  (resp.,  $U'_\vee(\mathfrak{g}_{\text{af}})$ ) denote the quantized universal enveloping algebra over  $\mathbb{C}(\vee)$  associated to  $\mathfrak{g}_{\text{af}}$  (resp.,  $[\mathfrak{g}_{\text{af}}, \mathfrak{g}_{\text{af}}]$ ), with  $E_i$  and  $F_i$ ,  $i \in I_{\text{af}}$ , the Chevalley generators corresponding to  $\alpha_i$ . We denote by  $U_\vee^-(\mathfrak{g}_{\text{af}})$  the negative part of  $U_\vee(\mathfrak{g}_{\text{af}})$ , that is, the  $\mathbb{C}(\vee)$ -subalgebra of  $U_\vee(\mathfrak{g}_{\text{af}})$  generated by the  $F_i$ ,  $i \in I_{\text{af}}$ .

**2.2 Parabolic semi-infinite Bruhat graph.** In this subsection, we take and fix an arbitrary subset  $J \subset I$ . We set  $Q_J := \bigoplus_{i \in J} \mathbb{Z} \alpha_i$ ,  $Q_J^\vee := \bigoplus_{i \in J} \mathbb{Z} \alpha_i^\vee$ ,  $Q_J^{\vee,+} := \sum_{i \in J} \mathbb{Z}_{\geq 0} \alpha_i^\vee$ ,  $\Delta_J := \Delta \cap Q_J$ ,  $\Delta_J^\pm := \Delta^\pm \cap Q_J$ ,  $W_J := \langle s_i \mid i \in J \rangle$ , and  $\rho_J := (1/2) \sum_{\alpha \in \Delta_J^+} \alpha$ ; we denote by

$$[\cdot]^J : Q^\vee \twoheadrightarrow Q_{I \setminus J}^\vee \quad (\text{resp., } [\cdot]_J : Q^\vee \twoheadrightarrow Q_J^\vee) \quad (2.4)$$

the projection from  $Q^\vee = Q_{I \setminus J}^\vee \oplus Q_J^\vee$  onto  $Q_{I \setminus J}^\vee$  (resp.,  $Q_J^\vee$ ) with kernel  $Q_J^\vee$  (resp.,  $Q_{I \setminus J}^\vee$ ). Let  $W^J$  denote the set of minimal(-length) coset representatives for the cosets in  $W/W_J$ ; we know from [BB, Sect. 2.4] that

$$W^J = \{w \in W \mid w\alpha \in \Delta^+ \text{ for all } \alpha \in \Delta_J^+\}. \quad (2.5)$$

For  $w \in W$ , we denote by  $[w] = [w]^J \in W^J$  the minimal coset representative for the coset  $wW_J$  in  $W/W_J$ . Also, we set

$$(\Delta_J)_{\text{af}} := \{\alpha + n\delta \mid \alpha \in \Delta_J, n \in \mathbb{Z}\} \subset \Delta_{\text{af}}, \quad (2.6)$$

$$(\Delta_J)_{\text{af}}^+ := (\Delta_J)_{\text{af}} \cap \Delta_{\text{af}}^+ = \Delta_J^+ \sqcup \{\alpha + n\delta \mid \alpha \in \Delta_J, n \in \mathbb{Z}_{>0}\}, \quad (2.7)$$

$$(W_J)_{\text{af}} := W_J \times \{t_\xi \mid \xi \in Q_J^\vee\} = \langle s_\beta \mid \beta \in (\Delta_J)_{\text{af}}^+ \rangle, \quad (2.8)$$

$$(W^J)_{\text{af}} := \{x \in W_{\text{af}} \mid x\beta \in \Delta_{\text{af}}^+ \text{ for all } \beta \in (\Delta_J)_{\text{af}}^+\}; \quad (2.9)$$

if  $J = \emptyset$ , then  $(W^\emptyset)_{\text{af}} = W_{\text{af}}$  and  $(W_\emptyset)_{\text{af}} = \{e\}$ . We know from [P] (see also [LaS, Lemma 10.6]) that for each  $x \in W_{\text{af}}$ , there exist a unique  $x_1 \in (W^J)_{\text{af}}$  and a unique  $x_2 \in (W_J)_{\text{af}}$  such that  $x = x_1x_2$ ; let

$$\Pi^J : W_{\text{af}} \twoheadrightarrow (W^J)_{\text{af}}, \quad x \mapsto x_1, \quad (2.10)$$

denote the projection, where  $x = x_1x_2$  with  $x_1 \in (W^J)_{\text{af}}$  and  $x_2 \in (W_J)_{\text{af}}$ .

**Lemma 2.1** (see, e.g., [NS4, Lemma 2.1]).

(1) *It holds that*

$$\begin{cases} \Pi^J(w) = [w]^J & \text{for all } w \in W, \\ \Pi^J(xt_\xi) = \Pi^J(x)\Pi^J(t_\xi) & \text{for all } x \in W_{\text{af}} \text{ and } \xi \in Q_J^\vee; \end{cases} \quad (2.11)$$

*in particular,  $(W^J)_{\text{af}} = \{w\Pi^J(t_\xi) \mid w \in W^J, \xi \in Q_J^\vee\}$ .*

(2) *For each  $\xi \in Q_J^\vee$ , the element  $\Pi^J(t_\xi) \in (W^J)_{\text{af}}$  is of the form:  $\Pi^J(t_\xi) = ut_{\xi+\gamma}$  for some  $u \in W_J$  and  $\gamma \in Q_J^\vee$ .*

(3) *For  $\xi, \zeta \in Q_J^\vee$ ,  $\Pi^J(t_\xi) = \Pi^J(t_\zeta)$  if and only if  $\xi - \zeta \in Q_J^\vee$ .*

**Definition 2.2.** Let  $x \in W_{\text{af}}$ , and write it as  $x = wt_\xi$  with  $w \in W$  and  $\xi \in Q_J^\vee$ . We define the semi-infinite length  $\ell^{\frac{\infty}{2}}(x)$  of  $x$  by:  $\ell^{\frac{\infty}{2}}(x) = \ell(w) + 2\langle \rho, \xi \rangle$ .

**Definition 2.3.**

(1) The (parabolic) semi-infinite Bruhat graph  $\text{BG}^{\frac{\infty}{2}}((W^J)_{\text{af}})$  is the  $\Delta_{\text{af}}^+$ -labeled directed graph whose vertices are the elements of  $(W^J)_{\text{af}}$ , and whose directed edges are of the form:  $x \xrightarrow{\beta} y$  for  $x, y \in (W^J)_{\text{af}}$  and  $\beta \in \Delta_{\text{af}}^+$  such that  $y = s_\beta x$  and  $\ell^{\frac{\infty}{2}}(y) = \ell^{\frac{\infty}{2}}(x) + 1$ . When  $J = \emptyset$ , we write  $\text{BG}^{\frac{\infty}{2}}(W_{\text{af}})$  for  $\text{BG}^{\frac{\infty}{2}}((W^\emptyset)_{\text{af}})$ .

(2) The (parabolic) semi-infinite Bruhat order is a partial order  $\preceq$  on  $(W^J)_{\text{af}}$  defined as follows: for  $x, y \in (W^J)_{\text{af}}$ , we write  $x \preceq y$  if there exists a directed path in  $\text{BG}^{\frac{\infty}{2}}((W^J)_{\text{af}})$  from  $x$  to  $y$ ; we write  $x \prec y$  if  $x \preceq y$  and  $x \neq y$ .

*Remark 2.4.* On the (finite) Weyl group  $W$ , the semi-infinite Bruhat order agrees with the ordinary Bruhat order.

Let us recall some of the basic properties of the semi-infinite Bruhat order. Take and fix  $\lambda \in P^+$  such that  $\{i \in I \mid \langle \lambda, \alpha_i^\vee \rangle = 0\} = J$ .

**Lemma 2.5** ([INS, Remark 4.1.3]). *Let  $x \in (W^J)_{\text{af}}$  and  $i \in I_{\text{af}}$ . Then,*

$$s_i x \in (W^J)_{\text{af}} \iff \langle x\lambda, \alpha_i^\vee \rangle \neq 0 \iff x^{-1}\alpha_i \in (\Delta \setminus \Delta_J) + \mathbb{Z}\delta. \quad (2.12)$$

Moreover, in this case,

$$\begin{cases} x \xrightarrow{\alpha_i} s_i x \iff \langle x\lambda, \alpha_i^\vee \rangle > 0 \iff x^{-1}\alpha_i \in (\Delta^+ \setminus \Delta_J^+) + \mathbb{Z}\delta, \\ s_i x \xrightarrow{\alpha_i} x \iff \langle x\lambda, \alpha_i^\vee \rangle < 0 \iff x^{-1}\alpha_i \in (\Delta^- \setminus \Delta_J^-) + \mathbb{Z}\delta. \end{cases} \quad (2.13)$$

*Remark 2.6.* Let  $x \in (W^J)_{\text{af}}$  and  $i \in I_{\text{af}}$ . Then,

$$\Pi^J(s_i x) = x \iff \langle x\lambda, \alpha_i^\vee \rangle = 0 \iff x^{-1}\alpha_i \in \Delta_J + \mathbb{Z}\delta. \quad (2.14)$$

**Lemma 2.7** ([NS3, Lemma 2.3.6]). *Let  $x, y \in (W^J)_{\text{af}}$  be such that  $x \preceq y$ , and let  $i \in I_{\text{af}}$ .*

- (1) *If  $\langle x\lambda, \alpha_i^\vee \rangle > 0$  and  $\langle y\lambda, \alpha_i^\vee \rangle \leq 0$ , then  $s_i x \preceq y$ .*
- (2) *If  $\langle x\lambda, \alpha_i^\vee \rangle \geq 0$  and  $\langle y\lambda, \alpha_i^\vee \rangle < 0$ , then  $x \preceq s_i y$ .*
- (3) *If  $\langle x\lambda, \alpha_i^\vee \rangle > 0$  and  $\langle y\lambda, \alpha_i^\vee \rangle > 0$ , or if  $\langle x\lambda, \alpha_i^\vee \rangle < 0$  and  $\langle y\lambda, \alpha_i^\vee \rangle < 0$ , then  $s_i x \preceq s_i y$ .*

**Lemma 2.8** ([NNS1, Lemmas 4.3.3–4.3.5]).

- (1) *Let  $w, v \in W^J$ , and  $\xi, \zeta \in Q^\vee$ . If  $w\Pi^J(t_\xi) \succeq v\Pi^J(t_\zeta)$ , then  $[\xi]^J \geq [\zeta]^J$ ; for the projection  $[\cdot]^J : Q^\vee \rightarrow Q_{I \setminus J}^\vee$ , see (2.4).*
- (2) *Let  $w \in W^J$ , and  $\xi, \zeta \in Q^\vee$ . Then,  $w\Pi^J(t_\xi) \succeq w\Pi^J(t_\zeta)$  if and only if  $[\xi]^J \geq [\zeta]^J$ .*
- (3) *Let  $x, y \in (W^J)_{\text{af}}$  and  $\beta \in \Delta_{\text{af}}^+$  be such that  $x \xrightarrow{\beta} y$  in  $\text{BG}^{\frac{\infty}{2}}((W^J)_{\text{af}})$ . Then,  $\Pi^J(xt_\xi) \xrightarrow{\beta} \Pi^J(yt_\xi)$  in  $\text{BG}^{\frac{\infty}{2}}((W^J)_{\text{af}})$  for all  $\xi \in Q^\vee$ . Therefore, if  $x \succeq y$ , then  $\Pi^J(xt_\xi) \succeq \Pi^J(yt_\xi)$  for all  $\xi \in Q^\vee$ .*

**Lemma 2.9** ([INS, Lemma 6.1.1]). *If  $x, y \in W_{\text{af}}$  satisfy  $x \succeq y$ , then  $\Pi^J(x) \succeq \Pi^J(y)$ .*

We denote by  $* : I \rightarrow I$ ,  $i \mapsto i^*$ , the Dynkin diagram automorphism (of order 1 or 2) induced by the longest element  $w_\circ \in W$  as:  $w_\circ(\alpha_i) = -\alpha_{i^*}$  for  $i \in I$ , and then define the  $\mathbb{Z}$ -linear automorphism  $* : P \oplus \mathbb{Z}\delta \rightarrow P \oplus \mathbb{Z}\delta$  (resp.,  $* : Q^\vee \rightarrow Q^\vee$ ) by:  $\varpi_i^* = \varpi_{i^*}$  for  $i \in I$  and  $\delta^* = \delta$  (resp.,  $(\alpha_i^\vee)^* = \alpha_{i^*}^\vee$ ). Also, we define the group automorphism  $* : W \rightarrow W$  by:  $s_i^* = s_{i^*}$  for  $i \in I$ .

We see by Lemma 2.1 that if  $x \in (W^J)_{\text{af}}$  is of the form  $x = w\Pi^J(t_\xi)$  for some  $w \in W^J$  and  $\xi \in Q^\vee$ , then

$$\Pi^{J^*}(xw_\circ) = [ww_\circ]^{J^*} \Pi^{J^*}(t_{-\xi^*}), \quad (2.15)$$

where  $J^* := \{j^* \mid j \in J\} \subset I$ ; notice that

$$\Pi^J(\Pi^{J^*}(xw_\circ)w_\circ) = x. \quad (2.16)$$

The next lemma follows from [INS, Proposition A.1.2] and [LNS<sup>3</sup>1, Proposition 4.3].

**Lemma 2.10.** *Let  $x, y \in (W^J)_{\text{af}}$ , and  $\beta \in \Delta_{\text{af}}^+$ . Then,  $x \xrightarrow{\beta} y$  in  $\text{BG}^{\frac{\infty}{2}}((W^J)_{\text{af}})$  if and only if  $\Pi^{J^*}(yw_{\circ}) \xrightarrow{\beta} \Pi^{J^*}(xw_{\circ})$  in  $\text{BG}^{\frac{\infty}{2}}((W^{J^*})_{\text{af}})$ . Therefore,  $x \succeq y$  in  $(W^J)_{\text{af}}$  if and only if  $\Pi^{J^*}(yw_{\circ}) \succeq \Pi^{J^*}(xw_{\circ})$  in  $(W^{J^*})_{\text{af}}$ .*

**2.3 Deodhar lifts.** Let  $J$  be a subset of  $I$ . For  $x \in (W^J)_{\text{af}}$ , let  $\text{Lift}(x)$  denote the set of lifts of  $x$  in  $W_{\text{af}}$  for the projection  $\Pi^J : W_{\text{af}} \rightarrow (W^J)_{\text{af}}$ , that is,  $\text{Lift}(x) := \{x' \in W_{\text{af}} \mid \Pi^J(x') = x\}$ ; we know from [KNS, Lemma B.1] that if  $x = w\Pi^J(t_{\xi}) \in (W^J)_{\text{af}}$  for  $w \in W^J$  and  $\xi \in Q^{\vee}$  (see Lemma 2.1 (1)), then  $\text{Lift}(x) = \{w't_{\xi+\gamma} \mid w' \in wW_J, \gamma \in Q_J^{\vee}\}$ .

**Proposition 2.11** ([KNS, Proposition 2.4]). *If  $y \in W_{\text{af}}$  and  $x \in (W^J)_{\text{af}}$  satisfy the condition that  $x \succeq \Pi^J(y)$ , then the set*

$$\text{Lift}_{\succeq y}(x) := \{x' \in \text{Lift}(x) \mid x' \succeq y\} \quad (2.17)$$

*has a (necessarily unique) minimum element with respect to the semi-infinite Bruhat order on  $W_{\text{af}}$ ; we denote this element by  $\min \text{Lift}_{\succeq y}(x)$ .*

**Proposition 2.12.** *If  $y \in W_{\text{af}}$  and  $x \in (W^J)_{\text{af}}$  satisfy the condition that  $\Pi^J(y) \succeq x$ , then the set*

$$\text{Lift}_{\preceq y}(x) := \{x' \in \text{Lift}(x) \mid y \succeq x'\} \quad (2.18)$$

*has a (necessarily unique) maximum element with respect to the semi-infinite Bruhat order on  $W_{\text{af}}$ ; we denote this element by  $\max \text{Lift}_{\preceq y}(x)$ .*

*Proof.* We first show that  $x'w_{\circ} \in \text{Lift}_{\succeq yw_{\circ}}(\Pi^{J^*}(xw_{\circ}))$  for all  $x' \in \text{Lift}_{\preceq y}(x)$ . We see by Lemma 2.10 that  $x'w_{\circ} \succeq yw_{\circ}$ . Hence it follows from Lemma 2.9 that  $\Pi^{J^*}(x'w_{\circ}) \succeq \Pi^{J^*}(yw_{\circ})$ . Also, since  $\Pi^J(x') = x$ , there exist  $z \in W_J$  and  $\gamma \in Q_J^{\vee}$  such that  $x' = xzt_{\gamma}$ ; note that  $x'w_{\circ} = xzt_{\gamma}w_{\circ} = xw_{\circ}z^*t_{-\gamma^*}$ , with  $z^* \in W_{J^*}$  and  $\gamma^* \in Q_{J^*}^{\vee}$ . Therefore, we deduce that  $\Pi^{J^*}(x'w_{\circ}) = \Pi^{J^*}(xw_{\circ})$ , which implies that  $x'w_{\circ} \in \text{Lift}_{\succeq yw_{\circ}}(\Pi^{J^*}(xw_{\circ}))$ , as desired. Similarly, we can show that if  $x'' \in \text{Lift}_{\succeq yw_{\circ}}(\Pi^{J^*}(xw_{\circ}))$ , then  $x''w_{\circ} \in \text{Lift}_{\preceq y}(x)$  (see also (2.16)). Recall from Proposition 2.11 that  $\min \text{Lift}_{\succeq yw_{\circ}}(\Pi^{J^*}(xw_{\circ}))$  denotes the minimum element of  $\text{Lift}_{\succeq yw_{\circ}}(\Pi^{J^*}(xw_{\circ}))$ . From Lemma 2.10 and the assertions above, we deduce that  $(\min \text{Lift}_{\succeq yw_{\circ}}(\Pi^{J^*}(xw_{\circ})))w_{\circ}$  is the maximum element of  $\text{Lift}_{\preceq y}(x)$ . This proves the lemma.  $\square$

**Corollary 2.13.**

(1) *If  $y \in W_{\text{af}}$  and  $x \in (W^J)_{\text{af}}$  satisfy the condition that  $\Pi^J(y) \succeq x$ , then  $\Pi^{J^*}(xw_{\circ}) \succeq \Pi^{J^*}(yw_{\circ})$  and*

$$\max \text{Lift}_{\preceq y}(x) = (\min \text{Lift}_{\succeq yw_{\circ}}(\Pi^{J^*}(xw_{\circ})))w_{\circ}. \quad (2.19)$$

(2) *If  $y \in W_{\text{af}}$  and  $x \in (W^J)_{\text{af}}$  satisfy the condition that  $x \succeq \Pi^J(y)$ , then  $\Pi^{J^*}(yw_{\circ}) \succeq \Pi^{J^*}(xw_{\circ})$  and*

$$\min \text{Lift}_{\succeq y}(x) = (\max \text{Lift}_{\preceq yw_{\circ}}(\Pi^{J^*}(xw_{\circ})))w_{\circ}. \quad (2.20)$$

**Lemma 2.14** (cf. [NNS3, Lemma 3.6]). *Let  $J$  be a subset of  $I$ . Let  $y \in W_{\text{af}}$  and  $x \in (W^J)_{\text{af}}$  be such that  $x \succeq \Pi^J(y)$ , and set  $\tilde{x} := \min \text{Lift}_{\succeq y}(x)$ . Let  $i \in I_{\text{af}}$  be such that  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ .*

- (1) Assume that  $x^{-1}\alpha_i \in (\Delta^+ \setminus \Delta_J^+) + \mathbb{Z}\delta$ ; note that  $s_i x \in (W^J)_{\text{af}}$  by Lemma 2.5. It holds that  $s_i x \succeq \Pi^J(s_i y)$  and  $\min \text{Lift}_{\succeq s_i y}(s_i x) = s_i \tilde{x}$ .
- (2) If  $x^{-1}\alpha_i \in (\Delta^- \setminus \Delta_J^-) + \mathbb{Z}\delta$ , then  $x \succeq \Pi^J(s_i y)$  and  $\min \text{Lift}_{\succeq s_i y}(x) = \tilde{x}$ .
- (3) If  $x^{-1}\alpha_i \in \Delta_J + \mathbb{Z}\delta$ , then  $\tilde{x}^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ . Moreover, we have  $x \succeq \Pi^J(s_i y)$ , and  $\min \text{Lift}_{\succeq s_i y}(x)$  is identical to  $\tilde{x}$  or  $s_i \tilde{x}$ .

*Proof.* Although we can show this lemma by combining [NNS3, Lemma 3.6] and Proposition 2.22 below, we give a direct proof below.

(1) Since  $x^{-1}\alpha_i \in (\Delta^+ \setminus \Delta_J^+) + \mathbb{Z}\delta$  by the assumption, and since  $\Delta^+ \setminus \Delta_J^+$  is stable under the action of  $W_J$ , we see that  $\tilde{x}^{-1}\alpha_i \in (\Delta^+ \setminus \Delta_J^+) + \mathbb{Z}\delta$ . Also, we have  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$  by the assumption, and  $\tilde{x} \succeq y$  by the definition of  $\tilde{x}$ . Therefore, it follows from Lemma 2.7 (3) that  $s_i \tilde{x} \succeq s_i y$ , and hence  $\Pi^J(s_i \tilde{x}) \succeq \Pi^J(s_i y)$  by Lemma 2.9. Here it is easily checked that  $\Pi^J(s_i \tilde{x}) = s_i x$ . From these, we deduce that  $s_i x \succeq \Pi^J(s_i y)$  and  $s_i \tilde{x} \in \text{Lift}_{\succeq s_i y}(s_i x)$ . If we set  $z := \min \text{Lift}_{\succeq s_i y}(s_i x)$ , then  $s_i \tilde{x} \succeq z \succeq s_i y$ . Since  $\Pi^J(z) = s_i x$  and  $(s_i x)^{-1}\alpha_i = -x^{-1}\alpha_i \in (\Delta^- \setminus \Delta_J^-) + \mathbb{Z}\delta$ , it is easily seen that  $z^{-1}\alpha_i \in (\Delta^- \setminus \Delta_J^-) + \mathbb{Z}\delta$ . Therefore, it follows from Lemma 2.7 (3) that  $\tilde{x} \succeq s_i z \succeq y$ . Here we note that  $\Pi^J(s_i z) = x$ , and hence  $s_i z \in \text{Lift}_{\succeq y}(x)$ . Since  $\tilde{x} = \min \text{Lift}_{\succeq y}(x)$ , we obtain  $\tilde{x} = s_i z$ , and hence  $z = s_i \tilde{x}$ , as desired.

(2) As in the proof of (1), we see that  $\tilde{x}^{-1}\alpha_i \in (\Delta^- \setminus \Delta_J^-) + \mathbb{Z}\delta$ . Also, we have  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$  by the assumption, and  $\tilde{x} \succeq y$  by the definition of  $\tilde{x}$ . Therefore, it follows from Lemma 2.7 (1) that  $\tilde{x} \succeq s_i y$ , which implies that  $x = \Pi^J(\tilde{x}) \succeq \Pi^J(s_i y)$  by Lemma 2.9. Thus we have  $\tilde{x} \in \text{Lift}_{\succeq s_i y}(x)$ . If we set  $z := \min \text{Lift}_{\succeq s_i y}(x)$ , then  $\tilde{x} \succeq z \succeq s_i y$ . Since  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , we deduce that  $s_i y \succ y$ , and hence  $\tilde{x} \succeq z \succeq y$ . Since  $\tilde{x} = \min \text{Lift}_{\succeq y}(x)$ , we obtain  $z = \tilde{x}$ , as desired.

(3) Suppose, for a contradiction, that  $\tilde{x}^{-1}\alpha_i \in \Delta^- + \mathbb{Z}\delta$ . Then we have  $\tilde{x} \succ s_i \tilde{x}$ . Since  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , it follows from Lemma 2.7 (2) that  $s_i \tilde{x} \succeq y$ . Also, since  $x^{-1}\alpha_i \in \Delta_J + \mathbb{Z}\delta$ , we see that  $\Pi^J(s_i \tilde{x}) = x$ . Hence we deduce that  $s_i \tilde{x} \in \text{Lift}_{\succeq y}(x)$ . However, since  $\tilde{x} \succ s_i \tilde{x}$  as seen above, this contradicts  $\tilde{x} = \min \text{Lift}_{\succeq y}(x)$ . Thus we conclude that  $\tilde{x}^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ .

Since  $\tilde{x}^{-1}\alpha_i, y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , and  $\tilde{x} \succeq y$ , it follows from Lemma 2.7 (3) that  $s_i \tilde{x} \succeq s_i y$ . Therefore, by Lemma 2.9, we have  $x = \Pi^J(s_i \tilde{x}) \succeq \Pi^J(s_i y)$ , and hence  $s_i \tilde{x} \in \text{Lift}_{\succeq s_i y}(x)$ . Here we set  $z := \min \text{Lift}_{\succeq s_i y}(x)$ . Assume first that  $\tilde{x} \succeq s_i y$ . Then we have  $\tilde{x} \succeq z \succeq s_i y$ . Since  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , we have  $s_i y \succ y$ , and hence  $\tilde{x} \succeq z \succeq y$ . Since  $\tilde{x} = \min \text{Lift}_{\succeq y}(x)$ , we obtain  $z = \tilde{x}$ . Assume next that  $\tilde{x} \not\succeq s_i y$ . Since  $s_i \tilde{x} \in \text{Lift}_{\succeq s_i y}(x)$  as seen above, we have  $s_i \tilde{x} \succeq z \succeq s_i y$ . Note that  $(s_i \tilde{x})^{-1}\alpha_i \in \Delta^- + \mathbb{Z}\delta$ . If  $z^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , then it follows from Lemma 2.7 (2) that  $\tilde{x} \succeq z \succeq s_i y$ , which contradicts the assumption that  $\tilde{x} \not\succeq s_i y$ . Hence we obtain  $z^{-1}\alpha_i \in \Delta^- + \mathbb{Z}\delta$ . Since  $(s_i \tilde{x})^{-1}\alpha_i$  and  $(s_i y)^{-1}\alpha_i$  are contained in  $\Delta^- + \mathbb{Z}\delta$ , we see by Lemma 2.7 (3) that  $\tilde{x} \succeq s_i z \succeq y$ . Since  $x^{-1}\alpha_i \in \Delta_J + \mathbb{Z}\delta$  and  $z \in \text{Lift}(x)$ , it is easily checked that  $\Pi^J(s_i z) = x$ . Thus,  $s_i z \in \text{Lift}_{\succeq y}(x)$ . Since  $\tilde{x} = \min \text{Lift}_{\succeq y}(x)$ , we obtain  $s_i z = \tilde{x}$ , and hence  $z = s_i \tilde{x}$ . This proves the lemma.  $\square$

**2.4 Quantum Bruhat graph.** We take and fix a subset  $J$  of  $I$ .

**Definition 2.15.** The (parabolic) quantum Bruhat graph  $\text{QBG}(W^J)$  is the  $(\Delta^+ \setminus \Delta_J^+)$ -labeled directed graph whose vertices are the elements of  $W^J$ , and whose directed edges are of the form:  $w \xrightarrow{\beta} v$  for  $w, v \in W^J$  and  $\beta \in \Delta^+ \setminus \Delta_J^+$  such that  $v = [ws_\beta]$ , and such that either of the following holds: (i)  $\ell(v) = \ell(w) + 1$ ; (ii)  $\ell(v) = \ell(w) + 1 - 2\langle \rho - \rho_J, \beta^\vee \rangle$ . An edge satisfying (i) (resp., (ii)) is called a Bruhat (resp., quantum) edge. When  $J = \emptyset$ , we write  $\text{QBG}(W)$  for  $\text{QBG}(W^\emptyset)$ .

*Remark 2.16.* We know from [LNS<sup>3</sup>1, Remark 6.13] that for each  $w, v \in W^J$ , there exists a directed path in  $\text{QBG}(W^J)$  from  $w$  to  $v$ .

Let  $w, v \in W^J$ , and let  $\mathbf{p} : w = v_0 \xrightarrow{\beta_1} v_1 \xrightarrow{\beta_2} \dots \xrightarrow{\beta_l} v_l = v$  be a directed path in  $\text{QBG}(W^J)$  from  $w$  to  $v$ . We define the weight  $\text{wt}^J(\mathbf{p})$  of  $\mathbf{p}$  by

$$\text{wt}^J(\mathbf{p}) := \sum_{\substack{1 \leq k \leq l; \\ v_{k-1} \xrightarrow{\beta_k} v_k \text{ is} \\ \text{a quantum edge}}} \beta_k^\vee \in Q^{\vee,+}; \quad (2.21)$$

when  $J = \emptyset$ , we write  $\text{wt}(\mathbf{p})$  for  $\text{wt}^\emptyset(\mathbf{p})$ . For  $w, v \in W^J$ , we take a shortest directed path  $\mathbf{p}$  in  $\text{QBG}(W^J)$  from  $w$  to  $v$ , and set  $\text{wt}^J(w \Rightarrow v) := [\text{wt}^J(\mathbf{p})]^J \in Q_{I \setminus J}^{\vee,+}$ ; we know from [LNS<sup>3</sup>2, Sect. 4.1] that  $\text{wt}^J(w \Rightarrow v)$  does not depend on the choice of a shortest directed path  $\mathbf{p}$ . When  $J = \emptyset$ , we write  $\text{wt}(u \Rightarrow v)$  for  $\text{wt}^\emptyset(u \Rightarrow v)$ .

**Lemma 2.17** ([LNS<sup>3</sup>2, Lemma 7.2]). *Let  $w, v \in W^J$ , and let  $w_1 \in wW_J, v_1 \in vW_J$ . Then we have  $\text{wt}^J(w \Rightarrow v) = [\text{wt}(w_1 \Rightarrow v_1)]^J$ .*

**Lemma 2.18** ([NNS1, Lemmas 4.3.6 and 4.3.7]). *Let  $w, v \in W^J$ , and  $\xi, \zeta \in Q^\vee$ . Then,*

$$w\Pi^J(t_\xi) \succeq v\Pi^J(t_\zeta) \iff [\xi]^J \geq [\zeta + \text{wt}(v \Rightarrow w)]^J. \quad (2.22)$$

**2.5 Tilted Bruhat order.** For  $w, v \in W$ , we denote by  $\ell(w \Rightarrow v)$  the length of a shortest directed path from  $w$  to  $v$  in  $\text{QBG}(W) = \text{QBG}(W^\emptyset)$ .

**Definition 2.19** (tilted Bruhat order). For each  $v \in W$ , we define the  $v$ -tilted Bruhat order  $\leq_v$  on  $W$  as follows: for  $w_1, w_2 \in W$ ,

$$w_1 \leq_v w_2 \iff \ell(v \Rightarrow w_2) = \ell(v \Rightarrow w_1) + \ell(w_1 \Rightarrow w_2). \quad (2.23)$$

Namely,  $w_1 \leq_v w_2$  if and only if there exists a shortest directed path in  $\text{QBG}(W)$  from  $v$  to  $w_2$  passing through  $w_1$ ; or equivalently, if and only if the concatenation of a shortest directed path from  $v$  to  $w_1$  and one from  $w_1$  to  $w_2$  is one from  $v$  to  $w_2$ .

**Proposition 2.20** ([LNS<sup>3</sup>1, Theorem 7.1]). *Let  $J$  be a subset of  $I$ , and let  $v \in W$ . Then each coset  $uW_J$  for  $u \in W$  has a unique minimal element with respect to  $\leq_v$ ; we denote it by  $\min(uW_J, \leq_v)$ .*

*Remark 2.21.* Let  $J$  be a subset of  $I$ , and let  $u, v \in W$ . It is obvious by the definition of the  $v$ -tilted Bruhat order that if  $uW_J = vW_J$ , then  $\min(uW_J, \leq_v) = v$ .

**Proposition 2.22.** *Let  $J$  be a subset of  $I$ . Let  $y \in W_{\text{af}}$  and  $x \in (W^J)_{\text{af}}$  be such that  $x \succeq \Pi^J(y)$ , and write these as:*

$$\begin{cases} y = v_y t_{\xi_y} & \text{with } v_y \in W \text{ and } \xi_y \in Q^\vee; \\ x = v_x \Pi^J(t_{\xi_x}) & \text{with } v_x \in W^J \text{ and } \xi_x \in Q^\vee, \end{cases}$$

respectively. Also, write  $\min \text{Lift}_{\succeq y}(x) \in W_{\text{af}}$  as:

$$\min \text{Lift}_{\succeq y}(x) = wt_\gamma \quad \text{with } w \in W \text{ and } \gamma \in Q^\vee.$$

Then,  $w = \min(v_x W_J, \leq_{v_y})$  and  $\gamma = [\xi_x]^J + [\xi_y + \text{wt}(v_y \Rightarrow w)]_J$ .

*Proof.* Since  $wt_\gamma \in \text{Lift}(x)$ , we see that  $w \in v_x W_J$  and  $\xi_x - \gamma \in Q_J^\vee$ . If we set  $w' := \min(v_x W_J, \leq_{v_y})$ , then we have  $w' \leq_{v_y} w$ . First we show that  $w' = w$ . Suppose, for a contradiction, that  $w' \neq w$ . Then there exists a shortest directed path in  $\text{QBG}(W)$  from  $v_y$  to  $w$  that passes through  $w'$ . Hence we have  $\text{wt}(v_y \Rightarrow w) = \text{wt}(v_y \Rightarrow w') + \text{wt}(w' \Rightarrow w)$ . Since  $w$  and  $w'$  are contained in the same coset  $v_x W_J$ , we deduce by Lemma 2.17 that

$$0 = \text{wt}^J(v_x \Rightarrow v_x) = \text{wt}^J([w'] \Rightarrow [w]) = [\text{wt}(w' \Rightarrow w)]^J,$$

which implies that  $\text{wt}(w' \Rightarrow w) \in Q_J^{\vee,+}$ . Hence we obtain  $w' t_{\gamma - \text{wt}(w' \Rightarrow w)} \in \text{Lift}(x)$ . Let us show that  $w' t_{\gamma - \text{wt}(w' \Rightarrow w)} \succeq y$ . Since  $wt_\gamma = \min \text{Lift}_{\succeq y}(x) \succeq y = v_y t_{\xi_y}$ , it follows from Lemma 2.18 that  $\gamma \geq \text{wt}(v_y \Rightarrow w) + \xi_y$ . Hence we deduce that

$$\gamma - \text{wt}(w' \Rightarrow w) \geq \text{wt}(v_y \Rightarrow w) + \xi_y - \text{wt}(w' \Rightarrow w) = \text{wt}(v_y \Rightarrow w') + \xi_y.$$

We see by Lemma 2.18 that  $w' t_{\gamma - \text{wt}(w' \Rightarrow w)} \succeq v_y t_{\xi_y} = y$ . Therefore, we conclude that  $w' t_{\gamma - \text{wt}(w' \Rightarrow w)} \in \text{Lift}_{\succeq y}(x)$ . However, by Lemma 2.18, we have  $\min \text{Lift}_{\succeq y}(x) = wt_\gamma \succeq w' t_{\gamma - \text{wt}(w' \Rightarrow w)}$ , which is a contradiction; note that  $w \neq w'$  by our assumption. Thus we obtain  $w = \min(v_x W_J, \leq_{v_y})$ , as desired.

Next, we set  $\gamma' := [\xi_x]^J + [\xi_y + \text{wt}(v_y \Rightarrow w)]_J$ , and show that  $\gamma = \gamma'$ . Since  $\xi_x - \gamma \in Q_J^\vee$  as seen above,  $\gamma = [\xi_x]^J + \zeta$  for some  $\zeta \in Q_J$ . Also, since  $\min \text{Lift}_{\succeq y}(x) = wt_\gamma \succeq v_y t_{\xi_y} = y$ , we see by Lemma 2.18 that  $[\xi_x]^J + \zeta = \gamma \geq \xi_y + \text{wt}(v_y \Rightarrow w)$ . It follows that  $\zeta = [\gamma']_J \geq [\xi_y + \text{wt}(v_y \Rightarrow w)]_J$ , and hence  $\gamma = [\xi_x]^J + \zeta \geq [\xi_x]^J + [\xi_y + \text{wt}(v_y \Rightarrow w)]_J = \gamma'$ . Let us show the opposite inequality  $\gamma' \geq \gamma$ . Since  $\xi_x - \gamma' \in Q_J^\vee$  and  $w \in v_x W_J$ , it follows that  $wt_{\gamma'} \in \text{Lift}(x)$ . Because  $v_x \Pi^J(t_{\xi_x}) = x \succeq \Pi^J(y) = [v_y]^J \Pi^J(t_{\xi_y})$  by the assumption, it follows from Lemmas 2.17 and 2.18 that  $[\xi_x]^J \geq [\xi_y + \text{wt}(v_y \Rightarrow w)]^J$ . Therefore, we deduce that  $\gamma' = [\xi_x]^J + [\xi_y + \text{wt}(v_y \Rightarrow w)]_J \geq \xi_y + \text{wt}(v_y \Rightarrow w)$ , which implies that  $wt_{\gamma'} \succeq v_y t_{\xi_y} = y$  by Lemma 2.18. Hence we obtain  $wt_{\gamma'} \in \text{Lift}_{\succeq y}(x)$ . Since  $wt_\gamma = \min \text{Lift}_{\succeq y}(x)$ , it follows from Lemma 2.8 (2) that  $\gamma' \geq \gamma$ . Thus we conclude that  $\gamma = \gamma'$ , as desired. This completes the proof of the proposition.  $\square$

**Corollary 2.23.** *Let  $y \in W_{\text{af}}$  and  $\xi \in Q^\vee$  be such that  $[w_\circ]^J \Pi^J(t_\xi) \succeq \Pi^J(y)$ . We write  $y$  as  $y = v_y t_{\xi_y}$  with  $v_y \in W$  and  $\xi_y \in Q^\vee$ . Then,*

$$\min \text{Lift}_{\succeq y}([w_\circ]^J \Pi^J(t_\xi)) = \min(w_\circ W_J, \leq_{v_y}) \cdot t_{[\xi]^J + [\xi_y]_J}. \quad (2.24)$$

*Proof.* We set  $w := \min(w_\circ W_J, \leq_{v_y})$ . By Proposition 2.22, we see that

$$\min \text{Lift}_{\succeq y}([\!|w_\circ|\!]^J \Pi^J(t_\xi)) = \text{wt}_{[\xi]^J + [\xi_y + \text{wt}(v_y \Rightarrow w)]_J}.$$

Since  $w_\circ \in w_\circ W_J$  and  $w = \min(w_\circ W_J, \leq_{v_y})$ , we have  $w \leq_{v_y} w_\circ$ . Therefore, there exists a shortest directed path in  $\text{QBG}(W)$  from  $v_y$  to  $w_\circ$  passing through  $w$ , which implies that

$$\text{wt}(v_y \Rightarrow w_\circ) = \text{wt}(v_y \Rightarrow w) + \text{wt}(w \Rightarrow w_\circ).$$

Because  $w_\circ$  is greater than or equal to  $v_y$  in the ordinary Bruhat order on  $W$ , there exists a shortest directed path from  $v_y$  to  $w_\circ$  in  $\text{QBG}(W)$  whose directed edges are all Bruhat edges. Hence it follows that  $\text{wt}(v_y \Rightarrow w_\circ) = 0$ . Similarly, we have  $\text{wt}(w \Rightarrow w_\circ) = 0$ . Thus we obtain  $\text{wt}(v_y \Rightarrow w) = 0$ . This proves the corollary.  $\square$

**Definition 2.24** (dual tilted Bruhat order). For each  $v \in W$ , we define the dual  $v$ -tilted Bruhat order  $\leq_v^*$  on  $W$  as follows: for  $w_1, w_2 \in W$ ,

$$w_1 \leq_v^* w_2 \iff \ell(w_1 \Rightarrow v) = \ell(w_1 \Rightarrow w_2) + \ell(w_2 \Rightarrow v). \quad (2.25)$$

Namely,  $w_1 \leq_v^* w_2$  if and only if there exists a shortest directed path in  $\text{QBG}(W)$  from  $w_1$  to  $v$  passing through  $w_2$ ; or equivalently, if and only if the concatenation of a shortest directed path from  $w_1$  to  $w_2$  and one from  $w_2$  to  $v$  is one from  $w_1$  to  $v$ .

We see by [NNS1, Lemma 2.1.3] that  $w_1 \leq_v^* w_2$  if and only if  $w_2 w_\circ \leq_{vw_\circ} w_1 w_\circ$  for  $w_1, w_2 \in W$  and  $v \in W$ . The next proposition follows from Proposition 2.20 and this observation.

**Proposition 2.25.** *Let  $v \in W$ , and let  $J$  be a subset of  $I$ . Then each coset  $uW_J$  for  $u \in W$  has a unique maximal element with respect to  $\leq_v^*$ ; we denote it by  $\max(uW_J, \leq_v^*)$ . Then we have*

$$\max(uW_J, \leq_v^*) = (\min(uw_\circ W_{J^*}, \leq_{vw_\circ}))w_\circ. \quad (2.26)$$

We can prove the following proposition by using Proposition 2.22, together with Corollary 2.13 (1) and (2.15).

**Proposition 2.26.** *Let  $J$  be a subset of  $I$ . Let  $y \in W_{\text{af}}$  and  $x \in (W^J)_{\text{af}}$  be such that  $\Pi^J(y) \succeq x$ , and write these as:*

$$\begin{cases} y = v_y t_{\xi_y} & \text{with } v_y \in W \text{ and } \xi_y \in Q^\vee; \\ x = v_x \Pi^J(t_{\xi_x}) & \text{with } v_x \in W^J \text{ and } \xi_x \in Q^\vee, \end{cases}$$

*respectively. We write  $\max \text{Lift}_{\preceq y}(x) \in W_{\text{af}}$  as:*

$$\max \text{Lift}_{\preceq y}(x) = \text{wt}_\gamma \quad \text{with } w \in W \text{ and } \gamma \in Q^\vee.$$

*Then,  $w = \max(v_x W_J, \leq_{v_y}^*)$  and  $\gamma = [\xi_x]^J + [\xi_y - \text{wt}(w \Rightarrow v_y)]_J$ .*



### 3 Main result.

**3.1 Semi-infinite Lakshmibai-Seshadri paths.** We fix  $\lambda \in P^+ \subset P_{\text{af}}^0$  (see (2.1) and (2.2)), and set

$$J = J_\lambda := \{i \in I \mid \langle \lambda, \alpha_i^\vee \rangle = 0\} \subset I. \quad (3.1)$$

**Definition 3.1.** For a rational number  $0 < a < 1$ , we define  $\text{BG}_{a\lambda}^{\frac{\infty}{2}}((W^J)_{\text{af}})$  to be the subgraph of  $\text{BG}^{\frac{\infty}{2}}((W^J)_{\text{af}})$  with the same vertex set but having only those edges of the form  $x \xrightarrow{\beta} y$  for which  $a\langle x\lambda, \beta^\vee \rangle \in \mathbb{Z}$  holds.

**Definition 3.2.** A semi-infinite Lakshmibai-Seshadri (LS for short) path of shape  $\lambda$  is a pair

$$\pi = (\mathbf{x}; \mathbf{a}) = (x_1, \dots, x_s; a_0, a_1, \dots, a_s), \quad s \geq 1, \quad (3.2)$$

of a strictly decreasing sequence  $\mathbf{x} : x_1 \succ \dots \succ x_s$  of elements in  $(W^J)_{\text{af}}$  and an increasing sequence  $\mathbf{a} : 0 = a_0 < a_1 < \dots < a_s = 1$  of rational numbers satisfying the condition that there exists a directed path from  $x_{u+1}$  to  $x_u$  in  $\text{BG}_{a_u\lambda}^{\frac{\infty}{2}}((W^J)_{\text{af}})$  for each  $u = 1, 2, \dots, s-1$ . We denote by  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$  the set of all semi-infinite LS paths of shape  $\lambda$ .

Following [INS, Sect. 3.1] (see also [NS3, Sect. 2.4]), we endow the set  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$  with a (regular) crystal structure with weights in  $P_{\text{af}}$  by the map  $\text{wt} : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \rightarrow P_{\text{af}}$  and the root operators  $e_i, f_i, i \in I_{\text{af}}$ , defined as follows. Let  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  be of the form (3.2). Define  $\bar{\pi} : [0, 1] \rightarrow \mathbb{R} \otimes_{\mathbb{Z}} P_{\text{af}}$  to be the piecewise-linear, continuous map whose ‘‘direction vector’’ on the interval  $[a_{u-1}, a_u]$  is  $x_u\lambda \in P_{\text{af}}$  for each  $1 \leq u \leq s$ , that is,

$$\bar{\pi}(t) := \sum_{k=1}^{u-1} (a_k - a_{k-1})x_k\lambda + (t - a_{u-1})x_u\lambda \quad \text{for } t \in [a_{u-1}, a_u], 1 \leq u \leq s; \quad (3.3)$$

we know from [INS, Proposition 3.1.3] that  $\bar{\pi}$  is an (ordinary) LS path of shape  $\lambda$ , introduced in [L2, Sect. 4]. We set

$$\text{wt}(\pi) := \bar{\pi}(1) = \sum_{u=1}^s (a_u - a_{u-1})x_u\lambda \in P_{\text{af}}. \quad (3.4)$$

Now, we set

$$\begin{cases} H_i^\pi(t) := \langle \bar{\pi}(t), \alpha_i^\vee \rangle & \text{for } t \in [0, 1], \\ m_i^\pi := \min\{H_i^\pi(t) \mid t \in [0, 1]\}. \end{cases} \quad (3.5)$$

As explained in [NS3, Remark 2.4.3], all local minima of the function  $H_i^\pi(t)$ ,  $t \in [0, 1]$ , are integers; in particular, the minimum value  $m_i^\pi$  is a nonpositive integer (recall that  $\bar{\pi}(0) = 0$ , and hence  $H_i^\pi(0) = 0$ ).

We define  $e_i\pi$  as follows. If  $m_i^\pi = 0$ , then we set  $e_i\pi := \mathbf{0}$ , where  $\mathbf{0}$  is an additional element not contained in any crystal. If  $m_i^\pi \leq -1$ , then we set

$$\begin{cases} t_1 := \min\{t \in [0, 1] \mid H_i^\pi(t) = m_i^\pi\}, \\ t_0 := \max\{t \in [0, t_1] \mid H_i^\pi(t) = m_i^\pi + 1\}; \end{cases} \quad (3.6)$$

notice that  $H_i^\pi(t)$  is strictly decreasing on the interval  $[t_0, t_1]$ . Let  $1 \leq p \leq q \leq s$  be such that  $a_{p-1} \leq t_0 < a_p$  and  $t_1 = a_q$ . Then we define  $e_i\pi$  by

$$\begin{aligned} e_i\pi := & (x_1, \dots, x_p, s_i x_p, s_i x_{p+1}, \dots, s_i x_q, x_{q+1}, \dots, x_s; \\ & a_0, \dots, a_{p-1}, t_0, a_p, \dots, a_q = t_1, \dots, a_s); \end{aligned} \quad (3.7)$$

if  $t_0 = a_{p-1}$ , then we drop  $x_p$  and  $a_{p-1}$ , and if  $s_i x_q = x_{q+1}$ , then we drop  $x_{q+1}$  and  $a_q = t_1$ .

Similarly, we define  $f_i\pi$  as follows. Note that  $H_i^\pi(1) - m_i^\pi$  is a nonnegative integer. If  $H_i^\pi(1) - m_i^\pi = 0$ , then we set  $f_i\pi := \mathbf{0}$ . If  $H_i^\pi(1) - m_i^\pi \geq 1$ , then we set

$$\begin{cases} t_0 := \max\{t \in [0, 1] \mid H_i^\pi(t) = m_i^\pi\}, \\ t_1 := \min\{t \in [t_0, 1] \mid H_i^\pi(t) = m_i^\pi + 1\}; \end{cases} \quad (3.8)$$

notice that  $H_i^\pi(t)$  is strictly increasing on the interval  $[t_0, t_1]$ . Let  $0 \leq p \leq q \leq s-1$  be such that  $t_0 = a_p$  and  $a_q < t_1 \leq a_{q+1}$ . Then we define  $f_i\pi$  by

$$\begin{aligned} f_i\pi := & (x_1, \dots, x_p, s_i x_{p+1}, \dots, s_i x_q, s_i x_{q+1}, x_{q+1}, \dots, x_s; \\ & a_0, \dots, a_p = t_0, \dots, a_q, t_1, a_{q+1}, \dots, a_s); \end{aligned} \quad (3.9)$$

if  $t_1 = a_{q+1}$ , then we drop  $x_{q+1}$  and  $a_{q+1}$ , and if  $x_p = s_i x_{p+1}$ , then we drop  $x_p$  and  $a_p = t_0$ . In addition, we set  $e_i\mathbf{0} = f_i\mathbf{0} := \mathbf{0}$  for all  $i \in I_{\text{af}}$ .

**Theorem 3.3** (see [INS, Theorem 3.1.5]).

- (1) The set  $\mathbb{B}^{\frac{\infty}{2}}(\lambda) \sqcup \{\mathbf{0}\}$  is stable under the action of the root operators  $e_i$  and  $f_i$ ,  $i \in I_{\text{af}}$ .
- (2) For each  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  and  $i \in I_{\text{af}}$ , we set

$$\begin{cases} \varepsilon_i(\pi) := \max\{n \geq 0 \mid e_i^n \pi \neq \mathbf{0}\}, \\ \varphi_i(\pi) := \max\{n \geq 0 \mid f_i^n \pi \neq \mathbf{0}\}. \end{cases}$$

Then, the set  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ , equipped with the maps  $\text{wt}$ ,  $e_i$ ,  $f_i$ ,  $i \in I_{\text{af}}$ , and  $\varepsilon_i$ ,  $\varphi_i$ ,  $i \in I_{\text{af}}$ , defined above, is a crystal with weights in  $P_{\text{af}}$ .

We denote by  $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$  the connected component of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$  containing  $\pi_\lambda := (e; 0, 1) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ ; for the description of the connected components of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ , see §4.1 below.

If  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  is of the form (3.2), then we deduce from Lemma 2.10 that

$$\pi^* := (\Pi^{J^*}(x_s w_\circ), \dots, \Pi^{J^*}(x_1 w_\circ); 1 - a_s, \dots, 1 - a_1, 1 - a_0) \quad (3.10)$$

is an element of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda^*) = \mathbb{B}^{\frac{\infty}{2}}(-w_\circ \lambda)$ ; we call  $\pi^*$  the dual path of  $\pi$  (cf. [L2, §2]). Notice that  $(\pi^*)^* = \pi$ , and

$$\begin{cases} \text{wt}(\pi^*) = -\text{wt}(\pi) & \text{for } \pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda), \\ (e_i\pi)^* = f_i\pi^*, \quad (f_i\pi)^* = e_i\pi^* & \text{for } \pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \text{ and } i \in I_{\text{af}}, \end{cases} \quad (3.11)$$

where  $\mathbf{0}^*$  is understood to be  $\mathbf{0}$ .

If  $\pi \in \mathbb{B}_{\succeq}^{\infty}(\lambda)$  is of the form (3.2), then we set

$$\iota(\pi) := x_1 \in (W^J)_{\text{af}}, \quad \kappa(\pi) := x_s \in (W^J)_{\text{af}}; \quad (3.12)$$

we call  $\iota(\pi)$  and  $\kappa(\pi)$  the initial direction and the final direction of  $\pi$ , respectively. For  $x \in W_{\text{af}}$ , we set

$$\mathbb{B}_{\succeq x}^{\infty}(\lambda) := \{\pi \in \mathbb{B}_{\succeq}^{\infty}(\lambda) \mid \kappa(\pi) \succeq \Pi^J(x)\}, \quad (3.13)$$

$$\mathbb{B}_{\preceq x}^{\infty}(\lambda) := \{\pi \in \mathbb{B}_{\succeq}^{\infty}(\lambda) \mid \Pi^J(x) \succeq \iota(\pi)\}. \quad (3.14)$$

*Remark 3.4* ([NS3, Lemma 5.3.1 and Proposition 5.3.2 (2)]). The set  $\mathbb{B}_{\succeq x}^{\infty}(\lambda) \cup \{\mathbf{0}\}$  is stable under the action of  $f_i$  for all  $i \in I_{\text{af}}$ . Moreover, for  $i \in I_{\text{af}}$  such that  $\langle x\lambda, \alpha_i^\vee \rangle \geq 0$ , the set  $\mathbb{B}_{\succeq x}^{\infty}(\lambda) \cup \{\mathbf{0}\}$  is stable under the action of  $e_i$ .

If  $\pi = (x_1, \dots, x_s; a_0, \dots, a_s) \in \mathbb{B}_{\succeq}^{\infty}(\lambda)$  and  $y \in W_{\text{af}}$  satisfy  $\Pi^J(y) \succeq x_1$ , that is, if  $\pi \in \mathbb{B}_{\succeq y}^{\infty}(\lambda)$ , then we define  $\kappa(\pi, y) \in W_{\text{af}}$  by the following recursive formula:

$$\begin{cases} \widehat{x}_0 := y, \\ \widehat{x}_u := \max \text{Lift}_{\preceq \widehat{x}_{u-1}}(x_u) \quad \text{for } 1 \leq u \leq s, \\ \kappa(\pi, y) := \widehat{x}_s; \end{cases} \quad (3.15)$$

we call  $\kappa(\pi, y) \in W_{\text{af}}$  the final direction of  $\pi$  with respect to  $y$ . Similarly, if  $\pi = (x_1, \dots, x_s; a_0, \dots, a_s) \in \mathbb{B}_{\succeq}^{\infty}(\lambda)$  and  $y \in W_{\text{af}}$  satisfy  $x_s \succeq \Pi^J(y)$ , that is, if  $\pi \in \mathbb{B}_{\succeq y}^{\infty}(\lambda)$ , then we define  $\iota(\pi, y) \in W_{\text{af}}$  by the following recursive formula (from  $u = s+1$  to  $u = 1$ ):

$$\begin{cases} \widetilde{x}_{s+1} := y, \\ \widetilde{x}_u := \min \text{Lift}_{\succeq \widetilde{x}_{u+1}}(x_u) \quad \text{for } 1 \leq u \leq s, \\ \iota(\pi, y) := \widetilde{x}_1; \end{cases} \quad (3.16)$$

we call  $\iota(\pi, y) \in W_{\text{af}}$  the initial direction of  $\pi$  with respect to  $y$ . The next lemma follows from Lemma 2.10 and Corollary 2.13.

**Lemma 3.5.** *Let  $y \in W_{\text{af}}$ .*

- (1) *If  $\pi \in \mathbb{B}_{\succeq y}^{\infty}(\lambda)$ , then  $\pi^* \in \mathbb{B}_{\succeq yw_{\circ}}^{\infty}(-w_{\circ}\lambda)$ , and  $\iota(\pi, y) = (\kappa(\pi^*, yw_{\circ}))w_{\circ}$ .*
- (2) *If  $\pi \in \mathbb{B}_{\preceq y}^{\infty}(\lambda)$ , then  $\pi^* \in \mathbb{B}_{\preceq yw_{\circ}}^{\infty}(-w_{\circ}\lambda)$ , and  $\kappa(\pi, y) = (\iota(\pi^*, yw_{\circ}))w_{\circ}$ .*

**3.2 Extremal weight modules and their Demazure submodules.** We take an arbitrary  $\lambda \in P^+ \subset P_{\text{af}}^0$ . Let  $V(\lambda)$  denote the (level-zero) extremal weight module of extremal weight  $\lambda$  over  $U_{\mathbf{v}}(\mathfrak{g}_{\text{af}})$ , which is defined to be the integrable  $U_{\mathbf{v}}(\mathfrak{g}_{\text{af}})$ -module generated by a single element  $v_\lambda$  with the defining relation that “ $v_\lambda$  is an extremal weight vector of weight  $\lambda$ ”. Here, recall from [Kas3, Sect. 3.1] and [Kas4, Sect. 2.6] that  $v_\lambda$  is an extremal weight vector of weight  $\lambda$  if and only if ( $v_\lambda$  is a weight vector of weight  $\lambda$  and) there exists a family  $\{v_x\}_{x \in W_{\text{af}}}$  of weight vectors in  $V(\lambda)$  such that  $v_e = v_\lambda$ , and such that for each  $i \in I_{\text{af}}$  and  $x \in W_{\text{af}}$  with  $n := \langle x\lambda, \alpha_i^\vee \rangle \geq 0$  (resp.,  $\leq 0$ ), the equalities  $E_i v_x = 0$  and  $F_i^{(n)} v_x = v_{s_i x}$  (resp.,  $F_i v_x = 0$  and  $E_i^{(-n)} v_x = v_{s_i x}$ ) hold, where for  $i \in I_{\text{af}}$  and  $k \in \mathbb{Z}_{\geq 0}$ , the  $E_i^{(k)}$  and  $F_i^{(k)}$  are the  $k$ -th divided powers of the Chevalley generators  $E_i$  and  $F_i$  of  $U_{\mathbf{v}}(\mathfrak{g}_{\text{af}})$ , respectively; note that the weight of  $v_x$  is  $x\lambda$ . Also, for each  $x \in W_{\text{af}}$ , we define the Demazure submodule  $V_x^-(\lambda)$  of  $V(\lambda)$  by

$$V_x^-(\lambda) := U_{\mathbf{v}}^-(\mathfrak{g}_{\text{af}})v_x. \quad (3.17)$$

We know from [Kas1, Proposition 8.2.2] that  $V(\lambda)$  has a crystal basis  $\mathcal{B}(\lambda)$  and the corresponding global basis  $\{G(b) \mid b \in \mathcal{B}(\lambda)\}$ . Also, we know from [Kas4, Sect. 2.8] (see also [NS3, Sect. 4.1]) that  $V_x^-(\lambda) \subset V(\lambda)$  is compatible with the global basis of  $V(\lambda)$ , that is, there exists a subset  $\mathcal{B}_x^-(\lambda)$  of the crystal basis  $\mathcal{B}(\lambda)$  such that

$$V_x^-(\lambda) = \bigoplus_{b \in \mathcal{B}_x^-(\lambda)} \mathbb{C}(\mathbf{v})G(b) \subset V(\lambda) = \bigoplus_{b \in \mathcal{B}(\lambda)} \mathbb{C}(\mathbf{v})G(b). \quad (3.18)$$

*Remark 3.6* ([NS3, Lemma 4.1.2]). For every  $x \in W_{\text{af}}$ , we have  $V_x^-(\lambda) = V_{\Pi^J(x)}^-(\lambda)$  and  $\mathcal{B}_x^-(\lambda) = \mathcal{B}_{\Pi^J(x)}^-(\lambda)$ .

Denote by  $u_\lambda$  the element of  $\mathcal{B}(\lambda)$  such that  $G(u_\lambda) = v_\lambda$ ; recall that  $\pi_\lambda = (e; 0, 1) \in \mathbb{B}_{\frac{\infty}{2}}(\lambda)$ . We know the following from [INS, Theorem 3.2.1] and [NS3, Theorem 4.2.1].

**Theorem 3.7.** *There exists an isomorphism of crystals from  $\mathcal{B}(\lambda)$  to  $\mathbb{B}_{\frac{\infty}{2}}(\lambda)$ , which sends  $u_\lambda$  to  $\pi_\lambda$ , and sends  $\mathcal{B}_x^-(\lambda)$  to  $\mathbb{B}_{\frac{\infty}{2}, x}(\lambda)$  for all  $x \in W_{\text{af}}$ .*

Recall that  $P_{\text{af}}^0 = P \oplus \mathbb{Z}\delta$ ; for  $\nu \in P_{\text{af}}^0$ , we write  $\nu = \text{fin}(\nu) + \text{nul}(\nu)\delta$  with  $\text{fin}(\nu) \in P$  and  $\text{nul}(\nu) \in \mathbb{Z}$ . Let  $\mathbf{e}^\nu$  denote the formal exponential for  $\nu \in P_{\text{af}}^0$ , and set  $q := \mathbf{e}^\delta$ ; note that the formal exponential  $\mathbf{e}^\nu$  for  $\nu \in P_{\text{af}}^0$  is identical to  $q^{\text{nul}(\nu)}\mathbf{e}^{\text{fin}(\nu)}$ . Following [KNS, Sect. 2.4], we define the graded character  $\text{gch } V_x^-(\lambda)$  of  $V_x^-(\lambda)$  by

$$\text{gch } V_x^-(\lambda) := \sum_{k \in \mathbb{Z}} \left( \sum_{\gamma \in Q} \dim(V_x^-(\lambda)_{\lambda + \gamma + k\delta}) \mathbf{e}^{\lambda + \gamma} \right) q^k; \quad (3.19)$$

if  $x = wt_\xi$  for  $w \in W$  and  $\xi \in Q^\vee$ , then  $\text{gch } V_x^-(\lambda)$  is an element of  $(\mathbb{Z}[P])[q^{-1}]q^{-(\lambda, \xi)}$  (see [KNS, (2.22)]). It follows from Theorem 3.7 that

$$\text{gch } V_x^-(\lambda) = \sum_{\pi \in \mathbb{B}_{\frac{\infty}{2}, x}(\lambda)} \mathbf{e}^{\text{wt}(\pi)} = \sum_{\pi \in \mathbb{B}_{\frac{\infty}{2}, x}(\lambda)} q^{\text{nul}(\text{wt}(\pi))} \mathbf{e}^{\text{fin}(\text{wt}(\pi))}. \quad (3.20)$$

**Proposition 3.8** ([KNS, Proposition D.1]). *For each  $x \in W_{\text{af}}$  and  $\xi \in Q^\vee$ , the equality  $\text{gch } V_{xt_\xi}^-(\lambda) = q^{-(\lambda, \xi)} \text{gch } V_x^-(\lambda)$  holds.*

**3.3 Statement of the main result in terms of semi-infinite LS paths.** Let  $\lambda, \mu \in P^+$  be such that  $\lambda - \mu \in P^+$ , and define  $J_\lambda, J_\mu, J_{\lambda-\mu} \subset I$  as in (3.1). Write  $\lambda$  and  $\mu$  as:  $\lambda = \sum_{i \in I} \lambda_i \varpi_i$  and  $\mu = \sum_{i \in I} \mu_i \varpi_i$ , respectively, where  $\lambda_i, \mu_i \in \mathbb{Z}_{\geq 0}$ , with  $\lambda_i - \mu_i \geq 0$  for all  $i \in I$ . The following theorem is one of the main results of this paper.

**Theorem 3.9.** *For  $x \in W_{\text{af}}$ , the following equality holds:*

$$\begin{aligned} & \prod_{i \in I} \prod_{k=\lambda_i-\mu_i+1}^{\lambda_i} \frac{1}{1-q^{-k}} \text{gch } V_x^-(\lambda - \mu) \\ &= \sum_{\substack{y \in W_{\text{af}} \\ y \succeq x}} \sum_{\substack{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\mu) \\ \iota(\pi) \preceq \Pi^J \mu(y), \kappa(\pi, y) = x}} (-1)^{\ell^{\frac{\infty}{2}}(y) - \ell^{\frac{\infty}{2}}(x)} \underbrace{q^{-\text{nul}(\text{wt}(\pi))} \mathbf{e}^{-\text{fn}(\text{wt}(\pi))}}_{= \mathbf{e}^{-\text{wt}(\pi)}} \text{gch } V_y^-(\lambda). \end{aligned} \quad (3.21)$$

The outline of our proof of this theorem is as follows. First, we prove (3.21) in the case that  $\mu$  is a fundamental weight and  $x$  is a translation element in  $W_{\text{af}}$  (see §5.2); we show a key formula in this case in Proposition 5.3 on the basis of standard monomial theory for semi-infinite LS paths, established in [KNS] (see §4). Next, by making use of Demazure operators (see §6.1), we show (3.21) in the case that  $\mu$  is a fundamental weight and  $x$  is an arbitrary element of  $W$  (see §7.2). Finally, using Proposition 4.4, we prove (3.21) in the general case (see §8.1).

**3.4 Quantum Lakshmibai-Seshadri paths.** In this subsection, we fix  $\mu \in P^+$ , and take  $J = J_\mu$  as in (3.1).

**Definition 3.10.** For a rational number  $0 < a < 1$ , we define  $\text{QBG}_{a\mu}(W^J)$  to be the subgraph of  $\text{QBG}(W^J)$  with the same vertex set but having only those directed edges of the form  $w \xrightarrow{\beta} v$  for which  $a\langle \mu, \beta^\vee \rangle \in \mathbb{Z}$  holds.

**Definition 3.11.** A quantum LS path of shape  $\mu$  is a pair

$$\eta = (\mathbf{w}; \mathbf{a}) = (w_1, \dots, w_s; a_0, a_1, \dots, a_s), \quad s \geq 1, \quad (3.22)$$

of a sequence  $w_1, \dots, w_s$  of elements in  $W^J$ , with  $w_u \neq w_{u+1}$  for any  $1 \leq u \leq s-1$ , and an increasing sequence  $0 = a_0 < a_1 < \dots < a_s = 1$  of rational numbers satisfying the condition that there exists a directed path in  $\text{QBG}_{a_u\mu}(W^J)$  from  $w_{u+1}$  to  $w_u$  for each  $u = 1, 2, \dots, s-1$ .

Denote by  $\text{QLS}(\mu)$  the set of all quantum LS paths of shape  $\mu$ . In the same manner as for  $\mathbb{B}^{\frac{\infty}{2}}(\mu)$ , we can endow the set  $\text{QLS}(\mu)$  with a crystal structure with weights in  $P \cong P_{\text{af}}^0 / \mathbb{Z}\delta \subset P_{\text{af}} / \mathbb{Z}\delta$ ; for the details, see [LNS<sup>3</sup>2, Sect. 4.2] and [NNS3, Sect. 2.5].

Define a projection  $\text{cl} : (W^J)_{\text{af}} \rightarrow W^J$  by  $\text{cl}(x) := w$  for  $x \in (W^J)_{\text{af}}$  of the form  $x = w\Pi^J(t_\xi)$  with  $w \in W^J$  and  $\xi \in Q^\vee$ . For  $\pi = (x_1, \dots, x_s; a_0, a_1, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\mu)$ , we define

$$\text{cl}(\pi) := (\text{cl}(x_1), \dots, \text{cl}(x_s); a_0, a_1, \dots, a_s); \quad (3.23)$$

here, for each  $1 \leq p < q \leq s$  such that  $\text{cl}(x_p) = \cdots = \text{cl}(x_q)$ , we drop  $\text{cl}(x_p), \dots, \text{cl}(x_{q-1})$  and  $a_p, \dots, a_{q-1}$ ; we set  $\text{cl}(\mathbf{0}) := \mathbf{0}$  by convention. We know from [NS3, Sect. 6.2] that  $\text{cl}(\pi) \in \text{QLS}(\mu)$  for all  $\pi \in \mathbb{B}_0^{\infty}(\mu)$ . Also, we know the following lemma from [NS3, Lemma 6.2.3]; recall that  $\mathbb{B}_0^{\infty}(\mu)$  denotes the connected component of  $\mathbb{B}^{\infty}(\mu)$  containing  $\pi_{\mu} = (e; 0, 1)$ .

**Lemma 3.12.** *For each  $\eta \in \text{QLS}(\mu)$ , there exists a unique  $\pi_{\eta} \in \mathbb{B}_0^{\infty}(\mu)$  such that  $\text{cl}(\pi_{\eta}) = \eta$  and  $\kappa(\pi_{\eta}) = \kappa(\eta) \in W^J$ .*

*Remark 3.13.* For  $\eta = (w_1, \dots, w_s; a_0, a_1, \dots, a_s) \in \text{QLS}(\mu)$ , define  $\xi_1, \dots, \xi_{s-1}, \xi_s \in Q^{\vee}$  by the following recursive formula (from  $u = s$  to  $u = 1$ ):

$$\xi_s = 0, \quad \xi_u = \xi_{u+1} + \text{wt}(w_{u+1} \Rightarrow w_u) \quad \text{for } 1 \leq u \leq s-1; \quad (3.24)$$

for the definition of  $\text{wt}(w_{u+1} \Rightarrow w_u)$ , see §2.4. Then we know from [NNS3, Proposition 2.32] that

$$\pi_{\eta} = (w_1 \Pi^J(t_{\xi_1}), \dots, w_{s-1} \Pi^J(t_{\xi_{s-1}}), w_s; a_0, a_1, \dots, a_s); \quad (3.25)$$

the weight  $\text{wt}(\pi_{\eta}) \in P_{\text{af}}^0 = P \oplus \mathbb{Z}\delta$  of  $\pi_{\eta}$  can be written as:  $\text{wt}(\pi_{\eta}) = \text{wt}(\eta) + \text{deg}(\eta)\delta$ , where  $\text{deg} : \text{QLS}(\mu) \rightarrow \mathbb{Z}_{\leq 0}$  denotes the (tail) degree function (see [LNS<sup>3</sup>2, Corollary 4.8]) given by:

$$\text{deg}(\eta) = - \sum_{u=1}^{s-1} a_u \langle \mu, \text{wt}(w_{u+1} \Rightarrow w_u) \rangle. \quad (3.26)$$

For  $\eta = (w_1, \dots, w_s; a_0, a_1, \dots, a_s) \in \text{QLS}(\mu)$  and  $v \in W$ , define  $\kappa(\eta, v) \in W$  by the following recursive formula (cf. (3.15) and Proposition 2.26):

$$\begin{cases} \widehat{w}_0 := v, \\ \widehat{w}_u := \max(w_u W_{J, \leq^* \widehat{w}_{u-1}}) \quad \text{for } 1 \leq u \leq s, \\ \kappa(\eta, v) := \widehat{w}_s; \end{cases} \quad (3.27)$$

we call  $\kappa(\eta, v) \in W$  the final direction of  $\eta$  with respect to  $v$ . We set

$$\zeta(\eta, v) := \text{wt}(\widehat{w}_1 \Rightarrow v) + \sum_{u=1}^{s-1} \text{wt}(\widehat{w}_{u+1} \Rightarrow \widehat{w}_u). \quad (3.28)$$

*Remark 3.14.* Keep the notation and setting of Remark 3.13. We see from Lemma 2.17 that  $[\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v)]^J = [\xi_1]^J$ . Hence, by Lemma 2.1(2), we have  $\iota(\pi_{\eta}) = w_1 \Pi^J(t_{\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v)})$ .

**3.5 Reformulation of the main result in terms of quantum LS paths.** Let  $\lambda, \mu \in P^+$  be such that  $\lambda - \mu \in P^+$ , and define  $J_\lambda, J_\mu, J_{\lambda-\mu} \subset I$  as in (3.1). The following is a corollary of Theorem 3.9.

**Corollary 3.15.** *For  $x \in W$ , the following equality holds:*

$$\begin{aligned} & \text{gch } V_x^-(\lambda - \mu) \\ &= \sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(\mu) \\ \kappa(\eta, v) = x}} (-1)^{\ell(v) - \ell(x)} \underbrace{q^{-\deg(\eta)} \mathbf{e}^{-\text{wt}(\eta)}}_{= \mathbf{e}^{-\text{wt}(\pi_\eta)}} \text{gch } V_{v\zeta(\eta, v)}^-(\lambda). \end{aligned} \quad (3.29)$$

We will give a proof of this corollary in §8.2.

**3.6 Case of minuscule weights.** Here we apply the formula (3.21) in Theorem 3.9 to the case that  $\mu \in P^+$  is a minuscule weight, i.e.,  $\langle \mu, \alpha^\vee \rangle \in \{0, 1\}$  for all  $\alpha \in \Delta^+$ ; we know that  $\mu$  is just a fundamental weight  $\varpi_r$  satisfying the condition that  $\langle \varpi_r, \theta^\vee \rangle = 1$ . We deduce from this condition that the subgraph  $\text{BG}_{a\varpi_r}^{\frac{\infty}{2}}((W^{J_{\varpi_r}})_{\text{af}})$  has no edge for any rational number  $0 < a < 1$ . Hence it follows from the definition of semi-infinite LS paths that

$$\mathbb{B}^{\frac{\infty}{2}}(\varpi_r) = \{(z; 0, 1) \mid z \in (W^{J_{\varpi_r}})_{\text{af}}\}. \quad (3.30)$$

Let  $x \in W_{\text{af}}$ , and let  $y \in W_{\text{af}}$  be such that  $y \succeq x$ . We see that  $\pi = (z; 0, 1) \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r)$  satisfies  $\iota(\pi) \preceq \Pi^{J_{\varpi_r}}(y)$  and  $\kappa(\pi, y) = x$  (note that  $\Pi^{J_{\varpi_r}}(\kappa(\pi, y)) = \kappa(\pi) = \iota(\pi) = z$  in this case) if and only if  $\pi = (\Pi^{J_{\varpi_r}}(x); 0, 1)$  and  $\max \text{Lift}_{\preceq y}(\Pi^{J_{\varpi_r}}(x)) = x$ ; we have  $\text{wt}(\pi) = x\varpi_r$  in this case. Therefore, we obtain

$$\begin{aligned} & \frac{1}{1 - q^{-\lambda_r}} \text{gch } V_x^-(\lambda - \varpi_r) \\ &= \sum_{\substack{y \in W_{\text{af}}, y \succeq x \\ \max \text{Lift}_{\preceq y}(\Pi^{J_{\varpi_r}}(x)) = x}} (-1)^{\ell^{\frac{\infty}{2}}(y) - \ell^{\frac{\infty}{2}}(x)} q^{-\text{nul}(x\varpi_r)} \mathbf{e}^{-\text{fin}(x\varpi_r)} \text{gch } V_y^-(\lambda). \end{aligned} \quad (3.31)$$

Similarly, let us apply the formula (3.29) in Corollary 3.15 to the case that  $\mu = \varpi_r$  is a minuscule weight; for examples in type  $A_2$ , see Appendix A. By the same reasoning as for  $\mathbb{B}^{\frac{\infty}{2}}(\varpi_r)$  above, we see that

$$\text{QLS}(\varpi_r) = \{(u; 0, 1) \mid u \in W^{J_{\varpi_r}}\}. \quad (3.32)$$

Let  $x, v \in W$ . We deduce that  $\eta = (u; 0, 1) \in \text{QLS}(\varpi_r)$  satisfies  $\kappa(\eta, v) = x$  (note that  $[\kappa(\eta, v)]^{J_{\varpi_r}} = \kappa(\eta) = \iota(\eta) = u$  in this case) if and only if  $\eta = ([x]^{J_{\varpi_r}}; 0, 1)$  and  $\max(xW_{J_{\varpi_r}}, \leq_v^*) = x$ ; observe that  $\text{wt}(\eta) = x\varpi_r$ ,  $\deg(\eta) = 0$ , and  $\zeta(\eta, v) = \text{wt}(x \Rightarrow v)$  in this case. Therefore, we obtain

$$\text{gch } V_x^-(\lambda - \varpi_r) = \sum_{\substack{v \in W \\ \max(xW_{J_{\varpi_r}}, \leq_v^*) = x}} (-1)^{\ell(v) - \ell(x)} \mathbf{e}^{-x\varpi_r} \text{gch } V_{v\text{wt}(x \Rightarrow v)}^-(\lambda). \quad (3.33)$$

## 4 Standard monomial theory for semi-infinite LS paths.

**4.1 Connected components of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ .** Let  $\lambda \in P^+$ , and write it as  $\lambda = \sum_{i \in I} \lambda_i \varpi_i$ , with  $\lambda_i \in \mathbb{Z}_{\geq 0}$ ; note that  $J = J_\lambda = \{i \in I \mid \lambda_i = 0\}$  (see (3.1)). We define  $\text{Par}(\lambda)$  to be the set of  $I$ -tuples of partitions  $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I}$  such that  $\chi^{(i)}$  is a partition of length (strictly) less than  $\lambda_i$  for each  $i \in I$ ; a partition of length less than 0 is understood to be the empty partition  $\emptyset$ . Also, for  $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I} \in \text{Par}(\lambda)$ , we set  $|\boldsymbol{\chi}| := \sum_{i \in I} |\chi^{(i)}|$ , where for a partition  $\chi = (\chi_1 \geq \chi_2 \geq \cdots \geq \chi_l \geq 0)$ , we set  $|\chi| := \chi_1 + \cdots + \chi_l$ .

Here we recall from [INS, Sect. 7] the parametrization of the set  $\text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$  of connected components of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$  in terms of the set  $\text{Par}(\lambda)$ . We set  $\text{Turn}(\lambda) := \{k/\lambda_i \mid i \in I \setminus J \text{ and } 0 \leq k \leq \lambda_i\}$ . By [INS, Proposition 7.1.2], each connected component of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$  contains a unique element of the form:

$$(\Pi^J(t_{\xi_1}), \dots, \Pi^J(t_{\xi_{s-1}}), e; a_0, a_1, \dots, a_{s-1}, a_s), \quad (4.1)$$

where  $s \geq 1$ ,  $\xi_1, \dots, \xi_{s-1}$  are elements of  $Q_{I \setminus J}^\vee$  such that  $\xi_1 > \cdots > \xi_{s-1} > 0 =: \xi_s$  (for the notation, see §2.1 and §2.2), and  $a_u \in \text{Turn}(\lambda)$  for all  $0 \leq u \leq s$ . For each element of the form (4.1) (or equivalently, each connected component of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ ), we define an element  $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I} \in \text{Par}(\lambda)$  as follows. First, let  $i \in I \setminus J$ ; note that  $\lambda_i \geq 1$ . For each  $1 \leq k \leq \lambda_i$ , take  $0 \leq u \leq s$  in such a way that  $a_u$  is contained in the interval  $((k-1)/\lambda_i, k/\lambda_i]$ . Then we define the  $k$ -th entry  $\chi_k^{(i)}$  of the partition  $\chi^{(i)}$  to be  $\langle \varpi_i, \xi_u \rangle$ , the coefficient of  $\alpha_i^\vee$  in  $\xi_u$ ; we know from (the proof of) [INS, Proposition 7.2.1] that  $\chi_k^{(i)}$  does not depend on the choice of  $u$  above. Since  $\xi_1 > \cdots > \xi_{s-1} > 0 = \xi_s$ , we see that  $\chi_1^{(i)} \geq \cdots \geq \chi_{\lambda_i-1}^{(i)} \geq \chi_{\lambda_i}^{(i)} = 0$ . Hence, for each  $i \in I \setminus J$ , we obtain a partition  $\chi^{(i)}$  of length less than  $\lambda_i$ . For  $i \in J$ , we set  $\chi^{(i)} := \emptyset$ . Thus we obtain an element  $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I} \in \text{Par}(\lambda)$ , and hence a map from  $\text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$  to  $\text{Par}(\lambda)$ . Moreover, we know from [INS, Proposition 7.2.1] that this map is bijective; we denote by  $\pi_{\boldsymbol{\chi}} \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  the element of the form (4.1) corresponding to  $\boldsymbol{\chi} \in \text{Par}(\lambda)$  under this bijection. For  $\boldsymbol{\chi} \in \text{Par}(\lambda)$ , we denote by  $\mathbb{B}_{\boldsymbol{\chi}}^{\frac{\infty}{2}}(\lambda)$  the connected component of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$  containing  $\pi_{\boldsymbol{\chi}}$ .

*Remark 4.1.* Let  $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I} \in \text{Par}(\lambda)$ , with  $\chi^{(i)} = (\chi_1^{(i)} \geq \cdots)$  for  $i \in I$ ; note that  $\chi_1^{(i)} = 0$  if  $\chi^{(i)} = \emptyset$ . We set

$$\iota(\boldsymbol{\chi}) := \sum_{i \in I} \chi_1^{(i)} \alpha_i^\vee \in Q^\vee; \quad (4.2)$$

we see from the definition that  $\iota(\pi_{\boldsymbol{\chi}}) = \Pi^J(t_{\iota(\boldsymbol{\chi})})$ .

**4.2 Affine Weyl group action.** Let  $\mathcal{B}$  be a regular crystal in the sense of [Kas3, Sect. 2.2] (or, a normal crystal in the sense of [HK, p.389]); for example,  $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$  for  $\lambda \in P^+$  is a regular crystal by Theorem 3.7, and hence so is  $\mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  for  $\lambda, \mu \in P^+$ . We know from [Kas1, Sect. 7] that the affine Weyl group  $W_{\text{af}}$  acts on  $\mathcal{B}$  as follows: for  $b \in \mathcal{B}$  and  $i \in I_{\text{af}}$ ,

$$s_i \cdot b := \begin{cases} f_i^n b & \text{if } n := \langle \text{wt}(b), \alpha_i^\vee \rangle \geq 0, \\ e_i^{-n} b & \text{if } n := \langle \text{wt}(b), \alpha_i^\vee \rangle \leq 0. \end{cases} \quad (4.3)$$



The following lemma is shown by induction on the (ordinary) length  $\ell(x)$  of  $x$  and the tensor product rule for crystals (see also [KNS, Lemma 7.2]).

**Lemma 4.2.**

- (1) Let  $\lambda \in P^+$ , and take  $J = J_\lambda$  as in (3.1). If  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  is of the form (4.1), then for  $x \in W_{\text{af}}$ ,

$$x \cdot \pi = (\Pi^J(xt_{\xi_1}), \dots, \Pi^J(xt_{\xi_{s-1}}), \Pi^J(x); a_0, a_1, \dots, a_{s-1}, a_s). \quad (4.4)$$

- (2) Let  $\lambda, \mu \in P^+$ . Let  $\sigma \in \text{Par}(\lambda)$ ,  $\chi \in \text{Par}(\mu)$ , and  $\xi, \zeta \in Q^\vee$ . Then, for  $x \in W_{\text{af}}$ ,

$$x \cdot ((t_\xi \cdot \pi_\sigma) \otimes (t_\zeta \cdot \pi_\chi)) = (xt_\xi \cdot \pi_\sigma) \otimes (xt_\zeta \cdot \pi_\chi). \quad (4.5)$$

**4.3 Standard monomial theory.** Let  $\lambda = \sum_{i \in I} \lambda_i \varpi_i$ ,  $\mu = \sum_{i \in I} \mu_i \varpi_i \in P^+$ , and define  $J_\lambda, J_\mu, J_{\lambda+\mu} \subset I$  as in (3.1). Following [KNS, Proposition 3.4], we define  $\mathbb{S}^{\frac{\infty}{2}}(\lambda + \mu)$  to be the subset of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  consisting of those elements  $\pi \otimes \eta$  satisfying the condition that there exists  $y \in W_{\text{af}}$  such that  $\kappa(\eta) \succeq \Pi^{J_\mu}(y)$  and  $\kappa(\pi) \succeq \Pi^{J_\lambda}(\iota(\eta, y))$ ; for the definition of  $\iota(\eta, y) \in W_{\text{af}}$ , see (3.16). We know from [KNS, Theorem 3.1] that  $\mathbb{S}^{\frac{\infty}{2}}(\lambda + \mu)$  is a subcrystal of  $\mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$ , and it is isomorphic as a crystal to  $\mathbb{B}^{\frac{\infty}{2}}(\lambda + \mu)$ . Now, we briefly recall from the proof of [KNS, Theorem 3.1] the description of the isomorphism from  $\mathbb{B}^{\frac{\infty}{2}}(\lambda + \mu)$  to  $\mathbb{S}^{\frac{\infty}{2}}(\lambda + \mu) \subset \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  (which we denote by  $\Phi_{\lambda\mu}$ ). Let  $\text{Par}(\lambda, \mu)$  be the subset of those elements  $(\sigma, \chi, \xi) \in \text{Par}(\lambda) \times \text{Par}(\mu) \times Q_{I \setminus (J_\lambda \cup J_\mu)}^\vee$  satisfying the condition that  $c_i \geq \chi_1^{(i)}$  for all  $I \setminus (J_\lambda \cup J_\mu)$ , where  $\chi = (\chi^{(i)})_{i \in I}$  with  $\chi^{(i)} = (\chi_1^{(i)} \geq \dots \geq \chi_{\mu_i-1}^{(i)} \geq 0)$  for  $i \in I$ , and  $\xi = \sum_{i \in I \setminus (J_\lambda \cup J_\mu)} c_i \alpha_i^\vee$ . By [KNS, Proposition 7.8], there exists a bijection from  $\text{Par}(\lambda, \mu)$  to the set of connected components of  $\mathbb{S}^{\frac{\infty}{2}}(\lambda + \mu)$ , which sends  $(\sigma, \chi, \xi) \in \text{Par}(\lambda, \mu)$  to the connected component of  $\mathbb{S}^{\frac{\infty}{2}}(\lambda + \mu)$  containing the element  $(t_\xi \cdot \pi_\sigma) \otimes \pi_\chi$ . Here, we define a map  $\Theta : \text{Par}(\lambda, \mu) \rightarrow \text{Par}(\lambda + \mu)$  as follows. Let  $(\sigma, \chi, \xi) \in \text{Par}(\lambda, \mu)$ , and write  $\sigma \in \text{Par}(\lambda)$ ,  $\chi \in \text{Par}(\mu)$ ,  $\xi \in Q_{I \setminus (J_\lambda \cup J_\mu)}^\vee$  as:

$$\begin{cases} \sigma = (\sigma^{(i)})_{i \in I}, & \text{with } \sigma^{(i)} = (\sigma_1^{(i)} \geq \dots \geq \sigma_{\lambda_i-1}^{(i)} \geq 0) \text{ for } i \in I, \\ \chi = (\chi^{(i)})_{i \in I}, & \text{with } \chi^{(i)} = (\chi_1^{(i)} \geq \dots \geq \chi_{\mu_i-1}^{(i)} \geq 0) \text{ for } i \in I, \\ \xi = \sum_{i \in I \setminus (J_\lambda \cup J_\mu)} c_i \alpha_i^\vee; & \text{recall that } c_i \geq \chi_1^{(i)} \text{ for all } i \in I \setminus (J_\lambda \cup J_\mu). \end{cases} \quad (4.6)$$

For each  $i \in I$ , we set

$$\omega^{(i)} := \overbrace{(\sigma_1^{(i)} + c_i \geq \dots \geq \sigma_{\lambda_i-1}^{(i)} + c_i \geq c_i \geq \chi_1^{(i)} \geq \dots \geq \chi_{\mu_i-1}^{(i)})}^{\text{Remove these parts if } i \in J_\lambda}, \quad (4.7)$$

Remove these parts  
and set  $c_i = 0$  if  $i \in J_\mu$

which is a partition of length less than  $\lambda_i + \mu_i$ . Define  $\Theta(\sigma, \chi, \xi) := (\omega^{(i)})_{i \in I} \in \text{Par}(\lambda + \mu)$ ; we can deduce that this map  $\Theta$  is bijective. We know from [KNS, Sect. 7] that there exists an isomorphism

$$\Phi_{\lambda\mu} : \mathbb{B}^{\frac{\infty}{2}}(\lambda + \mu) \xrightarrow{\sim} \mathbb{S}^{\frac{\infty}{2}}(\lambda + \mu) \quad (\hookrightarrow \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)) \quad (4.8)$$

of crystals, which sends  $\pi_\omega$  to  $(t_\xi \cdot \pi_\sigma) \otimes \pi_\chi$  if  $\Theta^{-1}(\omega) = (\sigma, \chi, \xi)$  for  $\omega \in \text{Par}(\lambda + \mu)$ .

**4.4 Standard monomial theory for Demazure crystals.** Let  $\lambda = \sum_{i \in I} \lambda_i \varpi_i$ ,  $\mu = \sum_{i \in I} \mu_i \varpi_i \in P^+$ , and define  $J_\lambda, J_\mu, J_{\lambda+\mu} \subset I$  as in (3.1). For  $\pi \otimes \eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$ , we set

$$(\pi \otimes \eta)^* := \eta^* \otimes \pi^* \in \mathbb{B}^{\frac{\infty}{2}}(\mu^*) \otimes \mathbb{B}^{\frac{\infty}{2}}(\lambda^*); \quad (4.9)$$

for the definition of the dual paths  $\eta^*$  and  $\pi^*$ , see (3.10). Then we have

$$\begin{cases} \text{wt}((\pi \otimes \eta)^*) = -\text{wt}(\pi \otimes \eta), \\ (e_i(\pi \otimes \eta))^* = f_i((\pi \otimes \eta)^*), \quad (f_i(\pi \otimes \eta))^* = e_i((\pi \otimes \eta)^*) \end{cases} \quad (4.10)$$

for  $\pi \otimes \eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  and  $i \in I_{\text{af}}$ ; cf. (3.11). Also, for  $y \in W_{\text{af}}$ , we set

$$\mathbb{S}_{\geq y}^{\frac{\infty}{2}}(\lambda + \mu) := \Phi_{\lambda\mu}(\mathbb{B}_{\geq y}^{\frac{\infty}{2}}(\lambda + \mu)), \quad \mathbb{S}_{\leq y}^{\frac{\infty}{2}}(\lambda + \mu) := \Phi_{\lambda\mu}(\mathbb{B}_{\leq y}^{\frac{\infty}{2}}(\lambda + \mu)). \quad (4.11)$$

**Theorem 4.3.** *Keep the notation and setting above. Let  $y \in W_{\text{af}}$ .*

- (1) *An element  $\pi \otimes \eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  is contained in  $\mathbb{S}_{\geq y}^{\frac{\infty}{2}}(\lambda + \mu)$  if and only if  $\kappa(\eta) \succeq \Pi^{J_\mu}(y)$  and  $\kappa(\pi) \succeq \Pi^{J_\lambda}(\iota(\eta, y))$ .*
- (2) *An element  $\pi \otimes \eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  is contained in  $\mathbb{S}_{\leq y}^{\frac{\infty}{2}}(\lambda + \mu)$  if and only if  $\iota(\pi) \preceq \Pi^{J_\lambda}(y)$  and  $\iota(\eta) \preceq \Pi^{J_\mu}(\kappa(\pi, y))$ .*

*Proof.* Part (1) follows from [KNS, Theorem 3.5]. Let us prove part (2). By using [KNS, Lemma 7.2 and Remark 7.3] and (3.11), (4.10), we deduce that the following diagram is commutative:

$$\begin{array}{ccc} \mathbb{B}^{\frac{\infty}{2}}(\lambda + \mu) & \xrightarrow{\Phi_{\lambda\mu}} & \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu) \\ * \downarrow & & \downarrow * \\ \mathbb{B}^{\frac{\infty}{2}}(\mu^* + \lambda^*) & \xrightarrow{\Phi_{\mu^*\lambda^*}} & \mathbb{B}^{\frac{\infty}{2}}(\mu^*) \otimes \mathbb{B}^{\frac{\infty}{2}}(\lambda^*), \end{array}$$

where  $\lambda^* = -w_\circ \lambda$  and  $\mu^* = -w_\circ \mu$ . Also, it follows from Lemma 2.10 that  $(\mathbb{B}_{\geq y}^{\frac{\infty}{2}}(\lambda + \mu))^* = \mathbb{B}_{\geq y w_\circ}^{\frac{\infty}{2}}(\mu^* + \lambda^*)$ . Therefore, we obtain

$$\mathbb{S}_{\geq x}^{\frac{\infty}{2}}(\lambda + \mu) = \underbrace{(\mathbb{S}_{\geq x w_\circ}^{\frac{\infty}{2}}(\mu^* + \lambda^*))^*}_{\subset \mathbb{B}^{\frac{\infty}{2}}(\mu^*) \otimes \mathbb{B}^{\frac{\infty}{2}}(\lambda^*)}.$$

Hence part (2) follows from part (1) and Lemma 3.5. This proves the theorem.  $\square$

**Proposition 4.4.** *Let  $\psi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda + \mu)$ , and write  $\Phi_{\lambda\mu}(\psi) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  as  $\Phi_{\lambda\mu}(\psi) = \pi \otimes \eta$ , with  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  and  $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\mu)$ . Let  $y \in W_{\text{af}}$ .*

- (1) *If  $\psi \in \mathbb{B}_{\geq y}^{\frac{\infty}{2}}(\lambda + \mu)$ , then  $\iota(\psi, y) = \iota(\pi, \iota(\eta, y))$ .*
- (2) *If  $\psi \in \mathbb{B}_{\leq y}^{\frac{\infty}{2}}(\lambda + \mu)$ , then  $\kappa(\psi, y) = \kappa(\eta, \kappa(\pi, y))$ .*

In order to prove this proposition, we need some technical lemmas.

**Lemma 4.5.** *Let  $\nu \in P^+$  and  $y \in W_{\text{af}}$ . Let  $\psi \in \mathbb{B}_{\sum y}^{\infty}(\nu)$  and  $i \in I$  be such that  $\psi' := f_i \psi \neq \mathbf{0}$ ; note that  $\psi' \in \mathbb{B}_{\sum y}^{\infty}(\nu)$  by Remark 3.4. If  $\varepsilon_i(\psi) \geq 1$ , then  $\iota(\psi', y) = \iota(\psi, y)$ . If  $\varepsilon_i(\psi) = 0$ , then  $\iota(\psi, y)^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$ . Moreover,  $\iota(\psi', y)$  is identical to  $\iota(\psi, y)$  or  $s_i \iota(\psi, y)$ .*

*Proof.* We set  $J = J_\nu \subset I$  as in (3.1). Write  $\psi \in \mathbb{B}_{\sum y}^{\infty}(\nu)$  as  $\psi = (x_1, \dots, x_s; a_0, a_1, \dots, a_s)$ , and define  $y = (\tilde{x}_{s+1}, \tilde{x}_s, \dots, \tilde{x}_2, \tilde{x}_1 = \iota(\psi, y)$  by the same formula as (3.16). Assume that  $\psi' = f_i \psi$  is of the form:

$$\begin{aligned} \psi' = f_i \psi = (x_1, \dots, x_p, s_i x_{p+1}, \dots, s_i x_q, x_q, x_{q+1}, \dots, x_s; \\ a_0, a_1, \dots, a_p = t_0, \dots, a_{q-1}, t_1, a_q, a_{q+1}, \dots, a_s) \end{aligned} \quad (4.12)$$

for some  $0 \leq p < q \leq s$  (see (3.8) and (3.9)); remark that  $\langle x_u \nu, \alpha_i^\vee \rangle > 0$  for all  $p+1 \leq u \leq q$  (see the comment after (3.8)), which implies that

$$x_u^{-1} \alpha_i \in (\Delta^+ \setminus \Delta_J^+) + \mathbb{Z}\delta, \quad \text{and hence} \quad \tilde{x}_u^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta \quad (4.13)$$

for all  $p+1 \leq u \leq q$ . We set  $(y_1, \dots, y_q) := (x_1, \dots, x_p, s_i x_{p+1}, \dots, s_i x_q)$ , and define  $\tilde{y}_q, \dots, \tilde{y}_1$  by

$$\begin{cases} \tilde{y}_q := \min \text{Lift}_{\succeq \tilde{x}_q}(y_q), \\ \tilde{y}_u := \min \text{Lift}_{\succeq \tilde{y}_{u+1}}(y_u) \quad \text{for } 1 \leq u \leq q-1; \end{cases}$$

we have  $\iota(\psi', y) = \tilde{y}_1$ . Since  $\langle x_q \nu, \alpha_i^\vee \rangle > 0$ , i.e.,  $x_q^{-1} \alpha_i \in (\Delta^+ \setminus \Delta_J^+) + \mathbb{Z}\delta$ , and since  $\tilde{x}_q^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$  by (4.13), we see by Lemma 2.14 (1) and (2) that  $\tilde{y}_q = \min \text{Lift}_{\succeq \tilde{x}_q}(y_q) = \min \text{Lift}_{\succeq \tilde{x}_q}(s_i x_q)$  is identical to  $\min \text{Lift}_{\succeq s_i \tilde{x}_q}(s_i x_q) = s_i (\min \text{Lift}_{\succeq \tilde{x}_q}(x_q)) = s_i \tilde{x}_q$ . Also, we deduce by Lemma 2.14 (1), together with (4.13), that  $\tilde{y}_u = s_i \tilde{x}_u$  for  $p+1 \leq u \leq q-1$ .

Assume first that  $\varepsilon_i(\psi) \geq 1$ ; in order to prove that  $\iota(\psi', y) = \tilde{y}_1 = \tilde{x}_1 = \iota(\psi, y)$ , it suffices to show that

$$\tilde{y}_u = \tilde{x}_u \quad \text{for some } 1 \leq u \leq p. \quad (4.14)$$

Since  $\varepsilon_i(\psi) \geq 1$ , we see by the definition of the root operator  $e_i$  (see (3.6)) that there exists  $1 \leq u \leq p$  such that  $\langle x_u \nu, \alpha_i^\vee \rangle < 0$ . We set  $k := \max\{1 \leq u \leq p \mid \langle x_u \nu, \alpha_i^\vee \rangle < 0\}$ ; note that

$$x_k^{-1} \alpha_i \in (\Delta^- \setminus \Delta_J^-) + \mathbb{Z}\delta. \quad (4.15)$$

By the definition of the root operator  $f_i$  (see (3.8)), we see that  $\langle x_u \nu, \alpha_i^\vee \rangle = 0$  for all  $k+1 \leq u \leq p$ . Suppose, for a contradiction, that  $\tilde{x}_u^{-1} \alpha_i \in \Delta^- + \mathbb{Z}\delta$  for some  $k+1 \leq u \leq p$ , and set  $m := \max\{k+1 \leq u \leq p \mid \tilde{x}_u^{-1} \alpha_i \in \Delta^- + \mathbb{Z}\delta\}$ ; remark that  $\tilde{x}_{m+1}^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$  (for the case  $m = p$ , see (4.13)). Also, we have  $x_m^{-1} \alpha_i \in \Delta_J + \mathbb{Z}\delta$  since  $\langle x_m \nu, \alpha_i^\vee \rangle = 0$ , and  $\tilde{x}_m^{-1} \alpha_i \in \Delta^- + \mathbb{Z}\delta$  by the definition of  $m$ . However, this is a contradiction by Lemma 2.14 (3) (applied to  $\tilde{x}_m = \min \text{Lift}_{\succeq \tilde{x}_{m+1}}(x_m)$ ). Therefore, we conclude that

$$\tilde{x}_u^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta \quad \text{for all } k+1 \leq u \leq p. \quad (4.16)$$

Here, it follows from Lemma 2.14 (3) that  $\tilde{y}_p = \min \text{Lift}_{\succeq \tilde{y}_{p+1}}(y_p) = \min \text{Lift}_{\succeq s_i \tilde{x}_{p+1}}(x_p)$  is identical to  $\tilde{x}_p$  or  $s_i \tilde{x}_p$ . If  $\tilde{y}_p = \tilde{x}_p$ , then (4.14) holds for  $u = p$ . Hence we may

assume that  $\tilde{y}_p = s_i \tilde{x}_p$ . In this case, it follows again from Lemma 2.14 (3) that  $\tilde{y}_{p-1} = \min \text{Lift}_{\succeq \tilde{y}_p}(y_{p-1}) = \min \text{Lift}_{\succeq s_i \tilde{x}_p}(x_{p-1})$  is identical to  $\tilde{x}_{p-1}$  or  $s_i \tilde{x}_{p-1}$ . If  $\tilde{y}_{p-1} = \tilde{x}_{p-1}$ , then (4.14) holds for  $u = p - 1$ . Hence we may assume that  $\tilde{y}_{p-1} = s_i \tilde{x}_{p-1}$ . By repeating this argument, we may assume that

$$\tilde{y}_u = s_i \tilde{x}_u \quad \text{for all } k+1 \leq u \leq p. \quad (4.17)$$

Then, by Lemma 2.14 (2) and (4.15), (4.16),  $\tilde{y}_k = \min \text{Lift}_{\succeq \tilde{y}_{k+1}}(y_k) = \min \text{Lift}_{\succeq s_i \tilde{x}_{k+1}}(x_k)$  is identical to  $\tilde{x}_k$ . Thus we obtain  $\iota(\psi', x) = \tilde{y}_1 = \tilde{x}_1 = \iota(\psi, x)$ .

Assume next that  $\varepsilon_i(\psi) = 0$ . Suppose, for a contradiction, that  $\iota(\psi, y)^{-1} \alpha_i = \tilde{x}_1^{-1} \alpha_i \in \Delta^- + \mathbb{Z}\delta$ , and set  $l := \min\{2 \leq u \leq p+1 \mid \tilde{x}_u^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta\}$  (recall (4.13)); note that  $\tilde{x}_u^{-1} \alpha_i \in \Delta^- + \mathbb{Z}\delta$  for all  $1 \leq u \leq l-1$ , which implies that  $\langle x_u \nu, \alpha_i^\vee \rangle = \langle \tilde{x}_u \nu, \alpha_i^\vee \rangle \leq 0$  for all  $1 \leq u \leq l-1$ . Since  $\varepsilon_i(\psi) = 0$ , we deduce by the definition of the root operator  $e_i$  that  $\langle x_u \nu, \alpha_i^\vee \rangle = 0$  for all  $1 \leq u \leq l-1$ , and hence  $x_u^{-1} \alpha_i \in \Delta_J + \mathbb{Z}\delta$  for all  $1 \leq u \leq l-1$ . In particular, we obtain  $x_{l-1}^{-1} \alpha_i \in \Delta_J + \mathbb{Z}\delta$  and  $\tilde{x}_{l-1}^{-1} \alpha_i \in \Delta^- + \mathbb{Z}\delta$ . However, since  $\tilde{x}_l^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , this is a contradiction by Lemma 2.14 (3) (applied to  $\tilde{x}_{l-1} = \min \text{Lift}_{\succeq \tilde{x}_l}(x_{l-1})$ ). Thus we obtain  $\iota(\psi, y)^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$ .

Now, recall that  $\psi' = f_i \psi$  is of form (4.12). Since  $\varepsilon_i(\psi) = 0$  by our assumption, we see by the definition of the root operators  $e_i$  and  $f_i$  that

$$H_i^\psi(t) \geq 0 \text{ for } t \in [a_0, a_p] = [0, t_0], \text{ with } H_i^\psi(0) = H_i^\psi(t_0) = 0. \quad (4.18)$$

If there exists  $1 \leq u \leq p$  such that  $\langle x_u \nu, \alpha_i^\vee \rangle < 0$ , then we deduce by the same argument as for (4.14) that  $\iota(\psi', x) = \tilde{y}_1 = \tilde{x}_1 = \iota(\psi, x)$ . Hence we may assume that  $\langle x_u \nu, \alpha_i^\vee \rangle \geq 0$  for all  $1 \leq u \leq p$ . In this case, we have  $\langle x_u \nu, \alpha_i^\vee \rangle = 0$  for all  $1 \leq u \leq p$  by (4.18). By the same argument as for (4.16), we deduce that  $\tilde{x}_u^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$  for all  $1 \leq u \leq p$ . Furthermore, by the same argument as for (4.17), we see that  $\iota(\psi', x)$  is identical to  $\iota(\psi, x)$  or  $s_i \iota(\psi, x)$ . This proves the lemma.  $\square$

**Lemma 4.6.** *Let  $\lambda, \mu \in P^+$ , and  $y \in W_{\text{af}}$ . Let  $\pi \otimes \eta \in \mathbb{S}_{\succeq y}^{\infty}(\lambda + \mu)$  and  $i \in I$  be such that  $\pi' \otimes \eta' := f_i(\pi \otimes \eta) \neq \mathbf{0}$ , where  $\pi' \in \mathbb{B}_{\succeq}^{\infty}(\lambda)$  and  $\eta' \in \mathbb{B}_{\succeq}^{\infty}(\mu)$ ; notice that  $\pi' \otimes \eta' \in \mathbb{S}_{\succeq y}^{\infty}(\lambda + \mu)$  by Remark 3.4. If  $\varepsilon_i(\pi \otimes \eta) \geq 1$ , then  $\iota(\pi', \iota(\eta', y)) = \iota(\pi, \iota(\eta, y))$ . If  $\varepsilon_i(\pi \otimes \eta) = 0$ , then  $\iota(\pi, \iota(\eta, y))^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$ . Moreover,  $\iota(\pi', \iota(\eta', y))$  is identical to  $\iota(\pi, \iota(\eta, y))$  or  $s_i \iota(\pi, \iota(\eta, y))$ .*

*Proof.* If  $\varepsilon_i(\pi \otimes \eta) = 0$ , then we see by the tensor product rule for crystals that  $\varepsilon_i(\pi) = 0$ . Since  $\pi \in \mathbb{B}_{\succeq \iota(\eta, y)}^{\infty}(\lambda)$  by Theorem 4.3 (1), we see by Lemma 4.5 that  $\iota(\pi, \iota(\eta, y))^{-1} \alpha_i \in \Delta^+ + \mathbb{Z}\delta$  in this case.

Assume first that  $f_i(\pi \otimes \eta) = f_i \pi \otimes \eta$ , i.e.,  $\pi' = f_i \pi$  and  $\eta' = \eta$ ; note that  $\varphi_i(\pi) > \varepsilon_i(\eta)$  by the tensor product rule for crystals, and that  $\iota(\eta', y) = \iota(\eta, y)$ . If  $\varepsilon_i(\pi \otimes \eta) \geq 1$ , then we see by the tensor product rule for crystals and the inequality  $\varphi_i(\pi) > \varepsilon_i(\eta)$  that  $\varepsilon_i(\pi) = \varepsilon_i(\pi \otimes \eta) \geq 1$ . Hence it follows from Lemma 4.5, applied to  $\pi \in \mathbb{B}_{\succeq \iota(\eta, y)}^{\infty}(\lambda)$ , that  $\iota(\pi', \iota(\eta', y)) = \iota(\pi', \iota(\eta, y)) = \iota(\pi, \iota(\eta, y))$ . If  $\varepsilon_i(\pi \otimes \eta) = 0$ , then we have  $\varepsilon_i(\pi) = 0$ , as seen above. Therefore, we see by Lemma 4.5, applied to  $\pi \in \mathbb{B}_{\succeq \iota(\eta, y)}^{\infty}(\lambda)$ , that  $\iota(\pi', \iota(\eta', y)) = \iota(\pi', \iota(\eta, y))$  is identical to  $\iota(\pi, \iota(\eta, y))$  or  $s_i \iota(\pi, \iota(\eta, y))$ .

Assume next that  $f_i(\pi \otimes \eta) = \pi \otimes f_i\eta$ , i.e.,  $\pi' = \pi$  and  $\eta' = f_i\eta$ . We see by Lemma 4.5 that  $\iota(\eta', y)$  is identical to  $\iota(\eta, y)$  or  $s_i\iota(\eta, y)$ . If  $\iota(\eta', y) = \iota(\eta, y)$ , then it is obvious that  $\iota(\pi', \iota(\eta', y)) = \iota(\pi, \iota(\eta, y))$ . Assume now that  $\iota(\eta', y) = s_i\iota(\eta, y)$ . We see by Lemma 4.5 that  $\iota(\eta', y) = s_i\iota(\eta, y)$  only if  $\varepsilon_i(\eta) = 0$ ; note that  $\iota(\eta, y)^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ . Write  $\pi \in \mathbb{B}_{\sum \iota(\eta, y)}^{\frac{\infty}{2}}(\nu)$  as  $\pi = (x_1, \dots, x_s; a_0, a_1, \dots, a_s)$ , and define

$$\begin{aligned}\iota(\eta, y) &= \tilde{x}_{s+1}, \tilde{x}_s, \dots, \tilde{x}_2, \tilde{x}_1 = \iota(\pi, \iota(\eta, y)), \\ \iota(\eta', y) &= s_i\iota(\eta, y) = \tilde{y}_{s+1}, \tilde{y}_s, \dots, \tilde{y}_2, \tilde{y}_1 = \iota(\pi, \iota(\eta', y)),\end{aligned}$$

by the same formula as (3.16). If  $\varepsilon_i(\pi \otimes \eta) \geq 1$ , then we see by the tensor product rule for crystals and the equality  $\varepsilon_i(\eta) = 0$  that  $\varepsilon_i(\pi) \geq 1$ . Hence, by the same argument as for (4.14) (with  $p$  replaced by  $s$ ), we deduce that  $\tilde{y}_u = \tilde{x}_u$  for some  $1 \leq u \leq s$ , which implies that  $\iota(\pi', \iota(\eta', y)) = \iota(\pi, \iota(\eta, y))$ . Also, if  $\varepsilon_i(\pi \otimes \eta) = 0$ , then we have  $\varepsilon_i(\pi) = 0$ , as seen above. By the same argument as in the last paragraph of the proof of Lemma 4.5, we deduce that  $\iota(\pi', \iota(\eta', y))$  is identical to  $\iota(\pi, \iota(\eta, y))$  or  $s_i\iota(\pi, \iota(\eta, y))$ . This proves the lemma.  $\square$

**Lemma 4.7.** *Let  $\lambda, \mu \in P^+$ , and  $v \in W$ . If we set  $v' := \min(w_\circ W_{J_\mu}, \leq_v)$ , then*

$$\min(w_\circ W_{J_{\lambda+\mu}}, \leq_v) = \min(w_\circ W_{J_\lambda}, \leq_{v'}). \quad (4.19)$$

*Proof.* We set  $w := \min(w_\circ W_{J_{\lambda+\mu}}, \leq_v)$  and  $w' := \min(w_\circ W_{J_\lambda}, \leq_{v'})$ . We prove the assertion by descending induction on  $\ell(v)$ . If  $v = w_\circ$ , then we see by Remark 2.21 that  $w = w_\circ = v' = w'$ . Assume now that  $\ell(v) < \ell(w_\circ)$ , and take  $i \in I$  such that  $\ell(s_i v) = \ell(v) + 1$ , or equivalently,  $v^{-1}\alpha_i \in \Delta^+$ . We set

$$w_1 := \min(w_\circ W_{J_{\lambda+\mu}}, \leq_{s_i v}), \quad v'_1 := \min(w_\circ W_{J_\mu}, \leq_{s_i v}), \quad w'_1 := \min(w_\circ W_{J_\lambda}, \leq_{v'_1});$$

by our induction hypothesis, we have  $w_1 = w'_1$ .

**Case 1.** Assume that  $w_\circ^{-1}\alpha_i \in \Delta^- \setminus \Delta_{J_{\lambda+\mu}}^-$ , or equivalently,  $i^* \notin J_{\lambda+\mu} = J_\lambda \cap J_\mu$ . We deduce from [NNS3, Lemma 3.6 (2)] that  $w_1 = w$ . If  $w_\circ^{-1}\alpha_i \in \Delta^- \setminus \Delta_{J_\mu}^-$ , or equivalently, if  $i^* \notin J_\mu$ , then we deduce from [NNS3, Lemma 3.6 (2)] that  $v'_1 = v'$ , and hence  $w'_1 = w'$ . Hence, by our induction hypothesis, we obtain  $w = w'$ . Thus we may assume that  $w_\circ^{-1}\alpha_i \in \Delta_{J_\mu}$ , or equivalently,  $i^* \in J_\mu$ . In this case, it follows from [NNS3, Lemma 3.6 (3)] that  $(v')^{-1}\alpha_i \in \Delta^+$ , and  $v'_1$  is identical to  $v'$  or  $s_i v'$ . If  $v'_1 = v'$ , then we obtain  $w = w'$  in exactly the same way as above. Assume now that  $v'_1 = s_i v'$ . Since  $i^* \notin J_{\lambda+\mu} = J_\lambda \cap J_\mu$  and since  $i^* \in J_\mu$ , we see that  $i^* \notin J_\lambda$ , which implies that  $w_\circ^{-1}\alpha_i \in \Delta^- \setminus \Delta_{J_\lambda}^-$ . Hence it follows from [NNS3, Lemma 3.6 (2)] that

$$w'_1 = \min(w_\circ W_{J_\lambda}, \leq_{v'_1}) = \min(w_\circ W_{J_\lambda}, \leq_{s_i v'}) = \min(w_\circ W_{J_\lambda}, \leq_{v'}) = w'.$$

Therefore, by our induction hypothesis, we obtain  $w = w'$ .

**Case 2.** Assume that  $w_\circ^{-1}\alpha_i \in \Delta_{J_{\lambda+\mu}}$ , or equivalently,  $i^* \in J_{\lambda+\mu} = J_\lambda \cap J_\mu$ . In this case, it follows from [NNS3, Lemma 3.6 (3)] that  $w^{-1}\alpha_i \in \Delta^+$ , and  $w_1$  is identical to  $w$  or  $s_i w$ . Similarly, since  $w_\circ^{-1}\alpha_i \in \Delta_{J_{\lambda+\mu}} \subset \Delta_{J_\mu}$  by the assumption, we have  $(v')^{-1}\alpha_i \in \Delta^+$ , and  $v'_1$  is identical to  $v'$  or  $s_i v'$ . Since  $(v')^{-1}\alpha_i \in \Delta^+$ , and since  $w_\circ^{-1}\alpha_i \in \Delta_{J_{\lambda+\mu}} \subset \Delta_{J_\lambda}$  by the assumption, it follows from [NNS3, Lemma 3.6 (3)] that  $(w')^{-1}\alpha_i \in \Delta^+$ . In addition, if  $v'_1 = s_i v'$ , then it follows from [NNS3, Lemma 3.6 (3)] that  $w'_1$  is identical to  $w'$  or  $s_i w'$ ; if  $v'_1 = v'$ , then  $w'_1 = w'$ . In both cases,  $w'_1$  is identical to  $w'$  or  $s_i w'$ . Since  $w_1 = w'_1$  by our induction hypothesis, we deduce that  $w$  is identical to  $w'$  or  $s_i w'$ . Since  $w^{-1}\alpha_i \in \Delta^+$  and  $(w')^{-1}\alpha_i \in \Delta^+$  as seen above, we obtain  $w = w'$ , as desired. This proves the lemma.  $\square$

*Proof of Proposition 4.4.* We give a proof only for part (1), since we can show part (2) by taking the dual paths  $\psi^*$  and  $(\pi \otimes \eta)^* = \eta^* \otimes \pi^*$ , and then by applying Lemma 3.5 and part (1) to them. First, we claim that there exists  $i_1, \dots, i_n \in I_{\text{af}}$  such that

$$f_{i_1} f_{i_2} \cdots f_{i_{n-1}} f_{i_n} \psi = (w_\circ t_\gamma) \cdot \pi_\omega$$

for some  $\gamma \in Q^\vee$  and  $\omega \in \text{Par}(\lambda + \mu)$ ; for the definition of  $\pi_\omega$ , see §4.1. Indeed, by [NS3, Lemma 5.4.1], there exist  $j_1, \dots, j_m \in I_{\text{af}}$  such that  $f_{j_1} f_{j_2} \cdots f_{j_{m-1}} f_{j_m} \psi = t_\gamma \cdot \pi_\omega$  for some  $\gamma \in Q^\vee$  and  $\omega \in \text{Par}(\lambda + \mu)$ . We deduce from definition (4.3) that the action of  $w_\circ \in W$  on the (extremal) element  $t_\gamma \cdot \pi_\omega$  is given only by the root operators  $f_i$ ,  $i \in I$ . Hence we have verified the claim above (see also [NNS3, Lemma 3.11]). Here it follows from Remark 3.4 that  $f_{i_k} f_{i_{k+1}} \cdots f_{i_{n-1}} f_{i_n} \psi \in \mathbb{B}_{\leq y}^{\frac{\infty}{2}}(\lambda + \mu)$  for all  $1 \leq k \leq n+1$ . Now we proceed by induction on  $n$ . Assume first that  $n = 0$ , that is,  $\psi = (w_\circ t_\gamma) \cdot \pi_\omega$ . We write  $\Theta^{-1}(\omega)$  as  $\Theta^{-1}(\omega) = (\sigma, \chi, \xi) \in \text{Par}(\lambda, \mu)$ ; see (4.6) and (4.7). By Lemma 4.2 (2),

$$\pi \otimes \eta = \Phi_{\lambda\mu}(\psi) = \Phi_{\lambda\mu}((w_\circ t_\gamma) \cdot \pi_\omega) = ((w_\circ t_{\xi+\gamma}) \cdot \pi_\sigma) \otimes ((w_\circ t_\gamma) \cdot \pi_\chi).$$

Write  $y$  as  $y = vt_\zeta$ , with  $v \in W$  and  $\zeta \in Q^\vee$ . By Lemma 4.2 (1), Corollary 2.23, and Remark 2.21, we deduce that

$$\iota(\psi, y) = \min(w_\circ W_{J_{\lambda+\mu}}, \leq_v) \cdot t_{[\iota(\omega)+\gamma]^{J_{\lambda+\mu}+[\zeta]_{J_{\lambda+\mu}}}};$$

for  $\iota(\omega) \in Q^\vee$ , see Remark 4.1. Similarly, we deduce that

$$\iota(\eta, y) = \underbrace{\min(w_\circ W_{J_\mu}, \leq_v)}_{=: v'} \cdot t_{[\iota(\chi)+\gamma]^{J_\mu+[\zeta]_{J_\mu}}},$$

and then that

$$\iota(\pi, \iota(\eta, y)) = \min(w_\circ W_{J_\lambda}, \leq_{v'}) \cdot t_{[\xi+\gamma+\iota(\sigma)]^{J_\lambda+[[\iota(\chi)+\gamma]^{J_\mu+[\zeta]_{J_\mu}}]_{J_\lambda}}}.$$

It follows from Lemma 4.7 that  $\min(w_\circ W_{J_{\lambda+\mu}}, \leq_v) = \min(w_\circ W_{J_\lambda}, \leq_{v'})$ . Also, we see by definitions (4.6) and (4.7) that

$$[\iota(\omega) + \gamma]^{J_{\lambda+\mu}} + [\zeta]_{J_{\lambda+\mu}} = [\xi + \gamma + \iota(\sigma)]^{J_\lambda} + [[\iota(\chi) + \gamma]^{J_\mu} + [\zeta]_{J_\mu}]_{J_\lambda}.$$

From these, we obtain  $\iota(\psi, y) = \iota(\pi, \iota(\eta, y))$  in the case  $n = 0$ , as desired.

Assume now that  $n > 0$ ; for simplicity of notation, we set  $\psi' := f_{i_n} \psi$  and  $i := i_n$ . Also, write  $\Phi_{\lambda\mu}(\psi') = f_i(\pi \otimes \eta) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  as  $\Phi_{\lambda\mu}(\psi') = \pi' \otimes \eta'$ , with  $\pi' \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  and  $\eta' \in \mathbb{B}^{\frac{\infty}{2}}(\mu)$ . By our induction hypothesis, we have  $\iota(\psi', y) = \iota(\pi', \iota(\eta', y))$ . If  $\varepsilon_i(\psi) = \varepsilon_i(\pi \otimes \eta) \geq 1$ , then it follows from Lemmas 4.5 and 4.6, together with our induction hypothesis, that  $\iota(\psi, y) = \iota(\psi', y) = \iota(\pi', \iota(\eta', y)) = \iota(\pi, \iota(\eta, y))$ . Assume that  $\varepsilon_i(\psi) = \varepsilon_i(\pi \otimes \eta) = 0$ . Then we see again by Lemmas 4.5 and 4.6, together with our induction hypothesis, that  $\iota(\psi, y)$  is identical to  $\iota(\pi, \iota(\eta, y))$  or  $s_i \iota(\pi, \iota(\eta, y))$ , and that  $\iota(\psi, y)^{-1} \alpha_i \in \Delta^+$  and  $\iota(\pi, \iota(\eta, y))^{-1} \alpha_i \in \Delta^+$ ; if  $\iota(\psi, y) = s_i \iota(\pi, \iota(\eta, y))$ , then either  $\iota(\psi, y)^{-1} \alpha_i$  or  $\iota(\pi, \iota(\eta, y))^{-1} \alpha_i$  is a negative root, which is a contradiction. Therefore, we obtain  $\iota(\psi, y) = \iota(\pi, \iota(\eta, y))$ , as desired. This completes the proof of Proposition 4.4.  $\square$

**Corollary 4.8** (cf. [NNS3, Lemma 3.12]). *Let  $\psi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda + \mu)$ , and write  $\Phi_{\lambda\mu}(\psi) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\mu)$  as  $\Phi_{\lambda\mu}(\psi) = \pi \otimes \eta$ , with  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  and  $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\mu)$ . Then,*

$$\iota(\pi) = \Pi^{J_\lambda}(\iota(\psi)) \quad \text{and} \quad \kappa(\eta) = \Pi^{J_\mu}(\kappa(\psi)). \quad (4.20)$$

*Proof.* We first remark that  $J_{\lambda+\mu} \subset J_\lambda$  and  $J_{\lambda+\mu} \subset J_\mu$ , which implies that  $\Pi^{J_\lambda} \circ \Pi^{J_{\lambda+\mu}} = \Pi^{J_\lambda}$  and  $\Pi^{J_\mu} \circ \Pi^{J_{\lambda+\mu}} = \Pi^{J_\mu}$ , respectively. We take  $y \in W_{\text{af}}$  such that  $\psi \in \mathbb{B}_{\geq y}^{\frac{\infty}{2}}(\lambda + \mu)$ . By the definitions, we have  $\Pi^{J_{\lambda+\mu}}(\iota(\psi, y)) = \iota(\psi)$  and  $\Pi^{J_\lambda}(\iota(\pi, \iota(\eta, y))) = \iota(\pi)$ . Since  $\iota(\psi, y) = \iota(\pi, \iota(\eta, y))$  by Proposition 4.4 (1), we see that

$$\begin{aligned} \Pi^{J_\lambda}(\iota(\psi)) &= \Pi^{J_\lambda}(\Pi^{J_{\lambda+\mu}}(\iota(\psi, y))) = \Pi^{J_\lambda}(\Pi^{J_{\lambda+\mu}}(\iota(\pi, \iota(\eta, y)))) \\ &= \Pi^{J_\lambda}(\iota(\pi, \iota(\eta, y))) = \iota(\pi), \end{aligned}$$

as desired. Similarly, we can prove that  $\kappa(\eta) = \Pi^{J_\mu}(\kappa(\psi))$ . This proves the corollary.  $\square$

## 5 Proof of Theorem 3.9: part 1.

**5.1 Formula for graded characters.** Let us take an arbitrary  $r \in I$ ; note that  $J_{\varpi_r} = I \setminus \{r\}$  (see (3.1)). It is easily checked that the map  $(W^{J_{\varpi_r}})_{\text{af}} \rightarrow W_{\text{af}} \varpi_r$ ,  $x \mapsto x \varpi_r$ , is bijective. The next lemma follows from [INS, Proposition 4.2.1] and [NS1, Lemma 2.1.5].

**Lemma 5.1.** *If  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r)$  satisfies  $\kappa(\pi) = \Pi^{J_{\varpi_r}}(t_\xi)$  for some  $\xi \in Q^\vee$ , then  $\pi = (\Pi^{J_{\varpi_r}}(t_\xi); 0, 1)$ .*

Let us fix  $\lambda \in P^+$ . Recall from (4.8) the following isomorphism of crystals:

$$\Phi_{\lambda\varpi_r} : \mathbb{B}^{\frac{\infty}{2}}(\lambda + \varpi_r) \xrightarrow{\sim} \mathbb{S}^{\frac{\infty}{2}}(\lambda + \varpi_r) \quad (\hookrightarrow \mathbb{B}^{\frac{\infty}{2}}(\lambda) \otimes \mathbb{B}^{\frac{\infty}{2}}(\varpi_r)).$$

**Lemma 5.2.** *Let  $\xi \in Q^\vee$ , and set*

$$\mathbb{B} := \{ \psi \in \mathbb{B}_{\geq t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r) \mid \Pi^{J_{\varpi_r}}(\kappa(\psi)) = \Pi^{J_{\varpi_r}}(t_\xi) \}. \quad (5.1)$$

*Then,*

$$\Phi_{\lambda\varpi_r}(\mathbb{B}) = \mathbb{B}_{\geq t_\xi}^{\frac{\infty}{2}}(\lambda) \otimes \{ (\Pi^{J_{\varpi_r}}(t_\xi); 0, 1) \}. \quad (5.2)$$

*Proof.* First, we prove the inclusion  $\subset$  in (5.2). Let  $\psi \in \mathbb{B}$ , and write  $\Phi_{\lambda\varpi_r}(\psi)$  as  $\Phi_{\lambda\varpi_r}(\psi) = \pi \otimes \eta$ , with  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$  and  $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r)$ . Then we see from Corollary 4.8 that  $\kappa(\eta) = \Pi^{J\varpi_r}(\kappa(\psi)) = \Pi^{J\varpi_r}(t_\xi)$ . Hence it follows from Lemma 5.1 that  $\eta = (\Pi^{J\varpi_r}(t_\xi); 0, 1)$ . Also, since  $\psi \in \mathbb{B}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$ , we see from Theorem 4.3(1) that  $\kappa(\pi) \succeq \Pi^{J\lambda}(\iota(\eta, t_\xi))$ ; it is obvious that  $\iota(\eta, t_\xi) = t_\xi$  since  $\eta = (\Pi^{J\varpi_r}(t_\xi); 0, 1)$  as shown above. Hence we obtain  $\pi \in \mathbb{B}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda)$ . Thus we have proved the inclusion  $\subset$ .

Next, we prove the opposite inclusion  $\supset$  in (5.2). Let  $\pi \in \mathbb{B}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda)$ , and set  $\eta = (\Pi^{J\varpi_r}(t_\xi); 0, 1)$ ; recall that  $\iota(\eta, t_\xi) = t_\xi$ . Therefore, by Theorem 4.3(1), we have  $\pi \otimes \eta \in \mathbb{S}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r) = \Phi_{\lambda\varpi_r}(\mathbb{B}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r))$ . Let  $\psi$  be the unique element of  $\mathbb{B}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$  such that  $\pi \otimes \eta = \Phi_{\lambda\varpi_r}(\psi)$ . Then, by Corollary 4.8,  $\Pi^{J\varpi_r}(\kappa(\psi)) = \kappa(\eta) = \Pi^{J\varpi_r}(t_\xi)$ , which implies that  $\psi \in \mathbb{B}$ , as desired. This proves the lemma.  $\square$

**Proposition 5.3.** *Let  $\lambda \in P^+$  and  $r \in I$ . For  $\xi \in Q^\vee$ , the following equality holds:*

$$\text{gch } V_{t_\xi}^-(\lambda + \varpi_r) = \mathbf{e}^{t_\xi\varpi_r} \text{gch } V_{t_\xi}^-(\lambda) + \text{gch } V_{s_r t_\xi}^-(\lambda + \varpi_r). \quad (5.3)$$

*Proof.* Define  $\mathbb{B}$  as in (5.1). By Lemma 5.2 and (3.20), it suffices to prove that

$$\mathbb{B}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r) = \mathbb{B} \sqcup \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r). \quad (5.4)$$

First, suppose, for a contradiction, that  $\mathbb{B} \cap \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r) \neq \emptyset$ . Let  $\psi \in \mathbb{B} \cap \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$ . Since  $\psi \in \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$ , we have  $\kappa(\psi) \succeq \Pi^{J\lambda+\varpi_r}(s_r t_\xi)$ . Since  $\psi \in \mathbb{B}$ , we have  $\Pi^{J\varpi_r}(\kappa(\psi)) = \Pi^{J\varpi_r}(t_\xi)$ . Also, it is checked by Lemma 2.1, together with  $J_{\lambda+\varpi_r} \subset J_{\varpi_r}$ , that  $\Pi^{J\varpi_r}(\Pi^{J\lambda+\varpi_r}(s_r t_\xi)) = s_r \Pi^{J\varpi_r}(t_\xi)$ . Therefore, by Lemma 2.9, we see that

$$\Pi^{J\varpi_r}(t_\xi) = \Pi^{J\varpi_r}(\kappa(\psi)) \succeq \Pi^{J\varpi_r}(\Pi^{J\lambda+\varpi_r}(s_r t_\xi)) = s_r \Pi^{J\varpi_r}(t_\xi).$$

However, since  $\langle \Pi^{J\varpi_r}(t_\xi)\varpi_r, \alpha_r^\vee \rangle = \langle t_\xi\varpi_r, \alpha_r^\vee \rangle = 1 > 0$ , it follows from Lemma 2.5 that  $s_r \Pi^{J\varpi_r}(t_\xi) \succ \Pi^{J\varpi_r}(t_\xi)$ , which is a contradiction. Thus we obtain  $\mathbb{B} \cap \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r) = \emptyset$ .

Next we prove the equality in (5.4). The inclusion  $\supset$  is obvious; note that  $s_r t_\xi \succ t_\xi$ . Let us prove the opposite inclusion  $\subset$ . Let  $\psi \in \mathbb{B}_{\succeq t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$ , and assume that  $\psi \notin \mathbb{B}$  to conclude that  $\psi \in \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$ . We write  $\kappa(\psi) \in (W^{J\lambda+\varpi_r})_{\text{af}}$  as  $\kappa(\psi) = w\Pi^{J\lambda+\varpi_r}(t_\gamma)$  with  $w \in W^{J\lambda+\varpi_r}$  and  $\gamma \in Q^\vee$ ; note that  $[\gamma]^{J\lambda+\varpi_r} \geq [\xi]^{J\lambda+\varpi_r}$  by Lemma 2.8(1). It is easily checked by Lemma 2.1 that  $\Pi^{J\varpi_r}(\kappa(\psi)) = \Pi^{J\varpi_r}(w\Pi^{J\lambda+\varpi_r}(t_\gamma)) = [w]^{J\varpi_r}\Pi^{J\varpi_r}(t_\gamma)$ ; since  $\psi \notin \mathbb{B}$ , it follows that  $[w]^{J\varpi_r} \neq e$  or  $\gamma - \xi \notin Q_{J_{\varpi_r}}^\vee$ . If  $[w]^{J\varpi_r} \neq e$ , then a reduced expression for  $w$  contains  $s_r$ , which implies that  $w \succeq s_r$  (note that  $s_r \in W^{J\lambda+\varpi_r}$ ). Therefore, by Lemma 2.8(3),  $\kappa(\psi) = w\Pi^{J\lambda+\varpi_r}(t_\gamma) \succeq s_r\Pi^{J\lambda+\varpi_r}(t_\gamma)$ . Also, since  $[\gamma]^{J\lambda+\varpi_r} \geq [\xi]^{J\lambda+\varpi_r}$  as seen above, we have  $s_r\Pi^{J\lambda+\varpi_r}(t_\gamma) \succeq s_r\Pi^{J\lambda+\varpi_r}(t_\xi)$  by Lemma 2.8(2); note that  $s_r\Pi^{J\lambda+\varpi_r}(t_\xi) = \Pi^{J\lambda+\varpi_r}(s_r t_\xi)$  by Lemma 2.1. Combining these inequalities, we obtain  $\kappa(\psi) \succeq \Pi^{J\lambda+\varpi_r}(s_r t_\xi)$ , and hence  $\psi \in \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$ . Assume now that  $\gamma - \xi \notin Q_{J_{\varpi_r}}^\vee$ , or equivalently,  $\langle \varpi_r, \gamma - \xi \rangle \neq 0$ . Since  $[\gamma]^{J\lambda+\varpi_r} \geq [\xi]^{J\lambda+\varpi_r}$  as seen above, and since  $r \in I \setminus J_{\lambda+\varpi_r}$ , it follows that  $\langle \varpi_r, \gamma - \xi \rangle > 0$ , which implies that  $[\gamma]^{J\lambda+\varpi_r} \geq [\xi + \alpha_r^\vee]^{J\lambda+\varpi_r}$ .



Therefore, by Lemma 2.8 (2),  $\kappa(\psi) = w\Pi^{J_{\lambda+\varpi_r}}(t_\gamma) \succeq w\Pi^{J_{\lambda+\varpi_r}}(t_{\xi+\alpha_r^\vee})$ . Also, since  $w \succeq e$ , it follows by Lemma 2.8 (3) that  $w\Pi^{J_{\lambda+\varpi_r}}(t_{\xi+\alpha_r^\vee}) \succeq \Pi^{J_{\lambda+\varpi_r}}(t_{\xi+\alpha_r^\vee})$ . Finally, notice that  $s_r t_\xi \prec t_{\xi+\alpha_r^\vee}$  since  $s_r t_\xi \xrightarrow{-\alpha_r+\delta} t_{\xi+\alpha_r^\vee}$  in  $\text{BG}^{\frac{\infty}{2}}(W_{\text{af}})$ . It follows from Lemma 2.9 that  $\Pi^{J_{\lambda+\varpi_r}}(t_{\xi+\alpha_r^\vee}) \succeq \Pi^{J_{\lambda+\varpi_r}}(s_r t_\xi)$ . Combining these inequalities, we obtain  $\kappa(\psi) \succeq \Pi^{J_{\lambda+\varpi_r}}(s_r t_\xi)$ , and hence  $\psi \in \mathbb{B}_{\succeq s_r t_\xi}^{\frac{\infty}{2}}(\lambda + \varpi_r)$ . This completes the proof of the proposition.  $\square$

**5.2 First step in the proof of Theorem 3.9.** We prove (3.21) in the case that  $\mu = \varpi_r$  for  $r \in I$ , and  $x = t_\xi$  for  $\xi \in Q^\vee$ . In this case, the right-hand side of (3.21) is identical to

$$\sum_{\substack{y \in W_{\text{af}} \\ y \succeq t_\xi}} \sum_{\substack{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r) \\ \iota(\pi) \preceq \Pi^{J_{\varpi_r}}(y), \kappa(\pi, y) = t_\xi}} (-1)^{\ell^{\frac{\infty}{2}}(y)} \mathbf{e}^{-\text{wt}(\pi)} \text{gch } V_y^-(\lambda). \quad (5.5)$$

Let  $y \in W_{\text{af}}$  be such that  $y \succeq t_\xi$ . If  $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r)$  satisfies  $\kappa(\pi, y) = t_\xi \in W_{\text{af}}$ , then we see that  $\kappa(\pi) = \Pi^{J_{\varpi_r}}(t_\xi)$ . Hence it follows from Lemma 5.1 that  $\pi = (\Pi^{J_{\varpi_r}}(t_\xi); 0, 1)$ . Therefore, for  $y \in W_{\text{af}}$  such that  $y \succeq t_\xi$ ,

$$\underbrace{\{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r) \mid \iota(\pi) \preceq \Pi^{J_{\varpi_r}}(y), \kappa(\pi, y) = t_\xi\}}_{\text{this equality holds iff } \max \text{Lift}_{\preceq y}(\Pi^{J_{\varpi_r}}(t_\xi)) = t_\xi} = \{(\Pi^{J_{\varpi_r}}(t_\xi); 0, 1)\} \quad \text{or } \emptyset. \quad (5.6)$$

**Lemma 5.4.** *Let  $y \in W_{\text{af}}$  be such that  $y \succeq t_\xi$ , and write it as  $y = vt_\zeta$  with  $v \in W$  and  $\zeta \in Q^{\vee,+}$ . Then,*

$$\max \text{Lift}_{\preceq y}(\Pi^{J_{\varpi_r}}(t_\xi)) = t_\xi \quad \iff \quad v \in \{e, s_r\} \text{ and } \zeta - \xi \in \mathbb{Z}_{\geq 0}\alpha_r^\vee.$$

*Proof.* We write  $\max \text{Lift}_{\preceq y}(\Pi^{J_{\varpi_r}}(t_\xi))$  as:  $\max \text{Lift}_{\preceq y}(\Pi^{J_{\varpi_r}}(t_\xi)) = wt_\gamma$  with  $w \in W$  and  $\gamma \in Q^\vee$ . We see from Proposition 2.26 that

$$w = \max(eW_{J_{\varpi_r}, \leq_v^*}), \quad \gamma = [\xi]^{J_{\varpi_r}} + [\zeta - \text{wt}(w \Rightarrow v)]_{J_{\varpi_r}}.$$

We first prove the implication  $\Rightarrow$ . Let us assume that  $w = e$  and  $\gamma = \xi$ . Because  $v$  is greater than or equal to  $e$  in the (ordinary) Bruhat order on  $W$ , there exists a shortest directed path from  $e$  to  $v$  in  $\text{QBG}(W)$  whose directed edges are all Bruhat edges. Therefore, we have  $\text{wt}(w \Rightarrow v) = \text{wt}(e \Rightarrow v) = 0$ . Hence we obtain  $\xi = [\xi]^{J_{\varpi_r}} + [\zeta]_{J_{\varpi_r}}$ , which implies that  $\zeta - \xi \in \mathbb{Z}_{\geq 0}\alpha_r^\vee$ . Also, suppose, for a contradiction, that  $s_j$  appears in a reduced expression for  $v$  for some  $j \in J_{\varpi_r} = I \setminus \{r\}$ . Then,  $v$  is greater than or equal to  $s_j$  in the Bruhat order on  $W$ , Therefore, there exists a shortest directed path from  $e$  to  $v$  in  $\text{QBG}(W)$  passing through  $s_j$  (whose directed edges are all Bruhat edges); in particular,  $\ell(e \Rightarrow v) > \ell(s_j \Rightarrow v)$ . However, since  $\max(eW_{J_{\varpi_r}, \leq_v^*}) = e$  by our assumption, and since  $s_j \in eW_{J_{\varpi_r}}$ , it follows from the definition of  $\leq_v^*$  that  $\ell(s_j \Rightarrow v) = \ell(s_j \Rightarrow e) + \ell(e \Rightarrow v)$ ; in particular,  $\ell(s_j \Rightarrow v) < \ell(e \Rightarrow v)$ , which is a contradiction. Hence we conclude that  $v \in \{e, s_r\}$ .

We next prove the implication  $\Leftarrow$ . Assume that  $v = e$ . Then it is obvious that  $w = e$ , and hence  $\text{wt}(w \Rightarrow v) = \text{wt}(e \Rightarrow e) = 0$ . Hence we obtain  $\gamma = [\xi]^{J_{\varpi_r}} + [\zeta]_{J_{\varpi_r}}$ . Since

$\zeta - \xi \in \mathbb{Z}_{\geq 0} \alpha_r^\vee$  by the assumption, we deduce that  $\gamma = \xi$ . Assume now that  $v = s_r$ . By the definition of  $\leq_v^*$ , we have  $\ell(e \Rightarrow s_r) = \ell(e \Rightarrow w) + \ell(w \Rightarrow s_r)$ . Since  $\ell(e \Rightarrow s_r) = 1$ , we see that  $\ell(w \Rightarrow s_r) = 0$  or  $1$ . Since  $w \in eW_{J_{\varpi_r}}$  and  $s_r \notin eW_{J_{\varpi_r}}$ , it follows that  $w \neq s_r$ , which implies that  $\ell(w \Rightarrow s_r) = 1$ . Therefore, we obtain  $\ell(e \Rightarrow w) = 0$ , i.e.,  $w = e$ , and hence the same argument as above shows that  $\gamma = \xi$ . This proves the lemma.  $\square$

By Lemma 5.4 and (5.6), we see that the right-hand side of (5.5) (and hence that of (3.21)) is identical to:

$$\begin{aligned} & \sum_{m \in \mathbb{Z}_{\geq 0}} \sum_{v \in \{e, s_r\}} (-1)^{\ell \frac{\infty}{2}(vt_\xi + m\alpha_r^\vee)} \mathbf{e}^{-t_\xi \varpi_r} \text{gch } V_{vt_\xi + m\alpha_r^\vee}^- (\lambda) \\ &= \sum_{m \in \mathbb{Z}_{\geq 0}} q^{-m\lambda_r} \mathbf{e}^{-t_\xi \varpi_r} (\text{gch } V_{t_\xi}^- (\lambda) - \text{gch } V_{s_r t_\xi}^- (\lambda)) \quad \text{by Proposition 3.8.} \end{aligned} \quad (5.7)$$

By Proposition 5.3, we have  $\text{gch } V_{t_\xi}^- (\lambda) - \text{gch } V_{s_r t_\xi}^- (\lambda) = \mathbf{e}^{t_\xi \varpi_r} \text{gch } V_{t_\xi}^- (\lambda - \varpi_r)$ . Substituting this equality into (5.7), we deduce that the right-hand side of (5.7) is identical to:

$$\sum_{m \in \mathbb{Z}_{\geq 0}} q^{-m\lambda_r} \text{gch } V_{t_\xi}^- (\lambda - \varpi_r) = \frac{1}{1 - q^{-\lambda_r}} \text{gch } V_{t_\xi}^- (\lambda - \varpi_r).$$

This proves Theorem 3.9 in the case that  $\mu = \varpi_r$  and  $x = t_\xi$ .

## 6 String property and Demazure operators.

### 6.1 Recursion formula for graded characters in terms of Demazure operators.

For each  $i \in I_{\text{af}}$ , we define a  $\mathbb{C}(q)$ -linear operator  $D_i$  on  $\mathbb{C}(q)[P]$  by

$$D_i \mathbf{e}^\nu := \frac{\mathbf{e}^{\nu - \rho} - \mathbf{e}^{s_i(\nu - \rho)}}{1 - \mathbf{e}^{\alpha_i}} \mathbf{e}^\rho = \frac{\mathbf{e}^\nu - \mathbf{e}^{\alpha_i} \mathbf{e}^{s_i \nu}}{1 - \mathbf{e}^{\alpha_i}} \quad \text{for } \nu \in P, \text{ where } q = \mathbf{e}^\delta;$$

note that  $D_i^2 = D_i$ , and that

$$D_i \mathbf{e}^\nu = \begin{cases} \mathbf{e}^\nu (1 + \mathbf{e}^{\alpha_i} + \mathbf{e}^{2\alpha_i} + \dots + \mathbf{e}^{-\langle \nu, \alpha_i^\vee \rangle \alpha_i}) & \text{if } \langle \nu, \alpha_i^\vee \rangle \leq 0, \\ 0 & \text{if } \langle \nu, \alpha_i^\vee \rangle = 1, \\ -\mathbf{e}^\nu (\mathbf{e}^{-\alpha_i} + \mathbf{e}^{-2\alpha_i} + \dots + \mathbf{e}^{-(\langle \nu, \alpha_i^\vee \rangle + 1)\alpha_i}) & \text{if } \langle \nu, \alpha_i^\vee \rangle \geq 2. \end{cases} \quad (6.1)$$

Also, we define a  $\mathbb{C}(q)$ -linear operator  $T_i$  on  $\mathbb{C}(q)[P]$  by

$$T_i := D_i - 1, \quad \text{that is,} \quad T_i \mathbf{e}^\nu = \frac{\mathbf{e}^{\alpha_i} (\mathbf{e}^\nu - \mathbf{e}^{s_i \nu})}{1 - \mathbf{e}^{\alpha_i}};$$

note that  $T_i^2 = -T_i$  and  $T_i D_i = D_i T_i = 0$ . We can easily verify the following lemma.

**Lemma 6.1** (Leibniz rule). *For  $\nu_1, \nu_2 \in P$  and  $i \in I_{\text{af}}$ , it holds that*

$$D_i (\mathbf{e}^{\nu_1} \mathbf{e}^{\nu_2}) = (D_i \mathbf{e}^{\nu_1 + \rho}) \mathbf{e}^{\nu_2 - \rho} + \mathbf{e}^{s_i \nu_1} (D_i \mathbf{e}^{\nu_2}), \quad (6.2)$$

$$T_i (\mathbf{e}^{\nu_1} \mathbf{e}^{\nu_2}) = (T_i \mathbf{e}^{\nu_1}) \mathbf{e}^{\nu_2} + \mathbf{e}^{s_i \nu_1} (T_i \mathbf{e}^{\nu_2}). \quad (6.3)$$

Now, we take an arbitrary  $\mu \in P^+$ , and define  $J_\mu \subset I$  as in (3.1). Fix  $i \in I_{\text{af}}$ . A subset  $S \subset \mathbb{B}_{\frac{\infty}{2}}(\mu)$  is called an  $i$ -string if  $S$  is of the form  $S = \{\pi, f_i \pi, \dots, f_i^{\varphi_i(\pi)-1} \pi, f_i^{\varphi_i(\pi)} \pi\}$  for some  $\pi \in \mathbb{B}_{\frac{\infty}{2}}(\mu)$  such that  $e_i \pi = \mathbf{0}$ ; in this case, we call  $\pi_{\text{H}} = \pi_{\text{H}}^S := \pi$  and  $\pi_{\text{L}}^S := f_i^{\varphi_i(\pi)} \pi$  the  $i$ -highest element and the  $i$ -lowest element in  $S$ , respectively. Note that  $\pi_{\text{H}} = \pi_{\text{L}}$  if and only if  $\#S = 1$ .

*Remark 6.2.* The crystal  $\mathbb{B}_{\frac{\infty}{2}}(\mu)$  decomposes into a disjoint union of (infinitely many)  $i$ -strings for each  $i \in I_{\text{af}}$ .

The next lemma follows from the definition of the root operator  $f_i$  (see also [NS3, Lemma 2.4.5 and its proof]; cf. [L1, 5.3 Lemma]).

**Lemma 6.3.** *Keep the notation and setting above. We set  $w := \kappa(\pi_{\text{H}}) \in (W^{J_\mu})_{\text{af}}$  and  $N := \varphi_i(\pi_{\text{H}}) \in \mathbb{Z}_{\geq 0}$ ; note that  $f_i^N \pi_{\text{H}} = \pi_{\text{L}}$ .*

- (1) *If  $N \geq 1$ , then  $\kappa(f_i^k \pi_{\text{H}}) = w$  for all  $0 \leq k \leq N - 1$ .*
- (2) *If  $\langle w\mu, \alpha_i^\vee \rangle \leq 0$ , then  $\kappa(f_i^k \pi_{\text{H}}) = w$  for all  $0 \leq k \leq N$ .*
- (3) *If  $\langle w\mu, \alpha_i^\vee \rangle > 0$ , then  $N \geq 1$ , and  $\kappa(f_i^N \pi_{\text{H}}) = s_i w \succ w$ .*

The following corollary is an immediate consequence of Lemma 6.3.

**Corollary 6.4** (cf. [L1, 5.4 Lemma]). *Let  $i \in I_{\text{af}}$  and  $y \in W_{\text{af}}$ . For each  $i$ -string  $S \subset \mathbb{B}_{\frac{\infty}{2}}(\mu)$ , the intersection  $\mathbb{B}_{\frac{\infty}{2}}(\mu) \cap S$  is identical to  $\emptyset$ ,  $S$ , or  $\{\pi_{\text{L}}^S\}$ .*

The proof of the next lemma is straightforward.

**Lemma 6.5.** *Let  $i \in I_{\text{af}}$ , and let  $S \subset \mathbb{B}_{\frac{\infty}{2}}(\mu)$  be an  $i$ -string, with  $\pi_{\text{H}} = \pi_{\text{H}}^S$  and  $\pi_{\text{L}} = \pi_{\text{L}}^S$  the  $i$ -highest element and the  $i$ -lowest element in  $S$ , respectively. Then,*

$$D_i \left( \sum_{\pi \in S} e^{\text{wt}(\pi)} \right) = \sum_{\pi \in S} e^{\text{wt}(\pi)}, \quad T_i \left( \sum_{\pi \in S} e^{\text{wt}(\pi)} \right) = 0, \quad (6.4)$$

$$D_i e^{\text{wt}(\pi_{\text{L}})} = \sum_{\pi \in S} e^{\text{wt}(\pi)}, \quad T_i e^{\text{wt}(\pi_{\text{L}})} = \sum_{\pi \in S \setminus \{\pi_{\text{L}}\}} e^{\text{wt}(\pi)}, \quad (6.5)$$

$$D_i e^{\text{wt}(\pi_{\text{H}})} = - \sum_{\pi \in S \setminus \{\pi_{\text{H}}, \pi_{\text{L}}\}} e^{\text{wt}(\pi)}, \quad T_i e^{\text{wt}(\pi_{\text{H}})} = - \sum_{\pi \in S \setminus \{\pi_{\text{L}}\}} e^{\text{wt}(\pi)}. \quad (6.6)$$

Let  $y \in W_{\text{af}}$  and  $i \in I_{\text{af}}$  be such that  $s_i y \prec y$ , or equivalently,  $y^{-1} \alpha_i \in \Delta^- + \mathbb{Z} \delta$  (see Lemma 2.5); note that  $\langle y\mu, \alpha_i^\vee \rangle \leq 0$ , and that  $\langle y\mu, \alpha_i^\vee \rangle = 0$  if and only if  $\Pi^{J_\mu}(s_i y) = \Pi^{J_\mu}(y)$ . It follows from [NS3, Proposition 5.3.2] that the set  $\mathbb{B}_{\frac{\infty}{2}}(\mu) \cup \{\mathbf{0}\}$  is stable under the action of the root operator  $e_i$ , and that

$$\mathbb{B}_{\frac{\infty}{2}}(\mu) = \{e_i^k \pi \mid \pi \in \mathbb{B}_{\frac{\infty}{2}}(\mu), 0 \leq k \leq \varepsilon_i(\pi)\}. \quad (6.7)$$

Using Corollary 6.4 and (6.7), (6.5), together with Remark 6.2, we can easily show the following proposition (see also [L1, Sect. 5.5] and [Kas2, Proposition 9.2.3]).

**Proposition 6.6.** *Let  $y \in W_{\text{af}}$  and  $i \in I_{\text{af}}$  be such that  $s_i y \prec y$ . Then,*

$$\text{gch } V_{s_i y}^-(\mu) = D_i \text{gch } V_y^-(\mu). \quad (6.8)$$

*Remark 6.7.* Since  $D_i^2 = D_i$ , we deduce that for  $y \in W_{\text{af}}$  such that  $s_i y \succ y$ ,

$$D_i \text{gch } V_y^-(\mu) = \text{gch } V_y^-(\mu). \quad (6.9)$$

Also, by (6.8) and (6.9), we have

$$\mathbb{T}_i \text{gch } V_y^-(\mu) = \begin{cases} \text{gch } V_{s_i y}^-(\mu) - \text{gch } V_y^-(\mu) & \text{if } s_i y \prec y, \\ 0 & \text{if } s_i y \succ y. \end{cases} \quad (6.10)$$

**6.2 String property for Demazure-like subsets of  $\mathbb{B}^{\frac{\infty}{2}}(\mu)$ .** We take an arbitrary  $\mu \in P^+$ , and define  $J_\mu \subset I$  as in (3.1). For  $u, v \in W_{\text{af}}$ , we set

$$\mathbb{B}_{\substack{\iota(\bullet, v)=u \\ \kappa(\bullet) \succeq \Pi^{J_\mu}(v)}}^{\frac{\infty}{2}}(\mu) := \{ \pi \in \mathbb{B}^{\frac{\infty}{2}}(\mu) \mid \kappa(\pi) \succeq \Pi^{J_\mu}(v), \iota(\pi, v) = u \}. \quad (6.11)$$

Fix  $i \in I_{\text{af}}$ . For an  $i$ -string  $S \subset \mathbb{B}^{\frac{\infty}{2}}(\mu)$ , we set

$$S_{u, v} := \mathbb{B}_{\substack{\iota(\bullet, v)=u \\ \kappa(\bullet) \succeq \Pi^{J_\mu}(v)}}^{\frac{\infty}{2}}(\mu) \cap S. \quad (6.12)$$

**Proposition 6.8.** *Keep the notation and setting above. In addition, assume that  $s_i u \succ u$  and  $s_i v \succ v$ . Then we have the following table:*

		Case 1		Case 2		Case 3
	$\#S$	$\geq 2$		$\geq 1$		$\geq 1$
(i)	$S_{u, v}$	$S$		$\{\pi_H\}$		$\emptyset$
(ii)	$S_{s_i u, v}$	$\emptyset$		$S \setminus \{\pi_H\}$		$\emptyset$
(iii)	$S_{u, s_i v}$	$\{\pi_L\}$	$S$	$\emptyset$	$\{\pi_H\}$	$\emptyset$
(iv)	$S_{s_i u, s_i v}$	$\emptyset$	$\emptyset$	$\{\pi_L\}$	$S \setminus \{\pi_H\}$	$\emptyset$
		Case 1.1	Case 1.2	Case 2.1	Case 2.2	Case 3

(6.13)

*Proof.* We set  $w := \kappa(\pi_H)$  and  $N := \varphi_i(\pi_H) = \varepsilon_i(\pi_L) \geq 0$ . Since  $s_i v \succ v$  by the assumption, we see by Corollary 6.4 and (6.7) that

$$\mathbb{B}_{\succeq v}^{\frac{\infty}{2}}(\mu) \cap S = S \quad \text{or} \quad \emptyset. \quad (6.14)$$

Also, we see by (6.7) that  $\mathbb{B}_{\succeq v}^{\frac{\infty}{2}}(\mu) \supset \mathbb{B}_{\succeq s_i v}^{\frac{\infty}{2}}(\mu)$ . Therefore,

$$\mathbb{B}_{\succeq v}^{\frac{\infty}{2}}(\mu) \cap S = \emptyset \quad \Rightarrow \quad \mathbb{B}_{\succeq s_i v}^{\frac{\infty}{2}}(\mu) \cap S = \emptyset. \quad (6.15)$$

Assume that  $\mathbb{B}_{\succeq v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ . If  $\langle w\mu, \alpha_i^\vee \rangle \leq 0$  (resp., if  $\langle w\mu, \alpha_i^\vee \rangle > 0$  and  $w \succeq \Pi^{J_\mu}(s_i v)$ ), then we deduce from Lemma 6.3 and Lemma 2.7(1) (resp., from Lemma 6.3) that  $\mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ , that is,

$$\left. \begin{array}{l} \langle w\mu, \alpha_i^\vee \rangle \leq 0, \quad \text{or} \\ \langle w\mu, \alpha_i^\vee \rangle > 0 \text{ and } w \succeq \Pi^{J_\mu}(s_i v) \end{array} \right\} \Rightarrow \mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}. \quad (6.16)$$

Also, it follows from Lemma 6.3 and Lemma 2.7(3) that

$$\langle w\mu, \alpha_i^\vee \rangle > 0 \text{ and } w \not\succeq \Pi^{J_\mu}(s_i v) \Rightarrow \mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \{\pi_L\}. \quad (6.17)$$

**Claim 6.8.1.** *Let  $y \in W_{\text{af}}$  be such that  $\mathbb{B}_{\succeq y}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ .*

- (1) *If  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , then  $\iota(\pi_H, y)^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ .*
- (2) *If  $N \geq 1$ , then  $\iota(\pi_H, y)^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$  and  $\iota(f_i \pi_H, y) \in \{\iota(\pi_H, y), s_i \iota(\pi_H, y)\}$ .*
- (3) *If  $N \geq 2$ , then*

$$\iota(f_i \pi_H, y) = \iota(f_i^2 \pi_H, y) = \cdots = \iota(f_i^{N-1} \pi_H, y) = \underbrace{\iota(f_i^N \pi_H, y)}_{=\pi_L}. \quad (6.18)$$

*Proof of Claim 6.8.1.* Since  $\varepsilon_i(\pi_H) = 0$ , part (2) follows from Lemma 4.5, applied to the case that  $\nu = \mu$  and  $\psi = \pi_H$ . Also, part (3) follows from Lemma 4.5, applied to the case that  $\nu = \mu$  and  $\psi = f_i^k \pi_H$  for each  $1 \leq k \leq N-1$ ; note that  $\varepsilon_i(f_i^k \pi_H) \geq 1$  if  $k \geq 1$ . Hence it remains to show part (1) in the case that  $N = 0$ . Let  $\Lambda \in P^+$  be an arbitrary regular and dominant weight, that is,  $\langle \Lambda, \alpha_i^\vee \rangle > 0$  for all  $i \in I$ , or equivalently,  $J_\Lambda = \emptyset$ . We set  $\eta := (y; 0, 1) \in \mathbb{B}_{\succeq y}^{\infty}(\Lambda)$ ; we see that  $\iota(\eta, y) = y$ , and hence  $\pi_H \otimes \eta \in \mathbb{S}_{\succeq y}^{\infty}(\mu + \Lambda) \subset \mathbb{B}_{\succeq y}^{\infty}(\mu) \otimes \mathbb{B}_{\succeq y}^{\infty}(\Lambda)$  by Theorem 4.3(1). Since  $y^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$  by the assumption, we see that  $\varepsilon_i(\pi_H \otimes \eta) = 0$  and  $\varphi_i(\pi_H \otimes \eta) \geq 1$  by the tensor product rule for crystals. Set  $\psi := \Phi_{\mu\Lambda}^{-1}(\pi_H \otimes \eta) \in \mathbb{B}_{\succeq y}^{\infty}(\mu + \Lambda)$ . Since  $\varepsilon_i(\psi) = 0$  and  $\varphi_i(\psi) \geq 1$ , it follows from Lemma 4.5 that  $\iota(\psi, y)^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ . By Proposition 4.4 and the fact that  $\iota(\eta, y) = y$  seen above, we deduce that  $\iota(\psi, y) = \iota(\pi_H, \iota(\eta, y)) = \iota(\pi_H, y)$ . Hence we obtain  $\iota(\pi_H, y)^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , as desired.  $\blacksquare$

**Claim 6.8.2.** *Assume that  $\mathbb{B}_{\succeq v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ ; note that  $\pi_L \in \mathbb{B}_{\succeq s_i v}^{\infty}(\mu)$  by (6.16) and (6.17). If  $N \geq 1$ , then  $\iota(\pi_L, s_i v) = \iota(\pi_L, v)$ , and hence*

$$\begin{aligned} & \overbrace{\iota(f_i \pi_H, v) = \iota(f_i^2 \pi_H, v) = \cdots = \iota(f_i^{N-1} \pi_H, v)}^{\text{if } N \geq 2; \text{ see (6.18)}} = \underbrace{\iota(f_i^N \pi_H, v)}_{=\pi_L} \\ & \underbrace{\iota(f_i \pi_H, s_i v) = \iota(f_i^2 \pi_H, s_i v) = \cdots = \iota(f_i^{N-1} \pi_H, s_i v)}_{\text{if } N \geq 2 \text{ and } \mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}; \text{ see (6.16) and (6.18)}} = \underbrace{\iota(f_i^N \pi_H, s_i v)}_{=\pi_L}. \end{aligned} \quad (6.19)$$

*Proof of Claim 6.8.2.* Take an arbitrary regular and dominant weight  $\Lambda \in P^+$ ; note that  $\langle v\Lambda, \alpha_i^\vee \rangle > 0$  since  $s_i v \succ v$  by the assumption. Set  $\pi_v := (v; 0, 1) \in \mathbb{B}_{\geq v}^{\infty}(\Lambda)$ . Since  $s_i v \succ v$ , we see by Theorem 4.3 (1) that  $\pi_L \otimes \pi_v \in \mathbb{S}_{\geq v}^{\infty}(\mu + \Lambda) \subset \mathbb{B}_{\geq v}^{\infty}(\mu) \otimes \mathbb{B}_{\geq v}^{\infty}(\Lambda)$ ; we set  $\psi := \Phi_{\mu\Lambda}^{-1}(\pi_L \otimes \pi_v) \in \mathbb{B}_{\geq v}^{\infty}(\mu + \Lambda)$ . Recall that  $N = \varepsilon_i(\pi_L) \geq 1$  by the assumption, which implies that  $\varepsilon_i(\psi) = \varepsilon_i(\pi_L \otimes \pi_v) \geq 1$  by the tensor product rule for crystals. Also, we have  $\varphi_i(\pi_v) = \langle v\Lambda, \alpha_i^\vee \rangle > 0$ , which implies that  $\varphi_i(\psi) \geq 1$ , and hence  $f_i \psi \neq \mathbf{0}$ . Therefore, by Lemma 4.5, we have

$$\iota(\psi, v) = \iota(f_i \psi, v). \quad (6.20)$$

Here it follows from Proposition 4.4 (1) that  $\iota(\psi, v) = \iota(\pi_L, \iota(\pi_v, v))$ ; it is easily checked that  $\iota(\pi_v, v) = v$ . Hence we obtain

$$\iota(\psi, v) = \iota(\pi_L, v). \quad (6.21)$$

Observe that  $\Phi_{\mu\Lambda}(f_i \psi) = f_i(\pi_L \otimes \pi_v) = \pi_L \otimes (f_i \pi_v)$  by the tensor product rule for crystals, where  $f_i \pi_v = (s_i v, v; 0, a, 1)$  for some  $0 < a < 1$ . It follows from Proposition 4.4 (1) that  $\iota(f_i \psi, v) = \iota(\pi_L, \iota(f_i \pi_v, v))$ ; it is easily checked that  $\iota(f_i \pi_v, v) = s_i v$ . Hence we obtain

$$\iota(f_i \psi, v) = \iota(\pi_L, s_i v). \quad (6.22)$$

From (6.20), (6.21), and (6.22), we conclude that  $\iota(\pi_L, v) = \iota(\pi_L, s_i v)$ , as desired.  $\blacksquare$

**Claim 6.8.3.** *Assume that  $\mathbb{B}_{\geq v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ . If  $N = 0$ , then  $\iota(\pi_L, s_i v)$  is identical to  $\iota(\pi_L, v)$  or  $s_i \iota(\pi_L, v)$ .*

*Proof of Claim 6.8.3.* The proof is similar to that of Claim 6.8.2; instead of (6.20), we can show by Lemma 4.5 that

$$\iota(f_i \psi, v) \in \{\iota(\psi, v), s_i \iota(\psi, v)\}. \quad (6.23)$$

Also, equations (6.21) and (6.22) hold also in this case. Substituting (6.21) and (6.22) into (6.23), we obtain  $\iota(\pi_L, s_i v) \in \{\iota(\pi_L, v), s_i \iota(\pi_L, v)\}$ , as desired.  $\blacksquare$

In order to show that  $\mathbf{S}_{u,v} = \mathbf{S}$ ,  $\{\pi_H\}$ , or  $\emptyset$ , let us assume that  $\mathbf{S}_{u,v} \neq \emptyset$ ; by (6.14), we have  $\mathbb{B}_{\geq v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ . If  $\#\mathbf{S} = 1$  (or equivalently,  $N = 0$ ), then it is obvious that  $\mathbf{S}_{u,v} = \mathbf{S} = \{\pi_H\}$ . Assume now that  $\#\mathbf{S} \geq 2$  (or equivalently,  $N \geq 1$ ). Since  $s_i u \succ u$  by the assumption, we see from Claim 6.8.1 that either (6.24) or (6.25) below holds:

$$\underbrace{\iota(\pi_H, v) = \iota(f_i \pi_H, v) = \cdots = \iota(f_i^N \pi_H, v)}_{=u}; \quad (6.24)$$

$$\underbrace{\iota(\pi_H, v)}_{=u} \prec \underbrace{\iota(f_i \pi_H, v) = \cdots = \iota(f_i^N \pi_H, v)}_{=s_i u}. \quad (6.25)$$

If (6.24) (resp., (6.25)) holds, then  $\mathbf{S}_{u,v} = \mathbf{S}$  (resp.,  $\mathbf{S}_{u,v} = \{\pi_H\}$ ). Thus we have proved that  $\mathbf{S}_{u,v} = \mathbf{S}$ ,  $\{\pi_H\}$ , or  $\emptyset$ , as desired. Hence we have the following possibilities (Cases 1, 2a, 2b, and 3).

**Case 1.** Assume that  $\#\mathbf{S} \geq 2$  and  $\mathbf{S}_{u,v} = \mathbf{S}$ . Since (6.24) holds in this case, it is obvious that  $\mathbf{S}_{s_i u, v} = \emptyset$ . Recall that  $w = \kappa(\pi_{\mathbf{H}})$ .

**Subcase 1.1.** Assume that  $\langle w\mu, \alpha_i^{\vee} \rangle > 0$  and  $w \not\geq \Pi^{J\mu}(s_i v)$ . It follows from (6.17) that  $\mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \{\pi_{\mathbf{L}}\}$ . By Claim 6.8.2 and (6.24), we have  $\iota(\pi_{\mathbf{L}}, s_i v) = u$ . Hence we conclude that  $\mathbf{S}_{u, s_i v} = \{\pi_{\mathbf{L}}\}$  and  $\mathbf{S}_{s_i u, s_i v} = \emptyset$ .

**Subcase 1.2.** Assume that  $\langle w\mu, \alpha_i^{\vee} \rangle \leq 0$ , or that  $\langle w\mu, \alpha_i^{\vee} \rangle > 0$  and  $w \succeq \Pi^{J\mu}(s_i v)$ . It follows from (6.16) that  $\mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ . By Claim 6.8.1 (2), (6.19), (6.24), and the assumption  $s_i u \succ u$ , we deduce that

$$\underbrace{\iota(\pi_{\mathbf{H}}, s_i v) = \iota(f_i \pi_{\mathbf{H}}, s_i v) = \cdots = \iota(f_i^N \pi_{\mathbf{H}}, s_i v)}_{=u}. \quad (6.26)$$

Therefore, we conclude that  $\mathbf{S}_{u, s_i v} = \mathbf{S}$  and  $\mathbf{S}_{s_i u, s_i v} = \emptyset$ .

**Case 2a.** Assume that  $\#\mathbf{S} \geq 2$  and  $\mathbf{S}_{u,v} = \{\pi_{\mathbf{H}}\}$ . Since (6.25) holds in this case, it is obvious that  $\mathbf{S}_{s_i u, v} = \mathbf{S} \setminus \{\pi_{\mathbf{H}}\}$ .

**Subcase 2a.1.** Assume that  $\langle w\mu, \alpha_i^{\vee} \rangle > 0$  and  $w \not\geq \Pi^{J\mu}(s_i v)$ . It follows from (6.17) that  $\mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \{\pi_{\mathbf{L}}\}$ . By Claim 6.8.2 and (6.25), we have  $\iota(\pi_{\mathbf{L}}, s_i v) = s_i u$ . Hence we conclude that  $\mathbf{S}_{u, s_i v} = \emptyset$  and  $\mathbf{S}_{s_i u, s_i v} = \{\pi_{\mathbf{L}}\}$ .

**Subcase 2a.2.** Assume that  $\langle w\mu, \alpha_i^{\vee} \rangle \leq 0$ , or that  $\langle w\mu, \alpha_i^{\vee} \rangle > 0$  and  $w \succeq \Pi^{J\mu}(s_i v)$ . It follows from (6.16) that  $\mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$ . By Claim 6.8.1 (2), (6.19), (6.25), and the assumption  $s_i u \succ u$ , we deduce that

$$\underbrace{\iota(\pi_{\mathbf{H}}, s_i v)}_{=u} \prec \underbrace{\iota(f_i \pi_{\mathbf{H}}, s_i v) = \cdots = \iota(f_i^N \pi_{\mathbf{H}}, s_i v)}_{=s_i u}. \quad (6.27)$$

Therefore, we conclude that  $\mathbf{S}_{u, s_i v} = \{\pi_{\mathbf{H}}\}$  and  $\mathbf{S}_{s_i u, s_i v} = \mathbf{S} \setminus \{\pi_{\mathbf{H}}\}$ .

**Case 2b.** Assume that  $\#\mathbf{S} = 1$  (or equivalently,  $N = 0$ ; in this case,  $\mathbf{S} = \{\pi_{\mathbf{H}}\} = \{\pi_{\mathbf{L}}\}$ ) and  $\mathbf{S}_{u,v} = \{\pi_{\mathbf{H}}\} = \mathbf{S}$ . It is obvious that  $\mathbf{S}_{s_i u, v} = \emptyset = \mathbf{S} \setminus \{\pi_{\mathbf{H}}\}$ .

Since  $\mathbf{S} = \{\pi_{\mathbf{H}}\} = \{\pi_{\mathbf{L}}\}$ , we have  $\mathbb{B}_{\succeq s_i v}^{\infty}(\mu) \cap \mathbf{S} = \mathbf{S}$  by (6.16) and (6.17). Also, it follows from Claim 6.8.3 that  $\iota(\pi_{\mathbf{L}}, s_i v)$  is identical to  $\iota(\pi_{\mathbf{L}}, v)$  or  $s_i \iota(\pi_{\mathbf{L}}, v)$ ; in this case,  $\pi_{\mathbf{L}} = \pi_{\mathbf{H}}$  and  $\iota(\pi_{\mathbf{H}}, v) = u$ . Therefore,  $\iota(\pi_{\mathbf{H}}, s_i v) \in \{u, s_i u\}$ .

**Subcase 2b.1.** If  $\iota(\pi_{\mathbf{H}}, s_i v) = s_i u$ , then  $\mathbf{S}_{u, s_i v} = \emptyset$  and  $\mathbf{S}_{s_i u, s_i v} = \{\pi_{\mathbf{H}}\} = \{\pi_{\mathbf{L}}\}$ .

**Subcase 2b.2.** If  $\iota(\pi_{\mathbf{H}}, s_i v) = u$ , then  $\mathbf{S}_{u, s_i v} = \{\pi_{\mathbf{H}}\}$  and  $\mathbf{S}_{s_i u, s_i v} = \emptyset = \mathbf{S} \setminus \{\pi_{\mathbf{H}}\}$ .

**Case 3.** Assume that  $S_{u,v} = \emptyset$ ; we show that  $S_{s_i u, v} = S_{u, s_i v} = S_{s_i u, s_i v} = \emptyset$ . By (6.14) and (6.15), we may (and do) assume that  $\mathbb{B}_{\succeq v}^{\infty}(\mu) \cap S = S$ . We set  $z := \iota(\pi_H, v)$ ; remark that  $z \neq u$  since  $S_{u,v} = \emptyset$ . We see from Claim 6.8.1 that  $z^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , and  $\iota(f_i^k \pi_H, v) \in \{z, s_i z\}$  for all  $0 \leq k \leq N$ . Therefore, if at least one of  $S_{s_i u, v}$ ,  $S_{u, s_i v}$ , and  $S_{s_i u, s_i v}$  is nonempty, then  $z$  is identical to  $u$  or  $s_i u$ . Since  $z^{-1}\alpha_i \in \Delta^+ + \mathbb{Z}\delta$ , and  $s_i u \succ u$  by the assumption, we must have  $z = u$ , which is a contradiction.

This completes the proof of Proposition 6.8.  $\square$

## 7 Proof of Theorem 3.9: part 2.

**7.1 Recursion formula for the right-hand side of (3.21).** Let us take an arbitrary  $\mu \in P^+$ . Recall from (3.10) and (3.11) the definition and some properties of the dual path  $\pi^* \in \mathbb{B}_{\succeq}^{\infty}(-w_\circ \mu)$  for  $\pi \in \mathbb{B}_{\succeq}^{\infty}(\mu)$ . We deduce by Lemmas 2.10 and 3.5 that for  $x, y \in W_{\text{af}}$  such that  $y \succeq x$ ,

$$\sum_{\substack{\pi \in \mathbb{B}_{\succeq}^{\infty}(\mu) \\ \iota(\pi) \preceq \Pi^J \mu(y), \kappa(\pi, y) = x}} e^{-\text{wt}(\pi)} = \sum_{\substack{\eta \in \mathbb{B}_{\succeq}^{\infty}(-w_\circ \mu) \\ \kappa(\eta) \succeq \Pi^J -w_\circ \mu(yw_\circ), \iota(\eta, yw_\circ) = xw_\circ}} e^{\text{wt}(\eta)} =: \mathbf{a}_\mu(x, y). \quad (7.1)$$

The next lemma follows from Proposition 6.8 (applied to the case that  $u = xw_\circ$  and  $v = yw_\circ$ ) and Lemma 6.5, together with Remark 6.2.

**Lemma 7.1.** *Let  $x, y \in W_{\text{af}}$  be such that  $y \succeq x$ , and  $s_i x \prec x$ ,  $s_i y \prec y$ . We have*

$$\mathbf{a}_\mu(s_i x, y) = -s_i \mathbf{T}_i \mathbf{a}_\mu(x, y), \quad (7.2)$$

$$\mathbf{a}_\mu(s_i x, s_i y) = -s_i \mathbf{T}_i \mathbf{a}_\mu(x, s_i y) + s_i (\mathbf{a}_\mu(x, y) - \mathbf{a}_\mu(x, s_i y)), \quad (7.3)$$

$$s_i \mathbf{a}_\mu(s_i x, y) + s_i \mathbf{a}_\mu(x, y) = \mathbf{a}_\mu(s_i x, y) + \mathbf{a}_\mu(x, y), \quad (7.4)$$

$$s_i \mathbf{a}_\mu(s_i x, s_i y) + s_i \mathbf{a}_\mu(x, y) = \mathbf{a}_\mu(s_i x, s_i y) + \mathbf{a}_\mu(x, y), \quad (7.5)$$

$$s_i \mathbf{a}_\mu(s_i x, y) - s_i \mathbf{a}_\mu(s_i x, s_i y) = \mathbf{a}_\mu(s_i x, y) - \mathbf{a}_\mu(s_i x, s_i y). \quad (7.6)$$

For simplicity of notation, we set

$$(W_{\text{af}})_{\succeq x} := \{y \in W_{\text{af}} \mid y \succeq x\} \quad \text{for } x \in W_{\text{af}},$$

$$\mathbf{v}_\lambda(x, y) := (-1)^{\ell_{\frac{\infty}{2}}(y) - \ell_{\frac{\infty}{2}}(x)} \text{gch } V_y^-(\lambda) \quad \text{for } \lambda \in P^+ \text{ and } x, y \in W_{\text{af}}.$$

Also, for  $\lambda, \mu \in P^+$  such that  $\lambda - \mu \in P^+$ , we denote by  $\mathbf{F}(x) = \mathbf{F}_{\lambda\mu}(x)$  the right-hand side of (3.21); we see that

$$\mathbf{F}(x) = \sum_{y \in (W_{\text{af}})_{\succeq x}} \mathbf{a}_\mu(x, y) \mathbf{v}_\lambda(x, y). \quad (7.7)$$

**Proposition 7.2** (cf. (6.9) and (6.10)). *Let  $\lambda, \mu \in P^+$  be such that  $\lambda - \mu \in P^+$ . For  $x \in W_{\text{af}}$  and  $i \in I_{\text{af}}$  such that  $s_i x \prec x$ , it holds that  $\mathbf{T}_i \mathbf{F}_{\lambda\mu}(x) = \mathbf{F}_{\lambda\mu}(s_i x) - \mathbf{F}_{\lambda\mu}(x)$ , and hence  $\mathbf{F}_{\lambda\mu}(s_i x) = \mathbf{D}_i \mathbf{F}_{\lambda\mu}(x)$ .*



*Proof.* We set

$$(W_{\text{af}})_{\succeq x}^+ := \{y \in (W_{\text{af}})_{\succeq x} \mid s_i y \succ y\}, \quad (W_{\text{af}})_{\succeq x}^- := \{y \in (W_{\text{af}})_{\succeq x} \mid s_i y \prec y\};$$

we have

$$\mathbf{F}(x) = \mathbf{F}^+(x) + \mathbf{F}^-(x), \quad \text{where} \quad \mathbf{F}^\pm(x) := \sum_{y \in (W_{\text{af}})_{\succeq x}^\pm} \mathbf{a}_\mu(x, y) \mathbf{v}_\lambda(x, y).$$

We compute as follows:

$$\begin{aligned} \mathbb{T}_i \mathbf{F}^+(x) &\stackrel{(6.3)}{=} \sum_{y \in (W_{\text{af}})_{\succeq x}^+} (\mathbb{T}_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, y) + \sum_{y \in (W_{\text{af}})_{\succeq x}^+} (s_i \mathbf{a}_\mu(x, y)) \underbrace{(\mathbb{T}_i \mathbf{v}_\lambda(x, y))}_{=0 \text{ by (6.10)}} \\ &\stackrel{(7.3)}{=} \sum_{y \in (W_{\text{af}})_{\succeq x}^+} (\mathbf{a}_\mu(x, s_i y) - \mathbf{a}_\mu(x, y) - s_i \mathbf{a}_\mu(s_i x, y)) \mathbf{v}_\lambda(x, y) \\ &= - \sum_{y \in (W_{\text{af}})_{\succeq x}^+} \mathbf{a}_\mu(x, y) \mathbf{v}_\lambda(x, y) + \sum_{y \in (W_{\text{af}})_{\succeq x}^+} (\mathbf{a}_\mu(x, s_i y) - s_i \mathbf{a}_\mu(s_i x, y)) \mathbf{v}_\lambda(x, y), \end{aligned}$$

and

$$\begin{aligned} \mathbb{T}_i \mathbf{F}^-(x) &\stackrel{(6.3)}{=} \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (\mathbb{T}_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, y) + \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(x, y)) (\mathbb{T}_i \mathbf{v}_\lambda(x, y)) \\ &= \sum_{y \in (W_{\text{af}})_{\succeq x}^-} \underbrace{(-s_i \mathbf{a}_\mu(s_i x, y))}_{\text{by (7.2)}} \mathbf{v}_\lambda(x, y) - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(x, y)) \underbrace{(\mathbf{v}_\lambda(x, s_i y) + \mathbf{v}_\lambda(x, y))}_{\text{by (6.10)}} \\ &= - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(s_i x, y) + s_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, y) - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, s_i y) \\ &= - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} \underbrace{(\mathbf{a}_\mu(s_i x, y) + \mathbf{a}_\mu(x, y))}_{\text{by (7.4)}} \mathbf{v}_\lambda(x, y) - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, s_i y) \\ &= - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(x, y) - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} \mathbf{a}_\mu(x, y) \mathbf{v}_\lambda(x, y) \\ &\quad - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, s_i y). \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \mathbb{T}_i \mathbf{F}(x) &= \mathbb{T}_i \mathbf{F}^+(x) + \mathbb{T}_i \mathbf{F}^-(x) \\ &= -\mathbf{F}(x) + \sum_{y \in (W_{\text{af}})_{\succeq x}^+} (\mathbf{a}_\mu(x, s_i y) - s_i \mathbf{a}_\mu(s_i x, y)) \mathbf{v}_\lambda(x, y) \end{aligned}$$

$$- \sum_{y \in (W_{\text{af}})_{\succeq x}^-} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(x, y) - \sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, s_i y). \quad (7.8)$$

Since  $s_i x \prec x$  by the assumption, we deduce by Lemma 2.7(1) that  $(W_{\text{af}})_{\succeq s_i x}^- = (W_{\text{af}})_{\succeq x}^-$ . Hence it follows that

$$- \sum_{y \in (W_{\text{af}})_{\succeq x}^-} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(x, y) = \sum_{y \in (W_{\text{af}})_{\succeq s_i x}^-} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(s_i x, y).$$

Also, we see by Lemma 2.7(3) that  $y \in (W_{\text{af}})_{\succeq x}^-$  if and only if  $s_i y \in (W_{\text{af}})_{\succeq s_i x}^+$ . Hence we deduce that

$$\sum_{y \in (W_{\text{af}})_{\succeq x}^-} (s_i \mathbf{a}_\mu(x, y)) \mathbf{v}_\lambda(x, s_i y) = \sum_{y \in (W_{\text{af}})_{\succeq s_i x}^+} (s_i \mathbf{a}_\mu(x, s_i y)) \mathbf{v}_\lambda(x, y).$$

Therefore, the right-hand side of (7.8) plus  $\mathbf{F}(x)$  is identical to

$$\begin{aligned} & \sum_{y \in (W_{\text{af}})_{\succeq s_i x}^-} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(s_i x, y) \\ & + \underbrace{\sum_{y \in (W_{\text{af}})_{\succeq x}^+} (\mathbf{a}_\mu(x, s_i y) - s_i \mathbf{a}_\mu(s_i x, y)) \mathbf{v}_\lambda(x, y) - \sum_{y \in (W_{\text{af}})_{\succeq s_i x}^+} (s_i \mathbf{a}_\mu(x, s_i y)) \mathbf{v}_\lambda(x, y)}_{=:\mathbf{G}(x)}. \end{aligned} \quad (7.9)$$

Because  $(W_{\text{af}})_{\succeq x}^+ \subset (W_{\text{af}})_{\succeq s_i x}^+$  by the assumption that  $s_i x \prec x$ , we see that

$$\begin{aligned} \mathbf{G}(x) &= \sum_{y \in (W_{\text{af}})_{\succeq x}^+} \overbrace{(\mathbf{a}_\mu(x, s_i y) - s_i \mathbf{a}_\mu(s_i x, y) - s_i \mathbf{a}_\mu(x, s_i y))}^{= -\mathbf{a}_\mu(s_i x, y) \text{ by (7.5)}} \mathbf{v}_\lambda(x, y) \\ & - \sum_{\substack{y \in (W_{\text{af}})_{\succeq s_i x}^+ \\ y \not\prec x}} (s_i \mathbf{a}_\mu(x, s_i y)) \mathbf{v}_\lambda(x, y). \end{aligned} \quad (7.10)$$

Here we claim that for  $y \in (W_{\text{af}})_{\succeq s_i x}^+$  such that  $y \not\prec x$ ,

$$s_i \mathbf{a}_\mu(x, s_i y) = \mathbf{a}_\mu(s_i x, y). \quad (7.11)$$

In order to show this equality, it suffices to prove the following claim.

**Claim 7.2.1.** *Let  $y \in (W_{\text{af}})_{\succeq s_i x}^+$  be such that  $y \not\prec x$ . Let  $S \subset \mathbb{B}^{\frac{\infty}{2}}(-w_\circ \mu)$  be an  $i$ -string such that*

$$S_{xw_\circ, s_i y w_\circ} := S \cap \mathbb{B}^{\frac{\infty}{2}}_{\substack{\iota(\bullet, s_i y w_\circ) = xw_\circ \\ \kappa(\bullet) \succeq \Pi^{J-w_\circ \mu}(s_i y w_\circ)}} \quad (-w_\circ \mu) \neq \emptyset.$$

Then we have

$$S_{xw_\circ, s_i y w_\circ} = \{\pi_{\text{H}}^S\} \quad \text{and} \quad S_{s_i x w_\circ, y w_\circ} = \{\pi_{\text{L}}^S\}. \quad (7.12)$$

*Proof of Claim 7.2.1.* Suppose, for a contradiction, that  $S_{xw_\circ, yw_\circ} \neq \emptyset$ . If  $\pi \in S_{xw_\circ, yw_\circ}$ , then  $xw_\circ = \iota(\pi, yw_\circ) \succeq yw_\circ$ , and hence  $y \succeq x$  by Lemma 2.10. However, this contradicts the assumption that  $y \not\succeq x$ . Hence we have  $S_{xw_\circ, yw_\circ} = \emptyset$ . Since  $s_i x w_\circ \succ x w_\circ$  and  $y w_\circ \succ s_i y w_\circ$  by the assumption, the set  $S_{xw_\circ, yw_\circ}$  is determined by row (iii) in table (6.13). Since  $S_{xw_\circ, yw_\circ} = \emptyset$  as seen above, we obtain (7.12) by rows (i) and (iv) in table (6.13). This proves the claim (and hence (7.11)).  $\blacksquare$

Substituting (7.11) into (7.10), we see that

$$\begin{aligned} \mathbf{G}(x) &= - \sum_{y \in (W_{\text{af}})_{\succeq x}^+} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(x, y) - \sum_{\substack{y \in (W_{\text{af}})_{\succeq s_i x}^+ \\ y \not\succeq x}} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(x, y) \\ &= \sum_{y \in (W_{\text{af}})_{\succeq x}^+} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(s_i x, y) + \sum_{\substack{y \in (W_{\text{af}})_{\succeq s_i x}^+ \\ y \not\succeq x}} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(s_i x, y) \\ &= \sum_{y \in (W_{\text{af}})_{\succeq s_i x}^+} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(s_i x, y). \end{aligned}$$

Hence the right-hand side of (7.8) plus  $\mathbf{F}(x)$  (see (7.9)) is identical to

$$\sum_{y \in (W_{\text{af}})_{\succeq s_i x}^-} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(s_i x, y) + \sum_{y \in (W_{\text{af}})_{\succeq s_i x}^+} \mathbf{a}_\mu(s_i x, y) \mathbf{v}_\lambda(s_i x, y) = \mathbf{F}(s_i x).$$

Substituting this equality into (7.8), we conclude that  $\mathbf{T}_i \mathbf{F}(x) = \mathbf{F}(s_i x) - \mathbf{F}(x)$ , as desired. This completes the proof of Proposition 7.2.  $\square$

**7.2 Second step in the proof of Theorem 3.9.** We prove (3.21) in the case that  $\mu = \varpi_r$  for  $r \in I$  (and  $x \in W_{\text{af}}$  is general). Let  $\lambda = \sum_{i \in I} \lambda_i \varpi_i \in P^+$  be such that  $\lambda - \varpi_r \in P^+$ , and let  $x \in W_{\text{af}}$ . We deduce from [AK, Lemma 1.4] (see also [NS3, (1a) and (2a) in the proof of Lemma 5.4.1]) that there exist  $i_1, \dots, i_n \in I_{\text{af}}$  and  $\xi \in Q^\vee$  such that

$$x = s_{i_n} s_{i_{n-1}} \cdots s_{i_2} s_{i_1} t_\xi \prec s_{i_{n-1}} \cdots s_{i_2} s_{i_1} t_\xi \prec \cdots \prec s_{i_2} s_{i_1} t_\xi \prec s_{i_1} t_\xi \prec t_\xi.$$

By Propositions 6.6 and 7.2, we see that

$$\begin{aligned} \text{gch } V_x^-(\lambda - \varpi_r) &= D_{i_n} D_{i_{n-1}} \cdots D_{i_2} D_{i_1} \text{gch } V_{t_\xi}^-(\lambda - \varpi_r), \\ \mathbf{F}_{\lambda \varpi_r}(x) &= D_{i_n} D_{i_{n-1}} \cdots D_{i_2} D_{i_1} \mathbf{F}_{\lambda \varpi_r}(t_\xi), \end{aligned}$$

respectively. Also, in §5.2, we proved that

$$\frac{1}{1 - q^{-\lambda_r}} \text{gch } V_{t_\xi}^-(\lambda - \varpi_r) = \mathbf{F}_{\lambda \varpi_r}(t_\xi).$$

From these equalities, we obtain

$$\frac{1}{1 - q^{-\lambda_r}} \text{gch } V_x^-(\lambda - \varpi_r) = \mathbf{F}_{\lambda \varpi_r}(x),$$

as desired.

## 8 Proof of Theorem 3.9: part 3.

**8.1 Final step in the proof of Theorem 3.9.** Keep the notation and setting of Theorem 3.9. We prove (3.21) by induction on  $|\mu| := \sum_{i \in I} \mu_i \in \mathbb{Z}_{\geq 0}$ . If  $|\mu| = 1$ , that is, if  $\mu = \varpi_r$  for some  $r \in I$ , then we proved (3.21) in §7.2. Assume now that  $|\mu| > 1$ , and take  $r \in I$  such that  $\mu_r \geq 1$ . We set  $\nu := \mu - \varpi_r \in P_+$ . We compute as follows:

$$\begin{aligned}
& \prod_{i \in I} \prod_{k=\lambda_i-\mu_i+1}^{\lambda_i} \frac{1}{1-q^{-k}} \text{gch } V_x^-(\lambda - \mu) = \prod_{i \in I} \prod_{k=\lambda_i-\mu_i+1}^{\lambda_i} \frac{1}{1-q^{-k}} \text{gch } V_x^-(\lambda - \varpi_r - \nu) \\
&= \frac{1}{1-q^{-\lambda_r}} \sum_{\substack{y \in W_{\text{af}} \\ y \succeq x}} \sum_{\substack{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\nu) \\ \iota(\pi) \preceq \Pi^{J\nu}(y), \kappa(\pi, y) = x}} (-1)^{\ell^{\frac{\infty}{2}}(y) - \ell^{\frac{\infty}{2}}(x)} \mathbf{e}^{-\text{wt}(\pi)} \text{gch } V_y^-(\lambda - \varpi_r) \\
&\hspace{15em} \text{by our induction hypothesis} \\
&= \sum_{\substack{y \in W_{\text{af}} \\ y \succeq x}} \sum_{\substack{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\nu) \\ \iota(\pi) \preceq \Pi^{J\nu}(y), \kappa(\pi, y) = x}} (-1)^{\ell^{\frac{\infty}{2}}(y) - \ell^{\frac{\infty}{2}}(x)} \mathbf{e}^{-\text{wt}(\pi)} \times \\
&\hspace{10em} \sum_{\substack{z \in W_{\text{af}} \\ z \succeq y}} \sum_{\substack{\eta \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r) \\ \iota(\eta) \preceq \Pi^{J\varpi_r}(z), \kappa(\eta, z) = y}} (-1)^{\ell^{\frac{\infty}{2}}(z) - \ell^{\frac{\infty}{2}}(y)} \mathbf{e}^{-\text{wt}(\eta)} \text{gch } V_z^-(\lambda) \\
&\hspace{15em} \text{by the formula shown in §7.2} \\
&= \sum_{\substack{z, y \in W_{\text{af}} \\ z \succeq y \succeq x}} \sum_{\substack{\eta \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r) \\ \iota(\eta) \preceq \Pi^{J\varpi_r}(z), \kappa(\eta, z) = y}} \sum_{\substack{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\nu) \\ \iota(\pi) \preceq \Pi^{J\nu}(y), \kappa(\pi, y) = x}} (-1)^{\ell^{\frac{\infty}{2}}(z) - \ell^{\frac{\infty}{2}}(x)} \mathbf{e}^{-(\text{wt}(\pi) + \text{wt}(\eta))} \text{gch } V_z^-(\lambda) \\
&=: \mathbf{H}(x).
\end{aligned}$$

We see by Theorem 4.3 (2) that for each  $z \in W_{\text{af}}$  with  $z \succeq x$ , the set

$$\coprod_{z \succeq y \succeq x} \left( \left\{ \eta \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r) \mid \iota(\eta) \preceq \Pi^{J\varpi_r}(z), \kappa(\eta, z) = y \right\} \times \left\{ \pi \in \mathbb{B}^{\frac{\infty}{2}}(\nu) \mid \iota(\pi) \preceq \Pi^{J\nu}(y), \kappa(\pi, y) = x \right\} \right)$$

is in bijection with the set  $\left\{ \eta \otimes \pi \in \mathbb{S}_{\succeq z}^{\frac{\infty}{2}}(\varpi_r + \nu) \mid \kappa(\pi, \kappa(\eta, z)) = x \right\}$  by the map  $(\eta, \pi) \mapsto \eta \otimes \pi$ . Hence we have

$$\mathbf{H}(x) = \sum_{\substack{z \in W_{\text{af}} \\ z \succeq x}} \sum_{\substack{\eta \otimes \pi \in \mathbb{S}_{\succeq z}^{\frac{\infty}{2}}(\varpi_r + \nu) \\ \kappa(\pi, \kappa(\eta, z)) = x}} (-1)^{\ell^{\frac{\infty}{2}}(z) - \ell^{\frac{\infty}{2}}(x)} \mathbf{e}^{-(\text{wt}(\pi) + \text{wt}(\eta))} \text{gch } V_z^-(\lambda).$$

Moreover, it follows from Proposition 4.4(2) that the set  $\left\{ \eta \otimes \pi \in \mathbb{S}_{\succeq z}^{\frac{\infty}{2}}(\varpi_r + \nu) \mid \kappa(\pi, \kappa(\eta, z)) = x \right\}$  is in bijection with the set  $\left\{ \psi \in \mathbb{B}_{\succeq z}^{\frac{\infty}{2}}(\varpi_r + \nu) \mid \kappa(\psi, z) = x \right\}$  by

the map  $\Phi_{\varpi_r, \nu} : \mathbb{B}_{\frac{\infty}{2}}(\mu) = \mathbb{B}_{\frac{\infty}{2}}(\varpi_r + \nu) \xrightarrow{\sim} \mathbb{S}_{\frac{\infty}{2}}(\varpi_r + \nu) \hookrightarrow \mathbb{B}_{\frac{\infty}{2}}(\varpi_r) \otimes \mathbb{B}_{\frac{\infty}{2}}(\nu)$ . Therefore, we conclude that

$$\mathbf{H}(x) = \sum_{\substack{z \in W_{\text{af}} \\ z \succeq x}} \sum_{\substack{\psi \in \mathbb{B}_{\frac{\infty}{2}}(\mu) \\ \iota(\psi) \preceq \Pi^{J\mu}(z), \kappa(\psi, z) = x}} (-1)^{\ell_{\frac{\infty}{2}}(z) - \ell_{\frac{\infty}{2}}(x)} \mathbf{e}^{-\text{wt}(\psi)} \text{gch } V_z^-(\lambda).$$

This completes the proof of Theorem 3.9.

**8.2 Proof of Corollary 3.15.** We take and fix  $x \in W$ . Let  $\mathbb{X}$  be the subset of  $\mathbb{B}_{\frac{\infty}{2}}(\mu) \times W_{\text{af}}$  consisting of those elements  $(\pi, y)$  satisfying the conditions that  $\iota(\pi) \preceq \Pi^{J\mu}(y)$  and  $\kappa(\pi, y) = x$ ; note that  $y \succeq \kappa(\pi, y) = x$  if  $(\pi, y) \in \mathbb{X}$ . Then the right-hand side of (3.21) can be rewritten as:

$$\sum_{(\pi, y) \in \mathbb{X}} (-1)^{\ell_{\frac{\infty}{2}}(y) - \ell_{\frac{\infty}{2}}(x)} \mathbf{e}^{-\text{wt}(\pi)} \text{gch } V_y^-(\lambda). \quad (8.1)$$

Also, let  $\mathbb{Y}$  be the subset of  $\text{QLS}(\mu) \times W$  consisting of those elements  $(\eta, v)$  satisfying the condition that  $\kappa(\eta, v) = x$ . We define the map  $\text{cl} : \mathbb{B}_{\frac{\infty}{2}}(\mu) \times W_{\text{af}} \rightarrow \text{QLS}(\mu) \times W$  by  $\text{cl}(\pi, y) := (\text{cl}(\pi), \text{cl}(y))$  for  $(\pi, y) \in \mathbb{B}_{\frac{\infty}{2}}(\mu) \times W_{\text{af}}$  (for the map  $\text{cl} : W_{\text{af}} \rightarrow W$ , see §3.4). We claim that  $\text{cl}(\mathbb{X}) = \mathbb{Y}$ .

Let  $(\eta, v) \in \text{QLS}(\mu) \times W$ , and let  $(\pi, y) \in \mathbb{B}_{\frac{\infty}{2}}(\mu) \times W_{\text{af}}$ . Write  $\pi \in \mathbb{B}_{\frac{\infty}{2}}(\mu)$  as:

$$\begin{cases} \pi = (x_1, \dots, x_s; a_0, a_1, \dots, a_s), & \text{with} \\ x_u = w_u \Pi^{J\mu}(t_{\xi_u}) & \text{for } w_u \in W^{J\mu} \text{ and } \xi_u \in Q^\vee, \quad 1 \leq u \leq s. \end{cases} \quad (8.2)$$

Assume that  $\text{cl}(\pi, y) = (\eta, v)$ . If we take  $1 \leq u_1 < \dots < u_r = s$  in such a way that

$$\begin{aligned} w_1 = \dots = w_{u_1} \neq w_{u_1+1} = \dots = w_{u_2} \neq w_{u_2+1} = \dots \\ \dots \neq w_{u_{r-1}+1} = \dots = w_{u_r} (= w_s), \end{aligned}$$

then  $\eta \in \text{QLS}(\mu)$  is of the form  $\eta = (w_{u_1}, \dots, w_{u_r}; a_0, a_{u_1}, \dots, a_{u_r})$ . Since  $\text{cl}(y) = v$ , we see that  $y \in W_{\text{af}}$  is of the form  $y = vt_\zeta$  for some  $\zeta \in Q^\vee$ . If  $\iota(\pi) \preceq \Pi^{J\mu}(y)$ , then we deduce by Proposition 2.26 that

$$\kappa(\pi, y) = \kappa(\eta, v) \cdot t_{[\xi_s]^{J\mu} + [\zeta - \zeta(\eta, v)]_{J\mu}}. \quad (8.3)$$

Moreover, if in addition  $\kappa(\pi, y) = x \in W$ , or equivalently, if  $(\pi, y) \in \mathbb{X}$ , then we see by (8.3) that  $\kappa(\eta, v) = x$  since  $\kappa(\eta, v) \in W$ . Thus we obtain  $(\eta, v) \in \mathbb{Y}$ , which implies that  $\text{cl}(\mathbb{X}) \subset \mathbb{Y}$ .

By exactly the same argument as for Lemma 3.12 (that is, as for [NS3, Lemma 6.2.3]; see also [LNS<sup>3</sup>3, Lemma 2.3.2]), we can show that for each  $\eta \in \text{QLS}(\mu)$  and  $\chi \in \text{Par}(\mu)$ , there exists a unique element  $\pi_{\chi, \eta} \in \mathbb{B}_{\frac{\infty}{2}}(\mu)$  such that  $\text{cl}(\pi_{\chi, \eta}) = \eta$  and  $\kappa(\pi_{\chi, \eta}) = \kappa(\eta) \in W^{J\mu}$ ; for the definitions of  $\text{Par}(\mu)$  and  $\mathbb{B}_{\frac{\infty}{2}}(\mu)$ , see §4.1. Notice that if  $\chi = (\emptyset)_{i \in I}$ ,

then  $\pi_{\boldsymbol{\chi}, \eta}$  is identical to  $\pi_\eta$  in Lemma 3.12. We deduce by [INS, Lemma 7.1.4] that if  $\iota(\pi_\eta) = w\Pi^{J_\mu}(t_\xi)$  for some  $w \in W^{J_\mu}$  and  $\xi \in Q^\vee$ , then  $\iota(\pi_{\boldsymbol{\chi}, \eta}) = w\Pi^{J_\mu}(t_{\xi + \iota(\boldsymbol{\chi})})$  (for  $\iota(\boldsymbol{\chi})$ , see Remark 4.1). Also, observe that  $\text{wt}(\pi_{\boldsymbol{\chi}, \eta}) = \text{wt}(\pi_\eta) - |\boldsymbol{\chi}|\delta$ .

Fix  $(\eta, v) \in \mathbb{Y}$ . We claim that for  $(\pi, y) \in \mathbb{B}^{\infty}(\mu) \times W_{\text{af}}$ ,

$$\begin{aligned} \text{cl}(\pi, y) = (\eta, v) \text{ and } (\pi, y) \in \mathbb{X} &\iff \\ \left\{ \begin{array}{l} \pi = \pi_{\boldsymbol{\chi}, \eta} \quad \text{for some } \boldsymbol{\chi} \in \text{Par}(\mu), \\ y = vt_{\zeta(\eta, v) + \gamma} \quad \text{for some } \gamma \in Q_{I \setminus J_\mu}^\vee \text{ such that } \gamma \geq \iota(\boldsymbol{\chi}); \end{array} \right. & \quad (8.4) \end{aligned}$$

notice that the opposite inclusion  $\text{cl}(\mathbb{X}) \supset \mathbb{Y}$  follows from this claim. First, we show the implication  $\Rightarrow$ . Write  $\pi \in \mathbb{B}^{\infty}(\mu)$  in the form (8.2). If we write  $y \in W_{\text{af}}$  as  $y = vt_\zeta$  with  $\zeta \in Q^\vee$  (recall that  $\text{cl}(y) = v$ ), then we see by (8.3) that

$$x = \kappa(\pi, y) = \kappa(\eta, v) \cdot t_{[\xi_s]^{J_\mu} + [\zeta - \zeta(\eta, v)]_{J_\mu}} = x \cdot t_{[\xi_s]^{J_\mu} + [\zeta - \zeta(\eta, v)]_{J_\mu}}; \quad (8.5)$$

recall that  $(\pi, y) \in \mathbb{X}$  and  $(\eta, v) \in \mathbb{Y}$ . Therefore, we deduce that  $[\xi_s]^{J_\mu} + [\zeta - \zeta(\eta, v)]_{J_\mu} = 0$ , and hence  $[\xi_s]^{J_\mu} = 0$  and  $[\zeta - \zeta(\eta, v)]_{J_\mu} = 0$ . By the equality  $[\xi_s]^{J_\mu} = 0$  and Lemma 2.1 (3), we have  $\kappa(\pi) = x_s = w_s \Pi^{J_\mu}(t_{\xi_s}) = w_s \in W^{J_\mu}$ . Since  $\text{cl}(\pi) = \eta$  by our assumption, we see by the definition that  $\pi = \pi_{\boldsymbol{\chi}, \eta}$  if  $\pi \in \mathbb{B}^{\infty}_{\boldsymbol{\chi}}(\mu)$  for  $\boldsymbol{\chi} \in \text{Par}(\mu)$  (see the previous paragraph). Since  $\iota(\pi_\eta) = w_1 \Pi^{J_\mu}(t_{\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v)})$  by Remark 3.14, it follows that

$$\iota(\pi) = \iota(\pi_{\boldsymbol{\chi}, \eta}) = w_1 \Pi^{J_\mu}(t_{\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v) + \iota(\boldsymbol{\chi})}). \quad (8.6)$$

Since  $(\pi, y) \in \mathbb{X}$ , we have  $\Pi^{J_\mu}(y) = \Pi^{J_\mu}(vt_\zeta) \succeq \iota(\pi)$ . Hence it follows from Lemmas 2.18 and 2.17 (recall that  $\widehat{w}_1 \in w_1 W_{J_\mu}$ ) that

$$[\zeta]^{J_\mu} \geq [\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v) + \iota(\boldsymbol{\chi}) + \text{wt}(w_1 \Rightarrow v)]^{J_\mu} = [\zeta(\eta, v) + \iota(\boldsymbol{\chi})]^{J_\mu};$$

recall that  $\iota(\boldsymbol{\chi}) \in Q_{I \setminus J_\mu}^\vee$ . Combining this inequality with the equality  $[\zeta - \zeta(\eta, v)]_{J_\mu} = 0$  shown above, we obtain  $\zeta = \zeta(\eta, v) + \gamma$  for some  $\gamma \in Q_{I \setminus J_\mu}^\vee$  such that  $\gamma \geq \iota(\boldsymbol{\chi})$ . Thus we have proved the implication  $\Rightarrow$ . Next, we show the implication  $\Leftarrow$ . It is obvious that  $\text{cl}(\pi) = \text{cl}(\pi_{\boldsymbol{\chi}, \eta}) = \eta$  and  $\text{cl}(y) = v$ . Also, we have  $\iota(\pi) = \iota(\pi_{\boldsymbol{\chi}, \eta}) = w_1 \Pi^{J_\mu}(t_{\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v) + \iota(\boldsymbol{\chi})})$  as shown in (8.6), where  $w_1 = \iota(\eta)$  and  $\widehat{w}_1 = \max(w_1 W_{J_\mu}, \leq_v^*)$ . Because

$$[\zeta(\eta, v) + \gamma]^{J_\mu} \geq [\zeta(\eta, v) + \iota(\boldsymbol{\chi})]^{J_\mu} = [\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v) + \iota(\boldsymbol{\chi}) + \text{wt}(w_1 \Rightarrow [v]^{J_\mu})]^{J_\mu}$$

by Lemma 2.17, it follows from Lemma 2.18 that

$$\begin{aligned} \Pi^{J_\mu}(y) &= \Pi^{J_\mu}(vt_{\zeta(\eta, v) + \gamma}) = [v]^{J_\mu} \Pi^{J_\mu}(t_{\zeta(\eta, v) + \gamma}) \\ &\succeq w_1 \Pi^{J_\mu}(t_{\zeta(\eta, v) - \text{wt}(\widehat{w}_1 \Rightarrow v) + \iota(\boldsymbol{\chi})}) = \iota(\pi_{\boldsymbol{\chi}, \eta}) = \iota(\pi). \end{aligned}$$

Finally, by the same argument as for (8.5) (applied to the case that  $\xi_s = 0$  and  $\zeta = \zeta(\eta, v) + \gamma$ ), we deduce that  $\kappa(\pi, y) = \kappa(\eta, v) \cdot t_{[0]^{J_\mu} + [(\zeta(\eta, v) + \gamma) - \zeta(\eta, v)]_{J_\mu}} = x$ . Thus we have shown the implication  $\Leftarrow$ , thereby completing the proof of (8.4).

By (8.4), we can rewrite (8.1) (which is identical to the right-hand side of (3.21)) as:

$$\begin{aligned}
& \sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(\mu) \\ \kappa(\eta, v) = x}} \sum_{\chi \in \text{Par}(\mu)} \sum_{\substack{\gamma \in Q_{I \setminus J_\mu}^\vee \\ \gamma \geq \iota(\chi)}} (-1)^{\ell \frac{\infty}{2} (vt_{\zeta(\eta, v)} + \gamma) - \ell \frac{\infty}{2} (x)} e^{-\text{wt}(\pi_{\chi, \eta})} \text{gch } V_{vt_{\zeta(\eta, v)} + \gamma}^- (\lambda) \\
&= \sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(\mu) \\ \kappa(\eta, v) = x}} \sum_{\chi \in \text{Par}(\mu)} \sum_{\substack{\gamma \in Q_{I \setminus J_\mu}^\vee \\ \gamma \geq \iota(\chi)}} (-1)^{\ell(v) - \ell(x)} e^{-\text{wt}(\pi_\eta) + |\chi| \delta} \underbrace{q^{-\langle \lambda, \gamma \rangle} \text{gch } V_{vt_{\zeta(\eta, v)}}^- (\lambda)}_{\text{by Proposition 3.8}} \\
&= \sum_{\chi \in \text{Par}(\mu)} \sum_{\substack{\gamma \in Q_{I \setminus J_\mu}^\vee \\ \gamma \geq \iota(\chi)}} q^{|\chi| - \langle \lambda, \gamma \rangle} \underbrace{\sum_{v \in W} \sum_{\substack{\eta \in \text{QLS}(\mu) \\ \kappa(\eta, v) = x}} (-1)^{\ell(v) - \ell(x)} e^{-\text{wt}(\pi_\eta)} \text{gch } V_{vt_{\zeta(\eta, v)}}^- (\lambda)}_{\text{the right-hand side of (3.29)}}.
\end{aligned}$$

Therefore, in order to prove (3.29) in Corollary 3.15, it suffices to show that

$$\sum_{\chi \in \text{Par}(\mu)} \sum_{\substack{\gamma \in Q_{I \setminus J_\mu}^\vee \\ \gamma \geq \iota(\chi)}} q^{|\chi| - \langle \lambda, \gamma \rangle} = \prod_{i \in I} \prod_{k = \lambda_i - \mu_i + 1}^{\lambda_i} \frac{1}{1 - q^{-k}}. \quad (8.7)$$

For each  $i \in I \setminus J_\mu$ , we denote by  $\text{Par}(\mu_i)$  the set of partitions of length less than  $\mu_i \in \mathbb{Z}_{\geq 1}$ . We rewrite (8.7) as:

$$\begin{aligned}
& \prod_{i \in I \setminus J_\mu} \underbrace{\sum_{\substack{\chi^{(i)} = (\chi_1^{(i)} \geq \dots \geq \chi_{\mu_i - 1}^{(i)} \geq 0) \\ \in \text{Par}(\mu_i)}} \sum_{\substack{c_i \in \mathbb{Z} \\ c_i \geq \chi_1^{(i)}}} q^{|\chi^{(i)}| - \lambda_i c_i}}_{= \sum_{(\chi^{(i)}, c_i) \in \mathbb{P}_1^{(i)}} q^{|\chi^{(i)}| - \lambda_i c_i}} = \prod_{i \in I \setminus J_\mu} \underbrace{\prod_{k = \lambda_i - \mu_i + 1}^{\lambda_i} \frac{1}{1 - q^{-k}}}_{= \sum_{\tau \in \mathbb{P}_2^{(i)}} q^{-|\tau|}}
\end{aligned}$$

where for each  $i \in I \setminus J_\mu$ , we set

$$\mathbb{P}_1^{(i)} := \{(\chi^{(i)}, c_i) \mid \chi^{(i)} = (\chi_1^{(i)} \geq \dots \geq \chi_{\mu_i - 1}^{(i)} \geq 0) \in \text{Par}(\mu_i) \text{ and } c_i \in \mathbb{Z}, c_i \geq \chi_1^{(i)}\},$$

$$\mathbb{P}_2^{(i)} := \{\tau = (\tau_1 \geq \tau_2 \geq \dots \geq \tau_m \geq 0) \mid m \geq 0, \lambda_i - \mu_i + 1 \leq \tau_m \leq \tau_1 \leq \lambda_i\};$$

recall that  $\lambda_i \geq \mu_i$  for all  $i \in I \setminus J_\mu$ . We show that for each  $i \in I \setminus J_\mu$ ,

$$\sum_{(\chi^{(i)}, c_i) \in \mathbb{P}_1^{(i)}} q^{|\chi^{(i)}| - \lambda_i c_i} = \sum_{\tau \in \mathbb{P}_2^{(i)}} q^{-|\tau|}. \quad (8.8)$$

Let  $(\chi^{(i)}, c_i) \in \mathbb{P}_1^{(i)}$ , and write  $\chi^{(i)} \in \text{Par}(\mu_i)$  as  $\chi^{(i)} = (\chi_1^{(i)} \geq \dots \geq \chi_{\mu_i - 1}^{(i)} \geq 0)$ . Observe that


$$\underbrace{(c_i, c_i, \dots, c_i, c_i, c_i - \chi_{\mu_i - 1}^{(i)}, \dots, c_i - \chi_2^{(i)}, c_i - \chi_1^{(i)})}_{(\lambda_i - \mu_i + 1) \text{ times}} \quad (8.9)$$

is a partition of length less than or equal to  $\lambda_i$ . We define  $\Psi(\chi^{(i)}, c_i)$  to be the conjugate (or transposed) partition of the partition (8.9); it is easily checked that  $\Psi(\chi^{(i)}, c_i) \in \mathcal{P}_2^{(i)}$ . Also, we can easily check that the map  $\Psi : \mathcal{P}_1^{(i)} \rightarrow \mathcal{P}_2^{(i)}$  is bijective, and  $|\Psi(\chi^{(i)}, c_i)| = -|\chi^{(i)}| + \lambda_i c_i$  for all  $(\chi^{(i)}, c_i) \in \mathcal{P}_1^{(i)}$ . Thus we have shown (8.8), as desired. This completes the proof of Corollary 3.15.

## Appendix.

### A Examples of formula (1.4).

We give some examples of formula (1.4) in the Introduction in the case that  $\mathfrak{g}$  is of type  $A_2$  and  $\mu = \varpi_1$ ; recall that  $\varpi_1$  is a minuscule weight. Note that  $J_\mu = J_{\varpi_1} = \{2\}$ , and  $W^{J_\mu} = W^{J_{\varpi_1}} = \{e, s_1, s_2 s_1\}$ ,  $W_{J_\mu} = W_{J_{\varpi_1}} = \{e, s_2\}$ . The quantum Bruhat graph in type  $A_2$  is as follows:



Note that  $\theta$  denotes the highest root  $\alpha_1 + \alpha_2$ . By direct calculation, we obtain

	$eW_{J_{\varpi_1}}$		$s_1W_{J_{\varpi_1}}$		$s_2s_1W_{J_{\varpi_1}}$	
$e$	$e$	0	$s_1$	$\alpha_1^\vee$	$w_\circ$	$\theta^\vee$
$s_2$	$s_2$	0	$s_1$	$\alpha_1^\vee$	$s_2s_1$	$\alpha_1^\vee$
$s_1$	$e$	0	$s_1$	0	$w_\circ$	$\theta^\vee$
$s_1s_2$	$s_2$	0	$s_1s_2$	0	$w_\circ$	$\alpha_1^\vee$
$s_2s_1$	$s_2$	0	$s_1$	0	$s_2s_1$	0
$w_\circ$	$s_2$	0	$s_1s_2$	0	$w_\circ$	0

(A.1)



where for  $v \in W$  and  $w \in W^{J_{\varpi_1}}$ , we have

$$\begin{array}{|c|c|} \hline & wW_{J_{\varpi_1}} \\ \hline v & \max(wW_{J_{\varpi_1}}, \leq_v^*) =: \widehat{w} \quad \text{wt}(\widehat{w} \Rightarrow v) \\ \hline \end{array}.$$

Since  $\mu = \varpi_1$  is a minuscule weight, we see by (3.32) that

$$\text{QLS}(\mu) = \text{QLS}(\varpi_1) = \{(e; 0, 1), (s_1; 0, 1), (s_2s_1; 0, 1)\}.$$

Let us recall formula (1.4) from the Introduction: for  $w \in W$ ,

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G(s_1)}] \cdot [\mathcal{O}_{\mathbf{Q}_G(w)}] \\ &= [\mathcal{O}_{\mathbf{Q}_G(w)}] - e^{\varpi_1 - w\varpi_1} \sum_{\substack{v \in W \\ \max(wW_{J_{\varpi_1} \setminus \{1\}}, \leq_v^*) = w}} (-1)^{\ell(v) - \ell(w)} [\mathcal{O}_{\mathbf{Q}_G(vt_{\text{wt}(w \Rightarrow v)})}]. \end{aligned} \quad (\text{A.2})$$

For example, if  $w = s_2$ , then it follows from (A.1) that

$$\{v \in W \mid \max(s_2W_{J_{\varpi_1}}, \leq_v^*) = s_2\} = \{s_2, s_1s_2, s_2s_1, w_\circ\},$$

and hence by (A.2)

$$[\mathcal{O}_{\mathbf{Q}_G(s_1)}] \cdot [\mathcal{O}_{\mathbf{Q}_G(s_2)}] = [\mathcal{O}_{\mathbf{Q}_G(s_1s_2)}] + [\mathcal{O}_{\mathbf{Q}_G(s_2s_1)}] - [\mathcal{O}_{\mathbf{Q}_G(w_\circ)}].$$

Similarly, if  $w = s_1$ , then

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G(s_1)}] \cdot [\mathcal{O}_{\mathbf{Q}_G(s_1)}] = \\ & [\mathcal{O}_{\mathbf{Q}_G(s_1)}] - e^{\varpi_1 - s_1\varpi_1} ([\mathcal{O}_{\mathbf{Q}_G(s_1)}] - [\mathcal{O}_{\mathbf{Q}_G(t_{\alpha_1^\vee})}] - [\mathcal{O}_{\mathbf{Q}_G(s_2s_1)}] + [\mathcal{O}_{\mathbf{Q}_G(s_2t_{\alpha_1^\vee})}]); \end{aligned}$$

if  $w = s_1s_2$ , then

$$[\mathcal{O}_{\mathbf{Q}_G(s_1)}] \cdot [\mathcal{O}_{\mathbf{Q}_G(s_1s_2)}] = [\mathcal{O}_{\mathbf{Q}_G(s_1s_2)}] - e^{\varpi_1 - s_1\varpi_1} ([\mathcal{O}_{\mathbf{Q}_G(s_1s_2)}] - [\mathcal{O}_{\mathbf{Q}_G(w_\circ)}]);$$

if  $w = s_2s_1$ , then

$$[\mathcal{O}_{\mathbf{Q}_G(s_1)}] \cdot [\mathcal{O}_{\mathbf{Q}_G(s_2s_1)}] = [\mathcal{O}_{\mathbf{Q}_G(s_2s_1)}] - e^{\varpi_1 - s_2s_1\varpi_1} ([\mathcal{O}_{\mathbf{Q}_G(s_2s_1)}] - [\mathcal{O}_{\mathbf{Q}_G(s_2t_{\alpha_1^\vee})}]);$$

if  $w = w_\circ$ , then

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G(s_1)}] \cdot [\mathcal{O}_{\mathbf{Q}_G(w_\circ)}] = \\ & [\mathcal{O}_{\mathbf{Q}_G(w_\circ)}] - e^{\varpi_1 - s_2s_1\varpi_1} ([\mathcal{O}_{\mathbf{Q}_G(w_\circ)}] - [\mathcal{O}_{\mathbf{Q}_G(t_{\theta^\vee})}] - [\mathcal{O}_{\mathbf{Q}_G(s_1s_2t_{\alpha_1^\vee})}] + [\mathcal{O}_{\mathbf{Q}_G(s_1t_{\theta^\vee})}]). \end{aligned}$$

## B Right-hand side of (3.21) in the case that $\lambda - \mu$ is not dominant.

Let  $\lambda, \mu \in P^+$ , and define  $J_\lambda, J_\mu \subset I$  as in (3.1). We write  $\lambda$  and  $\mu$  as  $\lambda = \sum_{i \in I} \lambda_i \varpi_i$  and  $\mu = \sum_{i \in I} \mu_i \varpi_i$ , respectively. We will prove that if  $\lambda - \mu$  is not dominant, then the right-hand side of (3.21) is identical to 0 for all  $x \in W_{\text{af}}$ ; we prove this assertion by induction on  $|\mu| = \sum_{i \in I} \mu_i$ . Assume that  $\lambda - \mu$  is not dominant. Also in this case, we denote by  $\mathbf{F}(x) = \mathbf{F}_{\lambda\mu}(x)$  the right-hand side of (3.21), as in §7.1.

Assume first that  $\mu = \varpi_r$  for some  $r \in I$  and  $x = t_\xi$  for some  $\xi \in Q^\vee$ ; since  $\lambda - \mu = \lambda - \varpi_r$  is not dominant, it follows that  $\lambda_r = 0$ . By the same argument as for (5.7), we deduce that

$$\mathbf{F}_{\lambda\varpi_r}(t_\xi) = \sum_{m \in \mathbb{Z}_{\geq 0}} q^{-m\lambda_r} \mathbf{e}^{-t_\xi \varpi_r} (\text{gch } V_{t_\xi}^-(\lambda) - \text{gch } V_{s_r t_\xi}^-(\lambda)).$$

Since  $\lambda_r = 0$  as seen above, we have  $V_{s_r t_\xi}^-(\lambda) = V_{t_\xi}^-(\lambda)$  (see Remark 3.6). Hence we obtain  $\mathbf{F}_{\lambda\varpi_r}(t_\xi) = 0$  in this case.

Also, we remark that the assertion of Proposition 7.2 and the arguments in its proof are still valid even if  $\lambda - \mu$  is not dominant.

Assume next that  $\mu = \varpi_r$  for some  $r \in I$  and  $x \in W_{\text{af}}$ . If we take a sequence  $i_1, \dots, i_n \in I_{\text{af}}$  and  $\xi \in Q^\vee$  as in §7.2, then we have

$$\mathbf{F}_{\lambda\varpi_r}(x) = D_{i_n} D_{i_{n-1}} \cdots D_{i_2} D_{i_1} \mathbf{F}_{\lambda\varpi_r}(t_\xi).$$

Since  $\mathbf{F}_{\lambda\varpi_r}(t_\xi) = 0$  as shown above, we obtain  $\mathbf{F}_{\lambda\varpi_r}(x) = 0$ . This proves the desired equality in the case that  $|\mu| = 1$ .

Assume now that  $|\mu| > 1$ , and take  $r \in I$  such that  $\mu_r \geq 1$ . We set  $\nu := \mu - \varpi_r \in P_+$ ; notice that  $|\nu| < |\mu|$ . By the same argument as for  $\mathbf{H}(x)$  in §8.1, we see that

$$\begin{aligned} \mathbf{F}_{\lambda\mu}(x) &= \sum_{\substack{z \in W_{\text{af}} \\ z \succeq x}} \sum_{\substack{\psi \in \mathbb{B}^{\frac{\infty}{2}}(\mu) \\ \iota(\psi) \preceq \Pi^{J_\mu}(z), \kappa(\psi, z) = x}} (-1)^{\ell^{\frac{\infty}{2}}(z) - \ell^{\frac{\infty}{2}}(x)} \mathbf{e}^{-\text{wt}(\psi)} \text{gch } V_z^-(\lambda) \\ &= \sum_{\substack{y \in W_{\text{af}} \\ y \succeq x}} \sum_{\substack{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\nu) \\ \iota(\pi) \preceq \Pi^{J_\nu}(y), \kappa(\pi, y) = x}} (-1)^{\ell^{\frac{\infty}{2}}(y) - \ell^{\frac{\infty}{2}}(x)} \mathbf{e}^{-\text{wt}(\pi)} \times \\ &\quad \underbrace{\sum_{\substack{z \in W_{\text{af}} \\ z \succeq y}} \sum_{\substack{\eta \in \mathbb{B}^{\frac{\infty}{2}}(\varpi_r) \\ \iota(\eta) \preceq \Pi^{J_{\varpi_r}}(z), \kappa(\eta, z) = y}} (-1)^{\ell^{\frac{\infty}{2}}(z) - \ell^{\frac{\infty}{2}}(y)} \mathbf{e}^{-\text{wt}(\eta)} \text{gch } V_z^-(\lambda)}_{=\mathbf{F}_{\lambda\varpi_r}(y)}. \end{aligned} \quad (\text{B.1})$$

If  $\lambda - \varpi_r$  is not dominant, then we have  $\mathbf{F}_{\lambda\varpi_r}(y) = 0$  by the argument above, and hence  $\mathbf{F}_{\lambda\mu}(x) = 0$ . Hence we assume that  $\lambda - \varpi_r$  is dominant. Then it follows from Theorem 3.9 that  $\mathbf{F}_{\lambda\varpi_r}(y) = (1 - q^{-\lambda_r})^{-1} \text{gch } V_y^-(\lambda - \varpi_r)$ . Substituting this equality into (B.1), we obtain

$$\mathbf{F}_{\lambda\mu}(x)$$

$$= (1 - q^{-\lambda_r})^{-1} \underbrace{\sum_{\substack{y \in W_{\text{af}} \\ y \succeq x}} \sum_{\substack{\pi \in \mathbb{B}_{\frac{\infty}{2}}(\nu) \\ \iota(\pi) \preceq \Pi^{J\nu}(y), \kappa(\pi, y) = x}} (-1)^{\ell_{\frac{\infty}{2}}(y) - \ell_{\frac{\infty}{2}}(x)} e^{-\text{wt}(\pi)} \text{gch } V_y^-(\lambda - \varpi_r)}_{= \mathbf{F}_{\lambda - \varpi_r, \nu}(x)},$$

and hence  $\mathbf{F}_{\lambda\mu}(x) = (1 - q^{-\lambda_r})^{-1} \mathbf{F}_{\lambda - \varpi_r, \nu}(x)$ . Since  $(\lambda - \varpi_r) - \nu = \lambda - \mu$  is not dominant, and since  $|\nu| < |\mu|$ , it follows by the induction hypothesis that  $\mathbf{F}_{\lambda - \varpi_r, \nu}(x) = 0$ , and hence  $\mathbf{F}_{\lambda\mu}(x) = 0$ . This completes the proof of the desired equality  $\mathbf{F}_{\lambda\mu}(x) = 0$  in the case that  $\lambda - \mu$  is not dominant.

## C Chevalley formula for dominant weights in terms of quantum LS paths.

We know from [KNS, (3.3)] that for  $\lambda, \mu \in P^+$  and  $x \in W$ ,

$$\text{gch } V_x^-(\lambda + \mu) = \sum_{\pi \in \mathbb{B}_{\frac{\infty}{2}}(\mu)} e^{\text{fin}(\text{wt}(\pi))} q^{\text{nul}(\text{wt}(\pi))} \text{gch } V_{\iota(\pi, x)}^-(\lambda), \quad (\text{C.1})$$

where  $\iota(\pi, x) \in W_{\text{af}}$  is defined as in (3.16) with  $y$  replaced by  $x$ . We will reformulate this formula in terms of quantum LS paths, as in Corollary 3.15 and §8.2.

We write  $\mu$  as  $\mu = \sum_{i \in I} \mu_i \varpi_i$ , and define  $\overline{\text{Par}(\mu)}$  to be the set of  $I$ -tuples of partitions  $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I}$  such that  $\chi^{(i)}$  is a partition of length less than or equal to  $\mu_i$  for each  $i \in I$ ; for  $\boldsymbol{\chi} \in \overline{\text{Par}(\mu)}$ , we define  $|\boldsymbol{\chi}| \in \mathbb{Z}_{\geq 0}$  and  $\iota(\boldsymbol{\chi}) \in Q^{\vee, +}$  as in §4.1. For  $\eta = (w_1, \dots, w_s; a_0, a_1, \dots, a_s) \in \text{QLS}(\mu)$  and  $x \in W$ , we define  $\iota(\eta, x) \in W$  by the following recursive formula (cf. (3.16) and Proposition 2.22):

$$\begin{cases} \tilde{w}_{s+1} := x, \\ \tilde{w}_u := \min(w_u W_{J_\mu}, \leq \tilde{w}_{u+1}) \quad \text{for } 1 \leq u \leq s, \\ \iota(\eta, x) := \tilde{w}_1. \end{cases} \quad (\text{C.2})$$

Also, we set

$$\xi(\eta, x) := \sum_{u=1}^s \text{wt}(\tilde{w}_{u+1} \Rightarrow \tilde{w}_u), \quad (\text{C.3})$$

and

$$\text{deg}_x(\eta) := - \sum_{u=1}^s a_u \langle \mu, \text{wt}(\tilde{w}_{u+1} \Rightarrow \tilde{w}_u) \rangle; \quad (\text{C.4})$$

notice that  $\text{deg}_x(\eta) = \text{deg}(\eta) - \langle \mu, \text{wt}(x \Rightarrow \kappa(\eta)) \rangle$  by (3.26) and Lemma 2.17.

**Corollary C.1.** *Keep the notation and setting above. For  $x \in W$ , we have*

$$\text{gch } V_x^-(\lambda + \mu) = \sum_{\eta \in \text{QLS}(\mu)} \sum_{\boldsymbol{\chi} \in \overline{\text{Par}(\mu)}} e^{\text{wt}(\eta)} q^{\text{deg}_x(\eta) - |\boldsymbol{\chi}|} \text{gch } V_{\iota(\eta, x) t_{\xi(\eta, x) + \iota(\boldsymbol{\chi})}}^-(\lambda). \quad (\text{C.5})$$

*Proof.* We see that

$$\mathbb{B}_{\geq x}^{\infty}(\mu) = \bigsqcup_{\eta \in \text{QLS}(\mu)} \text{cl}^{-1}(\eta) \cap \mathbb{B}_{\geq x}^{\infty}(\mu),$$

where  $\text{cl} : \mathbb{B}_{\geq x}^{\infty}(\mu) \rightarrow \text{QLS}(\mu)$  is the projection given in §3.4. We deduce from [NNS1, (5.4)], together with Lemma 2.17, that

$$\text{cl}^{-1}(\eta) \cap \mathbb{B}_{\geq x}^{\infty}(\mu) = \{ \pi_{\chi, \eta} \cdot t_{\text{wt}(x \Rightarrow \kappa(\eta)) + \gamma} \mid \chi \in \text{Par}(\mu), \gamma \in Q_{I \setminus J_{\mu}}^{\vee, +} \}; \quad (\text{C.6})$$

here,  $\pi_{\chi, \eta}$  is the (unique) element in the connected component  $\mathbb{B}_{\chi}^{\infty}(\mu)$  corresponding to  $\chi \in \text{Par}(\mu)$  such that  $\text{cl}(\pi_{\chi, \eta}) = \eta$  and  $\kappa(\pi_{\chi, \eta}) = \kappa(\eta) \in W$  (see §8.2), and for  $\pi = (x_1, \dots, x_s; a_0, a_1, \dots, a_s) \in \mathbb{B}_{\geq x}^{\infty}(\mu)$  and  $\xi \in Q^{\vee}$ , we set

$$\pi \cdot t_{\xi} := (\Pi^{J_{\mu}}(x_1 t_{\xi}), \dots, \Pi^{J_{\mu}}(x_s t_{\xi}); a_0, a_1, \dots, a_s) \in \mathbb{B}_{\geq x}^{\infty}(\mu)$$

(see [KNS, (7.7)]). In the same manner as in the calculation on page 43 of [NNS1], we see that if  $\chi = (\chi^{(i)})_{i \in I} \in \text{Par}(\mu)$  and  $\gamma = \sum_{i \in I \setminus J_{\mu}} \gamma_i \alpha_i^{\vee} \in Q_{I \setminus J_{\mu}}^{\vee}$ , then

$$\text{wt}(\pi_{\chi, \eta} \cdot t_{\text{wt}(x \Rightarrow \kappa(\eta)) + \gamma}) = \text{wt}(\eta) + (\deg_x(\eta) - |(\gamma_i + \chi^{(i)})_{i \in I}|) \delta,$$

where  $\gamma_i + \chi^{(i)}$  is defined to be the partition  $(\gamma_i + \chi_1^{(i)} \geq \gamma_i + \chi_2^{(i)} \geq \dots \geq \gamma_i + \chi_{\mu_i - 1}^{(i)} \geq \gamma_i)$  of length less than or equal to  $\mu_i$  for each  $i \in I \setminus J_{\mu}$ , and  $\emptyset$  for  $i \in J_{\mu}$ ; note that  $(\gamma_i + \chi^{(i)})_{i \in I} \in \overline{\text{Par}(\mu)}$ . Also, we deduce by Proposition 2.22, together with Lemma 2.17 and (3.25), that

$$\iota(\pi_{\chi, \eta} \cdot t_{\text{wt}(x \Rightarrow \kappa(\eta)) + \gamma}, x) = \iota(\eta, x) t_{\xi(\eta, x) + \iota(\chi) + \gamma}.$$

Therefore, the right-hand side of (C.1) is identical to:

$$\begin{aligned} & \sum_{\eta \in \text{QLS}(\mu)} \sum_{\chi \in \text{Par}(\mu)} \sum_{\gamma \in Q_{I \setminus J_{\mu}}^{\vee, +}} e^{\text{wt}(\eta)} q^{\deg_x(\eta) - |(\gamma_i + \chi^{(i)})_{i \in I}|} \text{gch } V_{\iota(\eta, x) t_{\xi(\eta, x) + \iota(\chi) + \gamma}}^{-}(\lambda) \\ &= \sum_{\eta \in \text{QLS}(\mu)} \sum_{\chi \in \overline{\text{Par}(\mu)}} e^{\text{wt}(\eta)} q^{\deg_x(\eta) - |\chi|} \text{gch } V_{\iota(\eta, x) t_{\xi(\eta, x) + \iota(\chi)}}^{-}(\lambda). \end{aligned}$$

This proves Corollary C.1.  $\square$

*Remark C.2.* In the sum on the right-hand side of (C.5), for each  $\eta \in \text{QLS}(\mu)$  and an arbitrary fixed  $\gamma \in Q^{\vee, +}$ , the number of the elements  $\chi \in \overline{\text{Par}(\mu)}$  contributing to the coefficient of  $\text{gch } V_{\iota(\eta, x) t_{\xi(\eta, x) + \gamma}}^{-}(\lambda)$  is finite.

By the same argument as for Theorem 1 in the Introduction, we obtain the following corollary, which is a rephrasing of [KNS, Theorem 5.10] in terms of quantum LS paths.

**Corollary C.3.** *Let  $\lambda \in P^+$  be a dominant integral weight, and let  $x = wt_{\xi} \in W_{\text{af}}^{\geq 0}$ . Then, in the  $H \times \mathbb{C}^*$ -equivariant  $K$ -group  $K_{H \times \mathbb{C}^*}(\mathbf{Q}_G)$ , we have*

$$\begin{aligned} & [\mathcal{O}_{\mathbf{Q}_G}(\lambda)] \cdot [\mathcal{O}_{\mathbf{Q}_G(x)}] \\ &= \sum_{\eta \in \text{QLS}(-w \circ \lambda)} \sum_{\chi \in \overline{\text{Par}(-w \circ \lambda)}} e^{\text{wt}(\eta)} q^{\deg_w(\eta) - \langle -w \circ \lambda, \xi \rangle - |\chi|} [\mathcal{O}_{\mathbf{Q}_G}(\iota(\eta, w) t_{\xi + \xi(\eta, w) + \iota(\chi)})]. \end{aligned} \quad (\text{C.7})$$

## References

- [AK] T. Akasaka and M. Kashiwara, Finite-dimensional representations of quantum affine algebras, *Publ. Res. Inst. Math. Sci.* **33** (1997), 839–867.
- [ACT] D. Anderson, L. Chen, and H.-H. Tseng, On the finiteness of quantum  $K$ -theory of a homogeneous space, arXiv:1804.04579.
- [BB] A. Björner and F. Brenti, “Combinatorics of Coxeter Groups”, Graduate Texts in Mathematics Vol. 231, Springer, New York, 2005.
- [BCMP] A.S. Buch, P.-E. Chaput, L. C. Mihalcea, and N. Perrin, A Chevalley formula for the equivariant quantum  $K$ -theory of cominuscle varieties, *Algebr. Geom.* **5** (2018), 568–595.
- [GR] S. Griffeth and A. Ram, Affine Hecke algebras and the Schubert calculus, *European J. Combin.* **25** (2004), 1263–1283.
- [HK] J. Hong and S.-J. Kang, “Introduction to Quantum Groups and Crystal Bases”, Graduate Studies in Mathematics Vol. 42, Amer. Math. Soc., Providence, RI, 2002.
- [IMT] H. Iritani, T. Milanov, and V. Tonita, Reconstruction and convergence in quantum  $K$ -theory via difference equations, *Int. Math. Res. Not.* **2015** (2015), 2887–2937.
- [INS] M. Ishii, S. Naito, and D. Sagaki, Semi-infinite Lakshmibai-Seshadri path model for level-zero extremal weight modules over quantum affine algebras, *Adv. Math.* **290** (2016), 967–1009.
- [Kac] V. G. Kac, “Infinite Dimensional Lie Algebras”, 3rd Edition, Cambridge University Press, Cambridge, UK, 1990.
- [Kas1] M. Kashiwara, Crystal bases of modified quantized enveloping algebra, *Duke Math. J.* **73** (1994), 383–413.
- [Kas2] M. Kashiwara, “Bases Cristallines des Groupes Quantiques” (Notes by Charles Cochet), Cours Spécialisés Vol. 9, Société Mathématique de France, Paris, 2002.
- [Kas3] M. Kashiwara, On level-zero representations of quantized affine algebras, *Duke Math. J.* **112** (2002), 117–175.
- [Kas4] M. Kashiwara, Level zero fundamental representations over quantized affine algebras and Demazure modules, *Publ. Res. Inst. Math. Sci.* **41** (2005), 223–250.
- [Kat1] S. Kato, Loop structure on equivariant  $K$ -theory of semi-infinite flag manifolds, arXiv:1805.01718.
- [Kat2] S. Kato, Frobenius splitting of Schubert varieties of semi-infinite flag manifolds, arXiv:1810.07106.
- [Kat3] S. Kato, On quantum  $K$ -groups of partial flag manifolds, arXiv:1906.09343.
- [KNS] S. Kato, S. Naito, and D. Sagaki, Pieri-Chevalley type formula for equivalent  $K$ -theory of semi-infinite flag manifolds, arXiv:1702.02408, to appear in *Duke Math. J.*
- [KoNS] T. Kouno, S. Naito, and D. Sagaki, Chevalley formula for the equivariant quantum  $K$ -group of partial flag manifolds of minuscule type, in preparation.
- [L<sup>2</sup>MS] T. Lam, C. Li, L.C. Mihalcea, and M. Shimozono, A conjectural Peterson isomorphism in  $K$ -theory, *J. Algebra* **513** (2018), 326–343.

- [LaS] T. Lam and M. Shimozono, Quantum cohomology of  $G/P$  and homology of affine Grassmannian, *Acta Math.* **204** (2010), 49–90.
- [L1] P. Littelmann, A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras, *Invent. Math.* **116** (1994), 329–346.
- [L2] P. Littelmann, Paths and root operators in representation theory, *Ann. of Math.* (2) **142** (1995), 499–525.
- [LNS] C. Lenart, S. Naito, and D. Sagaki, A Chevalley formula for semi-infinite flag manifolds and quantum  $K$ -theory, arXiv:1911.12773.
- [LNS<sup>3</sup>1] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, A uniform model for Kirillov-Reshetikhin crystals I: Lifting the parabolic quantum Bruhat graph, *Int. Math. Res. Not.* **2015** (2015), 1848–1901.
- [LNS<sup>3</sup>2] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, A uniform model for Kirillov-Reshetikhin crystals II: Alcove model, path model, and  $P = X$ , *Int. Math. Res. Not.* **2017** (2017), 4259–4319.
- [LNS<sup>3</sup>3] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, Explicit description of the degree function in terms of quantum Lakshmibai-Seshadri paths, *Toyama Math. J.* **37** (2015), 107–130.
- [LeP] C. Lenart and A. Postnikov, Affine Weyl groups in  $K$ -theory and representation theory, *Int. Math. Res. Not.* **2007** (2007), no. 12, Art. ID: rnm038, 65 pp.
- [LeS] C. Lenart and M. Shimozono, Equivariant  $K$ -Chevalley rules for Kac-Moody flag manifolds, *Amer. J. Math.* **136** (2014), 1175–1213.
- [NNS1] S. Naito, F. Nomoto, and D. Sagaki, Specialization of nonsymmetric Macdonald polynomials at  $t = \infty$  and Demazure submodules of level-zero extremal weight modules, *Trans. Amer. Math. Soc.* **370** (2018), 2739–2783.
- [NNS2] S. Naito, F. Nomoto, and D. Sagaki, Representation-theoretic interpretation of Cherednik-Orr’s recursion formula for the specialization of nonsymmetric Macdonald polynomials at  $t = \infty$ , *Transform. Groups* **24** (2019), 155–191.
- [NNS3] S. Naito, F. Nomoto, and D. Sagaki, Tensor product decomposition theorem for quantum Lakshmibai-Seshadri paths and standard monomial theory for semi-infinite Lakshmibai-Seshadri paths, *J. Combin. Theory Ser. A* **169** (2020), 105122, 36pp.
- [NS1] S. Naito and D. Sagaki, Path model for a level-zero extremal weight module over a quantum affine algebra. II, *Adv. Math.* **200** (2006), 102–124.
- [NS2] S. Naito and D. Sagaki, Lakshmibai-Seshadri paths of a level-zero weight shape and one-dimensional sums associated to level-zero fundamental representations, *Compos. Math.* **144** (2008), 1525–1556.
- [NS3] S. Naito and D. Sagaki, Demazure submodules of level-zero extremal weight modules and specializations of Macdonald polynomials, *Math. Z.* **283** (2016), 937–978.
- [NS4] S. Naito and D. Sagaki, Level-zero van der Kallen modules and specialization of nonsymmetric Macdonald polynomials at  $t = \infty$ , arXiv:1802.06339.
- [O] D. Orr, Equivariant  $K$ -theory of the semi-infinite flag manifold as a nil-DAHA module, arXiv:2001.03490.
- [P] D. Peterson, Quantum Cohomology of  $G/P$ , Lecture notes, Massachusetts Institute of Technology, Spring 1997.