EXISTENCE AND CONCENTRATION OF NONTRIVIAL SOLUTIONS FOR A FRACTIONAL MAGNETIC SCHRÖDINGER-POISSON TYPE EQUATION

VINCENZO AMBROSIO

Abstract. We consider the following fractional Schrödinger-Poisson type equation with magnetic fields $% \mathcal{A}_{\mathrm{S}}$

$$\varepsilon^{2s}(-\Delta)^{s}_{A/\varepsilon}u + V(x)u + \varepsilon^{-2t}(|x|^{2t-3} * |u|^{2})u = f(|u|^{2})u \quad \text{in } \mathbb{R}^{3},$$

where $\varepsilon > 0$ is a parameter, $s \in (\frac{3}{4}, 1)$, $t \in (0, 1)$, $(-\Delta)_A^s$ is the fractional magnetic Laplacian, $A : \mathbb{R}^3 \to \mathbb{R}^3$ is a smooth magnetic potential, $V : \mathbb{R}^3 \to \mathbb{R}$ is a positive continuous electric potential and $f : \mathbb{R}^3 \to \mathbb{R}$ is a continuous function with subcritical growth. By using suitable variational methods, we show the existence of families of nontrivial solutions concentrating around local minima of the potential V(x) as $\varepsilon \to 0$.

1. INTRODUCTION

In this paper we are interested in the existence of nontrivial solutions for the following fractional nonlinear Schrödinger-Poisson type equation

$$\varepsilon^{2s}(-\Delta)^{s}_{A/\varepsilon}u + V(x)u + \varepsilon^{-2t}(|x|^{2t-3} * |u|^{2})u = f(|u|^{2}) \quad \text{in } \mathbb{R}^{3},$$
(1.1)

where $\varepsilon > 0$ is a parameter, $s \in (\frac{3}{4}, 1)$, $t \in (0, 1)$, $A : \mathbb{R}^3 \to \mathbb{R}^3 \in C^{0,\alpha}$, with $\alpha \in (0, 1]$, is a magnetic potential, and $(-\Delta)_A^s$ is the so called fractional magnetic Laplacian which can be defined by setting

$$(-\Delta)_{A}^{s}u(x) := c_{3,s} \lim_{r \to 0} \int_{B_{r}^{c}(x)} \frac{u(x) - e^{i(x-y) \cdot A(\frac{x+y}{2})}u(y)}{|x-y|^{3+2s}} dy, \quad c_{3,s} := \frac{4^{s}\Gamma\left(\frac{3+2s}{2}\right)}{\pi^{3/2}|\Gamma(-s)|}, \quad (1.2)$$

for any $u \in C_c^{\infty}(\mathbb{R}^3, \mathbb{C})$; see [18,31] for more details. As showed in [47] (see also [43]), when $s \to 1$, the previous operator reduces to the magnetic Laplacian $-\Delta_A := \left(\frac{1}{i}\nabla - A\right)^2$ (see [34,36]) given by

$$-\Delta_A u = -\Delta u - \frac{2}{i}A(x) \cdot \nabla u + |A(x)|^2 u - \frac{1}{i}u\operatorname{div}(A(x)).$$

which appears in the study of the following Schrödinger equation with magnetic fields

$$-\Delta_A u + V(x)u = f(x, |u|^2)u \quad \text{in } \mathbb{R}^N.$$
(1.3)

Equation (1.3) has been widely investigated by several authors in the last thirty years; see for instance [1, 7, 12, 14, 22, 33].

Along the paper, we assume that $V : \mathbb{R}^3 \to \mathbb{R}$ is a continuous potential satisfying the following assumptions due to del Pino and Felmer [19]:

- $(V_1) \inf_{x \in \mathbb{R}^3} V(x) = V_0 > 0;$
- (V_2) there exists a bounded domain $\Lambda \subset \mathbb{R}^3$ such that

$$V_0 < \min_{\partial \Lambda} V$$
 and $M = \{x \in \Lambda : V(x) = V_0\} \neq \emptyset$, (1.4)

and $f : \mathbb{R} \to \mathbb{R}$ is a continuous function verifying the following conditions:

 $(f_1) \ f(t) = 0 \text{ for } t \le 0 \text{ and } \lim_{t \to 0} \frac{f(t)}{t} = 0;$

²⁰¹⁰ Mathematics Subject Classification. 35A15, 35R11, 35S05.

Key words and phrases. Fractional magnetic operators, variational methods, concentration phenomenon.

 (f_2) there exist $q \in (4, 2_s^*)$ such that

$$\lim_{t \to \infty} \frac{f(t)}{t^{\frac{q-2}{2}}} = 0;$$

(f₃) there exists $\theta \in (4, 2_s^*)$ such that $0 < \frac{\theta}{2}F(t) \le tf(t)$ for any t > 0, where $F(t) = \int_0^t f(\tau)d\tau$; (f₄) $t \mapsto \frac{f(t)}{t}$ is increasing for t > 0. Let us state our main theorem:

Theorem 1.1. Assume that (V_1) - (V_2) and (f_1) - (f_4) hold. Then there exists $\varepsilon_0 > 0$ such that, for any $\varepsilon \in (0, \varepsilon_0)$, problem (1.1) has a nontrivial solution. Moreover, if u_{ε} denotes one of these solutions and x_{ε} the global maximum point of $|u_{\varepsilon}|$, then we have

$$\lim_{\varepsilon \to 0} V(x_{\varepsilon}) = V_0$$

and

$$|u_{\varepsilon}(x)| \leq \frac{C \,\varepsilon^{3+2s}}{C \,\varepsilon^{3+2s} + |x - x_{\varepsilon}|^{3+2s}} \quad \forall x \in \mathbb{R}^3.$$

The above result is motivated by some works that appeared in the last years concerning fractional Schrödinger equations with magnetic fields of the type

$$\varepsilon^{2s}(-\Delta)^s_A u + V(x)u = f(x, |u|^2)u \quad \text{in } \mathbb{R}^N.$$
(1.5)

For instance, in the unperturbed case (that is $\varepsilon = 1$), d'Avenia and Squassina [18] studied via a constrained minimization argument the existence of solutions (1.5), V is constant and f is a subcritical or critical nonlinearity. Fiscella et al. [26] obtained a multiplicity result for a fractional magnetic problem with homogeneous boundary conditions. When $\varepsilon > 0$ is small, Zhang et al. [52] considered a fractional magnetic Schrödinger equation involving critical frequency and critical growth. Recently, in [6], the author and d'Avenia dealt with the existence and the multiplicity of solutions to (1.5) for small $\varepsilon > 0$, when the potential V satisfies the global condition due to Rabinowitz [44] and f has a subcritical growth.

In absence of a magnetic field (that is A = 0), the fractional magnetic Laplacian $(-\Delta)_A^s$ coincides with the fractional Laplacian $(-\Delta)^s$ and the equation (1.5) becomes the well-known fractional Schrödinger equation (see [35])

$$\varepsilon^{2s}(-\Delta)^s u + V(x)u = f(x,u) \text{ in } \mathbb{R}^N, \tag{1.6}$$

for which the existence and concentration phenomena of positive solutions have been considered by many mathematicians. For example, Dávila et al. [17] used a Lyapunov-Schmidt variational reduction to prove that (1.6) has a multi-peak solution when $V \in L^{\infty}(\mathbb{R}^N) \cap C^{1,\alpha}(\mathbb{R}^N)$ is a positive potential and f is a subcritical nonlinearity; see also [16] in which a concentration result has been established for a nonlocal problem with Dirichlet datum. Fall et al. [24] showed that the concentration points of the solutions of (1.6) must be the critical points for V, as ε tends to zero. Alves and Miyagaki [2] (see also [4,5]) used the penalization method in [19] to study the existence and concentration of positive solutions of (1.6) requiring that f satisfies (f_1) - (f_4) and V verifies (V_1) - (V_2) .

On the other hand, in these last years, several authors investigated fractional Schrödinger-Poisson systems of the type

$$\begin{cases} \varepsilon^{2s}(-\Delta)^s u + V(x)\phi u = g(x,u) & \text{in } \mathbb{R}^3\\ \varepsilon^{2t}(-\Delta)^t \phi = u^2 & \text{in } \mathbb{R}^3, \end{cases}$$
(1.7)

which can be seen as the nonlocal counterpart of the well-known Schrödinger-Poisson systems appearing in quantum mechanics models [10] and in semiconductor theory [38]. Such systems have been introduced in [9] to describe systems of identical charged particles interacting each other in the case that effects of magnetic field could be ignored and its solution represents, in particular, a

standing wave for such a system. We refer to [8, 15, 28, 29, 45, 49, 53] for some interesting existence and multiplicity results for classical perturbed and unperturbed Schrödinger-Poisson systems.

Concerning (1.7), Giammetta [27] considered the local and global well-posedness of a one dimensional fractional Schrödinger-Poisson system in which $\varepsilon = 1$ and the fractional diffusion appears only in the Poisson equation. Zhang et al. [51] dealt with the existence of positive solutions to (1.7) with $\varepsilon = 1$, $V(x) = \mu > 0$ and g is a general nonlinearity having subcritical or critical growth. Murcia and Siciliano [41] proved that, for suitably small ε , the number of positive solutions to a doubly singularly perturbed fractional Schrödinger-Poisson system is estimated below by the Ljusternick-Schnirelmann category of the set of minima of the potential. Liu and Zhang [37] studied multiplicity and concentration of solutions to (1.7) involving the critical exponent and under a global condition on the potential V. Teng [48], inspired by [29], used the penalization method due to Byeon and Wang [11] to analyze the existence and concentration of positive solutions to (1.7) under the conditions (V_1) - (V_2) and g is a C^1 subcritical nonlinearity.

Particularly motivated by [2, 6, 48], in this paper we investigate the existence and concentration behavior of nontrivial solutions to (1.1) with $A \neq 0$ and under the assumptions (V_1) - (V_2) and (f_1) - (f_4) . We note that when s = t = 1 in (1.1), the multiplicity and concentration for a Schrödinger-Poisson type equation with magnetic field and under a local condition on V, has been established in [54] by using some ideas developed in [1]. Anyway, their arguments work for C^1 -Nehari manifolds and we can not apply them in our situation because we are assuming the only continuity of f.

Since we don't have any information on the behavior of V at infinity, we adapt the penalization argument developed by del Pino and Felmer in [19], which consists in making an appropriate modification on f, solving a modified problem and then check that, for ε small enough, the solutions of the modified problem are indeed solutions of the original one. We point out that the penalization argument developed here is different from the one used in [48], in which the author does not assume the suplinear-4 growth on f but has to require $f \in C^1$ to apply the techniques developed in [11,29]. The existence of nontrivial solutions for the modified problem is obtained by using the Mountain Pass Theorem [3] to the functional J_{ε} associated to the modified problem. We note that the main issue in the study of J_{ε} concerns with the verification the Palais-Smale compactness condition. Indeed, the presence of the fractional magnetic Laplacian and the convolution term $(|x|^{2t-3} * |u|^2)$, make our study more complicated and intriguing, and some suitable arguments will be needed to achieve our purpose; see Lemma 3.2. The next step is to show that if u_{ε} is a solution of the modified problem, then u_{ε} is also a solution of the original one (1.1). In the case A = 0 (see [2, 48]), this is proved taking into account some fundamental estimates established in [25] concerning the Bessel operator. In the case $A \neq 0$, we don't have similar informations for the following fractional equation

$$(-\Delta)_{A}^{s}u + V_{0}u = h(|u|^{2})u \text{ in } \mathbb{R}^{3}.$$
(1.8)

For the above reason, we use a new approximation argument which allows us to deduce that if u_{ε} is a solution to the modified problem, then $|u_{\varepsilon}|$ is a subsolution for an autonomous fractional Schrödinger equation without magnetic field, and then we apply a comparison argument to deduce informations on the behavior at infinity of $|u_{\varepsilon}|$; see Lemma 4.1. We point out that, in the case s = 1, a similar reasoning works (see [13, 33]) in view of the following distributional Kato's inequality [32]

$$-\Delta |u| \le \Re(sign(u)(-\Delta_A u)).$$

Recently, in [30], under the restriction $s \in (0, 1/2]$, a fractional distributional Kato's inequality has been established for some fractional magnetic operators which also include $(-\Delta)_A^{1/2}$. We suspect that a fractional Kato's inequality is available in our setting for any fractional power s (indeed it is easily seen that a pointwise Kato's inequality holds for smooth functions), but we are not able to prove it. Again, we can not repeat the iteration done in [1] to obtain L^{∞} -estimates on the modulus of solutions, due to the nonlocal character of $(-\Delta)_A^s$. Anyway, in the present paper, we

V. AMBROSIO

introduce some new arguments which we believe to be useful to be applied in other situations to obtain L^{∞} -estimates for problems like (1.5). Now we give a sketch of our idea. Firstly, we show that the (translated) sequence $|u_n|$ of solutions of the modified problem is bounded in $L^{\infty}(\mathbb{R}^3, \mathbb{R})$ uniformly in $n \in \mathbb{N}$, by using an appropriate Moser iteration scheme [40]. After that, we prove that $|u_n|$ verifies

$$(-\Delta)^s |u_n| + V_0 |u_n| \le g(\varepsilon x, |u_n|^2) |u_n| \text{ in } \mathbb{R}^3,$$

by using $\frac{u_n}{u_{\delta,n}}\varphi$ as test function in the modified problem, where $u_{\delta,n} = \sqrt{|u_n|^2 + \delta^2}$ and φ is a real smooth nonnegative function with compact support in \mathbb{R}^3 , and then we take the limit as $\delta \to 0$. In some sense, we are going to prove a fractional Kato's inequality for the modified problem. At this point, by comparison, we can show that $|u_n(x)| \to 0$ as $|x| \to \infty$ uniformly with respect to $n \in \mathbb{N}$, taking into account the power type decay of solutions of autonomous fractional Schrödinger equations; see [25].

The paper is organized as follows. In Section 2 we give some results on fractional magnetic Sobolev spaces and we recall some useful lemmas. In Section 3, we introduce the modified problem and we show that the corresponding functional satisfies the assumptions of the Mountain Pass Theorem. In the last section we give the proof of Theorem 1.1.

2. Preliminaries and functional setting

Let us consider the fractional Sobolev space

$$H^s(\mathbb{R}^3,\mathbb{R}) = \{ u \in L^2(\mathbb{R}^3,\mathbb{R}) : [u] < \infty \}$$

where

$$[u]^{2} = \iint_{\mathbb{R}^{6}} \frac{|u(x) - u(y)|^{2}}{|x - y|^{3 + 2s}} dx dy.$$

It is well-known (see [20, 39]) that the embedding $H^s(\mathbb{R}^3, \mathbb{R}) \subset L^q(\mathbb{R}^3, \mathbb{R})$ is continuous for all $q \in [2, 2^*_s)$ and locally compact for all $q \in [1, 2^*_s)$.

Let $L^2(\mathbb{R}^3, \mathbb{C})$ be the space of complex-valued functions such that $\int_{\mathbb{R}^3} |u|^2 dx < \infty$ endowed with the inner product $\langle u, v \rangle_{L^2} = \Re \int_{\mathbb{R}^3} u \bar{v} dx$, where the bar denotes complex conjugation.

Let us denote by

$$[u]_A^2 := \frac{c_{3,s}}{2} \iint_{\mathbb{R}^6} \frac{|u(x) - e^{i(x-y) \cdot A(\frac{x+y}{2})} u(y)|^2}{|x-y|^{3+2s}} \, dx \, dy,$$

and we define

$$D^s_A(\mathbb{R}^3, \mathbb{C}) := \left\{ u \in L^{2^*_s}(\mathbb{R}^3, \mathbb{C}) : [u]^2_A < \infty \right\}.$$

In order to study our problem, we introduce the Hilbert space

$$H^s_{\varepsilon} := \left\{ u \in D^s_{A_{\varepsilon}}(\mathbb{R}^3, \mathbb{C}) : \int_{\mathbb{R}^3} V(\varepsilon \, x) |u|^2 \, dx < \infty \right\}$$

endowed with the scalar product

$$\begin{split} \langle u, v \rangle_{\varepsilon} &= \Re \int_{\mathbb{R}^3} V(\varepsilon \, x) u \bar{v} dx \\ &+ \frac{c_{3,s}}{2} \Re \iint_{\mathbb{R}^6} \frac{(u(x) - e^{\imath (x-y) \cdot A_{\varepsilon}(\frac{x+y}{2})} u(y)) \overline{(v(x) - e^{\imath (x-y) \cdot A_{\varepsilon}(\frac{x+y}{2})} v(y))}}{|x-y|^{3+2s}} dx dy \end{split}$$

and we set

$$||u||_{\varepsilon} := \sqrt{\langle u, u \rangle_{\varepsilon}}.$$

The space H^s_{ε} satisfies the following fundamental properties; see [6, 18] for more details. Lemma 2.1. [6, 18] The space H^s_{ε} is complete and $C^{\infty}_c(\mathbb{R}^3, \mathbb{C})$ is dense in H^s_{ε} . **Theorem 2.1.** [18] The space H^s_{ε} is continuously embedded in $L^r(\mathbb{R}^3, \mathbb{C})$ for $r \in [2, 2^*_s]$, and compactly embedded in $L^r_{\text{loc}}(\mathbb{R}^3, \mathbb{C})$ for $r \in [1, 2^*_s)$.

Lemma 2.2. [18] If $u \in H^s_A(\mathbb{R}^3, \mathbb{C})$ then $|u| \in H^s(\mathbb{R}^3, \mathbb{R})$ and we have $[|u|] \leq [u]_A.$

Lemma 2.3. [6] If $u \in H^s(\mathbb{R}^3, \mathbb{R})$ and u has compact support, then $w = e^{iA(0) \cdot x} u \in H^s_{\varepsilon}$.

We also recall the following vanishing lemma [25] which will be useful for our study:

Lemma 2.4. [25] Let $q \in [2, 2_s^*)$. If (u_n) is a bounded sequence in $H^s(\mathbb{R}^3, \mathbb{R})$ and if

$$\lim_{n \to \infty} \sup_{y \in \mathbb{R}^3} \int_{B_R(y)} |u_n|^q dx = 0$$

for some R > 0, then $u_n \to 0$ in $L^r(\mathbb{R}^3, \mathbb{R})$ for all $r \in (2, 2^*_s)$.

Now, let $s, t \in (0, 1)$ such that $4s + 2t \ge 3$. Since $H^s(\mathbb{R}^3, \mathbb{R}) \subset L^q(\mathbb{R}^3, \mathbb{R})$ for all $q \in [2, 2^*_s)$, we can deduce that

$$H^{s}(\mathbb{R}^{3},\mathbb{R}) \subset L^{\frac{12}{3+2t}}(\mathbb{R}^{3},\mathbb{R}).$$

$$(2.1)$$

For any $u \in H^s_{\varepsilon}$, we know that $|u| \in H^s(\mathbb{R}^3, \mathbb{R})$ in view of Lemma 2.2, and then we consider the linear functional $\mathcal{L}_{|u|} : D^{t,2}(\mathbb{R}^3, \mathbb{R}) \to \mathbb{R}$ given by

$$\mathcal{L}_{|u|}(v) = \int_{\mathbb{R}^3} |u|^2 v \, dx$$

By using Hölder inequality and (2.1) we can see that

$$|\mathcal{L}_{|u|}(v)| \le \left(\int_{\mathbb{R}^3} |u|^{\frac{12}{3+2t}} dx\right)^{\frac{3+2t}{6}} \left(\int_{\mathbb{R}^3} |v|^{2^*_t} dx\right)^{\frac{1}{2^*_t}} \le C ||u||^2_{D^{s,2}} ||v||_{D^{t,2}},\tag{2.2}$$

where

$$\|v\|_{D^{t,2}}^2 = \iint_{\mathbb{R}^6} \frac{|v(x) - v(y)|^2}{|x - y|^{3+2t}} dx dy,$$

and this shows that $\mathcal{L}_{|u|}$ is well defined and continuous. By using the Lax-Milgram Theorem, there exists a unique $\phi_{|u|}^t \in D^{t,2}(\mathbb{R}^3,\mathbb{R})$ such that

$$(-\Delta)^t \phi^t_{|u|} = |u|^2 \text{ in } \mathbb{R}^3.$$
 (2.3)

Then we have the following t-Riesz formula

$$\phi_{|u|}^t(x) = c_t \int_{\mathbb{R}^3} \frac{|u(y)|^2}{|x-y|^{3-2t}} \, dy \quad (x \in \mathbb{R}^3), \quad c_t = \pi^{-\frac{3}{2}} 2^{-2t} \frac{\Gamma(3-2t)}{\Gamma(t)}. \tag{2.4}$$

In the sequel, we will omit the constant c_t in order to lighten the notation. Finally, we prove some properties on the convolution term.

Lemma 2.5. Let us assume that $4s + 2t \ge 3$ and $u \in H^s_{\varepsilon}$. Then we have:

- (1) $\phi_{|u|}^t : H^s(\mathbb{R}^3, \mathbb{R}) \to D^{t,2}(\mathbb{R}^3, \mathbb{R})$ is continuous and maps bounded sets into bounded sets,
- (2) if $u_n \rightharpoonup u$ in H^s_{ε} then $\phi^t_{|u_n|} \rightharpoonup \phi^t_{|u|}$ in $D^{t,2}(\mathbb{R}^3, \mathbb{R})$,
- (3) $\phi_{|ru|}^t = r^2 \phi_{|u|}^t$ for all $r \in \mathbb{R}$ and $\phi_{|u(\cdot+y)|}^t(x) = \phi_{|u|}^t(x+y)$,
- (4) $\phi_{|u|}^t \ge 0$ for all $u \in H^s_{\varepsilon}$, and we have

$$\|\phi_{|u|}^t\|_{D^{t,2}} \le C\|u\|_{L^{\frac{12}{3+2t}}(\mathbb{R}^3)}^2 \le C\|u\|_{\varepsilon}^2 \text{ and } \int_{\mathbb{R}^3} \phi_{|u|}^t|u|^2 dx \le C\|u\|_{L^{\frac{12}{3+2t}}(\mathbb{R}^3)}^4 \le C\|u\|_{\varepsilon}^4$$

Proof. (1) Since $\phi_{|u|}^t \in D^{t,2}(\mathbb{R}^3,\mathbb{R})$ satisfies (2.3), that is

$$\int_{\mathbb{R}^3} (-\Delta)^{\frac{t}{2}} \phi_{|u|}^t (-\Delta)^{\frac{t}{2}} v \, dx = \int_{\mathbb{R}^3} |u|^2 v \, dx$$

for all $v \in D^{t,2}(\mathbb{R}^3, \mathbb{R})$, we can see that $\mathcal{L}_{|u|}$ is such that $\|\mathcal{L}_{|u|}\|_{\mathcal{L}(D^{t,2},\mathbb{R})} = \|\phi_{|u|}^t\|_{D^{t,2}}$ for all $u \in H^s_{\varepsilon}$. Hence, in order to prove the continuity of $\phi_{|u|}^t$, it is enough to show that the map $u \mapsto \mathcal{L}_{|u|}$ is continuous. Let $u_n \to u$ in H^s_{ε} . By using Lemma 2.2 and Theorem 2.1 we deduce that $|u_n| \to |u|$ in $L^{\frac{12}{3+2t}}(\mathbb{R}^3)$. Hence, for all $v \in D^{t,2}(\mathbb{R}^3, \mathbb{R})$ we have

$$\begin{aligned} |\mathcal{L}_{|u_n|}(v) - \mathcal{L}_{|u|}(v)| &= \left| \int_{\mathbb{R}^3} (|u_n|^2 - |u|^2) v \, dx \right| \\ &\leq \left(\int_{\mathbb{R}^3} ||u_n|^2 - |u|^2|^{\frac{6}{3+2t}} \, dx \right)^{\frac{3+2t}{6}} \|v\|_{L^{\frac{6}{3-2t}}(\mathbb{R}^3)} \\ &\leq C \left[\left(\int_{\mathbb{R}^3} ||u_n| - |u||^{\frac{12}{3+2t}} \, dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^3} ||u_n| + |u||^{\frac{12}{3+2t}} \, dx \right)^{\frac{1}{2}} \right]^{\frac{3+2t}{6}} \|v\|_{D^{t,2}} \\ &\leq C \||u_n| - |u|\|_{L^{\frac{12}{3+2t}}(\mathbb{R}^3)} \|v\|_{D^{t,2}} \end{aligned}$$

which implies that $\|\phi_{|u_n|}^t - \phi_{|u|}^t\|_{D^{t,2}} = \|\mathcal{L}_{|u_n|} - \mathcal{L}_{|u|}\|_{\mathcal{L}(D^{t,2},\mathbb{R})} \to 0$ as $n \to \infty$. (2) If $u_n \to u$ in H^s_{ε} , then Lemma 2.2 and Theorem 2.1 yield $|u_n| \to |u|$ in $L^q_{loc}(\mathbb{R}^3, \mathbb{R})$ for all $q \in [1, 2^*_s)$. Hence, for all $v \in C^\infty_c(\mathbb{R}^3, \mathbb{R})$ we get

$$\begin{aligned} \langle \phi_{|u_n|}^t - \phi_{|u|}^t, v \rangle &= \int_{\mathbb{R}^3} (|u_n|^2 - |u|^2) v \, dx \\ &\leq \left(\int_{supp(v)} ||u_n| - |u||^2 \, dx \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^3} ||u_n| + |u||^2 \, dx \right)^{\frac{1}{2}} \|v\|_{L^{\infty}(\mathbb{R}^3)} \\ &\leq C \||u_n| - |u|\|_{L^2(supp(v))} \|v\|_{L^{\infty}(\mathbb{R}^3)} \to 0. \end{aligned}$$

(3) and (4) are easily obtained by applying Hardy-Littlewood-Sobolev inequality (see Theorem 4.3 in [36]), Hölder inequality and Sobolev embedding. $\hfill \Box$

3. The Modified problem

By using the change of variable $x \mapsto \varepsilon x$, we can see that the study of (1.1) is equivalent to consider the following problem

$$(-\Delta)_{A_{\varepsilon}}^{s} u + V_{\varepsilon}(x)u + (|x|^{2t-3} * |u|^{2})u = f(|u|^{2})u \text{ in } \mathbb{R}^{3},$$
(3.1)

where $A_{\varepsilon}(x) = A(\varepsilon x)$ and $V_{\varepsilon}(x) = V(\varepsilon x)$.

As in [2,19], we fix $k > \frac{\theta}{\theta-2}$ and a > 0 such that $\frac{f(a)}{a} = \frac{V_0}{k}$, and we introduce the function

$$\tilde{f}(t) := \begin{cases} f(t) & \text{if } t \leq a \\ \frac{V_0}{k} & \text{if } t > a \end{cases}$$

Then we define the penalized nonlinearity $g: \mathbb{R}^3 \times \mathbb{R} \to \mathbb{R}$ by setting

$$g(x,t) = \chi_{\Lambda}(x)f(t) + (1 - \chi_{\Lambda}(x))\tilde{f}(t),$$

where χ_{Λ} is the characteristic function on Λ , and we set $G(x,t) = \int_0^t g(x,\tau) d\tau$.

From the assumptions (f_1) - (f_4) it is standard to check that g verifies the following properties:

(g₁)
$$\lim_{t \to 0} \frac{g(x,t)}{t} = 0$$
 uniformly in $x \in \mathbb{R}^3$;

(g₂) $\lim_{t\to\infty} \frac{g(x,t)}{t^{\frac{q-2}{2}}} = 0$ uniformly in $x \in \mathbb{R}^3$;

 $\begin{array}{l} (g_3) \ (i) \ 0 < \frac{\bar{g}}{2}G(x,t) \le g(x,t)t \ \text{for any } x \in \Lambda \ \text{and } t > 0, \\ (ii) \ 0 \le G(x,t) \le g(x,t)t \le \frac{V(x)}{k}t \ \text{and } 0 \le g(x,t) \le \frac{V(x)}{k} \ \text{for any } x \in \Lambda^c \ \text{and } t > 0; \\ (g_4) \ t \mapsto \frac{g(x,t)}{t} \ \text{is increasing for all } x \in \Lambda \ \text{and } t > 0. \end{array}$

Then, we consider the following modified problem

$$(-\Delta)^s_{A_{\varepsilon}}u + V(\varepsilon x)u + \phi^t_{|u|}u = g_{\varepsilon}(x, |u|^2)u \text{ in } \mathbb{R}^3,$$
(3.2)

where $g_{\varepsilon}(x,t) = g(\varepsilon x,t)$ and $\phi_{|u|}^t$ is given by (2.4).

Let us note that if u is a solution of (3.2) such that

$$|u(x)| \le a \text{ for all } x \in \Lambda_{\varepsilon}^{c}, \tag{3.3}$$

where $\Lambda_{\varepsilon} := \{x \in \mathbb{R}^N : \varepsilon x \in \Lambda\}$, then u is also a solution of the original problem (3.1).

In order to find weak solutions to (3.1), we look for critical points of the Euler-Lagrange functional $J_{\varepsilon}: H^s_{\varepsilon} \to \mathbb{R}$ defined as

$$J_{\varepsilon}(u) = \frac{1}{2} \int_{\mathbb{R}^3} |(-\Delta)_{A_{\varepsilon}}^{\frac{s}{2}} u|^2 + V(\varepsilon x) |u|^2 \, dx + \frac{1}{4} \int_{\mathbb{R}^3} \phi_{|u|}^t |u|^2 \, dx - \frac{1}{2} \int_{\mathbb{R}^3} G(\varepsilon x, |u|^2) \, dx.$$

We also consider the autonomous problem associated to (3.1), that is

$$(-\Delta)^{s} u + V_{0} u + \phi^{t}_{|u|} u = f(u^{2}) u \text{ in } \mathbb{R}^{3},$$
(3.4)

and we denote by $J_0: H^s(\mathbb{R}^3, \mathbb{R}) \to \mathbb{R}$ the corresponding energy functional

$$J_0(u) = \frac{1}{2} \int_{\mathbb{R}^3} |(-\Delta)^{\frac{s}{2}} u|^2 + V_0 |u|^2 \, dx + \frac{1}{4} \int_{\mathbb{R}^3} \phi^t_{|u|} u^2 \, dx - \frac{1}{2} \int_{\mathbb{R}^3} F(u^2) \, dx$$
$$= \frac{1}{2} ||u||_0^2 + \frac{1}{4} \int_{\mathbb{R}^3} \phi^t_{|u|} u^2 \, dx - \frac{1}{2} \int_{\mathbb{R}^3} F(u^2) \, dx$$

where we used the notation $\|\cdot\|_0$ to indicate the $H^s(\mathbb{R}^3,\mathbb{R})$ -norm (equivalent to the standard one). In what follows, we show that J_{ε} verifies the assumptions of the Mountain Pass Theorem [3].

Lemma 3.1. The functional J_{ε} possesses a Mountain Pass geometry: (i) $J_{\varepsilon}(0) = 0$;

- (ii) there exists $\alpha, \rho > 0$ such that $J_{\varepsilon}(u) \ge \alpha$ for any $u \in H^s_{\varepsilon}$ such that $||u||_{\varepsilon} = \rho$;
- (iii) there exists $e \in H^s_{\varepsilon}$ with $||e||_{\varepsilon} > \rho$ such that $J_{\varepsilon}(e) < 0$.

Proof. The condition (i) is obvious. By using (g_1) and (g_2) , and Theorem 2.1 we can see that for any $\delta > 0$ there exists $C_{\delta} > 0$ such that

$$J_{\varepsilon}(u) \geq \frac{1}{2} \|u\|_{\varepsilon}^2 - \delta C \|u\|_{\varepsilon}^4 - C_{\delta} \|u\|_{\varepsilon}^q.$$

Choosing $\delta > 0$ sufficiently small, we can see that (*ii*) holds. Regarding (*iii*), we can note that in view of (g_3) , we have for any $u \in H^s_{\varepsilon} \setminus \{0\}$ with $supp(u) \subset \Lambda_{\varepsilon}$ and T > 1

$$J_{\varepsilon}(Tu) \leq \frac{T^2}{2} \|u\|_{\varepsilon}^2 + \frac{T^4}{4} \int_{\mathbb{R}^3} \phi_{|u|}^t |u|^2 dx - \frac{1}{2} \int_{\Lambda_{\varepsilon}} G(\varepsilon x, T^2 |u|^2) dx$$
$$\leq \frac{T^4}{2} \left(\|u\|_{\varepsilon}^2 + \int_{\mathbb{R}^3} \phi_{|u|}^t |u|^2 dx \right) - CT^{\theta} \int_{\Lambda_{\varepsilon}} |u|^{\theta} dx + C$$

which implies that $J_{\varepsilon}(Tu) \to -\infty$ as $T \to \infty$ being $\theta > 4$.

Lemma 3.2. Let $c \in \mathbb{R}$. Then J_{ε} satisfies the Palais-Smale condition at the level c.

Proof. Let $(u_n) \subset H^s_{\varepsilon}$ be a (PS)-sequence. Then (u_n) is bounded in H^s_{ε} . Indeed, using (g_3) we have

$$\begin{split} c + o_n(1) \|u_n\|_{\varepsilon} &= J_{\varepsilon}(u_n) - \frac{1}{\theta} \langle J_{\varepsilon}'(u_n), u_n \rangle \\ &= \left(\frac{1}{2} - \frac{1}{\theta}\right) \|u_n\|_{\varepsilon}^2 + \left(\frac{1}{4} - \frac{1}{\theta}\right) \int_{\mathbb{R}^3} \phi_{|u_n|}^t |u_n|^2 dx \\ &+ \frac{1}{\theta} \int_{\mathbb{R}^3} g_{\varepsilon}(x, |u_n|^2) |u_n|^2 dx - \frac{1}{2} \int_{\mathbb{R}^3} G_{\varepsilon}(x, |u_n|^2) dx \\ &\geq \left(\frac{1}{2} - \frac{1}{\theta}\right) \|u_n\|_{\varepsilon}^2 + \frac{1}{\theta} \int_{\Lambda_{\varepsilon}} \left(g_{\varepsilon}(x, |u_n|^2) |u_n|^2 - \frac{\theta}{2} G(\varepsilon x, |u_n|^2)\right) dx \\ &+ \frac{1}{\theta} \int_{\Lambda_{\varepsilon}^c} g_{\varepsilon}(x, |u_n|^2) |u_n|^2 dx - \frac{1}{2} \int_{\mathbb{R}^3} \frac{V(\varepsilon x)}{k} |u_n|^2 dx \\ &\geq \frac{1}{2} \left(\frac{\theta - 2}{\theta} - \frac{1}{k}\right) \|u_n\|_{\varepsilon}^2, \end{split}$$

and recalling that $k > \frac{\theta}{\theta-2}$ we get the thesis. Now, we show that for any $\xi > 0$ there exists $R = R_{\xi} > 0$ such that $\Lambda_{\varepsilon} \subset B_R$ and

$$\limsup_{n \to \infty} \int_{B_R^c} |(-\Delta)_{A_\varepsilon}^s u_n|^2 + V_\varepsilon(x)|u_n|^2 \, dx \le \xi.$$
(3.5)

Assume for the moment that the above claim holds, and we show how this information can be used. By using $u_n \rightharpoonup u$ in H^s_{ε} , Theorem 2.1 and (g_1) - (g_2) , it is easy to see that

$$(u_n,\psi)_{\varepsilon} \to (u,\psi)_{\varepsilon} \text{ and } \Re\left(\int_{\mathbb{R}^3} g(\varepsilon x,|u_n|^2)u_n\bar{\psi}dx\right) \to \Re\left(\int_{\mathbb{R}^3} g(\varepsilon x,|u|^2)u\bar{\psi}dx\right).$$
 (3.6)

Moreover, by using (3.5) and Theorem 2.1 we can see that for all $\xi > 0$ there exists $R = R_{\xi} > 0$ such that for any n large enough

$$\begin{aligned} \|u_n - u\|_{L^q(\mathbb{R}^3)} &= \|u_n - u\|_{L^q(B_R)} + \|u_n - u\|_{L^q(B_R^c)} \\ &\leq \|u_n - u\|_{L^q(B_R)} + (\|u_n\|_{L^q(B_R^c)} + \|u\|_{L^q(B_R^c)}) \\ &\leq \xi + 2C\xi, \end{aligned}$$

where $q \in [2, 2_s^*)$, which gives

$$u_n \to u \text{ in } L^q(\mathbb{R}^3, \mathbb{C}) \quad \forall q \in [2, 2^*_s).$$
 (3.7)

Since $||u_n| - |u|| \leq |u_n - u|$ and $\frac{12}{3+2t} \in (2, 2^*_s)$, we also have $|u_n| \to |u|$ in $L^{\frac{12}{3+2t}}(\mathbb{R}^3, \mathbb{R})$. Then, recalling that $\phi_{|u|} : L^{\frac{12}{3+27}}(\mathbb{R}^3, \mathbb{R}) \to D^{t,2}(\mathbb{R}^3, \mathbb{R})$ is continuous (see Lemma 2.5) we can deduce

that

$$\phi_{|u_n|}^t \to \phi_{|u|}^t \text{ in } D^{t,2}(\mathbb{R}^3, \mathbb{R}).$$

$$(3.8)$$

Putting together (3.7), (3.8), Hölder inequality and Theorem 2.1 we obtain

$$\Re\left(\int_{\mathbb{R}^{3}}(\phi_{|u_{n}|}^{t}u_{n}-\phi_{|u|}^{t}u)\bar{\psi}dx\right) = \Re\left(\int_{\mathbb{R}^{3}}\phi_{|u_{n}|}^{t}(u_{n}-u)\bar{\psi}+\int_{\mathbb{R}^{3}}(\phi_{|u_{n}|}^{t}-\phi_{|u|}^{t})u\bar{\psi}dx\right)$$

$$\leq \|\phi_{|u_{n}|}^{t}\|_{L^{\frac{6}{3+2t}}(\mathbb{R}^{3})}\|u_{n}-u\|_{L^{\frac{12}{3+2t}}(\mathbb{R}^{3})}\|\psi\|_{L^{\frac{12}{3+2t}}(\mathbb{R}^{3})}$$

$$+ \|\phi_{|u_{n}|}^{t}-\phi_{|u|}^{t}\|_{\frac{6}{3+2t}}\|u\|_{L^{\frac{12}{3+2t}}(\mathbb{R}^{3})}\|\psi\|_{\frac{12}{3+2t}}$$

$$\leq C\|u_{n}-u\|_{L^{\frac{12}{3+2t}}(\mathbb{R}^{3})}+C\|\phi_{|u_{n}|}^{t}-\phi_{|u|}^{t}\|_{D^{t,2}} \to 0.$$
(3.9)

Now, we show that

$$\int_{\mathbb{R}^3} \phi_{|u_n|}^t |u_n|^2 dx \to \int_{\mathbb{R}^3} \phi_{|u|}^t |u|^2 dx.$$
(3.10)

Let us start proving that

$$|\mathbb{D}(u_n) - \mathbb{D}(u)| \le \sqrt{\mathbb{D}(||u_n|^2 - |u|^2|^{1/2})} \sqrt{\mathbb{D}(||u_n|^2 + |u|^2|^{1/2})},$$

where

$$\mathbb{D}(u) = \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} |u(x)|^2 |u(y)|^2 dx dy.$$

Indeed, taking into account $|x|^{-(3-2t)}$ is even and Theorem 9.8 in [36] (see Remark after Theorem 9.8 and recall that -3 < -(3-2t) < 0) we have

$$\begin{split} |\mathbb{D}(u_n) - \mathbb{D}(u)| &= \left| \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} |u_n(x)|^2 |u_n(y)|^2 dx dy - \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} |u(x)|^2 |u(y)|^2 dx dy \right| \\ &= \left| \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} |u_n(x)|^2 |u_n(y)|^2 dx dy + \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} |u_n(x)|^2 |u(y)|^2 dx dy \right| \\ &- \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} |u(x)|^2 |u_n(y)|^2 dx dy - \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} |u(x)|^2 |u(y)|^2 dx dy \right| \\ &= \left| \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} (|u_n(x)|^2 - |u(x)|^2 |) (|u_n(y)|^2 + |u(y)|^2) dx dy \right| \\ &\leq \iint_{\mathbb{R}^6} |x - y|^{-(3-2t)} ||u_n(x)|^2 - |u(x)|^2 |||u_n(y)|^2 + |u(y)|^2 |dx dy \\ &\leq C \sqrt{\mathbb{D}(||u_n|^2 - |u|^2|^{1/2})} \sqrt{\mathbb{D}(||u_n|^2 + |u|^2|^{1/2})}. \end{split}$$

Thus, by using Hardy-Littlewood-Sobolev inequality (see Theorem 4.3 in [36]), Hölder inequality, the boundedness of $(|u_n|)$ in $H^s(\mathbb{R}^3, \mathbb{R})$ and $|u_n| \to |u|$ in $L^{\frac{12}{3+2t}}(\mathbb{R}^3, \mathbb{R})$ we can see that

$$\begin{aligned} |\mathbb{D}(u_n) - \mathbb{D}(u)|^2 &\leq C ||||u_n|^2 - |u|^2 ||^{1/2} ||_{L^{\frac{12}{3+2t}}(\mathbb{R}^3)}^4 ||||u_n|^2 + |u|^2 ||^{1/2} ||_{L^{\frac{12}{3+2t}}(\mathbb{R}^3)}^4 \\ &\leq C |||u_n| - |u|||_{L^{\frac{12}{3+2t}}(\mathbb{R}^3)}^2 \to 0. \end{aligned}$$

Therefore, by using $\langle J'_{\varepsilon}(u_n), \psi \rangle = o_n(1)$ for all $\psi \in C_c^{\infty}(\mathbb{R}^3, \mathbb{C})$, and taking into account (3.6) and (3.9), we can check that $J'_{\varepsilon}(u) = 0$. In particular

$$||u||_{\varepsilon}^{2} + \int_{\mathbb{R}^{3}} \phi_{|u|}^{t} |u|^{2} dx = \int_{\mathbb{R}^{3}} g_{\varepsilon}(x, |u|^{2}) |u|^{2} dx.$$
(3.11)

Now, we know that $\langle J'_{\varepsilon}(u_n), u_n \rangle = o_n(1)$ is equivalent to

$$||u_n||_{\varepsilon}^2 + \int_{\mathbb{R}^3} \phi_{|u_n|}^t |u_n|^2 dx = \int_{\mathbb{R}^3} g_{\varepsilon}(x, |u_n|^2) |u_n|^2 dx + o_n(1).$$
(3.12)

From the growth assumptions (g_1) - (g_2) and using (3.5), we can see that

$$\int_{\mathbb{R}^3} g_{\varepsilon}(x, |u_n|^2) |u_n|^2 \, dx \to \int_{\mathbb{R}^3} g_{\varepsilon}(x, |u|^2) |u|^2 \, dx. \tag{3.13}$$

Then, taking into account (3.10), (3.11), (3.12) and (3.13) we can infer that

$$\lim_{n \to \infty} \|u_n\|_{\varepsilon}^2 = \|u\|_{\varepsilon}^2$$

V. AMBROSIO

It remains to prove that (3.5) holds. Let $\eta_R \in C^{\infty}(\mathbb{R}^3, \mathbb{R})$ be such that $0 \leq \eta_R \leq 1$, $\eta_R = 0$ in $B_{\frac{R}{2}}$, $\eta_R = 1$ in B_R^c and $|\nabla \eta_R| \leq \frac{C}{R}$ for some C > 0 independent of R. Since $(u_n \eta_R)$ is bounded, we can see that $\langle J_{\varepsilon}'(u_n), u_n \eta_R \rangle = o_n(1)$, that is

$$\Re\left(\iint_{\mathbb{R}^{6}}\frac{(u_{n}(x)-u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})\overline{(u_{n}(x)\eta_{R}(x)-u_{n}(y)\eta_{R}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})}}{|x-y|^{3+2s}}dxdy\right) + \int_{\mathbb{R}^{3}}\phi_{|u_{n}|}^{t}|u_{n}|^{2}\eta_{R}dx + \int_{\mathbb{R}^{3}}V_{\varepsilon}(x)\eta_{R}|u_{n}|^{2}dx = \int_{\mathbb{R}^{3}}g_{\varepsilon}(x,|u_{n}|^{2})|u_{n}|^{2}\eta_{R}dx + o_{n}(1).$$

From

$$\begin{split} \Re\left(\iint_{\mathbb{R}^{6}} \frac{(u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})\overline{(u_{n}(x)\eta_{R}(x) - u_{n}(y)\eta_{R}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})}}{|x-y|^{3+2s}} \, dxdy\right) \\ &= \Re\left(\iint_{\mathbb{R}^{6}} \overline{u_{n}(y)}e^{-iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \frac{(u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})(\eta_{R}(x) - \eta_{R}(y))}{|x-y|^{3+2s}} \, dxdy\right) \\ &+ \iint_{\mathbb{R}^{6}} \eta_{R}(x) \frac{|u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)}|^{2}}{|x-y|^{3+2s}} \, dxdy, \end{split}$$

and using (g_3) -(ii) and Lemma 2.5-(4), it follows that

$$\iint_{\mathbb{R}^{6}} \eta_{R}(x) \frac{|u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)}|^{2}}{|x-y|^{3+2s}} dxdy + \int_{\mathbb{R}^{3}} V_{\varepsilon}(x)\eta_{R}|u_{n}|^{2} dx \\
\leq -\Re\left(\iint_{\mathbb{R}^{6}} \overline{u_{n}(y)}e^{-iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \frac{(u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})(\eta_{R}(x) - \eta_{R}(y))}{|x-y|^{3+2s}} dxdy\right) \\
+ \frac{1}{k} \int_{\mathbb{R}^{3}} V_{\varepsilon}(x)\eta_{R}|u_{n}|^{2} dx + o_{n}(1).$$
(3.14)

Now, by using Hölder inequality and the boundedness of (u_n) in H^s_{ε} we get

$$\left| \Re \left(\iint_{\mathbb{R}^{6}} \overline{u_{n}(y)} e^{-iA_{\varepsilon}(\frac{x+y}{2}) \cdot (x-y)} \frac{(u_{n}(x) - u_{n}(y) e^{iA_{\varepsilon}(\frac{x+y}{2}) \cdot (x-y)})(\eta_{R}(x) - \eta_{R}(y))}{|x-y|^{3+2s}} dxdy} \right) \right| \\ \leq \left(\iint_{\mathbb{R}^{6}} \frac{|u_{n}(x) - u_{n}(y) e^{iA_{\varepsilon}(\frac{x+y}{2}) \cdot (x-y)}|^{2}}{|x-y|^{3+2s}} dxdy} \right)^{\frac{1}{2}} \left(\iint_{\mathbb{R}^{6}} |\overline{u_{n}(y)}|^{2} \frac{|\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x-y|^{3+2s}} dxdy} \right)^{\frac{1}{2}} \\ \leq C \left(\iint_{\mathbb{R}^{6}} |u_{n}(y)|^{2} \frac{|\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x-y|^{3+2s}} dxdy \right)^{\frac{1}{2}}.$$
(3.15)

In what follows, we show that

$$\limsup_{R \to \infty} \limsup_{n \to \infty} \iint_{\mathbb{R}^6} |u_n(y)|^2 \frac{|\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3 + 2s}} \, dx \, dy = 0.$$
(3.16)

Let us note that

$$\mathbb{R}^6 = ((\mathbb{R}^3 \setminus B_{2R}) \times (\mathbb{R}^3 \setminus B_{2R})) \cup ((\mathbb{R}^3 \setminus B_{2R}) \times B_{2R}) \cup (B_{2R} \times \mathbb{R}^3) =: X_R^1 \cup X_R^2 \cup X_R^3.$$

Therefore

$$\iint_{\mathbb{R}^{6}} \frac{|\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} |u_{n}(x)|^{2} dx dy = \iint_{X_{R}^{1}} \frac{|\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} |u_{n}(x)|^{2} dx dy + \iint_{X_{R}^{2}} \frac{|\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} |u_{n}(x)|^{2} dx dy + \iint_{X_{R}^{3}} \frac{|\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} |u_{n}(x)|^{2} dx dy.$$
(3.17)

Since $\eta_R = 1$ in $\mathbb{R}^3 \setminus B_{2R}$, we can see that

$$\iint_{X_R^1} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3+2s}} dx dy = 0.$$
(3.18)

Now, fix k > 4, and we observe that

 $X_R^2 = (\mathbb{R}^3 \setminus B_{2R}) \times B_{2R} \subset ((\mathbb{R}^3 \setminus B_{kR}) \times B_{2R}) \cup ((B_{kR} \setminus B_{2R}) \times B_{2R})$

If $(x, y) \in (\mathbb{R}^3 \setminus B_{kR}) \times B_{2R}$, then

$$|x - y| \ge |x| - |y| \ge |x| - 2R > \frac{|x|}{2}.$$

Therefore, using $0 \le \eta_R \le 1$, $|\nabla \eta_R| \le \frac{C}{R}$ and applying Hölder inequality we obtain

$$\begin{aligned} \iint_{X_{R}^{2}} \frac{|u_{n}(x)|^{2} |\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} dx dy \\ &= \int_{\mathbb{R}^{3} \setminus B_{kR}} \int_{B_{2R}} \frac{|u_{n}(x)|^{2} |\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} dx dy + \int_{B_{kR} \setminus B_{2R}} \int_{B_{2R}} \frac{|u_{n}(x)|^{2} |\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} dx dy \\ &\leq 2^{2 + 3 + 2s} \int_{\mathbb{R}^{3} \setminus B_{kR}} \int_{B_{2R}} \frac{|u_{n}(x)|^{2}}{|x|^{3 + 2s}} dx dy + \frac{C}{R^{2}} \int_{B_{kR} \setminus B_{2R}} \int_{B_{2R}} \frac{|u_{n}(x)|^{2}}{|x - y|^{3 + 2(s - 1)}} dx dy \\ &\leq CR^{3} \int_{\mathbb{R}^{3} \setminus B_{kR}} \frac{|u_{n}(x)|^{2}}{|x|^{3 + 2s}} dx + \frac{C}{R^{2}} (kR)^{2(1 - s)} \int_{B_{kR} \setminus B_{2R}} |u_{n}(x)|^{2} dx \\ &\leq CR^{3} \left(\int_{\mathbb{R}^{3} \setminus B_{kR}} |u_{n}(x)|^{2^{s}_{s}} dx \right)^{\frac{2}{s}} \left(\int_{\mathbb{R}^{3} \setminus B_{kR}} \frac{1}{|x|^{\frac{3^{2}}{2s} + 3}} dx \right)^{\frac{2s}{3}} + \frac{Ck^{2(1 - s)}}{R^{2s}} \int_{B_{kR} \setminus B_{2R}} |u_{n}(x)|^{2} dx \\ &\leq \frac{C}{k^{3}} \left(\int_{\mathbb{R}^{3} \setminus B_{kR}} |u_{n}(x)|^{2^{s}_{s}} dx \right)^{\frac{2}{s}} + \frac{Ck^{2(1 - s)}}{R^{2s}} \int_{B_{kR} \setminus B_{2R}} |u_{n}(x)|^{2} dx \\ &\leq \frac{C}{k^{3}} + \frac{Ck^{2(1 - s)}}{R^{2s}} \int_{B_{kR} \setminus B_{2R}} |u_{n}(x)|^{2} dx. \end{aligned}$$
(3.19)

Take $\varepsilon \in (0, 1)$, and we obtain

$$\iint_{X_{R}^{3}} \frac{|u_{n}(x)|^{2} |\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} dx dy \\
\leq \int_{B_{2R} \setminus B_{\varepsilon R}} \int_{\mathbb{R}^{3}} \frac{|u_{n}(x)|^{2} |\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} dx dy + \int_{B_{\varepsilon R}} \int_{\mathbb{R}^{3}} \frac{|u_{n}(x)|^{2} |\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} dx dy. \quad (3.20)$$

Since

and

$$\int_{B_{2R}\setminus B_{\varepsilon R}} \int_{\mathbb{R}^3 \cap \{y: |x-y| \ge R\}} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x-y|^{3+2s}} \, dx dy \le \frac{C}{R^{2s}} \int_{B_{2R}\setminus B_{\varepsilon R}} |u_n(x)|^2 dx$$
 see that

we can see that

$$\int_{B_{2R}\setminus B_{\varepsilon R}} \int_{\mathbb{R}^3} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3+2s}} \, dx \, dy \le \frac{C}{R^{2s}} \int_{B_{2R}\setminus B_{\varepsilon R}} |u_n(x)|^2 \, dx. \tag{3.21}$$

On the other hand, from the definition of η_R , $\varepsilon \in (0, 1)$, and $\eta_R \leq 1$ we obtain

$$\int_{B_{\varepsilon R}} \int_{\mathbb{R}^3} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3 + 2s}} \, dx dy = \int_{B_{\varepsilon R}} \int_{\mathbb{R}^3 \setminus B_R} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3 + 2s}} \, dx dy$$

$$\leq 4 \int_{B_{\varepsilon R}} \int_{\mathbb{R}^3 \setminus B_R} \frac{|u_n(x)|^2}{|x - y|^{3 + 2s}} \, dx dy$$

$$\leq C \int_{B_{\varepsilon R}} |u_n|^2 dx \int_{(1 - \varepsilon)R}^{\infty} \frac{1}{r^{1 + 2s}} dr$$

$$= \frac{C}{[(1 - \varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} |u_n|^2 dx \qquad (3.22)$$

where we use the fact that if $(x, y) \in B_{\varepsilon R} \times (\mathbb{R}^3 \setminus B_R)$, then $|x - y| > (1 - \varepsilon)R$. Then (3.20), (3.21) and (3.22) yield

$$\iint_{X_R^3} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3 + 2s}} \, dx \, dy$$

$$\leq \frac{C}{R^{2s}} \int_{B_{2R} \setminus B_{\varepsilon R}} |u_n(x)|^p \, dx + \frac{C}{[(1 - \varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} |u_n(x)|^2 \, dx. \tag{3.23}$$

In view of (3.17), (3.18), (3.19) and (3.23) we can infer

$$\iint_{\mathbb{R}^{6}} \frac{|u_{n}(x)|^{2} |\eta_{R}(x) - \eta_{R}(y)|^{2}}{|x - y|^{3 + 2s}} dx dy \\
\leq \frac{C}{k^{3}} + \frac{Ck^{2(1 - s)}}{R^{2s}} \int_{B_{kR} \setminus B_{2R}} |u_{n}(x)|^{2} dx + \frac{C}{R^{2s}} \int_{B_{2R} \setminus B_{\varepsilon R}} |u_{n}(x)|^{2} dx + \frac{C}{[(1 - \varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} |u_{n}(x)|^{2} dx \\$$
(3.24)

Since $(|u_n|)$ is bounded in $H^s(\mathbb{R}^3, \mathbb{R})$, by using Sobolev embedding $H^s(\mathbb{R}^3, \mathbb{R}) \subset L^{2^*_s}(\mathbb{R}^3, \mathbb{R})$ (see [20]), we may assume that $|u_n| \to u$ in $L^2_{loc}(\mathbb{R}^3, \mathbb{R})$ for some $u \in H^s(\mathbb{R}^3, \mathbb{R})$. Letting the limit as $n \to \infty$ in (3.24) we find

$$\begin{split} &\limsup_{n \to \infty} \iint_{\mathbb{R}^6} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3 + 2s}} \, dx dy \\ &\leq \frac{C}{k^3} + \frac{Ck^{2(1-s)}}{R^{2s}} \int_{B_{kR} \setminus B_{2R}} |u(x)|^2 dx + \frac{C}{R^{2s}} \int_{B_{2R} \setminus B_{\varepsilon R}} |u(x)|^2 dx + \frac{C}{[(1 - \varepsilon)R]^{2s}} \int_{B_{\varepsilon R}} |u(x)|^2 dx \\ &\leq \frac{C}{k^3} + Ck^2 \left(\int_{B_{kR} \setminus B_{2R}} |u(x)|^{2^*_s} dx \right)^{\frac{2}{2^*_s}} + C \left(\int_{B_{2R} \setminus B_{\varepsilon R}} |u(x)|^{2^*_s} dx \right)^{\frac{2}{2^*_s}} + C \left(\frac{\varepsilon}{1 - \varepsilon} \right)^{2s} \left(\int_{B_{\varepsilon R}} |u(x)|^{2^*_s} dx \right)^{\frac{2}{2^*_s}}, \end{split}$$

where in the last passage we used Hölder inequality. Since $u \in L^{2^*_s}(\mathbb{R}^3, \mathbb{R})$, k > 4 and $\varepsilon \in (0, 1)$ we can see that

$$\limsup_{R \to \infty} \int_{B_{kR} \setminus B_{2R}} |u(x)|^{2^*_s} dx = \limsup_{R \to \infty} \int_{B_{2R} \setminus B_{\varepsilon R}} |u(x)|^{2^*_s} dx = 0.$$

12

Thus, taking $\varepsilon = \frac{1}{k}$, we have

$$\begin{split} &\limsup_{R \to \infty} \limsup_{n \to \infty} \iint_{\mathbb{R}^6} \frac{|u_n(x)|^2 |\eta_R(x) - \eta_R(y)|^2}{|x - y|^{3 + 2s}} \, dx dy \\ &\leq \lim_{k \to \infty} \limsup_{R \to \infty} \left[\frac{C}{k^3} + Ck^2 \left(\int_{B_{kR} \setminus B_{2R}} |u(x)|^{2^*_s} dx \right)^{\frac{2}{2^*_s}} + C \left(\int_{B_{2R} \setminus B_{\frac{1}{k}R}} |u(x)|^{2^*_s} dx \right)^{\frac{2}{2^*_s}} \\ &+ C \left(\frac{1}{k - 1} \right)^{2s} \left(\int_{B_{\frac{1}{k}R}} |u(x)|^{2^*_s} dx \right)^{\frac{2}{2^*_s}} \right] \\ &\leq \lim_{k \to \infty} \frac{C}{k^3} + C \left(\frac{1}{k - 1} \right)^{2s} \left(\int_{\mathbb{R}^3} |u(x)|^{2^*_s} dx \right)^{\frac{2}{2^*_s}} = 0, \end{split}$$

which implies that (3.16) holds true. Putting together (3.14), (3.15) and (3.16) we can deduce that

$$\limsup_{R \to \infty} \limsup_{n \to \infty} \left(1 - \frac{1}{k} \right) \int_{B_R^c} |(-\Delta)_{A_\varepsilon}^{\frac{s}{2}} u_n|^2 + V_\varepsilon(x) |u_n|^2 \, dx = 0,$$

and this completes the proof of (3.5).

In view of Lemma 3.1, we can define the mountain pass level

$$c_{\varepsilon} = \inf_{\gamma \in \Gamma_{\varepsilon}} \max_{t \in [0,1]} J_{\varepsilon}(\gamma(t))$$

where

$$\Gamma_{\varepsilon} = \{ \gamma \in C([0,1], H^s_{\varepsilon}) : \gamma(0) = 0 \text{ and } J_{\varepsilon}(\gamma(1)) < 0 \}$$

By applying Mountain Pass Theorem [3], we can see that there exists $u_{\varepsilon} \in H^s_{\varepsilon} \setminus \{0\}$ such that $J_{\varepsilon}(u_{\varepsilon}) = c_{\varepsilon}$ and $J'_{\varepsilon}(u_{\varepsilon}) = 0$. In similar fashion, one can prove that also J_0 has a mountain pass geometry, and we denote by c_{V_0} the mountain pass level associated to J_0 . Now, let us introduce the Nehari manifold associated to (3.1), that is

$$\mathcal{N}_{\varepsilon} := \{ u \in H^s_{\varepsilon} \setminus \{0\} : \langle J'_{\varepsilon}(u), u \rangle = 0 \},\$$

and we denote by \mathcal{N}_0 the Nehari manifold associated to (3.4). It is standard to verify (see [50]) that c_{ε} can be also characterized as follows:

$$c_{\varepsilon} = \inf_{u \in H^s_{\varepsilon} \setminus \{0\}} \sup_{t \geq 0} J_{\varepsilon}(tu) = \inf_{u \in \mathcal{N}_{\varepsilon}} J_{\varepsilon}(u);$$

Next, we prove the existence of a ground state solution to (3.4).

Lemma 3.3. Let $(u_n) \subset \mathcal{N}_0$ be a sequence satisfying $J_0(u_n) \to c_{V_0}$. Then, up to subsequences, the following alternatives holds:

- (i) (u_n) strongly converges in $H^s(\mathbb{R}^3, \mathbb{R})$,
- (ii) there exists a sequence $(\tilde{y}_n) \subset \mathbb{R}^3$ such that, up to a subsequence, $v_n(x) = u_n(x + \tilde{y}_n)$ converges strongly in $H^s(\mathbb{R}^3, \mathbb{R})$.

In particular, there exists a minimizer $w \in H^s(\mathbb{R}^3, \mathbb{R})$ for J_0 with $J_0(w) = c_{V_0}$.

Proof. Since J_0 has a Mountain Pass geometry, we can use a version of the Mountain Pass Theorem without (PS) condition (see [50]), and we may suppose that (u_n) is a $(PS)_{c_{V_0}}$ sequence for J_0 . Arguing as in Lemma 3.2, it is easy to check that (u_n) is bounded in $H^s(\mathbb{R}^3, \mathbb{R})$ so we may assume

that $u_n \rightharpoonup u$ in $H^s(\mathbb{R}^3, \mathbb{R})$. The weak convergence is enough to deduce that $J'_0(u) = 0$. Now, we assume that $u \neq 0$. Since $u \in \mathcal{N}_0$, we can use (f_3) and Fatou's Lemma to see that

$$c_{V_0} \leq J_0(u) - \frac{1}{4} \langle J'_0(u), u \rangle$$

= $\frac{1}{4} ||u||_{\mu}^2 + \frac{1}{2} \int_{\mathbb{R}^3} \frac{1}{2} f(u^2) u - F(u^2) dx$
 $\leq \liminf_{n \to \infty} \left[J_0(u_n) - \frac{1}{4} \langle J'_0(u), u \rangle \right] = c_{V_0}$

which implies that $J_0(u) = c_{V_0}$.

Let us consider the case u = 0. Since $c_{V_0} > 0$ and J_0 is continuous, we can see that $||u_n||_0 \neq 0$. Then, in view of Lemma 2.4 and (f_1) - (f_2) , it is standard to prove that there are a sequence $(y_n) \subset \mathbb{R}^3$ and constants $R, \beta > 0$ such that

$$\liminf_{n \to \infty} \int_{B_R(y_n)} |u_n|^4 dx \ge \beta > 0.$$

Let us define $v_n = u_n(\cdot + y_n)$, and we note that v_n has a nontrivial weak limit v in $H^s(\mathbb{R}^3, \mathbb{R})$. It is clear that also (v_n) is a $(PS)_{c_{V_0}}$ sequence for J_0 , and arguing as before we can deduce that $J_0(v) = c_{V_0}$. In conclusion, problem (3.4) admits a ground state solution.

Now, let u be a ground state for (3.4). By using $\varphi = u^-$ as test function in $\langle J'_0(u), \varphi \rangle = 0$, it is easy to check that $u \ge 0$ in \mathbb{R}^3 . In particular, observing that $\phi^t_u \ge 0$ and f has a subcritical growth, we can argue as in Proposition 5.1.1 in [21] to see that $u \in L^{\infty}(\mathbb{R}^3, \mathbb{R})$. In particular, we have

$$\begin{split} \phi_u^t(x) &= \int_{|y-x| \ge 1} \frac{|u(y)|^2}{|x-y|^{3-2t}} dy + \int_{|y-x| < 1} \frac{|u(y)|^2}{|x-y|^{3-2t}} dy \\ &\leq \|u\|_{L^2(\mathbb{R}^3)}^2 + \|u\|_{L^\infty(\mathbb{R}^3)}^2 \int_{|y-x| < 1} \frac{1}{|x-y|^{3-2t}} dy \le C, \end{split}$$

so that $g(x) = f(u^2)u - \mu u - \phi_u^t u \in L^{\infty}(\mathbb{R}^3, \mathbb{R})$. By applying Proposition 2.9 in [46] we can deduce that $u \in C^{0,\gamma}(\mathbb{R}^3, \mathbb{R})$ for some $0 < \gamma < 1$. By using maximum principle (see Corollary 3.4 in [23]) we can see that u > 0 in \mathbb{R}^3 . Since $u \in C^{0,\alpha}(\mathbb{R}^3, \mathbb{R}) \cap L^p(\mathbb{R}^3, \mathbb{R})$ for all $p \in [2, \infty)$, we can deduce that $u(x) \to 0$ as $|x| \to \infty$, so we can find R > 0 such that $(-\Delta)^s u + \frac{V_0}{2}u \leq 0$ in |x| > R. By using Lemma 4.3 in [25] we know that there exists a positive function w such that for |x| > R (taking Rlarger if necessary) it holds $(-\Delta)^s w + \frac{V_0}{2}w \geq 0$ and $w(x) = \frac{C_0}{|x|^{3+2s}}$, for some $C_0 > 0$. In view of the continuity of u and w there exists some constant $C_1 > 0$ such that $z := u - C_1 w \leq 0$ on |x| = R. Moreover, we can see that $(-\Delta)^s z + \frac{V_0}{2} z \geq 0$ in $|x| \geq R$. Then, it follows by the maximum principle that $z \leq 0$ in $|x| \geq R$, that is $0 < u(x) \leq C_1 w(x) \leq \frac{C_2}{|x|^{3+2s}}$ for all |x| big enough.

Now we prove the following interesting relation between c_{ε} and c_{V_0} .

Lemma 3.4. The numbers c_{ε} and c_{V_0} satisfy the following inequality

$$\limsup_{\varepsilon \to 0} c_{\varepsilon} \le c_{V_0}$$

Proof. Let $w \in H^s(\mathbb{R}^3, \mathbb{R})$ be a positive ground state to the autonomous problem (3.4) (see [48,51]), so $J'_0(w) = 0$ and $J_0(w) = c_{V_0}$. We recall that $w \in C^{0,\gamma}(\mathbb{R}^3,\mathbb{R}) \cap L^{\infty}(\mathbb{R}^3,\mathbb{R})$. Moreover, using Lemma 4.3 in [25], it is easy to check that w satisfies the following decay estimate:

$$0 < w(x) \le \frac{C}{|x|^{3+2s}} \text{ for all } |x| > 1.$$
(3.25)

Let $\eta \in C_c^{\infty}(\mathbb{R}^3, [0, 1])$ be a cut-off function such that $\eta = 1$ in a neighborhood of zero $B_{\frac{\delta}{2}}$ and $\operatorname{supp}(\eta) \subset B_{\delta} \subset \Lambda$ for some $\delta > 0$.

Let us define $w_{\varepsilon}(x) := \eta_{\varepsilon}(x)w(x)e^{iA(0)\cdot x}$, with $\eta_{\varepsilon}(x) = \eta(\varepsilon x)$ for $\varepsilon > 0$, and we observe that $|w_{\varepsilon}| = \eta_{\varepsilon}w$ and $w_{\varepsilon} \in H^{s}_{\varepsilon}$ in view of Lemma 2.3. Now we prove that

$$\lim_{\varepsilon \to 0} \|w_{\varepsilon}\|_{\varepsilon}^{2} = \|w\|_{0}^{2} \in (0, \infty).$$

$$(3.26)$$

Since it is clear that $\int_{\mathbb{R}^3} V_{\varepsilon}(x) |w_{\varepsilon}|^2 dx \to \int_{\mathbb{R}^3} V_0 |w|^2 dx$, we only need to show that

$$\lim_{\varepsilon \to 0} [w_{\varepsilon}]^2_{A_{\varepsilon}} = [w]^2.$$
(3.27)

By using Lemma 5 in [42] we know that

$$[\eta_{\varepsilon}w] \to [w] \text{ as } \varepsilon \to 0. \tag{3.28}$$

On the other hand

$$\begin{split} [w_{\varepsilon}]^2_{A_{\varepsilon}} &= \iint_{\mathbb{R}^6} \frac{|e^{iA(0) \cdot x} \eta_{\varepsilon}(x)w(x) - e^{iA_{\varepsilon}(\frac{x+y}{2}) \cdot (x-y)} e^{iA(0) \cdot y} \eta_{\varepsilon}(y)w(y)|^2}{|x-y|^{3+2s}} dxdy \\ &= [\eta_{\varepsilon}w]^2 + \iint_{\mathbb{R}^6} \frac{\eta_{\varepsilon}^2(y)w^2(y)|e^{i[A_{\varepsilon}(\frac{x+y}{2}) - A(0)] \cdot (x-y)} - 1|^2}{|x-y|^{3+2s}} dxdy \\ &\quad + 2\Re \iint_{\mathbb{R}^6} \frac{(\eta_{\varepsilon}(x)w(x) - \eta_{\varepsilon}(y)w(y))\eta_{\varepsilon}(y)w(y)(1 - e^{-i[A_{\varepsilon}(\frac{x+y}{2}) - A(0)] \cdot (x-y)})}{|x-y|^{3+2s}} dxdy \\ &=: [\eta_{\varepsilon}w]^2 + X_{\varepsilon} + 2Y_{\varepsilon}. \end{split}$$

Then, in view of $|Y_{\varepsilon}| \leq [\eta_{\varepsilon}w]\sqrt{X_{\varepsilon}}$ and (3.28), it is suffices to prove that $X_{\varepsilon} \to 0$ as $\varepsilon \to 0$ to deduce that (3.27) holds. Let us note that for $0 < \beta < \alpha/(1 + \alpha - s)$,

$$X_{\varepsilon} \leq \int_{\mathbb{R}^{3}} w^{2}(y) dy \int_{|x-y| \geq \varepsilon^{-\beta}} \frac{|e^{i[A_{\varepsilon}(\frac{x+y}{2}) - A(0)] \cdot (x-y)} - 1|^{2}}{|x-y|^{3+2s}} dx$$

+
$$\int_{\mathbb{R}^{3}} w^{2}(y) dy \int_{|x-y| < \varepsilon^{-\beta}} \frac{|e^{i[A_{\varepsilon}(\frac{x+y}{2}) - A(0)] \cdot (x-y)} - 1|^{2}}{|x-y|^{3+2s}} dx$$

=:
$$X_{\varepsilon}^{1} + X_{\varepsilon}^{2}.$$
 (3.29)

Using $|e^{it} - 1|^2 \leq 4$ and $w \in H^s(\mathbb{R}^3, \mathbb{R})$, we get

$$X_{\varepsilon}^{1} \leq C \int_{\mathbb{R}^{3}} w^{2}(y) dy \int_{\varepsilon^{-\beta}}^{\infty} \rho^{-1-2s} d\rho \leq C \varepsilon^{2\beta s} \to 0.$$
(3.30)

Since $|e^{it} - 1|^2 \le t^2$ for all $t \in \mathbb{R}$, $A \in C^{0,\alpha}(\mathbb{R}^3, \mathbb{R}^3)$ for $\alpha \in (0, 1]$, and $|x + y|^2 \le 2(|x - y|^2 + 4|y|^2)$, we have

$$\begin{split} X_{\varepsilon}^{2} &\leq \int_{\mathbb{R}^{3}} w^{2}(y) dy \int_{|x-y| < \varepsilon^{-\beta}} \frac{|A_{\varepsilon} \left(\frac{x+y}{2}\right) - A(0)|^{2}}{|x-y|^{3+2s-2}} dx \\ &\leq C \varepsilon^{2\alpha} \int_{\mathbb{R}^{3}} w^{2}(y) dy \int_{|x-y| < \varepsilon^{-\beta}} \frac{|x+y|^{2\alpha}}{|x-y|^{3+2s-2}} dx \\ &\leq C \varepsilon^{2\alpha} \left(\int_{\mathbb{R}^{3}} w^{2}(y) dy \int_{|x-y| < \varepsilon^{-\beta}} \frac{1}{|x-y|^{3+2s-2-2\alpha}} dx \\ &+ \int_{\mathbb{R}^{3}} |y|^{2\alpha} w^{2}(y) dy \int_{|x-y| < \varepsilon^{-\beta}} \frac{1}{|x-y|^{3+2s-2-2\alpha}} dx \\ &=: C \varepsilon^{2\alpha} (X_{\varepsilon}^{2,1} + X_{\varepsilon}^{2,2}). \end{split}$$
(3.31)

Then

$$X_{\varepsilon}^{2,1} = C \int_{\mathbb{R}^3} w^2(y) dy \int_0^{\varepsilon^{-\beta}} \rho^{1+2\alpha-2s} d\rho \le C \varepsilon^{-2\beta(1+\alpha-s)} \,. \tag{3.32}$$

On the other hand, using (3.25), we infer that

$$\begin{aligned} X_{\varepsilon}^{2,2} &\leq C \int_{\mathbb{R}^{3}} |y|^{2\alpha} w^{2}(y) dy \int_{0}^{\varepsilon^{-\beta}} \rho^{1-2s} d\rho \\ &\leq C \, \varepsilon^{-2\beta(1-s)} \left[\int_{B_{1}(0)} w^{2}(y) dy + \int_{B_{1}^{c}(0)} \frac{1}{|y|^{2(3+2s)-2\alpha}} dy \right] \\ &\leq C \, \varepsilon^{-2\beta(1-s)} \,. \end{aligned}$$
(3.33)

Taking into account (3.29), (3.30), (3.31), (3.32) and (3.33) we can conclude that $X_{\varepsilon} \to 0$. Therefore (3.26) holds. Moreover, since $\eta_{\varepsilon} w$ strongly converges to w in $H^s(\mathbb{R}^3, \mathbb{R})$, we can use Lemma 2.4-(5) in [37] to see that

$$\lim_{\varepsilon \to 0} \int_{\mathbb{R}^3} \phi^t_{|w_\varepsilon|} |w_\varepsilon|^2 dx = \int_{\mathbb{R}^3} \phi^t_w w^2 dx.$$
(3.34)

Now, let $t_{\varepsilon} > 0$ be the unique number such that

$$J_{\varepsilon}(t_{\varepsilon}w_{\varepsilon}) = \max_{t \ge 0} J_{\varepsilon}(tw_{\varepsilon}).$$

Then t_{ε} verifies

$$\|w_{\varepsilon}\|_{\varepsilon}^{2} + t_{\varepsilon}^{2} \int_{\mathbb{R}^{3}} \phi_{|w_{\varepsilon}|}^{t} |w_{\varepsilon}|^{2} dx = \int_{\mathbb{R}^{3}} g(\varepsilon \, x, t_{\varepsilon}^{2} |w_{\varepsilon}|^{2}) |w_{\varepsilon}|^{2} dx = \int_{\mathbb{R}^{3}} f(t_{\varepsilon}^{2} |w_{\varepsilon}|^{2}) |w_{\varepsilon}|^{2} dx \tag{3.35}$$

where we used $supp(\eta) \subset \Lambda$ and g = f on Λ .

Let us prove that $t_{\varepsilon} \to 1$ as $\varepsilon \to 0$. Using $\eta = 1$ in $B_{\frac{\delta}{2}}$ and that w is a continuous positive function, we can see that (f_4) yields

$$\frac{1}{t_{\varepsilon}^2} \|w_{\varepsilon}\|_{\varepsilon}^2 + \int_{\mathbb{R}^3} \phi_{|w_{\varepsilon}|}^t |w_{\varepsilon}|^2 dx \ge \frac{f(t_{\varepsilon}^2 \alpha_0^2)}{t_{\varepsilon}^2 \alpha_0^2} \int_{B_{\frac{\delta}{2}}} |w|^2 dx,$$

where $\alpha_0 = \min_{\bar{B}_{\frac{\delta}{2}}} w > 0$. So, if $t_{\varepsilon} \to \infty$ as $\varepsilon \to 0$, we can use (f_3) , (3.34) and (3.26) to deduce that $\int_{\mathbb{R}^3} \phi_{|w|,t} |w|^2 dx = \infty$, that is a contradiction. On the other hand, if $t_{\varepsilon} \to 0$ as $\varepsilon \to 0$, we can use the growth assumptions on g, (3.34), (3.26) to infer that $||w||_0^2 = 0$ which gives an absurd. Therefore,

 $t_{\varepsilon} \to t_0 \in (0, \infty)$ as $\varepsilon \to 0$. Now, taking the limit as $\varepsilon \to 0$ in (3.35) and using (3.34), (3.26), we can deduce that

$$\frac{1}{t_0^2} \|w\|_0^2 + \int_{\mathbb{R}^3} \phi_{|w|}^t |w|^2 dx = \int_{\mathbb{R}^3} \frac{f(t_0^2 |w|^2)}{(t_0^2 |w|^2)} |w|^4 dx.$$
(3.36)

Then $t_0 = 1$ as a consequence of $w \in \mathcal{N}_0$ and (f_4) . By applying the Dominated Convergence Theorem, we know that

$$\int_{\mathbb{R}^3} F(|t_\varepsilon w_\varepsilon|^2) \, dx \to \int_{\mathbb{R}^3} F(|w|^2) \, dx$$

so we have $\lim_{\varepsilon \to 0} J_{\varepsilon}(t_{\varepsilon}w_{\varepsilon}) = J_0(w) = c_{V_0}$. Since $c_{\varepsilon} \leq \max_{t \geq 0} J_{\varepsilon}(tw_{\varepsilon}) = J_{\varepsilon}(t_{\varepsilon}w_{\varepsilon})$, we can infer that $\limsup_{\varepsilon \to 0} c_{\varepsilon} \leq c_{V_0}$.

Now, we prove the following useful compactness result.

Lemma 3.5. Let $\varepsilon_n \to 0$ and $(u_n) \subset H^s_{\varepsilon_n}$ such that $J_{\varepsilon_n}(u_n) = c_{\varepsilon_n}$ and $J'_{\varepsilon_n}(u_n) = 0$. Then there exists $(\tilde{y}_n) \subset \mathbb{R}^3$ such that $v_n(x) = |u_n|(x + \tilde{y}_n)$ has a convergent subsequence in $H^s(\mathbb{R}^3, \mathbb{R})$. Moreover, up to a subsequence, $y_n = \varepsilon_n \tilde{y}_n \to y_0$ for some $y_0 \in \Lambda$ such that $V(y_0) = V_0$.

Proof. Since $\langle J_{\varepsilon_n}'(u_n), u_n \rangle = 0$, $J_{\varepsilon_n}(u_n) = c_{\varepsilon_n}$ and using Lemma 3.4, we can see that (u_n) is bounded in $H_{\varepsilon_n}^s$. Then, there exists C > 0 (independent of n) such that $||u_n||_{\varepsilon_n} \leq C$ for all $n \in \mathbb{N}$. Moreover, from Lemma 2.2, we also know that $(|u_n|)$ is bounded in $H^s(\mathbb{R}^3, \mathbb{R})$.

Now, we prove that there exist a sequence $(\tilde{y}_n) \subset \mathbb{R}^3$, and constants R > 0 and $\gamma > 0$ such that

$$\liminf_{n \to \infty} \int_{B_R(\tilde{y}_n)} |u_n|^2 \, dx \ge \gamma > 0. \tag{3.37}$$

Assume by contradiction (3.37) does not hold, so that, for all R > 0 we get

$$\lim_{n \to \infty} \sup_{y \in \mathbb{R}^3} \int_{B_R(y)} |u_n|^2 \, dx = 0$$

By using the boundedness of $(|u_n|)$ and Lemma 2.4, we know that $|u_n| \to 0$ in $L^q(\mathbb{R}^3, \mathbb{R})$ for any $q \in (2, 2^*_s)$. This fact and (g_1) and (g_2) yield

$$\lim_{n \to \infty} \int_{\mathbb{R}^3} g(\varepsilon_n x, |u_n|^2) |u_n|^2 \, dx = 0 = \lim_{n \to \infty} \int_{\mathbb{R}^3} G(\varepsilon_n x, |u_n|^2) \, dx. \tag{3.38}$$

On the other hand, $|u_n| \to 0$ in $L^{\frac{12}{3+2t}}(\mathbb{R}^3,\mathbb{R})$, so by Lemma 2.5-(4) we deduce that

$$\int_{\mathbb{R}^3} \phi_{|u_n|}^t |u_n|^2 dx \to 0.$$
(3.39)

Taking into account $\langle J'_{\varepsilon_n}(u_n), u_n \rangle = 0$, (3.38) and (3.39) we can infer that $||u_n||_{\varepsilon_n} \to 0$ as $n \to \infty$. This is impossible because (g_1) , (g_2) and $\langle J'_{\varepsilon_n}(u_n), u_n \rangle = 0$ imply that there exists $\alpha_0 > 0$ such that $||u_n||_{\varepsilon_n}^2 \ge \alpha_0$ for all $n \in \mathbb{N}$. Now, we set $v_n(x) = |u_n|(x + \tilde{y}_n)$. Then (v_n) is bounded in $H^s(\mathbb{R}^3, \mathbb{R})$, and we may assume that $v_n \to v \not\equiv 0$ in $H^s(\mathbb{R}^3, \mathbb{R})$ as $n \to \infty$. Fix $t_n > 0$ such that $\tilde{v}_n = t_n v_n \in \mathcal{N}_0$. In view of Lemma 2.2, we have

$$c_{V_0} \le J_0(\tilde{v}_n) \le \max_{t \ge 0} J_{\varepsilon_n}(tv_n) = J_{\varepsilon_n}(u_n)$$

which together with Lemma 3.4 yields $J_0(\tilde{v}_n) \to c_{V_0}$. Then, $\tilde{v}_n \to 0$ in $H^s(\mathbb{R}^3, \mathbb{R})$. Since (v_n) and (\tilde{v}_n) are bounded in $H^s(\mathbb{R}^3, \mathbb{R})$ and $\tilde{v}_n \to 0$ in $H^s(\mathbb{R}^3, \mathbb{R})$, we deduce that $t_n \to t^* > 0$. From the uniqueness of the weak limit, we can deduce that $\tilde{v}_n \to \tilde{v} = t^*v \neq 0$ in $H^s(\mathbb{R}^3, \mathbb{R})$, and by using Lemma 3.3, we can infer that

$$\tilde{v}_n \to \tilde{v} \text{ in } H^s(\mathbb{R}^3, \mathbb{R}).$$
 (3.40)

Therefore, $v_n \to v$ in $H^s(\mathbb{R}^3, \mathbb{R})$ as $n \to \infty$.

V. AMBROSIO

Now, we define $y_n = \varepsilon_n \tilde{y}_n$ and we show that (y_n) admits a subsequence, still denoted by y_n , such that $y_n \to y_0$ for some $y_0 \in \Lambda$ such that $V(y_0) = V_0$. Firstly, we prove that (y_n) is bounded. Assume by contradiction that, up to a subsequence, $|y_n| \to \infty$ as $n \to \infty$. Take R > 0 such that $\Lambda \subset B_R(0)$. Since we may suppose that $|y_n| > 2R$, we have that for any $z \in B_{R/\varepsilon_n}$

$$|\varepsilon_n z + y_n| \ge |y_n| - |\varepsilon_n z| > R.$$

Now, by using $(u_n) \subset \mathcal{N}_{\varepsilon_n}$, (V_1) , Lemma 2.2, Lemma 2.5 and the change of variable $x \mapsto z + \tilde{y}_n$ we observe that

$$[v_{n}]^{2} + \int_{\mathbb{R}^{3}} V_{0} v_{n}^{2} dx [v_{n}]^{2} + \int_{\mathbb{R}^{3}} V_{0} v_{n}^{2} dx + \int_{\mathbb{R}^{3}} \phi_{|v_{n}|}^{t} v_{n}^{2} dx$$

$$\leq \int_{\mathbb{R}^{3}} g(\varepsilon_{n} x + y_{n}, |v_{n}|^{2}) |v_{n}|^{2} dx$$

$$\leq \int_{B_{\frac{R}{\varepsilon_{n}}}(0)} \tilde{f}(|v_{n}|^{2}) |v_{n}|^{2} dx + \int_{\mathbb{R}^{3} \setminus B_{\frac{R}{\varepsilon_{n}}}(0)} f(|v_{n}|^{2}) |v_{n}|^{2} dx.$$
(3.41)
(3.41)
(3.42)

Since $v_n \to v$ in $H^s(\mathbb{R}^3, \mathbb{R})$ as $n \to \infty$ and $\tilde{f}(t) \leq \frac{V_0}{k}$, we can see that (3.41) yields

$$\min\left\{1, V_0\left(1 - \frac{1}{k}\right)\right\} \left([v_n]^2 + \int_{\mathbb{R}^3} |v_n|^2 \, dx\right) = o_n(1)$$

that is $v_n \to 0$ in $H^s(\mathbb{R}^3, \mathbb{R})$ and this gives a contradiction. Thus, (y_n) is bounded and we may assume that $y_n \to y_0 \in \mathbb{R}^3$. If $y_0 \notin \overline{\Lambda}$, we can proceed as before to deduce that $v_n \to 0$ in $H^s(\mathbb{R}^3, \mathbb{R})$. Therefore $y_0 \in \overline{\Lambda}$. We observe that if $V(y_0) = V_0$, then $y_0 \notin \partial \Lambda$ in view of (V_2) . Then, it is enough to verify that $V(y_0) = V_0$. Otherwise, we suppose that $V(y_0) > V_0$, and putting together (3.40), Fatou's Lemma, the invariance of \mathbb{R}^3 by translations, Lemma 2.2 and Lemma 3.4, we have

$$\begin{aligned} c_{V_0} &= J_0(\tilde{v}) < \frac{1}{2} [\tilde{v}]^2 + \frac{1}{2} \int_{\mathbb{R}^3} V(y_0) \tilde{v}^2 \, dx + \frac{1}{4} \int_{\mathbb{R}^3} \phi_{|\tilde{v}|}^t \tilde{v}^2 dx - \frac{1}{2} \int_{\mathbb{R}^3} F(|\tilde{v}|^2) \, dx \\ &\leq \liminf_{n \to \infty} \left[\frac{1}{2} [\tilde{v}_n]^2 + \frac{1}{2} \int_{\mathbb{R}^3} V(\varepsilon_n \, x + y_n) |\tilde{v}_n|^2 \, dx + \frac{1}{4} \int_{\mathbb{R}^3} \phi_{|\tilde{v}_n|}^t |\tilde{v}_n|^2 dx - \frac{1}{2} \int_{\mathbb{R}^3} F(|\tilde{v}_n|^2) \, dx \right] \\ &\leq \liminf_{n \to \infty} \left[\frac{t_n^2}{2} [|u_n|]^2 + \frac{t_n^2}{2} \int_{\mathbb{R}^3} V(\varepsilon_n \, z) |u_n|^2 \, dz + \frac{t_n^4}{4} \int_{\mathbb{R}^3} \phi_{|u_n|}^t |u_n|^2 dx - \frac{1}{2} \int_{\mathbb{R}^3} F(|t_n u_n|^2) \, dz \right] \\ &\leq \liminf_{n \to \infty} J_{\varepsilon_n}(t_n u_n) \leq \liminf_{n \to \infty} J_{\varepsilon_n}(u_n) \leq c_{V_0} \end{aligned}$$

which is a contradiction. This ends the proof of this lemma.

4. Proof of Theorem 1.1

This section is devoted to the proof of the main theorem of this work. Firstly, we prove the following lemma which plays a fundamental role to show that the solutions of (3.1) are indeed solutions to (1.1).

Lemma 4.1. Let $\varepsilon_n \to 0$ and $u_n \in H^s_{\varepsilon_n}$ be a solution to (3.2). Then $v_n = |u_n|(\cdot + \tilde{y}_n)$ satisfies $v_n \in L^{\infty}(\mathbb{R}^3, \mathbb{R})$ and there exists C > 0 such that

$$||v_n||_{L^{\infty}(\mathbb{R}^3)} \leq C \text{ for all } n \in \mathbb{N}$$

where \tilde{y}_n is given by Lemma 3.5. Moreover it holds

$$\lim_{|x|\to\infty} v_n(x) = 0 \text{ uniformly in } n \in \mathbb{N}.$$

Proof. For any L > 0, we define $u_{L,n} := \min\{|u_n|, L\} \ge 0$ and we set $v_{L,n} = u_{L,n}^{2(\beta-1)}u_n$ where $\beta > 1$ will be chosen later. Taking $v_{L,n}$ as test function in (3.2) we can see that

$$\Re\left(\iint_{\mathbb{R}^{6}} \frac{(u_{n}(x) - u_{n}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)})}{|x-y|^{3+2s}} \overline{(u_{n}u_{L,n}^{2(\beta-1)}(x) - u_{n}u_{L,n}^{2(\beta-1)}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)})} \, dxdy\right)$$

$$= -\int_{\mathbb{R}^{3}} \phi_{|u_{n}|}^{t} |u_{n}|^{2} u_{L,n}^{2(\beta-1)} \, dx + \int_{\mathbb{R}^{3}} g(\varepsilon x, |u_{n}|^{2}) |u_{n}|^{2} u_{L,n}^{2(\beta-1)} \, dx - \int_{\mathbb{R}^{3}} V(\varepsilon x) |u_{n}|^{2} u_{L,n}^{2(\beta-1)} \, dx. \quad (4.1)$$

Let us observe that

$$\begin{aligned} &\Re\left[\left(u_{n}(x)-u_{n}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)}\right)\overline{\left(u_{n}u_{L,n}^{2(\beta-1)}(x)-u_{n}u_{L,n}^{2(\beta-1)}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)}\right)}\right] \\ &=\Re\left[|u_{n}(x)|^{2}v_{L}^{2(\beta-1)}(x)-u_{n}(x)\overline{u_{n}(y)}u_{L,n}^{2(\beta-1)}(y)e^{-iA(\frac{x+y}{2})\cdot(x-y)}-u_{n}(y)\overline{u_{n}(x)}u_{L,n}^{2(\beta-1)}(x)e^{iA(\frac{x+y}{2})\cdot(x-y)}\right) \\ &+|u_{n}(y)|^{2}u_{L,n}^{2(\beta-1)}(y)\right] \\ &\geq (|u_{n}(x)|^{2}u_{L,n}^{2(\beta-1)}(x)-|u_{n}(x)||u_{n}(y)|u_{L,n}^{2(\beta-1)}(y)-|u_{n}(y)||u_{n}(x)|u_{L,n}^{2(\beta-1)}(x)+|u_{n}(y)|^{2}u_{L,n}^{2(\beta-1)}(y) \\ &= (|u_{n}(x)|-|u_{n}(y)|)(|u_{n}(x)|u_{L,n}^{2(\beta-1)}(x)-|u_{n}(y)|u_{L,n}^{2(\beta-1)}(y)), \end{aligned}$$

from which we deduce that

$$\Re\left(\iint_{\mathbb{R}^{6}} \frac{(u_{n}(x) - u_{n}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)})}{|x-y|^{3+2s}} \overline{(u_{n}u_{L,n}^{2(\beta-1)}(x) - u_{n}u_{L,n}^{2(\beta-1)}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)})} \, dxdy\right)$$

$$\geq \iint_{\mathbb{R}^{6}} \frac{(|u_{n}(x)| - |u_{n}(y)|)}{|x-y|^{3+2s}} (|u_{n}(x)|u_{L,n}^{2(\beta-1)}(x) - |u_{n}(y)|u_{L,n}^{2(\beta-1)}(y)) \, dxdy. \tag{4.2}$$

For all $t \ge 0$, let us define

$$\gamma(t) = \gamma_{L,\beta}(t) = t t_L^{2(\beta-1)}$$

where $t_L = \min\{t, L\}$. Let us observe that, since γ is an increasing function, then it holds

$$(a-b)(\gamma(a)-\gamma(b)) \ge 0$$
 for any $a, b \in \mathbb{R}$.

Let us define the functions

$$\Lambda(t) = \frac{|t|^2}{2} \quad \text{and} \quad \Gamma(t) = \int_0^t (\gamma'(\tau))^{\frac{1}{2}} d\tau.$$

and we note that

$$\Lambda'(a-b)(\gamma(a)-\gamma(b)) \ge |\Gamma(a)-\Gamma(b)|^2 \text{ for any } a, b \in \mathbb{R}.$$
(4.3)

Indeed, for any $a, b \in \mathbb{R}$ such that a < b, and using the Jensen inequality we have

$$\Lambda'(a-b)(\gamma(a)-\gamma(b)) = (a-b)\int_b^a \gamma'(t)dt = (a-b)\int_b^a (\Gamma'(t))^2 dt \ge \left(\int_b^a \Gamma'(t)dt\right)^2 = (\Gamma(a)-\Gamma(b))^2$$

In view of (4.2) we can deduce that

In view of (4.3) we can deduce that

$$|\Gamma(|u_n(x)|) - \Gamma(|u_n(y)|)|^2 \le (|u_n(x)| - |u_n(y)|)((|u_n|u_{L,n}^{2(\beta-1)})(x) - (|u_n|u_{L,n}^{2(\beta-1)})(y)).$$
(4.4)

Putting together (4.2) and (4.4) we have

$$\Re\left(\iint_{\mathbb{R}^{6}} \frac{(u_{n}(x) - u_{n}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)})}{|x-y|^{3+2s}} \overline{(u_{n}u_{L,n}^{2(\beta-1)}(x) - u_{n}u_{L,n}^{2(\beta-1)}(y)e^{iA(\frac{x+y}{2})\cdot(x-y)})} \, dxdy\right) \ge [\Gamma(|u_{n}|)]^{2}$$

$$\tag{4.5}$$

Being $\Gamma(|u_n|) \geq \frac{1}{\beta} |u_n| u_{L,n}^{\beta-1}$ and recalling that $\mathcal{D}^{s,2}(\mathbb{R}^3,\mathbb{R}) \subset L^{2^*_s}(\mathbb{R}^3,\mathbb{R})$ (see [20]), we get

$$[\Gamma(|u_n|)]^2 \ge S_* \|\Gamma(|u_n|)\|_{L^{2^*_s}(\mathbb{R}^3)}^2 \ge \left(\frac{1}{\beta}\right)^2 S_* \||u_n|u_{L,n}^{\beta-1}\|_{L^{2^*_s}(\mathbb{R}^3)}^2.$$
(4.6)

Taking into account (4.1), (4.5) and (4.6) we obtain

$$\left(\frac{1}{\beta}\right)^{2} S_{*} \||u_{n}|u_{L,n}^{\beta-1}\|_{L^{2^{*}_{s}}(\mathbb{R}^{3})}^{2} + \int_{\mathbb{R}^{3}} V(\varepsilon x)|u_{n}|^{2} u_{L,n}^{2(\beta-1)} dx \leq \int_{\mathbb{R}^{3}} g(\varepsilon_{n} x, |u_{n}|^{2})|u_{n}|^{2} u_{L,n}^{2(\beta-1)} dx.$$
(4.7)

On the other hand, from the assumptions (g_1) and (g_2) , for any $\xi > 0$ there exists $C_{\xi} > 0$ such that

$$g(\varepsilon x, t^2)t^2 \le \xi |t|^2 + C_{\xi} |t|^{2^*_s} \text{ for all } t \in \mathbb{R}.$$
(4.8)

Taking $\xi \in (0, V_0)$ and using (4.7) and (4.8) and Lemma 2.5 we can obtain that

$$\|w_{L,n}\|_{L^{2^*_s}(\mathbb{R}^3)}^2 \le C\beta^2 \int_{\mathbb{R}^3} |u_n|^{2^*_s} u_{L,n}^{2(\beta-1)},$$
(4.9)

where we set $w_{L,n} := |u_n| u_{L,n}^{\beta-1}$. Now, take $\beta = \frac{2^*_s}{2}$ and fix R > 0. Observing that $0 \le u_{L,n} \le |u_n|$ and applying Hölder inequality we have

$$\begin{split} \int_{\mathbb{R}^{3}} |u_{n}|^{2^{*}_{s}} u_{L,n}^{2(\beta-1)} dx &= \int_{\mathbb{R}^{3}} |u_{n}|^{2^{*}_{s}-2} |u_{n}|^{2} u_{L,n}^{2^{*}_{s}-2} dx \\ &= \int_{\mathbb{R}^{3}} |u_{n}|^{2^{*}_{s}-2} (|u_{n}| u_{L,n}^{\frac{2^{*}_{s}-2}{2}})^{2} dx \\ &\leq \int_{\{|u_{n}|R\}} |u_{n}|^{2^{*}_{s}-2} (|u_{n}| u_{L,n}^{\frac{2^{*}_{s}-2}{2}})^{2} dx \\ &\leq \int_{\{|u_{n}|R\}} |u_{n}|^{2^{*}_{s}} dx\right)^{\frac{2^{*}_{s}-2}{2^{*}_{s}}} \left(\int_{\mathbb{R}^{3}} (|u_{n}| u_{L,n}^{\frac{2^{*}_{s}-2}{2}})^{2^{*}_{s}} dx\right)^{\frac{2^{*}_{s}}{2^{*}_{s}}}. \end{split}$$

$$(4.10)$$

Since $(|u_n|)$ is bounded in $H^s(\mathbb{R}^3, \mathbb{R})$, we can choose R sufficiently large such that

$$\left(\int_{\{|u_n|>R\}} |u_n|^{2^*_s} dx\right)^{\frac{2^*_s-2}{2^*_s}} \le \frac{1}{2\beta^2}.$$
(4.11)

In view of (4.9), (4.10) and (4.11) we can infer

$$\left(\int_{\mathbb{R}^3} (|u_n|u_{L,n}^{\frac{2^*_s-2}{2}})^{2^*_s}\right)^{\frac{2^*}{2^*_s}} \le C\beta^2 \int_{\mathbb{R}^3} R^{2^*_s-2} |u_n|^{2^*_s} dx < \infty$$

and letting the limit as $L \to \infty$ we obtain $|u_n| \in L^{\frac{(2_s^*)^2}{2}}(\mathbb{R}^3, \mathbb{R})$. Now, using $0 \le u_{L,n} \le |u_n|$ and taking the limit as $L \to \infty$ in (4.9) we have

$$|||u_n|||_{L^{\beta 2^*_s}(\mathbb{R}^3)}^{2\beta} \le C\beta^2 \int_{\mathbb{R}^3} |u_n|^{2^*_s + 2(\beta - 1)},$$

from which we deduce that

$$\left(\int_{\mathbb{R}^3} |u_n|^{\beta 2^*_s} dx\right)^{\frac{1}{(\beta-1)2^*_s}} \le C\beta^{\frac{1}{\beta-1}} \left(\int_{\mathbb{R}^3} |u_n|^{2^*_s+2(\beta-1)}\right)^{\frac{1}{2(\beta-1)}}$$

For $m \ge 1$ we define β_{m+1} inductively so that $2_s^* + 2(\beta_{m+1} - 1) = 2_s^*\beta_m$ and $\beta_1 = \frac{2_s^*}{2}$. Then we can see that

$$\left(\int_{\mathbb{R}^3} |u_n|^{\beta_{m+1}2^*_s} dx\right)^{\frac{1}{(\beta_{m+1}-1)2^*_s}} \le C\beta_{m+1}^{\frac{1}{\beta_{m+1}-1}} \left(\int_{\mathbb{R}^3} |u_n|^{2^*_s\beta_m}\right)^{\frac{1}{2^*_s(\beta_m-1)}}.$$

Let us define

$$D_m = \left(\int_{\mathbb{R}^3} |u_n|^{2^*_s \beta_m}\right)^{\frac{1}{2^*_s (\beta_m - 1)}}$$

and by using an iteration argument, we can find $C_0 > 0$ independent of m such that

$$D_{m+1} \le \prod_{k=1}^{m} C\beta_{k+1}^{\frac{1}{\beta_{k+1}-1}} D_1 \le C_0 D_1.$$

Passing to the limit as $m \to \infty$ we find

$$||u_n||_{L^{\infty}(\mathbb{R}^3)} \le C_0 D_1 =: K \text{ for all } n \in \mathbb{N}.$$
(4.12)

Clearly, by interpolation, we can deduce that $(|u_n|)$ strongly converges in $L^r(\mathbb{R}^3, \mathbb{R})$ for all $r \in (2, \infty)$. From the growth assumptions on g, also $g(\varepsilon x, |u_n|^2)|u_n|$ strongly converges in the same Lebesgue spaces.

In what follows, we show that $|u_n|$ is a weak subsolution to

$$\begin{cases} (-\Delta)^s v + V_0 v = g(\varepsilon x, v^2) v & \text{in } \mathbb{R}^3 \\ v \ge 0 & \text{in } \mathbb{R}^3. \end{cases}$$
(4.13)

Fix $\varphi \in C_c^{\infty}(\mathbb{R}^3, \mathbb{R})$ such that $\varphi \geq 0$, and we take $\psi_{\delta,n} = \frac{u_n}{u_{\delta,n}}\varphi$ as test function in (3.1), where we set $u_{\delta,n} = \sqrt{|u_n|^2 + \delta^2}$ for $\delta > 0$. We note that $\psi_{\delta,n} \in H^s_{\varepsilon_n}$ for all $\delta > 0$ and $n \in \mathbb{N}$. Indeed $\int_{\mathbb{R}^3} V(\varepsilon_n x) |\psi_{\delta,n}|^2 dx \leq \int_{\mathrm{supp}(\varphi)} V(\varepsilon_n x) \varphi^2 dx < \infty$. On the other hand, we can observe

$$\begin{split} \psi_{\delta,n}(x) - \psi_{\delta,n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} &= \left(\frac{u_n(x)}{u_{\delta,n}(x)}\right)\varphi(x) - \left(\frac{u_n(y)}{u_{\delta,n}(y)}\right)\varphi(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \\ &= \left[\left(\frac{u_n(x)}{u_{\delta,n}(x)}\right) - \left(\frac{u_n(y)}{u_{\delta,n}(x)}\right)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)}\right]\varphi(x) \\ &+ \left[\varphi(x) - \varphi(y)\right]\left(\frac{u_n(y)}{u_{\delta,n}(x)}\right)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \\ &+ \left(\frac{u_n(y)}{u_{\delta,n}(x)} - \frac{u_n(y)}{u_{\delta,n}(y)}\right)\varphi(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \end{split}$$

which gives

$$\begin{aligned} |\psi_{\delta,n}(x) - \psi_{\delta,n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)}|^{2} \\ &\leq \frac{4}{\delta^{2}}|u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)}|^{2}||\varphi||^{2}_{L^{\infty}(\mathbb{R}^{3})} + \frac{4}{\delta^{2}}|\varphi(x) - \varphi(y)|^{2}||u_{n}||^{2}_{L^{\infty}(\mathbb{R}^{3})} \\ &+ \frac{4}{\delta^{4}}||u_{n}||^{2}_{L^{\infty}(\mathbb{R}^{3})}||\varphi||^{2}_{L^{\infty}(\mathbb{R}^{3})}|u_{\delta,n}(y) - u_{\delta,n}(x)|^{2} \\ &\leq \frac{4}{\delta^{2}}|u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)}|^{2}||\varphi||^{2}_{L^{\infty}(\mathbb{R}^{3})} + \frac{4K^{2}}{\delta^{2}}|\varphi(x) - \varphi(y)|^{2} \\ &+ \frac{4K^{2}}{\delta^{4}}||\varphi||^{2}_{L^{\infty}(\mathbb{R}^{3})}||u_{n}(y)| - |u_{n}(x)||^{2} \end{aligned}$$

where we used

$$|z+w+k|^2 \le 4(|z|^2+|w|^2+|k|^2) \quad \forall z, w, k \in \mathbb{C},$$

 $|e^{it}| = 1$ for all $t \in \mathbb{R}$, $u_{\delta,n} \ge \delta$, $|\frac{u_n}{u_{\delta,n}}| \le 1$, (4.12) and the following inequality

$$|\sqrt{|z|^2 + \delta^2} - \sqrt{|w|^2 + \delta^2}| \le ||z| - |w|| \quad \forall z, w \in \mathbb{C}.$$

Since $u_n \in H^s_{\varepsilon_n}$, $|u_n| \in H^s(\mathbb{R}^3, \mathbb{R})$ (by Lemma 2.2) and $\varphi \in C^{\infty}_c(\mathbb{R}^3, \mathbb{R})$, we deduce that $\psi_{\delta,n} \in H^s_{\varepsilon_n}$. Therefore

$$\Re \left[\iint_{\mathbb{R}^6} \frac{(u_n(x) - u_n(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})}{|x-y|^{3+2s}} \left(\frac{\overline{u_n(x)}}{u_{\delta,n}(x)} \varphi(x) - \frac{\overline{u_n(y)}}{u_{\delta,n}(y)} \varphi(y)e^{-iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \right) dxdy \right] \\ + \int_{\mathbb{R}^3} V(\varepsilon x) \frac{|u_n|^2}{u_{\delta,n}} \varphi dx + \int_{\mathbb{R}^3} \phi_{|u_n|}^t \frac{|u_n|^2}{u_{\delta,n}} \varphi dx = \int_{\mathbb{R}^3} g(\varepsilon x, |u_n|^2) \frac{|u_n|^2}{u_{\delta,n}} \varphi dx.$$
(4.14)

Since $\Re(z) \leq |z|$ for all $z \in \mathbb{C}$ and $|e^{it}| = 1$ for all $t \in \mathbb{R}$, we get

$$\Re \left[(u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)}) \left(\frac{\overline{u_{n}(x)}}{u_{\delta,n}(x)} \varphi(x) - \frac{\overline{u_{n}(y)}}{u_{\delta,n}(y)} \varphi(y)e^{-iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \right) \right] \\
= \Re \left[\frac{|u_{n}(x)|^{2}}{u_{\delta,n}(x)} \varphi(x) + \frac{|u_{n}(y)|^{2}}{u_{\delta,n}(y)} \varphi(y) - \frac{u_{n}(x)\overline{u_{n}(y)}}{u_{\delta,n}(y)} \varphi(y)e^{-iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} - \frac{u_{n}(y)\overline{u_{n}(x)}}{u_{\delta,n}(x)} \varphi(x)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \right] \\
\ge \left[\frac{|u_{n}(x)|^{2}}{u_{\delta,n}(x)} \varphi(x) + \frac{|u_{n}(y)|^{2}}{u_{\delta,n}(y)} \varphi(y) - |u_{n}(x)| \frac{|u_{n}(y)|}{u_{\delta,n}(y)} \varphi(y) - |u_{n}(y)| \frac{|u_{n}(x)|}{u_{\delta,n}(x)} \varphi(x) \right].$$
(4.15)

Now, we can note that

$$\frac{|u_{n}(x)|^{2}}{u_{\delta,n}(x)}\varphi(x) + \frac{|u_{n}(y)|^{2}}{u_{\delta,n}(y)}\varphi(y) - |u_{n}(x)|\frac{|u_{n}(y)|}{u_{\delta,n}(y)}\varphi(y) - |u_{n}(y)|\frac{|u_{n}(x)|}{u_{\delta,n}(x)}\varphi(x) \\
= \frac{|u_{n}(x)|}{u_{\delta,n}(x)}(|u_{n}(x)| - |u_{n}(y)|)\varphi(x) - \frac{|u_{n}(y)|}{u_{\delta,n}(y)}(|u_{n}(x)| - |u_{n}(y)|)\varphi(y) \\
= \left[\frac{|u_{n}(x)|}{u_{\delta,n}(x)}(|u_{n}(x)| - |u_{n}(y)|)\varphi(x) - \frac{|u_{n}(x)|}{u_{\delta,n}(x)}(|u_{n}(x)| - |u_{n}(y)|)\varphi(y)\right] \\
+ \left(\frac{|u_{n}(x)|}{u_{\delta,n}(x)} - \frac{|u_{n}(y)|}{u_{\delta,n}(y)}\right)(|u_{n}(x)| - |u_{n}(y)|)\varphi(y) \\
= \frac{|u_{n}(x)|}{u_{\delta,n}(x)}(|u_{n}(x)| - |u_{n}(y)|)(\varphi(x) - \varphi(y)) + \left(\frac{|u_{n}(x)|}{u_{\delta,n}(x)} - \frac{|u_{n}(y)|}{u_{\delta,n}(y)}\right)(|u_{n}(x)| - |u_{n}(y)|)\varphi(y) \\
\geq \frac{|u_{n}(x)|}{u_{\delta,n}(x)}(|u_{n}(x)| - |u_{n}(y)|)(\varphi(x) - \varphi(y)) \qquad (4.16)$$

where in the last inequality we used the fact that

$$\left(\frac{|u_n(x)|}{u_{\delta,n}(x)} - \frac{|u_n(y)|}{u_{\delta,n}(y)}\right) (|u_n(x)| - |u_n(y)|)\varphi(y) \ge 0$$

because

$$h(t) = \frac{t}{\sqrt{t^2 + \delta^2}}$$
 is increasing for $t \ge 0$ and $\varphi \ge 0$ in \mathbb{R}^3 .

Observing that

$$\frac{|\frac{|u_n(x)|}{u_{\delta,n}(x)}(|u_n(x)| - |u_n(y)|)(\varphi(x) - \varphi(y))|}{|x - y|^{N+2s}} \le \frac{||u_n(x)| - |u_n(y)||}{|x - y|^{\frac{3+2s}{2}}} \frac{|\varphi(x) - \varphi(y)|}{|x - y|^{\frac{3+2s}{2}}} \in L^1(\mathbb{R}^6),$$

and $\frac{|u_n(x)|}{u_{\delta,n}(x)} \to 1$ a.e. in \mathbb{R}^3 as $\delta \to 0$, we can use (4.15), (4.16) and the Dominated Convergence Theorem to deduce that

$$\limsup_{\delta \to 0} \Re \left[\iint_{\mathbb{R}^{6}} \frac{(u_{n}(x) - u_{n}(y)e^{iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)})}{|x-y|^{3+2s}} \left(\frac{\overline{u_{n}(x)}}{u_{\delta,n}(x)}\varphi(x) - \frac{\overline{u_{n}(y)}}{u_{\delta,n}(y)}\varphi(y)e^{-iA_{\varepsilon}(\frac{x+y}{2})\cdot(x-y)} \right) dxdy \right] \\
\geq \limsup_{\delta \to 0} \iint_{\mathbb{R}^{6}} \frac{|u_{n}(x)|}{u_{\delta,n}(x)} (|u_{n}(x)| - |u_{n}(y)|)(\varphi(x) - \varphi(y)) \frac{dxdy}{|x-y|^{3+2s}} \\
= \iint_{\mathbb{R}^{6}} \frac{(|u_{n}(x)| - |u_{n}(y)|)(\varphi(x) - \varphi(y))}{|x-y|^{3+2s}} dxdy. \tag{4.17}$$

We can also see that the Dominated Convergence Theorem (we recall that $\frac{|u_n|^2}{u_{\delta,n}} \leq |u_n|$, Fatou's Lemma and $\varphi \in C_c^{\infty}(\mathbb{R}^3, \mathbb{R})$) yield

$$\lim_{\delta \to 0} \int_{\mathbb{R}^3} V(\varepsilon x) \frac{|u_n|^2}{u_{\delta,n}} \varphi dx = \int_{\mathbb{R}^3} V(\varepsilon x) |u_n| \varphi dx \ge \int_{\mathbb{R}^3} V_0 |u_n| \varphi dx \tag{4.18}$$

$$\liminf_{\delta \to 0} \int_{\mathbb{R}^3} \phi_{|u_n|}^t \frac{|u_n|^2}{u_{\delta,n}} \varphi dx \ge \int_{\mathbb{R}^3} \phi_{|u|}^t |u| \varphi dx \ge 0$$
(4.19)

and

$$\lim_{\delta \to 0} \int_{\mathbb{R}^3} g(\varepsilon x, |u_n|^2) \frac{|u_n|^2}{u_{\delta,n}} \varphi dx = \int_{\mathbb{R}^3} g(\varepsilon x, |u_n|^2) |u_n| \varphi dx.$$

$$(4.20)$$

Taking into account (4.14), (4.17), (4.19), (4.18) and (4.20) we can deduce that

$$\iint_{\mathbb{R}^6} \frac{(|u_n(x)| - |u_n(y)|)(\varphi(x) - \varphi(y))}{|x - y|^{3+2s}} dx dy + \int_{\mathbb{R}^3} V_0 |u_n|\varphi dx \le \int_{\mathbb{R}^3} g(\varepsilon x, |u_n|^2) |u_n|\varphi dx$$

for any $\varphi \in C_c^{\infty}(\mathbb{R}^3, \mathbb{R})$ such that $\varphi \ge 0$, that is $|u_n|$ is a weak subsolution to (4.13).

Now, we note that $v_n = |u_n|(\cdot + \tilde{y}_n)$ solves

$$(-\Delta)^{s} v_{n} + V_{0} v_{n} \le g(\varepsilon_{n} x + \varepsilon_{n} \tilde{y}_{n}, v_{n}^{2}) v_{n} \text{ in } \mathbb{R}^{3}.$$

$$(4.21)$$

Let us denote by $z_n \in H^s(\mathbb{R}^3, \mathbb{R})$ the unique solution to

$$(-\Delta)^s z_n + V_0 z_n = g_n \text{ in } \mathbb{R}^3, \tag{4.22}$$

where

$$g_n := g(\varepsilon_n x + \varepsilon_n \, \tilde{y}_n, v_n^2) v_n \in L^r(\mathbb{R}^3, \mathbb{R}) \quad \forall r \in [2, \infty].$$

Since (4.12) yields $||v_n||_{L^{\infty}(\mathbb{R}^3)} \leq C$ for all $n \in \mathbb{N}$, by interpolation we know that $v_n \to v$ strongly converges in $L^r(\mathbb{R}^3, \mathbb{R})$ for all $r \in (2, \infty)$, for some $v \in L^r(\mathbb{R}^3, \mathbb{R})$. From the growth assumptions on f, we have $g_n \to f(v^2)v$ in $L^r(\mathbb{R}^3, \mathbb{R})$ and $||g_n||_{L^{\infty}(\mathbb{R}^3)} \leq C$ for all $n \in \mathbb{N}$. In view of [25], we know that $z_n = \mathcal{K} * g_n$, where \mathcal{K} is the Bessel kernel, and proceeding as in [2], we can infer that $|z_n(x)| \to 0$ as $|x| \to \infty$ uniformly with respect to $n \in \mathbb{N}$. Since v_n solves (4.21) and z_n verifies (4.22), it is easy to use a comparison argument to deduce that $0 \leq v_n \leq z_n$ a.e. in \mathbb{R}^3 and for all $n \in \mathbb{N}$. Therefore $v_n(x) \to 0$ as $|x| \to \infty$ uniformly with respect to $n \in \mathbb{N}$.

Now, we are ready to give the proof of Theorem 1.1.

Proof of Thorem 1.1. In view of Lemma 3.5, we can find $(\tilde{y}_n) \subset \mathbb{R}^3$ such that $\varepsilon_n \tilde{y}_n \to y_0$ for some $y_0 \in \Lambda$ such that $V(y_0) = V_0$. Then there is r > 0 such that, for some subsequence still denoted by itself, it holds $B_r(\tilde{y}_n) \subset \Lambda$ for all $n \in \mathbb{N}$. Thus $B_{\frac{r}{\varepsilon_n}}(\tilde{y}_n) \subset \Lambda_{\varepsilon_n} n \in \mathbb{N}$, and we can deduce that $\mathbb{R}^3 \setminus \Lambda_{\varepsilon_n} \subset \mathbb{R}^3 \setminus B_{\frac{r}{\varepsilon_n}}(\tilde{y}_n)$ for any $n \in \mathbb{N}$. By using Lemma 4.1, we know that there exists R > 0 such that

$$v_n(x) < a \text{ for } |x| \ge R, n \in \mathbb{N},$$

where $v_n(x) = |u_{\varepsilon_n}|(x + \tilde{y}_n)$. Thus $|u_{\varepsilon_n}(x)| < a$ for any $x \in \mathbb{R}^N \setminus B_R(\tilde{y}_n)$ and $n \in \mathbb{N}$. Then there exists $\nu \in \mathbb{N}$ such that for any $n \ge \nu$ and $r/\varepsilon_n > R$ it holds

$$\mathbb{R}^3 \setminus \Lambda_{\varepsilon_n} \subset \mathbb{R}^3 \setminus B_{\frac{r}{\varepsilon_n}}(\tilde{y}_n) \subset \mathbb{R}^3 \setminus B_R(\tilde{y}_n),$$

which gives $|u_{\varepsilon_n}(x)| < a$ for any $x \in \mathbb{R}^3 \setminus \Lambda_{\varepsilon_n}$ and $n \ge \nu$.

Therefore, there exists $\varepsilon_0 > 0$ such that problem (3.1) admits a nontrivial solution u_{ε} for all $\varepsilon \in (0, \varepsilon_0)$. Setting $\hat{u}_{\varepsilon}(x) = u_{\varepsilon}(x/\varepsilon)$, we can see that \hat{u}_{ε} is a solution to the original problem (1.1). Finally, we investigate the behavior of the maximum points of $|u_{\varepsilon_n}|$. By using (g_1) , there exists $\gamma \in (0, a)$ such that

$$g(\varepsilon x, t^2)t^2 \le \frac{V_0}{2}t^2$$
, for all $x \in \mathbb{R}^3, |t| \le \gamma.$ (4.23)

Arguing as before, we can take R > 0 such that

$$\|u_{\varepsilon_n}\|_{L^{\infty}(B^c_R(\tilde{y}_n))} < \gamma.$$

$$(4.24)$$

Up to a subsequence, we may also assume that

$$\|u_{\varepsilon_n}\|_{L^{\infty}(B_R(\tilde{y}_n))} \ge \gamma. \tag{4.25}$$

Indeed, if (4.25) does not hold, we have $||u_{\varepsilon_n}||_{L^{\infty}(\mathbb{R}^3)} < \gamma$, and by using $J'_{\varepsilon_n}(u_{\varepsilon_n}) = 0$, (4.23) and Lemma 2.2 we can see that

$$[|u_{\varepsilon_n}|]^2 + \int_{\mathbb{R}^3} V_0 |u_{\varepsilon_n}|^2 dx \le ||u_{\varepsilon_n}||_{\varepsilon_n}^2 + \int_{\mathbb{R}^3} \phi_{|u_{\varepsilon_n}|}^t |u_{\varepsilon_n}|^2 dx = \int_{\mathbb{R}^3} g_{\varepsilon_n}(x, |u_{\varepsilon_n}|^2) |u_{\varepsilon_n}|^2 dx \le \frac{V_0}{2} \int_{\mathbb{R}^3} |u_{\varepsilon_n}|^2 dx$$

that is $||u_{\varepsilon_n}||_{H^s(\mathbb{R}^3)} = 0$ which is a contradiction. Therefore (4.25) holds true. In view of (4.24) and (4.25), we can see that the maximum points p_n of $|u_{\varepsilon_n}|$ belong to $B_R(\tilde{y}_n)$, that is $p_n = \tilde{y}_n + q_n$ for some $q_n \in B_R$. Since $\hat{u}_n(x) = u_{\varepsilon_n}(x/\varepsilon_n)$ is a solution to (1.1), we can deduce that a maximum point η_{ε_n} of $|\hat{u}_n|$ is of the type $\eta_{\varepsilon_n} = \varepsilon_n \tilde{y}_n + \varepsilon_n q_n$. Since $q_n \in B_R$, $\varepsilon_n \tilde{y}_n \to y_0$ and $V(y_0) = V_0$, we can use the continuity of V to deduce that

$$\lim_{n \to \infty} V(\eta_{\varepsilon_n}) = V_0.$$

Finally, we prove the power decay estimate of $|\hat{u}_n|$. By applying Lemma 4.3 in [25], we can find a function w such that

$$0 < w(x) \le \frac{C}{1 + |x|^{3+2s}},\tag{4.26}$$

and

$$(-\Delta)^s w + \frac{V_0}{2} w \ge 0 \text{ in } \mathbb{R}^3 \setminus B_{R_1}$$

$$(4.27)$$

for some suitable $R_1 > 0$. Invoking Lemma 4.1, we know that we can that $v_n(x) \to 0$ as $|x| \to \infty$ uniformly in $n \in \mathbb{N}$, so we can find there exists $R_2 > 0$ such that

$$h_n = g(\varepsilon_n x + \varepsilon_n \tilde{y}_n, v_n^2) v_n \le \frac{V_0}{2} v_n \text{ in } B_{R_2}^c.$$

$$(4.28)$$

Let w_n be the unique solution to

$$(-\Delta)^s w_n + V_0 w_n = h_n \text{ in } \mathbb{R}^3$$

Then $w_n(x) \to 0$ as $|x| \to \infty$ uniformly in $n \in \mathbb{N}$, and by comparison $0 \le v_n \le w_n$ in \mathbb{R}^3 . By using (4.28) we can see that

$$(-\Delta)^s w_n + \frac{V_0}{2} w_n = h_n - \frac{V_0}{2} w_n \le 0$$
 in $B_{R_2}^c$.

Set $R_3 = \max\{R_1, R_2\}$ and we define

$$a = \inf_{B_{R_3}} w > 0 \text{ and } \tilde{w}_n = (b+1)w - aw_n.$$
 (4.29)

where $b = \sup_{n \in \mathbb{N}} \|w_n\|_{L^{\infty}(\mathbb{R}^3)} < \infty$. We aim to prove that

$$\tilde{w}_n \ge 0 \text{ in } \mathbb{R}^3. \tag{4.30}$$

We begin observing that

$$\lim_{|x| \to \infty} \sup_{n \in \mathbb{N}} \tilde{w}_n(x) = 0, \tag{4.31}$$

$$\tilde{w}_n \ge ba + w - ba > 0 \text{ in } B_{R_3}, \tag{4.32}$$

$$(-\Delta)^s \tilde{w}_n + \frac{V_0}{2} \tilde{w}_n \ge 0 \text{ in } \mathbb{R}^3 \setminus B_{R_3}.$$
(4.33)

We assume by contradiction that there exists a sequence $(\bar{x}_{j,n}) \subset \mathbb{R}^3$ such that

$$\inf_{x \in \mathbb{R}^3} \tilde{w}_n(x) = \lim_{j \to \infty} \tilde{w}_n(\bar{x}_{j,n}) < 0.$$
(4.34)

Clearly, from (4.31), it follows that $(\bar{x}_{j,n})$ is bounded, and, up to subsequence, we may assume that there exists $\bar{x}_n \in \mathbb{R}^N$ such that $\bar{x}_{j,n} \to \bar{x}_n$ as $j \to \infty$. Then (4.34) implies that

$$\inf_{x \in \mathbb{R}^3} \tilde{w}_n(x) = \tilde{w}_n(\bar{x}_n) < 0.$$

$$(4.35)$$

By using the minimality of \bar{x}_n and the representation formula for the fractional Laplacian [20], we obtain that

$$(-\Delta)^{s}\tilde{w}_{n}(\bar{x}_{n}) = \frac{C(3,s)}{2} \int_{\mathbb{R}^{3}} \frac{2\tilde{w}_{n}(\bar{x}_{n}) - \tilde{w}_{n}(\bar{x}_{n} + \xi) - \tilde{w}_{n}(\bar{x}_{n} - \xi)}{|\xi|^{3+2s}} d\xi \le 0.$$
(4.36)

In view of (4.32) and (4.34), we have $\bar{x}_n \in \mathbb{R}^3 \setminus B_{R_3}$, and by using (4.35) and (4.36), we can conclude that

$$(-\Delta)^s \tilde{w}_n(\bar{x}_n) + \frac{V_0}{2} \tilde{w}_n(\bar{x}_n) < 0$$

which is impossible due to (4.33). Therefore, (4.30) is true and by using (4.26) and $v_n \leq w_n$ we have

$$0 \le v_n(x) \le w_n(x) \le \frac{(b+1)}{a} w(x) \le \frac{\tilde{C}}{1+|x|^{3+2s}} \text{ for all } n \in \mathbb{N}, x \in \mathbb{R}^3,$$

for some constant $\tilde{C} > 0$. Taking in mind the definition of v_n , we can infer that

$$\hat{u}_{n}|(x) = |u_{\varepsilon_{n}}|\left(\frac{x}{\varepsilon_{n}}\right) = v_{n}\left(\frac{x}{\varepsilon_{n}} - \tilde{y}_{n}\right)$$

$$\leq \frac{\tilde{C}}{1 + |\frac{x}{\varepsilon_{n}} - \tilde{y}_{\varepsilon_{n}}|^{3+2s}}$$

$$= \frac{\tilde{C}\varepsilon_{n}^{3+2s}}{\varepsilon_{n}^{3+2s} + |x - \varepsilon_{n}\,\tilde{y}_{\varepsilon_{n}}|^{3+2s}}$$

$$\leq \frac{\tilde{C}\varepsilon_{n}^{3+2s}}{\varepsilon_{n}^{3+2s} + |x - \eta_{\varepsilon_{n}}|^{3+2s}}.$$

V. AMBROSIO

References

- C.O. Alves, G.M. Figueiredo, M.F. Furtado, Multiple solutions for a nonlinear Schrödinger equation with magnetic fields, Comm. Partial Differential Equations 36 (2011), 1565–1586. 1, 3
- [2] C.O. Alves and O.H. Miyagaki, Existence and concentration of solution for a class of fractional elliptic equation in R^N via penalization method, Calc. Var. Partial Differential Equations 55 (2016), art. 47, 19 pp. 2, 3, 6, 23
- [3] A. Ambrosetti and P.H. Rabinowitz, Dual variational methods in critical point theory and applications, J. Funct. Anal. 14 (1973), 349–381. 3, 7, 13
- [4] V. Ambrosio, Multiplicity of positive solutions for a class of fractional Schrödinger equations via penalization method, Ann. Mat. Pura Appl. (4) 196 (2017), no. 6, 2043–2062. 2
- [5] V. Ambrosio, Concentrating solutions for a class of nonlinear fractional Schrödinger equations in R^N, to appear in Rev. Mat. Iberoam., arXiv:1612.02388.
- [6] V. Ambrosio and P. d'Avenia, Nonlinear fractional magnetic Schrödinger equation: existence and multiplicity, J. Differential Equations 264 (2018), no. 5, 3336–3368. 2, 3, 4, 5
- [7] G. Arioli and A. Szulkin, A semilinear Schrödinger equation in the presence of a magnetic field, Arch. Ration. Mech. Anal. 170 (2003), 277–295. 1
- [8] A. Azzollini, P. d'Avenia, and A. Pomponio, On the Schrödinger-Maxwell equations under the effect of a general nonlinear term, Ann. Inst. H. Poincaré Anal. Non Linéaire 27 (2010), no. 2, 779–791. 3
- [9] V. Benci and D. Fortunato, An eigenvalue problem for the Schrödinger-Maxwell equations, Topol. Methods Nonlinear Anal. 11 (1998), no. 2, 283–293. 2
- [10] R. Benguria, H. Brezis, E.H. Lieb, The Thomas-Fermi-von Weizsäker theory of atoms and molecules, Comm. Math. Phys. 79 (1981) 167–180. 2
- [11] J. Byeon and Z. Q. Wang, Standing waves with a critical frequency for nonlinear Schrödinger equations, II, Calc. Var. Partial Differential Equations, 18 (2003), 207–219. 3
- [12] S. Cingolani, Semiclassical stationary states of nonlinear Schrödinger equations with an external magnetic field J. Differential Equations 188 (2003), 52–79. 1
- [13] S. Cingolani, S. Secchi, Semiclassical states for NLS equations with magnetic potentials having polynomial growths, J. Math. Phys. 46, 19 (2005). 3
- [14] S. Cingolani, S. Secchi and M. Squassina, Semi-classical limit for Schrödinger equations with magnetic field and Hartree-type nonlinearities, Proc. Roy. Soc. Edinburgh Sect. A 140 (2010), no. 5, 973–1009. 1
- [15] T. D'Aprile and J. Wei, On bound states concentrating on spheres for the Maxwell-Schrödinger equation, SIAM J. Math. Anal. 37 (2005), no. 1, 321–342. 3
- [16] J. Dávila, M. del Pino, S. Dipierro and E. Valdinoci, Concentration phenomena for the nonlocal Schrödinger equation with Dirichlet datum, Anal. PDE 8 (2015), no. 5, 1165–1235. 2
- [17] J. Dávila, M. del Pino and J. Wei, Concentrating standing waves for the fractional nonlinear Schrödinger equation, J. Differential Equations 256 (2014), no. 2, 858–892. 2
- [18] P. d'Avenia, M. Squassina, Ground states for fractional magnetic operators, to appear in ESAIM Control Optim. Calc. Var., doi: 10.1051/cocv/2016071. 1, 2, 4, 5
- [19] M. del Pino and P. L. Felmer, Local Mountain Pass for semilinear elliptic problems in unbounded domains, Calc. Var. Partial Differential Equations, 4 (1996), 121–137. 1, 2, 3, 6
- [20] E. Di Nezza, G. Palatucci, E. Valdinoci, Hitchhiker's guide to the fractional Sobolev spaces, Bull. Sci. math. 136 (2012), 521–573. 4, 12, 20, 25
- [21] S. Dipierro, M. Medina and E. Valdinoci, Fractional elliptic problems with critical growth in the whole of Rⁿ, Appunti. Scuola Normale Superiore di Pisa (Nuova Serie) [Lecture Notes. Scuola Normale Superiore di Pisa (New Series)], 15. Edizioni della Normale, Pisa, 2017. viii+152 pp. 14
- [22] M. Esteban and P.L. Lions, Stationary solutions of nonlinear Schrödinger equations with an external magnetic field, Partial differential equations and the calculus of variations, Vol. I, 401–449, Progr. Nonlinear Differential Equations Appl., 1, Birkhäuser Boston, Boston, MA, 1989. 1
- [23] M. Fall and S. Jarohs, Overdetermined problems with fractional Laplacian, ESAIM Control Optim. Calc. Var. 21 (2015), no. 4, 924–938. 14
- [24] M. M. Fall, F. Mahmoudi and E. Valdinoci, Ground states and concentration phenomena for the fractional Schrödinger equation, Nonlinearity 28 (2015), no. 6, 1937–1961. 2
- [25] P. Felmer, A. Quaas, J.Tan, Positive solutions of the nonlinear Schrödinger equation with the fractional Laplacian, Proc. Roy. Soc. Edinburgh Sect. A 142 (2012), 1237–1262. 3, 4, 5, 14, 23, 24
- [26] A. Fiscella, A. Pinamonti and E. Vecchi, Multiplicity results for magnetic fractional problems, J. Differential Equations 263 (2017), 4617–4633. 2
- [27] A. R. Giammetta, Fractional Schrödinger-Poisson-Slater system in one dimension, preprint arXiv:1405.2796. 3

- [28] X. He, Multiplicity and concentration of positive solutions for the Schrödinger-Poisson equations, Z. Angew. Math. Phys. 62 6 (2011) 869–889. 3
- [29] Y. He and G. Li, Standing waves for a class of Schrödinger-Poisson equations in ℝ³ involving critical Sobolev exponents, Ann. Acad. Sci. Fenn. Math. 40 (2015), no. 2, 729–766. 3
- [30] F. Hiroshima, T. Ichinose and J. Lörinczi, Kato's Inequality for Magnetic Relativistic Schrödinger Operators, preprint arXiv:1604.03933. 3
- [31] T. Ichinose, Magnetic relativistic Schrödinger operators and imaginary-time path integrals, Mathematical physics, spectral theory and stochastic analysis, 247–297, Oper. Theory Adv. Appl. 232, Birkhäuser/Springer Basel AG, Basel, 2013. 1
- [32] T. Kato, Schrödinger operators with singular potentials, Proceedings of the International Symposium on Partial Differential Equations and the Geometry of Normed Linear Spaces (Jerusalem, 1972), Israel J. Math. 13, 135– 148 (1973). 3
- [33] K. Kurata, Existence and semi-classical limit of the least energy solution to a nonlinear Schrödinger equation with electromagnetic fields, Nonlinear Anal. 41 (2000), 763–778. 1, 3
- [34] L.D. Landau and E.M. Lifshitz, Quantum mechanics, Pergamon Press, (1977). 1
- [35] N. Laskin, Fractional quantum mechanics and Lévy path integrals, Physics Letters A 268 (2000) 298–305. 2
- [36] E. H. Lieb and M. Loss, Analysis. Graduate Studies in Mathematics, 14. American Mathematical Society, Providence, RI, 1997. xviii+278 pp. 1, 6, 9
- [37] Z. Liu and J. Zhang, Multiplicity and concentration of positive solutions for the fractional Schrödinger-Poisson systems with critical growth, ESAIM Control Optim. Calc. Var. 23 (2017), no. 4, 1515–1542. 3, 16
- [38] P. Markowich, C. Ringhofer, C. Schmeiser, Semiconductor Equations, Springer-Verlag, New York, 1990. 2
- [39] G. Molica Bisci, V. Rădulescu and R. Servadei, Variational Methods for Nonlocal Fractional Problems, Cambridge University Press, 162 Cambridge, 2016. 4
- [40] J. Moser, A new proof of De Giorgi's theorem concerning the regularity problem for elliptic differential equations, Comm. Pure Appl. Math. 13 (1960), 457–468. 4
- [41] E. Murcia and G. Siciliano, Positive semiclassical states for a fractional Schrödinger-Poisson system, Differential Integral Equations 30 (2017), no. 3-4, 231–258. 3
- [42] G. Palatucci, A. Pisante, Improved Sobolev embeddings, profile decomposition, and concentration-compactness for fractional Sobolev spaces, Calc. Var. Partial Differential Equations 50 (2014), 799–829. 15
- [43] A. Pinamonti, M. Squassina, E. Vecchi, Magnetic BV functions and the Bourgain-Brezis-Mironescu formula, preprint, https://arxiv.org/abs/1609.097142 2, 10. 1
- [44] P. H. Rabinowitz, On a class of nonlinear Schrödinger equations Z. Angew. Math. Phys. 43 (1992), 270–291. 2
 [45] D. Ruiz, The Schrödinger-Poisson equation under the effect of a nonlinear local term, J. Funct. Anal. 237 (2006), no. 2, 655–674. 3
- [46] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, Comm. Pure Appl. Math., 60 (2007), no. 1, 67–112. 14
- [47] M. Squassina and B. Volzone, Bourgain-Brezis-Mironescu formula for magnetic operators, C. R. Math. 354, 825–831 (2016). 1
- [48] K. Teng, Concentrating bounded states for fractional Schrödinger-Poisson system, preprint arXiv:1710.03495. 3, 14
- [49] J. Wang, L. Tian, J. Xu, F. Zhang, Existence and concentration of positive solutions for semilinear Schrödinger-Poisson systems in R³, Calc. Var. Partial Differential Equations 48 (2013) 275–276. 3
- [50] M. Willem, *Minimax theorems*, Progress in Nonlinear Differential Equations and their Applications 24, Birkhäuser Boston, Inc., Boston, MA, 1996. 13
- [51] J. Zhang, M. do Ó, and M. Squassina, Fractional Schrödinger-Poisson Systems with a General Subcritical or Critical Nonlinearity, Adv. Nonlinear Stud. 16 (2016), no. 1, 15D30. 3, 14
- [52] B. Zhang, M. Squassina, X. Zhang, Fractional NLS equations with magnetic field, critical frequency and critical growth, to appear in Manuscripta Math., doi:10.1007/s00229-017-0937-4. 2
- [53] L. Zhao and F. Zhao, On the existence of solutions for the Schrödinger-Poisson equations, J. Math. Anal. Appl. 346 (2008), no. 1, 155–169. 3
- [54] A. Zhu and X. Sun, Multiple solutions for Schrödinger-Poisson type equation with magnetic field, J. Math. Phys. 56 (2015), no. 9, 091504, 15 pp. 3

VINCENZO AMBROSIO DEPARTMENT OF MATHEMATICS EPFL SB CAMA STATION 8 CH-1015 LAUSANNE, SWITZERLAND *E-mail address*: vincenzo.ambrosio2@unina.it